



HAL
open science

Neonicotinoids: Still present in farmland birds despite their ban

Elva Fuentes, Agathe Gaffard, Anaïs Rodrigues, Maurice Millet, Vincent Bretagnolle, Jérôme Moreau, Karine Monceau

► To cite this version:

Elva Fuentes, Agathe Gaffard, Anaïs Rodrigues, Maurice Millet, Vincent Bretagnolle, et al.. Neonicotinoids: Still present in farmland birds despite their ban. *Chemosphere*, 2023, 321, pp.138091. 10.1016/j.chemosphere.2023.138091 . hal-03990116

HAL Id: hal-03990116

<https://hal.science/hal-03990116>

Submitted on 22 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Neonicotinoids: Still present in farmland birds despite their ban**

2

3

4 Elva Fuentes^{1‡}, Agathe Gaffard^{1‡}, Anaïs Rodrigues², Maurice Millet², Vincent Bretagnolle^{1,3},
5 Jérôme Moreau⁴ and Karine Monceau^{1*}

6

7

8 ¹ UMR 7372, Centre d'Études Biologiques de Chizé, La Rochelle Université & CNRS, 79360
9 Villiers en Bois, France

10 ² Université de Strasbourg, CNRS-UMR 7515, ICPEES, 67087 Strasbourg cedex 2, France

11 ³ LTSER "Zone Atelier Plaine & Val de Sèvre", CNRS, 79360 Villiers-en-Bois, France

12 ⁴ UMR CNRS 6282 Biogéosciences, Équipe Écologie Évolutive, Université de Bourgogne-
13 Franche-Comté, 21000 Dijon, France

14

15 ‡ Authors contributed equally (in alphabetical order)

16

17 *Corresponding author: karine.monceau@univ-lr.fr

18

19 **Abstract**

20

21 Neonicotinoids (neonics) are the most widely used insecticides worldwide and are considered
22 to be of low risk to non-target organisms such as vertebrates. Further, they are reported to
23 be rapidly excreted and metabolized, reducing their potential toxicity. Nevertheless, growing
24 evidence of adverse effects of neonics on farmland bird species raise questions about the
25 purported harmless nature of these pesticides. We attempted to search for pesticide residues
26 in species of different trophic levels and at different life stages, by using multiple bird
27 monitoring programs on a Long-Term Socio-Ecological Research (LTSER) platform. Three
28 passerine birds—the blackbird (*Turdus merula*), ciril bunting (*Emberiza cirilus*), and common
29 nightingale (*Luscinia megarhynchos*)—that feed on seeds and invertebrates were monitored
30 during their reproductive period, and the grey partridge (*Perdix perdix*) that feeds on seeds
31 was monitored during its wintering period. We also monitored chicks of an apex predator—
32 the Montagu’s harrier (*Circus pygargus*)—that preys mostly upon common voles but also upon
33 insects. We found that the birds’ blood samples showed presence of residues of five neonics:
34 three banned since 2018 in France—clothianidin, thiacloprid, and thiamethoxam—and two—
35 dinotefuran and nitenpyram—used for veterinary purposes only. While none of these neonics
36 was detected in blackbirds, all were present in grey partridges. Clothianidin was detected in
37 all species, except blackbirds. Concentrations of the three banned neonics were similar or
38 higher than concentrations found in birds monitored elsewhere before the ban. These findings
39 raise questions about the persistence of neonics within the environment and the mode of
40 exposure to wild fauna. Future investigations on the sublethal effects of these neonics on life-
41 history traits of these farmland birds may help in providing a better understanding of the

42 effects of exposure of bird populations to these insecticides, and also to the consequent effect
43 on human health.

44

45 Keywords:

46 Biomonitoring, farmland birds, passerines, pesticides, raptors.

47 **1. Introduction**

48

49 During the last century, the need for feeding the growing human population worldwide
50 has led to an intensification of agricultural practices, including an extensive use of pesticides.
51 Despite their use for millennia, several pieces of evidence implicate pesticides in the global
52 decline in biodiversity (Wood & Goulson, 2017; Stanton et al., 2018; Moreau et al., 2022a).
53 Moreover, other studies identify them as the cause of certain diseases in humans (Köhler &
54 Triebkorn, 2013). As all living organisms in a given area share the same environment,
55 protecting biodiversity by reducing pesticides inputs means also reducing humans' exposure
56 to pesticides, and this is necessary to ensure our own health and safety (*One Health* concept).
57 One way is thus to capitalize on wildlife monitoring schemes to better understand the risk of
58 pesticide exposure for humans (Moreau et al., 2022a). In that respect, wild bird species are
59 valuable candidates as they are well-monitored worldwide, being involved in long-term
60 banding programs over decades or more, which have highlighted a global declining trend in
61 several taxa and especially in farmland birds, e.g., in France, Sweden, United Kingdom, US,
62 Europe, and North America (Wretenberg et al., 2006; Comolet-Tirman et al., 2015; Stanton et
63 al., 2018; Rosenberg et al., 2019; Li et al., 2020; Burns et al., 2021; DEFRA, 2021). Pesticide use
64 has been often identified as a major component responsible for this decline (Campbell et al.,
65 1997; Geiger et al., 2010; Mineau & Whiteside, 2013; Chiron et al., 2014; Tassin de Montaignu
66 & Goulson, 2020). Recent studies, for instance, showed a negative relationship between the
67 abundance of northern bobwhites (*Colinus virginianus*) and their exposure to neonicotinoid
68 insecticides (neonics) from 1978 to 2012 (Ertl et al., 2018a). Similarly, a wide-panel dataset
69 regarding neonics use and birds' population trends revealed a significant negative impact of

70 this family of pesticides on avian biodiversity from 2003 to 2010 in the Netherlands (Hallmann
71 et al., 2014) and from 2008 to 2014 in USA (Li et al., 2020).

72 Neonics are insecticides developed in the 1970s, with the first patent dating back to
73 1977 for nithiazine (rapidly abandoned due to its poor stability), followed by patents for
74 imidacloprid and thiacloprid in 1985, nitenpyram in 1988, acetamiprid and clothianidin in
75 1989, thiamethoxam in 1992, and dinotefuran in 1994 (Tomizawa & Casida, 2005). Neonics
76 target the neural function and are competitive agonists of nicotinic acetylcholine receptors
77 (nAChR), which increase specificity to insects and not vertebrates (Tomizawa & Casida, 2003).
78 In fact, in contrast to other insecticides such as carbamates and organophosphorous that also
79 target the neural function by inhibiting the acetylcholinesterase (AChE) enzyme—an
80 ubiquitous enzyme in the animal kingdom (review in Grue et al., 1997; Story & Cox, 2001;
81 Walker, 2003; Mitra et al., 2011)—neonics were supposed to have minimal effects and a low
82 toxicological impact on vertebrates and consequently on birds, due to the lower number of
83 nAChRs they have and the lower sensitivity of their nAChRs compared to those of insects
84 (Tomizawa & Casida, 2003, 2005; Ihara et al., 2017; Casida, 2018). Moreover, they were
85 claimed to be rapidly metabolized and excreted, in a few hours (Bishop et al., 2018, 2020;
86 Casida, 2018; Bean et al., 2019; English et al., 2021; Pan et al., 2022), but some reports
87 question their fate along the trophic chain as they have been found in insectivorous birds,
88 granivorous birds, piscivorous birds, and birds of prey (see references in the supplementary
89 materials Table S1). Under experimental conditions, they seem to accumulate in the liver
90 (Lopez-Antia et al., 2015a) and to be detectable in different organs and tissues, although they
91 (at least for imidacloprid and thiamethoxam) seem to be rapidly cleared from birds' organism
92 (Bean et al., 2019; Pan et al., 2022). Several studies showed various effects of neonics
93 exposure on birds at different physiological levels (review in Gibbons et al., 2015 and Moreau

94 et al., 2022a). Although neonics do not act directly on AChE, they may induce neuronal
95 degeneration, which affects AChE activity (Abu Zeid et al., 2019; Rawi et al., 2019), altering
96 more complex functions such as learning and migration behaviour (Eng. et al., 2017) but also
97 having sublethal effects on important functions of the organism such as the haematocrit,
98 antioxidant defences, immunity, or fecundity (Lopez-Antia et al., 2013, 2015a, 2015b;
99 Tokumoto et al., 2013; Hoshi et al., 2014; Mohanty et al., 2017; Humann-Guillemot et al.,
100 2019; Lv et al., 2020).

101 Since the 1990s, the use of neonics has become widespread, making them the most
102 widely used class of insecticide worldwide, mostly for coating seeds, despite being shown to
103 impact non-target species, including humans (Tomizawa & Casida, 2005; Casida & Durkin,
104 2013; Gibbons et al., 2015; Simon-Delso et al., 2015; Henry et al., 2015; Wood & Goulson,
105 2017; Casida, 2018; Thompson et al., 2020; Zhang et al., 2022). In Europe, despite an EU
106 moratorium in 2014, neonics were still detectable in bee-attractive crop nectar until 2018 at
107 least (Wintermantel et al., 2020). Indeed, their degradation in soil (measured by DT₅₀ which is
108 the Detection Time 50% representing the time to detect a 50% decrease in pesticide
109 concentration) can take quite a long time, up to more than 6900 days (i.e., 19 years for
110 clothianidin; see Table 2 in Thompson et al., 2020). Therefore, banning harmful neonics does
111 not necessarily eradicate the problem of exposure. In EU, dinotefuran and nitenpyram have
112 never been considered for use in phytopharmaceutical products (PPP; July 2022, EU Pesticides
113 database: https://food.ec.europa.eu/plants/pesticides_en) but are commonly used in
114 veterinary medicine. In France, neonics are banned for outdoor use and in PPP since
115 September 2018 (Décret n° 2018-675, July 2018), except for emergency authorized use of
116 thiamethoxam and imidacloprid on sugar beet crops in production areas (EFSA, 2021).
117 Nonetheless, the use of neonics is still allowed in veterinary medicine.

118 In the present study, we use different bird monitoring programs on a long-term socio-
119 ecological research (LTSER) platform and a multiresidue analysis (Rodrigues et al., 2023) to
120 evaluate the presence of neonics in an intensive farmland area where acetamiprid,
121 clothianidin, imidacloprid, thiacloprid, and thiamethoxam were banned for 3 years for
122 agricultural use. Nitenpyram and dinotefuran were also included in the screening to control
123 for potential exposure of wild fauna to veterinary products. As the method was not specifically
124 developed for neonics detection, acetamiprid and imidacloprid, were not detectable among
125 the other 104 pesticide compounds. We selected five different species for their different
126 ecology: (i) three passerine birds during their reproductive period: the blackbird (*Turdus*
127 *merula*), ciril bunting (*Emberiza cirilus*), and common nightingale (*Luscinia megarhynchos*) that
128 feed on seeds and invertebrates; (ii) the grey partridge (*Perdix perdix*) caught during its
129 wintering period when it feeds only seeds; and (iii) an apex predator species, namely, the
130 Montagu's harrier (*Circus pygargus*) that preys mostly upon common voles but also upon
131 orthopterans. For this fifth species, we focused on chicks that are fed by their parents during
132 the rearing period. Beyond the interest of their contrasted ecologies, farmland bird species
133 including buntings, blackbirds or raptors have been shown to be exposed to neonics, however,
134 a limited number were subject of measures in blood samples (Lennon et al., 2020a).
135 Additionally, grey partridge is recognised to be a focal species for pesticide risk assessment
136 (Millot et al., 2017; Bonneris et al., 2018). Here, we aimed at monitoring potential exposure
137 of multiple bird species in an area where there is presumably no use of neonics for agricultural
138 purposes for 3 years and where nitenpyram used in veterinary medicine has never been
139 assessed to our knowledge. We thus collected blood samples from all individuals to determine
140 the presence and measure the level of exposure to neonics. We selected blood (whole blood,
141 i.e., red blood cells and plasma) for the analyses in order to focus on the effects of short-term

142 exposure only (Espín et al., 2016), so that potential exposure of the migratory species (i.e.,
143 common nightingales, Montagu's harriers, and blackbirds) at wintering areas can be
144 neglected.

145

146 **2. Materials and methods**

147

148 **2.1. Study area**

149

150 The study site is located in southwestern France (46°11'N, 0°28'W, Figure 1), in the Long-Term
151 Socio-Ecological Research Zone Atelier Plaine & Val de Sèvres (LTSER ZAPVS), a 450 km² area
152 where the soil occupancy and the agricultural practices are monitored each year since 1994
153 (Bretagnolle et al., 2018). In this intensive farming area, winter cereal crops accounted for
154 ~41% (wheat: 33.8% and corn: 9.6%) of the area under cultivation; in addition, there were
155 sunflower (10.4%), oilseed rape (8.3%), pea (2%), and meadows (13.5%) (average coverage
156 between 2009 and 2016, Bretagnolle et al., 2018). In this area, organic farming (no pesticide
157 use) is carried out in 18% of the agricultural area. Detailed data on pesticide applications was
158 not available, however, until 2018 imidacloprid was used on cereal crops in the study area,
159 thiacloprid, thiamethoxam and clothianidin for their part were mainly used in maize, oilseed
160 rape and cereal crops. Different monitoring studies performed in this area showed the
161 transfer of neonics in several compartments. For instance, imidacloprid was found in soils,
162 oilseed rape nectar, earthworms, and small mammals in this area (Henry et al., 2015;
163 Wintermantel et al., 2020; Pelosi et al., 2021; Fritsch et al. 2022). Thiacloprid was detected in
164 soils, earthworms and small mammals' hair while thiamethoxam was detected in nectar and

165 soils, and clothianidin in nectar and small mammals' hair (Wintermantel et al., 2020; Pelosi et
166 al., 2021; Fritsch et al. 2022). Acetamiprid has been detected in small mammals' hair (Fritsch
167 et al., 2022) despite its use is mainly for market gardening such as tomatoes, squash, and
168 melon cultures which are not present in a substantial surface of the study area. All five neonics
169 are banned for agricultural use in France since 2018, except imidacloprid and thiamethoxam
170 for emergency authorizations on sugar beet crops. Still, sugar beet crops are only present in
171 this area for their first year (seed production), and the use of neonics is banned for this
172 purpose. Therefore, at the moment when birds were caught (2020–2022, see section 2.2.1),
173 no neonics were being used for agricultural purposes. However, imidacloprid, dinotefuran,
174 and nitenpyram may be in domestic use as veterinary treatment for domestic animals (cats,
175 dogs, ferrets, and bunnies). There is no known screening of nitenpyram in the study area and
176 dinotefuran has only been investigated in one previous study but was not detected in small
177 mammals' hair sampled (Fritsch et al., 2022).

178

179 **2.2. Model species and blood collection**

180

181 2.2.1. Model species

182

183 2.2.1.1. *Adult passerine birds*

184

185 The passerine species were sampled during the reproductive period from mid-April to end of
186 June 2021. Birds were caught using net-trapping, following the same methodology as Moreau
187 et al. (2022b). Among the 17 species trapped in this area, we selected three, namely, ciril
188 buntings (N = 31), blackbirds (N = 64), and common nightingales (N = 34) because these species

189 were among the biggest of those captured, which allowed us to draw enough blood for
190 analysis without risking any individual's health. They were also the most caught in the study
191 area allowing us to make a large spatial screening.

192

193 2.2.1.2. *Adult grey partridges*

194

195 Grey partridges were caught during two consecutive years from November 2020 to February
196 2021 (winter 2020–2021) and from December 2021 to March 2022 (winter 2021–2022) in the
197 study area. These individuals, even when caught in the wild, are gamebirds and have been
198 probably raised in captivity before being released for hunting purposes; however, knowing
199 the proportion of captive-born vs. wild-born partridges is almost impossible as banding before
200 release is not mandatory. In winter, grey partridges are mostly herbivorous and granivorous,
201 living in large coveys in winter crops and sleeping in ploughed fields at night. Thermal
202 binoculars were used to spot them at nightfall and birds were then captured in the dark by
203 dazzling them and using a landing net (i.e., a method inspired by Eurasian Woodcock catching
204 technique; Williams, 2015). Blood samples were then collected, and birds were immediately
205 released. A total of 23 and 31 partridges were caught during the 2020–2021 and 2021–2022
206 winters, respectively.

207

208 2.2.1.3. *Montagu's harrier chicks*

209

210 The Montagu's harrier has been monitored in the ZAPVS since 1994 (Bretagnolle et al., 2018).
211 In this intensive agricultural area, they nest mainly on the ground of cereal crops and lay up
212 to eight eggs (Arroyo et al., 1998; Millon et al., 2008). The incubation period lasts 29 days, and

213 the rearing period is between 30 and 35 days (Arroyo et al., 2007). The mean productivity in
214 this site is 2.05 fledglings per breeding attempt (Arroyo et al., 2004), this success depending
215 mainly on the availability of its main prey, the common vole (*Microtus arvalis*) although in case
216 of poor vole availability, harriers may also feed on orthopterans (Salamolard et al., 2000; Butet
217 & Leroux, 2001). Blood samples of chicks that were 26 ± 2 days old were collected from June
218 2021 to early August 2021. Fifty-five chicks from 22 nests were sampled.

219

220 2.2.2. Blood sampling procedure

221

222 For all species, blood samples were collected in 2021, and in 2020 and 2022 for grey
223 partridges, on wild individuals included in different monitoring programs (see details in
224 section 2.2.1). For all of them, blood sampling was conducted by puncturing the brachial vein
225 using a sterile needle and using heparinized capillaries to collect 50 μ L of blood. Blood samples
226 were placed in Eppendorf tubes and kept refrigerated (0–5°C), before being returned to the
227 laboratory where they were stored at -20°C for further analyses.

228

229 **2.3. Neonics analysis**

230

231 Neonics extractions were conducted following the method reported by Rodrigues *et al.*
232 (2023; see also Table S2 in supplementary materials for a description of the neonics). Briefly,
233 blood samples were defrosted and weighed, and a mixture of 2 mL of dichloromethane and
234 ethyl acetate (1:1) was added to each sample, followed by homogenisation by using a vortex
235 for 1 min. Extracts were then sonicated for 10 min. This sonication step was repeated three
236 successive times. After each sonication step, a centrifugation step of 5 min was performed,

237 and supernatants were collected, pooled, and then gently evaporated under a fume hood until
238 a final volume of 500 μ L. The extract was collected and stored at -20°C until the analyses to
239 determine pesticide levels were performed by liquid chromatography coupled to tandem
240 mass spectrometry (LC/MSMS) using multiple reactions monitoring (MRM) for quantification.

241 LC/MSMS analyses were conducted with a Thermo Scientific TSQ Quantum Access
242 Triple Quadrupole Mass Spectrometer operating in heated positive electrospray ionization
243 mode (HESI+) coupled with a Thermo Accela 1250 pump and a Thermo Combi Pal
244 autosampler. Analyses were performed on a Nucleodur C₁₈ Pyramid column (150 mm \times 3 mm,
245 3 μ m). Samples were analysed in the gradient mode using a mobile phase composed of water
246 and acetonitrile with both containing 0.1% formic acid.

247 The multiresidue analysis comprised detection and quantification of 104 pesticide
248 molecules, including five neonics—clothianidin, dinotefuran, nitenpyram, thiacloprid, and
249 thiamethoxam—performed using the MRM detection mode. The source was operated in the
250 positive ionization mode with a spray voltage of 4500 V and the same spray and capillary
251 temperatures of 300 °C each. Nitrogen was used as the sheath and auxiliary gas (20 and 10
252 arbitrary units), while argon was used as the collision gas (1.5 arbitrary units). Two precursor
253 product ion transitions for each analyte and internal standards were used for quantification.
254 The transitions selected for MSMS analysis and retention times are shown in supplementary
255 materials (Table S3). Data were acquired and processed using Excalibur software.

256 The limits of detection (LOD) and quantification (LOQ) represent three and ten times
257 the ratio of the average noise height on either side of a known amount of a compound's peak
258 to the peak height, respectively. The objective was to determine the minimum peak heights
259 that can be used to distinguish a compound's peak from the noise on either side of the peak.

260 The LOD and LOQ determined for each sample type are presented in Table S3. LODs varied
261 from 0.001 to 0.012 $\text{pg}\cdot\mu\text{L}^{-1}$ and LOQs from 0.005 to 0.041 $\text{pg}\cdot\mu\text{L}^{-1}$.

262

263 **3. Results**

264

265 No neonics were detected in blackbirds. For the other species, all 5 neonics searched, namely,
266 clothianidin, dinotefuran, nitenpyram, thiacloprid, and thiamethoxam were detected. A
267 summary of the concentrations and occurrence of these neonics in our sampled bird species
268 is provided in Table 1, and detailed neonicotinoid distribution among each individual is
269 provided in Figure 2. Clothianidin was found in all species (except blackbirds), with average
270 concentrations ranging from 0.05 $\text{pg}\cdot\mu\text{L}^{-1}$ in common nightingales to 951.60 $\text{pg}\cdot\mu\text{L}^{-1}$ in
271 Montagu's harriers. The number of individuals that exhibited concentrations higher than the
272 LOQ ranged from ~5% among Montagu's harriers up to ~26% among grey partridges sampled
273 during the 2021–2022 winter. No clothianidin was detected in grey partridges sampled in the
274 2020–2021 winter. Thiacloprid was detected in all circl buntings and common nightingales but
275 always at concentrations below the LOQ. It was detected in one Montagu's harrier chick (89.58
276 $\text{pg}\cdot\mu\text{L}^{-1}$) and one grey partridge (0.07 $\text{pg}\cdot\mu\text{L}^{-1}$) in the 2020–2021 winter. Thiamethoxam was
277 detected in both passerine species and in grey partridges from both winters, with
278 concentrations ranging from 0.06 $\text{pg}\cdot\mu\text{L}^{-1}$ in a common nightingale to 23.73 $\text{pg}\cdot\mu\text{L}^{-1}$ in a grey
279 partridge from the 2020–2021 winter. The number of individuals that exhibited
280 concentrations higher than the LOD ranged from ~4% among common nightingales up to 12%
281 among circl buntings. Dinotefuran and nitenpyram were only detected in grey partridges: In
282 the first winter (2020–2021), ~13% and ~87% of the individuals exhibited dinotefuran and
283 nitenpyram concentrations higher than the LOD, with the average concentrations being 6.20

284 and 23.10 $\text{pg}\cdot\mu\text{L}^{-1}$, respectively. In the second winter (2021–2022), dinotefuran and
285 nitenpyram were detected with concentrations higher than the LOD in ~32 and ~94% of the
286 individuals, respectively, with the average concentrations being 7.85 and 18.29 $\text{pg}\cdot\mu\text{L}^{-1}$,
287 respectively.

288

289 **4. Discussion**

290

291 In the last few decades, neonics have attracted considerable interest (Klingelhöfer et al.,
292 2022), especially because of the rising concern about their effects on non-target species such
293 as honeybees (Wood & Goulson, 2017) and humans (Cimino et al., 2017; Han et al., 2018).
294 Because of their chemical properties such as their half-life in the soil (DT_{50}), solvability, and
295 leaching potential, and because of their extensive use for agricultural purposes, the
296 assessment of their presence in farmlands is imperative (Thompson et al., 2020). Considering
297 that no neonics were used on plants since 2018 in France and that the analytical method used
298 here provided good sensitivity results (Rodrigues et al., 2023), we expected to find no or very
299 low concentrations in bird blood samples as this matrix reflects short-term exposure (Espín et
300 al., 2016). However, in the present study, we not only detected five out of seven neonics,
301 three of which have been banned since 2018 in France (clothianidin, thiacloprid, and
302 thiamethoxam) and two others (dinotefuran and nitenpyram) that are supposedly being used
303 only for treating domestic pets (Table S2), but in some cases, at rather high concentrations.
304 The remaining two neonics (imidacloprid and acetamiprid) are not discussed here as these
305 compounds were not detectable with the multiresidue method used in this study. However,
306 knowing the historical background of the study area and results from studies in the same area
307 before neonics ban (see section 2.1), if titration had been possible, we would have expected

308 to find large amounts of imidacloprid in birds' blood and probably no or very low
309 concentrations of acetamiprid.

310

311 **4.1. Monitoring neonics in wild birds**

312

313 Neonics were increasingly studied during the last decade and several recent studies have
314 focused on quantifying the level of exposure in wild birds. However, these studies differ from
315 our work in several aspects. First, in these studies, several biological matrices—blood,
316 carcasses, eggs, faecal pellets and cloacal fluids, feathers, liver, and/or gizzard—were
317 considered (see Table S1 for detailed references). Correlations between neonics
318 quantifications from different matrices can be found but are not systematic (reviewed in Espín
319 et al., 2016). Indeed, pesticides are distributed in biological tissues with different temporal
320 patterns. For instance, pesticide molecules are integrated into feathers during the moulting
321 period (Humann-Guillemint et al., 2022). The distribution of such neonic molecules in
322 internal tissues such as the liver also depends on the chemical nature of the molecules and
323 can be degraded if samples are taken on carcasses (Espín et al., 2016, and references therein).
324 Then, a time lag might be observed between the exposure and the detection depending on
325 the matrices used (Lennon et al., 2020b). Therefore, the results obtained from different
326 matrices cannot be directly compared.

327 Our results can thus be reasonably compared to those of five previous studies that
328 tested for neonics in whole blood (or plasma) samples from wild birds (Taliensky-Chamudis et
329 al., 2017; Byholm et al., 2018; Hao et al., 2018; Lennon et al., 2020b; and Humann-Guillemint
330 et al., 2021; Table S4). For clothianidin, our concentration range was similar to those found by
331 Lennon et al. (2020b), although the maximum value was lower: they found concentrations

332 ranging from 0.5 (dunnocks) to 69300 $\text{pg}\cdot\mu\text{L}^{-1}$ (yellowhammers), whereas in our study, the
333 values were 0.009 (common nightingale) to 2451.78 $\text{pg}\cdot\mu\text{L}^{-1}$ (Montagu's harrier chick). The
334 maximum clothianidin concentration found by Humann-Guillemint et al. (2021) was 0.34
335 $\text{pg}\cdot\mu\text{L}^{-1}$ in alpine swifts, which is similar to the lower values in the concentration range we
336 obtained. Thiacloprid concentrations in all ciril buntings and common nightingales were below
337 the LOQ, but concentrations in the grey partridges and the Montagu's harrier chicks were far
338 higher than those found in honey buzzards by Byholm et al. (2018) and in white-crowned
339 sparrows by Hao et al. (2018). They ranged from 0.012 to 0.031 and 0.0025 to 0.0031 $\text{pg}\cdot\mu\text{L}^{-1}$
340 for honey buzzards and white-crowned sparrows, respectively, and from 0.073 to 89.58 $\text{pg}\cdot\mu\text{L}^{-1}$
341 ¹ in our study (Table S4). This result suggests that the grey partridges and the Montagu's
342 harrier chicks were exposed to high, or recent quantities of thiacloprid (see section 4.2 for
343 further details). For thiamethoxam, the concentrations we found were above those obtained
344 by Hao et al. (2018): they ranged from 0.06 (common nightingales) to 23.73 (grey partridges)
345 $\text{pg}\cdot\mu\text{L}^{-1}$, whereas Hao et al. reported 0.0051–0.0337 $\text{pg}\cdot\mu\text{L}^{-1}$. Taliany-Chamudis et al. (2017)
346 detected imidacloprid in one Eurasian eagle owl (*Bubo bubo*) chick among the 30 sampled,
347 but did not detect any other neonics, so comparisons with our results are not possible. The
348 difference in the results despite all studies using blood samples may be because some of the
349 studies considered whole blood samples (Taliany-Chamudis et al., 2017; Byholm et al., 2018;
350 our study), while others used only plasma (Hao et al., 2018; Lennon et al., 2020b; Humann-
351 Guillemint et al., 2021), inducing differences in the detectability of some molecules whose
352 levels vary according to their water/lipid solubility, and affinity to different proteins
353 (Rodrigues et al., 2023; Zhang et al., 2023). Moreover, all these studies fundamentally differ
354 in their sensitivity as they rely on their own developed chemical analysis methods [although
355 they are all derived from the same method, i.e., Quick, Easy, Cheap, Effective, Rugged, and

356 Safe (QuEChERS); Anastassiades et al., 2003]. If we consider clothianidin for instance, LOD and
357 LOQ in our study (0.01 and 0.04 $\text{pg}\cdot\mu\text{L}^{-1}$, respectively; Table S3) were 5 (LOQ) and 15 (LOD)
358 times lower, respectively, than those (0.15 and 0.21 $\text{pg}\cdot\mu\text{L}^{-1}$, respectively) of Lennon et al.
359 (2020b), while LOQ was similar to that (0.05 $\text{pg}\cdot\mu\text{L}^{-1}$) reported by Humann-Guillemot et al.
360 (2021). This may induce a difference in the number of neonic positive samples between
361 studies, hence differences in average concentrations as well. For instance, applying the LOD
362 of clothianidin from the study of Lennon et al. (2020b) would have led to non-detection of
363 positive samples in common nightingales while we found 8% of individuals with
364 concentrations above our LOD. It is not therefore easy to compare the results provided by
365 the authors, especially when not all statistical values, such as the percentage of neonic
366 detection among individuals or the mean concentration with its standard deviation, were
367 obtained on the same basis (Table S4). As the use of the same methodology in all studies does
368 not seem to be feasible, researchers should systematically report these values to allow direct
369 comparisons.

370

371 **4.2. Exposure of wild fauna to neonics**

372

373 Previous studies reported the limitations of feather and internal tissue samples for
374 determining timing of exposure to contaminants, so blood is considered most suitable for
375 determining recent exposure (Espín et al., 2016; Lennon et al., 2020b). Neonics are supposedly
376 “rapidly” excreted (in hummingbirds, English et al., 2021) and cleared from blood (24h in
377 quails), according to data regarding imidacloprid (Bean et al., 2019) and thiamethoxam (Pan
378 et al., 2022); however, extrapolation to the behaviour of other neonics may not be reliable.
379 Based on blood samples used for this study we cannot estimate long-term exposure,

380 therefore, possible exposure of the migratory species at their wintering areas may be
381 disregarded. However, thiamethoxam/clothianidin were found in grey partridges' eggs in
382 another study (Bro et al., 2016), so in the case of the Montagu's harrier chicks, we cannot
383 exclude the possibility of maternal transfer of neonics, i.e., from the mother to the eggs. In
384 other words, all tested positive individuals in the present study were probably recently
385 exposed to neonics. This is important because the birds were sampled three and four years
386 after EU banned neonics for PPP, and no derogation is known in our study area conversely to
387 other regions in France. Nonetheless, we found three out of the five PPP neonics in our
388 samples, with concentrations similar to those in birds sampled elsewhere before the ban
389 (Byholm et al., 2018). Taken together, our results strongly suggests that clothianidin,
390 thiacloprid, and thiamethoxam are still present in farmlands and raises questions about the
391 mode of exposure to birds. Thiamethoxam is metabolized into clothianidin in animals, plants,
392 and soil (Nauen et al., 2003, review in Simon-Delso et al., 2015; Pan et al., 2022) potentially
393 explaining why it is more commonly present than thiamethoxam. These neonics' DT₅₀ ranged
394 from few days for thiacloprid up to several years for clothianidin (Table S2) in the soil, meaning
395 that they can still be incorporated into the diet of several detritivores such as earthworms
396 (Pelosi et al., 2022). If so, at least some blackbirds, which are omnivorous and feed on different
397 invertebrate species during the breeding season, including earthworms which are known to
398 bioaccumulate pesticides (Pelosi et al., 2022; 2021), should have tested positive, but were not.
399 This surprising result could have been first seen as a technical failure; however, other pesticide
400 molecules than neonics were detected in blackbirds' blood, so that an analysis problem can
401 be discarded. The absence of neonics in blackbirds' blood might be the consequence of several
402 processes. One of them could be a higher detoxification capacity of blackbirds, mediated by
403 their ability at monopolizing high amounts of carotenoids, which are antioxidants, involved in

404 sexual selection and known for their role in reducing oxidative stress (Møller et al., 2000, see
405 Moreau et al., 2022a). Further analyses would be needed to explore this explanation.
406 Granivorous birds, including grey partridges (as they were caught during the winter), ciril
407 buntings, and common nightingales, which are omnivorous (mostly insectivorous during
408 reproduction), tested positive. This suggests that the mode of exposure may include
409 contaminated seeds from previous treated crops (Wintermantel et al., 2018) or organisms
410 feeding on contaminated seeds. The higher concentrations found in Montagu's harriers, which
411 are apex predators (mostly preying upon common voles and orthopterans), might be a result
412 of biomagnification, i.e., the accumulation of toxic neonics along the trophic chain (Badry et
413 al., 2020). In fact, clothianidin and thiacloprid, two compounds quantified in Montagu's
414 harriers' blood, were found in hairs of small mammals from the area, reinforcing that view
415 (Fritsch et al., 2022). Further investigations would be needed, and one way to do so would be
416 using chicks' food pellets. These differences among species may also be attributable to the
417 choices of individuals regarding their habitat and feeding resources for avoiding contaminated
418 sources, as has been highlighted previously (McKay et al., 1999; Ruuskanen et al., 2020; Addy-
419 Orduna et al., 2022). Another explanation relies on the detoxification processes in wild birds
420 that are still poorly investigated and deserved further investigations (Moreau et al., 2022a).
421 Indeed, some individuals could be more efficient than others at protecting themselves against
422 toxic substances (Arnold et al., 2015).

423 Another important issue observed in our results is the high prevalence of dinotefuran
424 and nitenpyram in grey partridges, although wildlife fauna is not supposed to be exposed to
425 these as they are used only for pets and not farm animals, and consequently not for outdoor
426 use in Europe, including France (Table S2). Even if grey partridges caught in this study area
427 may originate mainly from captive breeding stocks (released for hunting purpose), the six

428 registered veterinary medicines containing dinotefuran and nitenpyram neonics that are
429 authorized for use in France are not intended for treating any bird species or farm animal
430 (<http://www.ircp.anmv.anses.fr/index.aspx>). Imidacloprid is included in 36 different speciality
431 medications, making this neonic potentially more common even if not detectable here (see,
432 for example, Perkins et al., 2021). Neonics are either used for topical applications on the skin
433 (dinotefuran) or *per os* (nitenpyram) to treat flea infestations in cats and dogs typically;
434 further, nitenpyram is expected to be eliminated within 48 h (Jeschke & Nauen, 2005; Rust,
435 2017). In Europe, veterinary regulatory processes are governed by the European Medicines
436 Agency, which states that products for non-food animals are not supposed to be of major
437 environmental concerns since these animals are treated individually with low concentrations
438 of active neonics (CVMP/VICH, 2000). To our knowledge, these two neonics, investigated in
439 few studies (Table S1), have only been detected in one sample of hummingbirds' feather
440 rinsate (i.e., not within the organism; Graves et al., 2019), suggesting contact and not
441 ingestion. In the present study, the substantial concentrations of dinotefuran and nitenpyram
442 found in grey partridges but not in any other species may indicate possible exposure during
443 their stay in farms before release. This implies either an illegal use of these substances in
444 farms, as they are not included in veterinary medicines for farm animals, or an unintentional
445 contact of partridges through multiple potential pathways. One of these could be direct
446 contact and/or ingestion of farmer's dogs or cats' urine, or indirect through contaminated
447 drinking water. However, neonics being presumably rapidly excreted and blood reflecting
448 short-term exposure, individuals sampled here, if contaminated in farms before release,
449 should not test positive. Thus, we cannot exclude in our case that partridges once in the wild
450 could have directly ingested contaminated water with cat and/or dog urine and/or after pet
451 baths (Teerlink et al., 2017; Diepens et al., 2023). Besides, partridges that commonly use field

452 margins contrary to the other species, where pets walk and urinate, may have ingested these
453 neonics while preening as the external surface of feathers are often contaminated and might
454 even accumulate compounds (Pacyna-Kuchta, 2023). Indeed, there is growing evidence of
455 veterinary products' transfer to the environment from dogs' hair and urine, and of secondary
456 transfer to wildlife through nesting material or contaminated water (Diepens et al., 2023).
457 Even though we cannot ascertain by which route the partridges were exposed to these
458 neonics, our results indicate in line with Diepens et al. (2023), the urgent need for monitoring
459 all pesticide or medicine compounds, irrespective of their intended use. Although treating
460 domestic animals against ectoparasites might be of sanitary importance for public health
461 (human and animal), these treatments may have the same adverse effects on wildlife as PPP
462 (reviewed in Moreau et al., 2022a).

463 It should be noted that the interpretations given to the results provided in the present
464 study are only assumptions and that the origin of birds' contamination remain unknown.
465 Ongoing studies on neonics levels in soils and invertebrates of the study area since 2018
466 should help to provide further clarifications on birds' contamination pathways. Moreover, the
467 method of multiresidue analysis used here could be improved to allow the detection of all
468 neonics, including imidacloprid and acetamiprid, especially as imidacloprid was extensively
469 applied in the crops of the study area and is still highly used in veterinary medicine.

470

471 **4.3. Conclusions**

472

473 Our results highlight several problems with the use of a class of chemicals that are among the
474 most used worldwide. First, banning neonics for outdoor use does not prevent the exposure
475 of wildlife fauna to them, at least a few years after ban. Although illegal use cannot be

476 disregarded, it cannot explain on its own their ubiquity in wild birds from a wide ecological
477 range, caught at the scale of the study site. This is an important element to consider in
478 countries where neonics are still massively applied, considering their impact on both animals
479 and humans (reviewed in Moreau et al., 2022a). Second, the detection of neonics used for
480 domestic animals in wildlife fauna raises questions regarding the manner in which risk
481 assessment for such applications is performed (Perkins et al., 2021; Diepens et al., 2023). For
482 instance, to our knowledge, few studies have investigated the effects of dinotefuran and
483 nitenpyram on wildlife fauna (Wang et al., 2018). Although clothianidin, dinotefuran,
484 nitenpyram, and thiamethoxam are considered to be of low toxicity to birds, as indicated by
485 their acute oral LD₅₀ (Table S2), thiacloprid, which was also found to be quite ubiquitous, is
486 highly toxic to birds (Table S2). However, LD₅₀ is indicative of acute lethal toxicity under
487 laboratory conditions for model species and is not necessarily an appropriate estimate of
488 sublethal effects, as shown in the numerous studies that investigated the adverse effects of
489 neonics on birds (Moreau et al., 2022a). Considering that very low residual levels of pesticides
490 may have considerable sublethal effects on birds' reproduction (see Moreau et al., 2021, for
491 example), the consequences of these exposures on bird populations as well as on human
492 health (*One Health* concept) should be carefully considered.

493

494 **Ethical statement**

495 This study was conducted conforming the French guidelines for the ethical use of animals in
496 research (APAFIS#18557-2019010822312199v2, APAFIS#9465e201703101551625). We are
497 grateful to the Nouvelle-Aquitaine Regional Agency of the Environment, Development and
498 Housing for the official authorizations for the capture of passerines birds
499 (DREAL/2019D/2323). Because Montagu's harrier is a protected raptor species, its handling

500 was allowed and licensed by the Centre de Recherches sur la Biologie des Populations
501 d'Oiseaux – Museum National d'Histoire Naturelle (license #1308 for Montagu's harriers).

502

503 **Declaration of competing interests**

504 The authors declare that they have no conflicts of interest.

505

506 **Credit author statement**

507 EF, AG, AR, MM, VB, JM and KM conceived the ideas and designed the methodology; AG, JM
508 and KM collected the data; EF, AR, MM and KM analysed the data; EF, AG, VB, JM and KM led
509 the writing of the paper. All the authors contributed critically to the drafts and gave final
510 approval for publication.

511

512 **Funding**

513 This work was supported by the French National Research Agency (ANR JCJC PestiStress, grant
514 #19-CE34-0003-01), the French National Centre of Scientific Research (CNRS), the French
515 National Research Institute for Agriculture, Food and the Environment (INRAE), and the
516 University of La Rochelle. This study was partly funded by the BioBird project (regional
517 government of Nouvelle-Aquitaine), by the ACI (University of La Rochelle), BIRDPEST project
518 (RECOTOX 2019), AgriBioBird project (OSU Theta ISITE-BFC), the Contrat de Plan État-Région
519 (CPER) Econat and the French National program EC2CO (Ecosphère Continentale et Côtière).

520

521 **Acknowledgements**

522 We thank all the people who contributed to fieldwork and data collection: Fanny Rey and
523 Mélissa Desbonnes for passerine birds, Landry Boussac and Samuel Perroteau for grey

524 partridges, and Mazzarine Laboureau and Mona Moreau for Montagu's harriers. Further, we
525 are grateful to all farmers who allowed us access to their fields. We would also like to thank
526 the trainees Emilie Jemmi and Zülal Can who helped with blood extraction at the ICPEES
527 laboratory.
528

529 **References**

530

531 Abu Zeid, E. H., Alam, R. T., Ali, S. A., Hendawi, M. Y., 2019. Dose-related impacts of
532 imidacloprid oral intoxication on brain and liver of rock pigeon (*Columba livia domestica*),
533 residues analysis in different organs. *Ecotox. Environ. Safe.* 167, 60-68.
534 <https://doi.org/10.1016/j.ecoenv.2018.09.121>

535

536 Addy-Orduna, L., Cazenave, J., Mateo, R., 2022. Avoidance of neonicotinoid-treated seeds and
537 cotyledons by captive eared doves (*Zenaida auriculata*, Columbidae). *Environ. Pollut.* 304,
538 119237. <https://doi.org/10.1016/j.envpol.2022.119237>

539

540 Anastassiades, M., Lehotay, S. J., Štajnbaher, D., Schenck, F. J., 2003. Fast and easy
541 multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-
542 phase extraction” for the determination of pesticide residues in produce. *J. AOAC Int.* 86, 412-
543 431. <https://doi.org/10.1093/jaoac/86.2.412>

544

545 Arnold, K. E., Herborn, K. A., Adam, A., Alexander, L., 2015. Individual variation in the oxidative
546 costs of personality traits. *Funct. Ecol.* 29, 522-530. <https://doi.org/10.1111/1365-2435.12375>

547

548 Arroyo, B. E., Bretagnolle, V., Leroux, A., 2007. Interactive effects of food and age on breeding
549 in the Montagu's Harrier *Circus pygargus*. *Ibis.* 149, 806-813. [https://doi.org/10.1111/j.1474-
550 919X.2007.00716.x](https://doi.org/10.1111/j.1474-919X.2007.00716.x)

551

552 Arroyo, B. E., García, J. T., Bretagnolle, V., 2004. Montagu's harrier. – BWP update (The Journal
553 of the Birds of the Western Palearctic), Vol. 6, Nos 1/2. Oxford University Press, Oxford, pp.
554 41–55.

555

556 Arroyo, B., Leroux, A., Bretagnolle, V., 1998. Patterns of egg and clutch size variation in the
557 Montagu's Harrier. *J. Raptor Res.* 32, 136-142.

558

559 Badry, A., Krone, O., Jaspers, V. L., Mateo, R., García-Fernández, A., Leivits, M., Shore, R. F.,
560 2020. Towards harmonisation of chemical monitoring using avian apex predators:
561 Identification of key species for pan-European biomonitoring. *Sci. Total Environ.* 731, 139198.

562 <https://doi.org/10.1016/j.scitotenv.2020.139198>

563

564 Bean, T. G., Gross, M. S., Karouna-Renier, N. K., Henry, P. F., Schultz, S. L., Hladik, M. L., Kuivila,
565 K. M., Rattner, B. A., 2019. Toxicokinetics of imidacloprid-coated wheat seeds in Japanese
566 quail (*Coturnix japonica*) and an evaluation of hazard. *Environ. Sci. Technol.* 53, 3888-3897.

567 <https://doi.org/10.1021/acs.est.8b07062>

568

569 Bishop, C. A., Woundneh, M. B., Maisonneuve, F., Common, J., Elliott, J. E., Moran, A. J., 2020.
570 Determination of neonicotinoids and butenolide residues in avian and insect pollinators and
571 their ambient environment in Western Canada (2017, 2018). *Sci. Total Environ.* 737, 139386.

572 <https://doi.org/10.1016/j.scitotenv.2020.139386>

573

574 Bishop, C. A., Moran, A. J., Toshack, M. C., Elle, E., Maisonneuve, F., Elliott, J. E., 2018.

575 Hummingbirds and bumble bees exposed to neonicotinoid and organophosphate insecticides

576 in the Fraser Valley, British Columbia, Canada. *Environ. Toxicol. Chem.* 37, 2143-2152.
577 <https://doi.org/10.1002/etc.4174>
578

579 Bonneris, E., Gao, Z., Prosser, A., Barfknecht, R., 2019. Selecting appropriate focal species for
580 assessing the risk to birds from newly drilled pesticide-treated winter cereal fields in France.
581 *Integr. Environ. Asses.* 15, 422-436. <https://doi.org/10.1002/ieam.4112>
582

583 Bretagnolle, V., Berthet, E., Gross, N., Gauffre, B., Plumejeaud, C., Houte, S., Badenhausser, I.,
584 Monceau, K., Allier, F., Monestiez, P., Gaba, S., 2018. Towards sustainable and multifunctional
585 agriculture in farmland landscapes: lessons from the integrative approach of a French LTSER
586 platform. *Sci. Total Environ.* 627, 822-834. <https://doi.org/10.1016/j.scitotenv.2018.01.142>
587

588 Bro, E., Devillers, J., Millot, F., Decors, A., 2016. Residues of plant protection products in grey
589 partridge eggs in French cereal ecosystems. *Environ. Sci. Pollut. R.* 23, 9559-9573.
590 <https://doi.org/10.1007/s11356-016-6093-7>
591

592 Burns, F., Eaton, M. A., Burfield, I. J., Klvaňová, A., Šilarová, E., Staneva, A., Gregory, R. D.,
593 2021. Abundance decline in the avifauna of the European Union reveals cross-continental
594 similarities in biodiversity change. *Ecol. Evol.* 11, 16647-16660.
595 <https://doi.org/10.1002/ece3.8282>
596

597 Butet, A., Leroux, A. B., 2001. Effects of agriculture development on vole dynamics and
598 conservation of Montagu's harrier in western French wetlands. *Biol. Conserv.* 100, 289-295.
599 [https://doi.org/10.1016/S0006-3207\(01\)00033-7](https://doi.org/10.1016/S0006-3207(01)00033-7)

600

601 Byholm, P., Mäkeläinen, S., Santangeli, A., Goulson, D., 2018. First evidence of neonicotinoid
602 residues in a long-distance migratory raptor, the European honey buzzard (*Pernis apivorus*).
603 Sci. Total Environ. 639, 929-933. <https://doi.org/10.1016/j.scitotenv.2018.05.185>

604

605 Campbell, L.H., Avery, M.I., Donald, P., Evans, A.D., Green, R.E., Wilson, J.D., 1997. A review of
606 the indirect effects of pesticides on birds. JNCC Report No. 227. Joint Nature Conservation
607 Committee, Peterborough.

608

609 Casida, J. E., 2018. Neonicotinoids and other insect nicotinic receptor competitive modulators:
610 progress and prospects. Annu. Rev. Entomol. 63, 125-144. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-ento-020117-043042)
611 [ento-020117-043042](https://doi.org/10.1146/annurev-ento-020117-043042)

612

613 Casida, J. E., Durkin, K. A., 2013. Neuroactive insecticides: targets, selectivity, resistance, and
614 secondary effects. Annu. Rev. Entomol. 58, 99-117. [https://doi.org/10.1146/annurev-ento-](https://doi.org/10.1146/annurev-ento-120811-153645)
615 [120811-153645](https://doi.org/10.1146/annurev-ento-120811-153645)

616

617 Chiron, F., Chargé, R., Julliard, R., Jiguet, F., Muratet, A., 2014. Pesticide doses, landscape
618 structure and their relative effects on farmland birds. Agr. Ecosyst. Environ. 185, 153-160.
619 <https://doi.org/10.1016/j.agee.2013.12.013>

620

621 Cimino, A. M., Boyles, A. L., Thayer, K. A., Perry, M. J., 2017. Effects of neonicotinoid pesticide
622 exposure on human health: a systematic review. Environ. Health Persp., 125, 155-162.
623 <https://doi.org/10.1289/EHP515>

624

625 Comolet-Tirman, J., Siblet, J. P., Witté, I., Cadiou, B., Czajkowski, M. A., Deceuninck, B., Jiguet,
626 F., Landry, P., Quaintenne, G., Roché, J.E., Sarasa, M., Touroult, J., 2015. Statuts et tendances
627 des populations d'oiseaux nicheurs de France. Bilan simplifié du premier rapportage national
628 au titre de la Directive Oiseaux. *Alauda*, 83, 35-76.

629

630 CVMP/VICH (2000). Guideline on Environmental impact assessment (EIAs) for veterinary
631 medicinal products (VMPs) – Phase I, VICH GL 6 (CVMP/VICH/592/1998) [Online]. Available at:
632 [https://www.ema.europa.eu/en/vich-gl6-environmental-impact-assessment-eias-veterinary-](https://www.ema.europa.eu/en/vich-gl6-environmental-impact-assessment-eias-veterinary-medicinal-products-phase-i)
633 [medicinal-products-phase-i](https://www.ema.europa.eu/en/vich-gl6-environmental-impact-assessment-eias-veterinary-medicinal-products-phase-i) , accessed date: 12 October 2022.

634

635 DEFRA (2021). Wild bird populations in the UK, 1970 to 2019 – Updated for Wintering and
636 Waterbirds. Department for Environment, Food Rural Affairs [Online]. Available at:
637 <https://www.gov.uk/government/statistics/wild-bird-populations-in-the-uk> , accessed date:
638 12 October 2022.

639

640 Diepens, N. J., Belgers, D., Buijse, L., Roessink, I., 2023. Pet dogs transfer veterinary medicines
641 to the environment. *Sci. Total Environ.* 858, 159550.
642 <https://doi.org/10.1016/j.scitotenv.2022.159550>

643

644 EFSA (European Food Safety Authority) (2021). Evaluation of the emergency authorisations
645 granted by Member State France for plant protection products containing imidacloprid or
646 thiamethoxam. EFSA supporting publication 2021 : EN-6968. 13pp.
647 <https://doi.org/10.2903/sp.efsa.2021.EN-6968>

648

649 Eng, M. L., Stutchbury, B. J., Morrissey, C. A., 2017. Imidacloprid and chlorpyrifos insecticides
650 impair migratory ability in a seed-eating songbird. *Sci. Rep-UK.* 7, 1-9.
651 <https://doi.org/10.1038/s41598-017-15446-x>

652

653 English, S. G., Sandoval-Herrera, N. I., Bishop, C. A., Cartwright, M., Maisonneuve, F., Elliott, J.
654 E., Welch, K. C., 2021. Neonicotinoid pesticides exert metabolic effects on avian pollinators.
655 *Sci. Rep-UK.* 11, 1-11. <https://doi.org/10.1038/s41598-021-82470-3>

656

657 Ertl, H. M., Mora, M. A., Brightsmith, D. J., Navarro-Alberto, J. A., 2018a. Potential impact of
658 neonicotinoid use on Northern bobwhite (*Colinus virginianus*) in Texas: A historical
659 analysis. *PLoS One.* 13, e0191100. <https://doi.org/10.1371/journal.pone.0191100>

660

661 Ertl, H. M., Mora, M. A., Boellstorff, D. E., Brightsmith, D., Carson, K., 2018b. Potential effects
662 of neonicotinoid insecticides on northern bobwhites. *Wildlife Soc. B.* 42, 649-655.
663 <https://doi.org/10.1002/wsb.921>

664

665 Espín, S., García-Fernández, A.J., Herzke, D., Shore, R. F., van Hattum, B., Martínez-López, E.,
666 Coeurdassier, M., Eulaers, I., Fritsch, C., Gómez-Ramírez, P., Jaspers, V. L. B., Krone, O., Duke,
667 G., Helander, B., Mateo, R., Movalli, P., Sonne, C., van den Brink, N. W., 2016. Tracking pan-
668 continental trends in environmental contamination using sentinel raptors—what types of
669 samples should we use?. *Ecotoxicology.* 25, 777–801. [https://doi.org/10.1007/s10646-016-](https://doi.org/10.1007/s10646-016-1636-8)
670 [1636-8](https://doi.org/10.1007/s10646-016-1636-8)

671

672 Fritsch, C., Appenzeller, B., Burkart, L., Coeurdassier, M., Scheifler, R., Raoul, F., Driget, V.,
673 Powolny, T., Gagnaison, C., Rieffel, D., Afonso, E., Goydadin, A.-C., Hardy, E. M., Palazzi, P.,
674 Schaeffer, C., Gaba, S., Bretagnolle, V., Bertrand, C., Pelosi, C., 2022. Pervasive exposure of
675 wild small mammals to legacy and currently used pesticide mixtures in arable landscapes. *Sci.*
676 *Rep-UK*. 12, 1-22. <https://doi.org/10.1038/s41598-022-19959-y>

677

678 Geiger, F., Bengtsson, J., Berendse, F., Weisser, W. W., Emmerson, M., Morales, M. B.,
679 Ceryngier, P., Liira, J., Tschardtke, T., Winqvist, C., Eggers, S., Bommarco, R., Pärt, T.,
680 Bretagnolle, V., Plantegenest, M., Clement, L. W., Dennis, C., Palmer, C., Oñate, J. J., Guerrero,
681 I., Hawro, V., Aavik, T., Thies, C., Flohre, A., Hänke, S., Fischer, C., Goedhart, P. W., Inchausti,
682 P., 2010. Persistent negative effects of pesticides on biodiversity and biological control
683 potential on European farmland. *Basic Appl. Ecol.* 11, 97-105.
684 <https://doi.org/10.1016/j.baae.2009.12.001>

685

686 Gibbons, D., Morrissey, C., Mineau, P., 2015. A review of the direct and indirect effects of
687 neonicotinoids and fipronil on vertebrate wildlife. *Environ. Sci. Pollut. R.* 22, 103-118.
688 <https://doi.org/10.1007/s11356-014-3180-5>

689

690 Graves, E. E., Jelks, K. A., Foley, J. E., Filigenzi, M. S., Poppenga, R. H., Ernest, H. B., Melnicoe,
691 R., Tell, L. A., 2019. Analysis of insecticide exposure in California hummingbirds using liquid
692 chromatography-mass spectrometry. *Environ. Sci. Pollut. R.* 26, 15458-15466.
693 <https://doi.org/10.1007/s11356-019-04903-x>

694

695 Grue, C. E., Gibert, P. L., Seeley, M. E., 1997. Neurophysiological and behavioral changes in
696 non-target wildlife exposed to organophosphate and carbamate pesticides:
697 thermoregulation, food consumption, and reproduction. *Am. Zool.* 37, 369-388.
698 <https://www.jstor.org/stable/3884019>
699

700 Han, W., Tian, Y., Shen, X., 2018. Human exposure to neonicotinoid insecticides and the
701 evaluation of their potential toxicity: An overview. *Chemosphere.* 192, 59-65.
702 <https://doi.org/10.1016/j.chemosphere.2017.10.149>
703

704 Goutte, A., Molbert, N., 2022. Benefits of Parasitism in Polluted Environments: A Review and
705 Perspectives. *Front. Ecol. Evol.* 10, 847869. <https://doi.org/10.3389/fevo.2022.847869>
706

707 Hao, C., Eng, M. L., Sun, F., Morrissey, C. A., 2018. Part-per-trillion LC-MS/MS determination
708 of neonicotinoids in small volumes of songbird plasma. *Sci. Total Environ.* 644, 1080-1087.
709 <https://doi.org/10.1016/j.scitotenv.2018.06.317>
710

711 Hallmann, C. A., Foppen, R. P., Van Turnhout, C. A., De Kroon, H., Jongejans, E., 2014. Declines
712 in insectivorous birds are associated with high neonicotinoid concentrations. *Nature.* 511,
713 341-343. <https://doi.org/10.1038/nature13531>
714

715 Henry, M., Cerrutti, N., Aupinel, P., Decourtye, A., Gayrard, M., Odoux, J. F., Pissard, A., Rüger,
716 C., Bretagnolle, V., 2015. Reconciling laboratory and field assessments of neonicotinoid
717 toxicity to honeybees. *P. R. Soc. B-Biol. Sci.* 282, 20152110.
718 <https://doi.org/10.1098/rspb.2015.2110>

719

720 Hoshi, N., Hirano, T., Omotehara, T., Tokumoto, J., Umemura, Y., Mantani, Y., Tanida, T.,
721 Warita, K., Tabuchi, Y., Yokoyama, T., Kitagawa, H., 2014. Insight into the mechanism of
722 reproductive dysfunction caused by neonicotinoid pesticides. *Biol. Pharm. Bull.* 37, 1439-
723 1443. <https://doi.org/10.1248/bpb.b14-00359>

724

725 Humann-Guilleminot, S., Andreo, L., Blatti, E., Glauser, G., Helfenstein, F., Desprat, J., 2022.
726 Experimental evidence for clothianidin deposition in feathers of house sparrows after
727 ingestion of sublethal doses treated seeds. *Chemosphere.* 315, 137724.
728 <https://doi.org/10.1016/j.chemosphere.2022.137724>

729

730 Humann-Guilleminot, S., Laurent, S., Bize, P., Roulin, A., Glauser, G., Helfenstein, F., 2021.
731 Contamination by neonicotinoid insecticides in barn owls (*Tyto alba*) and Alpine swifts
732 (*Tachymarptis melba*). *Sci. Total Environ.* 785, 147403.
733 <https://doi.org/10.1016/j.scitotenv.2021.147403>

734

735 Humann-Guilleminot, S., de Montaignu, C. T., Sire, J., Grünig, S., Gning, O., Glauser, G., Vallat,
736 A. Helfenstein, F., 2019. A sublethal dose of the neonicotinoid insecticide acetamiprid reduces
737 sperm density in a songbird. *Environ. Res.* 177, 108589.
738 <https://doi.org/10.1016/j.envres.2019.108589>

739

740 Ihara, M., D Buckingham, S., Matsuda, K., B Sattelle, D., 2017. Modes of action, resistance and
741 toxicity of insecticides targeting nicotinic acetylcholine receptors. *Curr. Med. Chem.* 24, 2925-
742 2934. <https://doi.org/10.2174/0929867324666170206142019>

743

744 Jeschke, P., Nauen, R., 2005. "Neonicotinoid Insecticides, 53-105". In: Comprehensive
745 Molecular Insect Science (Eds. L. I. Gilbert, L. Iatrou S. S. Gill). Elsevier, Oxford, UK, 459 pp.

746

747 Klingelhöfer, D., Braun, M., Brüggmann, D., Groneberg, D. A., 2022. Neonicotinoids: A critical
748 assessment of the global research landscape of the most extensively used insecticide. Environ.
749 Res. 213, 113727. <https://doi.org/10.1016/j.envres.2022.113727>

750

751 Köhler, H. R., Triebkorn, R., 2013. Wildlife ecotoxicology of pesticides: can we track effects to
752 the population level and beyond? Science. 341, 759-765.
753 <https://doi.org/10.1126/science.1237591>

754

755 Lennon, R. J., Peach, W. J., Dunn, J. C., Shore, R. F., Pereira, M. G., Sleep, D., Dodd, S., Wheatley,
756 C. J., Arnold, K. E., Brown, C. D., 2020a. From seeds to plasma: Confirmed exposure of multiple
757 farmland bird species to clothianidin during sowing of winter cereals. Sci. Total Environ. 723,
758 138056. <https://doi.org/10.1016/j.scitotenv.2020.138056>

759

760 Lennon, R. J., Shore, R. F., Pereira, M. G., Peach, W. J., Dunn, J. C., Arnold, K. E., Brown, C. D.,
761 2020b. High prevalence of the neonicotinoid clothianidin in liver and plasma samples collected
762 from gamebirds during autumn sowing. Sci. Total Environ. 742, 140493.
763 <https://doi.org/10.1016/j.scitotenv.2020.140493>

764

765 Li, Y., Miao, R., Khanna, M., 2020. Neonicotinoids and decline in bird biodiversity in the United
766 States. Nat. Sustain. 3, 1027-1035. <https://doi.org/10.1038/s41893-020-0582-x>

767

768 Lopez-Antia, A., Ortiz-Santaliestra, M. E., Camarero, P. R., Mougeot, F., Mateo, R., 2015a.

769 Assessing the risk of fipronil-treated seed ingestion and associated adverse effects in the red-

770 legged partridge. Environ. Sci. Technol. 49, 13649-13657.

771 <https://doi.org/10.1021/acs.est.5b03822>

772

773 Lopez-Antia, A., Ortiz-Santaliestra, M. E., Mougeot, F., Mateo, R., 2015b. Imidacloprid-treated

774 seed ingestion has lethal effect on adult partridges and reduces both breeding investment and

775 offspring immunity. Environ. Res. 136, 97-107. <https://doi.org/10.1016/j.envres.2014.10.023>

776

777 Lopez-Antia, A., Ortiz-Santaliestra, M. E., Mougeot, F., Mateo, R., 2013. Experimental

778 exposure of red-legged partridges (*Alectoris rufa*) to seeds coated with imidacloprid, thiram

779 and difenoconazole. Ecotoxicology. 22, 125-138. <https://doi.org/10.1007/s10646-012-1009-x>

780

781 Lv, Y., Bing, Q., Lv, Z., Xue, J., Li, S., Han, B., Yang, Q., Wang, X., Zhang, Z., 2020. Imidacloprid-

782 induced liver fibrosis in quails via activation of the TGF- β 1/Smad pathway. Sci. Total Environ.

783 705, 135915. <https://doi.org/10.1016/j.scitotenv.2019.135915>

784

785 MacDonald, A. M., Jardine, C. M., Thomas, P. J., Nemeth, N. M., 2018. Neonicotinoid detection

786 in wild turkeys (*Meleagris gallopavo silvestris*) in Ontario, Canada. Environ. Sci. Pollut. R. 25,

787 16254-16260. <https://doi.org/10.1007/s11356-018-2093-0>

788

789 McKay, H. V., Prosser, P. J., Hart, A. D. M., Langton, S. D., Jones, A., McCoy, C., Chandler-Morris,
790 S. A., Pascual, J. A., 1999. Do wood-pigeons avoid pesticide-treated cereal seed?. J. Appl. Ecol.
791 36, 283-296. <https://doi.org/10.1046/j.1365-2664.1999.00398.x>

792

793 Millon, A., Arroyo, B. E., Bretagnolle, V., 2008. Variable but predictable prey availability affects
794 predator breeding success: natural versus experimental evidence. J. Zool. 275, 349-358.
795 <https://doi.org/10.1111/j.1469-7998.2008.00447.x>

796

797 Millot, F., Decors, A., Mastain, O., Quintaine, T., Berny, P., Vey, D., Lasseur, R., Bro, E., 2017.
798 Field evidence of bird poisonings by imidacloprid-treated seeds: a review of incidents reported
799 by the French SAGIR network from 1995 to 2014. Environ. Sci. Pollut. Res. 24, 5469–5485.
800 <https://doi.org/10.1007/s11356-016-8272-y>

801

802 Mineau, P., Whiteside, M., 2013. Pesticide acute toxicity is a better correlate of US grassland
803 bird declines than agricultural intensification. PloS One. 8, e57457.
804 <https://doi.org/10.1371/journal.pone.0057457>

805

806 Mitra, A., Chatterjee, C., Mandal, F. B., 2011. Synthetic chemical pesticides and their effects
807 on birds. Res. J. Environ. Toxicol. 5, 81-96. <https://doi.org/10.3923/rjet.2011.81.96>

808

809 Mohanty, B., Pandey, S. P., Tsutsui, K., 2017. Thyroid disrupting pesticides impair the
810 hypothalamic-pituitary-testicular axis of a wildlife bird, *Amandava amandava*. Reprod.
811 Toxicol. 71, 32-41. <https://doi.org/10.1016/j.reprotox.2017.04.006>

812

813 Møller, A. P., Biard, C., Blount, J. D., Houston, D. C., Ninni, P., Saino, N., Surai, P. F., 2000.
814 Carotenoid-dependent signals: indicators of foraging efficiency, immunocompetence or
815 detoxification ability?. *Poult. Avian Biol. Rev.* 11, 137-160.

816

817 Moreau, J., Rabdeau, J., Badenhauer, I., Giraudeau, M., Sepp, T., Crépin, M., Gaffard, A.,
818 Bretagnolle, V., Monceau, K., 2022a. Pesticide impacts on avian species with special reference
819 to farmland birds: a review. *Environ. Monit. Assess.* 194, 1-48.
820 <https://doi.org/10.1007/s10661-022-10394-0>

821

822 Moreau, J., Monceau, K., Gonnet, G., Pfister, M., Bretagnolle, V., 2022b. Organic farming
823 positively affects the vitality of passerine birds in agricultural landscapes. *Agr. Ecosyst.*
824 *Environ.* 336, 108034. <https://doi.org/10.1016/j.agee.2022.108034>

825

826 Moreau, J., Monceau, K., Crépin, M., Tochon, F. D., Mondet, C., Fraikin, M., Teixeira, M.,
827 Bretagnolle, V., 2021. Feeding partridges with organic or conventional grain triggers cascading
828 effects in life-history traits. *Environ. Pollut.* 278, 116851.
829 <https://doi.org/10.1016/j.envpol.2021.116851>

830

831 Nauen, R., Ebbinghaus-Kintscher, U., Salgado, V. L., Kausmann, M., 2003. Thiamethoxam is a
832 neonicotinoid precursor converted to clothianidin in insects and plants. *Pestic. Biochem. Phys.*
833 76, 55-69. [https://doi.org/10.1016/S0048-3575\(03\)00065-8](https://doi.org/10.1016/S0048-3575(03)00065-8)

834

835 Pacyna-Kuchta, A. D., 2023. What should we know when choosing feather, blood, egg or preen
836 oil as biological samples for contaminants detection? A non-lethal approach to bird sampling

837 for PCBs, OCPs, PBDEs and PFASs. Crit. Rev. Environ. Sci. Technol. 53, 625-649.

838 <https://doi.org/10.1080/10643389.2022.2077077>

839

840 Pan, Y., Chang, J., Xu, P., Xie, Y., Yang, L., Hao, W., Li, J., Wan, B., 2022. Twenty-four hours of

841 Thiamethoxam: In vivo and molecular dynamics simulation study on the toxicokinetic and

842 underlying mechanisms in quails (*Coturnix japonica*). J. Hazard. Mater. 427, 128159.

843 <https://doi.org/10.1016/j.jhazmat.2021.128159>

844

845 Pelosi, C., Bertrand, C., Bretagnolle, V., Coeurdassier, M., Delhomme, O., Deschamps, M.,

846 Gaba, S., Millet, M., Nélieu, S., Fritsch, C., 2022. Glyphosate, AMPA and glufosinate in soils and

847 earthworms in a French arable landscape. Chemosphere. 301, 134672.

848 <https://doi.org/10.1016/j.chemosphere.2022.134672>

849

850 Pelosi, C., Bertrand, C., Daniele, G., Coeurdassier, M., Benoit, P., Nélieu, S., Lafay, F.,

851 Bretagnolle, V., Gaba, S., Vulliet, E., Fritsch, C., 2021. Residues of currently used pesticides in

852 soils and earthworms: A silent threat?. Agr. Ecosyst. Environ. 305, 107167.

853 <https://doi.org/10.1016/j.agee.2020.107167>

854

855 Perkins, R., Whitehead, M., Civil, W., Goulson, D., 2021. Potential role of veterinary flea

856 products in widespread pesticide contamination of English rivers. Sci. Total Environ. 755,

857 143560. <https://doi.org/10.1016/j.scitotenv.2020.143560>

858

859 Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., Stanton,
860 J. C., Panjabi, A., Helft, L., Parr, M., Marra, P. P., 2019. Decline of the North American
861 avifauna. *Science*. 366, 120-124. <https://doi.org/10.1126/science.aaw1313>
862

863 Rawi, S. M., Al-Logmani, A. S., Hamza, R. Z., 2019. Neurological alterations induced by
864 formulated imidacloprid toxicity in Japanese quails. *Metab. Brain Dis.* 34, 443-450.
865 <https://doi.org/10.1007/s11011-018-0377-1>
866

867 Rodrigues, A., Gaffard, A., Moreau, J., Monceau, K., Delhomme, O., Millet, M., 2023. Analytical
868 Development for the Assessment of Pesticide Contaminations in Blood and Plasma of Wild
869 Birds: The Case of Grey Partridges (*Perdix Perdix*). *J. Chromatogr. A.* 1687, 463681.
870 <https://doi.org/10.1016/j.chroma.2022.463681>
871

872 Rust, M. K., 2017. The biology and ecology of cat fleas and advancements in their pest
873 management: a review. *Insects*. 8, 118. <https://doi.org/10.3390/insects8040118>
874

875 Ruuskanen, S., Rainio, M. J., Kuosmanen, V., Laihonen, M., Saikkonen, K., Saloniemi, I.,
876 Helander, M., 2019. Female preference and adverse developmental effects of glyphosate-
877 based herbicides on ecologically relevant traits in Japanese quails. *Environ. Sci. Technol.* 54,
878 1128-1135. <https://doi.org/10.1021/acs.est.9b07331>
879

880 Salamolard, M., Butet, A., Leroux, A., Bretagnolle, V., 2000. Responses of an avian predator to
881 variations in prey density at a temperate latitude. *Ecology*. 81, 2428-2441.
882 [https://doi.org/10.1890/0012-9658\(2000\)081\[2428:ROAAPT\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[2428:ROAAPT]2.0.CO;2)

883

884 Simon-Delso, N., Amaral-Rogers, V., Belzunces, L. P., Bonmatin, J. M., Chagnon, M., Downs, C.,
885 Furlan, L., Gibbons, D. W., Giorio, C., Girolami, V., Goulson, D., Kreutzweiser, D. P., Krupke, C.
886 H., Liess, M., Long, E., McField, M., Mineau, P., Mitchell, E. A. D., Morrissey, C. A., Noome, D.
887 A., Pisa, L., Settele, J., Stark, J. D., Tapparo, A., Van Dyck, H., Van Praagh, J., Van der Sluijs, J.
888 P., Whitehorn, P. R., Wiemers, M., 2015. Systemic insecticides (neonicotinoids and fipronil):
889 trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. R.* 22, 5-34.
890 <https://doi.org/10.1007/s11356-014-3470-y>

891

892 Stanton, R. L., Morrissey, C. A., Clark, R. G., 2018. Analysis of trends and agricultural drivers of
893 farmland bird declines in North America: A review. *Agr. Ecosyst. Environ.* 254, 244-254.
894 <https://doi.org/10.1016/j.agee.2017.11.028>

895

896 Story, P., Cox, M., 2001. Review of the effects of organophosphorus and carbamate
897 insecticides on vertebrates. Are there implications for locust management in
898 Australia?. *Wildlife Res.* 28, 179-193. <https://doi.org/10.1071/WR99060>

899

900 Taliansky-Chamudis, A., Gómez-Ramírez, P., León-Ortega, M., García-Fernández, A. J., 2017.
901 Validation of a QuEChERS method for analysis of neonicotinoids in small volumes of blood and
902 assessment of exposure in Eurasian eagle owl (*Bubo bubo*) nestlings. *Sci. Total Environ.* 595,
903 93-100. <https://doi.org/10.1016/j.scitotenv.2017.03.246>

904

905 Tassin de Montaigu, C., Goulson, D., 2020. Identifying agricultural pesticides that may pose a
906 risk for birds. *PeerJ.* 8, e9526. <https://doi.org/10.7717/peerj.9526>

907

908 Teerlink, J., Hernandez, J., Budd, R., 2017. Fipronil washoff to municipal wastewater from dogs
909 treated with spot-on products. *Sci. Total Environ.* 599, 960-966.
910 <https://doi.org/10.1016/j.scitotenv.2017.04.219>

911

912 Thompson, D. A., Lehmler, H. J., Kolpin, D. W., Hladik, M. L., Vargo, J. D., Schilling, K. E.,
913 LeFevre, G. H., Peeples, T. L., Poch, M. C., LaDuca, L. E., Cwiertny, D. M., Field, R. W., 2020. A
914 critical review on the potential impacts of neonicotinoid insecticide use: current knowledge
915 of environmental fate, toxicity, and implications for human health. *Environ. Sci.-Proc. Imp.* 22,
916 1315-1346. <https://doi.org/10.1039/C9EM00586B>

917

918 Tokumoto, J., Danjo, M., Kobayashi, Y., Kinoshita, K., Omotehara, T., Tatsumi, A., Hashiguchi,
919 M., Sekijima, T., Kamisoyama, H., Yokoyama, T., Kitagawa, H., Hoshi, N., 2013. Effects of
920 exposure to clothianidin on the reproductive system of male quails. *J. Vet. Med. Sci.* 75, 755-
921 760. <https://doi.org/10.1292/jvms.12-0544>

922

923 Tomizawa, M., Casida, J. E., 2003. Selective toxicity of neonicotinoids attributable to specificity
924 of insect and mammalian nicotinic receptors. *Annu. Rev. Entomol.* 48, 339-364.
925 <https://doi.org/10.1146/annurev.ento.48.091801.112731>

926

927 Tomizawa, M., Casida, J. E., 2005. Neonicotinoid insecticide toxicology: mechanisms of
928 selective action. *Annu. Rev. Pharmacol.* 45, 247-268.
929 <https://doi.org/10.1146/annurev.pharmtox.45.120403.095930>

930

931 Wang, Y., Han, Y., Xie, Y., Xu, P., Li, W., 2018. The metabolism distribution and effect of
932 dinotefuran in Chinese lizards (*Eremias argus*). *Chemosphere*. 211, 591-599.
933 <https://doi.org/10.1016/j.chemosphere.2018.07.181>
934

935 Walker, C. H., 2003. Neurotoxic pesticides and behavioural effects upon
936 birds. *Ecotoxicology*. 12, 307-316. <https://doi.org/10.1023/A:1022523331343>
937

938 Williams, O., 2015. The Woodcock Network. <http://www.ringwoodcock.net/index.shtml>
939 Accessed date: 12 October 2022.
940

941 Wintermantel, D., Odoux, J. F., Decourtye, A., Henry, M., Allier, F., Bretagnolle, V., 2020.
942 Neonicotinoid-induced mortality risk for bees foraging on oilseed rape nectar persists despite
943 EU moratorium. *Sci. Total Environ.* 704, 135400.
944 <https://doi.org/10.1016/j.scitotenv.2019.135400>
945

946 Wintermantel, D., Locke, B., Andersson, G. K., Semberg, E., Forsgren, E., Osterman, J.,
947 Pedersen, T. R., Bommarco, R., Smith, H. G., Rundlöf, M., de Miranda, J. R., 2018. Field-level
948 clothianidin exposure affects bumblebees but generally not their pathogens. *Nat. Commun.*
949 9, 1-10. <https://doi.org/10.1038/s41467-018-07914-3>
950

951 Wood, T. J., Goulson, D., 2017. The environmental risks of neonicotinoid pesticides: a review
952 of the evidence post 2013. *Sci. Pollut. R.* 24, 17285-17325. [https://doi.org/10.1007/s11356-](https://doi.org/10.1007/s11356-017-9240-x)
953 [017-9240-x](https://doi.org/10.1007/s11356-017-9240-x)
954

955 Wretenberg, J., Lindström, Å., Svensson, S., Thierfelder, T., Pärt, T., 2006. Population trends
956 of farmland birds in Sweden and England: similar trends but different patterns of agricultural
957 intensification. *J. Appl. Ecol.* 43, 1110-1120. [https://doi.org/10.1111/j.1365-
958 2664.2006.01216.x](https://doi.org/10.1111/j.1365-2664.2006.01216.x)

959

960 Zhang, Q., Hu, S., Dai, W., Gu, S., Ying, Z., Wang, R., & Lu, C. (2023). The partitioning and
961 distribution of neonicotinoid insecticides in human blood. *Environmental Pollution*, 121082.
962 <https://doi.org/10.1016/j.envpol.2023.121082>

963

964 Zhang, C., Li, X., Li, F., Li, G., Niu, G., Chen, H., Ying, G.-G., Huang, M., 2022. Accurate prediction
965 and further dissection of neonicotinoid elimination in the water treatment by CTS@ AgBC
966 using multihead attention-based convolutional neural network combined with the time-
967 dependent Cox regression model. *J. Hazard. Mater.* 423, 127029.
968 <https://doi.org/10.1016/j.jhazmat.2021.127029>

969 **Caption for figures**


970

971 **Figure 1.** Spatial distribution of birds' catching sites across the Long-Term Socio-Ecological
972 Research Zone Atelier Plaine & Val de Sèvres (LTSER ZAPVS). Landscape features provided
973 correspond to data available for 2021 from our GIS database. Infrastructures correspond to
974 buildings, sport fields, cemeteries, locks, and bridges present in the study area. The five bird
975 species monitored in the study area are blackbird (*Turdus merula*, $N_{2021} = 64$), ciril bunting
976 (*Emberiza cirilus*, $N_{2021} = 31$), common nightingale (*Luscinia megarhynchos*, $N_{2021} = 34$), grey
977 partridge (*Perdix perdix*, $N_{2020-2021} = 23$ and $N_{2021-2022} = 31$), and chicks of Montagu's harrier
978 (*Circus pygargus*, $N_{2021} = 55$). N corresponds to the number of adults sampled, except in the
979 case of Montagu's harrier where N corresponds to the number of chicks sampled.

980

981 **Figure 2.** Wild farmland birds' exposure to 5 neonicotinoids. Distribution of the different
982 neonicotinoid molecules for each individual of each species are represented as one stacked
983 bar. Distribution in percent was obtained from raw concentrations of each molecule in birds
984 blood sampled.

985



986


987

Figure 1.

988

Fuentes et al.

989



990

991

992

Figure 2.

Fuentes et al.

993 **Caption for tables**

994

995 **Table 1.** Summary of the mean concentrations [c] with standard deviations (SD) and range in $\text{pg}\cdot\mu\text{L}^{-1}$ with their sample size (n) obtained by LC-
 996 MS/MS above the limits of detection (LOD) for each species. The sample size for each species is also provided (N). The percentage of samples
 997 above the LOD (n/N) is given in brackets (rounded to the nearest percent). Blackbirds are not referenced as no neonic was detected in the
 998 sampled individuals.

	Clothianidin	Dinotefuran	Nitenpyram	Thiacloprid	Thiamethoxam
Cirl buntings (N = 34)					
n	5 (15%)	0	0	34 (100%)	4 (12%)
[c] ± SD	2.28±1.55	-	-	LOD < [c] < LOQ	2.59±1.37
range	1.04–4.87	-	-	-	1.61–4.56
Common nightingales (N = 25)					
n	2 (8%)	0	0	25 (100%)	1 (4%)
[c] ± SD	0.05±0.06	-	-	LOD < [c] < LOQ	0.06
range	0.009–0.093	-	-	-	-

Grey partridges (N = 54)

<i>Winter 2020/2021 (N = 23)</i>	n	0	3 (13%)	20 (87%)	1 (4%)	1 (4%)
	[c] ± SD	-	6.20±3.62	23.10±11.33	0.07	23.73
	range	-	2.14–9.10	1.24–41.03	-	-

<i>Winter 2021/2022 (N = 31)</i>	n	8 (26%)	10 (32%)	29 (94%)	0	3 (10%)
	[c] ± SD	5.26±4.10	7.85±4.07	18.29±10.84	-	1.64±0.46
	range	1.92–14.26	3.32–16.61	1.78–43.53	-	1.23–2.14

Montagu's harriers (N = 55)

	n	3 (5%)	0	0	1 (2%)	0
	[c] ± SD	951.60±1299.21	-	-	89.58	-
	range	194.48–2 451.78	-	-	-	-

999

1000