



**HAL**  
open science

## **Role of miRNA in the Transmission of Metabolic Diseases Associated With Paternal Diet-Induced Obesity**

Charlotte Dupont, Laurent Kappeler, Sarah Saget, Valérie Grandjean, Rachel Lévy

### ► **To cite this version:**

Charlotte Dupont, Laurent Kappeler, Sarah Saget, Valérie Grandjean, Rachel Lévy. Role of miRNA in the Transmission of Metabolic Diseases Associated With Paternal Diet-Induced Obesity. *Frontiers in Genetics*, 2019, 10 (4), pp.e0210770. <10.3389/fgene.2019.00337>. <hal-03992408>

**HAL Id: hal-03992408**

**<https://hal.science/hal-03992408v1>**

Submitted on 23 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# **Role of miRNA in the transmission of metabolic diseases associated with paternal diet-induced obesity**

Charlotte Dupont<sup>1</sup>, Laurent Kappeler<sup>2</sup>, Sarah Saget<sup>2</sup>, Valérie Grandjean<sup>3</sup>, Rachel Lévy<sup>1</sup>

## Address:

1. Sorbonne Université, Inserm, Centre de Recherche St-Antoine, CRSA, AP-HP, Hôpital Tenon, Service de biologie de la reproduction-CECOS, F-75020 Paris, France

2. Sorbonne Université, INSERM, Institute of Cardiometabolism, Centre de Recherche St-Antoine, CRSA, F-75012 Paris, France

3. INSERM U1065, Université Cote d'Azur, Team Control of gene expression(10), 151 Route de Ginestière BP 2 3194, F-06204 NICE Cedex 3, France

Key words : miRNA, inter-generational, epigenetics, DOHaD, POHaD, metabolic disorders, paternal transmission

Running title: miRNA and paternal transmission of metabolic diseases

## **Abstract**

The concept of DOHaD (Developmental Origins of Health and Diseases) recognizes that an unfavourable maternal environment alters the developmental trajectory of the fetus and can lead to long-term risk of developing chronic non-communicable diseases. More recently, the concept of a paternal transmission (POHaD) has emerged stressing the impact of paternal overweight or obesity on offspring's health and development. While very few examples of paternal epigenetic inheritance of metabolic disorders have been evidenced in human, many experimental mouse models based on high fat diet-induced paternal obesity have been developed to breakdown molecular mechanisms involved in the process. Besides DNA methylation and chromatin structure, sperm short non-coding RNAs have been considered as the main epigenetic vector of inheritance of paternally environmentally induced changes. Among them, sperm miRNA are one particular subspecies sensitive to environmental changes and obesity can modify the sperm miRNA profile. Once delivered into the zygote, these molecules might induce epigenetic modifications in the embryo, leading to consequences for fetus development and offspring physical and metabolic health latter on in life. Furthermore, some data also suggest that metabolic pathologies may be intergenerationally or transgenerationally transmitted.

## **Introduction**

In the 1990s, Barker's epidemiological studies have highlighted a predisposition of metabolic syndrome in adulthood in low birth weight individuals. This "Barker hypothesis" has been the starting point of many works (Hales and Barker, 1992; Barker et al., 2002). The concept of maternal nutritional footprint has emerged and has extended towards the concept of DOHaD (Developmental Origins of Health and Diseases) (Gluckman and Hanson, 2004). This concept recognizes that maternal environment can alter the trajectory of fetal development through various mechanisms such as hormonal imbalance, oxidative stress and epigenetic changes. Organs development and function may be affected, leading to a long-term increased risk of developing several pathologies such as metabolic syndrome, cardiovascular diseases, autoimmune diseases, neurodegenerative diseases, cancers, and infertility (Eriksson, 2016).

More recently, the concept of paternal transmission of environmental messages (POHaD : paternal origins of health and diseases) has emerged (Soubry, 2018). In humans, epidemiological studies in Sweden, have indicated that paternal grandparents' exposure to famine may predispose subsequent generations to obesity and cardiovascular diseases (Kaati et al., 2002; Kaati et al., 2007). More recently, it was shown that paternal grandfather's nutrition in pre-puberty may predict grandchildren's mortality from cancer (Vagero et al., 2018). This phenomenon has been more widely studied in rodent models, although numerous clinical studies are in progress. Indeed, male mice offspring of fathers fed a low-protein diet presented lower liver cholesterol esters, impaired liver gene expression involved in the biosynthesis of lipids and cholesterol and changes in cytosine methylation at several loci (Carone et al., 2010). In a paternal model of high fat diet (HFD), metabolic disorders and alteration of  $\beta$ -cell function

have been observed in adult female rats' offspring (Ng et al., 2010). Pancreatic islet and retroperitoneal white adipose tissue transcriptomes were both altered. Expression of genes involved in aging and non-communicable chronic disorders was particularly affected (Ng et al., 2014). Separately, in a mouse model of HFD-induced obesity, it was reported that male and female offspring developed glucose intolerance, insulin resistance and obesity as well as impaired reproductive functions in both male and female, F1 and F2 offspring (Fullston et al., 2013). However, the phenotypes were differentially expressed according to a sexual dimorphism: obesity and glucose intolerance were observed in both F1 males and females, but only in the F2 males from the F1 maternal line. Insulin resistance was found in all animals except the F2 males from the F1 male line. Gametes quantity and quality were affected in both F1 and F2, male and female (Fullston et al., 2013).

As indicated above, environment may affect several subsequent generations. Consequently, it is important to define the terms “transgenerational” and “intergenerational”. During pregnancy or gestation, the maternal (F0) environment can modulate the development of the fetus (F1), which can have long-term effects. Importantly, this may involve development of all fetal tissues including the germ cells in reproductive ones, which may directly affect the second (F2) generation. This is called multigenerational or intergenerational transmission. In the case of transmission to subsequent generations (F3 and above), the transmission is called transgenerational because there is no direct contact between the environment and the individual's cells. If the exposure is paternal, intergenerational transmission is limited to the F1 generation that has been exposed by paternal gametes to the environment. For subsequent generations (F2 and above) we speak of transgenerational transmission (Skinner, 2010; Heard and Martienssen, 2014; Miska and Ferguson-Smith, 2016). With the concept of intergenerational and transgenerational programming, the notion of sexual dimorphism has broadened. Indeed, males and females present strong differences in regard to their susceptibility

to unfavourable programming. Sexual dimorphism is observed according to the sex of the transmitting parent, the sex of the individual and the generation (Gabory et al., 2009).

### **The role of epigenetic mechanisms**

Mechanisms involved in the DOHaD and POHaD phenomena are numerous, complex and not completely elucidated. In animal models and in humans, it is known that an unfavourable pre-conceptional paternal environment such as obesity can alter conventional sperm parameters such as sperm concentration and motility (Sermondade et al., 2013; Raad et al., 2019). Importantly, it also alters the quality of gametes and molecular parameter, notably by modifying the integrity of sperm DNA (Dupont et al., 2013) and epigenetic marks (Grandjean et al., 2015). These modifications potentially alter the developmental trajectory of the future embryo (Raad et al., 2019) and consequences of paternal obesity can thus be noted very early in offspring development. Indeed, it was observed that paternal obesity is associated with altered preimplantation embryo development, notably a modified kinetics of cell division and a decreased blastocyst cell number (Bakos et al., 2011a; Binder et al., 2015; Schliep et al., 2015). Epigenetics describes processes of complex modifications of gene expression without changes to the DNA sequence. It's a crucial mechanism for cell specialization and fetus development and serves as an interface between the environment and cells. Moreover, epigenetic mechanisms have important roles in environmental adaptations and may then predict future health of the fetus (Day et al., 2016). The main mechanisms accounting for the epigenetic regulation of gene expression are DNA methylation, post-translational histone modifications and small non-coding RNA.

The epigenetic modulation of the intergenerational impacts of paternal obesity on offspring health and development is increasingly well-documented (Lane et al., 2014). In humans, paternal obesity has been associated with changes in DNA methylation of offspring's cord blood cells. Demethylation in the IGF2 region (Soubry et al., 2013) and in other genes involved in metabolic regulation, such as MEST, PEG3, NNAT, has been observed (Soubry et al., 2015). Moreover, paternal obesity has also been associated with changes in gene expression and global DNA methylation of extraembryonic tissue. PPARA (Peroxisome Proliferator-Activated Receptors) and CASP12 expression were significantly down-regulated in male placentas. In female placentas, significant increase in total DNA methylation was observed (Binder et al., 2015).

Sperm epigenetic modifications may play a critical role in the mechanism of inheritance of environmentally induced changes. The role of DNA methylation in paternal transmission has been widely assessed in animals and several studies have highlighted DNA methylation modifications in case of paternal transmission of diseases. Indeed, de Castro Barbosa *et al.* have observed a reprogramming of sperm DNA methylation profile in male rats fed with a HFD. Eighteen regions were found differentially methylated and 92 genes related to DMRs were found commonly regulated by HFD in both F0 and F1 spermatozoa (de Castro Barbosa et al., 2016). Likewise, it was observed that spermatozoa produced by the obese mice fed with a HFD were hypomethylated (Fullston et al., 2013). Finally, most of the studies show that a small number of sperm loci present differential methylation according to paternal feeding. However, only a limited subset is maintained in the offspring tissues (Sharma and Rando, 2017). Consequently, whether or not DNA methylation is involved in epigenetic inheritance is still debated (Shea et al., 2015). Indeed, sperm DNA is strongly demethylated after fertilization in the oocyte, although imprinted genes escape reprogramming, and undergoes progressive re-methylation in the embryo (Smallwood and Kelsey, 2012). But some genes are more likely to

respond to environmental stress, and some have been recently identified (Kazachenka et al., 2018).

In view of the specific chromatin conformation and notably the replacement of histones by protamines and the compaction in spermatozoa, post-translational histone modifications were not widely studied. Nevertheless, small amounts of histone are not replaced by protamines during spermatogenesis, suggesting their potential role in paternal transmission of non-communicable diseases (Yoshida et al., 2018). Thus, by overexpressing the human KDM1A histone lysine 4 demethylase during mouse spermatogenesis, a team have highlighted the importance of correct histone methylation during spermatogenesis for offspring development (Siklenka et al., 2015). Furthermore, it was shown in a drosophila model of paternal-diet-induced intergenerational metabolic reprogramming, the importance of sperm chromatin state (Ost et al., 2014).

In contrast, there is greater evidence that RNA molecules are strong epigenetic vectors of paternally-mediated inheritance of environmental exposures. Indeed, some studies indicate that male intergenerational epigenetic signals may be transmitted by the sperm itself and/or by seminal fluid (Rando, 2012; Gannon et al., 2014), which contain a high concentration of small RNA (Figure 1). Indeed, some data indicate that seminal plasma alone may influence offspring development, as it was observed in a model of mice offspring conceived by artificial insemination associated to a mating with vasectomized male fed a low protein diet (Watkins et al., 2018).

### **Implication of Small RNAs**

Male germ cells are rich in small non coding RNAs (sncRNAs) such as micro-RNA (miRNA), endogenous small interfering RNAs (endo-siRNAs), piwi-interacting RNAs (piRNAs) and

transfer RNAs (tRNAs). sncRNAs don't encode functional proteins and according to their type, their length varies; miRNAs are about 22 nucleotides in length, endo-siRNAs are 21 nucleotides in length and piRNAs are approximately 26-31 nucleotides in length (Song et al., 2011). tsRNAs can be divided into two main types : tRNA-derived stress-induced RNA (tiRNAs) with 28–36 nts length and tRNA-derived fragment (trfRNAs) with 14–30 nts length (Li et al., 2018). tRNA-derived small RNAs (tsRNAs) derive from tRNA and have attracted great attention as they can have many regulatory roles (Li et al., 2018). Thus, a difference in expression level of specific miRNAs, piRNAs and tRNA fragments between lean and obese men spermatozoa has been observed. The authors have suggested that the altered piRNAs expression may contribute to offspring predisposition to obesity (Donkin et al., 2016). More importantly, injection of tsRNA, isolated from spermatozoa collected from HFD male, into wild-type zygotes, led to metabolic disorders in the F1 offspring. Altered expression of genes involved in metabolic pathways was observed in the early embryos and islets of F1 offspring (Chen et al., 2016). Finally, it was, also, observed in mice sperm that tRNA-gly and tRNA-Pro were both altered by running. Moreover, in this study, paternal exercise was associated with modified behavioural phenotype in offspring (Short et al., 2017).

Among the small noncoding RNAs, the microRNAs (miRNA) are by far the most studied ones and we focus our review on them. miRNAs derive from long primary transcripts (pri-miRNA) and are modified in the nucleus by RNase III (Drosha) to form precursors (pre-miRNA). After the nucleocytoplasmic translocation *via* exportin 5, the pre-miRNAs are modified by the enzyme Dicer into mature miRNAs. miRNA regulate gene expression at a post-transcriptional level by inhibiting the translation of their respective target genes by homologous binding to RNA (Bartel, 2004). They regulate more than 60% of the genes encoding proteins (Friedman et al., 2009). miRNAs are involved in almost all cellular functions, and in particular, proliferation, differentiation and apoptosis (Bueno and Malumbres, 2011). They play a critical

role in the pathogenesis of diseases such as cancer, cardiovascular diseases, endocrine dysfunctions, and infectious illness. (Lekka and Hall, 2018).

### **miRNA and male reproductive functions**

miRNAs are abundantly present in spermatozoa and seminal plasma. Alteration of miRNA profile may challenge spermatogenesis and sperm maturation and may be involved in male fertility (Gunes et al., 2016). Indeed, it has been observed that some miRNAs, such as miR-17-92 cluster and miR-290-295 cluster, are highly expressed in primordial germ cells and spermatogonia. The crucial role of miRNAs in spermatogenesis has been shown in Dicer-Knockout mice that presented altered spermatogenesis (Hayashi et al., 2008). [MiRNAs modulates the onset of primordial germ cell and somatic cell differentiation during mouse embryonic gonadal development](#) (Miska and Ferguson-Smith, 2016). More recently, it was demonstrated that miRNAs are involved in gonadal sexual differentiation in both primordial germ cells and gonadal somatic cells (Fernandez-Perez et al., 2018).

Moreover, miRNAs are differentially expressed through the epididymis regions, thus playing a critical role in regulation of gene expression involved in epididymis functions (Browne et al., 2018). The sperm miRNA profile is modified along the journey through the epididymis, and epididymosomes that contain more than 350 miRNAs that play a critical role in this phenomenon (Reilly et al., 2016). Some miRNA profiles were established in testis, spermatozoa and seminal plasma. Numerous miRNAs have been found down-regulated or up-regulated in azoospermic or oligozoospermic patients (Gunes et al., 2016). It has been difficult to highlight one miRNA or a certain miRNA profile that could precisely identify impairment of the male reproductive system. However, certain miRNAs have been consistently observed

to be down-regulated in case of spermatogenic defects (Gunes et al., 2016). Indeed, miR-34b, miR-34c, miR-19a have been observed down regulated in testicular tissues, spermatozoa or seminal plasma in case of oligozoospermia or azoospermia (Lian et al., 2009; Wang et al., 2011; Abu-Halima et al., 2013; Abu-Halima et al., 2014). Some miRNAs have been also correlated with spermatogenic parameters and sperm DNA integrity. Thus, three miRNAs, were regularly found down-regulated in case of sperm alteration (miR34c-5p, miR-122, miR-34b-5p) and miR-429 was regularly found up-regulated (Abu-Halima et al., 2013; Abu-Halima et al., 2014; Salas-Huetos et al., 2014; Salas-Huetos et al., 2016). These miRNAs could represent new biomarkers of fertility, but due to the variety of miRNAs that exist, further validation is required to develop more accurate profiling panels (Gunes et al., 2016).

### **Role of miRNAs in intergenerational or transgenerational programming**

Sperm miRNA profiles are very sensitive to paternal body weight and dietary factors changes. Thus, it was observed by RNA-seq analysis that obese men have increased expression of miR-155 and miR-122 in plasma and sperm compared to control (Lopez et al., 2018).

In the mouse model with HFD-induced obesity, 13 miRNAs were found differentially expressed in spermatozoa by microRNA array. The authors have hypothesized that these miRNAs may be responsible for paternal transmission of metabolic and reproductive phenotype (Fullston et al., 2016). In the same way, F1, F2, F3 offspring of obese prediabetic male rats were assessed for metabolic functions. Glucose and lipid metabolism defects were observed in the three generations, with the phenotype in the F3 largely attenuated. The authors have highlighted that sperm miRNAs could be involved in the mechanism of inheritance, since 15 miRNAs were observed up-regulated and 9 miRNAs down-regulated, by RNA-seq analysis, in F1 male from obese fathers (Cropley et al., 2016). Offspring and the second generation (F2)

from male rats fed with a HFD were lighter at birth and had a decrease of pancreatic beta-cell mass and the phenotype was also inherited in the F2 (Cropley et al., 2016). Importantly, epigenome (DNA methylation and miRNA profiles) of F0 and F1 HFD-founders' sperm was modified. Using RNA-seq analysis, the authors found 15 miRNAs differentially expressed in HFD-rat sperms compared to control. In particular, the miRNA let-7c-5p was differentially expressed in both F0 and F1 sperms. When assessed in metabolic tissues (liver, gonadal white adipose tissue and EDL (Extensor Digitorum Longus) muscle), let-7c appeared to be altered in HFD-fathers' offspring. Let-7 miRNA family plays a role in the control of lipid and glucose metabolism and insulin signalling pathways (de Castro Barbosa et al., 2016). These results suggest strongly that sperm miRNA might be involved in inter and transgenerational inheritance.

Independently of the first publications about non-genetic paternal transmission of diseases (Carone et al., 2010; Ng et al., 2010; Ng et al., 2014), it was shown that modification of the zygote's miRNA pool could lead to long-term phenotype modifications or pathologies in offspring. Thus, it was observed that injection of miR-1 in fertilized mouse eggs led to higher expression of miR-1 targeted RNA with clinical cardiac consequences (i.e. cardiac hypertrophy). miR-1 was observed abundant in sperm, giving first evidence suggesting a non-genetic transmission of paternal pathologies (Wagner et al., 2008). In an other study, the injection of miR-124 (highly expressed in the brain and involved in central nervous system development) led to large body size associated with up-regulated miR-124-targeted-transcripts like Sox9 (Grandjean et al., 2009). Disorders were observed from the embryo stage, as duplication of inner cell mass and resulted in twin pregnancies. These studies provide first evidence of the role of paternal miRNA in fetal programming. Later, the relationship between sperm miRNA and their role in programming zygotes became clear. Indeed, it was observed

that injection of sperm RNA collected from obese male mice in normal healthy zygotes led to the same metabolic disorders observed in obese fathers' offspring. By small RNA-seq analysis, it was found 22 deregulated miRNA in male mice fed with an obesogenic diet, and direct injection of miR-19b induced similar metabolic disorders in offspring (Grandjean et al., 2015).

On the other hand, it was observed in non-obese mice that long-term physical activity alters the offspring metabolic status. The offspring of father exposed to physical activity exhibited increasing body weight and higher adiposity, glucose intolerance and insulin resistance when exposed to high fat diet compared to control. Fathers exposed to physical intervention presented differential methylation in sperm DNA for some genes and altered spermatocytic miRNA profile. Thus by RNA-seq analysis, miR-483-3p, miR431, miR-21 were found up-regulated in spermatozoa from exercised male mice while miR-221 was found down-regulated. The authors have hypothesized a higher susceptibility to high fat diet induced metabolic diseases in the offspring of fathers that exercised (Murashov et al., 2016). A balance between diet and physical activity is therefore essential.

Finally, miRNA expression is very variable and can change quickly according to the environment. Consequently, it is difficult to find a profile for a specific phenotype. Nevertheless, it may be this specificity that plays a role in the adaptation of tissues and individuals to the environment.

### **Potential pathways involved in miRNA-mediated epigenetic inheritance**

Sperm miRNA are transferred into oocyte cytoplasm during fertilization and can modulate gene expression during embryo development. miRNAs are crucial for early embryo-development. Indeed, sperm miRNA deficiency, induced by Dicer and Drosha conditional knockout (cKO) in mice, impaired preimplantation developmental potential, which can be

reversed by injection of RNA or small RNA (Yuan et al., 2016). Also, the blastocyst formation rate and the live birth rate of embryos resulting from intracytoplasmic sperm injection (ICSI) using sperm depleted of 90% of its content of RNA after RNase treatment were significantly decreased (Guo et al., 2017). On the other hand, it has been shown that sperm miRNAs modulate zygotic genome activation and might play a critical role in regulating maternal miRNA turnover during zygote first cleavage (Yuan et al., 2016). In this regards, paternal miR-34c has been shown to be essential for the first embryo cleavage (Liu et al., 2012). Furthermore, we hypothesized that miRNAs impact the development of placenta, as it was observed with DNA methylation (Mitchell et al., 2017).

Pathways by which miRNAs subsequently modulate the development of individuals are not well known. Grandjean *et al.* have hypothesised that sperm miRNAs firstly act as gene modulators, but that the maintenance of the phenotype could be caused by modification of the chromatin structure, modulating then gene expression (Grandjean et al., 2009). It should be also noted that other signals than those transmitted by the sperm cells may program offspring (Robertson, 2007). As in blood, extracellular vesicles are present in seminal fluid and carry miRNA. Extracellular vesicles (EVs) are membrane-bound vesicles that carry and transfer molecules mediating intercellular communications (Cho et al., 2018). Recent studies indicated that they seem to play a role in gametogenesis, post-testicular sperm maturation, fertilization and to contribute to embryo development (Machtinger et al., 2016). Seminal plasma is essential for sperm protection and maturation. It contains proteins that can modulate sperm functions through exosomes bound to spermatozoa surface (Samanta et al., 2018). Small extracellular vesicles (sEVs) of seminal fluid contain miRNA that are sensitive to physiological conditions. Indeed, in azoospermic men, miRNAs profiles of seminal plasma sEVs vary according the presence of spermatozoa obtained from the testicular biopsy (Barcelo et al., 2018).

Epididymosomes, secreted from the epididymal stroma are able to convey protein cargo to the sperm (Nixon et al., 2018). They play a critical role in post testicular sperm maturation and can transfer miRNA to mature spermatozoa (Reilly et al., 2016). Furthermore, circulating extracellular vesicles derived miRNAs have been identified as biomarker of early embryonic viability in cattle (Pohler et al., 2017), suggesting their role in early embryonic development. Consequently, miRNAs present in the extracellular vesicles, such as epididymosomes, and miRNAs free in seminal fluids may program offspring long-term health.

Concerning offspring phenotypes transmitted by paternal stress, the role of sncRNAs and more particularly the role of miRNAs has been highlighted. In addition, glucocorticoids such as corticosterone seem deeply involved in this phenomenon (Pang et al., 2017). Thus, Chan et al. have shown that the miRNAs present in the epididymosomes of the epididymis cauda are regulated by glucocorticoid receptors and play a crucial role in the transmission of paternal stress (Chan et al., 2018). These data suggest that the endocrine system may be involved in paternal programming. Other hormonal pathways may also be involved in the transmission of metabolic diseases.

## **Perspectives**

It is well known that paternal obesity or metabolic disorders may alter sperm parameters and ART outcomes (Campbell et al., 2015). There is now growing evidence that paternal obesity also programs a high susceptibility to develop metabolic syndrome in the offspring with potential further male infertility. In a model of HFD-fed mice, an alteration of sperm parameters has been reported in obese males (Bakos et al., 2011b; Mitchell et al., 2011) with an alteration of both metabolic and reproductive health of the offspring. Male offspring presented altered

sperm parameters and female offspring oocytes presented reduced meiotic competence and altered mitochondrial membrane potential (Fullston et al., 2012).

Although they are rare in human, some interventional studies have shown a beneficial effect of weight loss in overweight or obese men on sperm quality or male fertility (Hakonsen et al., 2011; Faure et al., 2014; Lee et al., 2019). These obese fathers improved metabolic health when subjected to diet and/or physical activity and this was also associated with an improvement of their sperm parameters (Palmer et al., 2012). This weight loss associated with physical activity had a beneficial effect on embryonic development and the offspring showed an improvement in the several metabolic features. miRNAs changes may be involved in this phenomenon. Indeed, a normalization of the sperm miRNA profile of the obese fathers subjected to physical intervention and an improvement of the expression of the pancreatic miRNA in the female offspring was observed (McPherson et al., 2015; McPherson et al., 2017). In humans, no data are available for miRNA, whereas deep changes of methylation profile have been reported after bariatric surgery (Donkin et al., 2016).

Data assessing the impact of paternal exercise on offspring health are scarce and controversial, but the effect of physical exercise before conception seems beneficial especially in case of paternal obesity. The model of fathers fed a standard diet and subjected to long-term physical exercise may be similar to models of fathers exposed to under-nutrition programs (Murashov et al., 2016). Thus, in a paternal under-nourished model, offspring presented small birth weight, associated with a catch-up growth and increased adiposity and dyslipidemia later-on. In this model, supplementation with vitamins and antioxidants (vitamin C, vitamin E, folate, lycopene, zinc, selenium and green tea extract) allowed normalization of sperm damages caused by under-nutrition (increase in oxidative lesions and decrease in overall methylation). Interestingly, it also prevented deleterious effects in the offspring (McPherson et al., 2016).

Other environmental disorders or factors affecting paternal lifestyle may have deleterious consequences on offspring health, such as paternal smoking (Beal et al., 2017). Furthermore, several studies have highlighted a transmission of paternal trauma and stress to the offspring through miRNAs (Gapp et al., 2014; Fullston et al., 2017). Moreover, other outcomes than metabolic ones may also be challenged by paternal environment, suggesting a wide range of consequences for the offspring, like breast cancer risk in daughter for example (Fontelles et al., 2016).

Thus multiple environmental signals are able to induce epigenetic changes in the male gamete. Importantly, these modifications are environment-specific since they are able to induce in the respective progenies the phenotype of the father. Which molecular mechanism is involved in this specific modulation of sperm epigenome and/or sperm miRNAs? This remains an open question. To answer this question, it might be interesting to discriminate and identify the specific epigenetic modifications induced by each environment. This could be useful not only to better understand the general molecular mechanism involved in the process of epigenetic inheritance but it could also help to identify specific stress-markers. Whatever the molecular mechanism involved in the establishment of new spermatogenic epigenetic marks, it seems likely that these signals once transferred into the zygote are able to induce modifications which can be transmitted throughout development. Which specific modifications are involved and how are they maintained? To tackle challenges arising from these questions, a global epigenomic analysis - ie ATAC-seq and DNA methylome analysis- on the modified embryos at a very early stage of development could be performed. This would lead us -on the long view – to identify loci epigenetically targeted by environmental signals.

## **Conclusion**

The periconceptual environment and lifestyle factors modify sperm epigenome. This alteration might be maintained in the zygote and throughout development, leading to the inheritance of newly acquired pathologies. The role of sperm miRNA, not only as innovative markers of fertility issues, but also as vectors involved in the inheritance of paternal diseases appears to be crucial. Overweight and obesity seem to alter sperm miRNA profile, leading to transmission of different miRNA profiles in zygote, with consequences on embryo development. In long term, metabolic disorders have been described in offspring F1 and F2, and other pathologies may originate from paternal environment.

In practice, this concept is important to know because it can help to predict and improve the periconceptual period, with a personalised management, and could be a powerful tool for infertility care. Indeed, paternal environment is too often neglected compared to the maternal environment. Identifying sperm miRNA profiles to predict risk of fertility alteration or non-communicable disease inheritance is now required, and needs further investigation, especially in randomized control trials in human. The impact of selected miRNA on IVF outcome, and the possibility to reverse or modify these miRNA profiles, requires our interest. Health professionals should pay more attention to the health of men of childbearing age.

#### **Author contribution**

C.D., L.K., S.S., V.G., R.L. participated in the manuscript's conception and design. C.D, L.K., V.G. participated in drafting of the manuscript. R.L. participated in critical revision of the manuscript for intellectual content and supervised the study.

## References :

- Abu-Halima, M., Hammadeh, M., Backes, C., Fischer, U., Leidinger, P., Lubbad, A.M., et al. (2014). Panel of five microRNAs as potential biomarkers for the diagnosis and assessment of male infertility. *Fertil Steril* 102(4), 989-997 e981. doi: 10.1016/j.fertnstert.2014.07.001.
- Abu-Halima, M., Hammadeh, M., Schmitt, J., Leidinger, P., Keller, A., Meese, E., et al. (2013). Altered microRNA expression profiles of human spermatozoa in patients with different spermatogenic impairments. *Fertil Steril* 99(5), 1249-1255 e1216. doi: 10.1016/j.fertnstert.2012.11.054.
- Bakos, H.W., Henshaw, R.C., Mitchell, M., and Lane, M. (2011a). Paternal body mass index is associated with decreased blastocyst development and reduced live birth rates following assisted reproductive technology. *Fertil Steril* 95(5), 1700-1704. doi: 10.1016/j.fertnstert.2010.11.044.
- Bakos, H.W., Mitchell, M., Setchell, B.P., and Lane, M. (2011b). The effect of paternal diet-induced obesity on sperm function and fertilization in a mouse model. *Int J Androl* 34(5 Pt 1), 402-410. doi: 10.1111/j.1365-2605.2010.01092.x.
- Barcelo, M., Mata, A., Bassas, L., and Larriba, S. (2018). Exosomal microRNAs in seminal plasma are markers of the origin of azoospermia and can predict the presence of sperm in testicular tissue. *Hum Reprod* 33(6), 1087-1098. doi: 10.1093/humrep/dey072.
- Barker, D.J., Eriksson, J.G., Forsen, T., and Osmond, C. (2002). Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol* 31(6), 1235-1239.
- Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116(2), 281-297.
- Beal, M.A., Yauk, C.L., and Marchetti, F. (2017). From sperm to offspring: Assessing the heritable genetic consequences of paternal smoking and potential public health impacts. *Mutat Res* 773, 26-50. doi: 10.1016/j.mrrev.2017.04.001.
- Binder, N.K., Beard, S.A., Kaitu'u-Lino, T.J., Tong, S., Hannan, N.J., and Gardner, D.K. (2015). Paternal obesity in a rodent model affects placental gene expression in a sex-specific manner. *Reproduction* 149(5), 435-444. doi: 10.1530/REP-14-0676.
- Browne, J.A., Leir, S.H., Eggenger, S.E., and Harris, A. (2018). Region-specific microRNA signatures in the human epididymis. *Asian J Androl* 20(6), 539-544. doi: 10.4103/aja.aja\_40\_18.
- Bueno, M.J., and Malumbres, M. (2011). MicroRNAs and the cell cycle. *Biochim Biophys Acta* 1812(5), 592-601. doi: 10.1016/j.bbadis.2011.02.002.
- Campbell, J.M., Lane, M., Owens, J.A., and Bakos, H.W. (2015). Paternal obesity negatively affects male fertility and assisted reproduction outcomes: a systematic review and meta-analysis. *Reprod Biomed Online* 31(5), 593-604. doi: 10.1016/j.rbmo.2015.07.012.
- Carone, B.R., Fauquier, L., Habib, N., Shea, J.M., Hart, C.E., Li, R., et al. (2010). Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell* 143(7), 1084-1096. doi: 10.1016/j.cell.2010.12.008.
- Chan, J., Nugent, B., Morrison, K., Jasarevic, E., Bhanu, N., Garcia, B., et al. (2018). Epididymal glucocorticoid receptors promote intergenerational transmission of paternal stress. *bioRxiv*

- Chen, Q., Yan, M., Cao, Z., Li, X., Zhang, Y., Shi, J., et al. (2016). Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 351(6271), 397-400. doi: 10.1126/science.aad7977.
- Cho, Y.E., Song, B.J., Akbar, M., and Baek, M.C. (2018). Extracellular vesicles as potential biomarkers for alcohol- and drug-induced liver injury and their therapeutic applications. *Pharmacol Ther* 187, 180-194. doi: 10.1016/j.pharmthera.2018.03.009.
- Cropley, J.E., Eaton, S.A., Aiken, A., Young, P.E., Giannoulatou, E., Ho, J.W.K., et al. (2016). Male-lineage transmission of an acquired metabolic phenotype induced by grand-paternal obesity. *Mol Metab* 5(8), 699-708. doi: 10.1016/j.molmet.2016.06.008.
- Day, J., Savani, S., Krempley, B.D., Nguyen, M., and Kitlinska, J.B. (2016). Influence of paternal preconception exposures on their offspring: through epigenetics to phenotype. *Am J Stem Cells* 5(1), 11-18.
- de Castro Barbosa, T., Ingerslev, L.R., Alm, P.S., Versteyshe, S., Massart, J., Rasmussen, M., et al. (2016). High-fat diet reprograms the epigenome of rat spermatozoa and transgenerationally affects metabolism of the offspring. *Mol Metab* 5(3), 184-197. doi: 10.1016/j.molmet.2015.12.002.
- Donkin, I., Versteyshe, S., Ingerslev, L.R., Qian, K., Mehta, M., Nordkap, L., et al. (2016). Obesity and Bariatric Surgery Drive Epigenetic Variation of Spermatozoa in Humans. *Cell Metab* 23(2), 369-378. doi: 10.1016/j.cmet.2015.11.004.
- Dupont, C., Faure, C., Sermondade, N., Boubaya, M., Eustache, F., Clement, P., et al. (2013). Obesity leads to higher risk of sperm DNA damage in infertile patients. *Asian J Androl* 15(5), 622-625. doi: 10.1038/aja.2013.65.
- Eriksson, J.G. (2016). Developmental Origins of Health and Disease - from a small body size at birth to epigenetics. *Ann Med* 48(6), 456-467. doi: 10.1080/07853890.2016.1193786.
- Faure, C., Dupont, C., Baraibar, M.A., Ladouce, R., Cedrin-Durnerin, I., Wolf, J.P., et al. (2014). In subfertile couple, abdominal fat loss in men is associated with improvement of sperm quality and pregnancy: a case-series. *PLoS One* 9(2), e86300. doi: 10.1371/journal.pone.0086300.
- Fernandez-Perez, D., Brieno-Enriquez, M.A., Isolser-Alcaraz, J., Larriba, E., and Del Mazo, J. (2018). MicroRNA dynamics at the onset of primordial germ and somatic cell sex differentiation during mouse embryonic gonad development. *RNA* 24(3), 287-303. doi: 10.1261/rna.062869.117.
- Fontelles, C.C., Carney, E., Clarke, J., Nguyen, N.M., Yin, C., Jin, L., et al. (2016). Paternal overweight is associated with increased breast cancer risk in daughters in a mouse model. *Sci Rep* 6, 28602. doi: 10.1038/srep28602.
- Friedman, R.C., Farh, K.K., Burge, C.B., and Bartel, D.P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 19(1), 92-105. doi: 10.1101/gr.082701.108.
- Fullston, T., McPherson, N.O., Zander-Fox, D., and Lane, M. (2017). The most common vices of men can damage fertility and the health of the next generation. *J Endocrinol* 234(2), F1-F6. doi: 10.1530/JOE-16-0382.
- Fullston, T., Ohlsson Teague, E.M., Palmer, N.O., DeBlasio, M.J., Mitchell, M., Corbett, M., et al. (2013). Paternal obesity initiates metabolic disturbances in two generations of mice with incomplete penetrance to the F2 generation and alters the transcriptional profile of testis and sperm microRNA content. *FASEB J* 27(10), 4226-4243. doi: 10.1096/fj.12-224048.


- Fullston, T., Ohlsson-Teague, E.M., Print, C.G., Sandeman, L.Y., and Lane, M. (2016). Sperm microRNA Content Is Altered in a Mouse Model of Male Obesity, but the Same Suite of microRNAs Are Not Altered in Offspring's Sperm. *PLoS One* 11(11), e0166076. doi: 10.1371/journal.pone.0166076.
- Fullston, T., Palmer, N.O., Owens, J.A., Mitchell, M., Bakos, H.W., and Lane, M. (2012). Diet-induced paternal obesity in the absence of diabetes diminishes the reproductive health of two subsequent generations of mice. *Hum Reprod* 27(5), 1391-1400. doi: 10.1093/humrep/des030.
- Gabory, A., Attig, L., and Junien, C. (2009). Sexual dimorphism in environmental epigenetic programming. *Mol Cell Endocrinol* 304(1-2), 8-18. doi: 10.1016/j.mce.2009.02.015.
- Gannon, J.R., Emery, B.R., Jenkins, T.G., and Carrell, D.T. (2014). The sperm epigenome: implications for the embryo. *Adv Exp Med Biol* 791, 53-66. doi: 10.1007/978-1-4614-7783-9\_4.
- Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., et al. (2014). Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nat Neurosci* 17(5), 667-669. doi: 10.1038/nn.3695.
- Gluckman, P.D., and Hanson, M.A. (2004). Living with the past: evolution, development, and patterns of disease. *Science* 305(5691), 1733-1736. doi: 10.1126/science.1095292.
- Grandjean, V., Fourre, S., De Abreu, D.A., Derieppe, M.A., Remy, J.J., and Rassoulzadegan, M. (2015). RNA-mediated paternal heredity of diet-induced obesity and metabolic disorders. *Sci Rep* 5, 18193. doi: 10.1038/srep18193.
- Grandjean, V., Gounon, P., Wagner, N., Martin, L., Wagner, K.D., Bernex, F., et al. (2009). The miR-124-Sox9 paramutation: RNA-mediated epigenetic control of embryonic and adult growth. *Development* 136(21), 3647-3655. doi: 10.1242/dev.041061.
- Gunes, S., Arslan, M.A., Hekim, G.N.T., and Asci, R. (2016). The role of epigenetics in idiopathic male infertility. *J Assist Reprod Genet* 33(5), 553-569. doi: 10.1007/s10815-016-0682-8.
- Guo, L., Chao, S.B., Xiao, L., Wang, Z.B., Meng, T.G., Li, Y.Y., et al. (2017). Sperm-carried RNAs play critical roles in mouse embryonic development. *Oncotarget* 8(40), 67394-67405. doi: 10.18632/oncotarget.18672.
- Hakonsen, L.B., Thulstrup, A.M., Aggerholm, A.S., Olsen, J., Bonde, J.P., Andersen, C.Y., et al. (2011). Does weight loss improve semen quality and reproductive hormones? Results from a cohort of severely obese men. *Reprod Health* 8, 24. doi: 10.1186/1742-4755-8-24.
- Hales, C.N., and Barker, D.J. (1992). Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 35(7), 595-601.
- Hayashi, K., Chuva de Sousa Lopes, S.M., Kaneda, M., Tang, F., Hajkova, P., Lao, K., et al. (2008). MicroRNA biogenesis is required for mouse primordial germ cell development and spermatogenesis. *PLoS One* 3(3), e1738. doi: 10.1371/journal.pone.0001738.
- Heard, E., and Martienssen, R.A. (2014). Transgenerational epigenetic inheritance: myths and mechanisms. *Cell* 157(1), 95-109. doi: 10.1016/j.cell.2014.02.045.
- Kaati, G., Bygren, L.O., and Edvinsson, S. (2002). Cardiovascular and diabetes mortality determined by nutrition during parents' and grandparents' slow growth period. *Eur J Hum Genet* 10(11), 682-688. doi: 10.1038/sj.ejhg.5200859.

- Kaati, G., Bygren, L.O., Pembrey, M., and Sjöström, M. (2007). Transgenerational response to nutrition, early life circumstances and longevity. *Eur J Hum Genet* 15(7), 784-790. doi: 10.1038/sj.ejhg.5201832.
- Kazachenka, A., Bertozzi, T.M., Sjöberg-Herrera, M.K., Walker, N., Gardner, J., Gunning, R., et al. (2018). Identification, Characterization, and Heritability of Murine Metastable Epialleles: Implications for Non-genetic Inheritance. *Cell* 175(5), 1259-1271 e1213. doi: 10.1016/j.cell.2018.09.043.
- Lane, M., Robker, R.L., and Robertson, S.A. (2014). Parenting from before conception. *Science* 345(6198), 756-760. doi: 10.1126/science.1254400.
- Lee, Y., Dang, J.T., Switzer, N., Yu, J., Tian, C., Birch, D.W., et al. (2019). Impact of Bariatric Surgery on Male Sex Hormones and Sperm Quality: a Systematic Review and Meta-Analysis. *Obes Surg* 29(1), 334-346. doi: 10.1007/s11695-018-3557-5.
- Lekka, E., and Hall, J. (2018). Noncoding RNAs in disease. *FEBS Lett* 592(17), 2884-2900. doi: 10.1002/1873-3468.13182.
- Li, S., Xu, Z., and Sheng, J. (2018). tRNA-Derived Small RNA: A Novel Regulatory Small Non-Coding RNA. *Genes (Basel)* 9(5). doi: 10.3390/genes9050246.
- Lian, J., Zhang, X., Tian, H., Liang, N., Wang, Y., Liang, C., et al. (2009). Altered microRNA expression in patients with non-obstructive azoospermia. *Reprod Biol Endocrinol* 7, 13. doi: 10.1186/1477-7827-7-13.
- Liu, W.M., Pang, R.T., Chiu, P.C., Wong, B.P., Lao, K., Lee, K.F., et al. (2012). Sperm-borne microRNA-34c is required for the first cleavage division in mouse. *Proc Natl Acad Sci U S A* 109(2), 490-494. doi: 10.1073/pnas.1110368109.
- Lopez, P., Castro, A., Florez, M., Miranda, K., Aranda, P., Sanchez-Gonzalez, C., et al. (2018). miR-155 and miR-122 Expression of Spermatozoa in Obese Subjects. *Front Genet* 9, 175. doi: 10.3389/fgene.2018.00175.
- Machtiger, R., Laurent, L.C., and Baccarelli, A.A. (2016). Extracellular vesicles: roles in gamete maturation, fertilization and embryo implantation. *Hum Reprod Update* 22(2), 182-193. doi: 10.1093/humupd/dmv055.
- McPherson, N.O., Fullston, T., Kang, W.X., Sandeman, L.Y., Corbett, M.A., Owens, J.A., et al. (2016). Paternal under-nutrition programs metabolic syndrome in offspring which can be reversed by antioxidant/vitamin food fortification in fathers. *Sci Rep* 6, 27010. doi: 10.1038/srep27010.
- McPherson, N.O., Lane, M., Sandeman, L., Owens, J.A., and Fullston, T. (2017). An Exercise-Only Intervention in Obese Fathers Restores Glucose and Insulin Regulation in Conjunction with the Rescue of Pancreatic Islet Cell Morphology and MicroRNA Expression in Male Offspring. *Nutrients* 9(2). doi: 10.3390/nu9020122.
- McPherson, N.O., Owens, J.A., Fullston, T., and Lane, M. (2015). Preconception diet or exercise intervention in obese fathers normalizes sperm microRNA profile and metabolic syndrome in female offspring. *Am J Physiol Endocrinol Metab* 308(9), E805-821. doi: 10.1152/ajpendo.00013.2015.
- Miska, E.A., and Ferguson-Smith, A.C. (2016). Transgenerational inheritance: Models and mechanisms of non-DNA sequence-based inheritance. *Science* 354(6308), 59-63. doi: 10.1126/science.aaf4945.
- Mitchell, M., Bakos, H.W., and Lane, M. (2011). Paternal diet-induced obesity impairs embryo development and implantation in the mouse. *Fertil Steril* 95(4), 1349-1353. doi: 10.1016/j.fertnstert.2010.09.038.

- Mitchell, M., Strick, R., Strissel, P.L., Dittrich, R., McPherson, N.O., Lane, M., et al. (2017). Gene expression and epigenetic aberrations in F1-placentas fathered by obese males. *Mol Reprod Dev* 84(4), 316-328. doi: 10.1002/mrd.22784.
- Murashov, A.K., Pak, E.S., Koury, M., Ajmera, A., Jeyakumar, M., Parker, M., et al. (2016). Paternal long-term exercise programs offspring for low energy expenditure and increased risk for obesity in mice. *FASEB J* 30(2), 775-784. doi: 10.1096/fj.15-274274.
- Ng, S.F., Lin, R.C., Laybutt, D.R., Barres, R., Owens, J.A., and Morris, M.J. (2010). Chronic high-fat diet in fathers programs beta-cell dysfunction in female rat offspring. *Nature* 467(7318), 963-966. doi: 10.1038/nature09491.
- Ng, S.F., Lin, R.C., Maloney, C.A., Youngson, N.A., Owens, J.A., and Morris, M.J. (2014). Paternal high-fat diet consumption induces common changes in the transcriptomes of retroperitoneal adipose and pancreatic islet tissues in female rat offspring. *FASEB J* 28(4), 1830-1841. doi: 10.1096/fj.13-244046.
- Nixon, B., De Iulius, G.N., Hart, H.M., Zhou, W., Mathe, A., Bernstein, I., et al. (2018). Proteomic profiling of mouse epididymosomes reveals their contributions to post-testicular sperm maturation. *Mol Cell Proteomics*. doi: 10.1074/mcp.RA118.000946.
- Ost, A., Lempradl, A., Casas, E., Weigert, M., Tiko, T., Deniz, M., et al. (2014). Paternal diet defines offspring chromatin state and intergenerational obesity. *Cell* 159(6), 1352-1364. doi: 10.1016/j.cell.2014.11.005.
- Palmer, N.O., Bakos, H.W., Owens, J.A., Setchell, B.P., and Lane, M. (2012). Diet and exercise in an obese mouse fed a high-fat diet improve metabolic health and reverse perturbed sperm function. *Am J Physiol Endocrinol Metab* 302(7), E768-780. doi: 10.1152/ajpendo.00401.2011.
- Pang, T.Y.C., Short, A.K., Bredy, T.W., and Hannan, A.J. (2017). Transgenerational paternal transmission of acquired traits: Stress-induced modification of the sperm regulatory transcriptome and offspring phenotypes. *Curr Opin Behav Sci* 14, 140-147. doi: 10.1016/j.cobeha.2017.02.007.
- Pohler, K.G., Green, J.A., Moley, L.A., Gunewardena, S., Hung, W.T., Payton, R.R., et al. (2017). Circulating microRNA as candidates for early embryonic viability in cattle. *Mol Reprod Dev* 84(8), 731-743. doi: 10.1002/mrd.22856.
- Raad, G., Azouri, J., Rizk, K., Zeidan, N.S., Azouri, J., Grandjean, V., et al. (2019). Adverse effects of paternal obesity on the motile spermatozoa quality. *PLoS One* 14(2), e0211837. doi: 10.1371/journal.pone.0211837.
- Rando, O.J. (2012). Daddy issues: paternal effects on phenotype. *Cell* 151(4), 702-708. doi: 10.1016/j.cell.2012.10.020.
- Reilly, J.N., McLaughlin, E.A., Stanger, S.J., Anderson, A.L., Hutcheon, K., Church, K., et al. (2016). Characterisation of mouse epididymosomes reveals a complex profile of microRNAs and a potential mechanism for modification of the sperm epigenome. *Sci Rep* 6, 31794. doi: 10.1038/srep31794.
- Robertson, S.A. (2007). Seminal fluid signaling in the female reproductive tract: lessons from rodents and pigs. *J Anim Sci* 85(13 Suppl), E36-44. doi: 10.2527/jas.2006-578.
- Salas-Huetos, A., Blanco, J., Vidal, F., Grossmann, M., Pons, M.C., Garrido, N., et al. (2016). Spermatozoa from normozoospermic fertile and infertile individuals convey a distinct miRNA cargo. *Andrology* 4(6), 1028-1036. doi: 10.1111/andr.12276.
- Salas-Huetos, A., Blanco, J., Vidal, F., Mercader, J.M., Garrido, N., and Anton, E. (2014). New insights into the expression profile and function of micro-ribonucleic acid in

- human spermatozoa. *Fertil Steril* 102(1), 213-222 e214. doi: 10.1016/j.fertnstert.2014.03.040.
- Samanta, L., Parida, R., Dias, T.R., and Agarwal, A. (2018). The enigmatic seminal plasma: a proteomics insight from ejaculation to fertilization. *Reprod Biol Endocrinol* 16(1), 41. doi: 10.1186/s12958-018-0358-6.
- Schliep, K.C., Mumford, S.L., Ahrens, K.A., Hotaling, J.M., Carrell, D.T., Link, M., et al. (2015). Effect of male and female body mass index on pregnancy and live birth success after in vitro fertilization. *Fertil Steril* 103(2), 388-395. doi: 10.1016/j.fertnstert.2014.10.048.
- Sermondade, N., Faure, C., Fezeu, L., Shayeb, A.G., Bonde, J.P., Jensen, T.K., et al. (2013). BMI in relation to sperm count: an updated systematic review and collaborative meta-analysis. *Hum Reprod Update* 19(3), 221-231. doi: 10.1093/humupd/dms050.
- Sharma, U., and Rando, O.J. (2017). Metabolic Inputs into the Epigenome. *Cell Metab* 25(3), 544-558. doi: 10.1016/j.cmet.2017.02.003.
- Shea, J.M., Serra, R.W., Carone, B.R., Shulha, H.P., Kucukural, A., Ziller, M.J., et al. (2015). Genetic and Epigenetic Variation, but Not Diet, Shape the Sperm Methyome. *Dev Cell* 35(6), 750-758. doi: 10.1016/j.devcel.2015.11.024.
- Short, A.K., Yeshurun, S., Powell, R., Perreau, V.M., Fox, A., Kim, J.H., et al. (2017). Exercise alters mouse sperm small noncoding RNAs and induces a transgenerational modification of male offspring conditioned fear and anxiety. *Transl Psychiatry* 7(5), e1114. doi: 10.1038/tp.2017.82.
- Siklenka, K., Erkek, S., Godmann, M., Lambrot, R., McGraw, S., Lafleur, C., et al. (2015). Disruption of histone methylation in developing sperm impairs offspring health transgenerationally. *Science* 350(6261), aab2006. doi: 10.1126/science.aab2006.
- Skinner, M.K. (2010). Metabolic disorders: Fathers' nutritional legacy. *Nature* 467(7318), 922-923. doi: 10.1038/467922a.
- Smallwood, S.A., and Kelsey, G. (2012). De novo DNA methylation: a germ cell perspective. *Trends Genet* 28(1), 33-42. doi: 10.1016/j.tig.2011.09.004.
- Song, R., Hennig, G.W., Wu, Q., Jose, C., Zheng, H., and Yan, W. (2011). Male germ cells express abundant endogenous siRNAs. *Proc Natl Acad Sci U S A* 108(32), 13159-13164. doi: 10.1073/pnas.1108567108.
- Soubry, A. (2018). POHaD: why we should study future fathers. *Environ Epigenet* 4(2), dvy007. doi: 10.1093/eep/dvy007.
- Soubry, A., Murphy, S.K., Wang, F., Huang, Z., Vidal, A.C., Fuemmeler, B.F., et al. (2015). Newborns of obese parents have altered DNA methylation patterns at imprinted genes. *Int J Obes (Lond)* 39(4), 650-657. doi: 10.1038/ijo.2013.193.
- Soubry, A., Schildkraut, J.M., Murtha, A., Wang, F., Huang, Z., Bernal, A., et al. (2013). Paternal obesity is associated with IGF2 hypomethylation in newborns: results from a Newborn Epigenetics Study (NEST) cohort. *BMC Med* 11, 29. doi: 10.1186/1741-7015-11-29.
- Vagero, D., Pinger, P.R., Aronsson, V., and van den Berg, G.J. (2018). Paternal grandfather's access to food predicts all-cause and cancer mortality in grandsons. *Nat Commun* 9(1), 5124. doi: 10.1038/s41467-018-07617-9.
- Wagner, K.D., Wagner, N., Ghanbarian, H., Grandjean, V., Gounon, P., Cuzin, F., et al. (2008). RNA induction and inheritance of epigenetic cardiac hypertrophy in the mouse. *Dev Cell* 14(6), 962-969. doi: 10.1016/j.devcel.2008.03.009.

- Wang, C., Yang, C., Chen, X., Yao, B., Yang, C., Zhu, C., et al. (2011). Altered profile of seminal plasma microRNAs in the molecular diagnosis of male infertility. *Clin Chem* 57(12), 1722-1731. doi: 10.1373/clinchem.2011.169714.
- Watkins, A.J., Dias, I., Tsuro, H., Allen, D., Emes, R.D., Moreton, J., et al. (2018). Paternal diet programs offspring health through sperm- and seminal plasma-specific pathways in mice. *Proc Natl Acad Sci U S A* 115(40), 10064-10069. doi: 10.1073/pnas.1806333115.
- Yoshida, K., Muratani, M., Araki, H., Miura, F., Suzuki, T., Dohmae, N., et al. (2018). Mapping of histone-binding sites in histone replacement-completed spermatozoa. *Nat Commun* 9(1), 3885. doi: 10.1038/s41467-018-06243-9.
- Yuan, S., Schuster, A., Tang, C., Yu, T., Ortogero, N., Bao, J., et al. (2016). Sperm-borne miRNAs and endo-siRNAs are important for fertilization and preimplantation embryonic development. *Development* 143(4), 635-647. doi: 10.1242/dev.131755.



**Figure 1 :** Role of miRNAs in the paternal transmission of environmentally-induced metabolic diseases