



HAL
open science

Ingestion of Diazotrophs Makes Corals More Resistant to Heat Stress

Valentine Meunier, Sophie Bonnet, Mercedes Camps, Mar Benavides, Jeff Dubosc, Riccardo Rodolfo-Metalpa, Fanny Houlbrèque

► **To cite this version:**

Valentine Meunier, Sophie Bonnet, Mercedes Camps, Mar Benavides, Jeff Dubosc, et al.. Ingestion of Diazotrophs Makes Corals More Resistant to Heat Stress. *Biomolecules*, 2022, 12 (4), pp.537. 10.3390/biom12040537 . hal-03993353

HAL Id: hal-03993353

<https://hal.science/hal-03993353>

Submitted on 17 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Article

Ingestion of Diazotrophs Makes Corals More Resistant to Heat Stress

Valentine Meunier^{1,*} , Sophie Bonnet², Mercedes Camps², Mar Benavides², Jeff Dubosc³, Riccardo Rodolfo-Metalpa¹ and Fanny Houllbrèque¹ 

¹ ENTROPIE UMR 9220 (CNRS, IRD, UR, UNC, IFREMER) Institut de Recherche pour le Développement, Nouméa 98848, New Caledonia; riccardo.rodolfo-metalpa@ird.fr (R.R.-M.); fanny.houllbreque@ird.fr (F.H.)

² Aix-Marseille University, CNRS, IRD, MIO UM 110, 13288 Marseille, France; sophie.bonnet@mio.osupytheas.fr (S.B.); mercedes.camps@mio.osupytheas.fr (M.C.); mar.benavides@mio.osupytheas.fr (M.B.)

³ Laboratory of Marine Biology and Ecology, Aquarium des Lagons, Noumea 98807, New Caledonia; jeff.dubosc@aquarium.nc

* Correspondence: valentine0meunier@gmail.com

Abstract: Over the past decade, coral bleaching events have continued to recur and intensify. During bleaching, corals expel millions of their symbionts, depriving the host from its main food source. One mechanism used by corals to resist bleaching consists in exploiting food sources other than autotrophy. Among the food sources available in the reefs, dinitrogen (N₂)-fixing prokaryotes or planktonic diazotrophs (hereafter called 'PD') have the particularity to reduce atmospheric dinitrogen (N₂) and release part of this nitrogen (diazotroph-derived nitrogen or DDN) in bioavailable form. Here, we submitted coral colonies of *Stylophora pistillata*, fed or not with planktonic diazotrophs, to a temperature stress of up to 31 ± 0.5 °C and measured their physiological responses (photosynthetic efficiency, symbiont density, and growth rates). Heat-unfed colonies died 8 days after the heat stress while heat-PD-fed corals remained alive after 10 days of heat stress. The supply of PD allowed corals to maintain minimal chlorophyll concentration and symbiont density, sustaining photosynthetic efficiency and stimulating coral growth of up to 48% compared to unfed ones. By providing an alternative source of bioavailable nitrogen and carbon, this specific planktonic diazotroph feeding may have a profound potential for coral bleaching recovery.

Keywords: coral; heat stress; coral bleaching; heterotrophy; diazotrophy; climate change



Citation: Meunier, V.; Bonnet, S.; Camps, M.; Benavides, M.; Dubosc, J.; Rodolfo-Metalpa, R.; Houllbrèque, F. Ingestion of Diazotrophs Makes Corals More Resistant to Heat Stress. *Biomolecules* **2022**, *12*, 537. <https://doi.org/10.3390/biom12040537>

Academic Editor: Anika Wagner

Received: 23 January 2022

Accepted: 13 March 2022

Published: 2 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The success of scleractinian corals in nutrient-poor waters surrounding coral reefs is due to their symbiosis with unicellular dinoflagellates belonging to the Symbiodiniaceae family [1]. These symbionts translocate a large fraction of photosynthetically fixed carbon (C) and amino acids to the host for its nutritional needs [2]. Coral reefs are experiencing increasingly frequent and devastating bleaching events [3]. During bleaching, corals expel millions of their symbionts and/or suffer a loss of photosynthetic pigments, depriving the host from its main nutrition source [4,5]. This process leads to coral death and to the decline of coral reefs if bleached corals are not able to rapidly recover their symbionts [3]. Bleaching susceptibility varies among species, depths, and locations (e.g., [6–9]), and is also influenced by coral morphology [10,11], physiological responses of both animal and symbionts [12,13], and symbiont types hosted by corals (e.g., [14–16]).

Another mechanism used to resist and recover from bleaching consists in exploiting food sources other than autotrophy. Indeed, corals are voracious predators that can feed on a wide range of prey ranging from pico-nanoplanktonic cells (with a size < 20 µm; [17]) to macrozooplankton [18,19]. Multiple laboratory experiments have shown that heterotrophy increases skeletal and tissue growth [17,20–23], allows corals to build up energy

reserves [24,25], increases fertility [26], and reduces sensitivity to acidification [27–29]. Coral ability to switch from an autotrophic to a heterotrophic diet by increasing their feeding rates on zooplankton [22,30,31], making them more resistant to bleaching [30,32–34], has been much less studied. Only two studies have shown that corals increase their consumption of pico-nanoplankton during heat stress [35,36]. Among these small preys, dinitrogen (N₂)-fixing prokaryotes (subsequently referred to as planktonic diazotrophs, hereafter called ‘PD’) have received little attention. Many coral reef ecosystems are characterized by high planktonic diazotroph abundance and activity; they are very widespread in the Western South Pacific (e.g., New Caledonia, Papua New Guinea, and the Australian Great Barrier Reef) [37–40] but also in Hawaii, in the Caribbean, and the Red Sea [41–43]. In New Caledonia, planktonic diazotrophs support ~80% of the primary production of other phytoplankton in summer [44,45]. The lagoon is dominated by picoplankton (non-diazotrophic, i.e., *Synechococcus* and *Prochlorococcus*) but the diazotrophs, even if they are less abundant, are much larger (nanoplankton and microplankton size) and thus represent an important part of the total biomass [46]. Planktonic diazotrophs reduce atmospheric N₂ into bioavailable ammonium (NH₄⁺), and release part of the recently fixed nitrogen (Diazotroph-Derived Nitrogen, DDN) in seawater, providing sufficient nitrogen stocks for the development of the planktonic food web in oligotrophic waters [47]. These planktonic diazotroph cells are thus, by definition, very rich in nitrogen [48,49]. Their ingestion by scleractinian corals was first demonstrated by [50] using the ¹⁵N₂ labelling method. Direct consumption of planktonic diazotrophs would provide $0.76 \pm 0.15 \mu\text{g N cm}^{-2} \text{ h}^{-1}$ to corals, which corresponds to six times the daily N supply by non-diazotrophic pico- and nanoplankton [17]. Bleached colonies of *S. pistillata* incorporate more PD than healthy colonies [36], but the consequences on coral metabolism are not known. By providing an alternative source of bioavailable N and C, this increased incorporation of PD may have a profound influence on coral bleaching recovery.

To improve our understanding of the physiological response of corals to heat stress in anticipation of more frequent and severe coral bleaching events, we investigated if and how nutrition on PD is involved in the resistance to high temperature of the coral species *S. pistillata*. We evaluated key physiological measurements (photosynthetic efficiency, growth rates) and tissue parameters (chlorophyll, protein concentrations, and symbiont densities) of colonies of *S. pistillata*, fed or not with PD during heat stress.

2. Materials and Methods

2.1. Diazotrophic Strains Acclimation

Before the experiment, 4 strains of PD isolated from the Western Tropical South Pacific (WTSP) (details in Table 1) were grown in an F/2 culture medium [51] modified without nitrate to ensure obligate diazotrophy. This N-free F/2 was made with 0.2 µm-filtered sterilized seawater collected in the New Caledonian Lagoon (the provenance of studied corals, 166.44° E, 22.48° S), to prevent any physiological stress when added to aquaria containing corals colonies. Strains were cultivated in 2 L Erlenmeyer flasks in semi-continuous conditions, i.e., they were maintained under an exponential growing phase for the entire duration of the experiment (21 days), by adding 300 mL of new medium every 2 days. In vivo chlorophyll, fluorescence was measured every 3 days (model Trilogy, Turner Designs). A mixture of these 4 strains was given to some of the corals (see description of the protocol below).

Table 1. Details of the 4 cultures of planktonic diazotrophs isolated from the Western Tropical South Pacific.

Genus	Morphological Type	Size	Location
<i>Xenococcus</i> spp.	Unicellular	~6 μm	Noumea lagoon
<i>Cyanothece</i> spp.	Unicellular	~5.5 μm	Noumea lagoon
<i>Calothrix</i> spp.	Filamentous heterocystous	~50 μm	WTSP Ocean
<i>Rivularia</i> spp.	Filamentous heterocystous	~30 μm	WTSP Ocean

2.2. Coral Preparation and Acclimation

One hundred and sixty terminal branch portions (3–5 cm) of *S. pistillata* were cut from 20 different parent colonies in the New Caledonian lagoon (22° 21' 339" S; 166° 15' 404" E; sampling license issued by the 'Province Sud', Government of New Caledonia). Coral fragments were transported from the collection site in a cooler and randomly assigned experimental tanks. They were hung on nylon wires and suspended in eight 6.5 L aquaria (around 20 coral fragments per aquarium), spaced apart from each other without ever touching. All physicochemical parameters (temperature, salinity, nutrient concentration, and pH) were rigorously controlled in each tank so that the only difference between the tanks was temperature. Therefore, despite the fact that this setup could be considered as a pseudoreplication, the results obtained are representative, and the most parsimonious explanation for the differences observed between the coral batches is clearly the feeding and not stochastic errors. Fragments were maintained in an open seawater system, continuously supplied with 40 μm -filtered seawater, renewed at a rate of 16.5 L h⁻¹, and mixed using a submersible pump (Aquarium system, micro-jet MC 320, Mentor, OH, USA). In each tank, coral colonies were spaced apart and never touched each other. Temperature (28 ± 0.2 °C) and salinity (35.80 ± 0.01) were kept constant using heaters connected to electronic controllers and monitored with a YSI MPS 556 probe (YSI, Yellow Springs, OH, USA). Corals received a constant irradiance of 120 ± 10 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ (photoperiod 12 h:12 h light:dark) using four Aquablue plus neon bulbs (blue–white, 15,000 K, Giesemann, Germany). All coral fragments were left to acclimate and grow in these conditions for three weeks before starting coral feeding and carrying out the first sampling of the colonies (T₀ sampling).

2.3. Experimental Setup and Conditions

After T₀ and during the 55 days of the experiment, the environmental parameters (temperature, irradiance) were the same as those described above and were measured in each of the 8 aquaria. In 4 aquaria, coral fragments were not fed (called unfed corals), while in the other 4, coral fragments were fed exclusively with PD (called PD-fed corals) (Figure 1). These corals were fed 3 times/week with a mix of the 4 strains of PD. To prepare the PD 'solution', the mix of the 4 strains (100 mL) was first centrifuged at 5000 × g for 4 min 30 s to obtain a pellet and remove the entire culture medium. The pellet was then gently resuspended in 0.2 μm -filtered seawater. Between 100 and 200 mL of strain mix were added in each tank to reach final concentrations of 10³ and 10⁶ per mL, corresponding to the average concentration measured in the NC lagoon [37,52,53]. Each feeding episode was carried out for 2 h after the lights were switched off. Seawater circulation in the tanks was cut off during this time to avoid the flushing of prey. Both PD-fed and unfed corals were maintained in these conditions for 55 days before they were exposed for 10 additional days to a gradual increase in temperature of 0.3 °C per day until reaching 31 ± 0.5 °C on day 65 of the experiment (hereafter called T₆₅). The same feeding protocol was used throughout the experiment, before and during the heat stress. PD-fed and unfed corals submitted to heat stress are referred to as 'heat-PD-fed' and 'heat-unfed' corals, respectively. PD-fed and unfed corals maintained at the initial temperature of 28 ± 0.2 °C are referred to as 'PD-fed' and 'unfed' corals under ambient conditions. For each measured parameter, 6 coral fragments per time (T₀ and T₆₅) and per condition were sampled. Therefore, measurements

of photosynthesis, Symbiodiniaceae densities, chlorophyll, and protein concentrations were performed on the same coral fragment.




Figure 1. Experimental design with two feeding regimes. Each condition was duplicated with 20 fragments per tank. Ambient conditions of 28 ± 0.2 °C. Heat stress conditions with a temperature increase of up to 31 ± 0.5 °C.

2.4. Symbiodiniaceae Density and Total Chlorophyll, and Protein Content

Six coral fragments per time (T_0 and T_{65}) and per condition were sampled. Tissue was removed from the skeleton using an air pick and homogenized with a Potter tissue grinder [54,55]. Symbiont cells were counted 3 times with light microscopy using a Neubauer's cell. Subsamples of 10 mL from each coral tissue solution were centrifuged at $8000 \times g$ for 10 min at 4 °C, and the supernatant was removed. The pellet containing the symbiont was resuspended in 100% acetone for 24 h (4 °C) in the dark to extract chlorophyll *a* and *c*₂. The extracts were centrifuged at $10,000 \times g$ for 15 min and absorbances were read at 630, 663, and 750 nm. Chlorophyll *a* and *c*₂ concentrations were computed according to the spectrometric equations from [56,57]. Chlorophyll *a* and *c*₂ are given as total chlorophyll. All measurements were normalized to the skeletal surface area (cm^2), estimated using the paraffin wax-dipping method [58,59]. Sub-samples were taken at the beginning and after 30 days of experiments to assess the total protein content (mg prot cm^{-2}) according to [60] and [57] with a BCAssay Kit (Uptima, Interchim).

2.5. Growth Rates

Growth rates were monitored every 15 days on every coral fragment using the buoyant weight (BW) technique [61]. BW gives an estimation of the calcification rates of corals over long time scales and considers both daytime and nighttime calcification rates. Samples hung on a nylon wire were weighed using a Mettler XP205 electronic balance (readability of 0.01 mg) in seawater of known density, and temperature and salinity were continuously measured. The net BW of the corals was converted into dry weight using the density of pure aragonite (2.94 g cm^{-3}). Their growth rates were calculated as the change in dry weight between the initial and the final weight and expressed in $\text{mg g}^{-1} \text{ d}^{-1}$.

2.6. Photosynthetic Efficiency Measurements

Photosynthetic efficiency (F_v/F_m) and the electron transport rate (ETR) of the Photosystem II (PSII) of symbiont in hospite were measured at T_0 and at T_{65} using a DIVING-PAM fluorometer (DIVING-PAM, Walz, Germany [62]). Measurements were performed on 12 coral fragments for each condition. Before the measurements, these coral fragments were

dark-adapted for 15 min (after [63]). The 8 mm optical fiber was maintained perpendicular to the fragments' surface using a black-jacket at a fixed distance of 5 mm guaranteeing a correct distance of the optical fiber from the coral. A weak-pulse red light (max. intensity $< 1 \text{ mol photon m}^{-2} \text{ s}^{-1}$, width 3 μs , frequency 0.6 kHz) and a saturating pulse of actinic light (max. intensity $8000 \text{ } \mu\text{mol photon m}^{-2} \text{ s}^{-1}$, pulse width 800 ms) were applied to the coral surface to measure the minimal (F_0) and maximal (F_m) Chl a fluorescence, respectively. ETR was obtained through the rapid light curves (RLCs). RLCs were collected by illuminating corals for 10-s periods with eight actinic light sequences from 0 to ca. $1500 \text{ } \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. Variable fluorescence (F_v) was calculated resulting in $F_m - F_0$ [64] and F_v / F_m values that may be used as a quantitative measure of photo-inactivation during coral bleaching [16]. F_v / F_m values of healthy corals ranged between 0.5 and 0.75 [65]. After the physiological measurements, corals were rinsed 6 times with filtered seawater to remove cells that potentially adhered to the coral surface [20], transferred to zip-lock bags, and stored at $-20 \text{ } ^\circ\text{C}$ until analysis.

2.7. Statistical Analyses

All tests were performed using R version 4.1.2 within RStudio (Version 2021.09.1+372). The non-parametric Mann–Whitney–Wilcoxon test was first used to test for significant differences in Symbiodiniaceae densities and total chlorophyll content between feeding regimes. A three-way ANOVA was performed to assess the differences in the measured parameters (photosynthetic efficiency, growth rates, Symbiodiniaceae densities, chlorophyll, and protein concentrations). The tank effect was included as a random factor nested in the feeding regime, with temperature treatment and feeding regime as a fixed factor. The tank effect was eliminated from the ANOVA analyses because it was not significant for all measured parameters. Therefore, data between the tanks were pooled and two-way ANOVAs were used to test the effects of the feeding regime and temperature on symbiont density, chlorophyll concentration, growth rate, photosynthetic efficiency (F_v / F_m), and electron transport rate (ETR). When the ANOVA determined a significant difference, a Tukey's honest significant difference test (HSD) was used to test for pairwise differences while taking the interactions between the different variables into account. The *ggplot2* package [66] was used to create the box plot figures. Throughout the manuscript, values given are expressed as mean \pm SE. Statistical significance was accepted at $p < 0.05$.

3. Results

3.1. Symbiodiniaceae Density, Chlorophyll, and Protein Concentrations

All the coral fragments remained healthy during the 55 days prior to heat stress. At T_0 and T_{65} of incubation under ambient conditions, coral fragments had similar symbiont densities ($8.78 \pm 1.06 \times 10^5 \text{ cell cm}^{-2}$; $N = 6$ for each tank) regardless of the feeding regime (Figure 2A,B). At T_0 , total chlorophyll concentrations ($a + c_2$) were similar ($7.109 \pm 0.705 \text{ } \mu\text{g cm}^{-2}$; $N = 6$ for each tank) for unfed and PD-fed corals ($7.109 \pm 0.705 \text{ } \mu\text{g cm}^{-2}$; $N = 6$ for each tank) (Figure 3A,D); however, at T_{65} in ambient conditions, these concentrations decreased by 33% in unfed corals and conversely increased by 37% in PD-fed corals (two-way ANOVA, Tukey's HSD test, *ajd.* p -value = 0.01) (Figure 3B,E).

Heat-unfed colonies died 8 days after the heat stress, which did not allow us to measure the related symbiont densities and chlorophyll concentrations. In contrast, heat-PD-fed corals remained alive after 10 days of heat stress with their symbiont densities and total chlorophyll concentrations decreased by 88% ($0.925 \times 10^6 \text{ cell cm}^{-2}$) and by 90% ($0.973 \text{ } \mu\text{g cm}^{-2}$), respectively (Figures 2C and 3C,F).




Figure 2. Symbiodiniaceae density (cell cm⁻²) in unfed and PD-fed colonies of *Stylophora pistillata* at the beginning of the experiment (T₀) (A) and at 65 days of the experiment under (B) ambient and (C) heat stress conditions (N = 6; mean ± SE for each condition).




Figure 3. Chlorophyll *a* (A–C) and *c*₂ (D–F) concentrations (µg cm⁻²) in unfed and PD-fed colonies of *Stylophora pistillata* at the beginning of the experiment (T₀) and at 65 days of the experiment under ambient and heat stress conditions (mean ± SE; N = 6). Asterisks indicated significant differences between treatments (* *p* < 0.05; ** *p* < 0.01).

At T_0 and after 30 days of incubation under ambient conditions, coral fragments showed similar protein concentrations (0.802 ± 0.139 mg prot cm^{-2} and 0.751 ± 0.171 mg prot cm^{-2} , respectively; $N = 12$) regardless of the feeding regime (two-way ANOVA, Tukey's HSD test, *ajd.* p -value = 1; $N = 6$ for each feeding regime). As specified above, heat-unfed coral colonies died 8 days after the heat stress, leaving us no time to measure protein concentrations.

3.2. Growth Rates

At T_{30} under ambient conditions and 25 days prior to heat stress, PD-fed corals showed higher growth rates than unfed corals (5.2732 ± 0.2787 $\text{mg g}^{-1} \text{d}^{-1}$ and 3.4362 ± 0.2745 $\text{mg g}^{-1} \text{d}^{-1}$, respectively, two-way ANOVA, Tukey's HSD test, *ajd.* p -value = 0.02). At T_{65} under ambient conditions, PD-fed coral growth rates were 48% higher than unfed corals (two-way ANOVA, Tukey's HSD test, *ajd.* p -value = 0.0002). At T_{65} , heat-PD-fed and PD-fed growth rates remained similar. Heat-PD-fed coral growth rates were still higher than those of heat-unfed corals, before the death of the latter 8 days after heat stress (two-way ANOVA, Tukey's HSD test; *ajd.* p -value = 0.03) (Figure 4).




Figure 4. Growth rates of unfed ($N = 40$) and PD-fed ($N = 44$) colonies of *Stylophora pistillata* exposed for 55 days to ambient conditions and 10 days to heat stress conditions (mean \pm SE). Asterisks indicated significant differences between treatments (* $p < 0.05$; *** $p < 0.001$).

3.3. F_v/F_m and Electron Transport Rates of Symbiodiniaceae

At the beginning of the experiment, corals had similar F_v/F_m (0.696 ± 0.016). At T_{65} in ambient conditions, the F_v/F_m values remained unchanged regardless of the feeding regime (0.697 ± 0.014 for unfed corals; 0.689 ± 0.024 for PD-fed corals). However, F_v/F_m

values of all coral colonies were impacted by the heat stress and significantly decreased by 29.5% and 28.3% for unfed and PD-fed corals, respectively.

At T_{65} in ambient conditions, ETR_{max} values were 1.15-fold higher in PD-fed corals compared to those of the unfed ones. After the heat stress, a decrease in ETR curves was observed in all coral colonies regardless of the feeding regime, but ETR_{max} values decreased more for heat-unfed corals (23%) than for the heat-PD-fed corals (9%) (Figure 5).




Figure 5. Electron transport rate (ETR) versus irradiance (PAR) for unfed and DP-fed colonies of *Stylophora pistillata* at the beginning of the experiment (T_0) and at 65 days of the experiment under ambient and heat stress conditions (mean \pm SE; N = 12).

4. Discussion

Some coral species are able to increase their heterotrophic nutrition during bleaching episodes, which supplies their energy reserves, providing them with greater resistance, improved chances of survival [25,30,34,67–69], and allows them to recover their symbionts [70,71]. These conclusions were drawn by focusing primarily on the ingestion of mesozooplankton, and no study had so far investigated smaller planktonic fractions. Our experiment provides the first evidence that heat-stressed PD-fed corals are able to retain part of their symbionts and chlorophyll and appear more resistant compared to heat-stressed unfed ones, which died after only 8 days. We also show that a supply in planktonic diazotrophs (particularly rich in N) also alleviates some of the negative effects of heat stress on coral growth and photosynthetic efficiency.

In ambient conditions, feeding on planktonic diazotrophs induced significant changes in most coral physiological parameters. After 9 weeks of a planktonic diazotrophs' diet, colonies significantly increased their chlorophyll contents. Corals benefit from these planktonic diazotrophs, directly either by grazing them [36,50], and/or indirectly by uptaking the DIN (dissolved inorganic nitrogen: NH_4^+ and DON) they released in surrounding waters [49,72,73]. This stimulation of the symbiotic compartment has been highlighted before for mesozooplankton-fed corals (e.g., [20,74–76]). As heterotrophy results in a significant release of NH_4^+ and as this NH_4^+ accumulates 14 to 23 times faster in the Symbiodiniaceae than in animal cells [77–79], planktonic diazotrophs' diet stimulated pigment synthesis and symbiont division. The richness of planktonic diazotroph cells in vitamin B12 [80], which is an essential nutrient for the coral holobiont [81], has also contributed to this stimulation of the symbiotic compartment. In our study, ETR_{max} and F_v/F_m values of fed corals were similar to those of the unfed ones, and not higher as it has been demonstrated in mesozooplankton-fed corals (e.g., [22,76]). The PD-fed corals having higher symbiont densities should result in higher photosynthetic rates per surface regardless [21,22,71,82]. Given the N richness of diazotrophic cells, this specific diet should modify the photosynthate quality, with the highest amounts of N-rich amino acids [83].

One of the major results of this study is that an exclusive nutritional diet of planktonic diazotrophs stimulated coral growth by 48% compared to the unfed corals. Our data confirm that feeding enhances skeletal growth, suggesting that corals allocate a high proportion of the energy brought by planktonic diazotrophs to skeleton growth processes [75,84–86]. Skeleton growth is a dual process, involving the secretion of an organic matrix and the deposition of a CaCO₃ fraction. The contribution of diazotrophic plankton, particularly rich in N, could thus participate in a stimulation of the production of this organic matrix, mainly composed of proteins, glycoproteins, and polysaccharides [87–89], as already demonstrated in mesozooplankton-fed corals [75]. This growth stimulation by planktonic diazotrophs could also be achieved indirectly by boosting photosynthesis as stated above. These two stimulating ways make that growth rates in fed corals, submitted to heat stress, remain stable and always higher than those of unfed ones in the same conditions. In [71], calcification rates of *S. pistillata* colonies fed for 9 weeks with *Artemia salina* were reduced by up to 65% compared to control temperatures. Unlike *A. salina* nauplii, planktonic diazotrophs release large amounts of NH₄⁺ and DON into the surrounding water [49,72,73]. This surplus of N provided by the planktonic diazotrophs might have reduced calcification sensitivity to heat stress, as demonstrated for corals submitted to moderate nutrient inputs [90,91].

In corals under heat stress, the supply in planktonic diazotrophs helped maintain a minimal chlorophyll concentration and symbiont density to prevent a large decrease in the RLC curves, and above all, the colonies' death. A decrease in chlorophyll content and symbiont densities was observed in heat-PD-fed corals (decrease of respectively 85% and 88%) but contrary to heat-unfed corals, all heat-PD-fed corals stayed alive, managed to keep some of their symbionts and pigments, and grew faster. As in the studies testing the effects of the ingestion of *Artemia salina* nauplii on the same species (*S. pistillata*; [21,22,92]), planktonic diazotrophs helped maintain photosynthetic efficiency of PSII. The input of this new N might have (i) facilitated the protein repair and re-synthesis of the PSII D1 protein (reviewed by [92,93]), (ii) increased the photosynthetic and photoprotective pigment (such as xanthophyll and peridinin) contents in coral tissue as in corals submitted to a moderate DIN enrichment [91], and/or (iii) reduced the degradation of the symbionts by increasing the synthesis of antioxidant compounds or heat-shock proteins [92,94].

Our results clearly show a positive effect of planktonic diazotroph feeding under heat stress, sustaining photosynthetic efficiency and growth of *S. pistillata* colonies. So far, heterotrophic feeding on planktonic diazotrophs has been little considered compared to the N supply provided by coral-associated diazotrophs. However, in the case of heat stress, Ref. [95] have recently shown that this N supply by coral-associated diazotrophs is compromised. A 10-day heat stress caused a change in coral-associated diazotroph communities, which led to a 30% decrease in fixed N. Conversely, our previous works highlighted that N supply by planktonic diazotrophs (i) can fulfill a large part of the coral N requirements, compared to the symbiotic fixation ([50,69,96,97]) and (ii) is significant for several coral species [69]. This last study finally demonstrates that, contrary to endosymbiotic diazotrophs, a supply of planktonic diazotrophs allows corals to be more resistant to bleaching. It is therefore particularly essential to consider the contribution of DDN by planktonic diazotrophs in the coral holobiont N cycling which appears to be one of the main strategies for coral recovery facing bleaching. In the context of climate change, marine heat waves are becoming more intense and frequent and coral reefs are among the most vulnerable ecosystems to this emerging threat [98,99]. Our very encouraging results, obtained in this study, imply that coral reefs where planktonic diazotrophs are abundant (e.g., Papua New Guinea, New Caledonia) could be more resistant to climate change. A supply of planktonic diazotrophs on the reefs could be included among innovative management approaches to improve the resilience of coral reefs with high conservation potential, during and after bleaching episodes [100,101].

Author Contributions: Conceptualization, V.M., F.H., R.R.-M. and S.B.; methodological design, V.M., F.H., R.R.-M., J.D., S.B., M.B. and M.C.; laboratory analyses, V.M. and F.H.; statistical analyses, V.M.; writing—original draft preparation, V.M. and F.H.; writing—review and editing, V.M., F.H., R.R.-M., S.B. and M.B.; supervision, F.H. and S.B. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded to F.H. by LabEx-Corail (FLAMENCO project). V.M. was the beneficiary of a PhD grant from LabEx-Corail (MACADAM project). This work was also funded by the EC2CO/BIOEFFECT program (TOUCAN project).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in the study are included in the article, further inquiries can be directed to the corresponding author/s.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. LaJeunesse, C.T.; Parkinson, J.; Barielson, P.; Jeong, H.; Reimer, J.; Voolstra, C.R.; Santos, S. Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr. Biol.* **2018**, *28*, 2570–2580. [[CrossRef](#)]
2. Muscatine, L.; Porter, J.W. Reef Corals: Mutualistic Symbioses Adapted to Nutrient-Poor Environments. *Bioscience* **1977**, *27*, 454–460. [[CrossRef](#)]
3. Hughes, T.; Kerry, J.; Álvarez-Noriega, M.; Álvarez-Romero, J.; Anderson, K.; Baird, A.; Babcock, R.; Bejer, M.; Bellwood, D.; Berkelmans, R.; et al. Global warming and recurrent mass bleaching of corals. *Nature* **2017**, *543*, 373–377. [[CrossRef](#)]
4. Muscatine, L. The role of symbiotic algae in carbon and energy flux in reef corals. In *Ecosystems of the World Coral Reefs*; Dubinsky, Z., Ed.; Elsevier: Amsterdam, The Netherlands, 1990; pp. 75–87.
5. Lesser, M.P.; Mazel, C.H.; Gorbunov, M.Y.; Falkowski, P.G. Discovery of symbiotic nitrogen-fixing cyanobacteria in corals. *Science* **2004**, *305*, 997–1000. [[CrossRef](#)] [[PubMed](#)]
6. Fisk, D.D.A.; Done, T.T.J. Taxonomic and bathymetric patterns of bleaching in corals, Myrmdon reef. In Proceedings of the 5th International Coral Reef Congress, Tahiti, France, 27 May–1 June 1985.
7. Grottoli, G.; Rodrigues, L.J.; Juarez, C. Lipids and stable carbon isotopes in two species of Hawaiian corals, *Porites compressa* and *Montipora verrucosa*, following a bleaching event. *Mar. Biol.* **2004**, *145*, 621–631. [[CrossRef](#)]
8. Marshall, P.A.; Baird, A.H. Bleaching of corals on the Great Barrier Reef: Differential susceptibilities among taxa. *Coral Reefs* **2000**, *19*, 155–163. [[CrossRef](#)]
9. Stimson, J.; Sakai, K.; Sembali, H. Interspecific comparison of the symbiotic relationship in corals with high and low rates of bleaching-induced mortality. *Coral Reefs* **2002**, *21*, 409–421. [[CrossRef](#)]
10. Wilkinson, C.; Hodgson, G. Coral reefs and the 1997–1998 mass bleaching and mortality. *Nat. Resour.* **1999**, *35*, 16–25.
11. Loya, Y.; Sakai, K.; Yamazato, K.; Nakano, Y.; Sambali, H.; van Woesik, R. Coral bleaching: The winners and the losers. *Ecol. Lett.* **2001**, *4*, 122–131. [[CrossRef](#)]
12. Fitt, W.K.; Gates, R.D.; Hoegh-Guldberg, O.; Bythell, J.C.; Jatkar, A.; Grottoli, A.G.; Gomez, M.; Fisher, P.; Lajuenesse, T.C.; Pantos, O.; et al. Response of two species of Indo-Pacific corals, *Porites cylindrica* and *Stylophora pistillata*, to short-term thermal stress: The host does matter in determining the tolerance of corals to bleaching. *J. Exp. Mar. Biol. Ecol.* **2009**, *373*, 102–110. [[CrossRef](#)]
13. Dunn, S.R.; Pernice, M.; Green, K.; Hoegh-Guldberg, O.; Dove, S.G. Thermal stress promotes host mitochondrial degradation in symbiotic cnidarians: Are the batteries of the reef going to run out? *PLoS ONE* **2012**, *7*, e39024. [[CrossRef](#)] [[PubMed](#)]
14. Glynn, P.W.; Maté, J.L.; Baker, A.C.; Calderón, M.O. Coral bleaching and mortality in Panama and Ecuador during the 1997–1998 El Niño–Southern Oscillation event: Spatial/temporal patterns and comparisons with the 1982–1983 event. *Bull. Mar. Sci.* **2001**, *69*, 79–109.
15. Howells, E.J.; Beltran, V.H.; Larsen, N.W.; Bay, L.K.; Willis, B.L.; van Oppen, M.J.H. Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat. Clim. Chang.* **2012**, *2*, 116–120. [[CrossRef](#)]
16. Warner, M.E.; Fitt, W.K.; Schmidt, G.W. Damage to photosystem II in symbiotic dinoflagellates: A determinant of coral bleaching. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 8007–8012. [[CrossRef](#)]
17. Houlbrèque, F.; Tambutté, E.; Richard, C.; Ferrier-pagès, C. Importance of a micro-diet for scleractinian corals. *Mar. Ecol. Prog. Ser.* **2004**, *282*, 151–160. [[CrossRef](#)]
18. Houlbrèque, F.; Ferrier-Pagès, C. Heterotrophy in Tropical Scleractinian Corals. *Biol. Rev.* **2009**, *84*, 1–17. [[CrossRef](#)] [[PubMed](#)]
19. Goldberg, W.M. Coral Food, Feeding, Nutrition, and Secretion: A Review. *Mar. Org. Model Syst. Biol. Med.* **2018**, *65*, 377–421.
20. Houlbrèque, F.; Tambutté, E.; Ferrier-Pagès, C. Effect of zooplankton availability on the rates of photosynthesis, and tissue and skeletal growth in the scleractinian coral *Stylophora pistillata*. *J. Exp. Mar. Biol. Ecol.* **2003**, *296*, 145–166. [[CrossRef](#)]
21. Borell, E.M.; Yuliantri, A.R.; Bischof, K.; Richter, C. The effect of heterotrophy on photosynthesis and tissue composition of two scleractinian corals under elevated temperature. *J. Exp. Mar. Biol. Ecol.* **2008**, *364*, 116–123. [[CrossRef](#)]

22. Ferrier-Pagès, C.; Rottier, C.; Beraud, E.; Levy, O. Experimental assessment of the feeding effort of three scleractinian coral species during a thermal stress: Effect on the rates of photosynthesis. *J. Exp. Mar. Biol. Ecol.* **2010**, *390*, 118–124. [[CrossRef](#)]
23. Ferrier-Pagès, C.; Peiranoet, A.; Abbate, M.; Cocito, S.; Negri, A.; Rottier, C.; Riera, P.; Rodolfo-Metalpaal, R.; Reynaud, S. Summer autotrophy and winter heterotrophy in the temperate symbiotic coral *Cladocora caespitosa*. *Limnol. Oceanogr.* **2011**, *56*, 1429–1438. [[CrossRef](#)]
24. Treignier, C.; Grover, R.; Ferrier-Pagès, C.; Tolosa, I. Effect of light and feeding on the fatty acid and sterol composition of zooxanthellae and host tissue isolated from the scleractinian coral *Turbinaria reniformis*. *Limnol. Oceanogr.* **2008**, *53*, 2702–2710. [[CrossRef](#)]
25. Anthony, K.R.N.; Hoogenboom, M.O.; Maynard, J.A.; Grottoli, A.G.; Middlebrook, R. Energetics approach to predicting mortality risk from environmental stress: A case study of coral bleaching. *Funct. Ecol.* **2009**, *23*, 539–550. [[CrossRef](#)]
26. Cox, E.F. Continuation of sexual reproduction in *Montipora capitata* following bleaching. *Coral Reefs* **2007**, *26*, 721–724. [[CrossRef](#)]
27. Edmunds, P.J. Zooplanktivory ameliorates the effects of ocean acidification on the reef coral *Porites* spp. *Limnol. Oceanogr.* **2011**, *56*, 2402–2410. [[CrossRef](#)]
28. Houlbrèque, F.; Reynaud, S.; Godinot, C.; Oberhänsli, F.; Rodolfo-Metalpa, R.; Ferrier-Pagès, C. Ocean acidification reduces feeding rates in the scleractinian coral *Stylophora pistillata*. *Limnol. Oceanogr.* **2015**, *60*, 89–99. [[CrossRef](#)]
29. Drenkard, E.J.; Cohen, A.L.; McCorkle, D.C.; de Putron, S.J.; Starczak, V.R.; Zicht, A.E. Calcification by juvenile corals under heterotrophy and elevated CO₂. *Coral Reefs* **2013**, *32*, 727–735. [[CrossRef](#)]
30. Grottoli, G.; Rodrigues, L.J.; Palardy, J.E. Heterotrophic plasticity and resilience in bleached corals. *Nature* **2006**, *440*, 1186–1189. [[CrossRef](#)]
31. Bessell-Browne, P.; Stat, M.; Thomson, D.; Clode, P.L. *Coscinaraea marshae* corals that have survived prolonged bleaching exhibit signs of increased heterotrophic feeding. *Coral Reefs* **2014**, *33*, 795–804. [[CrossRef](#)]
32. Palardy, J.E.; Rodrigues, L.J.; Grottoli, A.G. The importance of zooplankton to the daily metabolic carbon requirements of healthy and bleached corals at two depths. *J. Exp. Mar. Biol. Ecol.* **2008**, *367*, 180–188. [[CrossRef](#)]
33. Sangmanee, K.; Casareto, B.; Nguyen, T.D.; Sangsawang, L.; Toyoda, K.; Suzuki, T.; Suzuki, Y. Influence of thermal stress and bleaching on heterotrophic feeding of two scleractinian corals on pico-nanoplankton. *Mar. Pollut. Bull.* **2020**, *158*, 111405. [[CrossRef](#)] [[PubMed](#)]
34. Conti-Jerpe, E.; Thompson, P.; Wong, C.W.M.; Oliveira, N.; Duprey, N.; Moynihan, M.; Baker, D. Trophic strategy and bleaching resistance in reef-building corals. *Sci. Adv.* **2020**, *6*, eaaz5443. [[CrossRef](#)] [[PubMed](#)]
35. Tremblay, P.; Naumann, M.S.; Sikorski, S.; Grover, R.; Ferrier-Pagès, C. Experimental assessment of organic carbon fluxes in the scleractinian coral *Stylophora pistillata* during a thermal and photo stress event. *Mar. Ecol. Prog. Ser.* **2012**, *453*, 63–77. [[CrossRef](#)]
36. Meunier, V.; Bonnet, S.; Pernice, M.; Benavides, M.; Lorrain, A.; Grosso, O.; Lambert, C.; Houlbrèque, F. Bleaching forces coral's heterotrophy on diazotrophs and *Synechococcus*. *ISME J.* **2019**, *13*, 2882–2886. [[CrossRef](#)]
37. Turk-Kubo, K.A.; Frank, I.E.; Hogan, M.E.; Desnues, A.; Bonnet, S.; Zehr, J.P. Diazotroph community succession during the VAHINE mesocosms experiment (New Caledonia Lagoon). *Biogeosciences Discuss.* **2015**, *12*, 9043–9079. [[CrossRef](#)]
38. Messer, F.; Mahaffey, C.; Robinson, C.M.; Jeffries, T.C.; Baker, K.G.; Isaksson, J.B.; Ostrowski, M.; Doblin, M.A.; Brown, M.V.; Seymour, J.R. High levels of heterogeneity in diazotroph diversity and activity within a putative hotspot for marine nitrogen fixation. *ISME J.* **2016**, *10*, 1499–1513. [[CrossRef](#)]
39. Messer, F.; Brown, M.V.; Furnas, M.J.; Carney, R.L.; McKinnon, A.D.; Seymour, J.R. Diversity and activity of diazotrophs in great barrier reef surface waters. *Front. Microbiol.* **2017**, *8*, 967. [[CrossRef](#)]
40. Bonnet, S.; Caffin, M.; Berthelot, H.; Moutin, T. Hot spot of N₂ fixation in the western tropical South Pacific pleads for a spatial decoupling between N₂ fixation and denitrification. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E2800–E2801. [[CrossRef](#)]
41. Luo, Y.-W.; Doney, S.C.; Anderson, L.A.; Benavides, M.; Berman-Frank, I.; Bode, A.; Bonnet, S.; Boström, K.H.; Böttjer, D.; Capone, D.G.; et al. Database of diazotrophs in global ocean: Abundance, biomass and nitrogen fixation rates. *Earth Syst. Sci. Data* **2012**, *4*, 47–73. [[CrossRef](#)]
42. Foster, R.A.; Paytan, A.; Zehr, J.P. Seasonality of N₂ fixation and nifH gene diversity in the Gulf of Aqaba (Red Sea). *Limnol. Oceanogr.* **2009**, *54*, 219–233. [[CrossRef](#)]
43. Rahav, E.; Bar-Zeev, E.; Ohayon, S.; Elifantz, H.; Belkin, N.; Herut, B.; Mulholland, M.; Berman-Frank, I. Dinitrogen fixation in aphotic oxygenated marine environments. *Front. Microbiol.* **2013**, *4*, 227. [[CrossRef](#)] [[PubMed](#)]
44. Saulia, E.; Benavides, M.; Henke, B.; Turk-Kubo, K.; Cooperguard, H.; Grosso, O.; Desnues, A.; Rodier, M.; Dupouy, C.; Riemann, L.; et al. Seasonal Shifts in Diazotrophs Players: Patterns Observed Over a Two-Year Time Series in the New Caledonian Lagoon (Western Tropical South Pacific Ocean). *Front. Mar. Sci.* **2020**, *7*, 915. [[CrossRef](#)]
45. Berthelot, H.; Moutin, T.; L'Helguen, S.; Leblanc, K.; Hélias, S.; Grosso, O.; Leblond, N.; Charrière, B.; Bonnet, S. Dinitrogen fixation and dissolved organic nitrogen fueled primary production and particulate export during the VAHINE mesocosm experiment (New Caledonia lagoon). *Biogeosciences* **2015**, *12*, 4099–4112. [[CrossRef](#)]
46. Rodier, M.; Le Borgne, R. Population and trophic dynamics of *Trichodesmium thiebautii* in the SE lagoon of New Caledonia. Comparison with *T. erythraeum* in the SW lagoon. *Mar. Pollut. Bull.* **2010**, *61*, 349–359. [[CrossRef](#)]
47. Bonnet, S.; Berthelot, H.; Turk-Kubo, K.; Fawcett, S.; Rahav, E.; L'Helguen, S.; Berman-Frank, I. Dynamics of N₂ fixation and fate of diazotroph-derived nitrogen in a low-nutrient, low-chlorophyll ecosystem: Results from the VAHINE mesocosm experiment (New Caledonia). *Biogeosciences* **2016**, *13*, 2653–2673. [[CrossRef](#)]

48. White, E.; Spitz, Y.H.; Karl, D.M.; Letelier, R.M. Flexible elemental stoichiometry in *Trichodesmium* spp. and its ecological implications. *Limnol. Oceanogr.* **2006**, *51*, 1777–1790. [[CrossRef](#)]
49. Berthelot, H.; Bonnet, S.; Camps, M.; Grosso, O.; Moutin, T. Assessment of the dinitrogen released as ammonium and dissolved organic nitrogen by unicellular and filamentous marine diazotrophic cyanobacteria grown in culture. *Front. Mar. Sci.* **2015**, *2*, 80. [[CrossRef](#)]
50. Benavides, M.; Houlbreque, F.; Camps, M.; Lorrain, A.; Grosso, O.; Bonnet, S. Diazotrophs: A non-negligible source of nitrogen for the tropical coral *Stylophora pistillata*. *J. Exp. Biol.* **2016**, *219*, 2608–2612. [[CrossRef](#)]
51. Guillard, R.R.; Ryther, J.H. Studies on marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* (Cleve) Gran. *Can. J. Microbiol.* **1962**, *8*, 229–239. [[CrossRef](#)]
52. Biegala, I.; Raimbault, P. High abundance of diazotrophic picocyanobacteria (<3 μm) in a Southwest Pacific coral lagoon. *Aquat. Microb. Ecol.* **2008**, *51*, 45–53. [[CrossRef](#)]
53. Henke, B.A.; Turk-Kubo, K.A.; Bonnet, S.; Zehr, J.P. Distributions and abundances of sublineages of the N_2 -fixing cyanobacterium *Candidatus Atelocyanobacterium thalassa* (UCYN-A) in the New Caledonian coral lagoon. *Front. Microbiol.* **2018**, *9*, 554. [[CrossRef](#)]
54. Rodolfo-Metalpa, R.; Martin, S.; Ferrier-Pagès, C.; Gattuso, J.-P. Response of the temperate coral *Cladocora caespitosa* to mid- and long-term exposure to pCO_2 and temperature levels projected for the year 2100 AD. *Biogeosciences* **2010**, *7*, 289–300. [[CrossRef](#)]
55. Godinot, C.; Houlbrèque, F.; Grover, R.; Ferrier-Pagès, C. Coral uptake of inorganic phosphorus and nitrogen negatively affected by simultaneous changes in temperature and pH. *PLoS ONE* **2011**, *6*, e25024. [[CrossRef](#)]
56. Jeffrey, S.W.; Humphrey, G.F. New spectrophotometric equations for determining chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. *Biochem. Und Physiol. Der Pflanz.* **1975**, *167*, 191–194. [[CrossRef](#)]
57. Hoogenboom, M.; Beraud, E.; Ferrier-Pagès, C. Relationship between symbiont density and photosynthetic carbon acquisition in the temperate coral *Cladocora caespitosa*. *Coral Reefs* **2010**, *29*, 21–29. [[CrossRef](#)]
58. Stimson, J.; Kinzie, R.A., III. The temporal pattern and rate of release of zooxanthellae from the reef coral *Pocillopora damicornis* (Linnaeus) under nitrogen-enrichment and control conditions. *J. Exp. Mar. Biol. Ecol.* **1991**, *153*, 63–74. [[CrossRef](#)]
59. Naumann, M.S.; Richter, C.; El-Zibdah, M.; Wild, C. Coral mucus as an efficient trap for picoplanktonic cyanobacteria: Implications for pelagic-benthic coupling in the reef ecosystem. *Mar. Ecol. Prog. Ser.* **2009**, *385*, 65–76. [[CrossRef](#)]
60. Smith, K.; Krohn, R.I.; Hermanson, G.T.; Mallia, A.K.; Gartner, F.H.; Provenzano, M.D.; Fujimoto, E.K.; Goeke, N.M.; Olson, B.J.; Klenk, B.C. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **1985**, *150*, 76–85. [[CrossRef](#)]
61. Spencer Davies, P. Short-term growth measurements of corals using an accurate buoyant weighing technique. *Mar. Biol.* **1989**, *101*, 389–395. [[CrossRef](#)]
62. Schreiber, U.; Schliwa, U.; Bilger, W. Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.* **1986**, *10*, 51–62. [[CrossRef](#)]
63. Hoegh-Guldberg, O.; Jones, R.J. Photoinhibition and photoprotection in symbiotic dinoflagellates from reef-building corals. *Mar. Ecol. Prog. Ser.* **1999**, *183*, 73–86. [[CrossRef](#)]
64. Biscéré, T.; Rodolfo-Metalpa, R.; Lorrain, A.; Chauvaud, L.; Thébault, J.; Clavier, J.; Houlbrèque, F. Responses of two scleractinian corals to cobalt pollution and ocean acidification. *PLoS ONE* **2015**, *10*, e0122898. [[CrossRef](#)] [[PubMed](#)]
65. Fitt, W.; Brown, B.; Warner, M.; Dunne, R. Coral bleaching: Interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs* **2001**, *20*, 51–65. [[CrossRef](#)]
66. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer: Berlin/Heidelberg, Germany, 2016.
67. Grottoli, G.; Rodrigues, L.J. Bleached *Porites compressa* and *Montipora capitata* corals catabolize $\delta^{13}\text{C}$ -enriched lipids. *Coral Reefs* **2011**, *30*, 687. [[CrossRef](#)]
68. Baumann, J.; Grottoli, A.G.; Hughes, A.D.; Matsui, Y. Photoautotrophic and heterotrophic carbon in bleached and non-bleached coral lipid acquisition and storage. *J. Exp. Mar. Biol. Ecol.* **2014**, *461*, 469–478. [[CrossRef](#)]
69. Meunier, V.; Bonnet, S.; Benavides, M.; Ravache, A.; Grosso, O.; Lambert, C.; Houlbrèque, F. Diazotroph-Derived Nitrogen Assimilation Strategies Differ by Scleractinian Coral Species. *Front. Mar. Sci.* **2021**, *8*, 1018. [[CrossRef](#)]
70. Connolly, S.R.; Lopez-Yglesias, M.A.; Anthony, K.R.N. Food availability promotes rapid recovery from thermal stress in a scleractinian coral. *Coral Reefs* **2012**, *31*, 951–960. [[CrossRef](#)]
71. Tremblay, P.; Gori, A.; Maguer, J.F.; Hoogenboom, M.; Ferrier-Pagès, C. Heterotrophy promotes the re-establishment of photosynthate translocation in a symbiotic coral after heat stress. *Sci. Rep.* **2016**, *6*, 38112. [[CrossRef](#)] [[PubMed](#)]
72. Benavides, M.; Agawin, N.; Aristegui, J.; Peene, J.; Stal, L. Dissolved organic nitrogen and carbon release by a marine unicellular diazotrophic cyanobacterium. *Aquat. Microb. Ecol.* **2013**, *69*, 69–80. [[CrossRef](#)]
73. Glibert, P.M.; Bronk, D.A. Release of dissolved organic nitrogen by marine diazotrophic cyanobacteria, *Trichodesmium* spp. *Appl. Environ. Microbiol.* **1994**, *60*, 3996–4000. [[CrossRef](#)] [[PubMed](#)]
74. Muscatine, L.; Falkowski, P.G.; Dubinsky, Z.; Cook, P.A.; McCloskey, L.R. The effect of external nutrient resources on the population dynamics of zooxanthellae in a reef coral. *Proc. R. Soc. Lond.* **1989**, *236*, 311–324. [[CrossRef](#)]
75. Houlbrèque, F.; Tambutté, E.; Allemand, D.; Ferrier-Pagès, C. Interactions between zooplankton feeding, photosynthesis and skeletal growth in the scleractinian coral *Stylophora pistillata*. *J. Exp. Biol.* **2004**, *207*, 1461–1469. [[CrossRef](#)] [[PubMed](#)]
76. Lyndby, N.H.; Holm, J.B.; Wangpraseurt, D.; Grover, R.; Rottier, C.; Kühl, M.; Ferrier-Pagès, C. Effect of temperature and feeding on carbon budgets and O_2 dynamics in *Pocillopora damicornis*. *Mar. Ecol. Prog. Ser.* **2020**, *652*, 49–62. [[CrossRef](#)]

77. Grover, R.; Maguer, J.F.; Reynaud-Vaganay, S.; Ferrier-Pagès, C. Uptake of ammonium by the scleractinian coral *Stylophora pistillata*: Effect of feeding, light, and ammonium concentrations. *Limnol. Oceanogr.* **2002**, *47*, 782–790. [[CrossRef](#)]
78. Pernice, M.; Meibom, A.; Heuvel, A.V.D.; Kopp, C.; Domart-Coulon, I.; Hoegh-Guldberg, O.; Dove, S. A single-cell view of ammonium assimilation in coral-dinoflagellate symbiosis. *ISME J.* **2012**, *6*, 1314–1324. [[CrossRef](#)] [[PubMed](#)]
79. Tanaka, Y.; Grottoli, A.G.; Matsui, Y.; Suzuki, A.; Sakai, K. Partitioning of nitrogen sources to algal endosymbionts of corals with long-term ¹⁵N-labelling and a mixing model. *Ecol. Modell.* **2015**, *309–310*, 163–169. [[CrossRef](#)]
80. Bonnet, S.; Webb, E.A.; Panzeca, C.; Karl, D.M.; Capone, D.G.; Wilhelmy, S.A.S. Vitamin B 12 excretion by cultures of the marine cyanobacteria *Crocospaera* and *Synechococcus*. *Limnol. Oceanogr.* **2010**, *55*, 1959–1964. [[CrossRef](#)]
81. Agostini, S.; Suzuki, Y.; Higuchi, T.; Casareto, B.E.; Yoshinaga, K.; Nakano, Y.; Fujimura, H. Biological and chemical characteristics of the coral gastric cavity. *Coral Reefs* **2012**, *31*, 147–156. [[CrossRef](#)]
82. Dubinsky, Z.; Stambler, N.; Ben-Zion, M.; McCloskey, L.R.; Muscatine, L.; Falkowski, P.G. The effect of external nutrient resources on the optical properties and photosynthetic efficiency of *Stylophora pistillata*. *Proc. R. Soc. B Biol. Sci.* **1990**, *239*, 231–246. [[CrossRef](#)]
83. Davy, S.; Cook, C. The relationship between nutritional status and carbon flux in the zooxanthellate sea anemone *Aiptasia pallida*. *Mar. Biol.* **2001**, *139*, 999–1005. [[CrossRef](#)]
84. Ferrier-Pagès, C.; Witting, J.; Tambutté, E.; Sebens, K.P. Effect of natural zooplankton feeding on the tissue and skeletal growth of the scleractinian coral *Stylophora pistillata*. *Coral Reefs* **2003**, *22*, 229–240. [[CrossRef](#)]
85. Jacques, G.; Pilson, M.E.Q. Experimental ecology of the temperate scleractinian coral *Astrangia danae* I. Partition of respiration, photosynthesis and calcification between host and symbionts. *Mar. Biol.* **1980**, *60*, 167–178. [[CrossRef](#)]
86. Witting, J.H. Zooplankton capture and coral growth: The role of heterotrophy in Caribbean reef corals. Ph.D. Thesis, Northeastern University, Boston, MA, USA, 1999.
87. Simkiss, K.; Wilbur, K.M. *Biomineralization: Cell Biology and Mineral Deposition*; Academic Press: San Diego, CA, USA, 1989.
88. Puverel, S.; Tambutté, E.; Pereira-Mouriès, L.; Zoccola, D.; Allemand, D.; Tambutté, S. Soluble organic matrix of two Scleractinian corals: Partial and comparative analysis. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2005**, *141*, 480–487. [[CrossRef](#)]
89. Puverel, S.; Houllbrèque, F.; Tambutté, E.; Zoccola, D.; Payan, P.; Caminiti, N.; Tambutté, S.; Allemand, D. Evidence of low molecular weight components in the organic matrix of the reef building coral, *Stylophora pistillata*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **2007**, *147*, 850–856. [[CrossRef](#)]
90. Schoepf, V.; Grottoli, A.G.; Warner, M.E.; Cai, W.-J.; Melman, T.F.; Hoadley, K.D.; Pettay, D.T.; Hu, X.P.; Li, Q.; Xu, H.; et al. Coral Energy Reserves and Calcification in a High-CO₂ World at Two Temperatures. *PLoS ONE* **2013**, *8*, e75049. [[CrossRef](#)] [[PubMed](#)]
91. Béraud, E.; Gevaert, F.; Rottier, C.; Ferrier-Pagès, C. The response of the scleractinian coral *Turbinaria reniformis* to thermal stress depends on the nitrogen status of the coral holobiont. *J. Exp. Biol.* **2013**, *216*, 2665–2674. [[CrossRef](#)]
92. Borell, E.M.; Bischof, K. Feeding sustains photosynthetic quantum yield of a scleractinian coral during thermal stress. *Oecologia* **2008**, *157*, 593–601. [[CrossRef](#)] [[PubMed](#)]
93. Smith, D.J.; Suggett, D.J.; Baker, N.R. Is photoinhibition of zooxanthellae photosynthesis the primary cause of thermal bleaching in corals? *Glob. Chang. Biol.* **2005**, *11*, 1–11. [[CrossRef](#)]
94. Yakovleva, I.; Bhagooli, R.; Takemura, A.; Hidaka, M. Differential susceptibility to oxidative stress of two scleractinian corals: Antioxidant functioning of mycosporine-glycine. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2004**, *139*, 721–730. [[CrossRef](#)]
95. Rådecker, N.; Pogoreutz, C.; Gegner, H.M.; Cardenas, A.; Perna, G.; Geissler, L.; Roth, F.; Bougoure, J.; Guagliardo, P.; Struck, U.; et al. Heat stress reduces the contribution of diazotrophs to coral holobiont nitrogen cycling. *ISME J.* **2021**. [[CrossRef](#)]
96. Benavides, M.; Bednarz, V.N.; Ferrier-Pagès, C. Diazotrophs: Overlooked key players within the coral symbiosis and tropical reef ecosystems? *Front. Mar. Sci.* **2017**, *4*. [[CrossRef](#)]
97. Meunier, V.; Geissler, L.; Bonnet, S.; Rådecker, N.; Perna, G.; Grosso, O.; Lambert, C.; Rodolfo-Metalpa, R.; Voolstra, C.R.; Houllbrèque, F. Microbes support enhanced nitrogen requirements of coral holobionts in a high CO₂ environment. *Mol. Ecol.* **2021**, *30*, 5888–5899. [[CrossRef](#)] [[PubMed](#)]
98. Bindoff, L.N.; Cheung, W.W.; Kairo, G.J.; Aristegui, J.; Guinder, A.V.; Hallberg, R. Changing Ocean, marine ecosystems, and dependent communities. In *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*; Pörtner, H.O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegria, A., Nicolai, M., Okem, A., et al., Eds.; IPCC: Geneva, Switzerland, 2019.
99. Hughes, T.P.; Anderson, K.D.; Connolly, S.R.; Heron, S.F.; Kerry, J.T.; Baird, A.H.; Baum, J.K.; Lough, J.M.; Berumen, M.L.; Bridge, T.C.; et al. Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* **2018**, *359*, 80–83. [[CrossRef](#)] [[PubMed](#)]
100. Hoey, A.; Howells, E.; Johansen, J.; Hobbs, J.; Messmer, V.; McCowan, D.; Wilson, S.; Pratchett, M. Recent advances in understanding the effects of climate change on coral reefs. *Diversity* **2016**, *8*, 12. [[CrossRef](#)]
101. Jin, K.; Lundgren, P.; Lutz, A.; Raina, J.-B.; Howells, E.J.; Paley, A.S.; Willis, B.L.; van Oppen, M.J.H. Genetic markers for antioxidant capacity in a reef-building coral. *Sci. Adv.* **2016**, *2*, e1500842. [[CrossRef](#)] [[PubMed](#)]