



**HAL**  
open science

## Mangrove zonation mapping in West Africa, at 10-m resolution, optimized for inter-annual monitoring

Florent Lombard, Safietou Soumaré, Julien Andrieu, Didier Josselin

### ► To cite this version:

Florent Lombard, Safietou Soumaré, Julien Andrieu, Didier Josselin. Mangrove zonation mapping in West Africa, at 10-m resolution, optimized for inter-annual monitoring. *Ecological Informatics*, 2023, 75, pp.102027. 10.1016/j.ecoinf.2023.102027 . hal-03993578

**HAL Id: hal-03993578**

**<https://hal.science/hal-03993578v1>**

Submitted on 4 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1            **Mangrove zonation mapping in West Africa, at 10-meter**  
2            **resolution, optimized for inter-annual monitoring**

Florent Lombard<sup>a</sup>, Safietou Soumaré<sup>a,b</sup>, Julien Andrieu<sup>a,c</sup>, Didier Josselin<sup>d</sup>

*<sup>a</sup> Department of Geography, Université Côte d'Azur, CNRS UMR Espace Campus Valrose, Bâtiment L 28, avenue de Valrose 06108 Nice Cedex 2 France*

*<sup>b</sup> Laboratoire des Sciences et Techniques de l'Eau et de l'Environnement (LaSTEE) - Ecole Polytechnique de Thiès, BP A10 THIÈS, Senegal*

*<sup>c</sup> Department of Geomatics, French Institute of Pondicherry (IFP) UMIFRE 21 CNRS-MEAE / UAR 3330 11, St. Louis Street, P.B. 33 Pondicherry, 605001, India*

*<sup>d</sup> CNRS UMR Espace, 74 rue Louis Pasteur, 84029 Avignon CEDEX, France*

[florent.lombard@univ-cotedazur.fr](mailto:florent.lombard@univ-cotedazur.fr)

3

## **Mangrove zonation mapping in West Africa, at 10-meter resolution, optimized for inter-annual monitoring**

The surface area of mangrove ecosystems in Senegal have fluctuated substantially over several decades. Satellite data at 10 to 30-meter resolution, which has been available since the 1980s, has allowed the mapping and quantification of these dynamics. However, the plant formations have reorganized internally; this has not been well-documented, possibly because there is no established method for detecting zonation of the Senegalese mangrove. This paper proposes a two-step method for mapping the zonation of the Saloum Delta mangrove. First, mangrove surfaces were detected using machine learning methods from an object-based time series. Finally, a typology was developed through object-based clustering using time-series metrics derived from the harmonic regression modelling of the vegetation fraction. A comparison with field data allowed us to determine the number of classes and discriminating variables. The results showed that the selected method resulted in an overall accuracy of 97.55% (Kappa = 95.42) for the land cover at 4 classes (Water, Mangrove, Salt flats, Other land covers), with an F1-Score of 98.91% for the mangroves. Second, our results suggest that the annual trend of the vegetation fraction at the object scale is effective in differentiating mangrove zonation into three classes based on canopy density and stand height (HM: High mangrove; LDM: Low and dense mangrove; LOM: Low and open mangrove). Finally, the temporal stability of the classes and uncertainty around the magnitudes of the plant fraction values per class were assessed by Bayesian inference. An overall accuracy of 85.5% can be expected to identify the zonation typology on an inter-annual scale. This mapping technique can be used to characterize the rate of change in zonation in response to environmental changes and to guide management strategies.

29 **Key policy highlights**

30

31 • There is currently no method for detecting the zonation of the Senegalese  
32 mangrove.

33

34 • A typology of mangrove zonation was created based on canopy density and  
35 stand height.

36

37 • Analyzed temporal stability/uncertainty by time series synthesis & Bayesian  
38 modeling.

39

40

41 • Accurately identified typology (85.5% accuracy) in annual monitoring context

42

43 • Reduced uncertainty margins for monitoring using temporal segmentation  
44 algorithms.

45

46

47

48 **Keywords:** Mangrove; Sentinel-2; Sentinel-1; Times series; Machine Learning; OBIA

49

50

51

52

53

## 54 **1. Introduction**

55 Mangrove mapping is an important aspect of ecosystem management strategies (Giri et al.,  
56 2011). The availability of remotely sensed megadata at decametric spatial resolution allows the  
57 classification of different surface states over large continental areas (Mutanga and Kumar,  
58 2019), especially at a scale that can guide local and national environmental policies. Access to  
59 and processing of Landsat and Sentinel archives have been facilitated by freely available cloud  
60 platforms, such as the Google Earth Engine (GEE) (Alonso et al., 2016; Gorelick et al., 2017;  
61 Wang et al., 2019). Furthermore, the computational power allocated by these platforms  
62 provides an opportunity for tropical countries to set up complex processing chains with big  
63 data (Midekisa et al., 2017). However, their dimensionality remains an important issue with  
64 regards to reducing computation times, which decreases the redundancy of information from  
65 different sensors (Myint et al., 2011; Hu et al., 2019; Stromann et al., 2020) and ensures  
66 classification efficiency (Speiser et al., 2019). Moreover, the Sahelo-Sudanese coastal zone  
67 contains socio-ecosystems that generate a variety of landscapes (Temudo et al., 2015; Andrieu,  
68 2020) that can be difficult to map (Cabral and Costa, 2017; Asenso Barnieh et al., 2020) and  
69 which can cause spectral confusion between the mangrove and dense forests on dry land  
70 depending on the season. Therefore, extracting the most effective data and variables is essential  
71 (Schulz et al., 2021).

72 Mangrove forests cover large areas (Cabral and Costa, 2017; Andrieu, 2018) and provide  
73 various ecosystem services (Cormier-Salem, 1999; Walters et al., 2008; Mukherjee et al., 2014;  
74 Arumugam et al., 2021). Global issues of climate change mitigation through carbon  
75 sequestration and preservation of mangrove biodiversity must be considered alongside local  
76 issues such as access to resources (Cormier-Salem et Panfili, 2016). Monitoring mangrove  
77 dynamics on a regular basis could help better reconcile resource management within changing  
78 environments.

79 Mapping mangrove ecosystems in West Africa is crucial for developing effective  
80 ecosystem management strategies (Jia et al., 2018). With the help of high resolution satellite  
81 data from Landsat and Sentinel, as well as access to cloud platforms such as Google Earth  
82 Engine (GEE) (Alonso et al., 2016; Gorelick et al., 2017; Wang et al., 2019), it is now possible  
83 to use powerful computing resources and advanced processing methods to better understand  
84 the dynamics of mangroves in this region (Midekisa et al., 2017). These methods allow for the  
85 management of long and dense time series data, which can provide a comprehensive view of  
86 mangrove ecosystems. The mapping of mangroves in West Africa is particularly valuable due  
87 to the numerous ecosystem services they provide (Cormier-Salem, 1999; Walters et al., 2008;  
88 Mukherjee et al., 2014; Arumugam et al., 2021).

89 Although databases of mangrove areas exist (GWM, 2010; USGS, 2014), they are  
90 highly inaccurate, particularly in West Africa due to persistent cloud cover (Bunting et al.,  
91 2018). The authors also mention that fringing and fragmented mangroves are sometimes  
92 omitted, even though they are frequently encountered in West-Africa (Liu et al., 2021).  
93 Mapping the extent of mangroves has been the subject of many studies (Kuenzer et al., 2011;  
94 Wang et al., 2019), and various mapping methods have been developed to produce local data.  
95 SPOT data has successfully mapped the extent of mangroves in West Africa over several years  
96 with an object-based image analysis (OBIA) approach (Conchedda et al., 2008). Landsat data  
97 has also led to the production of accurate maps using a pixel approach combined with linear  
98 spectral unmixing (Han et al., 2018). These data are also advantageous due to the long time

99 series. Cloud cover has often been mentioned as an impediment to mapping mangroves in the  
100 tropics with passive sensors (Kuenzer et al., 2011); active sensors are therefore of interest for  
101 mapping mangrove surfaces. These data also present good precision scores, especially with  
102 OBIA approaches (Flores de Santiago et al., 2013), and most of these methods are fairly  
103 accurate, particularly with Sentinel-2 optical data (Mondal et al., 2019) and Sentinel-1 and 2  
104 radar-optical combinations (Liu et al., 2021). These data have the advantage of being freely  
105 accessible and easily available on GEE. Machine learning approaches, which typically combine  
106 a large volume of data to avoid the biases of single-date image-based mapping (Cardenas et  
107 al., 2017), are currently favored for map development (Chen et al., 2017; Mondal et al., 2019;  
108 Liu et al., 2021; Wang et al., 2018). Several studies have highlighted the contributions of cloud  
109 solutions, such as GEE, for processing time series and multi-source data (Cardenas et al., 2017;  
110 Chen et al., 2017; Liu et al., 2021; Xiao et al., 2021). Combining optical (Landat-8 and  
111 Sentinel-2) and radar (Sentinel-1) data was found to improve the discrimination of Land Use  
112 Land Cover and mangrove classes (Clerici et al., 2017; Zhao and Qin, 2020; Xiao et al., 2021).  
113 Decision trees, such as Classification And Regression Trees (CART) or Random Forests (RF)  
114 are commonly used for mangroves (Giri et al., 2015; Zhang et al., 2017; Mondal et al., 2019;  
115 Thomas et al., 2018; Chen et al., 2020; Xiao et al., 2021). Liu et al. (2021) combined three  
116 algorithms with a gradient boosting machine (GBM) and neural network (NN) to form a  
117 powerful machine learning package. They achieved accuracy scores above 95% using S2 and  
118 S1 images in West Africa, and developed the most accurate mapping for this area to date.

119 Although the available mapping of mangrove areas seems reliable, differentiating  
120 between the mangrove plant formations using optical and radar data is challenging in West  
121 Africa (Flores de Santiago et al., 2013). Indeed, these can be characterized by the coverage and  
122 stand height, and by estimating the specific species compositions. Thus, plant formations  
123 roughly represent the zonation of mangroves and the resultant spatial organization of  
124 population dynamics at the local scale.

125 Subdivision of mangroves into several facies at a decametric resolution is a complex  
126 process; although it has been performed in other mangrove areas, it has rarely been attempted  
127 in Senegal. Various approaches have been developed to subdivide mangroves. The OBIA  
128 method consistently achieves good scores (Wang et al., 2019). Myint et al. (2008) mapped  
129 mangroves at a specific level using Landsat images with an overall accuracy of 94.2% and a  
130 kappa coefficient of 0.91; the pixel classifiers had an overall accuracy of 62.8% and a kappa  
131 coefficient of 0.57. Wang et al. (2018) achieved an overall accuracy of 70.95% for species  
132 community discrimination using S2 images with a combination of the object approach and an  
133 Random Forest (RF) classifier. In addition, approaches using pixel-level mangrove phenology  
134 have been used to classify mangroves at specific levels (Bullock et al., 2017; Valderrama-  
135 Landeros et al., 2021). Vegetation indices from dense S2 time series also made it possible to  
136 classify mangroves according to a three-class typology (*Rhizophora mangle* fringe, basin, and  
137 *Avicennia germinans* shrub) with overall accuracy scores between 92 and 95% (Vizcaya-  
138 Martínez et al. 2022). The authors were able to map the evolution of this typology after  
139 disturbances and measured species recovery, which are similar to those of the mangroves in  
140 Senegal. Finally, linear spectral unmixing consistently shows good results for quantifying  
141 mangrove canopy cover (Monsef and Smith, 2017; Bullock et al., 2020; Lymburner et al.,  
142 2020) and for establishing a forest composition typology (Gudex-Cross et al., 2017). For  
143 example, linear spectral unmixing was used to establish a 3-class typology in Australia  
144 (woodland, open forest, closed forest) based on the mangrove cover rate from Global Mangrove

145 Watch (GMW) data (Lymburner et al., 2020). Finally, several studies have used RADAR data  
146 to map mangrove species (Flores de Santiago et al., 2013; Ghazali et Wikantika, 2021).  
147 Although RADAR data has shown good accuracy in discriminating mangrove, the species  
148 classification remains more delicate and obtains low (OA = 26%) to moderate (OA ≈ 65%)  
149 scores. The best scores were obtained by optimizing the scale parameter with an OBIA  
150 approach (Flores de Santiago et al., 2013) or by deriving a dual-polarity RADAR vegetation  
151 index (DpRVI) with Sentinel-1 (Ghazali et Wikantika, 2021).

152 Furthermore, significant fluctuation in mangrove forest area in West Africa has  
153 occurred in recent decades (Andrieu, 2018). Mangroves in Senegal have experienced surface  
154 regeneration since the mid-1990s (Conchedda et al., 2011; Dièye et al., 2013; Andrieu, 2018;  
155 Fent et al., 2019; Lombard and Andrieu, 2021). However, the respective shares of species in  
156 regeneration processes differ according to the hydrosystems (Andrieu et al., 2020; Lombard et  
157 al., 2020). Therefore, a map of plant formation is important to obtain an overview of mangrove  
158 composition and to understand mangrove resilience to environmental changes. Plant  
159 formations can estimate the specific composition of the environment, the density of the cover,  
160 and the size of the individuals making up the group. Moreover, as was pointed out by Wang et  
161 al. (2019), mapping mangrove communities is important for conservation strategies (Jia et al.,  
162 2018). We assume that fine-scale processes participate in the spatial organization of mangrove  
163 zonation. The establishment of propagules of different species is constrained by environmental  
164 variables such as salinity and hydroperiod (Krauss et al., 2008) or sedimentary processes (Balke  
165 et al., 2013). In addition, propagule establishment has been reorganized in the face of recent  
166 environmental fluctuations in Senegal (Lombard et al., 2020). Therefore, our objective is to be  
167 able to map mangrove zonation, leading to an understanding of the arrangement of species  
168 zonation resulting from reproduction and growth, which are processes sensitive to  
169 environmental fluctuations. Therefore, our objective was to develop an optimized and robust  
170 multilevel mapping method for mangrove plant formations in the Sahelo-Sudanese delta.

171

172 By utilizing the knowledge gained from the different approaches mentioned above, we  
173 developed a method for mapping mangroves at two levels. First, mangrove surfaces were  
174 detected by machine learning of objects and corrected by classification based on linear spectral  
175 unmixing. Second, linear spectral unmixing was reapplied to subdivide the mangroves into  
176 several plant formations using the OBIA method and classified based on a 1-year time series.


## 177 **2. Study areas and data**

### 178 **2.1. Study site**

179 The study area (Fig. 1) covers the Saloum Delta hydrosystem. Located in the Sahelo-Sudanian  
180 domain, the north-south gradient is characterized by important inter-annual rainfall variations  
181 (Nicholson, 2013; Descroix et al., 2015); mangroves develop under a rainfall of 560 mm in the  
182 north and 800 mm in the south (Lombard et al., 2020). The forest comprises six species  
183 (*Avicennia germinans*, *Conocarpus erectus*, *Laguncularia racemosa*, *Rhizophora racemosa*,  
184 *Rhizophora mangle*, *Rhizophora harissonii*) but three dominate almost the mangrove forest:  
185 *Rhizophora racemosa*, *Rhizophora mangle*, and *Avicennia germinans* (Sow et al., 1991; Ndour  
186 et al., 2012; Andrieu et al., 2020; Lombard et al., 2020). Species composition, in addition to  
187 vegetation cover and tree height, is spatially structured in zonation, and can be subdivided into  
188 plant formations. Mangroves in Senegal and Guinea-Bissau have the same zonation patterns

189 (Sow et al., 1991), but the studied hydrosystem showed unique features. In the Saloum Delta  
 190 (and the Casamance Estuary), the salinity increases from downstream to upstream (Barousseau  
 191 et al., 1985; Descroix et al., 2020), whereas the estuaries of Gambia and those located in  
 192 Guinea-Bissau, for example, have a river input and upstream salinity levels closer to those of  
 193 fresh water. For Saloum, this results in hypersalinity of the water even after the rainy season  
 194 (Diop, 1997). The salinity of the water throughout the Saloum Delta was higher than that of  
 195 the sea. For example, in Sine-Saloum in June, salinity is approximately 35‰ at the mouth of  
 196 the delta and up to 100‰ in Fatick (82 km from the mouth) (Descroix, 2018). Mangrove  
 197 ecosystems in coastal areas are affected by their interplay of soil, topography, and salinity,  
 198 which bring complexity to the fine scale. (Marius, 1985; Diop, 1990). In such an ecological  
 199 context, the vegetation shows two gradients, north–south and east–west, where the density and  
 200 height of the stands differ without changing the specific composition. This makes detecting  
 201 species and plant formations within the Saloum Delta challenging. Therefore, sixty fields  
 202 surveys were conducted (Figure 1). Using transects following the protocol developed by  
 203 Andrieu et al. (2020), the density of the vegetation cover, height of individuals, and rate of  
 204 cover of each species were inventoried, which facilitated the evaluation of vegetation formation  
 205 maps.

206  
 207



208  
 209 **Figure 1:** Extent study site and field-based inventory for mangrove zonation.  
 210 Sixty botanical surveys were performed and detailed the species, the extent of their relative abundance, and the  
 211 height of the vegetation community.  
 212



213 *2.2.Data collection and pre-processing*

214 Sentinel-1 and Sentinel-2 images were selected for this study (Table 1). These data have  
 215 been proven to be effective for mangrove mapping (Wang et al., 2018; Mondal et al., 2019;  
 216 Wang et al., 2019; Jia et al., 2021; Lu and Wang, 2021; Lui et al., 2021; Xiao et al., 2021). The  
 217 Sentinel-2 Level-2A data is available through Google Earth Engine. This product offers  
 218 orthorectified Bottom-Of-Atmosphere (BOA) reflectances for all 12 bands in a 16-bit unsigned  
 219 integer format that has been scaled by a factor of 10,000. We selected all available images (n  
 220 = 79) from both sensors between 01/05/2020 and 01/06/2021 on the GEE platform. For  
 221 Sentinel-2, the images were filtered with a value of 20% cloud cover. All processing of the S2  
 222 time series was performed on the unmasked pixels using ‘QA60’ quality layers. The range of  
 223 the number of clear observations is from 47 to 65 cloud-free dates (Table 2). All spectral bands  
 224 at 10 m and 20 m resolution of the Sentinel-2 sensor were retained. Bands at 20m were  
 225 resampled at 10m. B1 and B9 were not included because they depict atmospheric traits, like  
 226 aerosols, water vapor, and cirrus clouds, rather than reflecting surface characteristics of land-  
 227 based features. The annual median was calculated for the optical spectral bands.

228  
 229  
 230  
 231

Table 1: Characteristics of Sentinel-1 and Sentinel-2 data

Sensor	Processing Levels	Acquisition Period	Relative orbit	Tiles
Sentinel 1	Level - GRD	01/05/2020 - 01/06/2021 (33 scenes)	133	-
Sentinel 2	Level – 2A	01/05/2020 - 01/06/2021 (79 scenes)	-	T28PBA, T28PCA, T28 PCAB, T28PBB, T28PDA, T28PDB

232  
 233  
 234

Table : The dates of observation for Sentinel-2 pixels that cloud-free are presented. These dates may differ for each pixel, and the dates shown depict those pixels with the minimum and maximum number of cloud-free days within the analyzed region

Year	Month	Days	Days
		(65 cloud-free dates )	47 cloud-free dates
2020	May	02, 07, 12, 17, 22, 27	02, 12, 17, 22, 27
	June	11, 16, 21, 26	11, 16, 21, 26
	July	01, 06, 21, 26, 31	01, 21
	August	05, 15, 20, 25, 30	15, 25
	September	09, 19, 24, 29	19, 24, 29
	October	04, 09, 14, 19, 24, 29	09, 14, 19, 24, 29
	November	08, 13, 18, 23, 28	08, 13, 18, 28
	December	03, 08, 13, 28	03, 08, 13, 28
2021	January	02, 07, 12, 27	02, 07, 27
	February	01, 06, 11, 16, 21, 26	21, 26
	March	08, 13, 18, 23, 28	03, 08, 13, 18, 28

April	02, 07, 12, 17, 22, 27	02, 07, 12, 17
May	02, 07, 12, 22, 27	07, 12, 22, 27

235

236 Four spectral indices were calculated. The normalized differential vegetation index  
 237 (NDVI; Rouse et al., 1974) measures the active photosynthesis of vegetation using reflectance  
 238 differences in the red and near-infrared wavelengths; it can be used to distinguish vegetation  
 239 from other types of land cover. The near infrared surface water index (NDWI, McFeeters,  
 240 1996) measures the amount of water present on the surface of soils using reflectance  
 241 differences in the near-infrared and mid-infrared wavelengths; it can be used to detect flooded  
 242 areas and to distinguish mangroves from drier vegetation types (Gupta et al, 2018). The  
 243 normalized difference moisture index (NDMI, Gao 1996), in particular the DBSI (Dry and Wet  
 244 Soil Index) was designed to specifically identify bare areas in dry climates, according to Rasul  
 245 et al. (2018). They were selected to identify the wet and dry areas constituting the sometimes  
 246 semi-arid landscapes of the study area. The previously mentioned indices (NDVI, NDMI,  
 247 NDWI, and DBSI) were used to calculate the GLCM textural indices introduced by Haralick  
 248 et al. in 1973. Finally, time-series variables were used to complement reflectance and spectral  
 249 indexes dataset. The phase and amplitude of the harmonic model (Clinton, 2016) were  
 250 extracted from the annual time series of four spectral indices in the GEE.

251 The Sentinel-1 satellite imagery in the Earth Engine 'COPERNICUS/S1\_GRD' image  
 252 collection consists of Level-1 Ground Range Detected (GRD) scenes that have been processed  
 253 to calculate the backscatter coefficient ( $\sigma^0$ ) in decibels (dB). The backscatter coefficient is a  
 254 measure of how much microwave radiation is scattered back to the radar sensor by the target  
 255 area and is expressed as the target backscattering area (radar cross-section) per unit of ground  
 256 area. Since it can vary significantly, it is converted to dB by  $10 \cdot \log_{10} \sigma^0$ . The backscatter  
 257 coefficient can be used to determine physical characteristics of the terrain, including the  
 258 geometry and electromagnetic properties of the terrain elements.

259 To calculate the backscatter coefficient, Earth Engine uses a series of preprocessing  
 260 steps implemented with the Sentinel-1 Toolbox. These steps include applying orbit files to  
 261 update metadata, removing low intensity noise and invalid data on the edges of the scenes,  
 262 removing thermal noise in certain acquisition modes, applying radiometric calibration values,  
 263 and correcting the terrain data using either the SRTM 30 meter or ASTER DEM. The terrain  
 264 correction step converts the data from ground range geometry, which does not take terrain into  
 265 account, to  $\sigma^0$ , resulting in a more accurate representation of the terrain.

266 In the absence of top-down data for this area, only bottom-up data were selected, and the two  
 267 polarizations (VH and VV) were combined. A total of 33 images for the period from May 1st,  
 268 2020 to June 1st, 2021 have been processed. For the SAR data, the ratio of VH to VV  
 269 polarization (VH/VV) was calculated. Four variables were extracted from the VH, VV, and  
 270 VH/VV bands over an annual time series: the median, minimum, maximum, and coefficient of  
 271 variation (standard deviation divided by the mean). The dataset consisted of 102 predictor  
 272 variables before being optimized through selection. (Table 3).

273

274 Table 3: A list of initial variables for classification level-1 (mangrove and land cover)

Sensor	Description	Bands name	Wavelength range	Resolution
--------	-------------	------------	------------------	------------

Spectral bands			
	Band2 : Blue	458–523 nm	10 m
	Band3 : Green	543–578 nm	10 m
	Band4 : Red	650 – 680 nm	10 m
	Band5 : Red Edge 1	698–713 nm	20 m
	Band6 : Red Edge 2	733–748 nm	20 m
	Band7 : Red Edge 3	773–793 nm	20 m
	Band8 : NIR	785–900 nm	10 m
	Band8A : Red Edge 4	855–875 nm	20 m
	Band11 : SWIR 1	1565–1655 nm	20 m
	Band12 : SWIR 2	2100–2280 nm	20 m
Spectral index		Formula	Reference
	NDVI	$(B8-B4)/(B8+B4)$	Rouse et al. (1974)
	NDWI	$(B3 - B8) / (B3 + B8)$	McFeeters (1996)
	NDMI	$(B8 - B11) / (B8 + B11)$	Gao (1996)
	DBSI	$B11 - B3)/(B11 + B3) - NDVI$	
Harmonic index (calculated on the spectral indices)			
	Phase	displacement between the origin and the peak of the wave in the range of 0 to $2\pi$	Jakubauskas et al. (2002)
	Magnitude	half the value at which the function is maximized	Jakubauskas et al. (2002)
Texture information (calculated on the spectral indices)			Haralick et al. (1973)
	Angular Second Moment: measures the number of repeated pairs		
	Contrast: measures the local contrast of an image		
	Correlation: measures the correlation between pairs of pixels		
	Variance: measures how spread out the distribution of gray-levels is		
	Inverse Difference Moment: measures the homogeneity		
	Sum Average		
	Sum Variance		
	Sum Entropy		

Entropy: Measures the randomness of a gray-level distribution  
 Difference variance  
 Difference entropy  
 Information Measure of Corr. 1  
 Information Measure of Corr. 2  
 Dissimilarity  
 Inertia  
 Cluster Shade  
 Cluster prominence

Sensor	Description	Bands name	Centre frequency	Resolution
Sentinel-1	SAR	VH- ascending	5.405 GHz	X: 10 m Y:10 m
		VV- ascending	5.405 GHz	X: 10 m Y:10 m
		VH/VV		
	SAR statistics on the time series (calculated on <i>VHasc</i> , <i>VVasc</i> , <i>VHasc/VVasc</i> )	Minimum		
		Maximum		
		Median		
		coefficient of variation		


275

276 **3. Methods**

277 The complete methodological flowchart is shown in Figure 2. This method is divided into two  
 278 levels:

279 Level 1: Mapping the mangroves and land use required for linear spectral unmixing

280 Level 2: Mapping vegetation formations by object-based spectral-temporal feature  
 281 classification



**Figure 2:** Flowchart of the proposed approach.

At white background: treatments allowing to obtain a cartography of the mangrove.

At light green background, treatments allowed us to obtain a cartography of mangrove plant formations.


### 3.1. Mangrove and land cover mapping (Level-1)

First, the mangrove is mapped using a combination of OBIA and machine learning approaches, specifically Random Forest. Although the pixel approach has been extensively used for mangrove classification (Bunting et al., 2018; Mondal et al., 2019; Jia et al., 2021; Lu and Wang, 2021; Lui et al., 2021; Xiao et al., 2021), we chose an object approach (Myint et al., 2008; Wang et al., 2018; Wang et al., 2021) to reduce spectral confusion by smoothing the dataset features within objects and reducing the statistical noise of the images (Blaschke et al., 2008; Blaschke, 2010; Dronova, 2015; Hossain and Chen, 2019). This reduces the complexity of the images and the size and dimensionality of the dataset. In addition, the optimal features for classification were evaluated. With textural characteristics, objects appear as a privileged entity to evaluate the importance of texture indices (GLCM) in classifying inter-tidal areas.

#### 3.1.1. Segmentation

Segmentation is the first classification process. The superpixel clustering approach based on simple non-iterative clustering (SNIC) using the GEE was chosen. Bands with the highest optical spatial resolution only (B2, B3, B4, and B8) were implemented. The SNIC algorithm (Achanta and Süsstrunk, 2017) uses a regular grid of 'seeds' as the initial centroid of the superpixel. The parameterization of the algorithm requires the specification of superpixel seed spacing (in pixels). The larger the spacing, the larger the object size. Therefore, it is critical to scale parameters. Several methods have been developed to determine the optimal threshold for the scale parameters (Ma et al., 2017). The scale parameter estimation (ESP) initially proposed by Drăguț et al. (2010) and modified for multispectral imagery (Drăguț et al. 2014) was chosen because of its low computational cost, and therefore, could be applied to large areas. For each 2–50-pixel spacing step, the local variance was calculated for each band and then averaged

311 (Drăguț et al. 2014) for fixed values of connectivity at 8 and compactness at 1. The rate of  
 312 change (ROC) was then calculated for each scale change. The first break in the curve after  
 313 continuous decay is the threshold at which significant objects emerge. In other words, the scale  
 314 value showing for the first time a negative change (Figure 3) in the ROC curve was selected as  
 315 the optimal seed spacing value.



316

**Figure 3:** Estimate scale parameter.

317

318 The graphs show the changes in local variance (LV) (black) and rate of change (ROC) (gray dots) with  
 319 increasing scale parameters. Vertical lines in red indicate the optimal scaling parameters selected for Saloum

320

### 3.1.2. Training and validation samples

321

322 The training and validation of machine learning classifications require reference sample data  
 323 for land cover. Four classes were referenced: Water, Mangrove, Salt flats, Other land covers  
 324 (all continental surfaces, including bare soil, other vegetation, urban, and cropland). Reference  
 325 data collection followed a labeling procedure using ESRI (2017) and Google Earth (2018,  
 326 2021) high-resolution imagery, with a Sentinel-2 color composition check from 2021 to ensure  
 327 the class stability spotted on slightly older THRS imagery.

328 Stratified random sampling was applied to avoid bias during the validation (Olofsson  
 329 et al., 2014). Therefore, a random draw was performed in the nine classes of the ESA  
 330 WorldCover 10 m 2020 v100 map (Zanaga et al., 2021) with 400 points per class. This map  
 331 had an overall accuracy of 74.4%; therefore, we removed or corrected erroneous data. After  
 332 filtering, 2592 reference samples were collected, of which 75% and 25% of the samples were  
 333 retained for training and mapping validation, respectively. For model training, 223, 312, 322,  
 334 and 1,660 samples were retained for water, mangrove, salt flats, other land covers, respectively.  
 335 The objective was to obtain a statistical representation of the different land cover types within  
 336 the dry land class. For model validation, 74, 77, 80, and 414 samples from water, mangrove,  
 337 salt flats, other land covers, respectively, were selected.

338        *3.1.3. Classification processing and accuracy assessment*


339        The classification into water, mangrove, salt flats, other land covers was performed using the  
340        random forest algorithm (Breiman, 2001) on the GEE platform. Although the random forest  
341        model has the ability to handle many variables (Belgiu and Dragut, 2016), we chose to reduce  
342        the dimension of the dataset by retaining only the so-called optimal variables. To develop  
343        simple and parsimonious classification models, the use of a variable selection algorithm allows  
344        the maintenance of a small number of variables. This will allow for fewer data collection  
345        procedures in the future and a more efficient evaluation of the variable importance. Speiser et  
346        al. (2019) mentioned that reducing the number of variables in random forest models can also  
347        improve the efficiency of classification. To address these objectives, a backward elimination  
348        approach was used, based on model performance as a function of the rate of change in  
349        classification accuracy when adding or removing variables (Díaz-Uriarte and De Andres,  
350        2006). The VarSelRF package in R statistical software was used to select variables. The ‘Out  
351        of Bag’ (OOB) error was used as a minimization criterion and the least important variables  
352        were eliminated. In each iteration, 10% of the variables were eliminated. This procedure was  
353        repeated 20 times, as recommended by Li et al. (2016) and Wang et al. (2018), to select a subset  
354        of the most important features. To assess the stability of the classifications, a 10-fold repeated  
355        cross validation was undertaken. The accuracy scores for mangrove, inter-tidal, and upland  
356        classifications were obtained using a confusion matrix (Stehman 1997). Overall accuracy (OA),  
357        producer accuracy (PA), user accuracy (UA), and kappa coefficients were generated. Finally,  
358        an overall F1 score was used to obtain the accuracies for each land cover class.

359

360        *3.2. Mangrove zonation mapping (Level-2)*

361        We also used the OIBA approach to map mangrove zonation by discriminating its plant  
362        formations. Segmentation is a crucial step and is performed using a linear spectral unmixing  
363        image of the vegetation fraction. The first mapping allowed for the selection of the  
364        endmembers. Based on the model of Taureau et al. (2019), we decomposed the pixel intensity  
365        into three endmembers, each representing a component of the mangrove ecosystem.: water,  
366        mangrove and salt flats. The 10th decile of the mangrove NDVI was retained. Mudflats with  
367        the lowest coefficient of variation of NDWI were chosen for their spectral stability throughout  
368        the year. Non-turbid water spectral was obtained by unsupervised classification of the water  
369        mask into 30 classes and selection of the spectral signature with the lowest reflectance values  
370        in the Sentinel-2 optical bands. The spectral signatures of the endmembers are shown in Figure  
371        4.

372



373

374 **Figure 4:** Spectral reflectance curves for water, mangrove, and salt flats selected for linear spectral unmixing.

375

376 Four images were generated to represent the contribution of each endmember to each  
 377 pixel. Therefore, the pixels misclassified during the first step could be partially corrected by  
 378 considering the most important contribution of the endmember to definitively assign the classes  
 379 of mangrove, salt flats, and water. We preferred the OBIA approach in the first instance because  
 380 the optimal object size avoids confusion between mangroves and dense dryland vegetation that  
 381 was obtained using the pixel approach. Nevertheless, the optimal object size was calculated  
 382 globally, not locally. Therefore, objects that are too large in areas of mangrove vegetation that  
 383 combine water and salt flats may remain. Hence, Linear Spectral UnmixingLSU allows the  
 384 pixels to be reassigned to the right classes within the objects.

### 385 3.2.1. Segmentation and spatial structure of the mangrove

386 Intra-mangrove segmentation is a key step in determining homogeneous features that reflect  
 387 mangrove distribution along environmental gradients. The SNIC algorithm under GEE was  
 388 used again by varying the superpixel seed spacing in steps of 1 and up to a spacing of 20. We  
 389 considered that the ‘optimal’ intra-mangrove objects could not be larger than 20, because the  
 390 optimal scale parameter for land cover classification was 18. The choice of the segmentation  
 391 scale parameter is crucial because it will have important consequences on the clustering and  
 392 thus the typology of vegetation formations. To better detect transitions within the mangrove,  
 393 the selection of the optimal scale parameter must meet the criteria of intra-segment  
 394 homogeneity and inter-segment heterogeneity (Espindola et al., 2006; Johnson and Xie, 2011;  
 395 Johnson et al., 2015). For this purpose, the most common method is to calculate the weighted  
 396 variance (wVar) as an indicator of intra-segment spectral homogeneity (Eq. 1) and the Moran  
 397 Index (MI) to measure the inter-segment spectral heterogeneity (Eq. 2).

$$398 \quad wVar = \frac{\sum_{i=1}^n a_i \times v_i}{\sum_{i=1}^n a_i} \quad (1)$$

399 where  $n$  is the total number of objects

400  $v_i$  and  $a_i$  are the variance and area of the object  $i$ , respectively. Large objects have a greater  
 401 impact on the overall variance than small ones.



402

403

$$MI = \frac{n \sum_{i=1}^n \sum_{j=1}^n w_{ij} (y_i - \bar{y})(y_j - \bar{y})}{\sum_{i=1}^n (y_i - \bar{y})^2 (\sum_{i \neq j} \sum w_{ij})} \quad (2)$$

405

406 where  $n$  is the total number of objects

407  $y_i$  and  $\bar{y}$  are the average spectral values of the object  $i$  and the whole image

408  $w_{ij}$  is a measure of the spatial proximity between object  $i$  and  $j$ .

409

410 For the calculation, the objects were in a vector format. We considered the adjacency  
 411 of the segments as spatial proximity; if a single vertex of an entity is adjacent to another entity,  
 412 these two segments are neighbors. Therefore, we constructed a matrix of spatial weights.  
 413 Several studies (Johnson and Xie, 2011; Wang et al., 2021) have set the values of  $w_{ij}$  to zero  
 414 (non-neighbor) or one (neighbor). Binary matrices can induce statistical bias, generating  
 415 different weights for individuals depending on their neighbor numbers. To overcome this  
 416 problem, in-line standardization of the matrix was performed. Thus, the sum of the weights in  
 417 each row is equal to one. This operation was performed with the package ‘spdep’ in R (Bivand  
 et al., 2015).

418

419 The values of wVar and MI were normalized for comparison according to the following

420 normalization formula (eq. 3):

$$X_{norm} = \frac{(X - X_{min})}{(X_{max} - X_{min})} \quad (3)$$

421

422 Where  $X_{norm}$  is the normalization of the values of wVar or MI.

423  $X_{min}$  is the minimum value of wVar or MI.

424  $X_{max}$  is the maximum value of wVar or MI.

425

426 Finally, the F-measure was calculated (eq. 4) to capture the best compromise between  
 427 intra-homogenous and inter-heterogenous segmentation (; Johnson et al., 2015; Wang et al.,  
 428 2019).

$$F = \frac{1}{\alpha \frac{1}{wVar} + (1-\alpha) \frac{1}{MI}} \quad (4)$$

430

431 where  $\alpha$  (between 0 and 1) is the relative weight of the normalized values of wVar and MI.  
 432 Thus, intra-segmental homogeneity or inter-segmental heterogeneity can be assigned higher  
 433 importance. In this study, the weights of the two indices are considered the same. Therefore,  
 the weight was set as  $\alpha = 0.5$ . A large F-value indicates high-quality segmentation.

### 434 3.2.2. Clustering of mangrove zonation

435

436 To capture this cycle, the time series of the plant fraction was modelled using harmonic  
 437 regression in GEE (Clinton, 2016). This method has been shown to be relatively effective in  
 describing mangrove zonation patterns in semi-arid areas with Sentinel-2 images (Valderrama-

438 Landeros et al., 2021). Therefore, regression models with harmonics were used. We fitted and  
439 calculated the harmonic coefficients and extracted the amplitude (seasonal variability), annual  
440 trend (annual average), and RMSE (non-seasonal variability) to characterize the phenological  
441 patterns of the different objects (Pasquarella et al., 2018). Model fit was better for harmonic  
442 regressions based on fractional components from linear unmixing than for spectral indices,  
443 justifying the use of the vegetation fraction instead of NDVI. The objective here is to  
444 discriminate these zonations on an annual basis rather than revealing the variation itself. The  
445 annual average is indeed a feature that mostly influences classification.

446 The small sample size from the field made the training and validation process of a  
447 supervised classification difficult; therefore, unsupervised classification with the K-means  
448 algorithm was preferred. The objective was to obtain a typology aimed at maximizing the  
449 contrasts between classes while reflecting the zonation and physiognomy of the vegetation.  
450 Field data were used to interpret the clusters. To ensure the overall significance of the classes,  
451 the Kruskal-Wallis test was performed. In addition, pairwise multiple comparisons between  
452 groups were used to evaluate the significance between classes using the Wilcoxon test. These  
453 tests were performed with field-derived variables, such as cover rate, stand height, and species  
454 dominance. These tests were performed with all variables of the harmonic model describing  
455 seasonality and with different numbers of classes from two to six. To evaluate the significance  
456 of the differences between classes, the threshold p-value of 0.01 was retained.

### 457 3.2.3. Stability assessment of the typology

458 The detection of mangrove zonation typology must be reproducible over time to ensure its  
459 monitoring and reliably interpret its changes. Indeed, a typology in which the classes do not  
460 have enough contrast at certain times of the year could lead to errors in detecting changes.

461 To evaluate typology and temporal stability, we used a three-step process. First, the  
462 methodological chaining presented so far was repeated on the data from 01/11/2019 and  
463 01/11/2020. An intersection between the two classifications allowed us to isolate the objects  
464 assigned to the same class on both dates, which did not undergo any major changes. Finally,  
465 these were used as a basis for training (70% of the objects) and validation (30% of the objects)  
466 of the Bayesian model. For this stage, the probabilities of membership were calculated using  
467 Bayes' theorem.

$$468 \quad p(h | e) = \frac{p(e | h) \times p(h)}{\sum_i p(e | h_i) \times p(h_i)} \quad (5)$$

469 where  $p(h | e)$  is the probability that hypothesis (h) is true, given the proof (a posteriori  
470 probability noted e).

471  $p(e | h)$  is the probability of finding that the evidence provided by hypothesis (h) is true (from  
472 the training data).

473  $p(h)$  is the probability that the hypothesis is true, regardless of the evidence (a priori  
474 probability).

475 A series of images expressing the *a posteriori* probability of belonging to a set of plant  
476 formation classes were generated. The objective of this study was to evaluate the degree of  
477 belonging of an object to a class according to the vegetation fraction on both dates. Hard



478 classification was used to obtain a thematic map. For each object, the class with the highest  
479 probability was assigned.

480 Finally, the probable confusion over time was evaluated for each class by noting the  
481 frequency of false predictions of the Bayesian model for each value of the vegetation fraction.  
482 The objective of this experiment was to quantify the amplitude of the fluctuations in plant  
483 fractions that can be interpreted as a change in the spatial structure of mangrove zonation.

## 484 **4. Results**

### 485 ***4.1. Mangrove mapping***

486 At the end of the RF classification of the objects, the Saloum mangrove map (Figure 5a)  
487 presented relatively high accuracy scores (Table 4) using a reduced dataset of 11 discriminating  
488 variables. After ten cross-validations, the optical bands in the NIR and SWIR wavelengths  
489 proved to be the most effective for mapping the mangrove in this environment, followed by the  
490 SAR data (Figure 5b). From then on, the SAR data constituted more than 50% of the selected  
491 variables. Texture variables and time-series harmonic model coefficients were negligible for  
492 the desired land cover discrimination. Finally, 11 variables were selected: VHmin, VHmed,  
493 VHmax, VVmin, VVmed, VVmax for SAR data and B8, B8A, B11, B12, NDWI for optical  
494 data. The consistent appearance of SAR data in the group of optimal variables suggests that it  
495 is a valuable source of information for understanding and predicting the distribution and  
496 behavior of the studied system. In contrast, when optical bands like B8 or B8A were present in  
497 the group of variables, they tended to contribute more to the discrimination of the surface states  
498 of the selected typology.



499

500 **Figure 5a:** Random Forest classifications using Sentinel-1 and Sentinel-2 data (left) from an optimal dataset.

501 Locations refer to the following figures that focus on mangrove zonation

502 **Figure 5b:** Optimal dataset and variables importance.

503 The importance of the variables was measured by the average decrease in the Gini index of the nodes and leaves  
 504 in the resulting random forest. This expresses the degree of precision that the model loses by excluding each  
 505 variable.

506

507 Table 4 shows that the average overall accuracy is 97.12% and the Kappa index is  
 508 94.6%. Within the Sine-Saloum region, mangroves obtain the highest average relative accuracy  
 509 scores (F1-score = 98.55%) among the land cover types. Drylands has similar scores, and salt  
 510 flats had the lowest average accuracy with an F1-score of 88.31. On the other hand, the  
 511 classifications are relatively stable. Indeed, the standard deviation for the global accuracies and  
 512 Kappa index are 0.008 and 0.015, respectively. The surfaces and their fluctuations were  
 513 quantified according to the classifications. The water and mangrove areas varied respectively  
 514 by an average of 0.37% and 0.47%, that is, 721 ha and 317 ha. Variations of 0.1%, that is 756  
 515 ha, were observed on average for the drylands. Finally, the salt flats showed more important  
 516 fluctuations in the estimation of their surfaces, of the order of 2% or 1029 ha.

517

518 Table 4 shows that the contribution of post-classification linear spectral unmixing  
 519 brings a slight gain in overall precision (+ 0.99%) and (+ 1.88%) on the Kappa index. We also  
 520 observed a significant gain in relative precision class by class. Nevertheless, the LSU  
 521 contribution is sometimes better at the pixel level than at the object level, notably for water (F1  
 522 score = + 1.95 against 1.39 for RF-OBIA + LSU-OBIA) and mangroves (F1 score = + 0.36%  
 523 against -1.33 for RF-OBIA + LSU-OBIA). Given the scores and the purpose of this paper, the  
 524 mangrove areas from the RF-OBIA + LSU method were retained for mapping the vegetation  
 525 formations.

526

527 Table 4: Scores assessing the accuracy of the classifications and their impact on the quantification of Land  
 528 Cover types


<i>Accuracy</i>	<i>OA %</i>		<i>Kappa %</i>		<i>F1 - Score %</i>			
	<i>Mean</i>	<i>Std</i>	<i>Mean</i>	<i>Std</i>	<i>Water</i>	<i>Mangrove</i>	<i>Salt flats</i>	<i>Drylands</i>
(RF-OBIA)	97.12	0.8	94.6	1.5	96.30	98.55	88.31	98.53
(RF-OBIA + LSU- OBIA)	98.11	0.8	96.48	1.5	97.69	97.22	92.16	99.46
(RF-OBIA + LSU)	97.55	0.9	95.42	1.7	98.25	98.91	90.10	99.35
<i>Area</i>	<i>Area (ha)</i>				<i>Std Area (ha)</i>			
	<i>Water</i>	<i>Mangrove</i>	<i>Salt flats</i>	<i>Drylands</i>	<i>Water</i>	<i>Mangrove</i>	<i>Salt flats</i>	<i>Drylands</i>
(RF-OBIA)	193 214	67 762	50 380	750 382	721	317	1 029	756
(RF-OBIA + LSU- OBIA)	192 843	65 341	53 188	749 373	-	-	-	-
(RF-OBIA + LSU)	193 578	63 124	54 762	750 272	-	-	-	-

529

530 The linear spectral unmixing measuring, within a pixel, the fraction of each land cover  
 531 showed that the initial (object-based random forest) classification overestimated the mangrove  
 532 areas by an average of 4638 ha or 6.84%. The consequent decrease in mangrove surfaces with  
 533 the RF-OBIA + LSU approach coincided with the increase in salt flats surfaces. Therefore, the  
 534 main confusion concerns the mangroves and salt flats. Of the mangrove surfaces, 6.43% were

535 classified as salt flats using the RF-OBIA + LSU approach, and 0.42% as water. Therefore,  
536 93.5% of the classification differences between the two approaches concerned mangroves and  
537 saltflats (Figure 6).

538  
539  
540  
541



542  
543 **Figure 6:** Example of correction of the RF-OBIA +LSU approach. Red contours represent areas classified as  
544 mangrove before correction and as salt flats after correction. Most of the confusion concerns the mangrove-salt  
545 flats interface.

546

## 547 4.2. Mangrove zonation mapping


### 548 4.2.1. Optimal scale

549 Figure 7 shows the optimal scale obtained by maximizing inter-segment heterogeneity and  
550 intra-segment homogeneity. A spacing of nine pixels allows the SNIC approach to achieve an  
551 optimal global trade-off. The spacing from 2 to 11 pixels generated objects with a very different  
552 internal variance from one spacing to the other, increasing faster before the slope of the curve  
553 became smoother. The Moran index decreased consistently over the same interval, meaning  
554 that the objects diverge increasingly from their neighborhood. The F-metric allows the best  
555 global compromise to be synthesized. Therefore, the tested spacings from 2 to 8 induced a  
556 satisfactory internal homogeneity of the objects, but an extremely low global inter-object  
557 heterogeneity. A spacing of 2–8 induced over-segmentation; thus, the tested spacings from 10

558 to 20 induced extremely low intra-object homogeneity, but acceptable global inter-object  
 559 heterogeneity. Therefore, we concluded that a spacing of 10–20 induces under-segmentation.

560 Figure 7 shows the consequences of the chosen scale on vegetation fraction index segmentation  
 561 (Figures 7b and 7c). A THRS Google Earth image (Figure 7a) allows for apprehending the  
 562 grain of the vegetation, its retranscription by the vegetation fraction index (Figure 7b), and the  
 563 consideration of its spatial structure by segmentation (Figure 7c). Visually, the strong values  
 564 of the index (in red) synonymous with a high and dense mangrove, often at the edge of banks,  
 565 seem well-captured as a single object by the OBIA approach. On the other hand, some places  
 566 where the index is lower and where the vegetation cover appears less dense on the THRS image  
 567 appear oversegmented.

568



569

570

**Figure 7:** Optimal segmentation scale for detecting homogeneous plant formations.

571

a) High resolution Google Earth image of South Saloum

572

b) The vegetation fraction index from linear unmixing (Red values represent the largest vegetation fraction and can be equated with dense, high shoreline mangroves. The green gradient can be assimilated into the decrease in the density and height of the mangrove along the foreshore)

573

574

c) Optimal object segmentation

575

576

d) Results of optimal scale parameter detection. The 9 pixels spacing for SNIC segmentation is the best compromise between intra-homogenous and inter-heterogeneous segmentation.

577

578

579



580

4.2.2. Spectral Seasonality-Based Mangrove Zonation Classification Using Spatial Objects

581 The vegetation indices of these mangroves followed a seasonal pattern that could be captured  
582 by the vegetation fraction of a pixel. High values of the vegetation fraction index were  
583 observed during the rainy season, mainly from August onwards. A decrease in these values  
584 was observed at the end of the rainy season in November.

585 The discrimination of mangrove plant formations by time-series analysis meets the objective  
586 of developing a reliable and robust typology over time. Figure 8 shows that during the course  
587 of the year, the values of the indices do not always capture contrasts in the spatial structure of  
588 the vegetation. Indeed, at the end of the dry season, there were sometimes no differences in  
589 plant fraction index values between the two plant formations. The values obtained for  
590 18/06/2020 and 02/04/2021 (fig. 8) were similar between the high bank mangrove (HM) and  
591 the low and dense mangrove (LDM), and between the latter and the low and open mangrove  
592 (LOM). Therefore, single-date images and classifications based on the values alone may be  
593 inoperable.





594

595

Figure 8: Annual phenological patterns of mangrove zonation.

596 On the x-axis, the time axis ranges from 2020-05-01 to 2021-06-01. On the y-axis, the vegetation fraction values  
 597 from the linear spectral unmixing are shown.

598

The points represent the values within the time series for each class.


599

The solid lines represent the harmonic regression model for each class.

600 Solid line ellipses show the likely confusion between thematically different classes. The dashed ellipse  
601 represents the date when spectral sparsity is maximized.

602 HM : High mangrove – LDM : Low and dense mangrove - LOM : Low and open mangrove

603 Figure 9 shows the mangrove zonation clustering for 2021 and a visual comparison with  
604 Google Earth © images. The number of classes was determined from field data. The Kruskal-  
605 Wallis test was used to ensure the overall significance of the classes. The Wilcoxon test was  
606 used to determine whether the typology presented significant differences between classes with  
607 the parameters found in the field. The results of the different tests show that clustering based  
608 only on the annual average of the values modelled by harmonic regression and with three  
609 classes made the most thematic sense. These results indicate that the density of vegetation cover  
610 contributes significantly (p-value = 0.00057) to the discrimination of facies using the proposed  
611 approach. The three-class typology based on canopy height, canopy openness, and color  
612 appeared to fit well with the spatial resolution of the Sentinel-2 images (Figure 9). The  
613 differences in height between the second and third classes were not statistically significant (p-  
614 value = 0.102). Therefore, these two classes were denoted as low mangroves. Nevertheless, the  
615 proposed approach does not allow for the differentiation of mangroves at a specific level. It  
616 also does not allow for a significant determination of the abundance of species within the  
617 classes (Supplemental material). Therefore, a typology based on height, density of vegetation  
618 cover, and the presence/absence of plant species was adopted (Figure 9, Supplemental  
619 material). Finally, the morphology of the classes and their locations (Figure 9) seem to agree  
620 with the high-resolution data and with theoretical and empirical knowledge of the field (Sow  
621 et al., 1991; Ndour et al., 2012; Andrieu et al., 2020 ).



622

623

**Figure 9:** Mangrove zonation clustering

624

(1: High riverine mangrove dominated by *Rhizophora racemosa*; 2: Low et dense mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*; 3: Low and open mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*)

625

626

627

a) Significance of the differences between classes for the overlap rate.

628

(The results of the Kruskal-Wallis test are available at the top of the figures)

629

b) Significance of the differences between classes for the average height of the individuals

630

(Only the results of the significance tests appear in this figure).

631

c) Map of the mangrove zonation for the year 2021

632

The high mangrove facies have patches with rather long morphologies and are located at the edge of the channel. Patches of this class located in the middle of a mangrove area may indicate the presence of a small channel that is completely covered by a closed canopy. The low and

634

635 open mangrove facies showed a less elongated morphology and were consistently found at the  
 636 interface with the salt flats. Finally, the low and dense mangrove facies were generally located  
 637 between the other two classes, representing the transition between high-density and open  
 638 mangroves. The ground-truth data also confirmed that not all riverine mangrove facies were  
 639 necessarily high. In most concave meanders of channel banks, mangrove forests were medium  
 640 and dominated by the *Rhizophora mangle*. Conversely, in most convex meanders of banks,  
 641 mangrove forests were abundant and dominated by *Rhizophora racemosa*.

642 **4.3. Temporal accuracy**

643 Figure 10 shows the confounding probabilities for each class as a function of the plant fraction  
 644 values. These reflect the average values obtained by harmonic regression between 2019 and  
 645 2021. The overall expected accuracy for classifying the mangrove plant formations in the  
 646 Saloum Delta was 85.5% (Table 5). Confusion probabilities were the highest in classes that  
 647 often have spatial contiguity in zonation. Consequently, the class ‘Low and dense mangrove  
 648 with mixed *Rhizophora mangle* and *Avicennia germinans*’ had the highest prediction error rate.  
 649 This class is likely to be confused with the ‘High riverine mangrove dominated by *Rhizophora*  
 650 *racemosa*’ class with a probability of 5.8%, and with the ‘Low and open mangrove with mixed  
 651 *Rhizophora mangle* and *Avicennia germinans*’ class with a probabbility of 8.3%. Table 5  
 652 presents the confusion matrix from the Bayesian classification. Logically, the prediction errors  
 653 between the two classes located at the lowest and highest positions of the foreshore were  
 654 negligible.

655


656 Table 5: The error matrix for the period (2019-2021) using a Naïve Bayes classification approach on stable  
 657 objects (results are expressed as a proportion).

Class	Reference data		
	High riverine mangrove dominated by <i>Rhizophora racemosa</i>	Low and dense mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>	Low and open mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>
High riverine mangrove dominated by <i>Rhizophora racemose</i>	26.3 %	3 %	0.2 %
Low and dense mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>	2.8 %	28.5 %	4.6 %
Low and open mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>	0.1%	3.7 %	30.7 %
Overall accuracy (%)	85.5 %		

658

659

660 To evaluate the probability of confusion between real change in vegetation cover and seasonal  
 661 fluctuation observed by remote sensing, uncertainty bounds were calculated for each class of  
 662 zonation. Figure 10 shows the probability of confusion between classes as a function of the  
 663 vegetation fraction from linear spectral unmixing. The cumulative frequency curve quantifies  
 664 the proportion of the class subject to probable errors as a function of the plant fraction value.



665  
 666 **Figure 10:** Probability of confusion between classes according to the values of plant fraction from the linear  
 667 spectral unmixing.

668 Red represents ‘High riverine mangrove dominated by *Rhizophora racemosa*’

669 Dark green represents ‘Low and dense mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*’

670 Light green represents ‘Low and open mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*’

671

672 Probable confusion between «High riverine mangrove dominated by *Rhizophora racemosa*» and «Low and  
 673 dense mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*»

674 a) Probable confusion between “Low and dense mangroves with mixed *Rhizophora mangle* and *Avicennia*  
 675 *germinans*” et “High riverine mangrove dominated by *Rhizophora racemosa*”.

676 b) Probable confusion between “Low and dense mangrove with mixed *Rhizophora mangle* and *Avicennia*  
 677 *germinans*” et “Low and open mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*”.

678 c) Probable confusion between “Low and open mangrove with mixed *Rhizophora mangle* and *Avicennia*  
 679 *germinans*” et “Low and dense mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*”

680

681 Plant fraction values ranging from 61% to 100% were reliable for assigning to the class ‘High  
 682 riverine mangrove dominated by *Rhizophora racemosa*’. Next, the class ‘Low and dense  
 683 mangrove with mixed *Rhizophora mangle* and *Avicennia germinans*’ had a plant fraction range  
 684 between 38% and 52%, which does not seem to suffer from uncertainties. Finally, values below  
 685 33% seem to be reliably assigned to the class ‘Low and open mangrove with mixed *Rhizophora*  
 686 *mangle* and *Avicennia germinans*’. For clarity, Table 6 summarizes the uncertainty bounds of  
 687 the classes.

688

689 Table 6: Uncertainty intervals for assigning classes over time according to the plant fraction values.

690

The vegetation fraction is expressed here as a percentage.

	Class		
	High riverine mangrove dominated by <i>Rhizophora racemosa</i>	Low and dense mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>	Low and open mangrove with mixed <i>Rhizophora mangle</i> and <i>Avicennia germinans</i>
1st Quartile	58	42	29
Median	61	46	33
3rd Quartile	65	50	36
Incertitude	52 % < VF < 61 %	38 % > VF > 51 %	33 % < VF < 38 %

691

692

693 According to these results, the densification of the low mangroves can be reliably interpreted  
694 only if the class values cross the threshold of a vegetation fraction higher than 38% (or more).  
695 Conversely, the opening of the low mangrove can be reliably interpreted at a threshold of 33%  
696 (and below) vegetation fraction. A disturbance in the high shoreline mangrove can be credibly  
697 detected only if the values of this class fall below the threshold plant fraction of 52%.  
698 Conversely, a recovery can be reliably detected if the values of the low dense mangroves cross  
699 the 61% vegetation fraction threshold.

## 700 5. Discussion

### 701 5.1. Extraction of significant objects for mangrove monitoring

702 The databases on mangroves in West Africa cover only the surface area. Zonation maps are  
703 desirable for guiding conservation strategies (Jia et al., 2018; Wang et al., 2019). However,  
704 global or continental databases contain inaccuracies that bias the mapping of plant formations.  
705 Here, the differentiation of the mangroves approaches an accuracy of 98.91%, in line with those  
706 obtained by Andrieu et al., 2020 and Liu et al., 2021. We aimed to establish a map of mangroves  
707 because the quantifications of surfaces in the literature have significant discrepancies.

708 Fent et al. (2019) reported areas of 86 426 ha in 2018 for the same spatial coverage of 30 m.  
709 Liu et al. (2021) made their data freely available. For the same spatial extent, in 2017, the extent  
710 of areas reached 70 140 ha at 20 m. Andrieu et al. (2020) mentioned a surface area of 63 189  
711 ha at 30 m, in 2015, for the mangrove of the Saloum and Joal hydrosystems, but without the  
712 mangrove of the Gambia River tributary, which is within the administrative boundaries of  
713 Senegal. In this study, we obtained a surface area of 60 988 ha in 2021 at a spatial resolution  
714 of 10 m. A cartographic comparison showed that the areas mapped by Liu et al. (2021) were  
715 overestimated (Supplementary material). The mapping had inconsistencies; for example, some  
716 water channels were mapped as mangroves. However, this mapping was carried out over the  
717 entire West Africa, and currently, it is the most accurate database for all West African  
718 mangroves at this scale. Furthermore, the areas were likely overestimated by Fent et al. (2019)  
719 because the values were widely higher than those estimated by other studies (Andrieu et al.,  
720 2020; Liu et al., 2021).

721 Plant formations were mapped on a mangrove mask using the OBIA approach. The vegetation  
722 fraction was segmented by linear spectral unmixing. The OBIA approach is preferable for  
723 many reasons. First, the vegetation facies of Senegalese mangroves comprise one or more  
724 species. Owing to the global biogeography of mangroves (species rich in Southeast Asia but  
725 poor in Africa) and the occasionally arid climate, Senegalese mangroves show poor species  
726 richness, and are often monospecific in patches larger than a 10m pixel. Li et al. (2019)  
727 achieved overall accuracy scores of 83% for species mapping in China. Nevertheless, the  
728 species differentiated in this study have their own phenological patterns, which is not the case  
729 in the Saloum Delta. Indeed, even in monospecific pixels or objects, spectral contrasts in the  
730 time series appear essentially in the average annual trend and phenological patterns.  
731 Valderrama-Landeros et al. (2018) achieved scores ranging from 75% to 78% at a specific level  
732 and 10 m resolution. Vizcaya-Martínez et al. (2022) achieved accuracy scores between 92 and  
733 95% for the classification of species with VIs from S2 images. Using the time series, the authors  
734 were able to follow the recovery trajectories of mangrove zoning after disturbance, similar to  
735 the zoning presented in this work in Senegal. However, in our case, the upper mangrove was  
736 mixed within a pixel, and we did not claim to classify the species. Therefore, working on a  
737 plant formation scale is preferable. Transitions between plant formations are noticeable through  
738 changes in the relative density of the genera *Rhizophora* and *Avicennia*, and changes in  
739 vegetation physiognomy. Therefore, our typology converges with that of Valderrama-Landeros  
740 et al. (2021). These authors obtained an overall accuracy of 79% for similar mangroves, both  
741 on their specific content and on canopy density variations. Through remote sensing, these  
742 transitions are identifiable, and these plant formations are detectable using object-oriented  
743 approaches, as demonstrated by Flores De Santiago et al. (2013) and Wang et al. (2018). Plant  
744 formations are relevant units for detailed assessment of the impacts of environmental changes  
745 on mangroves. Monitoring the fluctuation of zonation over time is challenging, and a typology  
746 based on an object-oriented approach would allow for monitoring of plant formations with  
747 similar ecological properties. A pixel-based approach could lead to a heterogeneous  
748 cartographic result without an ecological context and, therefore, is not interpretable. Moreover,  
749 the zoning of the mangrove was unsupervised. Field observations have shown that the species  
750 abundance may be heterogenous within an entity equivalent to a pixel. In addition, in most  
751 pixels and surveys, one of the three species that were abundant in Saloum was absent. This  
752 generates zero inflation, which requires the use of other statistical methods to obtain the  
753 relevant pixel-scale information. The OBIA approach circumvents this problem and allows us  
754 to work on a scale where a specific mixture constitutes detectable and interpretable facies.

755

## 756 ***5.2.Relevance of typology***

757 The results show that the approach used in this work allows us to map the zonation of the  
758 mangrove only on spectral indices, acting as proxies of cover rate and stand height, but does  
759 not allow the extrapolation of information on species abundance for each class. These results  
760 are consistent with those of Lymburner et al. (2020), who obtained a three-class typology based  
761 on canopy openness (woodland (20-50%), open forest (50-80%), and closed forest (>80%)),  
762 using linear spectral unmixing. This typology allowed them to follow evolution over time,  
763 which is a future objective of this work.

764 However, our results suggest that the vegetation fraction per LSU is sensitive to mangrove  
765 seasonality. Therefore, the index can fluctuate without any changes in the canopy openness.

766 Therefore, following the evolution of classes with a known value amplitude and a quantified  
767 uncertainty will allow us to follow the intra-mangrove reorganization by reducing the errors of  
768 omissions and commissions of the changes.

769 Valderrama-Landeros et al. (2021) showed that typologies can be affected by mangrove  
770 phenology, and that certain dates of image acquisition are more conducive to the development  
771 of a reliable typology. In this study, we synthesized the phenological patterns of the classes and  
772 quantified the amplitude of their fluctuations over time, given the frequency of spectral  
773 similarities between classes during the year (Figure 8). Therefore, assessing changes in the  
774 future may be more appropriate, to apprehend signal breaks in the time series metrics of the  
775 mangrove rather than establishing a map with the same typology at each date. Indeed, the latter  
776 may be distorted by inter-annual climatic variability, which affects the spectral indices  
777 (Nicholson, 2013). In addition, Pastor-Guzman et al. (2018) showed that semi-arid mangroves  
778 with the same specific composition respond to the onset of the rainy season with a slight  
779 phenological delay that affects canopy reflectance values.

780 As was mentioned by Valderrama-Landeros et al. (2021), this could bias the classification of  
781 mangrove cover in areas with lower vegetation density. However, these authors showed that  
782 the period between the end of the dry season and beginning of the rainy season was the most  
783 favorable for producing accurate estimates of mangrove cover in Mexico. Our results indicated  
784 a slightly different optimal period. The end of the rainy season and beginning of the dry season  
785 showed the strongest contrasts between the classes on raw values and values modelled by  
786 harmonic regression. The climatic contexts of the two studies are similar; however, the  
787 ‘inverse’ character of Senegalese estuaries greatly influences salinity. A reduction in  
788 photosynthetic activity with increasing water salinity (George et al., 2020) or an impact on the  
789 seasonality of spectral indices (Celis-Hernandez et al., 2022) have been consistently observed.  
790 For Saloum, spectral contrasts appeared to be maximized when species faced lower salinities.  
791 Therefore, a typology based on objects that are more sensitive to spatial variations relative to  
792 neighboring pixels, rather than to fluctuations in pixel values, can partially overcome the  
793 constraints related to the phenology of mangroves mentioned above. Furthermore, the typology  
794 presented here and based on the annual phenological pattern allows us to follow the  
795 recommendations of Li et al. (2019) and Valderrama-Landeros et al. (2021) who advocate  
796 using images over a longer time series to potentially detect possible disturbances. For example,  
797 a similar typology has proven suitable for monitoring hurricane damage (Vizcaya-Martínez et  
798 al., 2022). Impacts of hydrological changes such as salinity could be monitored at least  
799 quarterly. This environmental parameter seems to be a major factor in the spatiotemporal  
800 dynamics of the Senegalese mangrove (Andrieu et al., 2020; Descroix et al., 2020; Lombard et  
801 al., 2020).

802

### 803 ***5.3.Perspectives: distinguishing stresses from disturbances***

804 The choice of typology based on a 1-year time series is justified with a view to developing  
805 classes that are less sensitive to seasonal fluctuations. When mapping changes in mangrove  
806 zonation, we must distinguish between stress and disturbance, which both showed a lasting  
807 effect on zonation. The harmonic regression model allowed us to understand seasonal  
808 fluctuations and annual trends. Here, we used only the mean of the regression model values to  
809 classify the mangrove zonation. The amplitude (seasonal variability), phase (travel time from



810 the origin to the peak of the wave), and RMSE (non-seasonal variability) did not contribute to  
811 the robust detection of zonation. Nevertheless, these indicators could be of great importance  
812 for understanding the response of mangroves to environmental fluctuations. Celis-Hernandez  
813 et al. (2022) captured the impact of physicochemical variables on seasonal fluctuations of  
814 mangroves through phenological metrics. Nevertheless, as these authors pointed out, the long-  
815 term effects of fluctuations in physicochemical parameters are still poorly understood.  
816 Therefore, we suggest that future applications should focus on the impact of climatic variables  
817 and salinity on the spatial organization of mangroves at several temporal scales. We also  
818 recommend characterizing the impact of these variables and their fluctuations at several scales  
819 (spatial and temporal) on the possible threshold effects inducing mangrove dynamics at decadal  
820 scales. The use of long-term time series such as Landsat (Bullock et al., 2017; Pirasteh et al.,  
821 2021) could provide insights into mangrove responses with this typology over time. In this  
822 study, we quantified the uncertainty around plant fraction values to characterize potential  
823 changes in mangrove zonation. In the future, we recommend using change detection  
824 approaches from temporal segmentations, such as LandTrendR (Kennedy et al., 2010) or  
825 CCDC (Zhu and Woodcock, 2014). For example, LandTrendr has been successfully used for  
826 mangroves (de Jong et al., 2021). The uncertainty bounds identified in this work allow the  
827 parameters to be calibrated in a robust manner to effectively detect zonation changes. As was  
828 pointed out by Pasquarella et al. (2022), calibrating the sensitivity of these models to change  
829 detection is a key step in mapping land-cover changes.

## 830 **6. Conclusion**

831 Mapping mangrove zonation is challenging yet important in the management of this ecosystem.  
832 Plant formations are organized along environmental gradients, and therefore allow us to  
833 approximate the outcome of population dynamics processes that operate at a finer scale. Three  
834 homogeneous features were mapped at 10 m spatial resolution from Sentinel-1 and Sentinel-2  
835 data. SAR and optical data have proved to be paramount for effective mangrove mapping. The  
836 vegetation fraction resulting from a linear spectral unmixing on optical data only allowed us to  
837 establish the following clustering: ‘High riverine mangrove dominated by *Rhizophora*  
838 *racemosa*’, ‘Low and dense mangrove with mixed *Rhizophora mangle* and *Avicennia*  
839 *germinans*’, ‘Low and open mangrove with mixed *Rhizophora mangle* and *Avicennia*  
840 *germinans*’. The results of this study show that the average of the values of the plant fraction,  
841 obtained by harmonic regressions on each object, allows robust mapping of the zonation of the  
842 mangrove on the criteria of canopy cover and height of the stands. In addition, the temporal  
843 stability of the typology was assessed. The results indicated that an overall margin of error of  
844 14.5% is expected if the plant formations are discriminated against the annual trend of the plant  
845 fraction. These acceptable results do not obscure the likelihood of confusion between the  
846 classes over time. This leads to probable commission errors and omissions regarding inter-  
847 annual zonation changes that must be addressed. Therefore, this study points out that low  
848 mangrove classes are most likely to be confused over time. Furthermore, from the perspective  
849 of mapping zonation changes on an inter-annual scale, the uncertainty bounds quantified in this  
850 study allowed us to estimate that the growth of dense mangrove stands can be reliably captured  
851 if the initial values of the low dense mangrove class increase between 16% and 45% for values  
852 between the 1st and 3rd quartiles. Conversely, a decrease in the height of the high mangrove  
853 stands while maintaining their density can be interpreted if a decrease of 10% to 20% of the  
854 vegetation fraction is observed. Second, a densification of the low mangrove canopy can be

855 interpreted if an increase of 13% to 28% in the vegetation fraction values was observed between  
856 the 1st and 3rd quartile. Finally, an opening of the low mangrove could be observed if the  
857 values of plant fraction of the initially dense mangrove decreased from 21% to 34% between  
858 the 1st and 3rd quartile of the class. Therefore, these limits were intended to reliably identify  
859 the response of mangrove zonation to environmental changes in the Saloum Delta. The  
860 amplitudes of each zonation class identified in this study could allow the characterization and  
861 distinction of stresses, disturbances, or recovery thresholds using temporal segmentation  
862 methods.

863

864 **Acknowledgments**

865 **We warmly thank Inssa Seydi for the many canoe trips, for having shared these**  
866 **observations with us for many years and for all the help provided in the field.**

867 **Funding details**

868 This work was supported under grant IDEX UCA JEDI and by the CNRS UMR 7300 ESPACE

869 **Disclosure statement**

870 The authors declare that they have no known competing financial interests or personal  
871 relationships that could have appeared to influence the work reported in this paper.

872 **Data availability statement**

873 Publicly available datasets were analyzed in this study. This data can be found here: The  
874 original satellite data from Sentinel-1 and Sentinel-2 are publically accessible from the open  
875 cloud platform Google Earth Engine.

876 Sentinel-1 Data Link: [https://developers.google.com/earth-](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S1_GRD)  
877 [engine/datasets/catalog/COPERNICUS\\_S1\\_GRD](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S1_GRD)

878 Sentinel-2 Data Link: [https://developers.google.com/earth-](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S2_SR_HARMONIZED)  
879 [engine/datasets/catalog/COPERNICUS\\_S2\\_SR\\_HARMONIZED](https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S2_SR_HARMONIZED)

880 The import of Sentinel-1 and Sentinel-2 data, the calculation of the number of images, the  
881 number of cloud-free pixels in the time series, and the harmonic model on the vegetation  
882 fraction derived from linear spectral unmixing are available in the following Google Earth  
883 Engine code : <https://code.earthengine.google.com/29061ee4b48e419baf491180809e7530>

884 The mangrove extent and zonation data products from this research project can be accessed  
885 through the following link: <https://doi.org/10.34847/nkl.c7ecc507>

886 **Supplemental online material**

887 Table: Significance of classes by p-value for the adopted typology based on height, canopy density and  
888 presence/absence of plant species.

889 HM: High mangrove – LDM: Low and dense mangrove - LOM: Low and open mangrove

890 a)

*Canopy Cover*

Class	p-value	p-value adjusted	significance
HM / LDM	0.011	0.021	*
HM / LOM	0.000456	0.001	**
LDM / LOM	0.016	0.021	*

891

892 b)

*Height canopy*

Class	p-value	p-value adjusted	significance
HM / LDM	0.002	0.005	**

HM / LOM	0.0000138	0.0000414	****
LDM / LOM	0.102	0.102	ns

893

894 c)

*Rhizophora  
racemosa*

Class	p-value	p-value adjusted	significance
HM / LDM	0.035	0.071	ns
HM / LOM	0.0000242	0.0000726	****
LDM / LOM	0.096	0.096	ns

d)

*Rhizophora  
mangle*

Class	p-value	p-value adjusted	significance
HM / LDM	0.864	1	ns
HM / LOM	0.807	1	ns
LDM / LOM	0.077	0.231	ns

895 e)

*Avicennia  
germinans*

Class	p-value	p-value adjusted	significance
HM / LDM	0.045	0.091	ns
HM / LOM	0.004	0.012	*
LDM / LOM	0.332	0.332	ns

896 **References**

897 R. Achanta and S. Ssstrunk. 2017. "Superpixels and Polygons Using Simple Non-Iterative  
898 Clustering." In 2017 IEEE Conference on Computer Vision and Pattern Recognition (CVPR),  
899 4895–4904. <https://doi.org/10.1109/CVPR.2017.520>.

900 Alonso, Alice, Rafael Muoz-Carpena, Robert E. Kennedy, and Carolina Murcia. 2016.  
901 "Wetland Landscape Spatio-Temporal Degradation Dynamics Using the New Google Earth  
902 Engine Cloud-Based Platform: Opportunities for Non-Specialists in Remote Sensing."  
903 Transactions of the ASABE 59 (5): 1331–42.

904 Andrieu, Julien. 2018. "Land Cover Changes on the West-African Coastline from the Saloum  
905 Delta (Senegal) to Rio Geba (Guinea-Bissau) between 1979 and 2015." European Journal of  
906 Remote Sensing 51 (1): 314–25.

907 Andrieu, Julien, Florent Lombard, Ababacar Fall, Mamadou Thior, Boubacar Demba Ba, and  
908 Barnabé Ephrem A. Dieme. 2020. “Botanical Field-Study and Remote Sensing to Describe  
909 Mangrove Resilience in the Saloum Delta (Senegal) after 30 Years of Degradation Narrative.”  
910 *Forest Ecology and Management* 461: 117963.

911 Andrieu, Julien. 2020. “Rice and Trees: Agrarian and Landscape Dynamics over 40 Years on  
912 the Coast of Southern Senegal and The Gambia.” *Dynamiques Environnementales. Journal*  
913 *International de Géosciences et de l’environnement*, no. 46: 2–23.

914 Arumugam, Manjula, Richard Niyomugabo, Farid Dahdouh-Guebas, and Jean Hugé. 2021.  
915 “The Perceptions of Stakeholders on Current Management of Mangroves in the Sine-Saloum  
916 Delta, Senegal.” *Estuarine, Coastal and Shelf Science* 248: 107160.  
917 <https://doi.org/10.1016/j.ecss.2020.107160>.

918 Asenso Barnieh, Beatrice, Li Jia, Massimo Menenti, Jie Zhou, and Yelong Zeng. 2020.  
919 “Mapping Land Use Land Cover Transitions at Different Spatiotemporal Scales in West  
920 Africa.” *Sustainability* 12 (20).

921 Balke, Thorsten, Edward L. Webb, Eva van den Elzen, Demis Galli, Peter M. J. Herman, and  
922 Tjeerd J. Bouma. 2013. “Seedling Establishment in a Dynamic Sedimentary Environment: A  
923 Conceptual Framework Using Mangroves.” *Journal of Applied Ecology* 50 (3): 740–47.  
924 <https://doi.org/10.1111/1365-2664.12067>.

925 Barousseau, J. P., E. H. S. Diop, and J. L. Saos. 1985. “Evidence of Dynamics Reversal in  
926 Tropical Estuaries, Geomorphological and Sedimentological Consequences (Salum and  
927 Casamance Rivers, Senegal).” *Sedimentology* 32 (4): 543–52. <https://doi.org/10.1111/j.1365-3091.1985.tb00469.x>.

929 Belgiu, Mariana, and Lucian Drăguț. 2016. “Random Forest in Remote Sensing: A Review of  
930 Applications and Future Directions.” *ISPRS Journal of Photogrammetry and Remote Sensing*  
931 114: 24–31.

932 Bivand, Roger, Micah Altman, Luc Anselin, Renato Assunção, Olaf Berke, Andrew Bernat,  
933 and Guillaume Blanchet. 2015. “Package ‘Spdep.’” *The Comprehensive R Archive Network*.

934 Blaschke, Thomas, Stefan Lang, and Geoffrey Hay. 2008. *Object-Based Image Analysis:  
935 Spatial Concepts for Knowledge-Driven Remote Sensing Applications*. Springer Science &  
936 Business Media.

937 Blaschke, Thomas. 2010. “Object Based Image Analysis for Remote Sensing.” *ISPRS Journal*  
938 *of Photogrammetry and Remote Sensing* 65 (1): 2–16.

939 Breiman, Leo. 2001. “Random Forests.” *Machine Learning* 45 (1): 5–32.

940 Bullock, Eric L., Sergio Fagherazzi, William Nardin, Phuoc Vo-Luong, Phong Nguyen, and  
941 Curtis E. Woodcock. 2017. “Temporal Patterns in Species Zonation in a Mangrove Forest in  
942 the Mekong Delta, Vietnam, Using a Time Series of Landsat Imagery.” *Continental Shelf*  
943 *Research* 147: 144–54.

944 Bullock, Eric L., Curtis E. Woodcock, and Pontus Olofsson. 2020. “Monitoring Tropical Forest  
945 Degradation Using Spectral Unmixing and Landsat Time Series Analysis.” *Remote Sensing of*  
946 *Environment* 238: 110968.

- 947 Bunting, Pete, Ake Rosenqvist, Richard M. Lucas, Lisa-Maria Rebelo, Lammert Hilarides,  
948 Nathan Thomas, Andy Hardy, Takuya Itoh, Masanobu Shimada, and C. Max Finlayson. 2018.  
949 "The Global Mangrove Watch—A New 2010 Global Baseline of Mangrove Extent" *Remote*  
950 *Sensing* 10, no. 10: 1669. <https://doi.org/10.3390/rs10101669>
- 951 Cabral, Ana I. R., and Fernando Lagos Costa. 2017. "Land Cover Changes and Landscape  
952 Pattern Dynamics in Senegal and Guinea Bissau Borderland." *Applied Geography* 82: 115–28.  
953 <https://doi.org/10.1016/j.apgeog.2017.03.010>.
- 954 Cárdenas, Nicolás Younes, Karen E. Joyce, and Stefan W. Maier. 2017. "Monitoring Mangrove  
955 Forests: Are We Taking Full Advantage of Technology?" *International Journal of Applied*  
956 *Earth Observation and Geoinformation* 63: 1–14.
- 957 Celis-Hernandez, Omar, Miguel Villoslada-Peciña, Raymond D. Ward, T.F. Bergamo, Rosela  
958 Perez-Ceballos, and María Patricia Girón-García. 2022. "Impacts of Environmental Pollution  
959 on Mangrove Phenology: Combining Remotely Sensed Data and Generalized Additive  
960 Models." *Science of The Total Environment* 810 (March): 152309.  
961 <https://doi.org/10.1016/j.scitotenv.2021.152309>.
- 962 Chen, Bangqian, Xiangming Xiao, Xiangping Li, Lianghao Pan, Russell Doughty, Jun Ma,  
963 Jinwei Dong, Yuanwei Qin, Bin Zhao, and Zhixiang Wu. 2017. "A Mangrove Forest Map of  
964 China in 2015: Analysis of Time Series Landsat 7/8 and Sentinel-1A Imagery in Google Earth  
965 Engine Cloud Computing Platform." *ISPRS Journal of Photogrammetry and Remote Sensing*  
966 131: 104–20.
- 967 Chen, Na. 2020. "Mapping Mangrove in Dongzhaigang, China Using Sentinel-2 Imagery."  
968 *Journal of Applied Remote Sensing* 14 (1): 014508.
- 969 Chen, Na. 2020. "Mapping Mangrove in Dongzhaigang, China Using Sentinel-2 Imagery."  
970 *Journal of Applied Remote Sensing* 14 (1): 014508.
- 971 Conchedda, Giulia, Laurent Durieux, and Philippe Mayaux. 2008. "An Object-Based Method  
972 for Mapping and Change Analysis in Mangrove Ecosystems." *Theme Issue: Remote Sensing*  
973 *of the Coastal Ecosystems* 63 (5): 578–89. <https://doi.org/10.1016/j.isprsjprs.2008.04.002>.
- 974 Conchedda, Giulia, Eric F. Lambin, and Philippe Mayaux. 2011. "Between land and sea:  
975 livelihoods and environmental changes in mangrove ecosystems of Senegal." *Annals of the*  
976 *Association of American Geographers* 101, no. 6 1259-1284.
- 977 Cormier-Salem, Marie-Christine. 1999. "Rivières du Sud." *Sociétés et mangroves ouest-*  
978 *africaines* 1: 2.
- 979 Cormier-Salem, Marie-Christine, and J. Panfili. 2016. "Mangrove reforestation: greening or  
980 grabbing coastal zones and deltas? Case studies in Senegal." *African Journal of Aquatic*  
981 *Science* 41, no. 1 :89-98.
- 982 Clinton, N. 2016. "Time Series Analysis in Earth Engine."
- 983 Jong, Steven M. de, Youchen Shen, Job de Vries, Ginny Bijnaar, Barend van Maanen, Pieter  
984 Augustinus, and Pita Verweij. 2021. "Mapping Mangrove Dynamics and Colonization Patterns  
985 at the Suriname Coast Using Historic Satellite Data and the LandTrendr Algorithm."

- 986 International Journal of Applied Earth Observation and Geoinformation 97 (May) : 102293.  
987 <https://doi.org/10.1016/j.jag.2020.102293>.
- 988 Descroix, Luc, Aïda Diongue Niang, Gérémy Panthou, Ansoumana Bodian, Youssouph Sane,  
989 Honoré Dacosta, Moussa Malam Abdou, Jean-Pierre Vandervaere, and Guillaume Quantin.  
990 2015. “Évolution Récente de La Pluviométrie En Afrique de l’ouest à Travers Deux Régions :  
991 La Sénégalie et Le Bassin Du Niger Moyen.” *Climatologie* 12: 25–43.  
992 <https://doi.org/10.4267/climatologie.1105>.
- 993 Luc Descroix. 2018. *Processus et enjeux d'eau en Afrique de l'Ouest soudano-sahélienne*. EAC,  
994 320 p.
- 995 Descroix, Luc, Yancouba Sané, Mamadou Thior, Sylvie-Paméla Manga, Boubacar D. Ba,  
996 Joseph Mingou, Victor Mendy, et al. 2020. “Inverse Estuaries in West Africa: Evidence of the  
997 Rainfall Recovery?” *Water* 12 (3). <https://doi.org/10.3390/w12030647>.
- 998 Díaz-Uriarte, Ramón, and Sara Alvarez de Andrés. 2006. "Gene selection and classification  
999 of microarray data using random forest." *BMC bioinformatics* 7, no. 1, 1-13.
- 1000 Balla Dieye, E. L., Amadou Tahirou Diaw, Tidiane Sané, and Ngor Ndour. 2013. "Dynamique  
1001 de la mangrove de l'estuaire du Saloum (Sénégal) entre 1972 et 2010." *Cybergeog: European*  
1002 *Journal of Geography*.
- 1003 Diop, E.S. 1990. *La côte Ouest-Africaine: du Saloum (Senegal). à la Mellacoree (Rep de*  
1004 *Guinee)*. Collection Études et Thésés, Éditions de l'ORSTOM, 2 volumes; Paris, 379 pp.
- 1005 Diop, E.S., A. Soumare, N. Diallo, and A. Guisse. 1997. “Recent Changes of the Mangroves  
1006 of the Saloum River Estuary, Senegal.” *Mangroves and Salt Marshes* 1 (3): 163–72.  
1007 <https://doi.org/10.1023/A:1009900724172>.
- 1008 Drăguț, Lucian, Dirk Tiede, and Shaun R. Levick. 2010. “ESP: A Tool to Estimate Scale  
1009 Parameter for Multiresolution Image Segmentation of Remotely Sensed Data.” *International*  
1010 *Journal of Geographical Information Science* 24 (6): 859–71.  
1011 <https://doi.org/10.1080/13658810903174803>.
- 1012 Drăguț, L., O. Csillik, C. Eisank, and D. Tiede. 2014. “Automated Parameterisation for Multi-  
1013 Scale Image Segmentation on Multiple Layers.” *ISPRS Journal of Photogrammetry and*  
1014 *Remote Sensing* 88 (February): 119–27. <https://doi.org/10.1016/j.isprsjprs.2013.11.018>.
- 1015 Dronova, Iryna. 2015. “Object-Based Image Analysis in Wetland Research: A Review.”  
1016 *Remote Sensing* 7 (5): 6380–6413. <https://doi.org/10.3390/rs70506380>.
- 1017 Espindola, G. M., G. Camara, I. A. Reis, L. S. Bins, and A. M. Monteiro. 2006. “Parameter  
1018 Selection for Region-growing Image Segmentation Algorithms Using Spatial  
1019 Autocorrelation.” *International Journal of Remote Sensing* 27 (14): 3035–40.  
1020 <https://doi.org/10.1080/01431160600617194>.
- 1021 Fent, Ashley, Rémi Bardou, Judith Carney, and Kyle Cavanaugh. 2019. “Transborder Political  
1022 Ecology of Mangroves in Senegal and The Gambia.” *Global Environmental Change* 54  
1023 (January): 214–26.

- 1024 Francisco Flores De Santiago, John M. Kovacs & Patrick Lafrance. 2013. An object-oriented  
1025 classification method for mapping mangroves in Guinea, West Africa, using multipolarized  
1026 ALOS PALSAR L-band data, *International Journal of Remote Sensing*, 34:2, 563-586, DOI:  
1027 10.1080/01431161.2012.715773
- 1028 Gao, Bo-cai. 1996. “NDWI—A Normalized Difference Water Index for Remote Sensing of  
1029 Vegetation Liquid Water from Space.” *Remote Sensing of Environment* 58 (3): 257–66.  
1030 [https://doi.org/10.1016/S0034-4257\(96\)00067-3](https://doi.org/10.1016/S0034-4257(96)00067-3).
- 1031 George, Rajee, Hitendra Padalia, S. K. Sinha, and A. Senthil Kumar. 2020. “Evaluating  
1032 Sensitivity of Hyperspectral Indices for Estimating Mangrove Chlorophyll in Middle Andaman  
1033 Island, India.” *Environmental Monitoring and Assessment* 191 (3): 785.  
1034 <https://doi.org/10.1007/s10661-019-7679-6>.
- 1035 Ghazali, Mochamad Firman, and Ketut Wikantika. 2021. “Pre-Assessment of the Potential of  
1036 Dual Polarization of Sentinel 1 Data for Mapping the Mangrove Tree Species Distribution in  
1037 South Bali, Indonesia.” In 2021 7th Asia-Pacific Conference on Synthetic Aperture Radar  
1038 (APSAR), 1–6. <https://doi.org/10.1109/APSAR52370.2021.9688441>.
- 1039 Giri, C., E. Ochieng, L. L. Tieszen, Z. Zhu, A. Singh, T. Loveland, J. Masek, and N. Duke.  
1040 2011. “Status and Distribution of Mangrove Forests of the World Using Earth Observation  
1041 Satellite Data.” *Global Ecology and Biogeography* 20 (1): 154–59.  
1042 <https://doi.org/10.1111/j.1466-8238.2010.00584.x>.
- 1043 Giri, Chandra, Jordan Long, Sawaid Abbas, R.Mani Murali, Faisal M. Qamer, Bruce Pengra,  
1044 and David Thau. 2015. “Distribution and Dynamics of Mangrove Forests of South Asia.” *Land  
1045 Cover/Land Use Change (LC/LUC) and Environmental Impacts in South Asia* 148 (January):  
1046 101–11. <https://doi.org/10.1016/j.jenvman.2014.01.020>.
- 1047 Gorelick, Noel, Matt Hancher, Mike Dixon, Simon Ilyushchenko, David Thau, and Rebecca  
1048 Moore. 2017. “Google Earth Engine : Planetary-Scale Geospatial Analysis for Everyone.” *Big  
1049 Remotely Sensed Data : Tools, Applications and 39xpériences* 202 (December) : 18–27.  
1050 <https://doi.org/10.1016/j.rse.2017.06.031>.
- 1051 Gudex-Cross, David, Jennifer Pontius, and Alison Adams. 2017. “Enhanced Forest Cover  
1052 Mapping Using Spectral Unmixing and Object-Based Classification of Multi-Temporal  
1053 Landsat Imagery.” *Remote Sensing of Environment* 196 (July): 193–204.
- 1054 Gupta, Kaushik, Anirban Mukhopadhyay, Sandip Giri, Abhra Chanda, Sayani Datta  
1055 Majumdar, Sourav Samanta, Debasish Mitra, Rabindro N. Samal, Ajit K. Pattnaik, and Sugata  
1056 Hazra. 2018. “An Index for Discrimination of Mangroves from Non-Mangroves Using  
1057 LANDSAT 8 OLI Imagery.” *MethodsX* 5 (January): 1129–39.  
1058 <https://doi.org/10.1016/j.mex.2018.09.011>.
- 1059 Han, Xingxing, Lian Feng, Chuanmin Hu, and Philip Kramer. 2018. “Hurricane-Induced  
1060 Changes in the Everglades National Park Mangrove Forest: Landsat Observations Between  
1061 1985 and 2017.” *Journal of Geophysical Research: Biogeosciences* 123 (11): 3470–88.  
1062 <https://doi.org/10.1029/2018JG004501>.



- 1063 Haralick, Robert M., K. Shanmugam, and Its' Hak Dinstein. 1973. "Textural Features for Image  
1064 Classification." *IEEE Transactions on Systems, Man, and Cybernetics SMC-3* (6): 610–21.  
1065 <https://doi.org/10.1109/TSMC.1973.4309314>.
- 1066 Hossain, Mohammad D., and Dongmei Chen. 2019. "Segmentation for Object-Based Image  
1067 Analysis (OBIA): A Review of Algorithms and Challenges from Remote Sensing Perspective."  
1068 *ISPRS Journal of Photogrammetry and Remote Sensing* 150 (April): 115–34.  
1069 <https://doi.org/10.1016/j.isprsjprs.2019.02.009>.
- 1070 Hu, Qiong, Damien Sulla-Menashe, Baodong Xu, He Yin, Huajun Tang, Peng Yang, and  
1071 Wenbin Wu. 2019. "A Phenology-Based Spectral and Temporal Feature Selection Method for  
1072 Crop Mapping from Satellite Time Series." *International Journal of Applied Earth Observation*  
1073 *and Geoinformation* 80: 218–29. <https://doi.org/10.1016/j.jag.2019.04.014>.
- 1074 Jakubauskas, Mark E., David R. Legates, and Jude H. Kastens. 2002. "Crop Identification  
1075 Using Harmonic Analysis of Time-Series AVHRR NDVI Data." *Computers and Electronics*  
1076 *in Agriculture* 37 (1): 127–39. [https://doi.org/10.1016/S0168-1699\(02\)00116-3](https://doi.org/10.1016/S0168-1699(02)00116-3).
- 1077 Jia, Mingming, Zongming Wang, Yuanzhi Zhang, Dehua Mao, and Chao Wang. 2018.  
1078 "Monitoring Loss and Recovery of Mangrove Forests during 42 Years: The Achievements of  
1079 Mangrove Conservation in China." *International Journal of Applied Earth Observation and*  
1080 *Geoinformation* 73 (December): 535–45. <https://doi.org/10.1016/j.jag.2018.07.025>.
- 1081 Jia, Mingming, Zongming Wang, Dehua Mao, Chunying Ren, Chao Wang, and Yeqiao Wang.  
1082 2021. "Rapid, Robust, and Automated Mapping of Tidal Flats in China Using Time Series  
1083 Sentinel-2 Images and Google Earth Engine." *Remote Sensing of Environment* 255 (March):  
1084 112285. <https://doi.org/10.1016/j.rse.2021.112285>.
- 1085 Jia, Mingming, Zongming Wang, Dehua Mao, Chunying Ren, Chao Wang, and Yeqiao Wang.  
1086 2021. "Rapid, Robust, and Automated Mapping of Tidal Flats in China Using Time Series  
1087 Sentinel-2 Images and Google Earth Engine." *Remote Sensing of Environment* 255 (March):  
1088 112285. <https://doi.org/10.1016/j.rse.2021.112285>.
- 1089 Johnson, Brian, and Zhixiao Xie. 2011. "Unsupervised Image Segmentation Evaluation and  
1090 Refinement Using a Multi-Scale Approach." *ISPRS Journal of Photogrammetry and Remote*  
1091 *Sensing* 66 (4): 473–83. <https://doi.org/10.1016/j.isprsjprs.2011.02.006>.
- 1092 Johnson, Brian A., Milben Bragais, Isao Endo, Damasa B. Magcale-Macandog, and Paula  
1093 Beatrice M. Macandog. 2015. "Image Segmentation Parameter Optimization Considering  
1094 Within- and Between-Segment Heterogeneity at Multiple Scale Levels: Test Case for Mapping  
1095 Residential Areas Using Landsat Imagery." *ISPRS International Journal of Geo-Information* 4  
1096 (4) .
- 1097 Kennedy, Robert E., Zhiqiang Yang, and Warren B. Cohen. 2010. "Detecting Trends in Forest  
1098 Disturbance and Recovery Using Yearly Landsat Time Series : 1. LandTrendr — Temporal  
1099 Segmentation Algorithms." *Remote Sensing of Environment* 114 (12) : 2897–2910.
- 1100 Kuenzer, Claudia, Andrea Bluemel, Steffen Gebhardt, Tuan Vo Quoc, and Stefan Dech. 2011.  
1101 "Remote Sensing of Mangrove Ecosystems: A Review" *Remote Sensing* 3, no. 5: 878-928.  
1102 <https://doi.org/10.3390/rs3050878>

- 1103 Krauss, Ken W., Catherine E. Lovelock, Karen L. McKee, Laura López-Hoffman, Sharon M.L.  
 1104 Ewe, and Wayne P. Sousa. 2008. “Environmental Drivers in Mangrove Establishment and  
 1105 Early Development: A Review.” *Mangrove Ecology – Applications in Forestry and Coastal*  
 1106 *Zone Management* 89 (2): 105–27. <https://doi.org/10.1016/j.aquabot.2007.12.014>.
- 1107 Li, Xianju, Weitao Chen, Xinwen Cheng, and Lizhe Wang. 2016. “A Comparison of Machine  
 1108 Learning Algorithms for Mapping of Complex Surface-Mined and Agricultural Landscapes  
 1109 Using ZiYuan-3 Stereo Satellite Imagery.” *Remote Sensing* 8 (6).  
 1110 <https://doi.org/10.3390/rs8060514>.
- 1111 Li, Huiying, Mingming Jia, Rong Zhang, Yongxing Ren, and Xin Wen. 2019. “Incorporating  
 1112 the Plant Phenological Trajectory into Mangrove Species Mapping with Dense Time Series  
 1113 Sentinel-2 Imagery and the Google Earth Engine Platform.” *Remote Sensing* 11 (21).  
 1114 <https://doi.org/10.3390/rs11212479>.
- 1115 Liu, Xue, Temilola E. Fatoyinbo, Nathan M. Thomas, Weihe Wendy Guan, Yanni Zhan, Pinki  
 1116 Mondal, David Lagomasino, et al. 2021. “Large-Scale High-Resolution Coastal Mangrove  
 1117 Forests Mapping Across West Africa With Machine Learning Ensemble and Satellite Big  
 1118 Data.” *Frontiers in Earth Science* 8.  
 1119 <https://www.frontiersin.org/articles/10.3389/feart.2020.560933>.
- 1120 Lu, Ying, and Le Wang. 2021. “How to Automate Timely Large-Scale Mangrove Mapping  
 1121 with Remote Sensing.” *Remote Sensing of Environment* 264 (October): 112584.  
 1122 <https://doi.org/10.1016/j.rse.2021.112584>.
- 1123 Lombard, Florent, and Andrieu Julien. 2021. “Mapping Mangrove Zonation Changes in  
 1124 Senegal with Landsat Imagery Using an OBIA Approach Combined with Linear Spectral  
 1125 Unmixing.” *Remote Sensing* 13 (10). <https://doi.org/10.3390/rs13101961>.
- 1126 Lombard, Florent, Andrieu Julien, and Descroix Luc. 2021. “La Population d’Avicennia  
 1127 Germinans Du Delta Du Saloum Est-Elle Relictuelle Depuis La Dernière Période Humide ?”  
 1128 *BOIS & FORETS DES TROPIQUES* 346 (0): 51–64.  
 1129 <https://doi.org/10.19182/bft2020.346.a36296>.
- 1130 Lymburner, Leo, Peter Bunting, Richard Lucas, Peter Scarth, Imam Alam, Claire Phillips,  
 1131 Catherine Ticehurst, and Alex Held. 2020. “Mapping the Multi-Decadal Mangrove Dynamics  
 1132 of the Australian Coastline.” *Time Series Analysis with High Spatial Resolution Imagery* 238  
 1133 (March): 111185. <https://doi.org/10.1016/j.rse.2019.05.004>.
- 1134 Ma, Lei, Manchun Li, Xiaoxue Ma, Liang Cheng, Peijun Du, and Yongxue Liu. 2017. “A  
 1135 Review of Supervised Object-Based Land-Cover Image Classification.” *ISPRS Journal of*  
 1136 *Photogrammetry and Remote Sensing* 130: 277–93.  
 1137 <https://doi.org/10.1016/j.isprsjprs.2017.06.001>.
- 1138 Marius, Claude. 1985. *Mangroves du Sénégal et de la Gambie : écologie, pédologie, géochimie,*  
 1139 *mise en valeur et aménagement. Travaux et Documents de l’ORSTOM. Paris: ORSTOM.*  
 1140 <https://www.documentation.ird.fr/hor/fdi:19338>.
- 1141 McFEETERS, S. K. 1996. “The Use of the Normalized Difference Water Index (NDWI) in the  
 1142 Delineation of Open Water Features.” *International Journal of Remote Sensing* 17 (7): 1425–  
 1143 32. <https://doi.org/10.1080/01431169608948714>.

- 1144 Midekisa, Alemayehu, Felix Holl, David J. Savory, Ricardo Andrade-Pacheco, Peter W.  
1145 Gething, Adam Bennett, and Hugh J. W. Sturrock. 2017. "Mapping Land Cover Change over  
1146 Continental Africa Using Landsat and Google Earth Engine Cloud Computing." PLOS ONE  
1147 12 (9): e0184926. <https://doi.org/10.1371/journal.pone.0184926>.
- 1148 Mondal, Pinki, Xue Liu, Temilola E. Fatoyinbo, and David Lagomasino. 2019. "Evaluating  
1149 Combinations of Sentinel-2 Data and Machine-Learning Algorithms for Mangrove Mapping  
1150 in West Africa." Remote Sensing 11 (24). <https://doi.org/10.3390/rs11242928>.
- 1151 Monsef, Hesham Abd-El, and Scot E. Smith. 2017. "A New Approach for Estimating  
1152 Mangrove Canopy Cover Using Landsat 8 Imagery." Computers and Electronics in Agriculture  
1153 135 (April): 183–94. <https://doi.org/10.1016/j.compag.2017.02.007>.
- 1154 Mukherjee, Nibedita, William J. Sutherland, Lynn Dicks, Jean Hugé, Nico Koedam, and Farid  
1155 Dahdouh-Guebas. 2014. "Ecosystem Service Valuations of Mangrove Ecosystems to Inform  
1156 Decision Making and Future Valuation Exercises." PLOS ONE 9 (9): e107706.  
1157 <https://doi.org/10.1371/journal.pone.0107706>.
- 1158 Mutanga, Onesimo, and Lalit Kumar. 2019. "Google Earth Engine Applications." Remote  
1159 Sensing 11 (5). <https://doi.org/10.3390/rs11050591>.
- 1160 Myint, Soe W., Chandra P. Giri, Le Wang, Zhiliang Zhu, and Shana C. Gillette. 2008.  
1161 "Identifying Mangrove Species and Their Surrounding Land Use and Land Cover Classes  
1162 Using an Object-Oriented Approach with a Lacunarity Spatial Measure." GIScience & Remote  
1163 Sensing 45 (2): 188–208. <https://doi.org/10.2747/1548-1603.45.2.188>.
- 1164 Myint, Soe W., Patricia Gober, Anthony Brazel, Susanne Grossman-Clarke, and Qihao Weng.  
1165 2011. "Per-Pixel vs. Object-Based Classification of Urban Land Cover Extraction Using High  
1166 Spatial Resolution Imagery." Remote Sensing of Environment 115 (5): 1145–61.  
1167 <https://doi.org/10.1016/j.rse.2010.12.017>.
- 1168 Ndour, Ngor, Sara Danièle Dieng, and Mamadou Fall. 2012. "Rôles Des Mangroves, Modes  
1169 et Perspectives de Gestion Au Delta Du Saloum (Sénégal)." Vertigo - la revue électronique en  
1170 sciences de l'environnement, Volume 11 Numéro 3 | décembre 2011. DOI :  
1171 <https://doi.org/10.4000/vertigo.11515>
- 1172 Nicholson, Sharon E. 2013. "The West African Sahel: A Review of Recent Studies on the  
1173 Rainfall Regime and Its Interannual Variability." Edited by D.-Y. Wang, I. Bordi, and F. Acs.  
1174 ISRN Meteorology 2013 (February): 453521. <https://doi.org/10.1155/2013/453521>.
- 1175 Olofsson, Pontus, Giles M. Foody, Martin Herold, Stephen V. Stehman, Curtis E. Woodcock,  
1176 and Michael A. Wulder. 2014. "Good Practices for Estimating Area and Assessing Accuracy  
1177 of Land Change." Remote Sensing of Environment 148: 42–57.  
1178 <https://doi.org/10.1016/j.rse.2014.02.015>.
- 1179 Pasquarella, Valerie J., Christopher E. Holden, and Curtis E. Woodcock. 2018. "Improved  
1180 Mapping of Forest Type Using Spectral-Temporal Landsat Features." Remote Sensing of  
1181 Environment 210: 193–207. <https://doi.org/10.1016/j.rse.2018.02.064>.
- 1182 Pasquarella, Valerie J., Paulo Arévalo, Kelsee H. Bratley, Eric L. Bullock, Noel Gorelick,  
1183 Zhiqiang Yang, and Robert E. Kennedy. 2022. "Demystifying LandTrendr and CCDC

- 1184 Temporal Segmentation.” *International Journal of Applied Earth Observation and*  
1185 *Geoinformation* 110 (June): 102806. <https://doi.org/10.1016/j.jag.2022.102806>.
- 1186 Pastor-Guzman, J., Jadunandan Dash, and Peter M. Atkinson. 2018. “Remote Sensing of  
1187 Mangrove Forest Phenology and Its Environmental Drivers.” *Remote Sensing of Environment*  
1188 205 (February): 71–84. <https://doi.org/10.1016/j.rse.2017.11.009>.
- 1189 Pirasteh, Saied, Eric K. Zenner, Davood Mafi-Gholami, Abolfazl Jaafari, Akram Nouri  
1190 Kamari, Guoxiang Liu, Qing Zhu, and Jonathan Li. 2021. “Modeling Mangrove Responses to  
1191 Multi-Decadal Climate Change and Anthropogenic Impacts Using a Long-Term Time Series  
1192 of Satellite Imagery.” *International Journal of Applied Earth Observation and Geoinformation*  
1193 102: 102390.
- 1194 Rasul, Azad, Heiko Balzter, Gaylan R. Faqe Ibrahim, Hasan M. Hameed, James Wheeler,  
1195 Bashir Adamu, Sa’ad Ibrahim, and Peshawa M. Najmaddin. 2018. "Applying Built-Up and  
1196 Bare-Soil Indices from Landsat 8 to Cities in Dry Climates" *Land* 7, no. 3: 81.  
1197 <https://doi.org/10.3390/land7030081>
- 1198 Rouse, Jr., J. W., R. H. Haas, J. A. Schell, and D. W. Deering. 1974. “Monitoring Vegetation  
1199 Systems in the Great Plains with ERTS.” In *NASA Special Publication*, 351:309.
- 1200 Schulz, Dario, He Yin, Bernhard Tischbein, Sarah Verleysdonk, Rabani Adamou, and Navneet  
1201 Kumar. 2021. “Land Use Mapping Using Sentinel-1 and Sentinel-2 Time Series in a  
1202 Heterogeneous Landscape in Niger, Sahel.” *ISPRS Journal of Photogrammetry and Remote*  
1203 *Sensing* 178: 97–111. <https://doi.org/10.1016/j.isprsjprs.2021.06.005>.
- 1204 Sow, M., Diallo, A., Diallo, N., Dixon, C. A., & Guissé, A. 1991. Formations végétales et sols  
1205 dans les mangroves des Rivières du Sud. In Cormier-Salem, M. (Ed.), *Dynamique et usages de*  
1206 *la mangrove dans les pays des rivières du Sud, du Sénégal à la Sierra Leone*. IRD Éditions.  
1207 doi:10.4000/books.irdeditions.3690
- 1208 Speiser, Jaime Lynn, Michael E. Miller, Janet Tooze, and Edward Ip. 2019. “A Comparison of  
1209 Random Forest Variable Selection Methods for Classification Prediction Modeling.” *Expert*  
1210 *Systems with Applications* 134: 93–101. <https://doi.org/10.1016/j.eswa.2019.05.028>.
- 1211 Stehman, Stephen V. 1997. “Selecting and Interpreting Measures of Thematic Classification  
1212 Accuracy.” *Remote Sensing of Environment* 62 (1): 77–89. [https://doi.org/10.1016/S0034-](https://doi.org/10.1016/S0034-4257(97)00083-7)  
1213 [4257\(97\)00083-7](https://doi.org/10.1016/S0034-4257(97)00083-7).
- 1214 Stromann, Oliver, Andrea Nascetti, Osama Yousif, and Yifang Ban. 2020. “Dimensionality  
1215 Reduction and Feature Selection for Object-Based Land Cover Classification Based on  
1216 Sentinel-1 and Sentinel-2 Time Series Using Google Earth Engine.” *Remote Sensing* 12 (1).  
1217 <https://doi.org/10.3390/rs12010076>.
- 1218 Taureau, Florent, Marc Robin, Christophe Proisy, François Fromard, Daniel Imbert, and  
1219 Françoise Debaine. 2019. “Mapping the Mangrove Forest Canopy Using Spectral Unmixing  
1220 of Very High Spatial Resolution Satellite Images.” *Remote Sensing* 11 (3).  
1221 <https://doi.org/10.3390/rs11030367>.
- 1222 Temudo, Marina Padrão, Rui Figueira, and Manuel Abrantes. 2015. “Landscapes of Bio-  
1223 Cultural Diversity: Shifting Cultivation in Guinea-Bissau, West Africa.” *Agroforestry Systems*  
1224 89 (1): 175–91. <https://doi.org/10.1007/s10457-014-9752-z>.

- 1225 Valderrama-Landeros, L., F. Flores-de-Santiago, J. M. Kovacs, and F. Flores-Verdugo. 2018.  
 1226 “An Assessment of Commonly Employed Satellite-Based Remote Sensors for Mapping  
 1227 Mangrove Species in Mexico Using an NDVI-Based Classification Scheme.” *Environmental*  
 1228 *Monitoring and Assessment* 190 (1): 23. <https://doi.org/10.1007/s10661-017-6399-z>.
- 1229 Valderrama-Landeros, Luis, Francisco Flores-Verdugo, Ranulfo Rodríguez-Sobreyra, John M.  
 1230 Kovacs, and Francisco Flores-de-Santiago. 2021. “Extrapolating Canopy Phenology  
 1231 Information Using Sentinel-2 Data and the Google Earth Engine Platform to Identify the  
 1232 Optimal Dates for Remotely Sensed Image Acquisition of Semiarid Mangroves.” *Journal of*  
 1233 *Environmental Management* 279 (February): 111617.  
 1234 <https://doi.org/10.1016/j.jenvman.2020.111617>.
- 1235 Vizcaya-Martínez, Diego Arturo, Francisco Flores-de-Santiago, Luis Valderrama-Landeros,  
 1236 David Serrano, Ranulfo Rodríguez-Sobreyra, León Felipe Álvarez-Sánchez, and Francisco  
 1237 Flores-Verdugo. 2022. “Monitoring Detailed Mangrove Hurricane Damage and Early  
 1238 Recovery Using Multisource Remote Sensing Data.” *Journal of Environmental Management*  
 1239 320 (October): 115830. <https://doi.org/10.1016/j.jenvman.2022.115830>.
- 1240 Walters, Bradley B., Patrik Rönnbäck, John M. Kovacs, Beatrice Crona, Syed Ainul Hussain,  
 1241 Ruchi Badola, Jurgenne H. Primavera, Edward Barbier, and Farid Dahdouh-Guebas. 2008.  
 1242 “Ethnobiology, Socio-Economics and Management of Mangrove Forests: A Review.”  
 1243 *Mangrove Ecology – Applications in Forestry and Coastal Zone Management* 89 (2): 220–36.  
 1244 <https://doi.org/10.1016/j.aquabot.2008.02.009>.
- 1245 Wang, Dezhi, Bo Wan, Penghua Qiu, Yanjun Su, Qinghua Guo, Run Wang, Fei Sun, and  
 1246 Xincui Wu. 2018. “Evaluating the Performance of Sentinel-2, Landsat 8 and Pléiades-1 in  
 1247 Mapping Mangrove Extent and Species.” *Remote Sensing* 10 (9).  
 1248 <https://doi.org/10.3390/rs10091468>.
- 1249 Wang, Le, Mingming Jia, Dameng Yin, and Jinyan Tian. 2019. “A Review of Remote Sensing  
 1250 for Mangrove Forests: 1956–2018.” *Remote Sensing of Environment* 231 (September):  
 1251 111223. <https://doi.org/10.1016/j.rse.2019.111223>.
- 1252 Wang, Xiaona, Le Wang, Jinyan Tian, and Chen Shi. 2021. “Object-Based Spectral-  
 1253 Phenological Features for Mapping Invasive *Spartina Alterniflora*.” *International Journal of*  
 1254 *Applied Earth Observation and Geoinformation* 101: 102349.  
 1255 <https://doi.org/10.1016/j.jag.2021.102349>.
- 1256 Xiao, Han, Fenzhen Su, Dongjie Fu, Vincent Lyne, Gaohuan Liu, Tingting Pan, and Jiakun  
 1257 Teng. 2021. “Optimal and Robust Vegetation Mapping in Complex Environments Using  
 1258 Multiple Satellite Imagery: Application to Mangroves in Southeast Asia.” *International Journal*  
 1259 *of Applied Earth Observation and Geoinformation* 99: 102320.  
 1260 <https://doi.org/10.1016/j.jag.2021.102320>.
- 1261 Zanaga, Daniele, Ruben Van De Kerchove, Wanda De Keersmaecker, Niels Souverijns,  
 1262 Carsten Brockmann, Ralf Quast, Jan Wevers, et al. 2021. “ESA WorldCover 10 m 2020 V100.”  
 1263 VITO. <https://edepot.wur.nl/576022>.
- 1264 Zhang, Xuehong, Paul M. Treitz, Dongmei Chen, Chang Quan, Lixin Shi, and Xinhui Li. 2017.  
 1265 “Mapping Mangrove Forests Using Multi-Tidal Remotely-Sensed Data and a Decision-Tree-

- 1266 Based Procedure.” *International Journal of Applied Earth Observation and Geoinformation* 62  
1267 (October): 201–14. <https://doi.org/10.1016/j.jag.2017.06.010>.
- 1268 Zhao, Chuanpeng, and Cheng-Zhi Qin. 2020. “10-m-Resolution Mangrove Maps of China  
1269 Derived from Multi-Source and Multi-Temporal Satellite Observations.” *ISPRS Journal of*  
1270 *Photogrammetry and Remote Sensing* 169 (November): 389–405.  
1271 <https://doi.org/10.1016/j.isprsjprs.2020.10.001>.
- 1272 Zhu, Zhe, and Curtis E. Woodcock. 2014. “Continuous Change Detection and Classification  
1273 of Land Cover Using All Available Landsat Data.” *Remote Sensing of Environment* 144: 152–  
1274 71. <https://doi.org/10.1016/j.rse.2014.01.011>.
- 1275