



HAL
open science

Natural photosynthetic microboring communities produce alkalinity in seawater whereas aragonite saturation state rises up to five

Aline Tribollet, Anne Chauvin, Pascale Cuet

► To cite this version:

Aline Tribollet, Anne Chauvin, Pascale Cuet. Natural photosynthetic microboring communities produce alkalinity in seawater whereas aragonite saturation state rises up to five. *Frontiers in Earth Science*, 2022, 10, p. 894501 [12 p.]. 10.3389/feart.2022.894501 . hal-03993602

HAL Id: hal-03993602

<https://hal.science/hal-03993602v1>

Submitted on 21 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY
Ana Santos,
University of Oviedo, Spain

REVIEWED BY
Chris Langdon,
University of Miami, United States
Francisco J. Rodriguez-Tovar,
University of Graanada, Spain

*CORRESPONDENCE
Aline Tribollet,
✉ aline.tribollet@ird.fr

SPECIALTY SECTION
This article was submitted to
Biogeoscience,
a section of the journal
Frontiers in Earth Science

RECEIVED 11 March 2022
ACCEPTED 25 November 2022
PUBLISHED 21 December 2022

CITATION
Tribollet A, Chauvin A and Cuet P (2022),
Natural photosynthetic microboring
communities produce alkalinity in
seawater whereas aragonite saturation
state rises up to five.
Front. Earth Sci. 10:894501.
doi: 10.3389/feart.2022.894501

COPYRIGHT
© 2022 Tribollet, Chauvin and Cuet. This
is an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).
The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Natural photosynthetic microboring communities produce alkalinity in seawater whereas aragonite saturation state rises up to five

Aline Tribollet^{1,2*}, Anne Chauvin¹ and Pascale Cuet³

¹IRD-Sorbonne University-CNRS-MNH, UMR LOCEAN-IPSL, Paris, France, ²HIMB, SOEST-University of Hawaii, Kaneohe, HI, United States, ³UMR ENTROPIE (Université de La Réunion-IRD-CNRS-Ifremer-Université de la Nouvelle Calédonie)—Labex Corail, Université de La Réunion, Réunion, France

Bioerosion, resulting from microbioerosion or biogenic dissolution, macrobioerosion and grazing, is one the main processes involved in reef carbonate budget and functioning. On healthy reefs, most of the produced carbonates are preserved and accumulate. But in the context of global change, reefs are increasingly degraded as environmental factors such as ocean warming and acidification affect negatively reef accretion and positively bioerosion processes. The recent 2019 SROCC report suggests that if CO₂ emissions in the atmosphere are not drastically reduced rapidly, 70%–99% of coral reefs will disappear by 2,100. However, to improve projections of coral reef evolution, it is important to better understand dynamics of bioerosion processes. Among those processes, it was shown recently that bioeroding microflora which actively colonize and dissolve experimental coral blocks, release significant amount of alkalinity in seawater both by day and at night under controlled conditions. It was also shown that this alkalinity production is enhanced under ocean acidification conditions (saturation state of aragonite comprised between 2 and 3.5) suggesting that reef carbonate accumulation will be even more limited in the future. To better understand the conditions of production of alkalinity in seawater by boring microflora and its possible consequences on reef resilience, we conducted a series of experiments with natural rubble maintained under natural or artificial light, and various saturation states of aragonite. We show here that biogenic dissolution of natural reef rubble colonized by microboring communities dominated by the chlorophyte *Ostreobium* sp., and thus the production of alkalinity in seawater, can occur under a large range of saturation states of aragonite, from 2 to 6.4 under daylight and that this production is directly correlated to the photosynthetic activity of microboring communities. We then discuss the possible implications of such paradoxical activities on reef resilience.

KEYWORDS

biogenic carbonate dissolution, microboring flora, euendoliths, production of seawater alkalinity, saturation state of aragonite, coral reef ecosystems

1 Introduction

Due to the exponentially rising atmospheric carbon dioxide partial pressure ($p\text{CO}_2$) and its partial absorption by the ocean ($\sim 30\%$, Sabine et al., 2004), the saturation state of surface seawater (Ω) with respect to calcium carbonate minerals (CaCO_3) will decrease together with seawater pH by the end of the century (-0.2 to -0.4 pH unit depending on the IPCC scenario and seasonality; Orr et al., 2005; Bindoff et al., 2019). Such decrease will greatly impact negatively major calcifying organisms and some coastal carbonate ecosystems (Guinotte and Fabry, 2008; Fabricius et al., 2011; Agostini et al., 2018) while enhancing carbonate dissolution (Andersson et al., 2007; Andersson et al., 2008; Krumins et al., 2013; Stubler and Peterson, 2016; Schönberg et al., 2017).

Cyronak et al. (2014) showed that the average $p\text{CO}_2$ may have increased faster in coral reefs than in the atmosphere and the open ocean over the past 2 decades (~ 3.5 -fold, i.e., $+6.6 \pm 1.4 \mu\text{atm.y}^{-1}$) due to additional local disturbances resulting from human activities (e.g., eutrophication), thus putting these ecosystems even more at risk under ocean acidification. Notwithstanding the fact that reef ecosystems will greatly be degraded by the end of the century due to this factor but also ocean warming and local disturbances (e.g., storm impacts and rising runoffs), the latest SROCC report (Bindoff et al., 2019) highlighted the lack of information regarding the sensitivity and adaptive capacity of coral reef organisms and ecosystems to climate change impacts inducing bias in projections.

While several studies highlighted the negative effect of ocean acidification (combined or not with other factors) on growth, abundance and calcification rates of the main reef framebuilders, i.e., corals and calcifying algae under more or less controlled conditions and over short term (e.g., Langdon and Atkinson, 2005; Kuffner et al., 2008; Pandolfi et al., 2011; Comeau et al., 2014; Johnson et al., 2014), at the global reef scale the relationship between net reef community calcification and aragonite saturation state varies greatly from one area to the next (e.g., Shamberger et al., 2011; Falter et al., 2012; Shaw et al., 2012; Page et al., 2016). Such variations may be explained by local biological adaptations of calcifiers to the natural variability of their environmental conditions (Vargas et al., 2022) and/or variability of carbonate dissolution processes in both reef sediments and hard substrates (Pandolfi et al., 2011; see summary Table 2 in Tribollet et al., 2019); reef budget depending on the equilibrium between reef calcification and dissolution. Eyre et al. (2014) showed that despite possible adaptations of calcifiers, changes in reef dissolution are more rapid than changes in reef calcification under ocean acidification, enhancing net reef dissolution (see also Andersson and Gledhill 2013).

Reef dissolution comprises the thermodynamically driven carbonate dissolution (seawater chemistry with $\Omega < 1$), the bacteria driven dissolution (release of CO_2 through organic matter remineralization) and the biogenic carbonate dissolution driven mainly by boring microflora and sponges (Tribollet and Golubic, 2011; Schönberg et al., 2017). Among those processes, the

spatial and temporal variability of biogenic dissolution rates due to microboring flora remains poorly known (Carreiro-Silva et al., 2005; Tribollet, 2008b; Grange et al., 2015; Schönberg et al., 2017), most probably because this process which occurs at the microscale is considered as negligible, or is ignored and/or too difficult to study. However, this process may explain an important part of the reef dissolution variability (Andersson and Gledhill 2013). Tribollet et al. (2019) showed indeed that biogenic dissolution due to boring microflora dissolve up to 20% of carbonates deposited by reef calcifiers under ambient conditions (dissolution estimated on average at $11 \text{ mmol CaCO}_3.\text{m}^{-2}$ of reef.d⁻¹ in both hard and sediment substrates *via* microscopy techniques). They also highlighted that mature microboring communities dominated by the chlorophyte *Ostreobium* sp. colonizing experimental blocks of dead coral skeleton can produce consequently, significant amount of seawater alkalinity both under low controlled light intensity (during photosynthesis) and at night, at ambient saturation states (at Ω_{Arag} 3–3.5). They finally showed that this production is enhanced by 50% under ocean acidification conditions (at $\Omega_{\text{Arag}} = 2$), confirming trends obtained for biogenic dissolution rates in dead corals measured by microscopy or buoyant weight by Tribollet et al. (2009), Reyes-Nivia et al. (2013) and Enochs et al. (2016). Meanwhile, Tribollet et al. (2019) suggested a potential negative feedback of this biogenic dissolution on ocean acidification; a feedback that could benefit to reef calcifiers and ecosystems, at least at the microscale. Thus, to better understand dynamics of the biogenic dissolution process resulting from microborers' metabolic activity and possible implications on reef resilience, it is necessary to further investigate the relationship between the production of seawater alkalinity by microborers and the saturation state of aragonite under natural conditions and at various spatial (from substrate to ecosystem scale) and temporal scales (day, season, year).

Here we studied the capacity of natural mature microboring communities dominated by *Ostreobium* sp. at dissolving natural coral rubble and consequently at producing seawater alkalinity under different light regimes and natural variations of aragonite saturation states. Two experiments were carried out, one in an indoor flume under controlled light and the other one outdoor under natural light, in Hawaii (Kaneohe Bay). Seawater DIC was not artificially modified as we did not test the effects of ocean acidification on the metabolism of natural microboring communities; this being reported elsewhere (see Tribollet et al., 2009; Reyes-Nivia et al., 2013; Enochs et al., 2016; Tribollet et al., 2019).

2 Material and methods

2.1 Experimental design

Two distinct experiments were carried out at the Hawaii Institute of Marine Biology (HIMB, HI, United States) in an indoor flume and in an outdoor tank to see the possible effects




FIGURE 1
(A) Brushed coral rubble originating from a massive *Porites* colony, colonized by at least a 12 months-old microboring community. Greenish patches of phototrophic endoliths dominated the microboring community. **(B)** Filaments of the bioeroding chlorophyte *Ostreobium* sp. observed under light microscopy. Chloroplasts are visible inside siphons (filaments without cross-walls) of the chlorophyte. Scale bar = 20 μm .

of light regimes and evolving aragonite saturation state on biogenic dissolution of carbonates by boring microflora, and thus on their production of seawater alkalinity. Substrates used in this study were rubbles mainly composed of *Porites lobata* scraps as one colony of this massive coral was used previously to cut experimental blocks for another bioerosion study (Figure 1A; see also Tribollet et al., 2006; Tribollet et al., 2009). The rest was a mix of *P. compressa*, *Pocillopora meandrina* and *Montipora capitata* rubble, which are commonly found in Kaneohe Bay, Oahu, Hawaii. Those rubbles were colonized by natural communities of microborers in the shallow fringing reef near the HIMB laboratory (<1 m depth) during at least 1 year in order to work with mature communities dominated by *Ostreobium* sp. (Chazottes et al., 1995; Gektidis, 1999; Tribollet, 2008b; Grange et al., 2015). The metabolism of colonized rubbles was then estimated in the two different settings by following the evolution of seawater pH and total alkalinity (see description below).

2.1.1 Indoor flume experiment under controlled light conditions

To study the impact of microborers' metabolism on the carbonate system under saturating light regime (Vooren, 1981; Tribollet et al., 2006) and natural seawater conditions (i.e., seawater from Kaneohe Bay with Ω_{Arag} varying between 3 and 4 during the experiment), we randomly collected natural rubble colonized by mature microboring communities near the HIMB laboratory to form a composite set of rubbles representative of those present on the natural reef flat. Rubbles were brushed gently to remove epiliths such as turfs, crustose coralline algae and serpulids, and were then affixed on a PVC plate with epoxy to obtain a planar surface area of 0.33 m^2 . The top surfaces of rubbles were positioned relative to light in the same way as they were on the reef. The plate was quickly (≈ 15 min) transported into the indoor flume facility at HIMB (see description in Falter et al., 2006) and kept in flow-through seawater (sand filtered seawater from Kaneohe Bay) during 1 day to allow acclimation. The flume was then closed and had a total volume of seawater of 1.1 m^3 . During 1 week, an oscillatory flow was maintained, as well as a constant temperature ($25 \pm 1^\circ\text{C}$) and a 12:12 h light to dark photoperiod (between 7 a.m. and 7 p.m. with a constant light intensity of $350 \mu\text{mol photons}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ delivered by two Solar simulator Arrays, Tailored lighting). The delivered light intensity was similar to that used in a previous experiment carried out at Biosphere (Tribollet et al., 2009). Seawater samples were taken every day at 7 a.m. and 7 p.m. for analysis of total alkalinity (A_T) and pH on the total hydrogen ion concentration scale (pH_T). After that first set of measurements, the flume was flushed to renew seawater to maintain natural reef conditions, and rubbles were brushed again to avoid any epilithic overgrowth. A_T and pH_T variations were then again surveyed during another week (2nd set of measurements). Slight differences in the carbonate system parameters between the initial conditions of the two sets of measurements (Table 1) were due to the time of the day the flume was filled, and therefore to the natural variation of seawater chemistry on the reef flat (1st set in the morning; 2nd set in the afternoon).

2.1.2 Outdoor incubations under natural daylight

To determine if biogenic dissolution by microborers could occur at high Ω_{Arag} values during daylight, i.e., when photosynthesis occurs and thus when carbonate dissolution is thermodynamically difficult, short incubations with another composite set of natural rubbles similar to those used in the indoor flume experiment and representative of the reef flat (Figure 1A), were performed in an outdoor tank. This time, the studied rubble planar surface area was 0.067 m^2 . Brushed rubbles were put in a 18 L tank with a marine seawater bilge pump (12 V, 250 rpm) to insure water motion, under natural daylight. The set of rubbles was incubated 3 times, 4 days apart between 9 a.m. and 6 p.m., allowing the microboring community

TABLE 1 Initial seawater chemistry conditions for each experiment (mean \pm SD).

	T <i>in situ</i> (C)	pH _T	A _T ($\mu\text{mol kg}^{-1}$)	DIC ($\mu\text{mol kg}^{-1}$)	pCO ₂ (μatm)	Ω_{Arag}
<i>Experiment 1</i>						
1st set	26.0	7.929	2144	1899	512	2.85
2nd set	25.5	7.976	2127	1863	447	3.02
<i>Experiment 2</i>						
1st incubation	27.6	7.772	1995	1823	726	2.09
2nd incubation	28.0	7.760	2029	1858	762	2.10
3rd incubation	26.8	7.722	1975	1829	822	1.81

pH_T (total hydrogen ion concentration scale) and total alkalinity (A_T, in units of $\mu\text{equiv.kg}^{-1}$) were used to calculate the other CO₂ chemistry parameters with the software CO₂Sys (Pierrot et al., 2006): dissolved inorganic carbon (DIC), partial pressure of CO₂ (pCO₂) and aragonite saturation state (Ω_{Arag}), using Roy et al. (1993) values for carbonic acid constants K₁ and K₂, and K_{SO₄} as determined by Dickson (1990).

to evolve naturally as in the natural reef flat. Rubbles were indeed kept in a flow-through seawater pumped on the reef flat (unfiltered) between two incubations. They were also gently brushed to remove epiliths' overgrowth, and to allow recruitment of new borers and epiliths, as well as growth of existing microborers up to their new depth of compensation (see Schneider and Le Campion-Alsumard, 1999; Tribollet, 2008a). During the experiments, mean light intensity measured at the weather station of HIMB varied between 750 and 1300 $\mu\text{mol photons.m}^{-2}.\text{s}^{-1}$ depending on the day of incubation. The 18 L tank with rubbles was partially submerged in flow-through seawater in an outdoor flume to maintain a constant temperature ($27 \pm 1^\circ\text{C}$). To ensure that Ω_{Arag} would not reach very high and unrealistic values in the tank (compare to coral reef conditions), a few ml of HCl (0.1 N) were added prior each of the 3 incubations to adjust an initial Ω_{Arag} comprised between 1.8 and 2.1 ($\Omega_{\text{Arag}} > 1$ to avoid dissolution thermodynamically driven). Incubations lasted for 5–9 h and A_T and pH_T were sampled every hour (see initial conditions in Table 1). A_T and pH_T of a control tank with just un-acidified seawater were also measured at the beginning and end of every working day to ensure that phytoplanktonic activity did not impact A_T and/or dissolved inorganic carbon concentration (DIC) during rubble incubations.

2.2 Seawater chemistry analysis

After collection, saturated mercuric chloride was added to seawater samples to prevent biological activity (Dickson et al., 2007). The potentiometric determination of pH_T was realized using Tris/HCl and 2-aminopyridine/HCl buffers in synthetic seawater to calibrate the Ross combination electrode (ORION 81-03) using a Thermo Scientific Orion 2 star Plus pH meter at room temperature. Temperature variation did not exceed 0.1°C during one set of measurements.

Potentiometric titration of A_T was carried out using 0.01 mol.L⁻¹ HCl in NaCl to approximate the ionic strength of seawater after filtration of each A_T sample through Whatman

GF/F (to remove possible suspended carbonate particles). The acid titrant concentration was determined each day of A_T measurements using Certified Reference Material from A. Dickson's laboratory (Scripps Institution of Oceanography). A manual Gran titration was performed with a precision of $\pm 2 \mu\text{equiv.L}^{-1}$.

Measured pH_T and A_T allowed calculating the other CO₂ chemistry parameters with the software CO₂SYS (Pierrot et al., 2006). The same parameters for seawater salinity, phosphates, silicates and seawater density were used as in Tribollet et al. (2019)'s study.

2.3 Net dissolution and organic carbon metabolism calculation

Net dissolution rates (G), expressed in $\text{mmol CaCO}_3.\text{m}^{-2}.\text{h}^{-1}$ were calculated according to the following Eq. 1:

$$G = \frac{1/2 \Delta A_T \times \text{seawater volume}}{\text{substrate surface area} \times \text{time}} \quad (1)$$

where G is positive when net dissolution occurs, ΔA_T is the difference in alkalinity between two measurements (final A_T minus initial A_T), *seawater volume* is the volume of the indoor flume (1st experiment) or the volume of the tank used during outdoor measurements (2nd experiment), *substrate surface area* is the planar surface area of rubbles (equivalent to m² of reef as rubble mimicked eroded reef pavement), and *time* was the duration of the incubation.

Net organic carbon metabolism (NP), expressed in $\text{mmol C.m}^{-2}.\text{h}^{-1}$ was calculated according to the following Eq. 2:

$$NP = \frac{\Delta \text{DIC} \times \text{seawater volume}}{\text{substrate surface area} \times \text{time}} - G \quad (2)$$

Where NP is the rate of net photosynthesis during daylight (negative value, although for simplicity it will be presented as positive in the results) or the rate of dark respiration at night (positive value) and ΔDIC is the difference in DIC between two measurements (final DIC minus initial DIC). The other symbols are as defined in Eq. 1.

For the indoor flume experiment, daily net dissolution was estimated from the slope of the A_T vs. time relationship. Daily excess production (the difference between gross production and daily respiration) was calculated from the slopes of the A_T and DIC vs. time relationships. Other calculations are explained in the Results section.

For the outdoor experiment in a 18 L tank (seawater surface area in contact with the atmosphere estimated to be $<0.1 \text{ m}^2$), NP were estimated without taking into account potential fluxes of CO_2 between air and the tank seawater. Because we did not observe much turbulence at the tank surface due to the bilge pump or wind during our incubations, we assume that CO_2 diffusion was probably limited. However we cannot exclude that NP rates were slightly overestimated at the start of incubations (due to an evasion of CO_2 from the tank to the atmosphere), or underestimated at the end of incubations (uptake of CO_2 atmospheric by the tank seawater).

2.4 Statistical analysis

Statistical tests were performed using STATISTICA 7.1 (Statsoft) and R version 2.15.1 (R Foundation for Statistical Computing). First, an analysis of covariance (ANCOVA) was performed to compare the slopes of the A_T vs. time regression lines between the two sets of measurements performed in the indoor flume (1st experiment). Regression lines were compared by studying the interaction of the categorical variable (i.e., 1st or 2nd set of measurements) with time. Beforehand, a Robust Jarque Bera test and a Levene test were used to test for normality and homogeneity of variances, respectively. Second, DIC concentration vs. A_T relationships were compared between the two sets of measurements using the (S)MATR software (version 1.0 2003, <http://www.bio.mq.edu.au/ecology/SMATR>) designed for making comparisons among lines fitted according to the standardized major axis method (SMA). In both cases, the slopes were not significantly different between the 2 sets of measurements in the indoor flume ($p > 0.05$), therefore a common slope was calculated. The common slope of the DIC concentration vs. A_T relationship was then compared to 0.5 using (S) MATR. Means are reported \pm SE.

3 Results

Observations of rubble showed that epiliths colonizing substrate surfaces were dominated by green short turfs. Rare crustose coralline algae and boring polychaetes (1 or 2) were observed on some rubble, as well as rare traces of grazing. Observations under light microscopy revealed that rubble were colonized by mature communities of microborers dominated by the chlorophyte of the genus *Ostreobium* (Figure 1B; Chazottes et al., 1995; Grange et al., 2015). Once epiliths were gently




FIGURE 2

Chemical parameters recorded during the first experiment, i.e., during the indoor flume experiment under controlled light ($350 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, Hawaii). (A) Total alkalinity (A_T) monitored over time. (B) Dissolved inorganic carbon (DIC) versus A_T . (C) Evolution of the saturation state of aragonite (Ω_{Arag}) over the course of the experiment. The equation of each slope is provided as well as its determination coefficient (R^2).

brushed off, rubble showed an intense yellowish-greenish colour indicative of the presence of a majority of phototrophic endoliths inside dead coral skeletons (Figure 1A).

3.1 Variability of rubble metabolism under constant light (indoor flume experiment)

In the indoor flume experiment ran with rubble exposed to constant light intensity ($350 \mu\text{mol photons.m}^{-2}.\text{s}^{-1}$ with a 12:12 light to dark photoperiod), temperature (-25.5°C) and water flow, total alkalinity of seawater (A_T) increased linearly with time (Figure 2A; $R^2 = 0.99$ and $R^2 = 0.98$ before and after water in the flume was renewed, respectively; $p < 0.0001$). The slope of A_T vs. time relationship remained the same when water in the flume was renewed after a few days of experiment, resulting in a constant mean net dissolution rate of $35.5 \pm 0.2 \text{ mmol CaCO}_3.\text{m}^{-2}.\text{d}^{-1}$ over the course of the experiment. At night, mean CaCO_3 dissolution was $2.4 \pm 0.3 \text{ mmol CaCO}_3.\text{m}^{-2}.\text{h}^{-1}$. A much lower CaCO_3

dissolution rate of about $[35.5 - (2.4 \times 12)]/12 = 0.5$ mmol $\text{CaCO}_3 \cdot \text{m}^{-2} \cdot \text{h}^{-1}$ was recorded under light.

DIC concentration increased linearly with time as well, although data were clearly more scattered than for A_T (Figure 2B; $R^2 = 0.91$; $p < 0.0001$, and $R^2 = 0.83$; $p < 0.005$ before and after water in the flume was renewed, respectively). The slope of the DIC vs. A_T relationships ($R^2 = 0.90$; $p < 0.0001$, and $R^2 = 0.79$; $p = 0.007$, respectively) did not change between the two parts of the experiment, and the mean slope (0.52 ± 0.02) was not significantly different from 0.5 ($p > 0.05$; Figure 2B). DIC increase in the flume was therefore mainly due to CaCO_3 dissolution while excess production (gross production minus respiration) remained on average close to zero. Night time respiration rate (R_n) was 1.8 ± 0.2 mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{h}^{-1}$, thus daily respiration (R) was assumed to be $1.8 \times 24 = 44$ mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{d}^{-1}$ (as daylight respiration could not be measured). Excess production, calculated from the mean slopes of the A_T and DIC vs. time relationships, was 1.9 mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{d}^{-1}$ (net release of DIC). Gross production (P_g) was therefore 42 mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{d}^{-1}$, resulting in a P_g/R ratio of 0.96. Net photosynthesis ($P_g - (12 \times R_n)$) was 20 mmol $\text{C} \cdot \text{m}^{-2}$ during the 12 h of light, or 1.7 mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{h}^{-1}$. Aragonite saturation state (Ω_{Arag}) varied between 2.85 and 3.02 (ambient) at the beginning of the two series of measurements and increased afterwards due to CaCO_3 dissolution (Figure 2C). Ω_{Arag} however either stabilized at about 3.7 or decreased at the end of the incubation period, suggesting that some increase in respiration rate or a decrease in photosynthesis occurred at that time. Nevertheless, these Ω_{Arag} variations did not affect the CaCO_3 dissolution rate.

3.2 Variability of rubble metabolism under natural daylight (outdoor tank)

In the outdoor experiment ran with another set of rubble exposed to natural daylight at constant temperature (-27.5°C ; Table 1), and after addition of a few ml of HCl to decrease Ω_{Arag} down to ~ 2 , we observed the increase of A_T over time in two of the three incubations (Figure 3A). Dissolution rates were 1.4 ± 0.2 and 1.5 ± 0.2 mmol $\text{CaCO}_3 \cdot \text{m}^{-2} \cdot \text{h}^{-1}$ during incubation 1 and 2, respectively. DIC decreased in the meantime (Figure 3B), showing that DIC removal linked to photosynthesis exceeded DIC release due to CaCO_3 dissolution. Net photosynthesis was 12.9 ± 1.6 and 16.4 ± 3.7 mmol $\text{C} \cdot \text{m}^{-2} \cdot \text{h}^{-1}$ during incubations 1 and 2, respectively. Due to both net photosynthesis and CaCO_3 dissolution, Ω_{Arag} increased greatly throughout the two incubations (Figure 3C), from an initial value of 2.1 to a final value of 5.0–5.1 after 5–6 h. In the meantime, $p\text{CO}_2$ decreased to about 150 μatm (Figure 3D). Interestingly these variations in seawater chemistry did not affect the CaCO_3 dissolution rate.




FIGURE 3

Chemical parameters recorded during the course of the second experiment, i.e., during the three outdoor incubations under natural daylight (between 9 a.m. and 6 p.m.; Hawaii). (A) Total alkalinity (A_T). (B) Dissolved inorganic carbon (DIC). (C) Saturation state of aragonite (Ω_{Arag}). (D) Aqueous partial pressure of CO_2 ($p\text{CO}_2$).

In contrast, during the third incubation, the A_T vs. time relationship was bell-shaped, suggesting that an initial phase of CaCO_3 dissolution was followed by some CaCO_3 precipitation in the afternoon (Figure 3A). While the initial value of Ω_{Arag} was very low (1.8; Figure 3C), Ω_{Arag} reached very high values at the end of the day: up to 7.9 after 9 h, corresponding to a $p\text{CO}_2$ of about 30 μatm . CaCO_3 precipitation clearly prevailed over CaCO_3 dissolution for Ω_{Arag} values above 6.4 (Figure 3A, C). Since such values are unlikely to be reached in most natural environments, CaCO_3 dissolution rate was calculated from the five first hours of the experiment (mean $\Omega_{\text{Arag}} = 4.1 \pm 0.7$, compared to 3.7 ± 0.4 and 3.8 ± 0.5 during the 1st and the 2nd experiment, respectively). Dissolution rate was slowing down from 2.8 to 0.2 mmol $\text{CaCO}_3 \cdot \text{m}^{-2} \cdot \text{h}^{-1}$ in the course of the experiment, and was on average 1.8 ± 0.6 mmol $\text{CaCO}_3 \cdot \text{m}^{-2} \cdot \text{h}^{-1}$. In the meantime, DIC decreased with time

(Figure 3B), and net photosynthesis was $24.6 \pm 2.2 \text{ mmol C.m}^{-2}.\text{h}^{-1}$.

4 Discussion


Although chasmo- and crypto-endoliths including phototrophs (see definitions in Golubic et al., 1981) are present in coral pores (Yang et al., 2019; Pernice et al., 2020), most of the biomass in dead corals or coral rubble is generally attributed to microboring flora (Wanders, 1977; Vooren 1981; Fine & Loya 2002). Our light microscopy observations of a few pieces of decalcified coral skeleton confirmed the abundance of the phototrophic eukaryote *Ostreobium* sp. within the natural microboring communities studied in our rubble set exposed to colonization for over 12 months (Figure 1B). The alga *Ostreobium* sp. is known to dominate mature microboring communities in dead coral skeletons after more than 6 months of exposure (Chazottes et al., 1995; Le Campion-Alsumard et al., 1995; le Bris et al., 1998; Tribollet et al., 2006; Tribollet, 2008a; Grange et al., 2015). The green turfs growing on rubble surfaces were also similar to those observed on experimental coral blocks exposed to colonization during at least 6 months at 3 m depth near Coconut Island (Tribollet et al., 2006; see also Hatcher and Larkum, 1983). The presence of only a few polychaetes in rubbles was not surprising as after 6 months of exposure, the experimental blocks studied by Tribollet et al. (2006) presented only a few traces of boring polychaetes at a lagoonal site nearby HIMB laboratory. Polychaetes are generally more abundant in dead coral after several years of exposure to colonization as shown by Kiene and Hutchings (1994) and Tribollet and Golubic (2005). Moreover, Hutchings et al. (1992) pointed out that polychaetes are generally less abundant in shallow lagoon reefs than in other environments.

In our experiments, we did not observe the formation of dense epilithic biofilms at the surface of our coral debris after brushing, unlike Leggat et al. (2019). These authors suggested that corals freshly killed by marine heat waves (i.e., denuded coral skeletons) are immediately and intensely colonized by epilithic biofilms dominated by *Ostreobium* sp. that originate from the interior of the skeleton (thus growing outward), generating strong carbonate dissolution of coral skeletons. This is unlikely as the green alga *Ostreobium* sp. is a cryptic sciaphile siphonale avoiding intense light intensities to prevent photoinhibition (Fine et al., 2005; Ralph et al., 2007). This alga has developed a large repertoire of specific pigments to live in extremely low light environments such as in coral skeletons (Shashar and Stambler, 1992; Koehne et al., 1999; Iha et al., 2021). It also actively dissolves carbonates by creating galleries that perfectly fit the shape of its filaments (e.g., Chazottes et al., 1995; Carreiro-Silva et al., 2005; Tribollet et al., 2009; Wisshak et al., 2011). Although Massé

et al. (2018) and Massé et al. (2020) showed that *Ostreobium* filaments can live freely in seawater and can be epilithic under certain conditions (see also Kobluk and Risk, 1977), they also never observed the formation of dense biofilms of *Ostreobium* sp. on dead corals within a few days. This is especially true since more exposure to light, if not too rapid (Fine et al., 2005), stimulates phototrophic microborers' growth inside carbonates (Schneider and Le Campion-Alsumard 1999; Fine and Loya, 2002; Tribollet and Golubic 2005; Iha et al., 2021). Here we thus present results obtained from typical mature microboring communities that developed in naturally dead corals in reef lagoonal shallow waters.

Brushed coral rubbles under a constant light in the indoor flume ($350 \mu\text{mol photons.m}^{-2}.\text{s}^{-1}$) had a Pg/R ratio of about 1, suggesting much less limitation of microborers' photosynthesis than in Tribollet et al. (2019)'s study (Pg/R ratio <1 in experimental coral blocks colonized by mature microboring communities dominated by *Ostreobium* sp. under $200 \mu\text{mol photons.m}^{-2}.\text{s}^{-1}$). Interestingly, net carbonate dissolution measured in the indoor flume was close to biogenic dissolution measured either by buoyant weight on recently killed corals colonized mostly by *Ostreobium* and regularly brushed ($27\text{--}42 \text{ mmol.m}^{-2}.\text{d}^{-1}$; Reyes-Nivia et al., 2013), or by microscopy techniques on 1 year or three years-old experimental coral blocks grazed by fishes ($8\text{--}30 \text{ mmol.m}^{-2} \text{ d}^{-1}$; Tribollet 2008a; see Table 2 in Tribollet et al., 2019). Removal of epiliths by brushing coral rubble may have indeed simulated a moderate scraping pressure. Grazers including scrapers and excavators, remove substrate surfaces (Clements et al., 2017) to feed on endolithic phototrophs (especially *Ostreobium* sp.). That way they allow more light reaching microborers and consequently, stimulate their growth inside substrates until they reach their new depth of compensation (Tribollet and Golubic, 2005).

The three incubations carried out under natural daylight showed that natural coral rubbles can release alkalinity in seawater at rates as high as $2.8\text{--}3.6 \text{ mequiv.m}^{-2}.\text{h}^{-1}$ (i.e., a CaCO_3 dissolution rate of $1.4\text{--}1.8 \text{ mmol.m}^{-2}.\text{h}^{-1}$). This is the first time that CaCO_3 biogenic dissolution is reported under natural daylight conditions, simultaneously with significant rates of net photosynthesis ($13\text{--}25 \text{ mmol C.m}^{-2}.\text{h}^{-1}$). Assuming a two-fold increase in CaCO_3 dissolution due to the use of HCl (consistent with results found during acidification experiments in New Caledonia as well as in literature; Tribollet et al., 2009; Tribollet et al., 2019; Reyes-Nivia et al., 2013), the alkalinity production by coral rubbles under ambient pCO_2 would still be high ($1.4\text{--}1.8 \text{ mequiv.m}^{-2}.\text{h}^{-1}$). Interestingly, net CaCO_3 biogenic dissolution rate and net photosynthesis increased with time in the three outdoor incubations (up to $\Omega_{\text{Arag}} < 6.4$), suggesting that constant brushing, combined with some environmental factors such as nutrient availability and temperature, certainly stimulated microborer growth (in depth and/or by branching) over the duration of the experiment. Rates of biogenic dissolution



in coral skeletons are known to be related to the biomass of microborers under natural light cycles (Reyes-Nivia et al., 2013; Grange et al., 2015).

By taking into account all experiments, the higher was net photosynthetic activity the higher was net biogenic dissolution (Figure 4), which is *a priori* counter intuitive as rising pH due to photosynthesis is not thermodynamically in favour of carbonate dissolution. This further supports the hypothesis that alkalinity release is related to the metabolic activity of phototrophic microborers (Tribollet et al., 2019). Light availability and DIC speciation did probably however, influenced pathways of carbon acquisition and thus, CaCO₃ dissolution by microborers. Massé et al. (2020) showed that seawater DIC (most probably CO₂ according to Tribollet et al., 2009; Tribollet et al., 2019) is the main source of C involved in *Ostreobium* photosynthesis. However, these authors also reported several other possible sources of C that could be used depending on the DIC concentration in seawater (i.e., HCO₃⁻ or CO₂ released during carbonate dissolution and organic C remineralized by *Ostreobium* associated bacteria). The recycling of inorganic C from dissolved CaCO₃ was also shown in the boring cyanobacteria, *Mastigocoleus testarum*, when DIC was limiting (Guida and Garcia-Pichel, 2016).

In the present study, the logarithmic relationship presented Figure 4 ($R_2 = 0.996$; $p < 0.0001$) shows that biogenic dissolution increased rapidly as a function of net photosynthesis at low light intensities ($\leq 350 \mu\text{mol photons.m}^{-2}.\text{s}^{-1}$), but was less dependent on net photosynthetic rates under high natural daylight intensities ($740\text{--}1300 \mu\text{mol photons.m}^{-2}.\text{s}^{-1}$). In the case of the outdoor incubations, it is unlikely that CO₂ escape into the atmosphere resulted in over-estimation of net photosynthesis because seawater

was on average under-saturated with respect to CO₂ (<90%). Although we cannot exclude that non-boring phototrophic chasmo- and cryptoendoliths may have benefited from increased amounts of light (Marcelino et al., 2018; Yang et al., 2019), we suggest that most probably the decreasing pCO₂ over the course of the outdoor incubations (Figure 3D) in combination with high light availability promoted the use of carbon concentrating mechanisms (CCMs) by *Ostreobium* filaments which dominated microboring communities. The use of CCM was suggested for the first time by Tribollet et al. (2009) in an acidification experiment, and more recently by Massé et al. (2020) depending on environmental conditions.

For four out of the five drift incubations (indoor and outdoor incubations combined), CaCO₃ dissolution rates did not change over time (Figure 2A and 3A) although Ω_{Arag} was clearly increasing (Figure 2C and 3C). Variations in Ω_{Arag} were driven by variations in A_T and DIC, which were in turn driven by organic carbon metabolism (net photosynthesis under natural daylight conditions) and/or CaCO₃ dissolution. It thus may not be appropriate to attempt to relate CaCO₃ fluxes to Ω_{Arag} variations (see Andersson and Gledhill, 2013; Chauvin, 2013). *Ostreobium* response to variations in the parameters of the carbonate system could also be non-linear, as found for some marine calcifiers (Maier et al., 2013) and bioeroding communities dominated by boring sponges (Wisshak et al., 2012; Stubler and Peterson, 2016). Total bioerosion rates (particulate + dissolved CaCO₃) for the boring clionaid sponge, *Cliona orientalis*, were similar under present (pCO₂ = $393 \pm 56 \mu\text{atm}$) and slightly lowered pCO₂ conditions (pCO₂ = $339 \pm 37 \mu\text{atm}$), suggesting that the sponge was able to compensate for the less favourable conditions (Wisshak et al., 2012). In a similar manner, the somewhat higher range in pCO₂ (pCO₂ = $339\text{--}512 \mu\text{atm}$; $\Omega_{\text{Arag}} = 2.85\text{--}3.78$) during the indoor flume experiment did not affect net biogenic dissolution rates, which were also similar at Ω_{Arag} 3 and 3.5 (pCO₂ = 570 and $437 \mu\text{atm}$ respectively) during the acidification experiment in New Caledonia conducted by Tribollet et al. (2019). It is therefore likely that a threshold somewhere between Ω_{Arag} 2 and 3 must be reached before promoting increased CaCO₃ dissolution by microboring communities dominated by *Ostreobium*. Above Ω_{Arag} 3, microboring communities dominated by *Ostreobium* seem to be able to maintain a constant carbonate dissolution rate.

Surprisingly, elevated CO₂ at the beginning of the incubations performed under natural daylight in the outdoor experiment ($757 \pm 31 \mu\text{atm}$, due to the initial addition of HCl) did not cause an increase of CaCO₃ dissolution early in the incubations like it was shown in controlled acidification experiments under low light intensities (Tribollet et al., 2009; Reyes-Nivia et al., 2013; Tribollet et al., 2019). It is hypothesized that high light availability promoted CO₂ fixation by RUBISCO, preventing acidification of the intracellular medium of boring

filaments. In contrast, during the third incubation performed under natural daylight, CaCO_3 dissolution was followed by some CaCO_3 precipitation when the tank seawater Ω_{Arag} reached values higher than 6.4 and pCO_2 decreased down to 30 μatm (Figure 3A and 3C). Conditions where pH is above 9, Ω is very high while pCO_2 is close to 13 μatm , are known to stimulate nucleation of CaCO_3 in seawater (Morse and He, 1993; Wurgaft et al., 2016). It is most probable here that chemical conditions at the interface (boundary layer) between carbonate rubble (behaving as carbonate nucleus) and seawater were even more extreme than those measured in the middle of the tank at the end of the third incubation, creating conditions for CaCO_3 precipitation. We also hypothesize that under such extreme conditions, either 1) the available energy was allocated by phototrophic microborers (mainly *Ostreobium* sp.) to CCM to support the Calvin cycle instead of supporting other processes such as the Ca^{2+} ATPase dependent pumps, and/or 2) the use of HCO_3^- as a source of CO_2 for photosynthesis via the implication of carbonic anhydrase enzymes in *Ostreobium* filaments (Shashar and Stambler 1992; Iha et al., 2021) may have produced significant amounts of OH^- (Giordano et al., 2005) which, combined to H^+ protons, could have stopped the biogenic dissolution process (this implies that *Ostreobium* filaments used $\text{Ca}^{2+}/\text{H}^+$ ATPase dependent pumps similar to those involved by the bioeroding cyanobacterium *Mastigocoleus testeterum*; (Garcia-Pichel et al., 2010), a hypothesis supported by Krause et al. (2019) and Iha et al. (2021)), and/or 3) the number of Ca^{2+} pumps was limited so despite an important photosynthetic activity (providing energy), biogenic dissolution could be limited or stopped. In the case of cultured euendolithic cyanobacteria, Garcia-Pichel et al. (2010) showed that calcium release at the substrate surface resulted in a local supersaturation with respect to CaCO_3 creating conditions for a micrite layer formation. More recently, Krause et al. (2019) highlighted the capacity of *Ostreobium* filaments to uptake Ca^{2+} at their apical thallus tip and to excrete it at the other end of its thallus, most probably in the coral pore spaces causing reprecipitation of CaCO_3 in the form of secondary aragonite. In contrast, in the indoor flume, micritization most probably did not contribute to the difference recorded between alkalinity production in the light and in the dark during the indoor flume experiment because Ω_{Arag} was at most 3.7 and no curvature of the A_T vs. time relationships was observed before Ω_{Arag} reached about 5 (Figure 3A and 3C). Nevertheless, this process could probably lower alkalinity production by microborers on some reef flats where extreme diurnal variability in carbonate chemistry is recorded (e.g. Ω_{Arag} up to 6.5 during the day at Lady Elliot reef, Great Barrier Reef; Shaw et al., 2012).

5 Conclusion and perspectives

We show here that microboring communities dominated by the phototrophic green alga *Ostreobium* sp. are most probably

the main drivers of reef dead coral skeleton dissolution observed under a large range of saturation states in seawater, at least between 2 and 6.4. In previous studies on microboring flora, no traces of random dissolution by endolithic bacteria inside coral skeletons were observed under scanning electron microscopy (Tribollet and Golubic, 2011; Tribollet et al., 2019) although bacteria are known to be associated to *Ostreobium* filaments (unpublished data) and the boring cyanobacterium *Solentia* sp. (see Figure 10 in Le Campion-Alsumard et al., 1996). Those bacteria are preferentially located inside filaments (personal observations) or in polysaccharide envelope layers of cells (Le Campion-Alsumard et al., 1996). We suggest that phototrophic microborers are thus able to keep dissolving reef carbonate substrates under higher saturation states of aragonite than those measured in our study as long as they have enough light and available C sources, as nitrogen source is not limiting (see Massé et al., 2020; Pernice et al., 2020). We thus strongly stress the importance for more investigations to better understand the roles of the phototrophic microboring flora in reef carbonate budget and resilience as they have paradoxical activities: they are able to rise pH, Ω_{Arag} , and seawater alkalinity under daylight while they dissolve actively dead carbonate substrates enhancing reef fragility and degradation. Those roles should be studied in more or less mature endolithic communities in rubble and dead coral substrates as for instance, the epilithic cover (Chazottes et al., 2002) and the presence of macroborers such as sponges and bivalves can greatly influence the composition of microboring flora communities and consequently, the biogenic dissolution rates (Grange et al., 2015; Schönberg et al., 2017; Fordyce et al., 2020). There is also a specific need to better understand the functional roles played by the various genotypes and phenotypes of *Ostreobium* as Massé et al. (2020) highlighted a large diversity in response to environmental constraints such as the type of habitat. Finally, we also suggest that microborer metabolic activity could counteract the effects of acidification at least at the local scale of very shallow reefs with long water residence time (Page et al., 2016) and in some living calcifiers such as massive corals which are heavily colonized by *Ostreobium* sp. compare to branching corals (Le Campion-Alsumard et al., 1995; Godinot et al., 2012; Massé et al., 2018) and crustose coralline algae (Tribollet and Payri, 2001; Reyes-Nivia et al., 2014). They could indeed benefit to their host through a potential transfer of photoassimilates (e.g. in Mediterranean subtropical corals: Fine and Loya, 2002; Sangsawang et al., 2017), photoprotection (Galindo-Martinez et al., 2022), and/or through the rise of pH and alkalinity at the microscale inside their host under various environmental conditions (Nothdurft and Webb, 2009; Garcia-Pichel et al., 2010; Reyes-Nivia et al., 2014; Krause et al., 2019; Tribollet et al., 2019 and present study).

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

Author contributions

AT designed the experiments. AT and PC collected samples and carried out the analyses. AC helped with statistical analysis. All authors (AT, AC, and PC) interpreted data, discussed trends and wrote the manuscript.

Funding

This study was supported by a Marie Curie Outgoing International Fellowship obtained as part of the FP6 of EU, as well as by the project MIDACOR funded by the French Ministry of Ecology (MEEDDM Ministry action: 23-190RECHERCINCITAT, program 190-090-THUR-BSAF) and IRD (Institut de Recherche pour le Développement).

References

- Agostini, S., Harvey, B. P., Wada, S., Kon, K., Milazzo, M., Inaba, K., et al. (2018). Ocean acidification drives community shifts towards simplified non-calcified habitats in a subtropical-temperate transition zone. *Sci. Rep.* 8, 11354. doi:10.1038/s41598-018-29251-7
- Andersson, A. J., Bates, N. R., and Mackenzie, F. T. (2007). Dissolution of carbonate sediments under rising pCO₂ and ocean acidification: Observations from Devil's Hole, Bermuda. *Aquat. Geochem.* 13, 237–264. doi:10.1007/s10498-007-9018-8
- Andersson, A. J., and Gledhill, D. (2013). Ocean acidification and coral reefs: Effects on breakdown, dissolution, and net ecosystem calcification. *Ann. Rev. Mar. Sci.* 5, 321–348. doi:10.1146/annurev-marine-121211-172241
- Andersson, A. J., Mackenzie, F. T., and Bates, N. R. (2008). Life on the margin: Implications of ocean acidification on Mg-calcite, high latitude and cold-water marine calcifiers. *Mar. Ecol. Prog. Ser.* 373, 265–273. doi:10.3354/meps07639
- Bindoff, N. L., Cheung, W. W. L., and Kairo, J. G. (2019). *IPCC SR ocean and cryosphere chapter 5: Changing ocean, marine ecosystems, and dependent communities*. Geneva, Switzerland: IPCC.
- Carreiro-Silva, M., McClanahan, T. R., and Kiene, W. E. (2005). The role of inorganic nutrients and herbivory in controlling microbioerosion of carbonate substratum. *Coral Reefs* 24, 214–221. doi:10.1007/s00338-004-0445-3
- Chauvin, A. (2013). "Photosynthèse et calcification sur les récifs coralliens face au changement global et aux impacts anthropiques: Du corail hermatypique *Acropora muricata* à l'écosystème," (University of Réunion). thesis.
- Chazottes, V., Le Campion-Alsumard, T., and Peyrot-Clausade, M. (1995). Bioerosion rates on coral reefs: Interactions between macroborers, microborers and grazers (moorea, French polynesia). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 113, 189–198. doi:10.1016/0031-0182(95)00043-1
- Chazottes, V., Le Campion-Alsumard, T., Peyrot-Clausade, M., and Cuët, P. (2002). The effects of eutrophication-related alterations to coral reef communities on agents and rates of bioerosion (Reunion Island, Indian Ocean). *Coral Reefs* 21, 375–390. doi:10.1007/s00338-002-0259-0
- Clements, K. D., German, D. P., Piché, J., Tribollet, A., and Choat, J. H. (2017). Integrating ecological roles and trophic diversification on coral reefs: Multiple lines of evidence identify parrotfishes as microphages. *Biol. J. Linn. Soc.* 120 (4), 729–751.
- Comeau, S., Edmunds, P. J., Spindel, N. B., and Carpenter, R. C. (2014). Fast coral reef calcifiers are more sensitive to ocean acidification in short-term laboratory incubations. *Limnol. Oceanogr.* 59, 1081–1091. doi:10.4319/lo.2014.59.3.1081
- Cyronak, T., Schulz, K. G., Santos, I. R., and Eyre, B. D. (2014). Enhanced acidification of global coral reefs driven by regional biogeochemical feedbacks. *Geophys. Res. Lett.* 41, 5538–5546. doi:10.1002/2014GL060849
- Dickson, A., Christian, J. R., and Sabine, C. L. (2007). *Guide to best practices for ocean CO₂ measurements*. Canada: PICES Special Publication.
- Dickson, A. G. (1990). Standard potential of the reaction: AgCl(s) + 1/2H₂(g) = Ag(s) + HCl(aq), and the standard acidity constant of the ion HSO₄⁻ in synthetic sea water from 273.15 to 318.15 K. *J. Chem. Thermodyn.* 22, 113–127. doi:10.1016/0021-9614(90)90074-z
- Enochs, I. C., Manzello, D. P., Tribollet, A., Valentino, L., Kolodziej, G., Donham, E. M., et al. (2016). Elevated colonization of microborers at a volcanically acidified coral reef. *PLoS ONE* 11, e0159818. doi:10.1371/journal.pone.0159818
- Eyre, B. D., Andersson, A. J., and Cyronak, T. (2014). Benthic coral reef calcium carbonate dissolution in an acidifying ocean. *Nat. Clim. Chang.* 4, 969–976. doi:10.1038/nclimate2380
- Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., et al. (2011). Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. *Nat. Clim. Chang.* 1, 165–169. doi:10.1038/nclimate1122
- Falter, J. L., Atkinson, M. J., Fleming, J. H., Bos, M. M., Lowe, R. J., Koseff, J. R., et al. (2006). A novel flume for simulating the effects of wave- and tide-driven water motion on the biogeochemistry of benthic reef communities. *Limnol. Oceanogr. Methods* 4, 68–79. doi:10.4319/lom.2006.4.68
- Falter, J. L., Lowe, R. J., Atkinson, M. J., and Cuët, P. (2012). Seasonal coupling and de-coupling of net calcification rates from coral reef metabolism and carbonate chemistry at Ningaloo Reef, Western Australia. *J. Geophys. Res.* 117, C05003. doi:10.1029/2011JC007268
- Fine, M., and Loya, Y. (2002). Endolithic algae: An alternative source of photoassimilates during coral bleaching. *Proc. R. Soc. Lond. B* 269, 1205–1210. doi:10.1098/rspb.2002.1983
- Fine, M., Meroz-Fine, E., and Hoegh-Guldberg, O. (2005). Tolerance of endolithic algae to elevated temperature and light in the coral *Montipora monasteriata* from the southern Great Barrier Reef. *J. Exp. Biol.* 208, 75–81. doi:10.1242/jeb.01381

Acknowledgments

The authors would like to warmly thank the late Marlin J. Atkinson and James H. Fleming, as well as Dan Schar without whom this study could not have been carried out at the Hawaii Institute of Marine Biology on Coconut Island.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Fordyce, A. J., Ainsworth, T. D., and Leggat, W. (2020). Microalgae, a boring bivalve and a coral – A newly described association between two coral reef bioeroders within their coral host. *Integr. Org. Biol.* 2 (1), obaa035. doi:10.1093/iob/obaa035
- Galindo-Martinez, C. T., Weber, M., Avila-Magna, V., Enriquez, S., Kitano, H., Medina, M., et al. (2022). The role of the endolithic alga *Ostreobium* spp. during coral bleaching recovery. *Sci. Rep.* 12, 2977. doi:10.1038/s41598-022-07017-6
- Garcia-Pichel, F., Ramirez-Reinat, E., and Gao, Q. (2010). Microbial excavation of solid carbonates powered by P-type ATPase-mediated transcellular Ca²⁺ transport. *Proc. Natl. Acad. Sci. U. S. A.* 107, 21749–21754. doi:10.1073/pnas.1011884108
- Gektidis, M. (1999). Development of microbial euendolithic communities: The influence of light and time. *bgsd.* 45, 147–150. doi:10.37570/bgsd-1998-45-18
- Giordano, M., Beardall, J., and Raven, J. A. (2005). CO₂ concentrating mechanisms in algae: Mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* 56, 99–131. doi:10.1146/annurev.arplant.56.032604.144052
- Godinot, C., Tribollet, A., Grover, R., and Ferrier-Pagès, C. (2012). Bioerosion by euendoliths decreases in phosphate-enriched skeletons of living corals. *Biogeosciences* 9, 2377–2384. doi:10.5194/bg-9-2377-2012
- Golubic, S., Friedmann, I., and Schneider, J. (1981). The lithobiontic ecological niche, with special reference to microorganisms. *J. Sediment. Petrology* 51, 0475–0478.
- Grange, J. S., Rybarczyk, H., and Tribollet, A. (2015). The three steps of the carbonate biogenic dissolution process by microborers in coral reefs (New Caledonia). *Environ. Sci. Pollut. Res.* 22, 13625–13637. doi:10.1007/s11356-014-4069-z
- Guida, B. S., and Garcia-Pichel, F. (2016). Draft genome assembly of a filamentous euendolithic (true boring) cyanobacterium, *Mastigocoleus testarum* strain BC008. *Genome Announc.* 4, 015744–15. doi:10.1128/genomeA.01574-15
- Guinotte, J. M., and Fabry, V. J. (2008). Ocean acidification and its potential effects on marine ecosystems. *Ann. N. Y. Acad. Sci.* 1134, 320–342. doi:10.1196/annals.1439.013
- Hatcher, B. G., and Larkum, A. W. D. (1983). An experimental analysis of factors controlling the standing crop of the epilithic algal community on a coral reef. *J. Exp. Mar. Biol. Ecol.* 69, 61–84. doi:10.1016/0022-0981(83)90172-7
- Hutchings, P. A., Kien, W. E., Cunningham, R. B., and Donnelly, C. (1992). Spatial and temporal patterns of non-colonial boring organisms (polychaetes, sipunculans and bivalve molluscs) in Porites at Lizard Island, Great Barrier Reef. *Coral Reefs* 11, 23–31. doi:10.1007/bf00291931
- Iha, C., Dougan, K. E., Varela, J. A., Avila, V., Jackson, C. J., Bogaert, K. A., et al. (2021). Genomic adaptations to an endolithic lifestyle in the coral-associated alga *Ostreobium*. *Curr. Biol.* 31, 1393–1402.e5. doi:10.1016/j.cub.2021.01.018
- Johnson, M. D., Price, N. N., and Smith, J. E. (2014). Contrasting effects of ocean acidification on tropical fleshy and calcareous algae. *PeerJ* 2, e411. doi:10.7717/peerj.411
- Kiene, W. E., and Hutchings, P. A. (1994). Bioerosion experiments at lizard Island, Great barrier reef. *Coral Reefs* 13, 91–98. doi:10.1007/bf00300767
- Kobluk, D. R., and Risk, M. J. (1977). Micritization and carbonate-grain binding by endolithic algae. *Am. Assoc. Petroleum Geol. Bull.* 61 (7), 1069–1082.
- Koehne, B., Elli, G., Jennings, R. C., Wilhelm, C., and Trissl, H.-W. (1999). Spectroscopic and molecular characterization of a long wavelength absorbing antenna of *Ostreobium* sp. *Biochimica Biophysica Acta - Bioenergetics* 1412, 94–107. doi:10.1016/S0005-2728(99)00061-4 Available at: www.elsevier.com/locate/bba.
- Krause, S., Liebetrau, V., Nehrke, G., Damm, T., Büsse, S., Leipe, T., et al. (2019). Endolithic algae affect modern coral carbonate morphology and chemistry. *Front. Earth Sci. (Lausanne)* 7, 1–19. doi:10.3389/feart.2019.00304
- Krumins, V., Gehlen, M., Arndt, S., van Cappellen, P., and Regnier, P. (2013). Dissolved inorganic carbon and alkalinity fluxes from coastal marine sediments: Model estimates for different shelf environments and sensitivity to global change. *Biogeosciences* 10, 371–398. doi:10.5194/bg-10-371-2013
- Kuffner, I. B., Andersson, A. J., Jokiel, P. L., Rodgers, K. S., and MacKenzie, F. T. (2008). Decreased abundance of crustose coralline algae due to ocean acidification. *Nat. Geosci.* 1, 114–117. doi:10.1038/ngeo100
- Langdon, C., and Atkinson, M. J. (2005). Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *J. Geophys. Res.* 110, C09S07–16. doi:10.1029/2004JC002576
- Le Bris, S., Le Campion-Alsumard, T., and Romano, J.-C. (1998). Caractéristiques du feutrage algal des récifs coralliens de Polynésie française soumis à différentes intensités de bioérosion. *Oceanol. Acta* 21, 695–708. doi:10.1016/s0399-1784(99)80025-5
- Le Campion-Alsumard, T., Golubic, S., and Hutchings, P. (1995). Microbial endoliths in skeletons of live and dead corals: *Porites lobata* (Moorea, French Polynesia). *Mar. Ecol. Prog. Ser.* 117, 149–157. doi:10.3354/meps117149
- Le Campion-Alsumard, T., Golubic, S., and Pantazidou, A. (1996). On the euendolithic genus *Solentia ERCEGOVIC* (Cyanophyta/Cyanobacteria). *Archiv. algolstud.* 83, 107–127. doi:10.1127/algol_stud/83/1996/107
- Leggat, W. P., Camp, E. F., Suggett, D. J., Heron, S. F., Fordyce, A. J., Gardner, S., et al. (2019). Rapid coral decay is associated with marine heatwave mortality events on reefs. *Curr. Biol.* 29, 2723–2730.e4. doi:10.1016/j.cub.2019.06.077
- Maier, C., Schubert, A., Berzunza Sánchez, M. M., Weinbauer, M. G., Watremetz, P., and Gattuso, J. P. (2013). End of the century pCO₂ levels do not impact calcification in mediterranean cold-water corals. *PLoS ONE* 8, e62655. doi:10.1371/journal.pone.0062655
- Marcelino, V. R., van Oppen, M. J. H., and Verbruggen, H. (2018). Highly structured prokaryote communities exist within the skeleton of coral colonies. *ISME J.* 12, 300–303. doi:10.1038/ismej.2017.164
- Massé, A., Domart-Coulon, I., Golubic, S., Duché, D., and Tribollet, A. (2018). Early skeletal colonization of the coral holobiont by the microboring Ulvophyceae *Ostreobium* sp. *Sci. Rep.* 8, 2293. doi:10.1038/s41598-018-20196-5
- Massé, A., Tribollet, A., Meziane, T., Bourguet-Kondracki, M. L., Yéprémian, C., Sève, C., et al. (2020). Functional diversity of microboring *Ostreobium* algae isolated from corals. *Environ. Microbiol.* 22, 4825–4846. doi:10.1111/1462-2920.15256
- Morse, J. W., and He, S. (1993). Influences of T, S and pCO₂ on the pseudo-homogeneous precipitation of CaCO₃ from seawater: Implications for whiting formation. *Mar. Chem.* 41, 291–297. doi:10.1016/0304-4203(93)90261-1
- Nothdurft, L. D., and Webb, G. E. (2009). Earliest diagenesis in scleractinian coral skeletons: Implications for palaeoclimate-sensitive geochemical archives. *Facies* 55, 161–201. doi:10.1007/s10347-008-0167-z
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., et al. (2005). Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437, 681–686. doi:10.1038/nature04095
- Page, H. N., Andersson, A. J., Jokiel, P. L., Rodgers, K. S., Lebrato, M., Yeakel, K., et al. (2016). Differential modification of seawater carbonate chemistry by major coral reef benthic communities. *Coral Reefs* 35, 1311–1325. doi:10.1007/s00338-016-1490-4
- Pandolfi, J. M., Connolly, S. R., Marshall, D. J., and Cohen, A. L. (2011). Projecting coral reef futures under global warming and ocean acidification. *Science* 333, 418–422. doi:10.1126/science.1204794 Available at: <http://paleodb.org/cgi-bin/bridge.pl>.
- Pernice, M., Raina, J. B., Räder, N., Cárdenas, A., Pogoreutz, C., and Voolstra, C. R. (2020). Down to the bone: The role of overlooked endolithic microbiomes in reef coral health. *ISME J.* 14, 325–334. doi:10.1038/s41396-019-0548-z
- Pierrot, D., Lewis, E., and Wallace, D. W. R. (2006). *MS Excel program developed for CO₂ system calculations*. Tennessee: ORNL/CDIAC-105 Carbon Dioxide Information Analysis Center.
- Ralph, P. J., Larkum, A. W. D., and Kühl, M. (2007). Photobiology of endolithic microorganisms in living coral skeletons: 1. Pigmentation, spectral reflectance and variable chlorophyll fluorescence analysis of endoliths in the massive corals *Cyphastrea serailia*, *Porites lutea* and *Goniastrea australensis*. *Mar. Biol.* 152, 395–404. doi:10.1007/s00227-007-0694-0
- Reyes-Nivia, C., Diaz-Pulido, G., and Dove, S. (2014). Relative roles of endolithic algae and carbonate chemistry variability in the skeletal dissolution of crustose coralline algae. *Biogeosciences* 11, 4615–4626. doi:10.5194/bg-11-4615-2014
- Reyes-Nivia, C., Diaz-Pulido, G., Kline, D., Guldberg, O. H., and Dove, S. (2013). Ocean acidification and warming scenarios increase microbioerosion of coral skeletons. *Glob. Chang. Biol.* 19, 1919–1929. doi:10.1111/gcb.12158
- Roy, R. N., Roy, L. N., Vogel, K. M., Porter-Moore, C., Pearson, T., Good, C. E., et al. (1993). The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Mar. Chem.* 44, 249–267. doi:10.1016/0304-4203(93)90207-5
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., et al. (2004). The oceanic sink for anthropogenic CO₂. *Science* 305, 367–371. doi:10.1126/science.1097403
- Sangsawang, L., Casareto, B. E., Ohba, H., Vu, H. M., Meekaew, A., Suzuki, T., et al. (2017). ¹³C and ¹⁵N assimilation and organic matter translocation by the endolithic community in the massive coral *Porites lutea*. *R. Soc. open Sci.* 4, 171201. doi:10.1098/rsos.171201
- Schneider, J., and Le Campion-Alsumard, T. (1999). Construction and destruction of carbonates by marine and freshwater cyanobacteria. *Eur. J. Phycol.* 34 (4), 417–426. doi:10.1080/09670269910001736472
- Schönberg, C. H. L., Fang, J. K. H., Carreiro-Silva, M., Tribollet, A., and Wisshak, M. (2017). Bioerosion: The other ocean acidification problem. *ICES J. Mar. Sci.* 74, 895–925. doi:10.1093/icesjms/fsw254

- Shamberger, K. E. F., Feely, R. A., Sabine, C. L., Atkinson, M. J., DeCarlo, E. H., Mackenzie, F. T., et al. (2011). Calcification and organic production on a Hawaiian coral reef. *Mar. Chem.* 127, 64–75. doi:10.1016/j.marchem.2011.08.003
- Shashar, N., and Stambler, N. (1992). Endolithic algae within corals - life in an extreme environment. *J. Exp. Mar. Biol. Ecol.* 163, 277–286. doi:10.1016/0022-0981(92)90055-f
- Shaw, E. C., McNeil, B. I., and Tilbrook, B. (2012). Impacts of ocean acidification in naturally variable coral reef flat ecosystems. *J. Geophys. Res.* 117, C03038. doi:10.1029/2011JC007655
- Stubler, A. D., and Peterson, B. J. (2016). Ocean acidification accelerates net calcium carbonate loss in a coral rubble community. *Coral Reefs* 35, 795–803. doi:10.1007/s00338-016-1436-x
- Tribollet, A., Chauvin, A., and Cuet, P. (2019). Carbonate dissolution by reef microbial borers: A biogeological process producing alkalinity under different pCO₂ conditions. *Facies* 65, 9. doi:10.1007/s10347-018-0548-x
- Tribollet, A. (2008a). Dissolution of dead corals by euendolithic microorganisms across the northern Great Barrier Reef (Australia). *Microb. Ecol.* 55, 569–580. doi:10.1007/s00248-007-9302-6
- Tribollet, A., Godinot, C., Atkinson, M., and Langdon, C. (2009). Effects of elevated pCO₂ on dissolution of coral carbonates by microbial euendoliths. *Glob. Biogeochem. Cycles* 23, GB3008. doi:10.1029/2008GB003286
- Tribollet, A., and Golubic, S. (2005). Cross-shelf differences in the pattern and pace of bioerosion of experimental carbonate substrates exposed for 3 years on the northern Great Barrier Reef, Australia. *Coral Reefs* 24, 422–434. doi:10.1007/s00338-005-0003-7
- Tribollet, A., and Golubic, S. (2011). "Reef bioerosion: Agents and processes," in *Coral reefs: An ecosystem in transition*. Editors Z. Dubinsky and N. Stambler (Berlin Heidelberg New York: Springer Science + Business Media), 435–449. doi:10.1007/978-94-007-0114-4_25
- Tribollet, A., Langdon, C., Golubic, S., and Atkinson, M. (2006). Endolithic microflora are major primary producers in dead carbonate substrates of Hawaiian coral reefs. *J. Phycol.* 42, 292–303. doi:10.1111/j.1529-8817.2006.00198.x
- Tribollet, A., and Payri, C. (2001). Bioerosion of the coralline alga *Hydrolithon onkodes* by microborers in the coral reefs of Moorea, French Polynesia. *Oceanol. Acta* 24, 329–342. doi:10.1016/s0399-1784(01)01150-1
- Tribollet, A. (2008b). "The boring microflora in modern coral reef ecosystems: A review of its roles," in *Current developments in bioerosion*. Editors M. Wisshak and L. Tapanila (Berlin Heidelberg New York: Springer), 67–94.
- Vargas, C. A., Cuevas, L. A., Broitman, B. R., San Martin, V. A., Lagos, N. A., Gaitan-Espitia, J. G., et al. (2022). Upper environmental pCO₂ drives sensitivity to ocean acidification in marine invertebrates. *Nat. Clim. Chang.* 12, 200–207. doi:10.1038/s41558-021-01269-2
- Vooren, C. M. (1981). Photosynthetic rates of benthic algae from the deep coral reef of Curaçao. *Aquat. Bot.* 10, 143–159. doi:10.1016/0304-3770(81)90017-6
- Wanders, J. B. W. (1977). The role of benthic algae in the shallow reef of Curaçao (Netherlands Antilles) III : The significance of grazing. *Aquat. Bot.* 3, 357–390. doi:10.1016/0304-3770(77)90040-7
- Wisshak, M., Schönberg, C. H. L., Form, A., and Freiwald, A. (2012). Ocean acidification accelerates reef bioerosion. *PLoS ONE* 7, e45124. doi:10.1371/journal.pone.0045124
- Wisshak, M., Tribollet, A., Golubic, S., Jakobsen, J., and Freiwald, A. (2011). Temperate bioerosion: Ichnodiversity and biodiversity from intertidal to bathyal depths (Azores). *Geobiology* 9, 492–520. doi:10.1111/j.1472-4669.2011.00299.x
- Wurgaft, E., Steiner, Z., Luz, B., and Lazar, B. (2016). Evidence for inorganic precipitation of CaCO₃ on suspended solids in the open water of the Red Sea. *Mar. Chem.* 186, 145–155. doi:10.1016/j.marchem.2016.09.006
- Yang, S. H., Tandon, K., Lu, C. Y., Wada, N., Shih, C. J., Hsiao, S. S. Y., et al. (2019). Metagenomic, phylogenetic, and functional characterization of predominant endolithic green sulfur bacteria in the coral *Isopora palifera*. *Microbiome* 7, 3–13. doi:10.1186/s40168-018-0616-z