



Simultaneous EEG-fNIRS explore somatosensory prediction in the premature neonate brain

A-L Marais, Victoria Dumont, Marie Anquetil, S Rossi, A.-S. Trentesaux,
Nadege Roche-Labarbe

► To cite this version:

A-L Marais, Victoria Dumont, Marie Anquetil, S Rossi, A.-S. Trentesaux, et al.. Simultaneous EEG-fNIRS explore somatosensory prediction in the premature neonate brain. fNIRS 2022, Oct 2022, Boston (MA), United States. 2022, 10.13140/RG.2.2.12566.01609 . hal-03994344

HAL Id: hal-03994344

<https://hal.science/hal-03994344>

Submitted on 17 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Simultaneous EEG-fNIRS to explore somatosensory prediction in the premature neonate brain

Poster
PS3-62-
335

Marais, A-L.^{1*}, Dumont, V.¹, Anquetil, M.², Rossi, S.², Trentesaux, A-S³, Roche-Labarbe, N.¹

* anne-lise.marais@unicaen.fr

¹ Normandie Univ, UNICAEN, INSERM, COMETE, GIP CYCERON, 14000 Caen, France;

² Normandie Univ, UNICAEN, LPCN, 14000 Caen, France; ³ Unité de Néonatalogie, CHU, Caen, France.


01 THEORETICAL BACKGROUND

Predictive coding

- Predictive coding** proposes that the brain is inherently predictive about the environment (Friston, 2005). These predictions can be observed through various phenomena such as:
 - sensory prediction**, the neural anticipation of future stimulation on the basis of previous sensory inputs
 - repetition suppression**, the decrease of cerebral activity to a repeated stimulus after a stable representation is formed, thus correctly predicted
- Predictions are **necessary to optimize** behavioral responses and the use of **attentional and energy resources**
- Sensory prediction and repetition suppression can be **altered in neurodevelopmental disorders** (NDD) like attention deficit disorders (Gonzalez-Gadea et al., 2015)
- Predictive abilities were recently described in preterm infants in the tactile modality (Dumont et al., 2022)


Tactile modality

- Somatosensory inputs are numerous from birth and touch ontogenetic anteriority makes it the **foundation of neurosensory integration, and motor, cognitive and affective development** (Cascio, 2010)
- The tactile modality and the cognitive functions relying on it are often **altered in NDD** (Zoelen & Delvenne, 2018)




04 PRELIMINARY RESULTS ($N_{EEG}=30$, $N_{fNIRS}=13$)


Somatosensory cortex




Repetition suppression



Deviance



Omission



REFERENCES

Basirat, A., Dehaene, S., Dehaene-Lambertz, G. (2014). A hierarchy of cortical responses to sequence violations in three-month-old infants. *Cognition*, 132(2), 137-50. doi: 10.1016/j.cognition.2014.03.013

Dehaene-Lambertz, G., & Dehaene, S. (1994). Speed and cerebral correlates of syllable discrimination in infants. *Nature*, 370(6487), 292-295. https://doi.org/10.1038/370292a0

Cascio, C. J. (2010). Somatosensory processing in neurodevelopmental disorders. *Journal of Neurodevelopmental Disorders*, 2(2), 62-69. https://doi.org/10.1007/s11689-010-9046-3

Dumont, V., Giovannella, M., Zuba, D., Clouard, R., Durduran, T., Guillois, B., & Roche-Labarbe, N. (2022). Somatosensory processing in the premature neonate brain. In *Developmental Cognitive Neuroscience* (Vol. 57, p. 101148). Elsevier BV. https://doi.org/10.1016/j.dcn.2022.101148

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1456), 815-836. https://doi.org/10.1098/rstb.2005.1622

Gonzalez-Gadea, M. L., Chennu, S., Bekinschtein, T. A., Rattazzi, A., Beraudi, A., Tripicchio, P., Moyano, B., Sofita, Y., Steinberg, L., Adolfi, F., Sigman, M., Marino, J., Manes, F., & Ibanez, A. (2015). Predictive coding in autism spectrum disorder and attention deficit hyperactivity disorder. *Journal of Neurophysiology*, 114(5), 2625-2636. https://doi.org/10.1152/jn.00543.2015

Nelson, C., Ellis, A., Collins, P., Lang, S. (1990). Infants' neuroelectric responses to missing stimuli: can missing stimuli be novel stimuli? *Developmental neuropsychology*, 6(4), 339-349.

Spackman, L. A., Towell, A., & Boyd, S. G. (2010). Somatosensory discrimination: An intracranial event-related potential study of children with refractory epilepsy. *Brain Research*, 1310, 68-76. https://doi.org/10.1016/j.brainres.2009.10.072

Zoelen, D., & Delvenne, V. (2018). Treatment of sensory information in neurodevelopmental disorders. *Revue Médicale de Bruxelles*, 39(1), 29-34. https://doi.org/10.30637/2018.17-073

02 AIM & HYPOTHESIS

Describe somatosensory prediction and repetition suppression in preterm neonates

Hypotheses


- Neonates will show **repetition suppression** in the **somatosensory and frontal cortices**: decrease of the neuronal activation between the **familiarization** and the **control** phases of the tactile stimulation sequence.
- Neonates will show a larger **early response** in the **somatosensory cortex** and a **late frontal response** to **deviance** (rare stimuli in the sequence).
- Neonates will have an **increase of HbO** during the **omission** in the somatosensory cortex, and a **late frontal response** on EEG.

03 MATERIAL & METHODS

Current population


$N = 30$ of 90 ($32,64 \pm 2,24$ SA at birth, 13 ♀), 13 had simultaneous EEG/fNIRS

Oddball-omission protocol




fNIRS acquisition and preprocessing

fNIRS at 690 & 830 nm (Imagent, ISS, Champaign IL, USA), 1 detector, 4 sources over fronto central areas



EEG acquisition and preprocessing

128 channels EEG (Magstim EGI, Eugene OR, USA), sampling rate : 1000 Hz



05 DISCUSSION

- Repeated stimulations elicit weaker HbO changes in the somatosensory cortex but increased HbO changes in the frontal cortex. We expected activation and not deactivation in the somatosensory cortex. In the frontal cortex, fNIRS and EEG show RS and are consistent with hypotheses.
- Deviance elicited a strong increase in HbO as well as an increased early electrical response in the frontal cortex. An increase in neuronal activity was observed only on EEG in the somatosensory cortex. Future measurements are needed to interpret fNIRS data.
- Omission elicited no change in the somatosensory cortex but elicited a decrease in HbO in the frontal cortex coupled with neuronal activity on EEG.


Conclusion

Preterm neonates form predictions and detect an error in the tactile environment.

The late negative components found on EEG during the deviance and the omission in the frontal cortex suggest that these prediction mechanisms could rely on a common frontal process as suggested by the predictive coding theory.

fNIRS measurement provide complementary information on tactile processing but more participants are needed to interpret fNIRS data further.

ABOUT US



Anne-Lise Marais, PhD student



@Annelisephd
<https://annelisemarais.github.io/>
<https://github.com/annelisemarais>
anne-lise.marais@unicaen.fr



Marie Anquetil,
PhD student



Victoria
Dumont,
PhD

DECODE

<https://decode.unicaen.fr/>

