



HAL
open science

Hydrodynamic limit of the Schelling model with spontaneous Glauber and Kawasaki dynamics

Florent Barret, Torri Niccolo

► **To cite this version:**

Florent Barret, Torri Niccolo. Hydrodynamic limit of the Schelling model with spontaneous Glauber and Kawasaki dynamics. 2023. hal-03996809v1

HAL Id: hal-03996809

<https://hal.science/hal-03996809v1>

Preprint submitted on 20 Feb 2023 (v1), last revised 14 Feb 2024 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hydrodynamic limit of the Schelling model with spontaneous Glauber and Kawasaki dynamics

Florent BARRET *

Niccolò TORRI *

Abstract

In the present article we consider the Shelling model, an agent-based model describing a segregation dynamics when we have a cohabitation of two social groups. As for several social models, the behavior of the Shelling model was analyzed by exploiting theoretical physics tools and computer simulations. This approach led to conjecture a phase diagram in which either different social groups were segregated in two large clusters or they were mixed. As far as the authors know, this model has never been considered by the mathematical community, a rigorous mathematical analysis of the model is one of the main novelties of the paper. We describe the Shelling model as a particle system model and, adding a Glauber and Kawasaki dynamics to the original Schelling dynamics, we prove the existence of an hydrodynamic limit described by a reaction-diffusion equation with a discontinuous non-linear reaction term. The existence and uniqueness of the solution is non trivial and the analysis of the limit PDE is interesting in its own. Based on our results, we conjecture the existence of a phase diagram in which we have a mixed, a segregated and a metastable segregation phase.

2010 Mathematics Subject Classification: 60K35, 82C22, 82D99.

Keywords: Shelling model, particle systems, hydrodynamics limit, reaction-diffusion equation, Ising model.

1 Introduction

Schelling's model of segregation was introduced by Thomas Schelling in 1971 [21, 22]. The original model is defined on a square grid of N sites (or, more generally, on a regular graph with N sites) where agents (individuals) belonging to two groups are randomly disposed. A local dynamic is defined by each agent, the agent located at a site of the grid compares its group with the group of its neighbors. More precisely, we fix a tolerance threshold $T \in [0, 1]$. We call r_x the fraction of neighbors belonging to the agent's group at site x and we say that the agent is satisfied if $r_x \geq T$. If the agent is unsatisfied, then he *moves* on a site that makes him satisfied. The dynamics is not unique, we briefly discuss this point here below. The process ends when all the agents are either satisfied or blocked (no move makes the agent satisfied). This model is thus characterized by a local dynamics which generates a global dynamics of the system. The study of this model enters naturally in the field of statistical physics which provides a large number of tools to study the properties of the model.

The parameters which characterize the model are

- (1) The size of the neighborhood,
- (2) The initial distribution of the agents,
- (3) The choice of the satisfaction condition,
- (4) The local dynamics between agents.

Several variants of Schelling's model have been considered in the recent literature in order to study the behavior of the model when the fundamental parameters are modified. We refer to [19] for a complete overview on the subject. The common result of the considered variants is the existence of three stationary states separated by two critical thresholds T_f and T_c (a *frozen* state $T < T_f$, a *segregated* state $T_f < T < T_c$ and a *mixed* state $T > T_c$) towards which the system evolves, suggesting an *universal* behavior of the model. When the system is open (agents can change the group) the behaviour of the system is expected to be symmetric with respect to $T = 1/2$, see [13].

In the present paper, we approach the model from a physical point of view, by interpreting the agent dynamics as a particle systems in interaction. This approach was adopted by the physical community to study this model,

*MODAL'X, UMR 9023, UPL, Univ. Paris Nanterre, F92000 Nanterre France. Email: florent.barret@parisnanterre.fr and niccolo.torri@parisnanterre.fr

see for instance [3, 9, 12, 13]. In particular we consider the setting where the points of the grid (a discrete torus $\mathbb{T}_N^d := (\mathbb{Z}/N\mathbb{Z})^d$, $d \geq 1$) are fully occupied and unsatisfied agents flip their state if it makes them satisfied (Glauber dynamics). The size of the neighborhood taken into account to compute the fraction r_x , grows at most logarithmically with N . Moreover we introduce random perturbations, either by flipping a state of an agent at rate β (spontaneous Glauber dynamics) or exchanging the position of two agents at rate αN^2 (accelerated Kawasaki dynamics).

Our main result is to prove an hydrodynamic limit described by a reaction-diffusion equation for the system and to give a complete description of the limit PDE that we get. In the case where the size of the neighborhood stays finite in the limit, we obtain a classical reaction-diffusion equation, this is the case where the size of the interaction term stays finite and thus microscopic. However, when the size of the neighborhood goes to infinity we get a non-linearity (the reaction term) which is discontinuous at two points. In this case, the interaction of the Schelling dynamics takes into account more and more agents but in the limit, the reaction term is still purely local.

In this ‘‘mesoscopic’’ limit, the existence and uniqueness of the solution of the reaction-diffusion equation with discontinuities is one of the major technical points of the paper. More precisely, the system does not have a unique solution for some class of initial condition, and some values of β , the parameter tuning the spontaneous Glauber dynamics. Finally, we conjecture the existence of a rich phase diagram in which beyond a mixed and a segregated phase there is a transition in between. In this phase, the system has the potential to show a metastable segregation: the mixing phase should be the most stable one but symmetrical stable segregation phases also exist. The critical points depend on the parameters β and T but not on α , see Figure 1. A rigorous proof of the phase diagram requires a delicate analysis of the local dynamics that goes beyond the techniques used in the present paper.

Our method of proof is based on the technique of the relative entropy method in the framework developed by Jara and Mezenes in [14], [15], and also used by Funaki and Tsunoda in [8]. However, in our setting, we needed to adapt their proofs to cover the case where the number of particles in the interaction would be going to infinity. We also prove existence and local uniqueness for a class of initial conditions of our limiting PDE, viewed as a differential inclusion.

A detailed plan of the method of proof and the article is given at the end of Section 3 containing the main result and assumptions. In the following Section, we define the model.

Acknowledgments

The authors thank Antoine Lucquiaud for the fruitful discussion on the model. This research has been conducted within the FP2M federation (CNRS FR 2036) and as part of the project Labex MME-DII (ANR11-LBX-0023-01).

2 The model

2.1 Configurations

For $N \in \mathbb{N} = \{1, 2, 3, \dots\}$ we let $\mathbb{T}_N^d = (\mathbb{Z}/N\mathbb{Z})^d$ be the discrete torus and let $\Omega_N = \{0, 1\}^{\mathbb{T}_N^d}$ be the space of all possible configurations. We call $\eta \in \Omega_N$ a configuration and $i \in \mathbb{T}_N^d$ a site. Let $\mathcal{V}_N \subset \mathbb{T}_N^d \setminus \{0\}$ be a bounded set. We say that two sites $i, j \in \mathbb{T}_N^d$ are neighbors if $i - j \in \mathcal{V}_N$. We denote $K_N = |\mathcal{V}_N|$ its cardinality. For a configuration $\eta \in \Omega_N$ and a site $i \in \mathbb{T}_N^d$ we let

$$r_i(\eta) = \frac{1}{K_N} \sum_{j \in \mathcal{V}_N} \mathbb{1}_{\{\eta_i = \eta_{i+j}\}} \quad \text{and} \quad \rho_i(\eta) = \frac{1}{K_N} \sum_{j \in \mathcal{V}_N} \eta_{i+j}. \quad (2.1)$$

The quantity $\rho_i(\eta)$ is the mean field of η on the neighborhood $\mathcal{V}_N + i$. Let us observe that $\rho_i(\eta)$ is independent of η_i . We note that

$$r_i(\eta) = \rho_i(\eta) \mathbb{1}_{\{\eta_i=1\}} + (1 - \rho_i(\eta)) \mathbb{1}_{\{\eta_i=0\}}. \quad (2.2)$$

For a given configuration, we now introduce the definition of *stable*, *unstable* and *potentially stable site*.

Definition 2.1. For a given site i , let us denote η^i the configuration where we change η_i to $1 - \eta_i$.

Let $T \in [0, 1]$. If $r_i(\eta) < T$, the site i is said *unstable* for η , otherwise if $r_i(\eta) \geq T$ the site is said *stable* for η . An *unstable site* i for η which is *stable* for η^i is said *potentially stable*.

Note that $r_i(\eta^i) = 1 - r_i(\eta)$. Thus

1. a site i is potentially stable if and only if $r_i(\eta) < T$ and $r_i(\eta) \leq 1 - T$. In particular if $T \leq \frac{1}{2}$ an unstable site for η is automatically potentially stable.
2. if $T > \frac{1}{2}$ and $1 - T < r_i(\eta) < T$, we have $r_i(\eta^i) < T$ and the site i is unstable for η and η^i .

2.2 Infinitesimal generator, construction of the process

Fix $\alpha > 0$ and $\beta > 0$. Let us consider the following dynamics : starting from a configuration η

1. if a site i is potentially stable, we flip the value at i with rate 1,
2. two nearest-neighbors i and j exchange their values with rate αN^2 (accelerated Kawasaki dynamics),
3. a site i can change its value at rate $\beta > 0$ (spontaneous Glauber dynamics).

This dynamics defines an infinitesimal generator \mathcal{L}_N defined for F cylinder function on Ω_N by

$$\mathcal{L}_N F(\eta) = \sum_{i \in \mathbb{T}_N^d} (F(\eta^i) - F(\eta)) (\mathbb{1}_{\{r_i(\eta) < T, r_i(\eta) \leq 1 - T\}} + \beta) + \alpha N^2 \sum_{\substack{i, j \in \mathbb{T}_N^d \\ |i - j| = 1}} (F(\eta^{ij}) - F(\eta)), \quad (2.3)$$

where η^{ij} is the configuration where the values at site i and j have been exchanged.

The following proposition states that the process is well defined, since the state space is finite.

Proposition 2.1. *Given an initial configuration η_0 , \mathcal{L}_N is the infinitesimal generator of a Feller process, denoted $(\eta^N(t))_{t \geq 0}$.*

We let μ_t^N be the distribution of $\eta^N(t)$.

Remark 2.1. *In this article we focus on the compact setting (torus) because a non compact framework, as \mathbb{R}^d , presents technical problems for the convergence of the process, nevertheless the discrete model can be well defined on \mathbb{Z}^d (cf. (2.3) and Proposition 2.1).*

3 Main results

We let $u_0^N(i) := \mathbb{E}_{\mu_N}[\eta_i^N(0)]$, $i \in \mathbb{T}_N^d$ be the initial distribution of our process, that is, $\eta_i^N(0)$ is distributed as a Bernoulli of parameter $u_0^N(i)$.

For a vector v , $|v|$ denotes its euclidean norm and $|v|_\infty$ its uniform norm.

Assumption 1. *Assumptions on \mathcal{V}_N .*

1. Let $\ell_{\mathcal{V}}$ be the diameter of \mathcal{V} . Then, $\ell_{\mathcal{V}} \leq \delta (\log N)^{\frac{1}{d}}$ for some $\delta > 0$.
2. If $d = 1$, suppose that $\mathcal{V}_N \subset \mathbb{Z} \setminus \mathbb{N}$.

Assumption 2. *Assumptions on u_0^N .*

1. There exists $\varepsilon > 0$ such that $\varepsilon \leq u_0^N(i) \leq 1 - \varepsilon$ uniformly on $i \in \mathbb{T}_N^d$ and $N \in \mathbb{N}$.
2. There exists $C_0 > 0$ independent of N such that $|\nabla u_0^N(i)|_\infty \leq \frac{C_0}{N}$, where $\nabla u_0^N(i) = (u_0^N(i + e_k) - u_0^N(i))_{k=1}^d$, with $e_k \in \mathbb{Z}^d$ the unit vector of direction k .
3. Let $\nu_{u_0^N}^N = \otimes_{i \in \mathbb{T}_N^d} \mathbb{B}(u_0^N(i))$ be the law of a sequence of independent Bernoulli of parameter $u_0^N(i)$. Suppose that $\mathcal{H}(\mu_0^N | \nu_{u_0^N(0)}^N) = O(N^{d-\varepsilon_0})$ for some $\varepsilon_0 > 0$ small, where $\mathcal{H}(\mu | \nu)$ is the relative entropy of $\mu \ll \nu$,

$$\mathcal{H}(\mu | \nu) := \int \frac{d\mu}{d\nu} \log \left(\frac{d\mu}{d\nu} \right) d\nu \quad (3.1)$$

4. Let $\tilde{u}_0^N(x)$ be the linear interpolation on $\mathbb{T}^d = (\mathbb{R}/\mathbb{Z})^d$, the d dimensional torus, of $u_0^N(i)$ such that $\tilde{u}_0^N(i/N) = u_0^N(i)$. Then, there exists $u_0 \in \mathcal{C}(\mathbb{T}^d)$ such that \tilde{u}_0^N converges to u_0 in $\mathcal{C}(\mathbb{T}^d)$.

Remark 3.1. *The assumption 1(2.) is only technical and it could be removed by considering the dimension $d = 1$ separately from the rest of the dimensions, cf. Remark 4.3.*

Define

$$\pi_t^N = \pi_t^N(\eta, dv) = \frac{1}{N^d} \sum_{i \in \mathbb{T}_N^d} \eta_i(t) \delta_{\frac{i}{N}}(dv) \quad (3.2)$$

the empirical measure associated to the Markov process η where the space is rescaled by $\frac{1}{N}$, which is a positive measure on \mathbb{T}^d .

We now state our main result, which concerns the convergence of subsequences of the empirical measure.

Theorem 3.1. *Under Assumptions 1 and 2, for every test function $\varphi : \mathbb{T}^d \rightarrow \mathbb{R}$ and for every $\varepsilon > 0$, there exists $\tau > 0$ and an increasing sequence (N_i) such that*

$$\lim_{i \rightarrow +\infty} \mu^{N_i} \left(\left| \langle \pi^{N_i}, \varphi \rangle - \langle u, \varphi \rangle \right| > \varepsilon \right) = 0, \quad \forall t \in [0, \tau],$$

where $\langle \pi^N, \varphi \rangle$ and $\langle u, \varphi \rangle$ denote the integral of φ with respect to the measure π^N or $u(x)dx$ respectively. The function $u = u(t, x)$ is a density solution of the following reaction-diffusion equation

$$\begin{cases} \partial_t u(t, x) = 2\alpha \Delta u(t, x) + \beta(1 - 2u(t, x)) + g_\infty(u(t, x)), \\ u(0, x) = u_0(x), \end{cases} \quad (3.3)$$

on the torus \mathbb{T}^d , with $g_\infty(p) := (1 - p)\mathbb{1}_{\{1-p < \min(T, 1-T)\}} - p\mathbb{1}_{\{p \leq \min(T, 1-T)\}}$. Moreover if u_0 (the limit of the initial condition, according to Assumption 2-4) is such that

1. $u_0 \in \mathcal{C}^1(\mathbb{T}^d)$, with ∇u_0 Lipschitz,
2. $\nabla u_0(p) \neq 0$ for $p = \min(T, 1 - T)$ and $p = 1 - \min(T, 1 - T)$,

then the solution of Equation (3.3) is unique on $[0, \tau]$ for some $\tau > 0$ and the convergence holds for the sequence itself.

3.1 Organisation of the paper

To prove Theorem 3.1 we first prove that the empirical measure is close to a discrete measure u^N which is a solution of a discrete analogous of (3.3), this is Theorem 4.2. Its proof is based on an entropy method approach in which the relative entropy between μ^N and $\nu_{u^N}^N$, see Theorem 4.1. Even if this technique is quite standard in the particle system theory, some new technical estimations arising from the geometry of the system are needed. In Section 5 we discuss some central technical estimations about u^N , in order to describe the behaviour of the discrete model. Then, in Section 6 we discuss the existence and uniqueness of (3.3) and in Section 7 we show the convergence of the u^N toward the density u by completing the proof of Theorem 3.1. We stress that the proof of the existence and uniqueness is not standard and the analysis of this PDE is interesting in its own.

In Section 3.2 we state our conjecture on the phase diagram of the model.

3.2 Conjecture on the phase diagram

In this Section, we discuss the phase diagram that describes the mixed and segregated phases. We start by setting the Equation (3.3) in a more convenient form. Set $p_0(T) := \min(T, 1 - T) \in [0, \frac{1}{2}]$. For $p \in [0, 1]$ we define

$$\gamma_{\infty, \beta}(p) = \begin{cases} \beta \left(p - \frac{1}{2} \right)^2 + \frac{1}{2} (p^2 - p_0(T)^2) & \text{for } 0 \leq p < p_0(T), \\ \beta \left(p - \frac{1}{2} \right)^2 & \text{for } p_0(T) \leq p < 1 - p_0(T), \\ \beta \left(p - \frac{1}{2} \right)^2 + \frac{1}{2} ((1 - p)^2 - p_0(T)^2) & \text{for } 1 - p_0(T) \leq p \leq 1. \end{cases} \quad (3.4)$$

Our conjecture is based on the analysis of $\gamma_{\infty, \beta}$ and it is represented in Figure 1. We observe that $\gamma_{\infty, \beta}$ is continuous and satisfies $\gamma_{\infty, \beta}(p) = \gamma_{\infty, \beta}(1 - p)$. For $p \neq p_0(T), 1 - p_0(T)$, we have that $\gamma'_{\infty, \beta}(p) = -\beta(1 - 2p) - g_\infty(p)$ and (3.3) can be written as

$$\partial_t u(t, x) = 2\alpha \Delta u(t, x) - \gamma'_{\infty, \beta}(u(t, x)). \quad (3.5)$$




Figure 1: Representation of the different phases of the system as function of the parameter $p_0(T) \in [0, \frac{1}{2}]$. When T is close to 0 or 1 ($p_0(T) \in (0, p^\ell)$) we do not have segregation (red parts) and typical configurations are provided by a mixing of 0 and 1. If T is close to 1/2 ($p_0(T) \in (p^m, \frac{1}{2})$) we have segregation (green parts): a very large parts of the configuration are composed of 0 or 1. We have also intermediate values of T ($p_0(T) \in (p^\ell, p^m)$) for which the segregation is metastable (yellow parts).

To discuss the phase transition we can look at the structure of $\gamma_{\infty, \beta}(p)$. The function $p \mapsto \beta \left(p - \frac{1}{2}\right)^2 + \frac{1}{2} (p^2 - p_0(T)^2)$ has a unique minimum at $p = p^\ell := \frac{\beta}{1+2\beta}$. Therefore, if $0 \leq p^\ell < p_0(T) < \frac{1}{2}$, the function $\gamma_{\infty, \beta}$ has three regular minima : $p^c := \frac{1}{2}$, p^ℓ and $p^r := 1 - p^\ell$. Note that

$$\gamma_{\infty, \beta}(p^c) = 0 \quad \text{and} \quad \gamma_{\infty, \beta}(p^r) = \gamma_{\infty, \beta}(p^\ell) = \frac{\beta}{4(1+2\beta)} - \frac{p_0(T)^2}{2}.$$

Then we get that if $p_0(T) < p^m := \sqrt{\frac{\beta}{2(1+2\beta)}}$, we have $\gamma_{\infty, \beta}(p^c) < \gamma_{\infty, \beta}(p^\ell)$ and if $p^m < p_0(T)$, we have that $\gamma_{\infty, \beta}(p^c) > \gamma_{\infty, \beta}(p^\ell)$. If $p^\ell > p_0(T)$, p^c is the only minimum.

The two thresholds for $p_0(T)$ are then p^ℓ and p^m , see Figure 1. Since $p^\ell < p^m$, we have the following picture: as T is close to 0 and below p^ℓ , we have a unique minimum of $\gamma_{\infty, \beta}$, so that typical configurations are close to $p = 1/2$ which is of lowest energy $\gamma_{\infty, \beta}$. It means that, at equilibrium, we expect a configuration balanced between 0 and 1 and we do not have segregation. Then, as T goes above the threshold p^ℓ but stays below p^m , other minima at $p = p^\ell$ and $p = p^r$ appear, and these two configurations are metastable since their energy is higher, so we can have segregation for a small proportion of the time. The next threshold is p^m , at which the two metastable configurations become stable and $p = 1/2$ is the metastable one so that we expect stable segregation. For T above $\frac{1}{2}$ the picture is symmetric.

Remark 3.2. *The hydrodynamic limit (3.3) has, at least, two different formulations as a gradient flow :*

1. *in the classical $L^2(\mathbb{T}^d)$ setting, with the potential \mathcal{F} defined for $u : \mathbb{T}^d \rightarrow [0, 1]$*

$$\mathcal{F}(u) = \int \alpha \|\nabla u\|^2 + \gamma_{\infty, \beta}(u), \quad (3.6)$$

Equation (6.2) can be written as $\partial_t u = -\delta \mathcal{F}(u)$ where $\delta \mathcal{F}$ denotes the Fréchet derivative of \mathcal{F} .

2. *in a Wasserstein-like setting defined in [18] with the entropy potential \mathcal{H} , for $u : \mathbb{T}^d \rightarrow]0, 1]$, and $\xi : \mathbb{T}^d \rightarrow \mathbb{R}$ (seen as an element in the tangent bundle)*

$$\mathcal{H}(u) = \frac{1}{2} \int (2u \log(2u) - 2u + 1) \geq 0, \quad \mathcal{K}(u)\xi = -2\alpha \nabla \cdot (u \nabla \xi) + \frac{\gamma'_{\infty, \beta}(u)}{\log(2u)} \xi, \quad (3.7)$$

Equation (3.3) can be written as

$$\partial_t u = -\mathcal{K}(u)(\delta \mathcal{H}).$$

Since $\delta \mathcal{H} = \log(2u)$ and $\mathcal{K}(u)(\delta \mathcal{H}) = -2\alpha \Delta u + \gamma'_{\beta, \infty}(u)$.

Both formulations could be useful to establish a rigorous proof of the phase diagram given in Figure 1. In particular, along a gradient flow the potential is non-increasing, thus for all $t > 0$, along a solution u we have $\mathcal{F}(u(t)) \leq \mathcal{F}(u(t=0))$ and if u converges to a stationary solution v , it must be a stationary point of \mathcal{F} (i.e. $\delta \mathcal{F}(v) = 0$).

For the second formulation, note that

$$\frac{\gamma'_{\infty, \beta}(u)}{\log(2u)} > 0, u \in]0, 1] \iff p_0(T) < p^\ell = \frac{\beta}{1+2\beta} \iff \beta > \frac{p_0(T)}{1-2p_0(T)}. \quad (3.8)$$

We are thus in the mixing phase of the diagram and $\mathcal{K}(u)$ is positive definite in the sense that

$$\int \xi \mathcal{K}(u) \xi = \int 2\alpha u |\nabla \xi|^2 + \frac{\gamma'_{\infty, \beta}(u)}{\log(2u)} \xi^2 \geq 0. \quad (3.9)$$

Then, one can prove that along a solution u , we get that :

$$\frac{d}{dt} \mathcal{H}(u) = \int \partial_t u \delta \mathcal{H}(u) \leq - \int \gamma'_{\infty, \beta}(u) \log(2u) \leq -c \mathcal{H}(u) \quad (3.10)$$

where $c = \beta - \frac{\rho_0(T)}{1-2\rho_0(T)} > 0$, and we get that $\mathcal{H}(u(t)) \leq \mathcal{H}(u(0))e^{-ct}$. Thus $\mathcal{H}(u(t))$ goes to 0 as $t \rightarrow \infty$, this entails that u converges to the only stationary point of \mathcal{H} which is the constant $\frac{1}{2}$. Heuristically, it suggests that an exponential relaxation is taking place in the mixing phase of Figure 1.

4 Relative entropy method

Using the *relative entropy* method, in this section we prove that the empirical measure $\pi^N(t)$ is close to $u^N(t) = (u^N(t, i))_{i \in \mathbb{T}_N^d}$, the solution of a suitable discrete PDE.

In order to state the main result of this section, we observe that the generator \mathcal{L}_N in (2.3) can be written as $\mathcal{L}_N = \mathcal{G}_N + 2\alpha N^2 \mathcal{K}_N$ where \mathcal{G}_N is the generator which describes the Glauber-Shelling dynamics and \mathcal{K}_N the generator which describes the Kawasaki dynamics, that is,

$$\mathcal{G}_N F(\eta) = \sum_{i \in \mathbb{T}_N^d} \left(\mathbb{1}_{\{r_i(\eta) < T, r_i(\eta) \leq 1-T\}} + \beta \right) (F(\eta^i) - F(\eta)), \quad (4.1)$$

$$\mathcal{K}_N F(\eta) = \frac{1}{2} \sum_{\substack{i, j \in \mathbb{T}_N^d \\ |i-j|=1}} (F(\eta^{ij}) - F(\eta)). \quad (4.2)$$

Let us stress that \mathcal{G}_N can be written as follows

$$\mathcal{G}_N F(\eta) = \sum_{i \in \mathbb{T}_N^d} (c_i(\eta) + \beta) (F(\eta^i) - F(\eta)), \quad (4.3)$$

where $c_i(\eta)$ is a local function which describes the dynamics (Glauber-Schelling dynamics). To be more precise, we write $c_i(\eta) = c_0(\tau_i \eta)$ where c_0 is the flipping rate of a particle at the origin, that is,

$$c_0(\eta) := \mathbb{1}_{\{r_0(\eta) < T, r_0(\eta) \leq 1-T\}} \quad (4.4)$$

and $(\tau_i \eta)_j = \eta_{i+j}$, likewise τ_i acts on u . Note that c_0 is a random variable and take the value 0 or 1.

We let

$$\kappa_N = \kappa_N(T) := \min \left\{ \lfloor K_N T \rfloor - 1; \lfloor K_N (1-T) \rfloor \right\}. \quad (4.5)$$

We observe that $\lim_{n \rightarrow +\infty} \frac{\kappa_N}{K_N} = \min(T, 1-T)$. Since $c_0(\eta) = \mathbb{1}_{\{r_0(\eta) \leq \frac{\kappa_N}{K_N}\}} = \mathbb{1}_{\{r_0(\eta) \leq \frac{\kappa_N}{K_N}, \eta_0=0\}} + \mathbb{1}_{\{r_0(\eta) \leq \frac{\kappa_N}{K_N}, \eta_0=1\}}$, we define

$$c_0^+(\eta) := \mathbb{1}_{\{\rho_0(\eta) \geq 1 - \frac{\kappa_N}{K_N}\}} \quad \text{and} \quad c_0^-(\eta) := \mathbb{1}_{\{\rho_0(\eta) \leq \frac{\kappa_N}{K_N}\}}, \quad (4.6)$$

so that $c_0(\eta) = c_0^+(\eta)(1 - \eta_0) + c_0^-(\eta)\eta_0$, by (2.2). The functions c^+ and c^- can be viewed as the rate of creation and annihilation of a particle at $i = 0$ respectively.

For any function $u = (u_i)_{i \in \mathbb{T}_N^d}$ we define

$$v_u(d\eta) = v_u^N(d\eta) := \bigotimes_{i \in \mathbb{T}_N^d} B(u_i) \quad (4.7)$$

and we let $c_0^+(u)$ and $c_0^-(u)$ be the expectation of $c_0^+(\eta)$ and $c_0^-(\eta)$ under v_u , that is,

$$c_0^+(u) := \mathbb{P}_{v_u} \left(\rho_0(\eta) \geq 1 - \frac{\kappa_N}{K_N} \right) \quad \text{and} \quad c_0^-(u) := \mathbb{P}_{v_u} \left(\rho_0(\eta) \leq \frac{\kappa_N}{K_N} \right). \quad (4.8)$$

We finally define

$$G(u) := c_0^+(u)(1 - u_0) - c_0^-(u)u_0 \quad \text{and} \quad G(i, u) := G(\tau_i u). \quad (4.9)$$

Let $u^N(t) = (u^N(t, i))_{i \in \mathbb{T}_N^d}$ be the solution of

$$\begin{cases} \partial_t u^N(t, i) = 2\alpha N^2 \Delta u^N(t, i) + \beta(1 - 2u^N(t, i)) + G(i, u^N) \\ u^N(t, i) = u_0^N(i), \end{cases} \quad (4.10)$$

where $\Delta u^N(t, i)$ is the discrete Laplacian on the torus. Note that (4.10) can be interpreted as a discretized version of (3.3).

In the following result we prove that the relative entropy of $v_{u^N(t)}^N$ (cf. (4.7)) with respect to μ_t^N stays small in time, if it is small at $t = 0$.

Theorem 4.1. *Under Assumptions 1 and 2, with $\delta > 0$ sufficiently small, we have that for some $\varepsilon > 0$ small*

$$\mathcal{H}(\mu_t^N | v_{u^N(t)}^N) = O(N^{d-\varepsilon}), \quad \forall t \in [0, \tau] \quad (4.11)$$

with $\tau = \tau(\delta) > 0$.

Theorem 4.1 implies Theorem 3.1 for u^N defined in (4.10). More precisely, we consider u^N as a measure on \mathbb{T}^d , that is,

$$u^N(t, dv) := \frac{1}{N^d} \sum_{i \in \mathbb{T}_N^d} u^N(t, i) \delta_{\frac{i}{N}}(dv), \quad (4.12)$$

so that we can state the following result.

Theorem 4.2. *Under Assumptions 1 and 2, for every test function $\varphi : \mathbb{T}^d \rightarrow \mathbb{R}$ and for every $\delta > 0$ there exists $\tau > 0$ such that*

$$\lim_{N \rightarrow +\infty} \mu^N \left(\left| \langle \pi^N, \varphi \rangle - \langle u^N, \varphi \rangle \right| > \delta \right) = 0, \quad \forall t \in [0, \tau].$$

Proof. We recall the entropy inequality stated for a set \mathcal{A} and two measures $\nu \ll \mu$, cf. A1.8.2 of [16] or Section 2.2 of [8],

$$\mu(\mathcal{A}) \leq \frac{\log 2 + \mathcal{H}(\mu | \nu)}{\log \left(1 + \frac{1}{\nu(\mathcal{A})} \right)}. \quad (4.13)$$

For a given test function φ and $\delta > 0$, we let

$$\mathcal{A}_{N,t,\varphi}^\delta := \{ \eta \in \Omega_N : |\langle \pi^N, \varphi \rangle - \langle u^N, \varphi \rangle| > \delta \}, \quad (4.14)$$

so that the proof follows by Theorem 4.1 and (4.13) if

$$v_{u^N(t)}^N(\mathcal{A}_{N,t,\varphi}^\delta) \leq e^{-C_\delta N^d}. \quad (4.15)$$

Since $u^N \in (0, 1)$ (cf. Proposition 5.2), the proof of (4.15) is model independent and follows line by line the proof of Proposition 2.2 in [8]. \square

Remark 4.1. *Let us note that c_0 can be expressed as a polynomial on the variables η_j 's, as in relation (1.5) of [8]. This remark will be useful in the sequel of the paper. For this purpose, let $A \subset \mathcal{V}_N$, we denote :*

$$c_A^+(\eta) = \prod_{j \in A} (1 - \eta_j) \prod_{j \in \bar{A} \cap \mathcal{V}_N} \eta_j \quad \text{and} \quad c_A^-(\eta) = \prod_{j \in A} \eta_j \prod_{j \in \bar{A} \cap \mathcal{V}_N} (1 - \eta_j) = c_A^+(1 - \eta) \quad (4.16)$$

where $1 - \eta$ is the configuration with $(1 - \eta)_i = 1 - \eta_i$ since $r_0(\eta) = r_0(1 - \eta)$. Note that

$$c_A^+(\eta) = \begin{cases} 1 & \text{if } \eta_j = 0 \text{ for } j \in A \text{ and } \eta_j = 1 \text{ for } j \in \bar{A} \cap \mathcal{V}_N \\ 0 & \text{otherwise.} \end{cases} \quad (4.17)$$

By an abuse of notation, for a function $u = (u_j)_{j \in \mathbb{T}_N^d}$ we let let also $c_A^+(u) = \prod_{j \in A} (1 - u_j) \prod_{j \in \bar{A} \cap \mathcal{V}_N} u_j$ and accordingly for $c_A^-(u)$. By (4.6) we have that

$$c_0^+(\eta) = \sum_{k=0}^{\kappa_N} \mathbb{1}_{\{r_0(\eta) = \frac{k}{\kappa_N}, \eta_0 = 0\}} = (1 - \eta_0) \sum_{\substack{A \subset \mathcal{V}_N \\ |A| \leq \kappa_N}} c_A^+(\eta). \quad (4.18)$$

Accordingly, we have

$$c_0^-(\eta) = \sum_{k=0}^{\kappa_N} \mathbb{1}_{\{r_0(\eta) = \frac{k}{\kappa_N}, \eta_0=1\}} = \eta_0 \sum_{\substack{A \subset \mathcal{V}_N \\ |A| \leq \kappa_N}} c_A^-(\eta). \quad (4.19)$$

In the rest of this section we prove Theorem 4.1. The strategy that we use follows the one used to prove the analogous result in [8] and [15], but some extra-technicality is required due to the geometry of our problem.

4.1 Proof of Theorem 4.1

To make the notation lighter, for $t \geq 0$, $j \in \mathbb{T}_N^d$, $p \in (0, 1)$ and u^N the solution of (4.10), we let

$$u_j(t) := u^N(t, j), \quad \chi(p) := p(1-p), \quad \omega_j := \frac{\eta_j - u_j}{\chi(u_j)}. \quad (4.20)$$

and, more generally, whenever the context is clear we omit the superscript N , so that $v_{u^N(t)}$ and μ_t^N will be denoted simply by $v_{u(t)}$ and μ_t respectively.

To compare μ and v_u we introduce

$$\vartheta_\alpha(d\eta) := \bigotimes_{i \in \mathbb{T}_N^d} B(\alpha) \quad (4.21)$$

be a sequence of independent Bernoulli of parameter $\alpha \in (0, 1)$ defined on the space of configurations. We define

$$f_t := \frac{d\mu_t}{dv_{u(t)}} \quad \text{and} \quad \psi_t := \frac{dv_{u(t)}}{d\vartheta_\alpha}. \quad (4.22)$$

We let $\mathcal{H}_N(t) := \mathcal{H}(\mu_t | v_{u(t)})$ be the relative entropy of $v_{u(t)}$ with respect to μ_t cf. (3.1). We have all the ingredients to state Yau's inequality in our context (cf. proof Lemma A.1 in [15]).

Proposition 4.3. *For any $t \geq 0$ we have that*

$$\partial_t \mathcal{H}_N(t) \leq - \int \Gamma_N(\sqrt{f_t(\eta)}) v_{u(t)}(d\eta) + \int f_t(\eta) \left[\mathcal{L}_N^{*, v_{u(t)}} \mathbf{1} - \partial_t \log \psi_t \right] v_{u(t)}(d\eta), \quad (4.23)$$

where $\mathcal{L}_N^{*, v_{u(t)}}$ is the adjoint of \mathcal{L}_N with respect to the measure $v_{u(t)}$ and $\Gamma_N(h)(\eta) = \mathcal{L}_N h^2(\eta) - 2h(\eta) \mathcal{L}_N h(\eta)$ is the carré du champ operator.

Proof. Let us observe that

$$\begin{aligned} \mathcal{H}_N(t) &= \int \frac{d\mu_t}{dv_{u(t)}}(\eta) \log \left(\frac{d\mu_t}{dv_{u(t)}}(\eta) \right) v_{u(t)}(d\eta) = \int f_t(\eta) \log \left(f_t(\eta) \right) v_{u(t)}(d\eta) \\ &= \int f_t(\eta) \psi_t(\eta) \log \left(\frac{f_t(\eta) \psi_t(\eta)}{\psi_t(\eta)} \right) \vartheta_\alpha(d\eta). \end{aligned} \quad (4.24)$$

The forward Fokker-Plank equation tells us that $f_t \psi_t = \frac{d\mu_t}{d\vartheta_\alpha}(\eta)$ is solution of $\frac{d}{dt}(f_t \psi_t) = \mathcal{L}_N^{*, \alpha}(f_t \psi_t)$, so that by deriving in time (4.24) we obtain

$$\begin{aligned} \partial_t \mathcal{H}_N(t) &= \int \left\{ \mathcal{L}_N^{*, \alpha}(f_t \psi_t)(\eta) \log(f_t(\eta)) \right. \\ &\quad \left. + \mathcal{L}_N^{*, \alpha}(f_t \psi_t)(\eta) - f_t(\eta) \psi_t(\eta) \partial_t \log(\psi_t(\eta)) \right\} \vartheta_\alpha(d\eta). \end{aligned} \quad (4.25)$$

Since $\int \mathcal{L}_N^{*, \alpha}(f_t \psi_t)(\eta) \vartheta_\alpha(d\eta) = \int (f_t \psi_t)(\eta) \mathcal{L}_N \mathbf{1} \vartheta_\alpha(d\eta) = 0$ we get that the right term of (4.25) equals

$$\int f_t(\eta) \left[\mathcal{L}_N \log(f_t(\eta)) - \partial_t \log(\psi_t(\eta)) \right] v_{u(t)}(d\eta). \quad (4.26)$$

The result is then a consequence of the following inequality: for any $a, b > 0$,

$$a(\log(b) - \log(a)) \leq 2\sqrt{a}(\sqrt{b} - \sqrt{a}) = (b - a) - (\sqrt{b} - \sqrt{a})^2.$$

Indeed, if we write $\mathcal{L}_N h(\eta) = \sum_{\eta'} r(\eta, \eta')(h(\eta') - h(\eta))$, we get,

$$\begin{aligned} h(\eta)\mathcal{L}_N \log(h(\eta)) &= \sum_{\eta'} r(\eta, \eta')h(\eta)(\log(h(\eta')) - \log(h(\eta))) \\ &\leq \sum_{\eta'} r(\eta, \eta') \left[(h(\eta') - h(\eta)) - (\sqrt{h(\eta')} - \sqrt{h(\eta)})^2 \right] = \mathcal{L}_N h(\eta) - \Gamma_N(\sqrt{h})(\eta). \end{aligned}$$

Applying this result to the first term of the integral in (4.26) we get (4.23). \square

We define the *current* $\mathcal{J}_t = \mathcal{J}_t^N(\eta)$ as

$$\mathcal{J}_t := \mathcal{L}_N^{*,v_{u(t)}} \mathbf{1} - \partial_t \log \psi_t. \quad (4.27)$$

Our main goal is to estimate the current \mathcal{J}_t to control the right hand side of (4.23). In particular we show that

$$\partial_t \mathcal{H}_N(t) \leq C \ell_V^d \left(\mathcal{H}_N(t) + O(N^{d-a}) \right) \quad (4.28)$$

with $a \in (0, 1)$ if $d \geq 2$ and $a \in (0, \frac{1}{2})$ if $d = 1$. Gronwall's inequality gives

$$\mathcal{H}_N(t) \leq \left(\mathcal{H}_N(0) + tO(N^{d-a}) \right) e^{C \ell_V^d t}. \quad (4.29)$$

Since $e^{C \ell_V^d t} \leq N^{C \delta t}$ by Assumption 1, the proof of Theorem 4.1 is complete.

4.1.1 The current \mathcal{J}_t^N

To control $\mathcal{L}_N^{*,v_{u(t)}} \mathbf{1}$ we have to compute the adjoint of \mathcal{K}_N and of \mathcal{G}_N , cf. (4.1) and (4.2). We follow the computations done in [8]. By Lemma 2.4 of [8], we get

$$\mathcal{K}_N^{*,v_{u(t)}} \mathbf{1} = -\frac{1}{2} \sum_{\substack{i,j \in \mathbb{T}_N^d, \\ |i-j|=1}} (u_i - u_j)^2 \omega_j \omega_i + \sum_{i \in \mathbb{T}_N^d} (\Delta u)_i \omega_i \quad (4.30)$$

where $(\Delta u)_i = \sum_{j \in \mathbb{T}_N^d, |i-j|=1} (u_j - u_i)$ is the discrete Laplacian. Since c_0 satisfies the condition (1.5) of [8] (see Remark 4.1), by Lemma 2.5 of [8] we get

$$\begin{aligned} \mathcal{G}_N^{*,v_{u(t)}} \mathbf{1} &= \sum_{i \in \mathbb{T}_N^d} (c_i^+(\eta)(1 - u_i) - c_i^-(\eta)u_i + \beta(1 - 2u_i)) \omega_i \\ &= \sum_{i \in \mathbb{T}_N^d} ((c_i^+(\eta) - c_i^+(u))(1 - u_i) - (c_i^-(\eta) - c_i^-(u))u_i) \omega_i \\ &\quad + \sum_{i \in \mathbb{T}_N^d} (c_i^+(u)(1 - u_i) - c_i^-(u)u_i + \beta(1 - 2u_i)) \omega_i. \end{aligned} \quad (4.31)$$

In the second equality we centered the variables $c_i^+(\eta)$ $c_i^-(\eta)$ since under v_u , $c_i^+(\eta)$ and $c_i^-(\eta)$ are Bernoulli random variables with expectation $c_i^+(u)$ and $c_i^-(u)$ respectively. Finally, by Lemma 2.6 of [8] we get

$$\partial_t \log \psi_t(\eta) = \sum_{i \in \mathbb{T}_N^d} (\partial_t u_i) \omega_i. \quad (4.32)$$

Summarizing, we obtain the following result.

Proposition 4.4. *The current $\mathcal{J}_t(\eta)$ satisfies*

$$\begin{aligned} \mathcal{J}_t(\eta) &= \sum_{i \in \mathbb{T}_N^d} (-\partial_t u_i + 2\alpha N^2 (\Delta u)_i + \beta(1 - 2u_i) + G(i, u)) \omega_i \\ &\quad - V(u, \eta) + V^+(u, \eta) - V^-(u, \eta), \end{aligned} \quad (4.33)$$

where G was defined in (4.9) and

$$V^+(u, \eta) = \sum_{i \in \mathbb{T}_N^d} (c_i^+(\eta) - c_i^+(u)) (1 - u_i) \omega_i, \quad (4.34)$$

$$V^-(u, \eta) = \sum_{i \in \mathbb{T}_N^d} (c_i^-(\eta) - c_i^-(u)) u_i \omega_i, \quad (4.35)$$

$$V(u, \eta) = -\alpha N^2 \sum_{\substack{i, j \in \mathbb{T}_N^d, \\ |i-j|=1}} (u_i - u_j)^2 \omega_j \omega_i. \quad (4.36)$$

In particular, if u satisfies (4.10) the current reduces to the second line.

In the rest of the section we provide estimates of V^+ , V^- and V .

4.1.2 Estimates of V^+ and V^-

Let us denote $\{e_1, e_2, \dots, e_d\}$ the canonical basis of \mathbb{Z}^d . For $\varphi : \mathbb{T}_N^d \rightarrow \mathbb{R}$ and $i \in \mathbb{T}_N^d$ and $k \in \{1, \dots, d\}$, let $\nabla_k \varphi(i) = \varphi(i + e_k) - \varphi(i)$. We denote $\|\nabla \varphi\|_\infty = \max_{i, k} |\nabla_k \varphi(i)|$.

We note that

$$V^-(u, \eta) = -V^+(1 - u, 1 - \eta), \quad (4.37)$$

so that the bound for V^+ can be transferred to V^- , see Remark 4.2. In the following we get an upper-bound for V^+ . Denote $\vartheta_i^+ = c_i^+(\eta) - c_i^+(u)$ and $\omega_i^+ = (1 - u_i) \omega_i = \frac{\eta_i - u_i}{u_i}$.

Then

$$V^+(u, \eta) = \sum_{i \in \mathbb{T}_N^d} \vartheta_i^+ \omega_i^+ \quad (4.38)$$

To bound $V^+(u, \eta)$ we follow the method used by Jara and Menezes [15] and by Funaki and Tsuneda [8]. For this purpose let us observe that the carré du champ referred to the generator \mathcal{L}_N , namely, $\Gamma_N(h) = \mathcal{L}_N h^2 - 2h \mathcal{L}_N h$ can be decomposed as

$$\Gamma_N(h) = \Gamma_N^{\mathcal{G}}(h) + 2\alpha N^2 \Gamma_N^{\mathcal{K}}(h), \quad (4.39)$$

where $\Gamma_N^{\mathcal{G}}(h)$ and $\Gamma_N^{\mathcal{K}}(h)$ are the carré du champ related to the generator \mathcal{G}_N and \mathcal{K}_N respectively (cf. (4.1) and (4.2)). In particular,

$$\Gamma_N^{\mathcal{K}}(h) = \frac{1}{2} \sum_{\substack{i, j \in \mathbb{T}_N^d: \\ |i-j|=1}} (h(\eta^{i,j}) - h(\eta))^2. \quad (4.40)$$

In the next result we provide the control that we need for V^+ .

Proposition 4.5. *Under Assumption 1, we have that for any $u : \mathbb{T}_N^d \rightarrow [0, 1]$ such that*

1. $\varepsilon := 1 - \|u\|_\infty > 0$ with $\|u\|_\infty := \min \{ \|u\|_\infty, \|1 - u\|_\infty \}$,

2. $\|\nabla u(i)\|_\infty \leq \frac{C_0}{N}$ with C_0 independent of N ,

and for any density f with respect to v_u we have that

$$\int V^+(u, \eta) f(\eta) v_u(d\eta) \leq \delta N^2 \int \Gamma_N^{\mathcal{K}}(\sqrt{f})(\eta) v_u(d\eta) + C_1 \ell_{\mathcal{V}}^d \mathcal{H}(f v_u | v_u) + C_2 N^{d-a}, \quad (4.41)$$

for any $\delta > 0$ and $C_1 = 3^d d(d+1) \frac{6}{\varepsilon^2} (\frac{1}{8} + \frac{C_0}{\varepsilon^2})$, $C_2 = \frac{6C_0}{\varepsilon^4}$ for $N > \frac{2C_0}{\varepsilon^2}$.

Remark 4.2. *Note that if the estimate (4.41) holds, then it holds for V^- . Indeed, using (4.37), the fact that $\|u\|_\infty = \|1 - u\|_\infty = 1 - \varepsilon$ and that, under $v_{1-u}(d\eta)$, $1 - \eta$ has for law v_u we get*

$$\begin{aligned} \int V^-(u, \eta) f(\eta) v_u(d\eta) &= - \int V^+(1 - u, 1 - \eta) f(\eta) v_u(d\eta) = \int V^+(1 - u, \eta) f(1 - \eta) v_{1-u}(d\eta) \\ &\leq \delta N^2 \int \Gamma_N^{\mathcal{K}}(\sqrt{f})(1 - \eta) v_{1-u}(d\eta) + C_1 \ell_{\mathcal{V}}^d \mathcal{H}(f(1 - \cdot) v_{1-u} | v_{1-u}) + C_2 N^{d-a} \\ &= \delta N^2 \int \Gamma_N^{\mathcal{K}}(\sqrt{f})(\eta) v_u(d\eta) + C_1 \ell_{\mathcal{V}}^d \mathcal{H}(f v_u | v_u) + C_2 N^{d-a}. \end{aligned}$$

Proof of Proposition 4.5. The proof will proceed in several steps.

Note that, according to Hoeffding Inequality (Lemma A.4), under the probability measure ν_u , ϑ_i^+ is sub-Gaussian with variance parameter $\frac{1}{4}$ and ω_i^+ is sub-Gaussian with variance parameter $\frac{1}{4u_i^2}$.

As in Jara and Menezes [15], we proceed to use an averaged version of V^+ . Let $\ell > 0$ and consider $\Lambda_\ell = \{0 \dots, \ell - 1\}^d$ the cube of size ℓ starting at 0. Let $p_\ell(i) = \ell^{-d} \mathbb{1}_{\{i \in \Lambda_\ell\}}$ and $\hat{p}_\ell(i) = \ell^{-d} \mathbb{1}_{\{i \in -\Lambda_\ell\}} = p_\ell(-i)$. Then for φ defined on \mathbb{T}_N^d , we set

$$\overleftarrow{\varphi}(i) := p_\ell * \varphi(i) = \sum_{j+k=i} p_\ell(j) \varphi(k) = \sum_j p_\ell(j) \varphi(i-j) = \ell^{-d} \sum_{j \in \Lambda_\ell} \varphi(i-j) \quad (4.42)$$

$$\overrightarrow{\varphi}(i) := \hat{p}_\ell * \varphi(i) = \sum_{j+k=i} p_\ell(-j) \varphi(k) = \sum_j p_\ell(-j) \varphi(i-j) = \ell^{-d} \sum_{j \in \Lambda_\ell} \varphi(i+j). \quad (4.43)$$

Let us denote $q_\ell = p_\ell * p_\ell$. We have $q_\ell(i) = \ell^{-2d} |\Lambda_\ell \cap (i - \Lambda_\ell)|$. Thus, $0 \leq q_\ell(i) \leq \ell^{-d}$ and $q_\ell(i) = 0$ if and only if $i \notin \Lambda_{2\ell-2}$. All sum being finite, we get for φ and ψ defined on \mathbb{T}_N^d :

$$\begin{aligned} \sum_{i,j} \varphi(i) \psi(i+j) q_\ell(j) &= \sum_{i,j,k} \varphi(i) \psi(i+j) p_\ell(k) p_\ell(j-k) = \sum_{i,k} \varphi(i) p_\ell(k) \sum_j \psi(i+j) p_\ell(j-k) \\ &= \sum_{i,k} \varphi(i) p_\ell(k) (\hat{p}_\ell * \psi)(k+i) = \sum_j \sum_i \varphi(i) p_\ell(j-i) (\hat{p}_\ell * \psi)(j) \\ &= \sum_j (p_\ell * \varphi)(j) (\hat{p}_\ell * \psi)(j) = \sum_j \overleftarrow{\varphi}(j) \overrightarrow{\psi}(j). \end{aligned}$$

Thus let $V^{+,\ell} := \sum_{i \in \mathbb{T}_N^d} \vartheta_i^+ \omega_i^{+,\ell}$ with $\omega_i^{+,\ell} = \sum_j \omega_{i+j}^+ q_\ell(j)$. We have that

$$V^{+,\ell} = \sum_{i,j \in \mathbb{T}_N^d} \vartheta_i^+ \omega_{i+j}^+ q_\ell(j) = \sum_{i \in \mathbb{T}_N^d} \overleftarrow{\vartheta^+}_i \overrightarrow{\omega^+}_i. \quad (4.44)$$

Then Proposition 4.5 is proved with the following two estimates.

Lemma 4.6. *Suppose that $\|u\|_\infty < 1$ and let $\varepsilon = 1 - \|u\|_\infty$. Then for any $\ell > \ell_\nu$*

$$\int V^{+,\ell} f \nu_u(d\eta) \leq C \varepsilon^{-1} \ell_\nu^{d/2} \left(\mathcal{H}(f \nu_u | \nu_u) + \frac{N^d}{\ell^d} \right). \quad (4.45)$$

The constant C depends only on the dimension, namely it can be taken as $C = 3^d 2^{d/2+2}$.

Lemma 4.7. *Suppose $\varepsilon = 1 - \|u\|_\infty > 0$ and $\|\nabla u(i)\|_\infty \leq \frac{C_0}{N}$ for some $C_0 > 0$ independent of N , where $\nabla u(i) = (\nabla_k u(i))_{k=1}^d$. Then, for any $\ell > \ell_\nu$, with $\ell = N^\kappa$ for some $\kappa > 0$, and $\delta > 0$*

$$\begin{aligned} \int (V^+ - V^{+,\ell}) f \nu_u(d\eta) &\leq \delta N^2 \int \Gamma_N^\kappa(\sqrt{f}) \nu_u(d\eta) \\ &\quad + C_1 \ell_\nu^d \frac{g_d(\ell) \ell^d}{N} \left(\left(2 + \frac{\ell_\nu}{\ell}\right)^d \mathcal{H}(f \nu_u | \nu_u) + \frac{N^d}{\ell^d} \right) + C_2 N^{d-1} \end{aligned} \quad (4.46)$$

where $C_2 = \frac{3C_0}{\varepsilon^4} \left(1 + \frac{2C_0}{N\varepsilon^2}\right)$ and $C_1 = 3^d d(d+1) \left(\frac{2}{\delta\varepsilon^2} \left(1 + \frac{C_0}{N\varepsilon^2}\right) + C_2\right)$ depends only on the dimension and $g_d(\ell)$ is defined in (4.50).

Indeed, we choose ℓ such that $\frac{\ell^d g_d(\ell)}{N} \leq C_0$: more precisely, for any $\delta_0 \in (0, 1)$ we take

$$\ell = N^{\frac{1}{2}(1-\delta_0)} \quad \text{for } d = 1, \quad \text{and} \quad \ell = N^{\frac{1}{d}(1-\delta_0)} \quad \text{for } d \geq 2. \quad (4.47)$$

Since ℓ_ν^d has a log-growth (cf. Assumption 1), this choice of ℓ together with Lemmas 4.6 and 4.7 concludes the proof of Proposition 4.5. \square

We now prove Lemma 4.6 and Lemma 4.7.

Proof of Lemma 4.6. We start by recalling that $V^{+, \ell} = \sum_{i \in \mathbb{T}_N^d} \overleftarrow{\vartheta^+}_i \overrightarrow{\omega^+}_i$. Note that, under v_u , using Lemma A.3,

$\overrightarrow{\omega^+}_i$ is a sub-Gaussian variable with variance parameter $\sum_{i \in \Lambda_\ell} \frac{1}{4u_i^2 \ell^{2d}} \leq \frac{1}{4\varepsilon^2 \ell^d}$.

Since $c_i^+(\eta)$ is a function of $(\eta_{i+j})_{j \in \mathcal{V}_N}$, then for i and j such that $|i-j|_\infty > \ell_{\mathcal{V}}$, ϑ_i^+ and ϑ_j^+ are independent,

and sub-Gaussian with variance parameter $\frac{1}{4}$. Using Lemma A.3, $\overleftarrow{\vartheta^+}_i$ is a sub-Gaussian variable with variance parameter $\frac{\ell_{\mathcal{V}}^d}{4\ell^d} \left(1 + \frac{\ell_{\mathcal{V}}}{\ell}\right)^d \leq 2^{d-2} \frac{\ell_{\mathcal{V}}^d}{\ell^d}$.

We note that all the sites involved in the averages $\overleftarrow{\vartheta^+}_i \overrightarrow{\omega^+}_i$ are in $i + Q_{\ell_{\mathcal{V}}/2 + \ell}$, where $Q_m = \{-m, \dots, m\}^d$ is the d -dimensional cube centered at 0. In particular, for i and j such that $|i-j|_\infty > \ell_{\mathcal{V}} + 2\ell$, the corresponding averages are independent (under v_u). Then, we can take a partition of \mathbb{T}_N^d into independent sites by letting $i = j + (\ell_{\mathcal{V}} + 2\ell)k$ where $j \in \Lambda_{\ell_{\mathcal{V}} + 2\ell}$ and $k \in \Lambda_{\lfloor N/(\ell_{\mathcal{V}} + 2\ell) \rfloor}$. The entropy inequality (cf. (B.3) in [15]) gives

$$\begin{aligned} \int V^{+, \ell} f v_u(d\eta) &= \sum_{j \in \Lambda_{\ell_{\mathcal{V}} + 2\ell}} \int \sum_{k \in \Lambda_{\lfloor N/(\ell_{\mathcal{V}} + 2\ell) \rfloor}} \overleftarrow{\vartheta^+}_{j + (\ell_{\mathcal{V}} + 2\ell)k} \overrightarrow{\omega^+}_{j + (\ell_{\mathcal{V}} + 2\ell)k} f v_u(d\eta) \\ &\leq \sum_{j \in \Lambda_{\ell_{\mathcal{V}} + 2\ell}} \frac{1}{\gamma} \left(\mathcal{H}(f v_u | v_u) + \sum_{k \in \Lambda_{\lfloor N/(\ell_{\mathcal{V}} + 2\ell) \rfloor}} \log \int \exp \left\{ \gamma \overleftarrow{\vartheta^+}_{j + (\ell_{\mathcal{V}} + 2\ell)k} \overrightarrow{\omega^+}_{j + (\ell_{\mathcal{V}} + 2\ell)k} \right\} v_u(d\eta) \right) \\ &\leq \frac{(\ell_{\mathcal{V}} + 2\ell)^d}{\gamma} \mathcal{H}(f v_u | v_u) + \frac{1}{\gamma} \sum_{i \in \mathbb{T}_N^d} \log \int \exp \left\{ \gamma \overleftarrow{\vartheta^+}_i \overrightarrow{\omega^+}_i \right\} v_u(d\eta) \end{aligned}$$

According to Lemma A.2, for $\gamma^{-1} = 2^{d/2+2} \ell^{-d} \varepsilon^{-1} \ell_{\mathcal{V}}^{d/2}$, we have $\log \int \exp \left\{ \gamma \overleftarrow{\vartheta^+}_i \overrightarrow{\omega^+}_i \right\} v_u(d\eta) \leq \log 3$. Thus we obtain

$$\int V^{+, \ell} f v_u(d\eta) \leq 2^{d/2+2} \varepsilon^{-1} \ell_{\mathcal{V}}^{d/2} \left(\left(2 + \frac{\ell_{\mathcal{V}}}{\ell}\right)^d \mathcal{H}(f v_u | v_u) + \frac{N^d}{\ell^d} \log 3 \right).$$

□

Proof of Lemma 4.7. We first prove the Lemma by also assuming that $\mathcal{V}_N \subset \mathbb{Z}^d \setminus \mathbb{N}^d$. In Remark 4.3 at the end of the proof we show how to remove this assumption in dimension $d > 1$.

We then proceed as in Jara and Menezes [15]. We use the fact that

$$\begin{aligned} V^+ - V^{+, \ell} &= \sum_{i \in \mathbb{T}_N^d} \vartheta_i^+ (\omega_i^+ - \omega_i^{+, \ell}) \\ &= \sum_{i, j \in \mathbb{T}_N^d} \vartheta_i^+ \omega_{i+j}^+ (\mathbb{1}_{\{0\}}(j) - q_\ell(j)). \end{aligned} \tag{4.48}$$

We now use Lemma 3.2 in [15] stating that there exists a function $\Phi_\ell : \mathbb{Z}^d \times \mathbb{Z}^d \rightarrow \mathbb{R}$ which is a flow connecting the distribution $\mathbb{1}_{\{0\}}$ to q_ℓ , i.e.:

- $\Phi_\ell(i, j) = -\Phi_\ell(j, i)$
- $\sum_{j: |i-j|=1} \Phi_\ell(i, j) = \mathbb{1}_{\{0\}}(i) - q_\ell(i)$
- $\Phi_\ell(i, j) = 0$ for $i, j \notin \Lambda_{2\ell-1}$
- there is a constant $C = C(d)$ independent of ℓ such that

$$\sum_{|i-j|=1} |\Phi_\ell(i, j)|^2 < C g_d(\ell) \quad \text{and} \quad \sum_{|i-j|=1} |\Phi_\ell(i, j)| < C \ell \tag{4.49}$$

where

$$g_d(\ell) = \begin{cases} \ell & \text{for } d = 1 \\ \log(\ell) & \text{for } d = 2 \\ 1 & \text{for } d \geq 3 \end{cases} \tag{4.50}$$

Using the flow Φ_ℓ we therefore have

$$\begin{aligned}
V^+ - V^{+, \ell} &= \sum_{i, j \in \mathbb{T}_N^d} \vartheta_i^+ \omega_{i+j}^+ \sum_{k: |j-k|=1} \Phi_\ell(j, k) \\
&= \sum_{i, j \in \mathbb{T}_N^d} \vartheta_i^+ \omega_{i+j}^+ \sum_{k=1}^d \left\{ \Phi_\ell(j, j + e_k) - \Phi_\ell(j - e_k, j) \right\} \\
&= \sum_{k=1}^d \sum_{i, j \in \mathbb{T}_N^d} \vartheta_i^+ \Phi_\ell(j, j + e_k) (\omega_{i+j}^+ - \omega_{i+j+e_k}^+) \\
&= \sum_{k=1}^d \sum_{i, j \in \mathbb{T}_N^d} \vartheta_{i-j}^+ \Phi_\ell(j, j + e_k) (\omega_i^+ - \omega_{i+e_k}^+) \\
&= \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} h_i^k (\omega_i^+ - \omega_{i+e_k}^+) \tag{4.51}
\end{aligned}$$

where

$$h_i^k = h_i^{\ell, k} := \sum_{j \in \mathbb{T}_N^d} \vartheta_{i-j}^+ \Phi_\ell(j, j + e_k) = \sum_{j \in \Lambda_{2\ell-1}} \vartheta_{i-j}^+ \Phi_\ell(j, j + e_k). \tag{4.52}$$

To complete the result we need to estimate (4.51). We apply Lemma 3.5 of [8]. Note that the hypothesis of Lemma 3.5 are satisfied in our case: $u_- = \varepsilon$ and $u_+ = 1 - \varepsilon$ and $h_i^k(\eta^{i, i+e_k}) = h_i^k(\eta)$ for any configuration η inasmuch h_i^k is only a function of the sites η_{i-j+s} , for $s \in \mathcal{V}_N$, $j \in \Lambda_{2\ell-1}$, so that it does not depend on η_i and η_{i+e_k} since $\mathcal{V}_N \subset \mathbb{Z}^d \setminus \mathbb{N}^d$. Moreover, we observe that Lemmas 3.4 and 3.5 apply to our case replacing $\chi(u_i)$ by u_i . Therefore, for any $\alpha' > 0$ we have that

$$\int h_i^k (\omega_i^+ - \omega_{i+e_k}^+) f v_u(d\eta) \leq \frac{\alpha'}{2} \int (\sqrt{f(\eta^{i, i+e_k})} - \sqrt{f(\eta)})^2 v_u(d\eta) + \frac{C_{1, \varepsilon}}{\alpha'} \int (h_i^k)^2 f v_u(d\eta) + R_i^k. \tag{4.53}$$

with $C_{1, \varepsilon} = \frac{2}{\varepsilon^2} (1 + \frac{C_0}{N\varepsilon^2})$ and where the rest term R_i^k is controlled as

$$R_i^k \leq C_{2, \varepsilon} |\nabla_k u_i| \int |h_i^k(\eta)| f v_u(d\eta), \tag{4.54}$$

with, for $\varepsilon < \frac{1}{2}$, $C_{2, \varepsilon} = \frac{3}{\varepsilon^4} (1 + \frac{2C_0}{N\varepsilon^2})$. We now take $\alpha' = \delta N^2$, with $\delta > 0$. We get (cf. (4.40))

$$\begin{aligned}
\int (V^+ - V^{+, \ell}) f v_u(d\eta) &= \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \int h_i^k (\omega_i^+ - \omega_{i+e_k}^+) f v_u(d\eta) \\
&\leq \delta N^2 \int \Gamma_N^K(\sqrt{f}) v_u(d\eta) + \frac{C_{1, \varepsilon}}{\delta N^2} \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta) + \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} R_i^k. \tag{4.55}
\end{aligned}$$

The first term is the same of (4.46), then to conclude the proof we have to upper bound the second and the third term of (4.55).

Let us start from the third one. Using that $|\nabla_k u_i| \leq \frac{C_0}{N}$ and $|h_i^k| \leq 1 + (h_i^k)^2$ in (4.54) we get

$$R_i^k \leq \frac{C_0 C_{2, \varepsilon}}{N} \int (1 + (h_i^k)^2) f v_u(d\eta).$$

So that, the last term of (4.55) is bounded by

$$C_0 C_{2, \varepsilon} \left(N^{d-1} + \frac{1}{N} \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta) \right).$$

Let us note that this last term dominates the second term of (4.55). To conclude the proof we need to upper-bound

$$\sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta).$$

As we noted above, h_i^k depends only on the sites η_{i-j+s} , for $s \in \mathcal{V}_N$ and $j \in \Lambda_{2\ell-1}$, so that the random variables h_i^k and $h_{i'}^k$ are independent if $|i - i'|_\infty > (2\ell + \ell_\mathcal{V})$. We then decompose \mathbb{T}_N^d as a disjoint union of cubes of size $2\ell + \ell_\mathcal{V}$, that is, we write $i = j + (\ell_\mathcal{V} + 2\ell)z$ where $j \in \Lambda_{\ell_\mathcal{V}+2\ell}$ and $z \in \Lambda_{\lfloor N/(\ell_\mathcal{V}+2\ell) \rfloor}$,

$$\sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta) = \sum_{j \in \Lambda_{2\ell+\ell_\mathcal{V}}} \sum_{z \in \Lambda_{\lfloor N/(\ell_\mathcal{V}+2\ell) \rfloor}} \int (h_{j+z}^k)^2 f v_u(d\eta).$$

We then apply the entropy inequality and we get

$$\begin{aligned} \sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta) &\leq \frac{1}{\gamma} \sum_{j \in \Lambda_{2\ell+\ell_\mathcal{V}}} \left(\mathcal{H}(f v_u | v_u) + \log \int \exp \left\{ \gamma \sum_{z \in \Lambda_{\lfloor N/(\ell_\mathcal{V}+2\ell) \rfloor}} (h_{j+z}^k)^2 \right\} v_u(d\eta) \right) \\ &= \frac{(2\ell + \ell_\mathcal{V})^d}{\gamma} \mathcal{H}(f v_u | v_u) + \frac{1}{\gamma} \sum_{i \in \mathbb{T}_N^d} \log \int \exp \left\{ \gamma (h_i^k)^2 \right\} v_u(d\eta) \end{aligned} \quad (4.56)$$

To conclude we use a concentration inequality. We have that h_i^k is sub-Gaussian random variable, let σ^2 be its variance parameter. By Proposition A.1 we have that for any $\gamma \leq \frac{1}{4\sigma^2}$,

$$\int \exp \left\{ \gamma (h_i^k)^2 \right\} v_u(d\eta) \leq \log 3.$$

Moreover, to get an upper bound on the variance parameter, we use the same decomposition of the sum (4.52) into subsets which are independent (since ϑ_j^+ is a function of η_{i+j} , for $i \in \mathcal{V}_N$ and is sub-Gaussian with variance parameter $\frac{1}{4}$). This is done in Lemma F.12 in [15]. We then have that $\sigma^2 \leq C_d \ell_\mathcal{V}^d g_d(\ell)$, where C_d is a constant which depends only on the dimension. By getting γ as large as possible, namely $\gamma^{-1} = (d+1)\ell_\mathcal{V}^d g_d(\ell)$ we obtain that

$$\sum_{i \in \mathbb{T}_N^d} \int (h_i^k)^2 f v_u(d\eta) \leq (d+1)\ell_\mathcal{V}^d g_d(\ell) \ell^d \left(\left(2 + \frac{\ell_\mathcal{V}}{\ell}\right)^d \mathcal{H}(f v_u | v_u) + \frac{N^d}{\ell^d} \log 3 \right). \quad (4.57)$$

To conclude the proof we have to remove the assumption $\mathcal{V}_N \subset \mathbb{Z}^d \setminus \mathbb{N}^d$. We show that in Remark 4.3. \square

Remark 4.3. By (4.51) we recall that $V^+ - V^{+, \ell} = \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} h_i^k (\omega_i^+ - \omega_{i+e_k}^+)$ where h_i^k is defined in (4.52). We also observe that, by assumption, $\ell > \ell_\mathcal{V}$. Since h_i^k is a function of the sites η_{i-j+s} , for $s \in \mathcal{V}_N$, $j \in \Lambda_{2\ell-1}$, whenever $|j| > \ell_\mathcal{V}$, then ϑ_{i-j}^+ does not depend on η_i and η_{i+e_k} . Therefore we split h_i^k into the sum of two functions h_i' and h_i'' . The first one is independent of η_i and η_{i+e_k} , while the second one depends on these sites,

$$h_i' := \sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| > 2\ell_\mathcal{V}}} \vartheta_{i-j}^+ \Phi_\ell(j, j + e_k), \quad \text{and} \quad h_i'' := \sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| \leq 2\ell_\mathcal{V}}} \vartheta_{i-j}^+ \Phi_\ell(j, j + e_k). \quad (4.58)$$

In such a way

$$V^+ - V^{+, \ell} = (V^+ - V^{+, \ell})' + (V^+ - V^{+, \ell})'', \quad (4.59)$$

where

$$(V^+ - V^{+, \ell})' := \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} h_i' (\omega_i^+ - \omega_{i+e_k}^+) \quad \text{and} \quad (V^+ - V^{+, \ell})'' := \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} h_i'' (\omega_i^+ - \omega_{i+e_k}^+) \quad (4.60)$$

We can apply the method used above to $(V^+ - V^{+, \ell})'$ obtaining that (4.57) holds. We control $(V^+ - V^{+, \ell})''$ by showing that $\int (V^+ - V^{+, \ell})'' f v_u(d\eta) = O(N^{d-\varepsilon})$, for some $\varepsilon > 0$. For this purpose we apply Cauchy-Swartz inequality to the measure $f v_u(d\eta)$, which gives

$$\int h_i'' (\omega_i^+ - \omega_{i+e_k}^+) f v_u(d\eta) \leq \left(\int (h_i'')^2 f v_u(d\eta) \right)^{\frac{1}{2}} \left(\int (\omega_i^+ - \omega_{i+e_k}^+)^2 f v_u(d\eta) \right)^{\frac{1}{2}} \quad (4.61)$$

Therefore

$$\begin{aligned}
\int (V^+ - V^{+, \ell})'' f v_u(d\eta) &= \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \int h_i''(\omega_i^+ - \omega_{i+e_k}^+) f v_u(d\eta) \\
&\leq \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (h_i'')^2 f v_u(d\eta) \right)^{\frac{1}{2}} \left(\int (\omega_i^+ - \omega_{i+e_k}^+)^2 f v_u(d\eta) \right)^{\frac{1}{2}} \\
&\leq \left(\sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (h_i'')^2 f v_u(d\eta) \right) \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (\omega_i^+ - \omega_{i+e_k}^+)^2 f v_u(d\eta) \right) \right)^{\frac{1}{2}}, \tag{4.62}
\end{aligned}$$

where in the last inequality we used again Cauchy-Swartz inequality.

We observe that $(\omega_i^+ - \omega_{i+e_k}^+) = \left(\frac{\eta_i}{u_i} - \frac{\eta_{i+e_k}}{u_{i+e_k}} \right) = \eta_i \left(\frac{1}{u_i} - \frac{1}{u_{i+e_k}} \right) + \frac{1}{u_{i+e_k}} (\eta_i - \eta_{i+e_k})$, so that

$$\begin{aligned}
\int (\omega_i^+ - \omega_{i+e_k}^+)^2 f v_u(d\eta) &\leq 2 \int \eta_i \left(\frac{1}{u_i} - \frac{1}{u_{i+e_k}} \right)^2 f v_u(d\eta) + 2 \int \frac{1}{u_{i+e_k}^2} (\eta_i - \eta_{i+e_k})^2 f v_u(d\eta) \\
&\leq C_\varepsilon |\nabla u_k|_\infty^2 + C_\varepsilon \int (\eta_i - \eta_{i+e_k})^2 f v_u(d\eta) \leq C'_\varepsilon,
\end{aligned}$$

uniformly on i and N . So that,

$$\sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (\omega_i^+ - \omega_{i+e_k}^+)^2 f v_u(d\eta) \right) \leq C'_\varepsilon N^d. \tag{4.63}$$

Therefore, by (4.62), to conclude the proof it is enough to show that

$$\sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (h_i'')^2 f v_u(d\eta) \right) = O(N^{d-\varepsilon}), \tag{4.64}$$

for some $\varepsilon > 0$ small. For this purpose we have to look more carefully at the function $\Phi_\ell(j, j + e_k)$ which defines the flow. We recall (cf. Appendix G of [15]) that in the construction of the flow connecting $\mathbb{1}_{\{0\}}$ and $p_\ell * p_\ell$, we first define a flow Ψ_ℓ connecting $\mathbb{1}_{\{0\}}$ and p_ℓ supported in Λ_ℓ which satisfies (4.49) and then we define

$$\Phi_\ell(j, j + e_k) := \sum_{i \in \mathbb{T}_N^d} \Psi_\ell(i, i + e_k) p_\ell(j - i),$$

therefore

$$\begin{aligned}
\sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| \leq 2\ell_\nu}} |\Phi_\ell(j, j + e_k)| &\leq \sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| \leq 2\ell_\nu}} \sum_{i \in \mathbb{T}_N^d} |\Psi_\ell(i, i + e_k)| p_\ell(j - i) \\
&\leq \sum_{i \in \mathbb{T}_N^d} |\Psi_\ell(i, i + e_k)| \sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| \leq 2\ell_\nu}} p_\ell(j - i) \leq C \ell \frac{\ell_\nu^d}{\ell^d},
\end{aligned}$$

where we used that $\sum_{\substack{j \in \Lambda_{2\ell-1} \\ |j| \leq 2\ell_\nu}} p_\ell(j - i) \leq \frac{\ell_\nu^d}{\ell^d}$ uniformly on i and (4.49) applied to Ψ_ℓ . We deduce that, since $|\vartheta_{i-j}^+| \leq 2$,

$$\sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left(\int (h_i'')^2 f v_u(d\eta) \right) \leq C N^d \frac{\ell_\nu^{2d}}{\ell^{2(d-1)}}.$$

Since $\ell = N^\kappa$ for some $\kappa > 0$ and ℓ_ν^d grows logarithmically, the result follows (in dimension $d \geq 2$).

4.1.3 Estimate of V

We show that under the hypothesis of Proposition 4.5, $V(u, \eta)$ satisfies (4.41).

For $i \in \mathbb{T}_N^d$ and $k \in \{1, \dots, d\}$ we let

$$\tilde{\omega}_i^k := -\alpha N^2 (u_i - u_{i+e_k})^2 \omega_i.$$

In such a way we get

$$\begin{aligned} V(u, \eta) &= -\alpha N^2 \sum_{\substack{i, j \in \mathbb{T}_N^d \\ |i-j|=1}} (u_i - u_j)^2 \omega_i \omega_j \\ &= -\alpha N^2 \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left\{ (u_i - u_{i+e_k})^2 \omega_i \omega_{i+e_k} + (u_i - u_{i-e_k})^2 \omega_i \omega_{i-e_k} \right\} \\ &= \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \left\{ \tilde{\omega}_i^k \omega_{i+e_k} + \tilde{\omega}_{i-e_k}^k \omega_i \right\} = 2 \sum_{k=1}^d \sum_{i \in \mathbb{T}_N^d} \tilde{\omega}_{i-e_k}^k \omega_i. \end{aligned}$$

For $k \in \{1, \dots, d\}$ we let $\tilde{V}^k := \sum_{i \in \mathbb{T}_N^d} \tilde{\omega}_{i-e_k}^k \omega_i$ and $\overleftarrow{\tilde{V}^{k, \ell}} := \sum_{i \in \mathbb{T}_N^d} \overleftarrow{\tilde{\omega}_{i-e_k}^k} \omega_i$. We estimate $\overleftarrow{\tilde{V}^{k, \ell}}$ and $\tilde{V}^k - \overleftarrow{\tilde{V}^{k, \ell}}$ separately by showing that they satisfy (4.45) and (4.46) respectively.

For $\overleftarrow{\tilde{V}^{k, \ell}}$, by Lemma A.4 we have that $\overleftarrow{\tilde{\omega}_i^k}$ is a sub-Gaussian random variable with variance parameter $\sigma_i^2 \leq \frac{C^2}{\varepsilon^2}$ (we used the assumption on $|\nabla_k u_i|$ and that $\chi(u) > \frac{\varepsilon}{2}$), while we recall that ω_i is a sub-Gaussian random

variable with variance parameter $\frac{1}{4}$. Therefore, $\overleftarrow{\tilde{\omega}_i^k}$ is a sub-Gaussian random variable with variance parameter $\sum_{i \in \Lambda_\ell} \frac{\sigma_i^2}{\ell^{2d}} \leq \frac{C_0^2}{\varepsilon^2 \ell^d}$ and $\overleftarrow{\omega}_i$ is also a sub-Gaussian random variable with variance parameter $\frac{1}{4\ell^d}$.

Note that the averages $\overleftarrow{\tilde{\omega}_{i-e_k}^k}$ are in $i + Q_\ell$, so that for i and j such that $|i - j|_\infty > 2\ell$, the corresponding averages are independent (under v_u). Then, we can partition \mathbb{T}_N^d into independent blocks by letting $i = j + (2\ell)k$ where $j \in \Lambda_{2\ell}$ and $k \in \Lambda_{\lfloor N/(2\ell) \rfloor}$. Therefore, as in the proof of Lemma 4.6, the entropy inequality gives

$$\int \overleftarrow{\tilde{V}^{k, \ell}} f v_u(d\eta) \leq \frac{C \ell^d}{\gamma} \mathcal{H}(f v_u | v_u) + \frac{C}{\gamma} \sum_{i \in \mathbb{T}_N^d} \log \int \exp \left\{ \gamma \overleftarrow{\tilde{\omega}_{i-e_k}^k} \omega_i \right\} v_u(d\eta),$$

where $C = C(\alpha, \varepsilon, d)$. So that, by taking $\gamma^{-1} = \frac{C_0}{2\varepsilon \ell^d}$ and using Lemma A.2 we get

$$\int \overleftarrow{\tilde{V}^{k, \ell}} f v_u(d\eta) \leq C_1 \left(\mathcal{H}(f v_u | v_u) + \frac{N^d}{\ell^d} \right),$$

with $C_1 = C_1(d, \varepsilon, \alpha) > 0$. In particular V satisfies (4.45).

To control $\tilde{V}^k - \overleftarrow{\tilde{V}^{k, \ell}}$ we write the difference using the flow Φ_ℓ to get

$$\tilde{V}^k - \overleftarrow{\tilde{V}^{k, \ell}} = \sum_{i, j \in \mathbb{T}_N^d} \tilde{\omega}_{i-e_k}^k \omega_{i+j} \sum_{k: |j-k|=1} \Phi_\ell(j, k) = \sum_{m=1}^d \sum_{i \in \mathbb{T}_N^d} \tilde{h}_i^m (\omega_i - \omega_{i+e_m}),$$

where

$$\tilde{h}_i^m = \tilde{h}_i^{\ell, k, m} := \sum_{j \in \mathbb{T}_N^d} \tilde{\omega}_{i-e_k-j}^k \Phi_\ell(j, j+e_m) = \sum_{j \in \Lambda_{2\ell-1}} \tilde{\omega}_{i-e_k-j}^k \Phi_\ell(j, j+e_m).$$

We observe that we can apply Lemma 3.5 of [8] since $\tilde{h}_i^m(\eta^{i, i+e_m}) = \tilde{h}_i^m(\eta)$ (\tilde{h}_i^m depends on η_{i-e_k-j} , $j \in \Lambda_{2\ell-1}$), obtaining that

$$\begin{aligned} \int (\tilde{V}^k - \overleftarrow{\tilde{V}^{k, \ell}}) f v_u(d\eta) &= 2 \sum_{m=1}^d \sum_{i \in \mathbb{T}_N^d} \int \tilde{h}_i^m (\omega_i - \omega_{i+e_m}) f v_u(d\eta) \\ &\leq c N^2 \int \Gamma_N^{\mathcal{K}}(\sqrt{f}) v_u(d\eta) + \frac{c}{N} \sum_{m=1}^d \sum_{i \in \mathbb{T}_N^d} \int (\tilde{h}_i^m)^2 f v_u(d\eta) + N^{d-1}. \end{aligned}$$

To conclude the proof, for $m \in \{1, \dots, d\}$, we bound $\sum_{i \in \mathbb{T}_N^d} \int (\tilde{h}_i^m)^2 f v_u(d\eta)$. An application of the entropy inequality gives

$$\sum_{i \in \mathbb{T}_N^d} \int (\tilde{h}_i^m)^2 f v_u(d\eta) \leq \frac{c \ell^d}{\gamma} \mathcal{H}(f v_u | v_u) + \frac{c}{\gamma} \sum_{i \in \mathbb{T}_N^d} \log \int \exp \left\{ \gamma (\tilde{h}_i^m)^2 \right\} v_u(d\eta).$$

By Lemma F.12 in [15] we then have that \tilde{h}_i^m is a sub-Gaussian random variable with $\sigma^2 \leq C_{d,\varepsilon} g_d(\ell)$. We can take $\gamma^{-1} = C_{d,\varepsilon} g_d(\ell)$ and conclude that

$$\sum_{i \in \mathbb{T}_N^d} \int (\tilde{h}_i^m)^2 f v_u(d\eta) \leq C g_d(\ell) \ell^d \left(\mathcal{H}(f v_u | v_u) + \frac{N^d}{\ell^d} \log 3 \right),$$

in particular (4.46) is satisfied. This implies that $V(u, \eta)$ satisfies (4.41).

4.1.4 Conclusion: Gronwall's inequality (4.28)

We observe that since $\Gamma_N^K(\sqrt{f}) \leq \Gamma_N(\sqrt{f})$, cf. (4.39) and that the carré du champ operator is non-negative, (4.28) is a consequence of Propositions 4.3 and 4.5 with $u(t) = u^N(t)$, the solutions of (4.10), $f = \frac{d\mu_t}{dv_{u(t)}}$ if we show that u^N satisfies Assumption 2 for any $t \in [0, \tau]$. This is one of the goal of the Section 5, see Propositions 5.2 and 5.3.

5 Estimates on the solutions (u^N) of (4.10)

Note that (4.10) is a first order ordinary differential equation in $\mathbb{R}^{\mathbb{T}_N^d}$. By standard theory, we have a solution, locally in time, starting from every initial condition.

We say that u is supersolution of (4.10) if $\partial_t u_i \geq 2\alpha N^2 (\Delta u)_i + \beta(1 - 2u_i) + G(i, u)$ and that it is a subsolution if $\partial_t u_i \leq 2\alpha N^2 (\Delta u)_i + \beta(1 - 2u_i) + G(i, u)$.

Note that any solution is both a super- and a subsolution. We have a comparaison lemma between super and subsolutions.

Proposition 5.1. *Let u be a supersolution, and v be a subsolution such that $u(0, i) \geq v(0, i)$ for all $i \in \mathbb{T}_N^d$. Then, for all $t \geq 0$ and all $i \in \mathbb{T}_N^d$, $u(t, i) \geq v(t, i)$.*

Proof. Since u and v have derivative in time they are continuous, consider i and t such that $u(t, i) = v(t, i)$ and $u(t, j) \geq v(t, j)$. Then, we have $G(i, u(t)) \geq G(i, v(t))$ by Proposition B.1, and thus

$$\partial_t(u(i, t) - v(i, t)) = 2\alpha N^2 \sum_{j, |i-j|=1} (u(j, t) - v(j, t)) + G(i, u(t)) - G(i, v(t)) \geq 0. \quad (5.1)$$

This proves that u stays above v at all time. □

From the two previous propositions we conclude :

Proposition 5.2. *Let $\delta = \frac{\beta}{1+4\beta}$ and suppose $0 < \delta < \frac{1}{4e}$. Let also $4\delta \leq T \leq 1 - 4\delta$ and $0 \leq \varepsilon < 2\delta$. For all $K_N > \frac{|\log(\varepsilon/2)|}{\varepsilon^2}$, and all N , consider the solution $(u^N(t, i))_{i \in \mathbb{T}_N^d}$ of Equation (4.10) starting from $u_0 \in [\varepsilon, 1 - \varepsilon]$, then we have that $u^N(t, i) \in [\varepsilon, 1 - \varepsilon]$ for all i and $t \geq 0$.*

Note that the proposition holds for $\varepsilon = 0$.

Proof. Let $p \in [0, 1]$, and set $u(i) = p$, for all i . Using that $g_{K_N}(p) = G(i, p)$, we have that

- if $g_{K_N}(p) + \beta(1 - 2p) \geq 0$, then u is a subsolution;
- if $g_{K_N}(p) + \beta(1 - 2p) \leq 0$, then u is a supersolution.

Then, by using the result of Proposition B.2 on the analysis of $g_K(p)$ close to $p = 0$ and $p = 1$, it is easy to check that $u(i) = \varepsilon$ is a subsolution and $u(i) = 1 - \varepsilon$ is supersolution, for ε satisfying the hypothesis of the proposition.

In particular, for $p = 1$, $g_{K_N}(1) = 0$, we have a supersolution. For $p = 0$, $g_{K_N}(0) = 0$, we have a subsolution. □

In the next result, we show that if u solves (4.10) and $\|\nabla u(0, i)\|_\infty \leq \frac{C}{N}$, then $\|\nabla u(t, i)\|_\infty \leq \frac{C\sqrt{t}}{N}$ for any $t > 0$.

Proposition 5.3. *Let u be a solution of (4.10) with $u(0) = u_0$ such that, there exists $C_0 > 0$ for which $|\nabla u_0|_\infty \leq \frac{C_0}{N}$. Then, there exists $C > 0$ such that $\|\nabla u(t, i)\|_\infty \leq \frac{C_0 + C\sqrt{t}}{N}$ for all $t \geq 0$.*

To prove Proposition 5.3 we follow [8] and the reference therein, in particular [7]. Using the notation of [7] we let $p(t, x, z)$ be the heat kernel of discrete Laplacian

$$\Delta u(t, i) = \sum_{j \in \mathbb{Z}^d} a(t, i, i+j)[u(t, i+j) - u(t, i)], \quad (5.2)$$

with $a(t, i, j) = \mathbb{1}_{\{j \in \Gamma\}}$ and $\Gamma = \{\pm e_i, i = 1, \dots, d\}$, that is,

$$\begin{cases} p(\cdot, i_0, i) = \delta_{i_0}(\cdot), \\ \partial_t p(t, i_0, i) = \Delta p(t, i_0, i). \end{cases}$$

In this case (comments below (1.2) of [7]) we have that $a^*(t, i, j) = a(t, i, j)$ so that $p^*(t, i, j) = p(t, i, j)$. Let $\nabla_k u(t, i) = u(t, i + e_k) - u(t, i)$, then since p is a uniform transitions function, there exist $c, C > 0$ independent of t, k such that (cf. (1.3) of [7])

$$|\nabla_k p(t, 0, i)| \leq C \frac{p(ct, 0, i)}{\sqrt{1 \vee t}}. \quad (5.3)$$

We refer to [7], Section 4 and the reference therein for a proof. We stress that the delicate point is to extend the classical theory of E. De Giorgi, J. Nash and J. Moser to discrete operators. The authors follow mainly [6], but similar results can be also found in [10] (Appendix B) and [23].

Proof of Proposition 5.3. Using Duhamel's formula (i.e., variation of constant) we get that

$$u(t, i) = \sum_{j \in \mathbb{T}_N^d} u(0, j) p_N(t, i, j) + \int_0^t ds \sum_{j \in \mathbb{T}_N^d} (\beta(1 - 2u_j) + G_j(u)) p_N(t-s, i, j), \quad (5.4)$$

where $p_N(t, i, j) = \sum_{z \in \mathbb{N}\mathbb{Z}^d} p(2\alpha N^2 t, i, j+z)$ is the heat kernel of discrete Laplacian on the torus speeds up by a factor $2\alpha N^2$ and $p(t, i, j)$ is the heat kernel introduced above. We observe that (5.3) gives

$$|\nabla_k p_N(t, 0, i)| \leq \frac{C}{N} \frac{p_N(ct, 0, i)}{\sqrt{t}}. \quad (5.5)$$

For the first term, using that $p_N(t, i, j) = p_N(t, i+z, j+z)$ for any $i, j, z \in \mathbb{T}_N^d$ and the assumption on ∇u_0 (cf. Assumptions 2), an integration by parts gives

$$\begin{aligned} \left| \nabla_k \left\{ \sum_{j \in \mathbb{T}_N^d} u(0, j) p_N(t, i, j) \right\} \right| &= \left| \sum_{j \in \mathbb{T}_N^d} u(0, j) (p_N(t, i + e_k, j) - p_N(t, i, j)) \right| \\ &= \left| \sum_{j \in \mathbb{T}_N^d} u(0, j) p_N(t, i, j - e_k) - \sum_{j \in \mathbb{T}_N^d} u(0, j) p_N(t, i, j) \right| \\ &= \left| \sum_{j \in \mathbb{T}_N^d} (u(0, j + e_k) - u(0, j)) p_N(t, i, j) \right| \leq \frac{C}{N}. \end{aligned}$$

By using that $(\beta(1 - 2u_j) + G_j(u))$ is bounded and (5.5) we get that for any $k = 1, \dots, d$,

$$\left| \nabla_k \left\{ \int_0^t ds \sum_{j \in \mathbb{T}_N^d} (\beta(1 - 2u_j) + G_j(u)) p_N(t-s, i, j) \right\} \right| \leq \frac{C}{N} \int_0^t \frac{1}{\sqrt{t-s}} ds = \frac{C}{N} \sqrt{t}.$$

□

6 Existence and uniqueness of reaction-diffusion PDE

In Section 7, we prove that in the limit $N \rightarrow \infty$, the solution of the discretized Equation 4.10, with $u^N(t=0) = u_0^N \in [0, 1]^{\mathbb{T}_N^d}$ satisfying Assumption 2, converges to a solution of the scalar nonlinear reaction diffusion equation

- when $K_N \rightarrow K$ as $N \rightarrow \infty$:

$$\partial_t u(t, x) = 2\alpha \Delta u(t, x) + \beta(1 - 2u(t, x)) + g_K(u(t, x)), \quad (6.1)$$

- when $K_N \rightarrow \infty$ as $N \rightarrow \infty$:

$$\partial_t u(t, x) = 2\alpha \Delta u(t, x) + \beta(1 - 2u(t, x)) + g_\infty(u(t, x)). \quad (6.2)$$

The main difference between the two equations is that in the first case ($K < +\infty$), g_K is a C^1 function on $[0, 1]$ (thus Lipschitz) so the reaction diffusion equation (6.1) is very classical, whereas for the second case ($K = +\infty$), since g_∞ is not even continuous we need to consider (6.2) as a subdifferential inclusion.

The main results of this section are

- Proposition 6.3 which proves existence of a solution, in a suitable sense, for the equation (6.2),
- Proposition 6.4 which proves local uniqueness of the solution, in a suitable sense, for the equation (6.2), starting from a suitable class of initial conditions,
- and Theorem 7.1 which proves that all accumulation points of $(u^N)_N$ is a solution, in a suitable sense, for the equation (6.2).

In the rest of this section we change our notations and define $v = 2u - 1$. We center the solution around the constant steady state $u = \frac{1}{2}$. It simplifies the presentation and proofs of our results. The original form of our equations can be retrieved by letting $u = \frac{1}{2}(v + 1)$. In such a way (6.1) takes the form

$$\partial_t v(t, x) - 2\alpha \Delta v(t, x) + 2\beta v - 2g_K\left(\frac{1}{2}(v + 1)\right) = 0. \quad (6.3)$$

6.1 Solution of (6.1)

Let us denote $s(t, x, y)$ the semigroup of the operator $\frac{1}{2}\Delta$ on \mathbb{T}^d , that is,

$$s(t, x, y) = \frac{1}{(2\pi t)^{d/2}} \sum_{k \in \mathbb{Z}^d} \exp\left(-\frac{\|x - y - k\|^2}{2t}\right). \quad (6.4)$$

Denote also $s_0(t, x, y)$ the semigroup of the operator $\frac{1}{2}\Delta$ on \mathbb{R}^d ,

$$s_0(t, x, y) = \frac{1}{(2\pi t)^{d/2}} \exp\left(-\frac{\|x - y\|^2}{2t}\right). \quad (6.5)$$

Note that $\xi \mapsto s_0(t, 0, \xi)$ is the density of d independent normal random variables with variance t .

Let us consider $(S_t^{\lambda, \gamma})$ the semigroup on $L^1(\mathbb{T}^d)$ defined by, for $f \in L^1(\mathbb{T}^d)$, $\lambda \geq 0$ and $\gamma > 0$

$$S_t^{\lambda, \gamma} f(x) = \int_{\mathbb{T}^d} e^{-\lambda t} s(\gamma t, x, y) f(y) dy = \int_{\mathbb{R}^d} e^{-\lambda t} s_0(\gamma t, x, y) \tilde{f}(y) dy, \quad (6.6)$$

where for a measurable function f on \mathbb{T}^d , we denoted \tilde{f} its extension on \mathbb{R}^d defined by $\tilde{f}(x) = f(x - [x])$.

Another way to define $S_t^{\lambda, \gamma}$ is to use the Brownian motion: denote by X a Brownian motion on \mathbb{R}^d starting from x on some probability space $(\Omega, \mathcal{F}, \mathbb{P}_x)$, indeed we have $S_t^{\lambda, \gamma} f(x) = e^{-\lambda t} \mathbb{E}_x(\tilde{f}(X_{\gamma t}))$, and for all $\lambda \geq 0$, and $\gamma > 0$, $S^{\lambda, \gamma}$ is a C_0 -contraction semigroup on $L^p(\mathbb{T}^d)$ for $p \in [1, +\infty]$.

As we will look at (6.1) in its mild form, the following result is crucial to study the regularity of the solution.

Proposition 6.1. For $v_0 \in L^\infty(\mathbb{T}^d)$ and $g \in L^\infty([0, T] \times \mathbb{T}^d)$, define

$$v(t, x) := S_t^{\lambda, \gamma} v_0(x) + \int_0^t S_{t-s}^{\lambda, \gamma}(g(s, \cdot))(x) ds. \quad (6.7)$$

then $v \in C([0, T], \mathbb{T}^d)$. We have the following estimates, for all $(t, x) \in \mathbb{R}^+ \times \mathbb{T}^d$,

$$|v(t, x)| \leq e^{-\lambda t} \|v_0\|_\infty + \frac{1}{\lambda} (1 - e^{-\lambda t}) \|g\|_\infty \leq \|v_0\|_\infty + \frac{1}{\lambda} \|g\|_\infty \quad (6.8)$$

and for all $T > 0$, there exists a constant C depending only on T, γ, λ and d , such that for all $(t, x), (s, y) \in [1/T, T] \times \mathbb{T}^d$ with $s < t$

$$|v(t, x) - v(s, y)| \leq C((t-s) |\log(t-s)| + \|x-y\|)(\|g\|_\infty + \|v_0\|_\infty). \quad (6.9)$$

Remark 6.1. v is called a mild solution of the equation $\partial_t v - \frac{\gamma}{2} \Delta v + \lambda v = g$ with initial value v_0 . The fact that a mild solution is a classical solution if g is sufficiently regular is a result from Pazy ([20], Corollary 4.2.5).

Proof of Proposition 6.1. The fact that $v \in C([0, T], L^\infty(\mathbb{T}^d))$ is a consequence of the fact that $S^{\lambda, \gamma}$ is a C_0 contraction semigroups on L^∞ . For the first estimate (6.8), we have that :

$$\begin{aligned} |v(t, x)| &\leq |S_t^{\lambda, \gamma} v_0(x)| + \left| \int_0^t S_{t-u}^{\lambda, \gamma}(g(u, \cdot))(x) du \right| \\ &\leq e^{-\lambda t} \|v_0\|_\infty \int_{\mathbb{R}^d} s_0(\gamma t, x, z) dz + \|g\|_\infty \int_0^s e^{-\lambda(t-u)} \int_{\mathbb{R}^d} s_0(\gamma(t-u), x, z) dz du \\ &= e^{-\lambda t} \|v_0\|_\infty + \frac{1}{\lambda} (1 - e^{-\lambda t}) \|g\|_\infty. \end{aligned}$$

For the estimate (6.9), we start by letting $s < t$, we have

$$\begin{aligned} |v(t, x) - v(s, y)| &\leq |S_t^{\lambda, \gamma} v_0(x) - S_s^{\lambda, \gamma} v_0(y)| + \left| \int_0^t S_{t-u}^{\lambda, \gamma}(g(u, \cdot))(x) du - \int_0^s S_{s-u}^{\lambda, \gamma}(g(u, \cdot))(y) du \right| \\ &\leq I_1 \|v_0\|_\infty + (I_2 + I_3) \|g\|_\infty \end{aligned}$$

where

$$\begin{aligned} I_1 &= \int_{\mathbb{R}^d} |e^{-\lambda t} s_0(\gamma t, x, z) - e^{-\lambda s} s_0(\gamma s, y, z)| dz, \\ I_2 &= \int_0^s \int_{\mathbb{R}^d} |e^{-\lambda(t-u)} s_0(\gamma(t-u), x, z) - e^{-\lambda(s-u)} s_0(\gamma(s-u), y, z)| dz du, \\ I_3 &= \int_s^t \int_{\mathbb{R}^d} e^{-\lambda(t-u)} s_0(\gamma(t-u), x, z) dz du. \end{aligned}$$

We have that $I_3 \leq t - s$. For I_1 and I_2 , we use the fact that, for $i = 1 \dots d$

$$\begin{aligned} \partial_t s_0(t, 0, \xi) &= \frac{1}{2} \Delta_\xi s_0(t, 0, \xi) = \frac{1}{2} \left(\frac{\|\xi\|^2}{t^2} - \frac{d}{t} \right) s_0(t, 0, \xi) \\ \partial_{\xi_i} s_0(t, 0, \xi) &= -\frac{\xi_i}{t} s_0(t, 0, \xi). \end{aligned}$$

First for I_1 , denote $c_1(r) = rt + (1-r)s$ and $c_2(r) = r(z-x) + (1-r)(z-y) = z - (rx + (1-r)y)$ for $r \in [0, 1]$, We have that

$$\begin{aligned} |(\partial_t s_0)(\gamma c_1(r), 0, c_2(r))| &\leq \frac{1}{2} \left(\frac{\|c_2(r)\|^2}{\gamma^2 c_1(r)^2} + \frac{d}{\gamma c_1(r)} \right) s_0(\gamma c_1(r), 0, c_2(r)) \\ |(\partial_{\xi_i} s_0)(\gamma c_1(r), 0, c_2(r))| &\leq \frac{|c_2(r)_i|}{\gamma c_1(r)} s_0(\gamma c_1(r), 0, c_2(r)) \end{aligned}$$

Therefore

$$\begin{aligned}
I_1 &= \int_{\mathbb{R}^d} |e^{-\lambda t} s_0(\gamma t, \mathbf{0}, z - x) - e^{-\lambda s} s_0(\gamma s, \mathbf{0}, z - y)| dz \\
&\leq \int_{\mathbb{R}^d} \int_0^1 |c_1'(r) \partial_t (e^{-\lambda t} s_0(\gamma t, \mathbf{0}, \xi))|_{t=c_1(r), \xi=c_2(r)}| dr dz \\
&\quad + \int_{\mathbb{R}^d} \int_0^1 \sum_{i=1}^d |c_2'(r)_i \partial_{\xi_i} (e^{-\lambda t} s_0(\gamma t, \mathbf{0}, \xi))|_{t=c_1(r), \xi=c_2(r)}| dr dz \\
&\leq I_{1,1} + I_{1,2},
\end{aligned}$$

where we have, by applying Fubini and a change of variable

$$\begin{aligned}
I_{1,1} &= (t-s) \int_{\mathbb{R}^d} \int_0^1 \left[\lambda e^{-\lambda c_1(r)} + \frac{\gamma}{2} \left(\frac{\|c_2(r)\|^2}{c_1(r)^2} + \frac{d}{c_1(r)} \right) \right] s_0(\gamma c_1(r), \mathbf{0}, c_2(r)) dr dz \\
&\leq (t-s) \int_0^1 \left[\lambda e^{-\lambda c_1(r)} + \frac{\gamma}{2} \left(\frac{d \gamma c_1(r)}{\gamma^2 c_1(r)^2} + \frac{d}{\gamma c_1(r)} \right) \right] dr \\
&\leq (t-s) \int_0^1 \left[\lambda e^{-\lambda c_1(r)} + \frac{d}{c_1(r)} \right] dr = \int_s^t \left(\lambda e^{-\lambda u} + \frac{d}{u} \right) du = e^{-\lambda s} (1 - e^{-\lambda(t-s)}) + d \log \left(\frac{t}{s} \right) \\
&\leq \left(\lambda + \frac{d}{s} \right) (t-s).
\end{aligned}$$

For the term $I_{1,2}$, we get, using Cauchy-Schwarz, Fubini and a change of variable

$$\begin{aligned}
I_{1,2} &= \int_{\mathbb{R}^d} \int_0^1 \sum_{i=1}^d |y_i - x_i| e^{-\lambda c_1(r)} \frac{|c_2(r)_i|}{\gamma c_1(r)} s_0(\gamma c_1(r), \mathbf{0}, c_2(r)) dr dz \\
&\leq \|x - y\| \int_0^1 \frac{e^{-\lambda c_1(r)}}{\gamma c_1(r)} \int_{\mathbb{R}^d} \|c_2(r)\| s_0(\gamma c_1(r), \mathbf{0}, c_2(r)) dz dr = \|x - y\| \int_0^1 \frac{e^{-\lambda c_1(r)}}{\sqrt{\gamma c_1(r)}} dr C_1 \\
&\leq C_1 \|x - y\| \int_s^t \frac{e^{-\lambda u}}{\sqrt{\gamma u}} \frac{du}{t-s} \leq \frac{C_1}{\sqrt{\gamma s}} \|x - y\|
\end{aligned}$$

where C_1 is the expectation of the quadratic norm of $X = (X_1, X_2, \dots, X_d)$ of d independent standard normal variables : $C_1 = \mathbb{E}(\|X\|) \leq \mathbb{E}(\|X\|^2)^{1/2} = \sqrt{d}$ (we also have $C_1 = \sqrt{2} \frac{\Gamma((d+1)/2)}{\Gamma(d/2)} \sim \sqrt{d}$). We get that, for $s \geq \frac{1}{T}$,

$$I_1 \leq \left(\lambda + \frac{d}{s} \right) (t-s) + \sqrt{\frac{d}{\gamma s}} \|x - y\| \leq (\lambda + dT) (t-s) + \sqrt{\frac{dT}{\gamma}} \|x - y\|.$$

For I_2 , we make the same computations with $c_1(r) = r(t-u) + (1-r)(s-u) = rt + (1-r)s - u$ where $u \in [0, s]$, we have, since $c_1'(r) = t-s$,

$$I_2 \leq \int_0^s \int_{s-u}^{t-u} \left(\lambda e^{-\lambda v} + \frac{d}{v} \right) dv du + C_1 \|x - y\| \int_0^s \int_{s-u}^{t-u} \frac{e^{-\lambda v}}{\sqrt{\gamma v}} \frac{dv}{t-s} du$$

For the first integral, we have

$$\begin{aligned}
\int_0^s \int_{s-u}^{t-u} \left(\lambda e^{-\lambda v} + \frac{d}{v} \right) dv du &= (1 - e^{-\lambda(t-s)}) \int_0^s e^{-\lambda(s-u)} du + d \int_0^s \log \left(\frac{t-u}{s-u} \right) du \\
&= \frac{1}{\lambda} (1 - e^{-\lambda(t-s)}) (1 - e^{-\lambda s}) + d [t \log(t) - s \log(s) - (t-s) \log(t-s)] \\
&\leq \lambda s (t-s) + d [s (\log(t) - \log(s)) + (t-s) (\log(t) - \log(t-s))] \\
&\leq \lambda s (t-s) + d (t-s) + d (t-s) (|\log(t)| + |\log(t-s)|) \\
&\leq (\lambda s + d + |\log(t)|) (t-s) + d (t-s) |\log(t-s)|.
\end{aligned}$$

For the second integral, we get

$$\int_0^s \int_{s-u}^{t-u} \frac{e^{-\lambda v}}{\sqrt{\gamma v}} dv du \leq (t-s) \int_0^s \frac{1}{\sqrt{\gamma(s-u)}} du = (t-s) \sqrt{\frac{2s}{\gamma}}.$$

Then we obtain, for $1/T \leq s < t \leq T$

$$\begin{aligned} I_2 &\leq (\lambda s + d + |\log(t)|)(t-s) + d(t-s)|\log(t-s)| + C_1 \|x-y\| \sqrt{\frac{2s}{\gamma}} \\ &\leq (\lambda T + d + |\log(T)|)(t-s) + d(t-s)|\log(t-s)| + \sqrt{\frac{2dT}{\gamma}} \|x-y\|. \end{aligned}$$

At last, we get the following estimate

$$\begin{aligned} |v(t,x) - v(s,y)| &\leq \left[(\lambda + dT)(t-s) + \sqrt{\frac{dT}{\gamma}} \|x-y\| \right] \|v_0\|_\infty \\ &\quad + \left[(\lambda T + |\log(T)| + d + 1)(t-s) + d(t-s)|\log(t-s)| + \sqrt{\frac{2dT}{\gamma}} \|x-y\| \right] \|g\|_\infty. \end{aligned}$$

□

We modify a little our equation (6.1), both in order to obtain a sharper estimate on the uniform norm of the solution and to get a coherent notation with the solution of the limit equation when $K \rightarrow +\infty$.

We define for $q \in [-1, 1]$,

$$r_K(q) := - \int_{\frac{1}{2}}^{\frac{1}{2}(q+1)} 4g_K(s) ds. \quad (6.10)$$

We have that

$$r'_K(q) = -2g_K \left(\frac{1}{2}(q+1) \right) = -(1-q)\mathbb{P}_{\frac{1-q}{2}}[X < \kappa(K, T)] + (1+q)\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)]. \quad (6.11)$$

So we let h_K be

$$h_K(q) = \begin{cases} -1 & \text{for } q < -1 \\ -r'_K(q) + q & \text{for } q \in [-1, 1] \\ 1 & \text{for } q > 1 \end{cases} \quad (6.12)$$

Since $r'_K(1) = 0$ and $r'_K(-1) = 0$, h_K is continuous on \mathbb{R} . We have that, for $q \in [-1, 1]$,

$$h_K(q) = -r'_K(q) + q = (1-q)\mathbb{P}_{\frac{1-q}{2}}[X < \kappa(K, T)] - (1+q)\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + q.$$

We now solve the following equation

$$\partial_t v(t, x) - 2\alpha \Delta v(t, x) + (2\beta + 1)v = h_K(v(t, x)). \quad (6.13)$$

Note that, this equation and (6.3) are exactly the same with the term v added on both sides if $\|v\|_\infty \leq 1$, since for $q \notin [-1, 1]$, $h_K(q) \neq q$. Thus, a solution v of (6.3) with $\|v\|_\infty \leq 1$ will also be a solution of (6.13) and reciprocally.

Proposition 6.2. *For $v_0 \in L^\infty(\mathbb{T}^d)$ with $\|v_0\|_\infty \leq 1$, there exists a unique solution $(v(t, x), t \geq 0, x \in \mathbb{T}^d)$ to the problem*

- v is continuous from \mathbb{R}_+^* to $L^\infty(\mathbb{T}^d)$
- v satisfies, for all $t > 0$ and $x \in \mathbb{T}^d$

$$v(t, x) = S_t^{2\beta+1, 4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} [h_K(v(s, \cdot))](x) ds. \quad (6.14)$$

We say that v is a mild solution to (6.13). We have also that $\|v\|_\infty \leq 1$ and v satisfies

$$v(t, x) = S_t^{2\beta, 4\alpha} v_0(x) - \int_0^t S_{t-s}^{2\beta, 4\alpha} [r'_K(v(s, \cdot))](x) ds \quad (6.15)$$

and thus is a mild solution of (6.3).

Proof. The first part of the proposition comes from a fixed point argument (see also Pazy [20], Theorem 6.1.2) applied to the following functional let $T > 0$ and define the functional $F : C([0, T], L^\infty(\mathbb{T}^d)) \rightarrow C([0, T], L^\infty(\mathbb{T}^d))$ defined by

$$F(v)(t, x) := S_t^{2\beta+1, 4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} [h_K(v(s, \cdot))](x) ds. \quad (6.16)$$

We equip $C([0, T], L^\infty(\mathbb{T}^d))$ with the uniform topology on all compact subset. We can apply the Banach fixed point Theorem to F (see the proof of Pazy [20], Theorem 6.1.2. Moreover, the mapping $v_0 \mapsto v$ is Lipschitz continuous from L^∞ to $C([0, T], L^\infty(\mathbb{T}^d))$.

An application of Proposition 6.1 proves that $\|v\|_\infty \leq 1$.

Since h_K is differentiable, then, if $v_0 \in C^2(\mathbb{T}^d)$, v_0 is in the domain of Δ and thus v is classical solution of (6.13) (Theorem 6.1.5 [20]). Thus v is a classical solution of (6.3) since $\|v\|_\infty \leq 1$ and a mild solution of (6.3). Now consider an approximating sequence $(v_{0,n})$ in $C^2(\mathbb{T}^d)$ of $v_0 \in L^\infty$, and (v_n) the sequence of mild solutions with initial value $v_{0,n}$ and v the mild solution of (6.13) with initial value v . Then, since $v_0 \mapsto v$ is Lipschitz continuous, by the dominated convergence theorem, we get that, uniformly on $[t_0, T] \times \mathbb{T}^d$ for all $t_0 > 0$, the right hand side of

$$v_n(t, x) = S_t^{2\beta, 4\alpha} v_{0,n}(x) - \int_0^t S_{t-s}^{2\beta, 4\alpha} [r'_K(v_n(s, \cdot))](x) ds \quad (6.17)$$

converges to $S_t^{2\beta, 4\alpha} v_0(x) - \int_0^t S_{t-s}^{2\beta, 4\alpha} [r'_K(v(s, \cdot))](x) ds$, whereas the left hand side converges to v . So we obtain that v is a mild solution of (6.3). \square

6.2 Solution of (6.2)

6.2.1 Existence of a solution

We use the same transform as before, and let h_∞ be the pointwise limit of h_K . The equation (6.2) is now formally

$$\partial_t v(t, x) - 2\alpha \Delta v(t, x) + (2\beta + 1)v = h_\infty(v(t, x)). \quad (6.18)$$

For the limiting equation, we prove first that the family $(v_K)_K$ of solutions associated to h_K with common initial value $v_0 \in L^\infty(\mathbb{T}^d)$ in $C(\mathbb{R}_*^+ \times \mathbb{T})$ is compact (with uniform norm on all compact subset). Then, by taking the limit, any accumulation point v_∞ of the sequence satisfy the mild formulation of the limiting equation relaxed as a subdifferential inclusion.

In order to prove this, we set some notations :

$$r_\infty(q) := - \int_{\frac{1}{2}}^{\frac{1}{2}(q+1)} 4g_\infty(s) ds \quad (6.19)$$

h_∞ , the pointwise limit of h_K , is the function on \mathbb{R}

$$h_\infty(q) = 2g_\infty\left(\frac{1+q}{2}\right) + q = -\mathbb{1}_{q \leq -2\rho} + q\mathbb{1}_{-2\rho < q \leq 2\rho} + \mathbb{1}_{q > 2\rho}. \quad (6.20)$$

Then h_∞ is non-decreasing and is the left-derivative of the convex function H_∞

$$\begin{aligned} H_\infty(q) &:= -r_\infty(q) + \frac{q^2}{2} \\ &= [-q + 2\rho + 2\rho^2]\mathbb{1}_{\{q \leq -2\rho\}} + \frac{1}{2}q^2\mathbb{1}_{\{-2\rho < q \leq 2\rho\}} + [q - 2\rho + 2\rho^2]\mathbb{1}_{\{2\rho < q\}} \end{aligned} \quad (6.21)$$

The subdifferential of H_∞ at q is defined as

$$\partial H_\infty(q) = \{p \in \mathbb{R}, H_\infty(q') - H_\infty(q) \geq p(q' - q), \text{ for all } q' \in [-1, 1]\}.$$

In particular we have,

$$\partial H_\infty(q) = \begin{cases} \{-1\} & \text{for } q < -2\rho, \\ [-1, -2\rho] & \text{for } q = -2\rho, \\ \{q\} & \text{for } -2\rho < q < 2\rho, \\ [2\rho, 1] & \text{for } q = 2\rho, \\ \{1\} & \text{for } q > 2\rho. \end{cases} \quad (6.22)$$

We adopt the following definition for a solution of the equation :

$$\partial_t v(t, x) - 2\alpha \Delta v(t, x) + (2\beta + 1)v \in \partial H_\infty(v(t, x)). \quad (6.23)$$

Definition 6.1. We say that v is a mild solution of Equation (6.23) if it satisfies

$$v(t, x) = S_t^{2\beta+1, 4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} [w(s, \cdot)](x) ds. \quad (6.24)$$

where $w \in L^2([0, T] \times \mathbb{T}^d)$, with $w(t, x) \in \partial H_\infty(v(t, x))$ almost everywhere.

Proposition 6.3. For $v_0 \in L^\infty(\mathbb{T}^d)$, any accumulation point (in $C(\mathbb{R}_+^* \times \mathbb{T}^d)$ equipped with uniform norm on each compact set), of the sequence (v_K) of solutions given by Proposition 6.2 is a mild solution of (6.23).

As a consequence of the proposition, there exists $(v(t, x), t \geq 0, x \in \mathbb{T}^d)$ a mild solution of (6.23) such that v is continuous from \mathbb{R}_+^* to $L^\infty(\mathbb{T}^d)$, and $\|v\|_\infty \leq 1$.

The existence of a solution for a given initial condition v_0 is not difficult and can be prove in different ways (see remark 6.3). Here we adopt some kind of regularization procedure, since we have a natural family of differentiable functions (namely the (h_K)) approximating h_∞ and we use the convergence of the sequence (v_K) in the next section to prove the convergence of the stochastic process. In the appendix, we present another construction of solution(s) using the monotonicity of h_∞ which is interesting since it gives an insight on the problem of non-uniqueness.

Proof. For each K , we have a mild solution v_K from Proposition 6.2. From Proposition 6.1, we have that each solution is uniformly bounded, uniformly continuous on $[1/T, T] \times \mathbb{T}^d$, and the modulus of continuity only depends on $T > 1$ (since the others parameters are fixed).

Therefore, by the Arzela-Ascoli Theorem, the sequence $(v_K)_K$ is compact on $C([1/T, T] \times \mathbb{R}^d)$ and we can extract a subsequence converging uniformly in $C([1/T, T] \times \mathbb{R}^d)$, and then by a diagonal argument, a sequence converging to a limit v_∞ in $C([0, \infty[\times \mathbb{R}^d)$, uniformly on each compact. Note that since $\|v_K\| \leq 1$ for all K , we also have $\|v_\infty\| \leq 1$. We show that v_∞ satisfies (6.18).

Let us assume that $\rho > 0$. Denote for any $s > 0$, $\mathcal{A}_\rho(s) := \{y \in \mathbb{R}^d : |v_\infty(s, y) \pm 2\rho| = 0\}$. Note that since h_∞ is uniformly continuous on $[-1, -2\rho[,]-2\rho, 2\rho[$ and $]2\rho, 1]$, we have that for any $y \notin \mathcal{A}_\rho(s)$, $\lim_{K \rightarrow \infty} h_K(v_K(s, y)) = h_\infty(v_\infty(s, y))$. Let $y \in \mathcal{A}_\rho(s)$ and assume $v_\infty(s, y) = 2\rho$ without loss of generality, since $v_K(s, y)$ converges to $v_\infty(s, y)$, for all $\varepsilon > 0$ such that $2\rho - \varepsilon > 0$, there exists K_0 such that for all $K \geq K_0$, $2\rho - \varepsilon < v_K(s, y) < 2\rho + \varepsilon$. Thus, using Lemma B.5, we have that, for all $K \geq K_0$, $2\rho - 2\varepsilon \leq h_K(v_K(s, y)) \leq 1$. Then taking the limits in K and $\varepsilon \rightarrow 0$, we get

$$h_\infty(2\rho) = 2\rho \leq \liminf_{K \rightarrow \infty} h_K(v_K(s, y)) \leq \limsup_{K \rightarrow \infty} h_K(v_K(s, y)) \leq 1 = h_\infty(2\rho^+). \quad (6.25)$$

For $\rho = 0$, we have the same inequality since then $h_\infty(0) = -1$ and $h_\infty(0^+) = 1$.

Therefore, let $w_+(s, y) = \limsup_{K \rightarrow \infty} h_K(v_K(s, y))$ and $w_-(s, y) = \liminf_{K \rightarrow \infty} h_K(v_K(s, y))$, we have that for all $(s, y) \in]0, +\infty[\times \mathbb{T}^d$:

$$h_\infty(v_\infty(s, y)) \leq w_-(s, y) \leq w_+(s, y) \leq h_\infty(v_\infty(s, y)^+). \quad (6.26)$$

Since $h_K(v_K)$ is bounded, by the Banach Alaoglu Theorem, the sequence is weakly compact in $L^2([0, T] \times \mathbb{T}^d)$, and we have a subsequence of $(h_K(v_K))_K$ converging weakly to $w \in L^2_{loc}([0, +\infty[\times \mathbb{T}^d)$. Since the density of the semigroup $S^{2\beta+1, 4\alpha}$ is in $L^2([0, T] \times \mathbb{R}^d)$ for all $T > 0$, we have as $K \rightarrow +\infty$,

$$v_\infty(t, x) = S_t^{2\beta+1, 4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} w(s, \cdot)(x) ds. \quad (6.27)$$

Moreover, w_- and w_+ are bounded and therefore in $L^2([0, T] \times \mathbb{T}^d)$. Let $\varphi \in L^2([0, T] \times \mathbb{T}^d)$ and $\varphi \geq 0$, by the Fatou Lemma we get

$$0 = \int_{]0, T] \times \mathbb{T}^d} \liminf_{K \rightarrow +\infty} (h_K(v_K) - w_-) \varphi \leq \liminf_{K \rightarrow +\infty} \int_{]0, T] \times \mathbb{T}^d} (h_K(v_K) - w_-) \varphi = \int_{]0, T] \times \mathbb{T}^d} (w - w_-) \varphi. \quad (6.28)$$

We also have

$$0 = \int_{]0, T] \times \mathbb{T}^d} \liminf_{K \rightarrow +\infty} (w_+ - h_K(v_K)) \varphi \leq \liminf_{K \rightarrow +\infty} \int_{]0, T] \times \mathbb{T}^d} (w_+ - h_K(v_K)) \varphi = \int_{]0, T] \times \mathbb{T}^d} (w_+ - w) \varphi. \quad (6.29)$$

Thus, almost everywhere on $]0, +\infty[\times \mathbb{T}^d$, we have that

$$h_\infty(v_\infty) \leq w_- \leq w \leq w_+ \leq h_\infty(v_\infty^+). \quad (6.30)$$

Therefore, $w \in \partial H_\infty(v_\infty)$ a.e. \square

6.2.2 Uniqueness of solution of (6.23)

The main problem concerns the uniqueness of a solution.

Remark 6.2. We prove first that, we do not have uniqueness for a constant initial condition $v_0(x) = 2\rho$ when $2\rho < \frac{1}{1+2\beta}$, so we are in the case of segregation or metastable segregation described by Figure 1. We describe three possible solutions starting from this initial condition.

Note that

$$S_t^{2\beta+1,4\alpha} v_0(x) = \int_{\mathbb{R}^d} e^{-(2\beta+1)t} s_0(t, x, y) 2\rho dy = 2\rho e^{-(2\beta+1)t}. \quad (6.31)$$

Suppose that v does not depend on x , $v(t, x) = c(t)$ for all $x \in \mathbb{T}^d$, we have

$$\int_0^t S_{t-s}^{2\beta+1,4\alpha} [h_\infty(v(s, \cdot))](x) ds = \int_0^t h_\infty(c(s)) e^{-(2\beta+1)(t-s)} ds. \quad (6.32)$$

Let us consider the functions

$$v^1(t, x) = c^1(t) = 2\rho e^{-2\beta t} \quad (6.33)$$

$$v^2(t, x) = c^2(t) = 2\rho e^{-(2\beta+1)t} + \frac{1}{1+2\beta} \left(1 - e^{-(2\beta+1)t}\right) = 2\rho + \left(\frac{1}{1+2\beta} - 2\rho\right) \left(1 - e^{-(2\beta+1)t}\right). \quad (6.34)$$

Since, $c^1(t) \in [0, 2\rho]$, for $t > 0$, we have $h_\infty(c^1(t)) = c^1(t)$ and then

$$\int_0^t h_\infty(c^1(s)) e^{-(2\beta+1)(t-s)} ds = \int_0^t 2\rho e^{-2\beta s} e^{-(2\beta+1)(t-s)} ds = 2\rho e^{-(2\beta+1)t} (e^t - 1). \quad (6.35)$$

Therefore

$$S_t^{2\beta+1,4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1,4\alpha} [h_\infty(v^1(s, \cdot))](x) ds = 2\rho e^{-(2\beta+1)t} + 2\rho e^{-(2\beta+1)t} (e^t - 1) = 2\rho e^{-2\beta t} = v^1(t, x). \quad (6.36)$$

Since $2\rho < \frac{1}{1+2\beta}$, $c^2(t) \in]2\rho, 1]$, for $t > 0$, we have $h_\infty(c^2(t)) = 1$ and then by the same computation,

$$\int_0^t h_\infty(c^2(s)) e^{-(2\beta+1)(t-s)} ds = \frac{1}{1+2\beta} (1 - e^{-(2\beta+1)t}). \quad (6.37)$$

Therefore, we also have

$$S_t^{2\beta+1,4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1,4\alpha} [h_\infty(v^2(s, \cdot))](x) ds = v^2(t, x). \quad (6.38)$$

Thus, both v^1 and v^2 are mild solutions to (6.18) and thus to (6.23) with the same initial conditions. Note that at $t = 0$, we have $v^1(0, x) = v^2(0, x) = 2\rho$ and

$$\partial_t v^1(0, x) = -4\beta\rho = -(2\beta+1)2\rho + 2\rho = -(2\beta+1)2\rho + h_\infty(2\rho^-) \quad (6.39)$$

$$\partial_t v^2(0, x) = -2\rho(2\beta+1) + 1 = -(2\beta+1)2\rho + h_\infty(2\rho^+). \quad (6.40)$$

We see that non uniqueness really comes from the fact that at $t = 0$, where $v(t, x) = 2\rho$, we have two choices for the derivative due to the fact that h_∞ is not continuous. Therefore, we cannot expect uniqueness for this problem for all initial condition, we have to impose some condition on the initial condition if we want a unique solution.

Note that if we consider the mild solution to the subdifferential inclusion (6.23), then we have at least a third solution : $v^3(t, x) = 2\rho$. We consider $w(t, x) = 2\rho(2\beta+1)$, we have

$$\begin{aligned} S_t^{2\beta+1,4\alpha} v_0(x) + \int_0^t S_{t-s}^{2\beta+1,4\alpha} [w(s, \cdot)](x) ds &= 2\rho e^{-(2\beta+1)t} + 2\rho(2\beta+1) \int_0^t e^{-(2\beta+1)(t-s)} ds \\ &= 2\rho. \end{aligned} \quad (6.41)$$

Since, $2\rho < \frac{1}{1+2\beta}$, we have that $2\rho < 2\rho(2\beta+1) < 1$ so $w(t, x) \in \partial H_\infty(2\rho)$.

In the literature, we can find different conditions ensuring that the solution of Equation (6.23) is unique. Following [11] and [4], we prove that the regularity of the initial condition at the levels where the non-linearity H_∞ is not differentiable is sufficient.

Definition 6.2. A function $v_0 : \mathbb{T}^d \rightarrow [-1, 1]$ in $C^1(\mathbb{T}^d)$ is regular at level $q \in]-1, 1[$ if for all $x \in \mathbb{T}^d$, such that $v_0(x) = p$, we have $\nabla v_0(x) \neq 0$.

Proposition 6.4. For $v_0 \in C^1(\mathbb{T}^d)$, such that ∇v_0 is Lipschitz on \mathbb{T}^d and regular at levels 2ρ and -2ρ , the solution v to Equation (6.23) is locally unique. Moreover, the Lebesgue measure of the set $\mathcal{A}_\rho(s) := \{y \in \mathbb{R}^d : |v(s, y) \pm 2\rho| = 0\}$ is zero.

We adapt two arguments by [11] and [4].

Lemma 6.5. If v is a mild solution of (6.23) with $v_0 \in C^1(\mathbb{T}^d)$, and such that ∇v_0 is Lipschitz on \mathbb{T}^d , then, for all $T > 0$, there exists a constant $C > 0$ such that, for all $t \in [0, T]$

$$\|v(t) - v_0\|_\infty \leq Ct^{1/2}, \quad \|\nabla v(t) - \nabla v_0\|_\infty \leq Ct^{1/2}. \quad (6.42)$$

Proof. Since v is a mild solution of (6.23) there exists $w \in L_{loc}^\infty([0, \infty[\times \mathbb{T}^d)$ with $\|w\|_\infty \leq 1$ since $w \in \partial H_\infty(v)$. Thus we get

$$v(t) - v_0 = S_t^{2\beta+1, 4\alpha} v_0 - v_0 + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} w(s) ds. \quad (6.43)$$

Then, we have for the last integral

$$\left\| \int_0^t S_{t-s}^{2\beta+1, 4\alpha} w(s) ds \right\|_\infty \leq \int_0^t e^{-(2\beta+1)(t-s)} ds \leq t.$$

We have also

$$\begin{aligned} |S_t^{2\beta+1, 4\alpha} v_0(x) - v_0(x)| &\leq e^{-(2\beta+1)t} \int_{\mathbb{T}^d} s(4\alpha t, x, y) |v_0(y) - v_0(x)| dy \\ &\leq L e^{-(2\beta+1)t} \int_{\mathbb{R}^d} s_0(4\alpha t, x, y) \|y - x\|_2 dy \\ &\leq L e^{-(2\beta+1)t} \sqrt{4\alpha t d} \leq L \sqrt{4\alpha d} t^{1/2} \end{aligned}$$

where L is the Lipschitz constant of v_0 and the third inequality comes from the computation of the upper bound of the quadratic norm of d independent random variables with common variance $4\alpha t$. Therefore we obtain

$$\|v(t) - v_0\|_\infty \leq (L\sqrt{4\alpha d} + \sqrt{T})t^{1/2} \quad (6.44)$$

For the second bound, we use the fact that $\frac{d}{dx_i} s_0(t, x, y) = -\frac{d}{dy_i} s_0(t, x, y)$, thus we have, using an integration by parts

$$\partial_{x_i} S_t^{2\beta+1, 4\alpha}(v_0) = S_t^{2\beta+1, 4\alpha}(\partial_{x_i} v_0). \quad (6.45)$$

Then, we have

$$\partial_{x_i} v(t) - \partial_{x_i} v_0 = S_t^{2\beta+1, 4\alpha}(\partial_{x_i} v_0) - (\partial_{x_i} v_0) + \int_0^t \int_{\mathbb{R}^d} e^{-(2\beta+1)(t-s)} \partial_{x_i} s_0(4\alpha(t-s), x, y) w(s, y) dy ds. \quad (6.46)$$

We can treat both integrals as before, for the last integral :

$$\begin{aligned} \left| \int_0^t \int_{\mathbb{R}^d} e^{-(2\beta+1)(t-s)} \partial_{x_i} s_0(4\alpha(t-s), x, y) w(s, y) dy ds \right| &\leq \int_0^t \frac{e^{-(2\beta+1)(t-s)}}{4\alpha(t-s)} \int_{\mathbb{R}^d} |x_i - y_i| s_0(4\alpha(t-s), x, y) dy ds \\ &\leq \int_0^t \frac{e^{-(2\beta+1)(t-s)}}{4\alpha(t-s)} \sqrt{2\alpha(t-s)} ds \\ &\leq \frac{1}{\sqrt{2\alpha}} \int_0^{\sqrt{t}} e^{-(2\beta+1)u^2} du \leq \frac{\sqrt{t}}{\sqrt{2\alpha}}. \end{aligned}$$

For the first integral, we have the same estimates as before

$$|S_t^{2\beta+1,4\alpha}(\partial_{x_i}v_0)(x) - \partial_{x_i}v_0(x)| \leq L'\sqrt{4\alpha d}t^{1/2} \quad (6.47)$$

where L' is the maximum of the Lipschitz constants of $(\partial_{x_i}v_0)_i$. Thus, we get

$$\|\nabla v(t) - \nabla v_0\|_\infty \leq \left(L'\sqrt{4\alpha d} + \frac{1}{\sqrt{2\alpha}} \right) t^{1/2} \quad (6.48)$$

□

We now prove Proposition 6.4.

Proof. Let us assume that we have two solutions, v_1 and v_2 and let $e(t) = \|v_1 - v_2\|_{L^\infty([0,t] \times \mathbb{R}^d)}$. Note that the previous Lemma entails that, for all $T > 0$, there exists C , such that for $t < T$, we have $e(t) \leq C\sqrt{t}$.

We define $I_{s,t}^+ = \{(s, y), s \leq t, |v_1(s, y) - 2\rho| \leq e(t)\}$ and $I_{s,t}^- = \{(s, y), s \leq t, |v_1(s, y) + 2\rho| \leq e(t)\}$.

Since v_1 and v_2 are solutions of (6.23), there exists w_1 and w_2 such that $w_1 \in \partial H_\infty(v_1)$ a.e. and $w_2 \in \partial H_\infty(v_2)$. We can decompose each w_i as $w_i(t, x) = f_\infty(v_i(t, x)) + g_i(t, x)$ where f_∞ is the continuous part of ∂H_∞ :

$$f_\infty(q) = \begin{cases} -2\rho & \text{for } q \in [-1, -2\rho] \\ q & \text{for } q \in [-2\rho, 2\rho] \\ 2\rho & \text{for } q \in [2\rho, 1] \end{cases} \quad (6.49)$$

and $g_i(t, x) = w_i(t, x) - f_\infty(v_i(t, x))$. Note that

$$\begin{aligned} g_i(t, x) &= -1 + 2\rho \text{ a.e. on } \{(t, x), v_i(t, x) < -2\rho\}, \\ g_i(t, x) &= 0 \text{ a.e. on } \{(t, x), -2\rho < v_i(t, x) < 2\rho\}, \\ g_i(t, x) &= 1 - 2\rho \text{ a.e. on } \{(t, x), v_i(t, x) > 2\rho\}, \end{aligned} \quad (6.50)$$

since $w_i = h_\infty(v_i)$ a.e. on $\{(t, x), |v_i(t, x)| = 2\rho\}$.

As a consequence we have that, up to a negligible set, $\{(s, y), s \leq t, g_1(s, y) \neq g_2(s, y)\} \subset I_{s,t}^+ \cup I_{s,t}^-$ since, $g_1(s, y) \neq g_2(s, y)$ entails that one of the following inequalities is true $v_2(s, y) < 2\rho < v_1(s, y)$ or $v_2(s, y) < -2\rho < v_1(s, y)$ or $v_1(s, y) < 2\rho < v_2(s, y)$ or $v_1(s, y) < -2\rho < v_2(s, y)$. For each case, the inclusion is true : for the first one for example, if $s \leq t$ and $v_2(s, y) < 2\rho < v_1(s, y)$

$$e(t) \geq v_1(s, y) - v_2(s, y) = v_1(s, y) - 2\rho + 2\rho - v_2(s, y) \geq v_1(s, y) - 2\rho \geq 0 \quad (6.51)$$

thus $(s, y) \in I_{s,t}^+$. The same is true for the other cases.

Therefore we obtain the following expression for the difference $v_1 - v_2$:

$$\begin{aligned} v_1(t, x) - v_2(t, x) &= \int_0^t \int_{\mathbb{T}^d} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) (f_\infty(v_1(s, y)) - f_\infty(v_2(s, y))) dy ds \\ &\quad + \int_0^t \int_{I_{s,t}^+ \cup I_{s,t}^-} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) (g_1(s, y) - g_2(s, y)) dy ds. \end{aligned} \quad (6.52)$$

For the first integral in (6.52) we note that f_∞ is 1-Lipschitz, thus

$$\begin{aligned} &\left| \int_0^t \int_{\mathbb{T}^d} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) (f_\infty(v_1(s, y)) - f_\infty(v_2(s, y))) dy ds \right| \\ &\leq \int_0^t \int_{\mathbb{T}^d} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) e(t) dy ds \leq te(t). \end{aligned}$$

For the second integral in (6.52) we note that $|g_i| \leq 1 - 2\rho$, then we first have

$$\begin{aligned} &\left| \int_0^t \int_{I_{s,t}^+ \cup I_{s,t}^-} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) (g_1(s, y) - g_2(s, y)) dy ds \right| \\ &\leq 2(1 - 2\rho) \int_0^t \int_{I_{s,t}^+ \cup I_{s,t}^-} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) dy ds. \end{aligned}$$

Let $s \leq t$, since v_0 is regular on the level set $\{v_0 = 2\rho\}$ which is compact (since \mathbb{T}^d is) and ∇v_0 is a Lipschitz function, we can find $\delta, \eta > 0$ such that on $\{v_0 = 2\rho\} + B_\delta(0)$, $|\nabla v_0(x)| > \eta$. Using the second part of Lemma 6.5, and since $e(t) \leq C\sqrt{t}$, there exists $T > 0$ such that for $s \leq t \leq T$, $I_{s,t}^+ \subset \{v_0 = 2\rho\} + B_\delta(0)$ and on $I_{s,t}^+$, $|\nabla v_1(s)| > \eta/2$.

Since $I_{s,t}^+$ is compact and $\nabla v_1(s) \neq 0$, by the implicit function theorem, we can find a finite cover by open balls $(B_i)_{1 \leq i \leq N}$ centered on points on $I_{s,t}^+$ such that locally on each ball B_i , the level set $\{v_1(s, y) = 2\rho\}$ is the graph of a function, e.g. $y_1 = \varphi(y_2, \dots, y_d)$. Note that since $\{v_0 = 2\rho\}$ is compact, N is uniform in $s \leq T$, since by the lemma, we can make the cover of open balls on $\{v_0 = 2\rho\}$ and take their traces on $\{v_1(s, y) = 2\rho\}$.

By the mean value theorem on the first coordinate y_1 of $v_1(s)$, we have $I_{s,t}^+ \cap B_i \subset [-2e(t)/\nu, 2e(t)/\nu] \times \Pi_1(B_i)$, where Π_1 is the projection along the first coordinate. Thus,

$$\begin{aligned} & \int_0^t \int_{I_{s,t}^+ \cap B_i} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) dy ds \\ & \leq \int_0^t e^{-(2\beta+1)(t-s)} \frac{4e(t)}{\nu\sqrt{4\alpha(t-s)}} \int_{\Pi_1(B_i)} s(4\alpha(t-s), 0, (0, y_2, \dots, y_d)) dy_2 \cdots dy_d ds \\ & \leq \int_0^t e^{-(2\beta+1)(t-s)} \frac{2e(t)}{\nu\sqrt{\alpha(t-s)}} ds \leq \frac{2e(t)t^{1/2}}{\nu\sqrt{\alpha}}. \end{aligned}$$

Thus

$$\int_0^t \int_{I_{s,t}^+} e^{-(2\beta+1)(t-s)} s(4\alpha(t-s), x, y) dy ds \leq \frac{2Ne(t)t^{1/2}}{\nu\sqrt{\alpha}}. \quad (6.53)$$

Since the same holds for $I_{s,t}^-$, we obtain, that for some constant $C > 0$, and all $t < T$

$$|v_1(t, x) - v_2(t, x)| \leq (t + Ct^{1/2})e(t) \leq (T + CT^{1/2})e(T). \quad (6.54)$$

Then $e(T) \leq (T + CT^{1/2})e(T)$, and taking T small enough, we obtain $e(T) = 0$ thus $v_1 = v_2$ on $[0, T] \times \mathbb{T}^d$. \square

Remark 6.3. Maximal and minimal solutions. Another approach to (6.2) is to use a monotone construction of solutions, which arises from a comparison principle close to the one developed in Proposition 5.1. This was done initially in [2] and also in [5] Define $\overline{h_\infty}$ the right continuous version of h_∞ (Equation 6.20) by

$$\overline{h_\infty}(q) := -\mathbb{1}_{q < -2\rho} + q\mathbb{1}_{-2\rho \leq q < 2\rho} + \mathbb{1}_{q \geq 2\rho}. \quad (6.55)$$

Note that h_∞ and $\overline{h_\infty}$ are non decreasing (recall that $2\rho \in [0, 1]$). Recall that (S_t) is the semigroup on $L^1(\mathbb{T}^d)$ associated to $-2\alpha\Delta$, we denote

$$\underline{F}(v)(t, x) := e^{-(2\beta+1)t} S_t v_0(x) + \int_0^t e^{-(2\beta+1)(t-s)} S_{t-s} [h_\infty(v(s, \cdot))](x) ds \quad (6.56)$$

$$\overline{F}(v)(t, x) := e^{-(2\beta+1)t} S_t v_0(x) + \int_0^t e^{-(2\beta+1)(t-s)} S_{t-s} [\overline{h_\infty}(v(s, \cdot))](x) ds. \quad (6.57)$$

Then, fixed points of the maps above are mild solutions of these two formulations of our subdifferential inclusion:

$$\partial_t v(t, x) - 2\alpha\Delta v(t, x) + (2\beta + 1)v(t, x) = h_\infty(v(t, x)) \quad (6.58)$$

$$\partial_t v(t, x) - 2\alpha\Delta v(t, x) + (2\beta + 1)v(t, x) = \overline{h_\infty}(v(t, x)) \quad (6.59)$$

Since h_∞ (resp. $\overline{h_\infty}$) is non decreasing and that $h_\infty(p) \leq \overline{h_\infty}(p)$ for all $p \in [-1, 1]$, we have that, for u, v two functions such that $-1 \leq v \leq u \leq 1$,

1. $\underline{F}(v)(t, x) \leq \underline{F}(u)(t, x)$
2. $\overline{F}(v)(t, x) \leq \overline{F}(u)(t, x)$
3. $\underline{F}(u)(t, x) \leq \overline{F}(u)(t, x)$

We define the sequences $(V^n)_n$ and $(W^n)_n$ of functions on $\mathbb{R}^+ \times \mathbb{T}^d$: $V^0(t, x) = 1$, $W^0(t, x) = -1$ and for all $n \geq 1$

$$V^n(t, x) = e^{-(2\beta+1)t} S_t v_0(x) + \int_0^t e^{-(2\beta+1)(t-s)} S_{t-s} [\bar{f}(V^{n-1}(s, \cdot))](x) ds \quad (6.60)$$

$$W^n(t, x) = e^{-(2\beta+1)t} S_t v_0(x) + \int_0^t e^{-(2\beta+1)(t-s)} S_{t-s} [\underline{f}(W^{n-1}(s, \cdot))](x) ds. \quad (6.61)$$

Thus, for $-1 \leq v_0(x) \leq 1$, we can prove by induction that the sequences (V^n) and (W^n) satisfy, for all n

$$-1 \leq W^1 \leq W^2 \dots \leq W^n \leq V^n \leq \dots \leq V^2 \leq V^1 \leq 1. \quad (6.62)$$

By a compactness and monotony argument, one can prove that (W^n) and (V^n) converge to functions w and v which are mild solutions of the subdifferential inclusion. These are the minimal and maximal solutions of the subdifferential inclusion, in the sense that any other solution (Definition 6.1) must be bounded below by w and above by v . Uniqueness follows if one can prove that $w = v$.

7 Convergence of the discrete PDE

In analogy with the continuous setting, we define $v^N = 2u^N - 1$ where u^N is the solution of the discretized Equation (4.10) and $H(i, v^N) = 2G(i, \frac{v^N+1}{2}) + v_i^N$.

In such a way (4.10) becomes

$$\begin{cases} \partial_t v^N(t, i) = 2\alpha N^2 \Delta u^N(t, i) - (2\beta + 1)v^N(t, i) + H(i, v^N) \\ v^N(t, i) = 2u_0^N - 1(i), \end{cases} \quad (7.1)$$

We main goal of this section is to prove the following result which states the convergence of v^N .

Theorem 7.1. *Let v^N be the solution of (7.1). Then there exists a sub-sequence k_N such that*

$$\lim_{N \rightarrow +\infty} \|v^{k_N} - v_\infty\|_{\mathbb{T}^d \times]0, +\infty[} = 0 \quad (7.2)$$

where v_∞ is a solution of (6.23) and $\|v^N - v_\infty\|_{\mathbb{T}^d \times]0, +\infty[}$ means the uniform norm on all compact sets of $\mathbb{T}^d \times]0, +\infty[$.

In particular, whenever the solution of (7.1) is a.e. unique, v_∞ is also (the) mild solution of (6.2) and the whole sequence v^N converges to v_∞ , uniformly on all compact sets of $\mathbb{T}^d \times]0, +\infty[$.

To prove Theorem 7.1 we need some technical results. Let consider the semigroup of the discrete Laplacian $\frac{1}{2}N^2 \Delta^N$ on \mathbb{T}_N^d and \mathbb{Z}^d , denoted by $s^N(t, i, j)$ and $s_0^N(\gamma t, i, j)$ respectively. In particular we have that $s^N(t, i, j) = p_N(t, i, j)$, where $p_N(t, i, j)$ is heat kernel of discrete Laplacian on the discrete torus, cf. (5.2).

For any $\lambda, \gamma \geq 0$ and $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, we let $(S_t^{N, \lambda, \gamma})$ be the semigroup defined by

$$S_t^{N, \lambda, \gamma} f(x) = \sum_{y \in \frac{1}{N} \mathbb{T}_N^d} e^{-\lambda t} s^N(\gamma t, Nx, Ny) f(y) = \sum_{y \in \frac{1}{N} \mathbb{Z}^d} e^{-\lambda t} s_0^N(\gamma t, Nx, Ny) \tilde{f}(y) \quad (7.3)$$

where, as in (6.6), \tilde{f} is the periodic extension of f on $\frac{1}{N} \mathbb{Z}^d$.

In the remaining part of this article we will consider $S_t^{N, \lambda, \gamma} f(x)$ with $f \in \mathcal{C}(\mathbb{T}^d)$. In that case we mean that the function f is restricted on $\frac{1}{N} \mathbb{T}_N^d \subset \mathbb{T}^d$, which is equivalent to consider $f^N(x) := f(\lfloor Nx \rfloor / N)$. We observe that if f is also Lipschitz, then $\|f - f^N\|_{\mathbb{T}^d} \leq \frac{c}{N^d}$ for some $c > 0$ and $\|f^N\|_{\mathbb{T}^d} \leq \|f\|_{\mathbb{T}^d}$. Then, with the same extension to \mathbb{T}^d for s^N we can write, for any $x \in \mathbb{T}^d$

$$S_t^{N, \lambda, \gamma} f(x) = \int_{\mathbb{T}^d} e^{-\lambda t} N^d s^N(\gamma t, Nx, Ny) f^N(y) dy. \quad (7.4)$$

By a slight abuse of notation, we still denote by v^N the linear interpolation on \mathbb{T}^d such that $v^N(t, \frac{i}{N}) = v_i^N(t)$. We also redefine the function H on the torus \mathbb{T}^d by the linear interpolation such that $H(\frac{i}{N}, v^N) = H(i, v^N)$ and we define H^N as f^N in (7.4).

Definition 7.1. Let $N \in \mathbb{N}$ and $v_0^N \in L^\infty(\mathbb{T}^d)$. We say that $(v^N(t, x), t \geq 0, x \in \mathbb{T}^d)$ is a mild solution of (7.1) if

- for any N , v^N is continuous from \mathbb{R}_+^* to $L^\infty(\mathbb{T}^d)$,
- for all $t > 0$ and $x \in \mathbb{T}^d$

$$v^N(t, x) = S_t^{N, 2\beta+1, 4N^2\alpha}(v_0^N)(x) + \int_0^t S_{t-s}^{N, 2\beta+1, 4N^2\alpha} \left[H^N(\cdot, v^N(s, \cdot)) \right](x) ds. \quad (7.5)$$

Let u^N be the unique solution of (4.10), so that $v^N = 2u^N - 1$ satisfies (7.1). Of course, for any N the solution v^N of (7.5) exists and it is unique.

The proof of Theorem 7.1 is based on the representation of v^N as in (7.5). We define \tilde{v}^N as a slight modification of (7.5), that is,

$$\tilde{v}^N(t, x) := S_t^{2\beta+1, 4\alpha}(v_0)(x) + \int_0^t S_{t-s}^{2\beta+1, 4\alpha} \left[H(\cdot, v^N(s, \cdot)) \right](x) ds. \quad (7.6)$$

Lemma 7.2. For any $T > 0$

$$\lim_{N \rightarrow +\infty} \|\tilde{v}^N - v^N\|_{[1/T, T] \times \mathbb{T}^d} = 0. \quad (7.7)$$

Proof. Let $T > 0$, then

$$\begin{aligned} \sup_{t \in [\frac{1}{T}, T], x \in \mathbb{T}^d} \left| \tilde{v}^N(t, x) - v^N(t, x) \right| &\leq \sup_{t \in [\frac{1}{T}, T], x \in \mathbb{T}^d} \left| S_t^{N, \lambda, N^2\gamma} v_0^N(x) - S_t^{\lambda, \gamma} v_0(x) \right| \\ &+ \sup_{t \in [\frac{1}{T}, T], x \in \mathbb{T}^d} \left| \int_0^t \left\{ S_{t-s}^{N, 2\beta+1, 4N^2\alpha} \left[H^N(\cdot, v^N(s, \cdot)) \right](x) - S_{t-s}^{2\beta+1, 4\alpha} \left[H(\cdot, v^N(s, \cdot)) \right](x) \right\} ds \right|. \end{aligned} \quad (7.8)$$

We show that the right hand side of (7.8) converges to 0. We detail the convergence of the second term, which is more delicate. The argument can be adapted to the first term by using Assumption 2 which ensures that v_0^N converges to v_0 in $C(\mathbb{T}^d)$.

We fix $\varepsilon \in (0, \frac{1}{T})$ and we get that

$$\begin{aligned} &\left| \int_0^t S_{t-s}^{N, 2\beta+1, 4N^2\alpha} \left[H^N(\cdot, v^N(s, \cdot)) \right](x) - S_{t-s}^{2\beta+1, 4\alpha} \left[H(\cdot, v^N(s, \cdot)) \right](x) ds \right| \\ &\leq \left| \int_0^{t-\varepsilon} S_{t-s}^{N, 2\beta+1, 4N^2\alpha} \left[H^N(\cdot, v^N(s, \cdot)) \right](x) - S_{t-s}^{2\beta+1, 4\alpha} \left[H(\cdot, v^N(s, \cdot)) \right](x) ds \right| + C\varepsilon, \end{aligned} \quad (7.9)$$

where we used that Lemma B.6 which implies that $H(\cdot, v^N)$ is bounded by 1 uniformly on N . The integral on the right hand side of (7.9) is bounded from above by

$$\begin{aligned} &\int_0^{t-\varepsilon} e^{-(2\beta+1)(t-s)} \left| \int_{\mathbb{T}^d} \left(N^d s^N (4N^2\alpha(t-s), Nx, Ny) - s(4\alpha(t-s), x, y) \right) H^N(y, v^N(s, y)) dy \right| ds \\ &+ \int_0^{t-\varepsilon} e^{-(2\beta+1)(t-s)} \int_{\mathbb{T}^d} s(4\alpha(t-s), x, y) \left| H^N(y, v^N(s, y)) - H(y, v^N(s, y)) \right| dy ds. \end{aligned} \quad (7.10)$$

Since $\sup_{s \geq 0, y \in \mathbb{T}^d} \left| H^N(y, v^N(s, y)) - H(y, v^N(s, y)) \right| \leq \frac{1}{N}$, the second integral is smaller than $\frac{c_{\alpha, \beta}}{N}$. For the first integral, we first use that $H(\cdot, v^N)$ is bounded by 1 uniformly on N and then we operate the change of variable $u = t - s$ which gives that it is bounded from above by

$$\int_\varepsilon^t e^{-(2\beta+1)u} \int_{\mathbb{T}^d} \left| N^d s^N (4N^2\alpha u, Nx, Ny) - s(4\alpha u, x, y) \right| dy du \quad (7.11)$$

We now use the local central limit theorem (cf. Theorem 2.1.1 and (2.5) in [17]): let ρ be the Gaussian Kernel and $\rho(u, x, y) = \frac{1}{u^{d/2}} \rho\left(\frac{x-y}{u^{1/2}}\right)$, then

$$\psi_{\varepsilon, T, N} := \sup_{u \in [\varepsilon, T], y \in \frac{1}{N}\mathbb{Z}^d} \left| N^d p(uN^2, Nx, Ny) - \rho(u, x, y) \right| \xrightarrow{N \rightarrow +\infty} 0. \quad (7.12)$$

We also observe that by symmetry the supremum in (7.12) is independent of x . Moreover, by Proposition 2.4.6 in [17] we have that there exist $c_1, c_2 > 0$ independent of x, y, u such that

$$\left| N^d p(uN^2, Nx, Ny) - \rho(u, x, y) \right| \leq \frac{c_1}{u^{\frac{d}{2}}} e^{-c_2 \frac{\|x-y\|^2}{u}}. \quad (7.13)$$

In such a way, for any $M > 0$ fixed we write $\mathbb{T}^d = B_x(M) \cup B_x(M)^c$ and we get that (7.11) is smaller than

$$c_d \int_{\varepsilon}^t e^{-(2\beta+1)u} \left(M^d \psi_{N,\varepsilon,T} + \frac{c_1}{M^d u^{\frac{d}{2}}} e^{-c_2 \frac{M^2 d}{u}} \right) ds \leq C_d \left(M^d \psi_{N,\varepsilon,T} + \frac{1}{M^d} \right), \quad (7.14)$$

where $c_d, C_d > 0$ are two positive constants that depend only on the dimension d .

We conclude that the right hand side of (7.9) is bounded by $\frac{c_{\alpha,\beta}}{N} + cM^d \psi_{N,\varepsilon,T} + \frac{c}{M^d} + C\varepsilon$, uniformly on $x \in \mathbb{T}^d$ and $t \in [\frac{1}{T}, T]$. Therefore, by taking the limit on $N \rightarrow +\infty$ and then on $M \rightarrow +\infty$ and $\varepsilon \rightarrow 0$ we conclude the proof. \square

Proof of Theorem 7.1. We control \tilde{v}^N to get the convergence of v^N . We observe that since $H(\cdot, v^N(s, \cdot))$ is uniformly bounded so that by Proposition 6.1 \tilde{v}^N is uniformly bounded in N , uniformly continuous on $[1/T, T] \times \mathbb{T}^d$, and the modulus of continuity only depends on $T > 1$. By the Ascoli-Arzelà Theorem, the sequence $(\tilde{v}^N)_N$ is compact on $C([1/T, T] \times \mathbb{R}^d)$ and we can extract (by a diagonal argument) a subsequence converging uniformly to a limit v_∞ in $C([0, \infty[\times \mathbb{R}^d)$, uniformly on each compact. By Lemma (7.2) we get that v^N admits a sub-sequence converging to v_∞ .

Using Corollary B.4, we can adapt the argument used in the proof of Proposition 6.3 (6.25–6.30) to get that v_∞ is a mild solution of (6.23), we omit the details. \square

7.1 Proof of Theorem 3.1

Theorem 3.1 is now a consequence of Theorems 4.2 and 7.1.

A Concentration inequalities

We follow the definitions in Jara and Menezes [15] and [14] and Boucheron, Lugosi, Massart [1], Section 2.3. We omit the proofs since there are present in the references.

Definition A.1 ([15] and [1], Section 2.3). *Let X be a real random variable. X is said to be sub-Gaussian with variance parameter σ^2 if, for all $t \in \mathbb{R}$*

$$\psi_X(t) := \log \mathbb{E}(\exp(tX)) \leq \sigma^2 \frac{t^2}{2}. \quad (\text{A.1})$$

We denote $\mathcal{G}(\sigma^2)$ the set of real sub-Gaussian random variables with variance parameter σ^2 .

Proposition A.1. *[[1] and [14], Proposition F.7] The following statements are equivalent :*

1. $X \in \mathcal{G}(\sigma^2)$
2. $\mathbb{P}(|X| > t) \leq 2 \exp(-\frac{t^2}{2\sigma^2})$
3. $\mathbb{E}(\exp(\gamma X^2)) \leq 3$ for all $0 \leq \gamma < \frac{1}{4\sigma^2}$.

Let us complete our family of inequality :

Lemma A.2. *[[14], Proposition F.8] Let $X \in \mathcal{G}(\sigma_1^2)$ and $Y \in \mathcal{G}(\sigma_2^2)$, then for all $0 \leq \gamma < \frac{1}{4\sigma_1\sigma_2}$,*

$$\mathbb{E}(\exp(\gamma XY)) \leq 3$$

Lemma A.3. *[[14], Proposition F.12] Let X_1, \dots, X_n be random variables with $X_i \in \mathcal{G}(\sigma_i^2)$, such that there is a partition of K subsets P_1, \dots, P_K each containing L variables that $\Sigma_k = \sigma(X_i, i \in P_k)$ are independent σ -algebra then, for all real $\alpha_1, \dots, \alpha_n$, the random variable $Y = \sum_i \alpha_i X_i$ is sub-Gaussian with variance parameter $L \sum_i \alpha_i^2 \sigma_i^2$.*

Note that if $L = 1$, the variables are independent.

Lemma A.4 (Hoeffding Inequality, [1], Section 2.3). *Let X be a bounded random variable with $X \in [a, b]$, then $X - \mathbb{E}X \in \mathcal{G}\left(\frac{(b-a)^2}{4}\right)$.*

B Controls for the non-linearities

In this section, we collect some results about the specifics of our model. Recall the notations: for $\eta \in \{0, 1\}^{\mathbb{T}_N^d}$

$$c_0^+(\eta) := \mathbb{1}_{\{\rho_0(\eta) \geq 1 - \frac{\kappa_N}{K_N}\}}, \quad c_0^-(\eta) := \mathbb{1}_{\{\rho_0(\eta) \leq \frac{\kappa_N}{K_N}\}},$$

and $\kappa_N = \kappa_N(T) := \min \left\{ [K_N T] - 1; [K_N(1-T)] \right\}$. For any function $u = (u_i)_{i \in \mathbb{T}_N^d}$, we define $v_u(d\eta) = v_u^N(d\eta) := \otimes_{i \in \mathbb{T}_N^d} B(u_i)$ where $B(u_i)$ denote a Bernoulli distribution with parameter u_i . and we let $c_0^+(u)$ and $c_0^-(u)$ be the expectations of $c_0^+(\eta)$ and $c_0^-(\eta)$ under v_u . We set $(\tau_i \eta)_j = \eta_{i+j}$, and likewise τ_i acts on u . Then,

$$G(u) := (1 - u_0)c_0^+(u) - u_0c_0^-(u) = (1 - u_0)\mathbb{P}_{v_u} \left[\rho_0(\eta) \geq 1 - \frac{\kappa_N}{K_N} \right] - u_0\mathbb{P}_{v_u} \left[\rho_0(\eta) \leq \frac{\kappa_N}{K_N} \right] \quad (\text{B.1})$$

and $G(i, u) := G(\tau_i u)$. We start with some results on the non linearity G :

Proposition B.1. *Let $u, v \in [0, 1]^{\mathbb{T}_N^d}$ such that $u_i \geq v_i$ for all $i \in \mathcal{V}_N$ and $u_0 = v_0$. Then $G(0, u) \geq G(0, v)$.*

Proof. We construct a coupling between v_u and v_v : let (U_i) be independent and identically distributed random variables uniform on $[0, 1]$. Define $\eta_i = \mathbb{1}_{\{U_i \leq u_i\}}$ and $\eta'_i = \mathbb{1}_{\{U_i \leq v_i\}}$. We have $\eta_i \geq \eta'_i$ for all $i \in \mathcal{V}_N$, therefore $\rho_0(\eta) \geq \rho_0(\eta')$. This proves that $c_0^+(u) \geq c_0^+(v)$ and $c_0^-(u) \leq c_0^-(v)$. The results follows since $u_0 = v_0$. \square

For $p \in [0, 1]$, $T \in [0, 1]$, $K \in \mathbb{N}^*$, we let $\kappa(K, T) = \min \left\{ [KT] - 1; [K(1-T)] \right\} \leq K/2$, and we define $g_K(p)$ as

$$g_K(p) := (1-p)\mathbb{P}_p[X > K - \kappa(K, T)] - p\mathbb{P}_p[X \leq \kappa(K, T)]$$

where X is a random variable with binomial distribution with parameter (K, p) under \mathbb{P}_p . In particular, we have that $g_{K_N}(p) = G(i, u)$ for $u(t, i) = p$ for all $i \in \mathbb{T}_N^d$. Note that g_K is $C^\infty([0, 1])$. We also have

$$\begin{aligned} g_K(p) &= (1-p)\mathbb{P}_{1-p}[X < \kappa(K, T)] - p\mathbb{P}_p[X \leq \kappa(K, T)] \\ &= \sum_{k=0}^{\kappa(K, T)-1} \binom{K}{k} \left[(1-p)^{k+1} p^{K-k} - p^{k+1} (1-p)^{K-k} \right] - \binom{K}{\kappa(K, T)} p^{\kappa(K, T)+1} (1-p)^{K-\kappa(K, T)}. \end{aligned} \quad (\text{B.2})$$

We recall $g_\infty(p) := (1-p)\mathbb{1}_{\{1-p < p_0(T)\}} - p\mathbb{1}_{\{p \leq p_0(T)\}}$, where $p_0(T) = \min(T, 1-T)$.

The following proposition estimates the convergence of g_K to g_∞ , in particular we prove that close to $p = 0$ (resp. $p = 1$), g_K is negative (resp. positive).

Proposition B.2. *For all $K \in \mathbb{N}^* \cup \{\infty\}$, we have $g_K(0) = g_K(1) = 0$. For all $K \in \mathbb{N}^*$ and $p \in [0, 1]$, we have, for $|p - p_0(T)| > \frac{1}{K}$ and $|(1-p) - p_0(T)| > \frac{1}{K}$*

$$|g_K(p) - g_\infty(p)| \leq 2e \exp(-2K(p_0(T) - p)^2) + 2e \exp(-2K(p_0(T) - (1-p))^2) \quad (\text{B.3})$$

In particular, for any $0 < \delta < \frac{1}{4e}$, $T \in [4\delta, 1-4\delta]$, and $K > \frac{|\log(\delta)|}{4\delta^2}$, we have that $g_K(2\delta) < -\delta$ and $g_K(1-2\delta) > \delta$.

Proof. We consider g_K written as in (B.2). Then, the values at $p = 0$ and $p = 1$ are obvious.

Note that, under \mathbb{P}_p , $\frac{X}{K} - p$ converges to 0 (in $L^2(\Omega_N)$) and is sub-Gaussian with variance parameter $\frac{1}{4K}$, thus, for any $t > 0$,

$$\mathbb{P}_p \left(\left| \frac{X}{K} - p \right| > t \right) \leq 2 \exp(-2Kt^2).$$

We also have that

$$0 \leq p_0(T) - \frac{\kappa(K, T)}{K} \leq \frac{1}{K}.$$

Then, for $p > p_0(T)$,

$$\begin{aligned} \mathbb{P}_p[X \leq \kappa(K, T)] &= \mathbb{P}_p \left[p - \frac{X}{K} \geq p - p_0(T) + p_0(T) - \frac{\kappa(K, T)}{K} \right] \leq \mathbb{P}_p \left[p - \frac{X}{K} \geq p - p_0(T) \right] \\ &\leq 2 \exp(-2K(p - p_0(T))^2), \end{aligned}$$

and for $p < p_0(T) - \frac{1}{K}$,

$$\begin{aligned}\mathbb{P}_p[X > \kappa(K, T)] &= \mathbb{P}_p\left[\frac{X}{K} - p > \frac{\kappa(K, T)}{K} - p_0(T) + p_0(T) - p\right] \leq \mathbb{P}_p\left[\frac{X}{K} - p \geq p_0(T) - p - \frac{1}{K}\right] \\ &\leq 2\exp(-2K(p_0(T) - p - 1/K)^2) \leq 2e^{2(p_0(T) - p)} \exp(-2K(p_0(T) - p)^2) \\ &\leq 2e \exp(-2K(p_0(T) - p)^2).\end{aligned}$$

Thus, we have the following, for $|p - p_0(T)| > \frac{1}{K}$,

$$|\mathbb{P}_p[X \leq \kappa(K, T)] - \mathbb{1}_{\{p \leq p_0(T)\}}| < 2e \exp(-2K(p_0(T) - p)^2).$$

The results follows with the same estimates with $(1 - p)$ instead of p . This prove (B.3).

Let $\delta > 0$ and set $p = 2\delta$, and $T \in [4\delta, 1 - 4\delta]$, then $|2\delta - p_0(T)| \leq 2\delta$ for $K > \frac{1}{2\delta}$ and we have the same for $1 - p = 1 - 2\delta$. Applying the result, we have

$$|g_K(2\delta) - g_\infty(2\delta)| = |g_K(2\delta) + 2\delta| \leq 4e \exp(-8K\delta^2)$$

Then, if $4e \exp(-8K\delta^2) < \delta$, we get the result. This happens if $\exp(-8K\delta^2) < \delta^2$ and $\delta < \frac{1}{4e}$, which gives the condition on K . \square

Proposition B.3. *Let $v : \mathbb{T}^d \rightarrow [0, 1]$ be a continuous fixed density on the torus. For any $i \in \mathbb{T}_N^d$ we let $u_i := v_{i/N}$. Then as $i/N \rightarrow x$, we have that u_i converges to v_x . Let $v_u = \bigotimes_{i \in \mathbb{T}_N^d} \mathcal{B}(u_i)$. Then, $\rho_0(\eta)$ converges to v_0 in probability.*

Proof. We use the coupling introduced in Lemma B.1: we let (U_i) be i.i.d uniform random variables on $[0, 1]$ so that under v_u we have that $\rho_0(\eta)$ and $\frac{1}{K_N} \sum_{i \in \mathcal{V}_N} \mathbb{1}_{\{U_i < u_i\}}$ are equal in law. By the Tchebychev inequality, we have that for any $\varepsilon > 0$

$$\mathbb{P}\left(\left|\frac{\sum_{i \in \mathcal{V}_N} \mathbb{1}_{\{U_i < u_i\}}}{K_N} - \frac{\sum_{i \in \mathcal{V}_N} u_i}{K_N}\right| > \varepsilon\right) \leq \frac{1}{4K_N \varepsilon^2}.$$

Moreover, the sequence $K_N^{-1} \sum_{i \in \mathcal{V}_N} u_i$ converges to v_0 because for any $i \in \mathcal{V}_N$, $i/N \rightarrow 0$. \square

We have the following corollary.

Corollary B.4. *Let $u = (u_i)_{i \in \mathbb{T}_N^d}$ as in Proposition B.3. Then, $G(u)$ and $g_{K_N}(v_0)$ converge both to $g_\infty(v_0)$ as $N \rightarrow +\infty$.*

Recall that, for $q \in [-1, 1]$,

$$h_K(q) = 2g_K\left(\frac{1}{2}(q+1)\right) + q = (1-q)\mathbb{P}_{\frac{1-q}{2}}[X < \kappa(K, T)] - (1+q)\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + q.$$

Recall the critical parameter $\rho := \rho(T) = |T - \frac{1}{2}| = \frac{1}{2} - p_0(T) \in [0, \frac{1}{2}]$. Note that h_K converges pointwise to the function $h_\infty(q) = 2g_\infty\left(\frac{1+q}{2}\right) + q = -\mathbb{1}_{q \leq -2\rho} + q\mathbb{1}_{-2\rho < q \leq 2\rho} + \mathbb{1}_{q > 2\rho}$. The points $q = \pm 2\rho$ are the discontinuities of h_∞ , and compare the Lemma to the fact that $2\rho = h_\infty(2\rho^-)$ and $1 = h_\infty(2\rho^+)$. A similar estimate holds at $q = -2\rho$.

Lemma B.5. *For all $q \in [-1, 1]$, $|h_K(q)| \leq 1$. Moreover, for all $\varepsilon > 0$ such that $2\rho - \varepsilon > 0$, there is $K_0 > 0$ such that for $K \geq 0$,*

$$\begin{aligned}2\rho - 2\varepsilon = h_\infty(2\rho^-) - 2\varepsilon \leq h_K(q) \leq h_\infty(2\rho^+) = 1 & \quad \text{for } q \in [2\rho - \varepsilon, 2\rho + \varepsilon] \\ -1 = h_\infty(-2\rho^-) \leq h_K(q) \leq h_\infty(-2\rho^+) + 2\varepsilon = -2\rho + 2\varepsilon & \quad \text{for } q \in [-2\rho - \varepsilon, -2\rho + \varepsilon]\end{aligned}$$

Proof. We start by observing that for all $q \in [-1, 1]$, $\mathbb{P}_{\frac{1-q}{2}}[X < \kappa(K, T)] = \mathbb{P}_{\frac{1+q}{2}}[X > K - \kappa(K, T)]$. We also have that

$$1 = \mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + \mathbb{P}_{\frac{1+q}{2}}[\kappa(K, T) < X \leq K - \kappa(K, T)] + \mathbb{P}_{\frac{1+q}{2}}[X > K - \kappa(K, T)].$$

Thus, using the definition of h_K , we get

$$h_K(q) = -\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + q\mathbb{P}_{\frac{1+q}{2}}[\kappa(K, T) < X \leq K - \kappa(K, T)] + \mathbb{P}_{\frac{1+q}{2}}[X > K - \kappa(K, T)].$$

This gives us the result. Indeed, for $q \in [0, 1]$, we have

$$\begin{aligned} h_K(q) &\leq q\mathbb{P}_{\frac{1+q}{2}}[\kappa(K, T) < X \leq K - \kappa(K, T)] + \mathbb{P}_{\frac{1+q}{2}}[X > K - \kappa(K, T)] \\ &\leq \mathbb{P}_{\frac{1+q}{2}}[\kappa(K, T) < X] \leq 1 \end{aligned}$$

and

$$\begin{aligned} h_K(q) &\geq -\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + \mathbb{P}_{\frac{1+q}{2}}[X > K - \kappa(K, T)] \\ &\geq \mathbb{P}_{\frac{1-q}{2}}[X < \kappa(K, T)] - \mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] \geq -\mathbb{P}_{\frac{1+q}{2}}[X = \kappa(K, T)]. \end{aligned}$$

The last inequality comes from the fact that, by a coupling argument, $p \mapsto \mathbb{P}_p[X < \kappa(K, T)]$ is decreasing on $[0, 1]$, and since $q \geq 0$, $\frac{1-q}{2} \leq \frac{1+q}{2}$.

In particular, for $q \in [2\rho - \varepsilon, 2\rho + \varepsilon]$ such that $2\rho - \varepsilon > 0$, we have the same upper bound as before for $h_K(q)$, and for the lower bound:

$$\begin{aligned} h_K(q) &\geq -\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + q\mathbb{P}_{\frac{1+q}{2}}[X > \kappa(K, T)] \\ &\geq -\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] + (2\rho - \varepsilon) \left(1 - \mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)]\right) \\ &\geq 2\rho - \varepsilon - 2\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)]. \end{aligned}$$

From the proof of Proposition B.2, we have that, for $K \geq K_0$:

$$\mathbb{P}_{\frac{1+q}{2}}[X \leq \kappa(K, T)] \leq 2 \exp(-K(q + 2\rho)^2/2) \leq 2e^{-2K_0\rho^2}$$

Choosing K_0 large enough such that the right hand side is less than $\varepsilon/2$, we get the result.

For $q \in [-1, 0]$, the proof is completely similar. □

Lemma B.6. *Let $u = (u_i)_{i \in \mathbb{T}_N^d}$, with $u_i \in [0, 1]$ and let $v = 2u - 1$. Then, $|H(i, v)| \leq 1$, uniformly on $i \in \mathbb{T}_N^d$.*

Proof. We recall (4.9), in particular that $G(i, u) = G(\tau_i u)$. So that we only prove that $|H(v)| \leq 1$, where $H(v) = G((v+1)/2) + v_0$. The proof is similar to Lemma B.5. Indeed, again by (4.9) we have that

$$\begin{aligned} 2G\left(\frac{v+1}{2}\right) + v_0 &= c_0^+ \left(\frac{v+1}{2}\right) (1 - v_0) - c_0^- \left(\frac{v+1}{2}\right) (1 + v_0) + v_0 \\ &= c_0^+ \left(\frac{v+1}{2}\right) - c_0^- \left(\frac{v+1}{2}\right) + v_0 \left(1 - c_0^+ \left(\frac{v+1}{2}\right) - c_0^- \left(\frac{v+1}{2}\right)\right). \end{aligned}$$

We observe that, by definition, $1 - c_0^+ \left(\frac{v+1}{2}\right) - c_0^- \left(\frac{v+1}{2}\right) > 0$. This implies that $-1 \leq H(v) \leq 1$. □

References

- [1] S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities. A nonasymptotic theory of independence*. Oxford: Oxford University Press, corrected paperback edition edition, 2016.
- [2] S. Carl. The monotone iterative technique for a parabolic boundary value problem with discontinuous nonlinearity. *Nonlinear Anal., Theory Methods Appl.*, 13(12):1399–1407, 1989.
- [3] C. Claudio, F. Santo, and L. Vittorio. Statistical physics of social dynamics. *Rev. Mod. Phys.*, 81:591–647, 2009.
- [4] H. Deguchi. Existence, uniqueness and non-uniqueness of weak solutions of parabolic initial-value problems with discontinuous nonlinearities. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, 135(6):1139–1167, 2005.
- [5] H. Deguchi. Existence, uniqueness and non-uniqueness of weak solutions of parabolic initial-value problems with discontinuous nonlinearities. *Proc. R. Soc. Edinb., Sect. A, Math.*, 135(6):1139–1167, 2005.
- [6] T. Delmotte. Parabolic harnack inequality and estimates of markov chains on graphs. *Revista Matemática Iberoamericana*, 15:181–232, 1999.

- [7] T. Delmotte and J.-D. Deuschel. On estimating the derivatives of symmetric diffusions in stationary random environment, with applications to $\nabla\varphi$ interface model. *Probab. Theory Related Fields*, 133(3):358–390, 2005.
- [8] T. Funaki and K. Tsunoda. Motion by mean curvature from Glauber-Kawasaki dynamics. *J. Stat. Phys.*, 177(2):183–208, 2019.
- [9] L. Gauvin, J. Vannimenus, and J. Nadal. Phase diagram of a schelling segregation model. *Eur. Phys. J., B* 70:293–304, 2009.
- [10] G. Giacomin, S. Olla, and H. Spohn. Equilibrium fluctuations for $\nabla\varphi$ interface model. *Ann. Probab.*, 29:1138–1172, 2001.
- [11] R. Gianni. Existence of the free boundary in a multi-dimensional combustion problem. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, 125(3):525–544, 1995.
- [12] O. Hamed and F. Massimo. Evolution and limiting configuration of a long-range schelling-type spin system. arXiv: 1804.00358, 2018.
- [13] O. Hamed and F. Massimo. Self-organized segregation on the grid. *J. Stat. Phys.*, 170:748–783, 2018.
- [14] M. Jara and O. Menezes. Non-equilibrium fluctuations of interacting particle systems, 2018.
- [15] M. Jara and O. Menezes. Non-equilibrium fluctuations for a reaction-diffusion model via relative entropy. *Markov Process. Related Fields*, 26(1):95–124, 2020.
- [16] C. Kipnis and C. Landim. *Scaling limits of interacting particle systems*, volume 320 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [17] G. Lawler and V. Limic. *Random Walk: A Modern Introduction*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2010.
- [18] M. Liero and A. Mielke. Gradient structures and geodesic convexity for reaction-diffusion systems. *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 371(2005):20120346, 28, 2013.
- [19] A. Lucquiaud. *Modéliser l'émergence de la ségrégation sociospatiale et détecter ses structures*. PhD thesis, École doctorale n. 386 Sciences Mathématiques de Paris Centre, Statistique, Analyse et Modélisation Multidisciplinaire, 2022.
- [20] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Applied Mathematical Sciences. Springer New York, 2012.
- [21] T. Schelling. Dynamic models of segregation. *The Journal of Mathematical Sociology*, 1(2):143–186, 1971.
- [22] T. Schelling. *Micromotives and Macrobehavior*. Norton Company, 2006 (first published 1978).
- [23] D. W. Stroock and W. Zheng. Markov chain approximations to symmetric diffusions. *Annales de l'Institut Henri Poincaré (B) Probability and Statistics*, 33(5):619–649, 1997.