



**HAL**  
open science

# An appraisal of the frequency and severity of noninfectious manifestations in primary immunodeficiencies: A study of a national retrospective cohort of 1375 patients over 10 years

Mickaël Alligon, Nizar Mahlaoui, Virginie Courteille, Laurence Costes, Veronica Afonso, Philippe Randrianomenjanahary, Nathalie de Vergnes, Anja Ranohavimparany, Duy Vo, Inès Hafsa, et al.

## ► To cite this version:

Mickaël Alligon, Nizar Mahlaoui, Virginie Courteille, Laurence Costes, Veronica Afonso, et al.. An appraisal of the frequency and severity of noninfectious manifestations in primary immunodeficiencies: A study of a national retrospective cohort of 1375 patients over 10 years. *Journal of Allergy and Clinical Immunology*, 2022, 149 (6), pp.2116-2125. 10.1016/j.jaci.2021.12.790 . hal-03997839

**HAL Id: hal-03997839**

**<https://hal.science/hal-03997839>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1           **An appraisal of the frequency and severity of non-infectious manifestations in primary**  
2           **immunodeficiencies. A study of a national retrospective cohort of 1375 patients over 10 years**  
3  
4           **Mickaël Alligon, BS <sup>1,\*</sup>, Nizar Mahlaoui, MD, PhD <sup>1,2,\*</sup>, Virginie Courteille, BS <sup>1</sup>, Laurence Costes,**  
5           **BS <sup>1</sup>, Veronica Afonso, BS <sup>1</sup>, Philippe Randrianomenjanahary, BS <sup>1</sup>, Nathalie de Vergnes, BS <sup>1</sup>, Anja**  
6           **Ranohavimparany, BS <sup>1</sup>, Duy Vo, BS<sup>1</sup>, Inès Hafsa, BS <sup>1</sup>, Perrine Bach, BS <sup>1</sup>, Vincent Benoit, PhD <sup>3</sup>,**  
7           **Nicolas Garcelon, PhD <sup>3</sup>, Alain Fischer, MD, PhD<sup>1,2,4,5</sup>,**  
8           **& the members of the CEREDIH French PID study group <sup>§</sup>**

9  
10           **\*: Equal contribution**

- 11  
12           1. French National Reference Center for Primary Immune Deficiencies (CEREDIH), Necker  
13           Enfants Malades University Hospital, Assistance Publique-Hôpitaux de Paris (AP-HP), Paris,  
14           France.
- 15           2. Pediatric Immuno-Hematology and Rheumatology Unit, Necker Enfants Malades  
16           University Hospital, Assistance Publique-Hôpitaux de Paris (AP-HP), Paris, France.
- 17           3. Université de Paris, Imagine Institute, Data Science Platform, INSERM UMR 1163, F-75015,  
18           Paris, France
- 19           4. Paris Descartes University, Sorbonne Paris Cité, Imagine Institute, INSERM UMR 1163,  
20           Paris, France.
- 21           5. Collège de France, Paris, France

22  
23

24 § *(by alphabetical order)*:

25 **Wadih Abou-Chahla, Daniel Adoue, Nathalie Aladjidi, Corinne Armari-Alla, Vincent Barlogis,**  
26 **Sophie Bayart, Yves Bertrand, Stéphane Blanche, Damien Bodet, Bernard Bonnotte, Raphaël**  
27 **Borie, Patrick Boutard, David Boutboul, Claire Briandet, Jean-Paul Brion, Jacques Brouard,**  
28 **Liana Carausu, Martin Castelle, Pascal Cathebras, Emilie Catherinot, Nathalie Cheikh,**  
29 **Morgane Cheminant, Sarah Cohen-Beaussant, Thibault Comont, Louis-Jean Couderc, Pierre**  
30 **Cougoul, Gérard Couillault, Lionel Crevon, Elisa Demonchy, Anne Deville, Catherine**  
31 **Devoldere, Eric Dore, Fabienne Dulieu, Isabelle Durieu, Natacha Entz-Werle, Claire Fieschi,**  
32 **Fanny Fouyssac, Pierre Frange, Vincent Gajdos, Lionel Galicier, Virginie Gandemer, Martine**  
33 **Gardembas, Catherine Gaud, Bernard Grosbois, Aurélien Guffroy, Corinne Guitton, Gaëlle**  
34 **Guillerm, Eric Hachulla, Mohamed Hamidou, Sophie Haro, Yves Hatchuel, Olivier Hermine,**  
35 **Cyrille Hoarau, Arnaud Hot, Sébastien Humbert, Arnaud Jaccard, Jean-Philippe Jais, Sarah**  
36 **Jannier, Serge Jacquot, Roland Jaussaud, Pierre-Yves Jeandel, Eric Jeziorski, Kamila Kebaili,**  
37 **Anne-Sophie Korganow, Olivier Lambotte, Fanny Lanternier, Claire Larroche, David Launay,**  
38 **Emmanuelle Le Moigne, Alain Le Quellec, Emmanuelle Le Moigne, Vincent Le Moing, Yvon**  
39 **Lebranchu, Marc Lecuit, Guillaume Lefèvre, Jean-Daniel Lelièvre, Richard Lemal, Valérie Li-**  
40 **Thiao-Te, Olivier Lortholary, Luminita Luca, Coralie Mallebranche, Marion Malphettes, Aude**  
41 **Marie-Cardine, Nicolas Martin-Silva, Agathe Masseur, Françoise Mazingue, Etienne Merlin,**  
42 **Gérard Michel, Frédéric Millot, Charline Miot, Béatrice Monlibert, Fabrice Monpoux,**  
43 **Despina Moshous, Luc Mouthon, Martine Münzer, Robert Navarro, Bénédicte Neven, Dalila**  
44 **Nouar, Raphaële Nove-Josserand, Eric Oksenhendler, Marie Ouachée-Chardin, Anne**  
45 **Pagnier, Marlène Pasquet, Isabelle Pellier, Yves Perel, Antoinette Perlat, Christophe Piguet,**  
46 **Dominique Plantaz, Sophie Rivière, Pascal Roblot, Pierre-Simon Rohrlich, Bruno Royer,**  
47 **Valéry Salle, Françoise Sarrot-Reynauld, Amélie Servettaz, Jean-Louis Stephan, Nicolas**

48 **Schleinitz, Harry Sokol, Felipe Suarez, Laure Swiader, Sophie Taque, Caroline Thomas, Olivier**

49 **Tournilhac, Caroline Thumerelle, Jean-Pierre Vannier, Jean- François Viillard.**

50

51

52 **Corresponding author**

53 Alain Fischer

54 Address: Institut Imagine, 24 Boulevard du Montparnasse – 75015 Paris, France

55 Phone: +331 42 75 42 43 - Fax: +331 42 75 42 21 - E-mail: [alain.fischer@aphp.fr](mailto:alain.fischer@aphp.fr)

56

57

58 **Keywords**

59 Primary Immunodeficiencies; Inborn errors of immunity; epidemiology; cohort study;  
60 registry; non-infectious complications; autoimmunity; inflammation; malignancies; allergy

61

62 **Conflict-of-interest disclosure**

63 The authors declare no competing financial interests.

64

65 **Word count**

66 Abstract word count: 242

67 Main manuscript word count: 6822

68 Number of figures: 5

69 Number of tables: 3

70 Number supplementary figures/ table: 16

71 Number of references: 35

72

73 **Abbreviations (by alphabetical order)**

74 AT: Ataxia-Telangiectasia

75 CGD: Chronic granulomatous disease

76 CID: Combined Immunodeficiency

77 CVID: Common Variable Immunodeficiency

78 ESID: European Society for Immunodeficiencies

79 HIES: Hyper-IgE syndrome

80 HLH: Hemophagocytic lymphohistiocytosis

81 HSCT: Hematopoietic stem cell transplantation

82 IgRT: Immunoglobulin replacement therapy

83 PID: Primary Immunodeficiencies

84 WAS: Wiskott-Aldrich syndrome

85

86 **Acknowledgement**

87 We thank Dr. Jean-Philippe Jais and Dr. Olivier Bouaziz for the biostatistical support.

88 CEREDIH uses the European Society for Immunodeficiencies (ESID) registry platform to  
89 collect patient data.

90 CEREDIH receives unrestricted grants from the following pharmaceutical companies: LFB  
91 Biomédicaments, Takeda, Grifols, CSL Behring, Binding Site, Octapharma and from the  
92 following patients' associations: IRIS and AT Europe.

93

94 **Capsule summary (35 words)**

95 Non-infectious manifestations, i.e. allergy, autoimmunity/inflammation, lymphoproliferation  
96 and malignancies are frequent and severe manifestations in various PID categories across all  
97 age groups, emphasizing the importance of care by a multidisciplinary team of healthcare  
98 professionals for every single patient.

99

100 **Clinical Implications (29 words)**

101 Non-infectious manifestations, i.e. allergy, autoimmunity/inflammation, lymphoproliferation  
102 and malignancies in PID are more frequent than previously thought and should be assessed  
103 carefully throughout the life course of all PID patients.

104

105



106 **Abstract**

107 **Background:** Non-infectious manifestations, i.e. allergy, autoimmunity/inflammation,  
108 lymphoproliferation and malignancies are known to be observed in many primary  
109 immunodeficiency diseases (PID) and to participate to their prognosis.

110 **Objective:** In order to have a global view on their occurrence, we retrieved data from a  
111 retrospective cohort of 1375 patients included in the French national registry of PID (CEREDIH)  
112 for whom we had a 10-year follow-up since inclusion in the registry.

113 **Methods:** These patients were followed for 10 years (2009-2018) by specialized centers in  
114 University Hospitals. This study shows that 20.1% of patients without prior curative therapy  
115 (n=1163) developed at least one manifestation (event) encompassing 277 events.

116 **Results:** Autoimmune/inflammatory events (n=138) and malignancies (n=85) affected all age  
117 classes and virtually all PID diagnostic groups. They were associated with a risk of death that  
118 occurred in 14.2% of them (n=195), being found as causal in 43% of cases. Malignancies (OR:  
119 5.62 [3.66 – 8.62]) and autoimmunity (OR: 1.9 [1.27 – 2.84]) were clearly identified as risk  
120 factors for lethality. Patients who underwent curative therapy (i.e. mostly allogeneic  
121 hematopoietic stem cell transplantation, a few cases of gene therapy or thymic  
122 transplantation) prior to the 10-year study period (n=212) had comparatively reduced but still  
123 detectable clinical manifestations (n=16) leading to death in 9.4% of them.

124 **Conclusion:** This study points to the frequency and severity of non-infectious manifestations  
125 in various PID groups across all age groups. These results warrant further prospective analysis  
126 to better assess their consequences and to adapt therapy, notably indication of curative  
127 therapy.

128

129 **Introduction**

130 Primary immunodeficiencies (PID) represent a group of over 430 inherited diseases, with an  
131 estimated incidence of 1:3000 per year (1,2). Most are life-threatening diseases because these  
132 many conditions cause vulnerability to severe and recurrent infections but also because they  
133 are associated with immunopathology and cancer (3,4). This is justifying the efforts to collect  
134 as much as possible information on the mechanisms and pathophysiology of these various  
135 PIDs. Dedicated clinical centers caring from such complex cases have accordingly been  
136 developed all over the world.

137 In France, a network of pediatric and adult clinical units along with laboratories in every  
138 university medical center dedicated to the management of patients with PID (CEREDIH) has  
139 been created in 2005. A national registry documenting information on patients with PID has  
140 been created since then (5). National guidelines for care of both pediatric and adult patients  
141 with PID have been developed and updated, while regular videoconferences ensure  
142 harmonization on procedures and therapeutic indications at the national level (6). On this  
143 background, and given the paucity of overall information on the occurrence of non-infectious  
144 manifestations in PID patients over time, we decided to analyze data from a retrospective  
145 cohort of 1375 unselected patients with PID followed over a 10-year period (i.e. 2009 to 2018)  
146 in France. The analysis shows that non-infectious complications, i.e.  
147 allergy/autoimmunity/inflammation and cancer occur with a relatively high frequency across  
148 all ages and PID categories. These results call for further perspective assessment of their  
149 occurrence in various PIDs and potential therapeutic actions.

150

151

152 **Methods**

153

154 Since December 2005, the French National Reference Center for PID (CEREDIH) runs a national  
155 registry for documenting prevalent, incident as well as deceased patients diagnosed with a  
156 PID with the exception of severe congenital neutropenia that is covered by an independent  
157 registry (see Ref 5 for detailed methodology) (5). Relevant information is collected from  
158 patients' electronic charts on local visits to University Hospital covering all France by full-time  
159 dedicated and trained data entry clerks and then documented in the European Society for  
160 Immunodeficiencies (ESID) registry (7). Follow-up documentations are performed on a regular  
161 basis, once every 2 years on average.

162 The current study included patients recorded in the registry before 2009 January 1<sup>st</sup>  
163 and alive by this date and living in France with a follow up to 2018 December 31<sup>st</sup>. Lost to  
164 follow-up patients, defined as patients with no news after 2015 December 31<sup>st</sup>, were excluded  
165 (**Suppl. Fig1**). Information including patient demographics, way to diagnosis, clinical diagnosis,  
166 genetic diagnosis, occurrence of major clinical events such as allergy,  
167 autoimmunity/inflammation or malignancies, therapies (Immunoglobulin replacement  
168 therapy (IgRT) or curative such as Hematopoietic stem cell transplantation (HSCT), thymic  
169 transplantation or gene therapy) and causes of deaths were retrieved from the database.  
170 Infectious events were not analyzed in detail in this study.

171 Patients were classified in broad PID groups including (i) B cell deficiencies (common  
172 variable immunodeficiency CVID and non CVID), (ii) T cell deficiencies including severe  
173 combined immunodeficiencies (SCID), combined immunodeficiencies (CID), Wiskott-Aldrich  
174 Syndrome (WAS), Hyper-IgE syndrome (HIES), Ataxia-Telangiectasia (AT), (iii) Hemophagocytic  
175 lymphohistiocytosis (HLH) or other PID with predominant Immune dysregulation features and  
176 (iv) Innate immune deficiencies (defined as all PID affecting the innate immune system except  
177 chronic granulomatous disease (CGD) which is specifically described). Of note, patients could  
178 have multiple causes concurring to death. Ages at events are ages at first occurrence of events  
179 in the defined period of time. The national registry database (7,973 patients) was exported on  
180 March 9<sup>th</sup>, 2020, for data management and quality control. All statistical analyses were  
181 performed on R software version 3.6.1 (R Core Team (2018)). Categorical variables and  
182 continuous variables were compared by Chi2 test or Mann-Whitney test, respectively. Risk

183 factors of death were analyzed using logistic regression method. Ninety-five percent  
184 confidence intervals were calculated using the profile-likelihood approach.

185 From a population of 1,577 patients alive on January 1<sup>st</sup>, 2009, 202 were excluded for  
186 lack of completion of follow up information. No striking difference in terms of age and  
187 diagnosis was noticed between this group and the other cases (***Suppl Figure 1 and Suppl Table***  
188 ***1***). Overall, a population of 1375 patients met the inclusion criteria. They include 1163 patients  
189 who did not undergo a curative therapy (encompassing HSCT, gene therapy and thymic  
190 transplantation) and 212 who underwent a curative therapy prior to the study starting point  
191 (January 1<sup>st</sup>, 2009). These populations were accordingly analyzed separately. Global  
192 information on these cohorts is given in ***Table 1 and Supplementary Table 2, Supplementary***  
193 ***Table 3a and 3b***, including age, diagnosis, frequency at which these patients underwent  
194 relevant events for the study (i.e., cancer, autoimmunity/inflammation, allergy events) prior  
195 to January 1<sup>st</sup>, 2009.

196

197 **Results**

198 As shown in **Figures 1a-c**, and **Table 2**, 37% of patients with no previous curative  
199 therapy from the cohort experienced before 2009 an "event", i.e. allergy, autoimmunity,  
200 inflammation or cancer during their life time. In the 10-year period study, 20.1% patients  
201 experienced a non-infectious event (**Table 2**). These data illustrate how frequent are such non-  
202 infectious manifestations in PID patients. Together 16.1% of patients who did not have any  
203 event prior to 2009 (n=97) developed such a complication within the 10-year period (**Figure**  
204 **1b**). These events have occurred at all ages with a relatively similar frequency (**Figure 1c**). All  
205 PID groups are affected although frequency varies somewhat between them (**Figure 1d**). In  
206 total, 277 events occurred in 223 patients within the 10-year period, leading to a risk of 0.024  
207 event/year/patient (**Figure 1e-g**). A summary of events that occurred during the 10-year  
208 period is provided in **Table 2** and a detailed account as a function of PID diagnosis is depicted  
209 in **Suppl Table 4**.

210 A total of 195 patients (14.2%) died within this 10-year period including 175 of patients  
211 without a curative therapy and 20 (9.4%) of patients who previously received a curative  
212 therapy. Kinetics of death events of patients with no previous curative therapy is shown in  
213 **Figure 2**. Death rate was unsurprisingly higher in older patients but was quite high (9 up to  
214 16.8%) over all age groups (**Figure 2b**). Risk of death was particularly high in the group of  
215 patients with AT because of the very high incidence of cancer (see below). However, fatal  
216 events occurred in all groups of patients (**Figure 2c**). Age and underlying cause of deaths as a  
217 function of PID group are depicted in **Table 3**. Of note, non-infectious events accounted for  
218 43% of deaths with an identified cause. Malignancy accounted to 22.4% of cases (34.2% of  
219 deaths with an identified cause), which is a considerable frequency. Besides AT, cancer  
220 occurred in 11 CVID patients out of 329 (3%) and 5 CID patients out of 193 (2.5%).

221 Fourteen percent of patients from the cohort had allergic events prior to 2009  
222 confirming the known high frequency of allergy in PID patients. This was incremented by  
223 another 2% of patients who developed allergy within the 10-year study period (**Figure 3 and**  
224 **Suppl figure 2**). Onset of allergic events spanned all age classes showing that these might  
225 occur during all adulthood (**Figure 3b**). Distribution as a function of PID diagnosis was broad,  
226 although known PID groups prone to allergies (HIES, WAS, CID) were slightly overrepresented  
227 together with CGD (**Suppl table 2**). The different categories of allergic events and their  
228 frequency are indicated in **Suppl Table 5**.

229 Autoimmunity and inflammation are well-established features of PID. This is illustrated  
230 by the fact that 22.8% patients had such manifestations prior to 2009 (**Figure 4**). Strikingly,  
231 another 6.5% of patients developed these manifestations (**Suppl figure 3**) over the 10-year  
232 follow up. Altogether 135 events were noted in this 10-year period. Age distribution shows an  
233 apparent peak in the 11-30 years age category with a decline thereafter. Patients from all PID  
234 groups have developed such manifestations, with a notable peak of inflammation events,  
235 mostly digestive in CGD patients illustrating how frequent these complications are. Categories  
236 and frequency of autoimmune/inflammatory manifestations are depicted in **Suppl Table 6**. As  
237 shown in **Suppl Figure 4a,b**, occurrence of autoimmune/inflammatory manifestations either  
238 before 2009 or between 2009-2018 was a death risk factor ( $p=0.00045$ ).

239 This cohort study clearly shows the high risk of cancer in patients with PID since 6.2%  
240 (85 patients) have developed a cancer during the 10-year study period (**Figure 5, Suppl figure**  
241 **5**). Detailed list of malignancies is given in **Suppl Table 7**. There was an equal risk of  
242 hematological malignancies and solid tumors. Interestingly enough, the risk of the former was  
243 equally distributed overall all ages while the risk of solid tumors sharply increased with age  
244 (**Figure 5b**). AT patients were, as expected, mostly affected by the occurrence of cancer  
245 (**Figure 5c**). Nevertheless, virtually all PID groups were at risk for hematological malignancies  
246 (mostly lymphomas) with the notable exception of patients with CGD. Solid tumors occurred  
247 in patients with CVID, but also T-cell immunodeficiencies (CID). Not surprisingly, cancer was  
248 associated with a higher death toll (**Suppl figure 4c,d**).

249 Malignancy (OR: 5.62 [3.66 – 8.62],  $p<0.001$ ) and autoimmunity (OR: 1.90 [1.27 – 2.84],  
250  $p=0.002$ ) were found to be significantly associated with an increased risk of death. Close to  
251 40% of patients with malignancy died within the 10-year period (**Suppl figure 4c**). No  
252 significant association has been observed between allergy and risk of death (OR: 0.85 [0.50 –  
253 1.40],  $p=0.536$ ).

254

### 255 **Curative therapy**

256 When analyzing occurrence of events in patients who underwent a curative treatment  
257 of PID before 2009 ( $n=212$ , out of whom 180 patients with documentation, see **Suppl. Table**  
258 **8**), an event occurred in 8.9% of them ( $n=16$ ) including cancer ( $n=7$ ), autoimmunity ( $n=9$ ) and  
259 allergy ( $n=2$ ) (**Table 2**). Overall, 20 patients died, causes being shown in **Suppl Table 8, Suppl**  
260 **Fig 3 and Suppl Fig 6**.

261            In part, because of occurrence of these serious complications and as a function of PID  
262 diagnosis 71 patients (5.6%) underwent HSCT in this 10-year period (**Table 2**). PID categories  
263 are as follows: CVID: 1, CID: 26, WAS: 9, immune dysregulation: 1, HLH: 9, SCID: 6 and innate  
264 deficiencies: 19 out of whom 18 CGD. Median age at HSCT was 15.6 years (Q1;Q3: 8.4;23.3).  
265 Median age at last news was 17.6 years (Q1;Q3: 13.7;27.8). Median post HSCT follow-up is 2.3  
266 years (Q1;Q3: 0.7;6.2). Eight patients (11.3%) died after HSCT.  
267  
268

269 **Discussion**

270 In this study, we report on the occurrence of non-infectious complications in a large,  
271 unselected cohort of patients with PID who were under care in France for a continuous 10  
272 year-period. To our knowledge, this is the first study of this kind. It was made possible by the  
273 continuous and updated documentation in a national registry of patients with PID who  
274 underwent homogenous care and management including diagnostic methods, treatment of  
275 complications as well as indication and implementation of curative therapy based on  
276 indication on guidelines (6,8–10) and regular multidisciplinary meetings including transition  
277 care (11–13).

278 There are several limitations to such a study since a small fraction of patients were lost  
279 to follow up or information could not be retrieved over the 10-year period. Nevertheless, it  
280 appears that the study population did not differ extensively from the overall PID population  
281 of the national registry, thus making unlikely a selection bias. This study focuses on non-  
282 infectious complications named as events in this manuscript, and their impact on survival. It  
283 would be of obvious interest to perform similar studies encompassing all patients' medical  
284 history including infectious events and also including quality of life assessments – a key issue  
285 for patients with PID (14,15). Based on the CEREDIH registry, there is an ongoing collection of  
286 these pieces of information to be later analyzed.

287 Nevertheless, as it stands, this study provides information on the relative frequency of  
288 non-infectious events, as a function of age and PID diagnosis. It is an extension to previous  
289 reports (16–20) including ones on the advent of allergy (21), autoimmune and inflammatory  
290 disorders (3,22,23) and cancer across various PID (24–27). All age classes are affected albeit  
291 distinctly and almost none of the PID diseases are spared for the advent of such complications  
292 all along life span. It is translated into a significant risk of deaths that should definitively be  
293 considered for therapeutic decision. The literature is rich in reports about the occurrence of  
294 major non-infectious events and risk of deaths in patients with CVID (16,17,25,26,28,29).  
295 Although there is a heterogeneity of PID category distribution of patients with  
296 autoimmunity/inflammation across age classes, the present report nevertheless shows that  
297 many more PID during childhood and adulthood carry a risk of immunopathological  
298 comorbidities as a consequence of their disease. Of course, this type of study cannot replace  
299 disease-specific assessment of clinical outcome as exemplified (13,30,31), that can much  
300 better detect the specific features of the many known and less well known PID. However, this



301 study provides an estimation of the global burden of such complications in the setting of  
302 specialized care of PID and the payload in terms of fatal risk.

303 Not surprisingly, cancer appears as a key prognostic factor for death as previously  
304 reported for CVID (16,24,25,27). Most patients experienced one malignancy. Malignancies  
305 occurred at all ages and in all PID categories. Median age at first malignancy is 35.3 years. PID  
306 categories with the highest frequency of malignancies were B cell deficiencies (both CVID and  
307 nonCVID) and AT. Hematological malignancies were prominent, with a higher frequency of  
308 Non-Hodgkin lymphoma, followed by Hodgkin lymphomas, leukemias, myelodysplastic  
309 syndromes and plasma cell disorders. Solid tumors were mostly skin cancers, digestive tract  
310 cancers, breast cancers, thymoma, genital organ cancers and urinary tract cancers. These  
311 findings call for a prospective surveillance of relevant patients for occurrence of such  
312 malignancies. Mechanisms underlying cancer risk in PID are numerous, likely accounting for  
313 the age distribution and type of cancer. They include defective DNA repair of pro-oncogenic  
314 somatic mutations (AT, NBS...), poor handling of oncogenic viruses (i.e. EBV, HHV8, and  
315 possibly HPVs) mostly in patients with T-cell immunodeficiencies, defective control of cell  
316 death (ALPS...), repeated and/or protracted immune activation by micro-organisms leading to  
317 the generation of DNA errors along cycles of cell division (potentially all PID causing  
318 vulnerability to infections). Immunosenescence on top of immunodeficiency itself may  
319 contribute as well as accumulation of somatic mutation in various tissues (24). Data indicate  
320 that type of cancer, i.e. hematological malignancies, mostly lymphomas and solid tumor, occur  
321 at different ages as a function of underlying disease and likely pathophysiological mechanism,  
322 but in any case with a frequency much higher than in the study cohort compared to the whole  
323 registry ( $p=0.003$ , See **Suppl. Table 1**).

324 The study also provides a picture of the related proportion at a nationwide level of  
325 patients who underwent a curative therapy (mostly HSCT) in this 10-year period, i.e., in 5.6%  
326 of the cohort. As expected, HSCT was performed in classical PID categories i.e., T-cell  
327 immunodeficiencies, HLH and CGD and was mostly performed in children although some adult  
328 patients were also transplanted (manuscript in preparation) (32,33).

329 Another interesting piece of data retrieved from this study is the follow-up of patients  
330 who underwent curative therapy (mostly HSCT) prior to the study period. Amongst the 212  
331 patients followed for 10 years, a fraction 8.9% (versus 20.1% in the cohort of patients who did  
332 not receive curative therapy before 2009) still had some significant events that were not all

333 consequences of recent transplantation. This shows that late complication of HSCT, persisting  
334 symptoms of PID, sequelae of pre HSCT manifestations or syndromic features of complex PID  
335 can contribute significantly to post therapeutic morbidity and mortality (34,35). Such an  
336 observation is an incentive to plan a prospective study carefully assessing the long-term  
337 outcome of HSCT for PID as a function of all these parameters. Indeed, we propose that  
338 patients with PID developing autoimmunity should be considered for curative therapy as very  
339 significant advances in these procedures have been made in the past decades, including for  
340 teenagers and adults with CGD, CID or WAS, amongst other PID categories. For patients with  
341 PIDs such as CTLA4, LRBA, APDS1/2, targeted therapies now play a very important role in the  
342 management of these patients (32). Hence, performing a genetic documentation (using NGS  
343 techniques, including panels) is of utmost importance in providing the best level of care for  
344 patients with PID (8).

345         The aim of our manuscript was to provide a descriptive observational analysis of the  
346 major comorbidities occurring in a large cohort of patients with PID followed up in a national  
347 reference network. This study reflects on the severity and complexity of care of PID patients.  
348 It calls a) to emphasize the need that every single PID patient benefits from the care from a  
349 multidisciplinary team of experienced healthcare professionals and b) to design prospective  
350 refined studies analyzing as a function of time the PID patients' history under optimal care,  
351 including comparison with control groups.

352

353 **References**

- 354 1. Tangye SG, Al-Herz W, Bousfiha A, Cunningham-Rundles C, Franco JL, Holland SM, et  
355 al. The Ever-Increasing Array of Novel Inborn Errors of Immunity: an Interim Update  
356 by the IUIS Committee. *J Clin Immunol* [Internet]. 2021 Apr 1 [cited 2021 Jun  
357 11];41(3):666–79. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/33598806/)  
358 [gov.proxy.insermbiblio.inist.fr/33598806/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/33598806/)
- 359 2. Tangye SG, Al-Herz W, Bousfiha A, Chatila T, Cunningham-Rundles C, Etzioni A, et al.  
360 Human Inborn Errors of Immunity: 2019 Update on the Classification from the  
361 International Union of Immunological Societies Expert Committee. *J Clin Immunol*.  
362 2020 Jan 1;40(1):24–64.
- 363 3. Fischer A, Provot J, Jais J-P, Alcais A, Mahlaoui N, Adoue D, et al. Autoimmune and  
364 inflammatory manifestations occur frequently in patients with primary  
365 immunodeficiencies. *J Allergy Clin Immunol* [Internet]. 2017 Nov [cited 2017 Apr  
366 9];140(5):1388-1393.e8. Available from:  
367 <http://www.sciencedirect.com.gate2.inist.fr/science/article/pii/S0091674917302269>
- 368 4. Thalhammer J, Kindle G, Nieters A, Rusch S, Seppänen MRJ, Fischer A, et al. Initial  
369 presenting manifestations in 16,486 patients with Inborn Errors of Immunity include  
370 infections and non-infectious manifestations. *J Allergy Clin Immunol* [Internet]. 2021  
371 Apr 23 [cited 2021 May 6]; Available from:  
372 <https://linkinghub.elsevier.com/retrieve/pii/S0091674921006540>
- 373 5. CEREDIH. The French PID Study Group. The French national registry of primary  
374 immunodeficiency diseases. *Clin Immunol* [Internet]. 2010 May [cited 2014 Aug  
375 25];135(2):264–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20399414>
- 376 6. Aguilar C, Malphettes M, Donadieu J, Chandesris O, Coignard-Biehler H, Catherinot E,  
377 et al. Prevention of infections during primary immunodeficiency. *Clin Infect Dis*. 2014;
- 378 7. Scheible R, Rusch S, Guzman D, Mahlaoui N, Ehl S, Kindle G. The NEW ESID online  
379 database network. *Bioinformatics* [Internet]. 2019 Dec 15 [cited 2021 Jun  
380 11];35(24):5367–9. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31263866/)  
381 [gov.proxy.insermbiblio.inist.fr/31263866/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31263866/)
- 382 8. Fusaro M, Rosain J, Grandin V, Lambert N, Hanein S, Fourrage C, et al. Improving the  
383 diagnostic efficiency of primary immunodeficiencies with targeted next-generation  
384 sequencing. *J Allergy Clin Immunol* [Internet]. 2021 Feb 1 [cited 2021 Jun

- 385 11];147(2):734–7. Available from: <https://pubmed-ncbi-nlm-nih->  
386 [gov.proxy.insermbiblio.inist.fr/32531373/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/32531373/)
- 387 9. Goussault H, Salvator H, Catherinot E, Chabi ML, Tcherakian C, Chabrol A, et al.  
388 Primary immunodeficiency-related bronchiectasis in adults: Comparison with  
389 bronchiectasis of other etiologies in a French reference center. *Respir Res* [Internet].  
390 2019 Dec 4 [cited 2021 Jun 11];20(1). Available from: <https://pubmed-ncbi-nlm-nih->  
391 [gov.proxy.insermbiblio.inist.fr/31801528/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31801528/)
- 392 10. Seidel MG, Kindle G, Gathmann B, Quinti I, Buckland M, van Montfrans J, et al. The  
393 European Society for Immunodeficiencies (ESID) Registry Working Definitions for the  
394 Clinical Diagnosis of Inborn Errors of Immunity. *J Allergy Clin Immunol Pract*. 2019;
- 395 11. Salvator H, Mahlaoui N, Catherinot E, Rivaud E, Pilmis B, Borie R, et al. Pulmonary  
396 manifestations in adult patients with chronic granulomatous disease. *Eur Respir J*.  
397 2015;
- 398 12. Dunogué B, Pilmis B, Mahlaoui N, Elie C, Coignard-Biehler H, Amazzough K, et al.  
399 Chronic Granulomatous Disease in Patients Reaching Adulthood: A Nationwide Study  
400 in France. *Clin Infect Dis*. 2017;64(6).
- 401 13. Chandesris M-O, Melki I, Natividad A, Puel A, Fieschi C, Yun L, et al. Autosomal  
402 dominant STAT3 deficiency and hyper-IgE syndrome: molecular, cellular, and clinical  
403 features from a French national survey. *Medicine (Baltimore)* [Internet]. 2012 Jul  
404 [cited 2013 Dec 28];91(4):e1-19. Available from:  
405 <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3680355&tool=pmcentr>  
406 [ez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3680355&tool=pmcentr&rendertype=abstract)
- 407 14. Barlogis V, Mahlaoui N, Auquier P, Pellier I, Fouyssac F, Vercasson C, et al. Physical  
408 health conditions and quality of life in adults with primary immunodeficiency  
409 diagnosed during childhood: A French Reference Center for PIDs (CEREDIH) study. *J*  
410 *Allergy Clin Immunol*. 2017;139(4).
- 411 15. Barlogis V, Mahlaoui N, Auquier P, Fouyssac F, Pellier I, Vercasson C, et al. Burden of  
412 Poor Health Conditions and Quality of Life in 656 Children with Primary  
413 Immunodeficiency. *Journal of Pediatrics*. 2017;
- 414 16. Resnick ES, Moshier EL, Godbold JH, Cunningham-Rundles C. Morbidity and mortality  
415 in common variable immune deficiency over 4 decades. *Blood* [Internet]. 2012 Feb 16  
416 [cited 2021 Jun 11];119(7):1650–7. Available from: <https://pubmed-ncbi-nlm-nih->

- 417 gov.proxy.insermbiblio.inist.fr/22180439/
- 418 17. Farmer JR, Ong M-S, Barmettler S, Yonker LM, Fuleihan R, Sullivan KE, et al. Common  
419 Variable Immunodeficiency Non-Infectious Disease Endotypes Redefined Using  
420 Unbiased Network Clustering in Large Electronic Datasets. *Front Immunol* [Internet].  
421 2017 Jan 9 [cited 2018 Sep 30];8:1740. Available from:  
422 <http://journal.frontiersin.org/article/10.3389/fimmu.2017.01740/full>
- 423 18. Gathmann B, Mahlaoui N, Gérard L, Oksenhendler E, Warnatz K, Schulze I, et al.  
424 Clinical picture and treatment of 2212 patients with common variable  
425 immunodeficiency. *J Allergy Clin Immunol*. 2014;134(1):116-126.e11.
- 426 19. Odnoletkova I, Kindle G, Quinti I, Grimbacher B, Knerr V, Gathmann B, et al. The  
427 burden of common variable immunodeficiency disorders: a retrospective analysis of  
428 the European Society for Immunodeficiency (ESID) registry data. *Orphanet J Rare Dis*.  
429 2018;13(1).
- 430 20. Coignard-Biehler H, Mahlaoui N, Pilmis B, Barlogis V, Brosselin P, De Vergnes N, et al.  
431 A 1-Year Prospective French Nationwide Study of Emergency Hospital Admissions in  
432 Children and Adults with Primary Immunodeficiency. *J Clin Immunol* [Internet]. 2019  
433 Oct 1 [cited 2021 Jun 11];39(7):702–12. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31401750/)  
434 [gov.proxy.insermbiblio.inist.fr/31401750/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31401750/)
- 435 21. Tuano KS, Orange JS, Sullivan K, Cunningham-Rundles C, Bonilla FA, Davis CM. Food  
436 allergy in patients with primary immunodeficiency diseases: prevalence within the US  
437 Immunodeficiency Network (USIDNET). *J Allergy Clin Immunol* [Internet]. 2015 Jan  
438 [cited 2015 May 3];135(1):273–5. Available from:  
439 <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4324505&tool=pmcentr>  
440 [ez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4324505&tool=pmcentr)
- 441 22. Leung J, Sullivan KE, Perelygina L, Icenogle JP, Fuleihan RL, Lanzieri TM. Prevalence of  
442 Granulomas in Patients With Primary Immunodeficiency Disorders, United States:  
443 Data From National Health Care Claims and the US Immunodeficiency Network  
444 Registry. *J Clin Immunol* [Internet]. 2018 Aug 24 [cited 2018 Sep 23];38(6):717–26.  
445 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30043271>
- 446 23. Feuille EJ, Anooshiravani N, Sullivan KE, Fuleihan RL, Cunningham-Rundles C.  
447 Autoimmune Cytopenias and Associated Conditions in CVID: a Report From the  
448 USIDNET Registry. *J Clin Immunol* [Internet]. 2018 Jan 1 [cited 2021 Jun 11];38(1):28–

- 449 34. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/29080979/)  
450 [gov.proxy.insermbiblio.inist.fr/29080979/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/29080979/)
- 451 24. Mayor PC, Eng KH, Singel KL, Abrams SI, Odunsi K, Moysich KB, et al. Cancer in primary  
452 immunodeficiency diseases: Cancer incidence in the United States Immune Deficiency  
453 Network Registry. *J Allergy Clin Immunol* [Internet]. 2018 Mar [cited 2018 Sep  
454 30];141(3):1028–35. Available from:  
455 <https://linkinghub.elsevier.com/retrieve/pii/S0091674917309259>
- 456 25. Quinti I, Agostini C, Tabolli S, Brunetti G, Cinetto F, Pecoraro A, et al. Malignancies are  
457 the major cause of death in patients with adult onset common variable  
458 immunodeficiency. *Vol. 120, Blood*. 2012. p. 1953–4.
- 459 26. Yakaboski E, Fuleihan RL, Sullivan KE, Cunningham-Rundles C, Feuille E.  
460 Lymphoproliferative Disease in CVID: a Report of Types and Frequencies from a US  
461 Patient Registry. *J Clin Immunol* [Internet]. 2020 Apr 1 [cited 2021 Jun 11];40(3):524–  
462 30. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/32185577/)  
463 [gov.proxy.insermbiblio.inist.fr/32185577/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/32185577/)
- 464 27. Cheminant M, Mahlaoui N, Desconclois C, Canioni D, Ysebaert L, Dupré L, et al.  
465 Lymphoproliferative disease in patients with Wiskott-Aldrich syndrome: Analysis of  
466 the French Registry of Primary Immunodeficiencies. *J Allergy Clin Immunol* [Internet].  
467 2019 Jun 1 [cited 2021 Jun 12];143(6):2311-2315.e7. Available from: [https://pubmed-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/30796981/)  
468 [ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/30796981/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/30796981/)
- 469 28. Pensieri MV, Pulvirenti F, Schiepati A, Maimaris S, Lattanzio S, Quinti I, et al. The high  
470 mortality of patients with common variable immunodeficiency and small bowel villous  
471 atrophy. *Scand J Gastroenterol* [Internet]. 2019 Feb 1 [cited 2021 Jun 12];54(2):164–8.  
472 Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31006294/)  
473 [gov.proxy.insermbiblio.inist.fr/31006294/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31006294/)
- 474 29. Pulvirenti F, Pecoraro A, Cinetto F, Milito C, Valente M, Santangeli E, et al. Gastric  
475 Cancer Is the Leading Cause of Death in Italian Adult Patients With Common Variable  
476 Immunodeficiency. *Front Immunol* [Internet]. 2018 Nov 5 [cited 2018 Nov 21];9:2546.  
477 Available from: <https://www.frontiersin.org/article/10.3389/fimmu.2018.02546/full>
- 478 30. Conley ME, Park CL, Douglas SD. Childhood common variable immunodeficiency with  
479 autoimmune disease. *J Pediatr* [Internet]. 1986 [cited 2021 Jun 12];108(6):915–22.  
480 Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/31006294/)

- 481 gov.proxy.insermbiblio.inist.fr/2423668/
- 482 31. Schubert D, Bode C, Kenefeck R, Hou TZ, Wing JB, Kennedy A, et al. Autosomal  
483 dominant immune dysregulation syndrome in humans with CTLA4 mutations. *Nat*  
484 *Med.* 2014;20(12).
- 485 32. Fox TA, Chakraverty R, Burns S, Carpenter B, Thomson K, Lowe D, et al. Successful  
486 outcome following allogeneic hematopoietic stem cell transplantation in adults with  
487 primary immunodeficiency. *Blood* [Internet]. 2018 Feb 22 [cited 2018 Sep  
488 29];131(8):917–31. Available from:  
489 <http://www.bloodjournal.org/lookup/doi/10.1182/blood-2017-09-807487>
- 490 33. Chiesa R, Wang J, Blok HJ, Hazelaar S, Neven B, Moshous D, et al. Hematopoietic cell  
491 transplantation in chronic granulomatous disease: a study of 712 children and adults.  
492 *Blood* [Internet]. 2020 Sep 3 [cited 2021 Jun 12];136(10):1201–11. Available from:  
493 <https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/32614953/>
- 494 34. Neven B, Leroy S, Decaluwe H, Le Deist F, Picard C, Moshous D, et al. Long-term  
495 outcome after hematopoietic stem cell transplantation of a single-center cohort of 90  
496 patients with severe combined immunodeficiency. *Blood* [Internet]. 2009 [cited 2021  
497 Jun 11];113(17):4114–24. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/19168787/)  
498 [gov.proxy.insermbiblio.inist.fr/19168787/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/19168787/)
- 499 35. Heimall J, Puck J, Buckley R, Fleisher TA, Gennery AR, Neven B, et al. Current  
500 Knowledge and Priorities for Future Research in Late Effects after Hematopoietic Stem  
501 Cell Transplantation (HCT) for Severe Combined Immunodeficiency Patients: A  
502 Consensus Statement from the Second Pediatric Blood and Marrow Transplant  
503 Consortium International Conference on Late Effects after Pediatric HCT. In: *Biology of*  
504 *Blood and Marrow Transplantation* [Internet]. Elsevier Inc.; 2017 [cited 2021 Jun 12].  
505 p. 379–87. Available from: [https://pubmed-ncbi-nlm-nih-](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/28068510/)  
506 [gov.proxy.insermbiblio.inist.fr/28068510/](https://pubmed-ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/28068510/)
- 507
- 508
- 509

510 **List of figures**

511 **Figure 1: Frequency at which patients developed events during the 10-year follow-up period**  
512 **and frequency of events**

- 513 a. Cumulative percentage of patients who experienced at least one event before 2009  
514 and since January, 1st 2009 (n=1163)
- 515 b. Cumulative percentage of patients who experienced a first event during the study  
516 period (2009-2018) (Percentages are calculated on the 603 patients who never  
517 experienced any event before 2009)
- 518 c. Fraction of patients who experienced at least one event during the study period (2009-  
519 2018) according to age class in 2009 (n=1163)
- 520 d. Fraction of patients who experienced at least one event during the study period (2009-  
521 2018) by PID category (n=1163)
- 522 e. Cumulative number of events during the study period (2009-2018)
- 523 f. Number of events that occurred during the study period (2009-2018) according to age  
524 class in 2009
- 525 g. Number of events that occurred during the study period (2009-2018) by PID category

526

527 **Abbreviations:**

528 AT: ataxia telangiectasia; CGD: chronic granulomatous disease; CID: combined  
529 immunodeficiency; CVID: common variable immunodeficiency; HIES: hyper immunoglobulin  
530 E syndrome; HLH: hemophagocytic lymphohistiocytosis; ImDysReg: immune dysregulation;  
531 SCID: severe combined immunodeficiency; WAS: Wiskott Aldrich syndrome

532

533 **Figure 2: Death events**

- 534 a. Cumulative percentage of deaths (cohort, n=1163)
- 535 b. Fraction of deaths according to age class in 2009
- 536 c. Fraction of deaths by PID category

537

538 **Figure 3: Allergy events**

- 539 a. Cumulative percentage of patients who experienced at least one allergy event during  
540 the study period (2009-2018) among 1023 evaluable patients



541 b. Fraction of patients who experienced at least one allergy event during the study period  
542 (2009-2018) according to age class in 2009

543 c. Fraction of patients who experienced at least one allergy during the study period  
544 (2009-2018) event by PID category

545 Only pre-curative therapy events are considered.

546

547 **Figure 4: Autoimmunity/inflammation events**

548 a. Cumulative percentage of patients who experienced at least one  
549 autoimmunity/inflammation event during the study period (2009-2018) among 1025  
550 evaluable patients

551 b. Fraction of patients who experienced at least one autoimmunity/inflammation event  
552 during the study period (2009-2018) according to age class in 2009

553 c. Fraction of patients who experienced at least one autoimmunity/inflammation event  
554 during the study period (2009-2018) by PID category

555 Only pre-curative therapy events are considered.

556

557 **Figure 5: Malignancy events**

558 a. Cumulative percentage of patients who experienced at least one hematological  
559 malignancy and/or solid tumor during the study period (2009-2018) among 1151  
560 evaluable patients

561 b. Fraction of patients who experienced at least one hematological malignancy and/or  
562 solid tumor during the study period (2009-2018) according to age class in 2009

563 c. Fraction of patients who experienced at least one hematological malignancy and/or  
564 solid tumor during the study period (2009-2018) by PID category

565 Only pre-curative therapy events are considered.

566

567

568

569

570

571

572 **List of tables**

573

574 **Table 1: Patients characteristics at baseline, January, 1st 2009.**


575

576 **Table 2: Patients with events during the study period (2009-2018).**

577


578 **Table 3: Causes of deaths.**

a. At least one event




Years

b. Occurrence of a first event




Years

c. At least one event




Age class in 2009 (years)

d. At least one event




PID category

e. Cumulative number of events




Years

f. Number of events




Age class in 2009 (years)

g. Number of events




PID category


a. Cumulative percentage of deaths




b. Fraction of deaths




c. Fraction of deaths




a. Cumulative percentage of allergy




b. Fraction of allergy




c. Fraction of allergy




a. Cumulative percentage of AI/Infla




b. Fraction of AI/Infla




c. Fraction of AI/Infla



a. Cumulative percentage of malignancy



b. Fraction of malignancy



c. Fraction of malignancy




Table 1

Covariate	Values	Whole cohort	No curative therapy	Curative therapy	
		N=1375	N=1163	N=212	
Gender	F	516 (37.5%)	440 (37.8%)	76 (35.8%)	
	M	859 (62.5%)	723 (62.2%)	136 (64.2%)	
PID diagnosis	B CVID	329 (23.9%)	328 (28.2%)	1 (0.5%)	
	B nonCVID	321 (23.3%)	321 (27.6%)	0 (0%)	
	CID	193 (14%)	160 (13.8%)	33 (15.6%)	
	AT	96 (7%)	96 (8.3%)	0 (0%)	
	SCID	98 (7.1%)	5 (0.4%)	93 (43.9%)	
	WAS	70 (5.1%)	39 (3.4%)	31 (14.6%)	
	HIES	24 (1.7%)	24 (2.1%)	0 (0%)	
	HLH	63 (4.6%)	24 (2.1%)	39 (18.4%)	
	ImDysReg	41 (3%)	38 (3.3%)	3 (1.4%)	
	Innate non CGD	35 (2.5%)	32 (2.8%)	3 (1.4%)	
	CGD	105 (7.6%)	96 (8.3%)	9 (4.2%)	
	Age at last news		26.9 [18.8;42.3]	28.4 [19.4;47.2]	21.2 [15;28.2]
	Malignancy*	N	1271 (93.5%)	1072 (93.1%)	199 (95.7%)
Y		88 (6.5%)	79 (6.9%)	9 (4.3%)	
Hematol. malignancy*	N	1328 (96.6%)	1123 (96.6%)	205 (96.7%)	
	Y	47 (3.4%)	40 (3.4%)	7 (3.3%)	
Solid tumor*	N	1331 (96.8%)	1121 (96.4%)	210 (99.1%)	
	Y	44 (3.2%)	42 (3.6%)	2 (0.9%)	
Allergy	N	1008 (83.9%)	858 (83.9%)	150 (83.8%)	
	Y	194 (16.1%)	165 (16.1%)	29 (16.2%)	
Autoimmunity	N	915 (75.5%)	760 (74.1%)	155 (82.9%)	
	Y	297 (24.5%)	265 (25.9%)	32 (17.1%)	
All events	N	731 (60%)	603 (58.2%)	128 (69.9%)	
	Y	488 (40%)	433 (41.8%)	55 (30.1%)	
Ig replacement	N	528 (47.8%)	408 (43.1%)	120 (76.4%)	
	Y	576 (52.2%)	539 (56.9%)	37 (23.6%)	



Table 2

Covariate	Values	Whole cohort	No curative therapy (2009)	Curative therapy (2009)
		N=1375	N=1163	N=212
Living status	Alive	1180 (85.8%)	988 (85%)	192 (90.6%)
	Deceased	195 (14.2%)	175 (15%)	20 (9.4%)
Malignancy*	N	1260 (92.7%)	1058 (91.9%)	202 (97.1%)
	Y	99 (7.3%)	93 (8.1%)	6 (2.9%)
Hematol. Malignancy*	N	1312 (96.5%)	1106 (96.1%)	206 (99%)
	Y	47 (3.5%)	45 (3.9%)	2 (1%)
Solid tumor*	N	1303 (95.9%)	1100 (95.6%)	203 (97.6%)
	Y	56 (4.1%)	51 (4.4%)	5 (2.4%)
Allergy	N	1173 (97.6%)	996 (97.4%)	177 (98.9%)
	Y	29 (2.4%)	27 (2.6%)	2 (1.1%)
Autoimmunity	N	1096 (90.4%)	918 (89.6%)	178 (95.2%)
	Y	116 (9.6%)	107 (10.4%)	9 (4.8%)
All events**	N	987 (81.6%)	823 (79.9%)	164 (91.1%)
	Y	223 (18.4%)	207 (20.1%)	16 (8.9%)
Ig replacement	N	652 (91.8%)	602 (91.8%)	50 (92.6%)
	Y	58 (8.2%)	54 (8.2%)	4 (7.4%)
Route of Ig replacement***	IV	26 (55.3%)	25 (56.8%)	1 (33.3%)
	SC	21 (44.7%)	19 (43.2%)	2 (66.7%)
Curative therapy	N	1304 (94.8%)	1098 (94.4%)	206 (97.2%)
	Y	71 (5.2%)	65 (5.6%)	6 (2.8%)

Table 3

Covariate	Whole cohort	B CVID	B nonCVID	CID	AT	SCID	WAS	HIES	HLH	ImmDysReg	Innate nonCGD	CGD
	N=175	N=54	N=19	N=23	N=46	N=1	N=3	N=3	N=5	N=6	N=2	N=13
Age at death (y)	28.9 [18.6;57.6]	65 [50.2;74.2]	44.7 [20.7;72.5]	20.8 [14.6;26.6]	20.7 [16.4;26.8]	2.7 [2.7;2.7]	13 [12;19]	33.6 [31.3;34.6]	34 [23.1;35.1]	29.6 [23.5;48.1]	41.2 [33.7;48.7]	16.4 [14.8;31.2]
Infection	49 (28.2%)	11 (20.8%)	4 (21.1%)	9 (39.1%)	11 (23.9%)	1 (100%)	1 (33.3%)	2 (66.7%)	1 (20%)	0 (0%)	0 (0%)	9 (69.2%)
Immune dysr.	7 (4%)	2 (3.8%)	0 (0%)	2 (8.7%)	0 (0%)	0 (0%)	1 (33.3%)	0 (0%)	0 (0%)	1 (16.7%)	0 (0%)	1 (7.7%)
Malignancy	39 (22.4%)	11 (20.8%)	4 (21.1%)	5 (21.7%)	14 (30.4%)	0 (0%)	1 (33.3%)	0 (0%)	1 (20%)	1 (16.7%)	2 (100%)	0 (0%)
Transplantation related	14 (8%)	0 (0%)	0 (0%)	6 (26.1%)	0 (0%)	1 (100%)	0 (0%)	0 (0%)	3 (60%)	1 (16.7%)	0 (0%)	3 (23.1%)
Other	29 (16.8%)	9 (17.3%)	3 (15.8%)	4 (17.4%)	7 (15.2%)	0 (0%)	0 (0%)	2 (66.7%)	0 (0%)	1 (16.7%)	1 (50%)	2 (15.4%)
Unknown	61 (35.5%)	27 (50%)	9 (47.4%)	4 (19%)	17 (37%)	0 (0%)	0 (0%)	0 (0%)	1 (20%)	2 (40%)	0 (0%)	1 (7.7%)