



**HAL**  
open science

## **Debris-covered glacier systems and associated glacial lake outburst flood hazards : challenges and prospects**

Adina Racoviteanu, L. Nicholson, N.F. Glasser, E. Miles, S. Harrison, J.M. Reynolds

### ► **To cite this version:**

Adina Racoviteanu, L. Nicholson, N.F. Glasser, E. Miles, S. Harrison, et al.. Debris-covered glacier systems and associated glacial lake outburst flood hazards : challenges and prospects. *Journal of the Geological Society*, 2022, 179 (3), jgs2021-084 [21 p.]. <10.1144/jgs2021-084>. <hal-04010859>

**HAL Id: hal-04010859**

**<https://hal.science/hal-04010859v1>**

Submitted on 17 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

# Debris-covered glacier systems and associated glacial lake outburst flood hazards: challenges and prospects



A.E. Racoviteanu<sup>1,2\*</sup>, L. Nicholson<sup>3</sup>, N.F. Glasser<sup>2</sup>, Evan Miles<sup>4</sup>, S. Harrison<sup>1</sup> and J.M. Reynolds<sup>5</sup>


<sup>1</sup> Department of Geography, Exeter University, Penryn, Cornwall TR10 9FE, UK

<sup>2</sup> Department of Geography and Earth Sciences, Aberystwyth University, SY23 3DB, UK

<sup>3</sup> Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innrain 52, 6020 Innsbruck, Austria

<sup>4</sup> Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse, 111CH-8903 Birmensdorf, Switzerland

<sup>5</sup> Reynolds International Ltd, Wrexham Road, Mold, Flintshire CH7 1HP, UK

 AER, 0000-0003-4954-1871; LN, 0000-0003-0430-7950; NFG, 0000-0002-8245-2670; EM, 0000-0001-5446-8571; SH, 0000-0003-2820-7313; JMR, 0000-0003-4372-0510

\* Correspondence: [a.racoviteanu@exeter.ac.uk](mailto:a.racoviteanu@exeter.ac.uk)



**Abstract:** Glaciers respond sensitively to climate variability and change, with associated impacts on meltwater production, sea-level rise and geomorphological hazards. There is a strong societal interest in understanding the current response of all types of glacier systems to climate change and how they will continue to evolve in the context of the whole glacierized landscape. In particular, understanding the current and future behaviour of debris-covered glaciers is a ‘hot topic’ in glaciological research because of concerns for water resources and glacier-related hazards. The state of these glaciers is closely related to various hazardous geomorphological processes which are relatively poorly understood. Understanding the implications of debris-covered glacier evolution requires a systems approach. This includes the interplay of various factors such as local geomorphology, ice ablation patterns, debris characteristics and glacier lake growth and development. Such a broader, contextualized understanding is prerequisite to identifying and monitoring the geohazards and hydrologic implications associated with changes in the debris-covered glacier system under future climate scenarios. This paper presents a comprehensive review of current knowledge of the debris-covered glacier landsystem. Specifically, we review state-of-the-art field-based and the remote sensing-based methods for monitoring debris-covered glacier characteristics and lakes and their evolution under future climate change. We advocate a holistic process-based framework for assessing hazards associated with moraine-dammed glacio-terminal lakes that are a projected end-member state for many debris-covered glaciers under a warming climate.

Received 1 July 2021; revised 17 October 2021; accepted 1 November 2021

Glaciers respond sensitively to climate change and variability, with associated impacts on meltwater production (Kaser *et al.* 2010; Huss 2011; Immerzeel *et al.* 2012), sea-level rise (Berthier *et al.* 2010; Leclercq *et al.* 2011; Marzeion *et al.* 2012) and geomorphological hazards (Kääb *et al.* 2005; Benn *et al.* 2012; Harrison *et al.* 2018). Glacier behaviour has potential knock-on effects for valley-scale sediment fluxes, surface energy balance, water storage and geomorphological hazards. Therefore, there is a keen societal interest in understanding how the different types of glacier systems are currently responding to climate change and how they will evolve in the context of the whole landscape. In particular, the role of debris-covered glacier landsystems and associated lakes in water supply and related hazards is less well understood. Although they account for only *c.* 4 to 7% of the global glacierized area (Scherler *et al.* 2018; Herreid and Pellicciotti 2020), debris-covered glaciers are a prominent feature of high-relief orogenic belts where high denudation rates supply abundant rock debris to the glacier surface, often producing debris-covered glacier tongues up to tens of kilometres long such as Baltoro Glacier (62 km) in the Karakoram (Mihalcea *et al.* 2006) or Ngozumpa Glacier (*c.* 25 km) in the Nepal Himalaya (Casey and Kääb 2012) (Fig. 1). They are an especially well-developed feature in the Hindu Kush Himalayan region, where *c.* 13% of the total glacierized area is debris-covered, ranging from 9% in the Karakoram to 15% in the eastern Nepalese and Bhutanese Himalaya (Kääb *et al.* 2012). They are also found in the Tien Shan (Hagg *et al.* 2008), Caucasus (Stokes *et al.* 2007),

Alaska (Berthier *et al.* 2010), New Zealand (Anderson and Mackintosh 2012), parts of the Andes (Racoviteanu *et al.* 2008) and the Alps (Deline 2005), Greenland, and the Dry Valleys of Antarctica. Supraglacial debris varies in thickness from several centimetres up to 2 m or more (Benn and Evans 1998; Anderson and Anderson 2018).

Satellite-derived inventories show that glacier-wide ice mass loss from debris-covered glacier tongues over recent decades is substantial and increasing (Bolch *et al.* 2011; Kääb *et al.* 2012; Brun *et al.* 2019; Maurer *et al.* 2019; King *et al.* 2020b). As mountain glaciers continue to diminish in the coming decades, an increasing proportion of the remaining ice is expected to become debris-covered (Herreid and Pellicciotti 2020). This makes it critical to understand how debris cover impacts glacier meltwater production in order to make projections of regional water resources and global sea-level rise. Furthermore, mass loss from debris-covered glaciers in particular is closely associated with the formation of ice-contact and moraine-dammed lakes (Reynolds 2000; Benn *et al.* 2012; Sakai 2012; King *et al.* 2019). Their impact of lake evolution on local hazard potential in the context of future climate projections is still unclear (Harrison *et al.* 2018).

Over the last decade, debris-covered glaciers have become a ‘hot topic’ in glaciological research following concerns about the fate of glaciers, particularly in High Mountain Asia (Cogley *et al.* 2010; Bolch *et al.* 2012). During this time, several satellite remote sensing studies showed that thickly debris-covered glaciers display high



**Fig. 1.** Surface of a debris-covered glacier: Ngozumpa Glacier in the Nepal Himalaya. Photo taken in 2008; credit: A. Racoviteanu.


rates of surface lowering, comparable to those of clean ice glaciers (Kääb *et al.* 2012; Nuimura *et al.* 2012; Gardelle *et al.* 2013; Pellicciotti *et al.* 2015; Brun *et al.* 2019), even though a thick debris mantle has been conclusively shown to locally reduce ablation compared to that of clean ice (Østrem 1959; Kayastha *et al.* 2000; Nicholson and Benn 2006). This mass loss has been attributed to modified ice dynamics (Vincent *et al.* 2016; Brun *et al.* 2018; Anderson *et al.* 2021; Rounce *et al.* 2021) and to localized ice ablation rates related to ice cliffs and ponds (Sakai *et al.* 2000b; Miles *et al.* 2018b; Buri *et al.* 2021). The complex surface topography of debris-covered tongues exhibits exposed ice cliffs (Steiner *et al.* 2015; Buri and Pellicciotti 2018), surface ponds of various sizes (Sakai and Fujita 2010; Watson *et al.* 2016; Miles *et al.* 2018b), debris cones/hummocks (Moore 2018; Bartlett *et al.* 2021), medial moraines (Anderson 2000), lateral and terminal moraines (Hewitt and Shroder 1993; Owen *et al.* 2003; Benn *et al.* 2004), supraglacial streams (Fyffe *et al.* 2019a; Miles *et al.* 2020), surface depressions (Mertes *et al.* 2016; Benn *et al.* 2017; Miles *et al.* 2017a), relict englacial conduits (Gulley and Benn 2007; Gulley *et al.* 2009b), base-level lakes (terminal, proglacial or supraglacial, and proto-lakes) (Benn *et al.* 2012) and supraglacial vegetation (Fickert *et al.* 2007; Tampucci *et al.* 2016; Anderson *et al.* 2020) (Fig. 2). Certain supraglacial features act as ‘hot spots’ for ice melt, particularly ice cliffs (Sakai *et al.* 2002; Han *et al.* 2010; Steiner *et al.* 2015; Buri *et al.* 2016, 2021) and supraglacial ponds (Sakai *et al.* 2000b; Miles *et al.* 2016, 2018b; Salerno *et al.* 2017). In spite of the prevalence of these features, the insulating effect of a thick debris cover is still predominant on many glaciers (Vincent *et al.* 2016; Brun *et al.* 2018; Anderson *et al.* 2021), and is evidenced at a mountain-range scale by debris-covered glaciers having lower terminus elevations than clean ice glaciers.

Local, regional and global patterns of glacier thinning and mass loss are coupled with an increase in debris cover extent as the upper

limit of the debris cover migrates upglacier with the equilibrium line of the glacier (Deline 2005; Anderson and Mackintosh 2012; Herreid and Pellicciotti 2020; Xie *et al.* 2020a). Debris thickness increases due to cumulative exposure of englacial debris as glaciers thin due to surface ice ablation. While progress has been made in understanding local glacier-surface dynamics related to these supraglacial features, the extent to which the evolution of the debris-covered glacier surface influences overall glacier behaviour remains uncertain.

Insights into the surface characteristics of debris-covered glaciers and the enhanced local ablation rates gave rise to another concern relevant to both the scientific community and local communities, namely the accelerated growth of supraglacial and proglacial lakes associated with glacier thinning and recession documented around the world (Paul *et al.* 2007; Komori 2008; Gardelle *et al.* 2011; Thompson *et al.* 2012; Wang *et al.* 2014; Nie *et al.* 2017; Shukla *et al.* 2018; Chen *et al.* 2020; Shugar *et al.* 2020). Proglacial lakes exhibit an effect on glacier ice dynamics through enhanced ablation at the glacier margins via mechanical and thermal stresses; they modify meltwater routing and sediment fluxes through sedimentation (Carrivick and Tweed 2013). While proglacial lakes and their hazard potential have been addressed in several studies (Reynolds 1999; Watanabe *et al.* 2009; Aggarwal *et al.* 2017; Haritashya *et al.* 2018; Wilson *et al.* 2019), the link between supraglacial lakes and glacial hazards is less well studied.

There has been an increased interest in understanding the conditions for the formation and evolution of supraglacial ponds on debris-covered glaciers (Sakai and Fujita 2010; Sakai 2012). Some are highly dynamic, quickly evolving and growing, while others are persistent; some are short-lived (Miles *et al.* 2018a), while others may coalesce through time to create a larger moraine-dammed lake at the terminus of the glacier, adjoining a calving glacier front (Benn *et al.* 2012). The formation of moraine-dammed



**Fig. 2.** Components of a landsystem model for debris-covered valley glaciers. Relative positions of different surface features are indicative, as supraglacial features can exist in numerous configurations (credit: Gareth Evans and Naomi Lefroy).

lakes is associated with glacier retreat and downwasting patterns (Korup and Tweed 2007; Carrivick and Tweed 2013) and is favoured by low glacier slope and velocities (Quincey *et al.* 2007) and/or changes in supraglacial debris flux (Benn *et al.* 2012). Water stored behind a weak moraine has the potential to breach the moraine dam, resulting in glacier lake outburst floods (GLOFs). These involve the rapid release of large volumes of water and sediments, with disastrous consequences for communities downstream. The Dig Tsho flood event in the Khumbu Himalaya in 1985 was one of the largest such events recorded (Vuichard and Zimmermann 1987; Mool *et al.* 2002).

The evolution of moraine-dammed lakes associated with debris-covered glaciers has been addressed in few studies (e.g. Bolch *et al.* 2008b; Thompson *et al.* 2012, 2017; Harrison *et al.* 2018). Assessing the hazard potential of these lakes presents significant challenges both in the field due to the highly dynamic environment and in remote sensing due to limited multi-temporal, high-resolution data needed to estimate glacier surface evolution in some areas (Wang *et al.* 2020). Another important challenge is posed by the lack of systematic approaches for classifying and ranking proglacial and supraglacial glacier lakes in terms of their hazard potential. This results in significant gaps in glaciological and geomorphological characteristics of debris-covered glaciers and associated lakes, hindering the understanding of the future evolution of these glaciers and its implications for glacier hazards and water resources. A deeper and broader, contextualized understanding is prerequisite to identifying and monitoring the geohazards and evaluating the hydrologic implications associated with changes in the debris-covered glacier system under future climate scenarios.

Addressing this research gap requires a more complete estimate of glacier responses to climate change and their impacts, notably a better understanding of the debris-covered glacier landsystem, its

components and the interplay of various factors such as local geomorphology, ice ablation patterns, debris characteristics, glacier lake growth and development. In order to achieve this, a systems approach is needed. Traditionally, debris-covered glacier processes are addressed by a single discipline such as geomorphology, glaciology or hydrology; very few studies have adopted the required holistic, systemic approach. Although glaciology is an increasingly interdisciplinary field, most scientists are driven to specialize and only study one or few aspects of the glacier system (e.g. melt processes, hydrology or ice dynamics). Benn *et al.* (2012) is one example of a holistic study that links climate, mass balance, ice dynamics, topographic evolution and hydrology in the Everest region of Nepal and explores how observed behaviour and hazard potential emerge from interactions between these process domains. Lake hazard assessments are often conducted from a remote sensing or geophysical perspective (Pant and Reynolds 2000; Rana *et al.* 2000; Richardson and Reynolds 2000; Reynolds 2006; Hambrey *et al.* 2008; Thompson *et al.* 2012, 2017). A systems approach combines various perspectives to provide a more complete picture of the debris-cover-lake system, i.e. the interplay between the glacier surface topography, the lake dynamics and the ablation patterns related to supraglacial debris cover and its characteristics. Furthermore, this type of holistic perspective allows a better understanding of the past, present and future states of debris-covered glaciers, as well as their position, role and consequences within landscapes. This is key for various applications, but in this paper we focus on its value for estimating the hazards related to rapidly evolving moraine-dammed glacier lakes and their impacts on populations. We identify misunderstandings related to these concepts and failures in the way they have been communicated, and suggest ways to bridge these gaps to develop an understanding of resilience to climate change.

## The debris-covered glacier landsystem: concept and components

In this section we consider the question ‘what is a debris-covered glacier?’ from a *landsystem perspective*. In doing so, our aim is to widen the focus to understand how the glacier and landscape interact. In light of this, we must consider a debris-covered glacier as a system with a particular assemblage of features, associated with the higher rate of debris loading and exhumation than a typical valley glacier system. For example, while a simple and commonly used definition for identifying a debris-covered glacier is one where debris material covers the full width of part of its ablation zone (Kirkbride 2011), here we define a debris-covered glacier as one whose surface mass balance is sufficiently affected by supraglacial debris as to alter the glacier geometry, ice dynamics, surface features and hydrology (or some subset of the above) compared to that of a clean-ice glacier. The relative debris-richness of the glacier system controls its propensity to become debris-covered (Kirkbride 1989). Placing debris-covered glaciers on a continuum of mixed ice/debris landforms in this way ties the degree of debris cover to the relative abundance of snow v. debris supply. From this viewpoint, negative glacier mass balance conditions in the presence of abundant debris are expected to lead to development of a debris cover.

Rock debris is supplied to the glacier surface by gravitational mass movements from the surrounding terrain (e.g. Nakawo *et al.* 1986; Hambrey *et al.* 2008; Nagai *et al.* 2013), which generally occur as isolated events in space and time, and are poorly represented by the scant available measurements (Deline 2009; Hewitt 2009; Reznichenko *et al.* 2011) or by estimates derived from long-term headwall retreat rates (Heimsath and McGlynn 2008; Seong *et al.* 2009). Debris deposits onto the glacier surface within the accumulation zone are buried and transported englacially until they are exhumed to the surface by ice melt within the ablation zone (Kirkbride and Deline 2013). Deposits directly onto the ablation zone remain at the surface. Theoretical considerations and modelling studies (e.g. Rowan *et al.* 2015; Anderson and Anderson 2016; Wirbel *et al.* 2018; Scherler and Egholm 2020) highlight that the specific location of debris inputs strongly influences the spatial pattern of supraglacial debris. A constant rate over a longer time period cannot produce a localized, high debris concentration within the glacier, but will lead to an extended zone of lower concentration, which will produce distinctly different surface debris patterns compared to an initially localized zone of high concentration (Wirbel *et al.* 2018). The emergence of debris in the ablation zone is governed by the debris input location, englacial transport and melt-out rates. Thus, to accurately compute the point of emergence and thickness of debris at this melt-out location, the englacial transport pathways and deformation must be captured in some way (Kirkbride and Deline 2013; Wirbel *et al.* 2018). Once on the glacier surface, debris is transported by advection with underlying ice flow, where gradients in the ice surface velocity and resulting zones of compressive or extensional ice flow will thicken or thin the debris cover layer. In addition, as soon as debris emerges on the glacier surface, it is affected by other processes such as gravitational reworking (Anderson 2000; Kirkbride and Deline 2013; Moore 2018; Fyffe *et al.* 2020). Irregular englacial debris concentration and subsequent surface reworking causes local debris thickness variability (Moore 2018; Nicholson *et al.* 2018; Westoby *et al.* 2020), leading to strong small-scale variability in ablation rates and the formation of pronounced surface relief and features (Benn *et al.* 2001; Mertes *et al.* 2016).

The supraglacial debris mantle has a profound influence on the underlying ice ablation rate. Early studies (Østrem 1959) showed that glacier surfaces with patchy or very thin cover of supraglacial debris experience accelerated ablation, whereas a continuous debris cover of thickness greater than a few centimetres inhibits the

underlying ice ablation. The specific effect of debris on ice melt is influenced by its individual characteristics such as debris thickness, debris material, porosity, grain size, moisture and liquid water content and the prevailing meteorological conditions. However, field studies (e.g. Fujii 1977; Mattson *et al.* 1993), laboratory experiments (Reznichenko *et al.* 2010) and modelling studies (e.g. Nakawo and Young 1982; Nicholson and Benn 2006; Reid and Brock 2010) demonstrate that debris thickness is the primary determinant of how sub-debris ice ablation rates differ to clean-ice melt rates, with the properties of the debris layer playing secondary roles (Reznichenko *et al.* 2010; Nicholson and Benn 2012; Collier *et al.* 2014). As surface debris is continuously conveyed down-glacier with ice flow, debris cover thickness increases towards the glacier terminus (Rowan *et al.* 2015; Anderson and Anderson 2018). This profoundly alters the glacier-scale ablation regime, in principle causing an inversion of the ablation gradient toward the terminus such that maximum ablation occurs some distance upglacier instead of at the terminus as is the case for clean-ice glaciers (Benn and Lehmkuhl 2000; Bisset *et al.* 2020). This in turn has consequences for ice dynamics as the ablation gradient influences the development of the glacier surface longitudinal profile and thereby the driving stresses through the ablation zone.

Debris can be removed from the system by marginal deposition or by surface meltwater. Some debris-covered glaciers form large, impounding latero-terminal moraine complexes (e.g. Benn and Owen 2002; Hambrey *et al.* 2008) while other debris-covered glacier termini end in outwash plains without substantial terminal moraines (e.g. Mayer *et al.* 2006). Large terminal moraines affect the englacial water table, and increase the potential for water to be stored behind this impounding moraine (e.g. Benn *et al.* 2012). In the absence of impounding latero-terminal moraines, terminal lakes can only form by external geomorphological processes, for example where water courses are impounded by advances of neighbouring glaciers, or slope failures damming the valley downstream (Rashid *et al.* 2020). Large latero-terminal moraines also inhibit the evacuation of debris from the glacier surface by gravitational processes, and exert a physical constraint on the glacier terminus position and upstream ice dynamics. The sedimentological, geomorphological and dynamic context of debris-covered glaciers has been discussed by Hambrey *et al.* (2008). They presented a conceptual model for the eastern Himalaya applicable to other glaciers to explain the development of large, lateral-terminal moraine complexes and associated moraine dams. The presence or absence of confining moraine dams may play a decisive role in determining the end-member of glacier development under declining ice content. If present, they facilitate the formation of terminal lakes, while their absence may allow the transition of debris-covered glaciers to rock glaciers or other ice-debris landforms (Whalley and Martin 1992; Jones *et al.* 2019).

## Tools for observing and monitoring the debris-covered glacier landsystem and its components

Understanding the future states of the debris-covered glacier landsystem requires the knowledge of the location of such glaciers, as well as their current extent and state. This has been the subject of previous mapping efforts, including regional and global estimates of supraglacial debris cover (e.g. Scherler *et al.* 2018; Herreid and Pellicciotti 2020). However, a complete understanding of the system also requires information on the fundamental debris surface characteristics in order to understand the ice ablation processes, the velocity and dynamics, the evolution of ice cliffs and ponds and their importance to hydrology and hazard events. In this section we focus on state-of-the-art techniques for mapping and monitoring these characteristics, focusing in particular on supraglacial debris

cover extent, debris thickness, physical properties and associated surface features. For each, we present both remote sensing and field methods, we outline current advances, remaining gaps and challenges, and offer recommendations to overcome these.

### ***Delimiting debris cover extent***

#### *Remote sensing*

Mapping of debris-covered glaciers received considerable attention in the late 2000s and early 2010s, with important improvements in monitoring capacity as satellite imagery improved in both spatial and temporal resolution and coverage. The release of the Randolph Glacier Inventory (RGI; Pfeffer *et al.* 2014; RGI\_Consortium 2017) and subsequently the global supraglacial debris cover datasets constructed on the basis of the RGI (Scherler *et al.* 2018; Herreid and Pellicciotti 2020) enabled a step-change in understanding the distribution of debris-covered glaciers at the large scale. However, while these global databases provide an initial and global perspective of glacier and supraglacial debris extent, they suffer from several limitations. The RGI's composition from distinct sources means that both datasets suffer from inconsistent methods between and often within regions, varying representative dates, user-subjective post-processing and manual delineation (e.g. Paul *et al.* 2013), geolocation or projection errors, and occasional inclusion of spatially descriptive but not explicit sources (e.g. World Glacier Inventory). These problems were partially mitigated in a manual revision limited to glaciers larger than 1 km<sup>2</sup> (Herreid and Pellicciotti 2020), but such an effort is laborious for repeat application at the global scale. Consequently, although debris-covered glaciers may be accurately represented in available databases for well-known sites that have been mapped carefully, their representation may be inconsistent and may occasionally commit and/or omit entire features for areas that are less well surveyed (Racoviteanu *et al.* 2021). Thus, while acknowledging that the mapping of supraglacial debris within the bounds of the RGI (e.g. Scherler *et al.* 2018; Herreid and Pellicciotti 2020) constitutes an important advance in high-level understanding of debris-covered glacier distribution globally, we consider that the current representation of debris-covered ice within the RGI is not sufficiently robust or consistent for understanding debris-covered glacier processes and change. Accurate, large-scale mapping of debris-covered glacier tongues at multi-temporal resolution remains a gap.

In optical remote sensing, identifying the glacier boundary is difficult due to the spectral similarity of supraglacial debris to the surrounding moraines (Racoviteanu and Williams 2012). Previous remote sensing studies have used a combination of terrain information, spectral information and terrain curvature (Bishop *et al.* 2001; Paul *et al.* 2004; Bolch *et al.* 2007; Shukla *et al.* 2010a; Kamp *et al.* 2011) to map debris-covered glaciers. Recent studies combined these criteria in machine-learning algorithms in order to automate the mapping process (Robson *et al.* 2015; Zhang *et al.* 2019; Xie *et al.* 2020b; Holobăcă *et al.* 2021). Opportunities for method development have greatly improved in the past decade with the increased availability of new operational, rapid-repeat, and public satellite imagery (e.g. Sentinel). In addition to optical remote sensing, several other proxies offer promise and overcome its limitations, among which we mention the following.

- **Surface motion** derived from pairs of satellite optical or radar images helps identify active debris-covered areas (Gardner *et al.* 2018; Dehecq *et al.* 2019).

- Sequential **synthetic aperture radar (SAR) coherence** images which indicate changes in surface backscatter between repeat observations; SAR helps to identify the active parts of the debris-covered glaciers due to their motion-related decorrelation compared to the highly coherent surrounding areas (Strozzi *et al.*

2010; Frey *et al.* 2012; Robson *et al.* 2015; Lippl *et al.* 2018; Holobăcă *et al.* 2021). While SAR coherence images are widely available and overcome the limitations posed by cloud cover in optical remote sensing, wide application of SAR techniques is hindered by complex processing.

- **Satellite thermal imaging** helps distinguish the debris underlined by glacier ice and the surrounding non-ice moraines based on the brightness temperature difference (Taschner and Ranzi 2002; Shukla *et al.* 2010b; Bhabri *et al.* 2011; Racoviteanu and Williams 2012; Alifu *et al.* 2015).

- **Digital elevation model (DEM) differencing**, derived from topographic maps for example (GLAMOS 2020; Linsbauer *et al.* 2020) serves to identify surface lowering. This is based on the concept that even where debris cover is thick, some heat is generally transmitted through the debris layer seasonally, leading to a small amount of ice melt and thus resulting in surface lowering.

- **Surface roughness and characteristics** including pitted, hummocky topography with sharp breaks in slope, incised channels, etc. from a high-resolution DEM (King *et al.* 2020a) help to identify debris characteristics that may differ quantitatively from other land surfaces;

- **Object-oriented and machine-learning techniques (OBIA;** Robson *et al.* 2015; Khan *et al.* 2020) based on shape is a complement to debris-cover mapping procedures.

The challenge in remote sensing mapping of debris cover is how to capitalize on new monitoring tools to develop targeted repeat monitoring in a systematic manner. Currently, most of the methods presented above still need manual post-processing of glacier shapes to ensure that glacier outlines conform to our understanding of ice flow and known landforms. These studies remain limited to specific regions; there remains a need for an automated tool, workflow or set of methods to delimit debris-covered ice independent of the bounds of given glacier polygons. In order to achieve this, the above proxies should be used in addition to optical remote sensing to construct a method that should be transferable to different sites, should rely on freely available, global coverage data sources, and should be validated against the best available field data.

#### *Field methods*

Field-based delineation of debris cover extent is generally difficult. Many debris-covered glaciers occupy remote, rugged domains, with limited field access and the precise boundary of a debris-covered glacier is challenging to identify in the field. Validating the above remote sensing methods is particularly difficult as field methods for debris-covered glacier extent mapping are also not in an advanced state and detailed field studies are site-specific. Promising field-based methods based on ground temperatures, geophysics, drone-deployed optical and thermal imaging and time-lapse photography are time consuming, site-specific and difficult to extrapolate over larger areas. Simple recognition of debris-covered glacier surface features in the field (cliffs, ponds, etc.) can often be helpful to identify the presence of sub-debris ice, but it is often easier to (subjectively) identify debris-covered glacier extent from a planimetric perspective (e.g. high-resolution optical satellite data, ideally with multitemporal data) than in the field.

### ***Determining the spatial distribution of debris cover thickness***

#### *Remote sensing*

Surface debris thickness modulates the surface temperature and exerts a physical control over sub-debris melt rates (Østrem 1959; Nicholson and Benn 2006); it is probably the most crucial but also the most difficult debris cover property to quantify and monitor; field measurements of debris cover thickness are difficult to obtain

in the field due to the rugged terrain. Therefore, satellite remote sensing approaches have been increasingly used in recent decades to overcome this challenge.

○ **Thermal imagery:** a variety of approaches of varying complexity have used satellite thermal data to estimate debris thickness (Boxall *et al.* 2021). These range from simple band thresholding (Ranzi *et al.* 2004) to exponential curve fitting based on the empirical relationship between surface temperature and thickness (Juen *et al.* 2014; Kraaijenbrink *et al.* 2018) or energy-balance inversion, often requiring model spin-up (Mihalcea *et al.* 2008; Zhang *et al.* 2011; Foster *et al.* 2012; Rounce and McKinney 2014; Schauwecker *et al.* 2015; Rounce *et al.* 2021; Stewart *et al.* 2021). An intercomparison of these methods is needed and has been identified as a research target for the IACS Debris-covered Glaciers Working Group (<https://cryosphericsscience.org/activities/wgdebris/>).

○ **Elevation change/surface mass balance:** since the thickness of debris moderates energy transfer to the ice, it also controls ice melt rates (Østrem 1959; Nicholson and Benn 2012). Surface mass balance data can thus be used to invert an energy mass balance model for debris thickness (Ragetti *et al.* 2015; Rounce *et al.* 2018), although this often requires careful consideration of ice dynamics to estimate surface mass balance from elevation change, and the long-duration melt modelling is computationally expensive (Rounce *et al.* 2021).

○ **Polarimetric SAR:** as certain wavelengths of radar can penetrate into the debris surface, the attenuation of radar signals is indicative of debris thickness. Debris cover thickness can be estimated based on inversion of the volume scattering power and other parameters after target decomposition. This method is in its infancy, but shows promise for an independent assessment of debris thickness (Huang *et al.* 2017).

Debris thickness can be inferred from proxy data in remote sensing studies. Regional and global applications of these methods (e.g. Rounce *et al.* 2021) represent a key advance towards including explicit inclusion of the effects of debris cover in global glacier simulations. However, as with debris-covered glacier extent, methods for determining debris thickness require validation, which is usually achieved through field measurements. We thus recommend the continued investigation of local-scale debris thickness, and the compilation of a database of available debris thickness measurements. This is another important aspect of the IACS Debris-covered Glacier Working Group, which has begun to assembly of a repository of debris-related measurements via the Zenodo data repository (<https://zenodo.org/communities/iacswwgondcgs>). Each dataset is assigned a unique DOI to ensure that the responsible parties receive appropriate credit.

### Field methods

Field measurements of debris cover thickness are all spatially limited and labour-intensive to obtain. Manual excavation (e.g. Zhang *et al.* 2011) is practically limited to debris cover thickness < 0.8 m; high-frequency ground-penetrating radar (GPR) (e.g. McCarthy *et al.* 2017) is hindered by the difficulty of deploying GPR equipment at remote, high-elevation sites; estimates based on oblique terrestrial photography (e.g. Nicholson and Mertes 2017) require a number of crude geometric assumptions; estimates based on unmanned aerial vehicle (UAV) or terrestrial thermal imagery (Steiner and Pellicciotti 2016; Kraaijenbrink *et al.* 2018) have large uncertainties associated with image processing, due to local surface temperature variations as a result of shading or moisture and signal saturation at thickness >~30 cm (Steiner *et al.* 2021); calculation from field measurements of surface lowering by any terrestrial data acquisition (e.g. structure from motion DEMs, TLS/LIDAR,

terrestrial radar) are difficult to upscale to the glacier extent. In general, while these methods all have great potential to complement regional remote sensing studies, they currently have limited application due to logistical difficulties and limited spatial extent.

### Estimating surface velocity on debris-covered glaciers

Remote sensing methods are key for estimating ice velocities, and can be applied to the surface of debris-covered glaciers. Several studies used remote-sensing surface velocities to show the deceleration of ice flow downglacier leading to stagnation at the snout (Quincey *et al.* 2007; Hambrey *et al.* 2008; Haritashya *et al.* 2015). Flow velocities can be derived by feature tracking using satellite imagery such as ASTER, Landsat series or Sentinel (e.g. Berthier *et al.* 2003, 2005; Kääb 2005; Scherler *et al.* 2008; Dehecq *et al.* 2015, 2019; Millan *et al.* 2019) and established image coregistration methods (Leprince *et al.* 2007). Methodological advances and data availability led to the globally comprehensive and temporally dense multi-sensor record of land ice velocity from the Inter-mission Time Series of Land Ice Velocity and Elevation project. However, the spatial resolution of these data (120–240 m) remains an issue, as the data have limited application for monitoring narrow debris-covered glacier tongues. Recent databases with improved spatial resolution (50 m) (Millan *et al.* 2019) offer promise for monitoring of debris-covered surfaces, but this is limited by cloud cover. As for the debris thickness estimates, SAR can provide high-accuracy measurements of the direction and intensity of glacier flow in all weathers (Kumar *et al.* 2011) provided that corrections are applied to mitigate attitude effects and sensor distortions (Scherler *et al.* 2008).

### Other debris properties and features of interest

Beyond debris extent and thickness, debris properties such as lithology, grain size, porosity, stratification and stability (Table 1) (Casey *et al.* 2012; Casey and Kääb 2012; Juen *et al.* 2013) are important for specific applications. Local field mapping of these properties is difficult; thus measurements are scarce. At a glacier or

**Table 1.** Debris properties of interest for various applications, and remote sensing techniques used on previous studies to estimate them

Property	Data source/technique	Existing studies
Debris lithology	Medium to high optical remote sensing and hyperspectral data combined with <i>in situ</i> field spectrometry	Casey <i>et al.</i> 2012; Casey and Kääb 2012
Debris grain size	High-resolution imagery, SfM	Miles <i>et al.</i> 2017b; Detert and Weitbrecht 2020
Debris layer porosity or bulk density	No remote sensing method known yet; may be possible with polarimetric SAR	
Water content	No remote sensing method known yet; may be possible with passive microwave	Collier <i>et al.</i> 2014; Giese <i>et al.</i> 2020
Supraglacial vegetation	Normalized Difference Vegetation Index, NDVI; spectral unmixing	Fickert <i>et al.</i> 2007; Racoviteanu <i>et al.</i> 2021
Thermal conductivity	No remote sensing method known yet	—
Broadband albedo	Formulae for anisotropy correction and narrow to broadband conversion, though not developed for rock debris specifically	Knap <i>et al.</i> 1999; Liang 2001; Greuell and Orlemans 2004; Naegeli <i>et al.</i> 2017; Xu <i>et al.</i> 2020

regional scale, many debris cover properties and features are more easily mapped using high-resolution satellite imagery than in the field. Therefore, in this section we only discuss the remote sensing mapping and monitoring of these features as summarized in Table 1, with a focus on ice cliffs, ponds and streams.

### Ice cliffs

Although no formal definition for ice cliffs exists (Kneib *et al.* 2020) these are readily identifiable in the field as high-relief bare-ice areas interrupting the supraglacial debris layer and are often associated with a supraglacial pond. An increasing number of studies are using remote sensing techniques to identify ice cliffs from satellite data (Table 2). However, robust and transferable methods for mapping ice cliffs in a consistent manner are in their infancy. In addition, more studies are needed simply to assess the long-term changes in prevalence of ice cliffs, as well as the spatial differences in ice cliff occurrence. Of critical consideration for the above methods is the spatial resolution, and how resolved elevation models need to be to sufficiently represent ice cliffs. For example, high-resolution DEMs (c. 10 m spatial resolution) are available now at regional or global scales, some at no cost. These include the High Mountain Asia (HMA) DEM at 8 m (Shean 2017), ArcticDEM at 2 m (Noh and Howat 2015) or the TanDEM-X DEM at 12–30 m (Wessel *et al.* 2018) spatial resolutions. However, the spatial resolution of some of these DEMs, particularly TanDEM-X as well as other commonly available ASTER GDEMs or SRTMs (30–90 m) is not sufficient for mapping ice cliffs, which are often only a few metres wide. One possible way forward might be to validate a topographic proxy for ice cliff density and area, as nadir-view satellite imagery will have difficulty representing the total area of steep ice cliffs accurately.

### Supraglacial ponds

These small superficial water bodies are important indicators of the debris-covered glacier's drainage system, i.e. they control the rate at which meltwater derived from the melting ice flows downstream (Irvine-Fynn *et al.* 2017), and they contribute to ice mass losses themselves (Sakai *et al.* 2000b; Miles *et al.* 2018b). Supraglacial ponds are considerably easier to identify in satellite imagery than ice cliffs, meaning that a number of properties can be targeted, including: (i) surface temperature (from satellite, UAV or terrestrial thermal imagery); (ii) lake volume (via sonar or topographic sink analyses to derive volume-area relationships); (iii) lake turbidity (blue index or with sub-pixel spectral analyses); (iv) changes in elevation (high-accuracy DEMs). Supraglacial ponds and their

properties have received focused study over the past few years, largely with satellite data, and the optical methods to map them are well established (Table 3). Overall, few detailed field studies of supraglacial ponds exist, and more direct observations of ponds, their characteristics and their dynamics are still needed using a combination of the methods briefly outlined here. Contemporary satellite imagery can answer some of the current questions related to supraglacial ponds, including their seasonality and persistence, but efforts are needed to assess both properties and processes at the local scale, as well as their prevalence and change at the regional scale.

### Supraglacial streams

The inverted ablation gradient and low longitudinal gradient of debris-covered glaciers can have a strong impact on the structure and function of the glacier's entire drainage system (see the review by Miles *et al.* 2020). Supraglacial hydrology is directly observable in the field and with satellite data. Areas of thicker debris and lower debris are typically characterized by small catchments and discontinuous, low-efficiency drainage systems conducive to formation of supraglacial ponds (Miles *et al.* 2017b; Fyffe *et al.* 2019b) whereas areas of thinner debris and higher surface gradient can support larger catchments and efficient supraglacial stream systems (Gulley *et al.* 2009a; Miles *et al.* 2019). The relative extent of these domains is indicative of the glacier's decay and progression to stagnation (Benn *et al.* 2017; King *et al.* 2020a), but also important for understanding the diurnal and seasonal evolution of glacial discharge (Fyffe *et al.* 2019a). As supraglacial streams exist only where stream incision exceeds the background ablation rate (Marston 1983), they by definition directly contribute to melt; they also contribute indirectly to melt by promoting ice cliff development (Mölg *et al.* 2020; Kneib *et al.* 2021). Streams can be mapped using hydrologic analysis tools on high-precision, high-resolution topographic datasets derived from satellite stereo or UAV images (Benn *et al.* 2017; Miles *et al.* 2017b; Fyffe *et al.* 2020). Other efforts have mapped streams manually from satellite images or by walking their length in the field (Miles *et al.* 2019). Mapping supraglacial streams with DEM drainage analysis and optical imagery is similarly challenged by apparent stream discontinuities due to ice arches and flow through debris. Despite their importance for characterizing glacier drainage systems and debris-covered glacier stagnation, supraglacial streams have been addressed in fewer detailed studies than englacial conduits. Newly available high-resolution satellite images and DEMs offer the potential to better characterize supraglacial streams, but additional field investigations are needed to produce a generalized quantitative model of debris-covered glacier drainage efficiency.

**Table 2.** Remote sensing techniques used to map ice cliffs on debris-covered glaciers

Technique	Data source	Notes	Existing studies
Manual delineation of ice cliff crest or area	Satellite imagery (high resolution) including UAV	Time consuming and subjective; limited application at large scales	Sakai <i>et al.</i> 2002; Han <i>et al.</i> 2010; Brun <i>et al.</i> 2016; Steiner <i>et al.</i> 2019; Stefaniak <i>et al.</i> 2021
Use of thermal imagery	ASTER, Landsat	Spatial resolution is challenging	Herreid and Pellicciotti 2018
Feature detection (OBIA)	High-resolution imagery including DEM	Somewhat time consuming to set up the rules and may need some post-processing	Kraaijenbrink <i>et al.</i> 2016; Watson <i>et al.</i> 2017; Mölg <i>et al.</i> 2019
DEM slope thresholding/topography relief metrics	High-resolution DEM	Can be used as proxy for ice cliff presence; high resolution DEMs limited in some areas	Reid and Brock 2014; Herreid and Pellicciotti 2018; King <i>et al.</i> 2020a
SAR intensity tracking	High-resolution SAR, e.g. TerraSAR-X, IceEye	Potentially useful in areas where cloud-free optical data not available; other confounding surfaces	no studies yet
Optical broadband adaptive thresholding	High-resolution optical imagery	Fast, requires optimization of threshold	Anderson <i>et al.</i> 2021
Multispectral thresholding; adapted linear spectral unmixing	High-resolution multispectral optical imagery	Accurate, requires optimization	Kneib <i>et al.</i> 2020

**Table 3.** Existing remote sensing techniques to map supraglacial ponds on debris-covered glaciers

Technique	Data source	Notes	Existing studies
Manual delineation	Any high-resolution data (<10 m)	Time consuming and subjective but probably most accurate	Iwata <i>et al.</i> 2000; Salemo <i>et al.</i> 2012; Thompson <i>et al.</i> 2012; Watson <i>et al.</i> 2016; Miles <i>et al.</i> 2017b
Band ratios or normalized-difference such as NDWI	Landsat, Sentinel-2, ASTER, etc. (>10 m)	Manual/semi-automatic thresholding; can be improved using Planet microsatellite constellation which enables rapid-repeat monitoring at 3 m resolution	Wessels <i>et al.</i> 2002; Bolch <i>et al.</i> 2008a; Gardelle <i>et al.</i> 2011; Liu <i>et al.</i> 2015; Narama <i>et al.</i> 2017; Miles <i>et al.</i> 2017a; Watson <i>et al.</i> 2018a; Kneib <i>et al.</i> 2020
SAR backscatter intensity	Sentinel-1, TerraSAR-X, ENVISAT ASAR, ALOS PALSAR, ERS-1, 2 etc.	The oblique geometry of radar imaging creates challenges for pond identification (and especially for monitoring) due to the highly variable topography of debris-covered glaciers	Strozzi <i>et al.</i> 2012; Wangchuk and Bolch 2020; Zhang <i>et al.</i> 2021
Feature extraction via decision-trees and/or OBIA	Landsat, Sentinel-2, Pleiades etc.	Rules needed are subjective; it often commercial software	Panday <i>et al.</i> 2011; Liu <i>et al.</i> 2015; Kraaijenbrink <i>et al.</i> 2016
Sub-pixel spectral analysis	Landsat, Sentinel, ASTER	Upon careful end-member collection, ideally from field spectrometry; can be applied automatically at large scales	Panday <i>et al.</i> 2011; Scherler <i>et al.</i> 2018; Kneib <i>et al.</i> 2020; Racoviteanu <i>et al.</i> 2021
Thermal imaging	Landsat, ASTER, UAV	<ul style="list-style-type: none"> <li>Supraglacial ponds are usually considerably cooler than surrounding debris (during the day, debris can reach 25°C or more)</li> <li>•• Application of thermal imagery is limited by relatively coarse resolution (90–100 m), potentially most suitable at present for UAV thermal imagery when existent</li> </ul>	Suzuki <i>et al.</i> 2007

### Proglacial lakes


Proglacial lakes have been studied through long-term and regional-scale monitoring efforts (e.g. Zhang *et al.* 2015; Nie *et al.* 2017; Shugar *et al.* 2020) and have been mapped in a systematic manner using established methods based on historical multispectral (optical) imagery (Fig. 3) using various water indices (Zhang *et al.* 2018; Zhao *et al.* 2018) or manual digitization (Wilson *et al.* 2018). In general, proglacial lakes are easier to map from remote sensing than supraglacial lakes due to their larger size. A number of these lakes have bathymetric field surveys undertaken to assess to their hazard potential (Worni *et al.* 2013; Haritashya *et al.* 2018). Here we note a few specific aspects of proglacial lake mapping that are important to consider for future studies.

- **Shadows** cast across the lake surface have been historically problematic for automatic lake mapping efforts, but indices have emerged recently that show improved performance (e.g. Chen *et al.* 2013). More problematic are the shadows that are cast across non-water surfaces, which sometimes alias as water (this is also a challenge for supraglacial pond mapping) (see Gardelle *et al.* 2011; Miles *et al.* 2017a). Some strategies to mitigate this include topographic shadow casting for correction or the use of multi-temporal data to filter out shadows not associated with water.
- **Water turbidity** can cause a varying spectral signal across the surface of a glacial lake, and is also useful to observe as indicative of water circulation patterns, discharge plumes and bulk suspended sediment (Wessels *et al.* 2002; Kraaijenbrink *et al.* 2016). However, the combination of surface ice and varying turbidity can cause problems for automated algorithms. Nonetheless, automated determination of surface water turbidity from satellite imagery has been accomplished in other regions (Matta *et al.* 2017), and could be transferred to glacial lake assessments.
- **Surface ice** (lake ice and icebergs) are extremely useful indicators of environmental conditions and processes, but are usually confounding factors for automated methods. Calving rates in particular have been a key target of study in marine environments and for emergent lakes, but have received relatively little attention compared with debris-covered glacier or high-mountain glacier lakes.
- **Surface temperature** can be useful for the delineation of large lakes, as resolution matters less, and is also a useful property itself as a controlling factor for lake water stratification and circulation (along with turbidity) and to understand the energy balance of the ice–lake–stream domain.

### Applications of the remote sensing methods and further considerations

Of the methods mentioned in Table 1, we note here that SAR intensity mapping is extremely promising for glacial lake monitoring efforts, especially in cases where lakes are undergoing rapid change. Unlike optical data, SAR intensity mapping is insensitive to clouds and shadows, and less sensitive to turbidity and surface ice factors (e.g. Strozzi *et al.* 2012; Wangchuk and Bolch 2020). Furthermore, due to the size of most glacial lakes, the SAR intensity method is less affected by topographic and resolution issues than for supraglacial ponds and ice cliffs.

While we recognize that supraglacial features (ice cliffs and supraglacial ponds) and their dynamics are important to understand, it is useful to consider each in terms of their causal or controlling processes. Some features are associated with debris dynamics (e.g. differential ablation, debris deposition or emergence), while others are hydrologically/fluviually associated. We consider that mapping



**Fig. 3.** Decadal evolution of lakes in the Zemu basin of Sikkim Himalaya based on remote sensing: (a) 1962 panchromatic Corona KH4 imagery (7.5 m); (b) 2001 ASTER image; (c) 2006 Quickbird image (2.4 m) (d) 2020 PlanetScope image (3 m). All multispectral images are shown as colour composites (bands 3, 2 and 1) (revised and expanded from Racoviteanu *et al.* 2015).

strategies should be driven by a specific research question. For example, ice cliffs may be studied as an indicator of debris redistribution or they can be regarded as exposed ice within the debris-covered domain in order to more accurately represent ablation rates. If areas of high meltwater production are of interest, identifying all the exposed ice on the debris cover might be more important than specifically delineating only ‘ice cliffs’. This would include other ice surface features such as ice sails (Evatt *et al.* 2017).

For GLOF hazard assessments, there is clearly a need for widespread screening of proglacial lakes and supraglacial ponds on a regional scale using at least semi-automated if not fully automated techniques. At local scales, proglacial lakes can be monitored in the field using bathymetric surveys (Cook and Quincey 2015; Watson *et al.* 2018b). So far, relatively few lakes have been the subject of such surveys and this needs further addressing. Multiple recent, current and future satellite altimeters (IceSat, Cryosat, IceSat2, SWOT) are promising avenues of operational workflow development. IceSat2 penetrates within many shallow-water bodies and might be suitable for bathymetric mapping of some proglacial lakes, although water turbidity of proglacial lakes limits the optical penetration depth; this needs direct analysis to test its feasibility.

### Response of the debris-covered glacier landsystem to climate change

The global pattern of glacier recession (e.g. Kargel *et al.* 2014) and the global nature of climate warming indicate a clear attribution to climate change (e.g. Marzeion *et al.* 2014; Zemp *et al.* 2015; Roe *et al.* 2017). Clearly, the mass balance of a glacier is causally linked to changes in temperature and precipitation, with accelerated negative trends of mass loss in the 21st century (Zemp *et al.* 2015; Solomina *et al.* 2016; Hugonnet *et al.* 2021). However, at regional scales, glaciers exhibit contrasting patterns in their response to climate changes (Sakai and Fujita 2017) due to differences in local topo-climatic factors (Salerno *et al.* 2017; Brun *et al.* 2019). Furthermore, local meteorology is usually not precisely known due to scarce measurements, leading to simplified modelling optimization schemes (Hock 2003). Complicating variables for mass accumulation include the addition of snow avalanches to mass balance and the importance of wind-blown snow from surrounding catchments.

With regard to ice ablation, site-specific losses occur via dynamic processes such as calving related to ice flow and glacier surface characteristics. In addition, surface energy balance and related melt and sublimation losses are driven by spatiotemporally varying fields of potential insolation, temperature, cloudiness, relative humidity and wind, all of which can manifest very differently depending on glacier settings and surface conditions and which are not easily

characterized (Huo *et al.* 2021). However, these processes apply primarily in cases where the glacier surface is predominantly composed of exposed bare ice. It is observed in many mountain ranges that thickly debris-covered glacier termini persist at lower elevations than clean-ice glaciers. This indicates that the behaviour of a glacier terminus position in response to any given set of climate conditions is markedly different when the glacier has a surface debris cover compared to a clean-ice surface (Anderson and Anderson 2016). The geomorphological sensitivity of debris-covered glaciers is therefore an important and relevant concept (see Harrison 2009). While the geomorphological sensitivity of a clean-ice glacier could be established as its mass balance change over time and related to local climate change, it is not clear how we might assess the sensitivity of a debris-covered glacier, nor which metrics might be important.

The full response of debris-covered glaciers to climate forcing remains poorly understood in relation to that of clean-ice glaciers. One possible response of some mountain glaciers to climate change will be a transition from clean glaciers to debris-covered glaciers, and a further transition to rock glaciers in response to paraglacial processes increasing debris fluxes to glacier surfaces (see Monnier and Kinnard 2017; Jones *et al.* 2019). The long-term consequences of this transition are still largely unknown. In general, debris-covered glaciers pose a complicated case, where their behaviour and evolution are additionally related to non-climatic processes such as changes in debris flux from surrounding mountain sides or the presence of surface features such as ponds and ice cliffs. As a result, the system controlling the evolution of the debris-covered glacier system is not solely climatic in origin, but one in which paraglacial processes play an important role (Ballantyne 2002; Knight and Harrison 2014). Numerical models of glacier response to climate forcing under negative mass balance conditions suggest that debris-covered glaciers initially respond slower than clean-ice glaciers. However, the ultimate response time of debris-covered glaciers might be greater, as eventually the stagnant remnant of the glacier tongue detaches and decays *in situ* (Banerjee and Shankar 2013). The climate response of debris-covered glaciers is thought to be markedly asymmetrical between negative and positive mass balance conditions, with glacier adjustment rates to positive conditions matching those of clean-ice glaciers (Banerjee and Shankar 2013), such that the glacier length preferentially remembers positive mass balance phases over negative ones (Ferguson and Vieli 2020). There are very few observations of debris-covered glacier response to positive mass balance conditions (e.g. Deline 2005; Mölg *et al.* 2019), so the understanding gleaned from these modelling studies is unverified. However, it has been observed that substantial glacier advances can be triggered by extensive rockfall onto the glacier ablation zone. For example, following a large rock avalanche in

1920, the Brenva Glacier advanced 490 m between 1920 and 1941, whereas neighbouring glaciers in the Mont Blanc massif receded from the mid-1920s (Deline *et al.* 2015). This further highlights that the length of a debris-covered glacier is not a simple proxy for climate conditions alone. A better understanding of such processes is needed for long-term regional and global projections of glacier behaviour that form the basis of understanding trajectories of future meltwater availability and sea-level contribution from mountain glaciers (e.g. Kraaijenbrink *et al.* 2017; Rowan *et al.* 2017; Shannon *et al.* 2019).

### Evolution of debris-covered glacier systems

Our understanding of how debris-covered glaciers and related landforms will evolve in the future remains limited. This means that the impact of climate change on these ice-debris systems will vary as the systems change. Viewed from the landsystem perspective, a debris-covered glacier landsystem incorporates numerous processes that respond to climate in different ways over time. This process transience of the system components presents a key challenge in simulating coupled glacier–climate behaviour (Nicholson *et al.* in press). For example, warming might be expected to cause a monotonic shift in precipitation phase from solid to liquid (i.e. more precipitation falls as rain rather than snow), starving the glaciers of snow accumulation while simultaneously enhancing ablation by rainfall. However, debris supply rates may show a complex non-linear response to the same warming over time. For a debris-covered glacier, the debris cover characteristics change in time as a function of supply, transfer, melt-out, thickness distribution and removal. These processes all co-evolve over time in a manner that is dependent on how the glacier geometry and ice flow dynamics adjust to the debris-modified spatial pattern of ablation. As a result, inter-relationships between these system components observed thus far might not hold into the future, and this non-stationarity means that such relationships are subject to both lags in response as well as gradual and thresholding process change, which are challenging to incorporate into a model system capable of reproducing system development over time.

Surface debris supply rate on debris-covered glaciers can be enhanced by debuitressing of rockwalls exposed by glacier recession, which can cause weakening of the valley walls and slopes. The timescale and duration of this effect is difficult to constrain and contingent on many structural, lithological and geomorphological conditions (Knight and Harrison 2018; Mancini and Lane 2020). In the longer term, debris supply may be more controlled by the rockwall area that lies within the freeze–thaw zone (Nagai *et al.* 2013; Banerjee and Wani 2018) and can also be influenced by heatwaves and heavy rainfall events. Secondary debris supply from debuitressed lateral moraines is an additional non-stationarity that is interesting to grapple with (van Woerkom *et al.* 2019). The system debris content is also affected by debris evacuation rates, which is primarily governed by the nature of the terminal deposition environment. Debris-covered glacier termini ending in outwash plains (e.g. Mayer *et al.* 2006) can export sediment to the foreland, while those with large, impounding latero-terminal moraine complexes (Benn and Owen 2002; Hambrey *et al.* 2008) cannot readily do so. Changing debris load over time will influence, together with changing ice inputs and losses, how and when debris-covered glaciers can form, and when they might transition to ice-cored rock glaciers, for example, due to increasingly inefficient supraglacial sediment evacuation (Monnier and Kinnard 2017; Jones *et al.* 2019; Knight *et al.* 2019).


The characteristic downglacier increase in debris thickness (Anderson and Anderson 2018), and the associated ablation gradient inversion toward the glacier terminus implies that maximum ablation occurs at the upper part of the debris cover

and is reduced downglacier (Benn and Lehmkuhl 2000; Bisset *et al.* 2020). This favours glacier mass adjustment to negative mass balance conditions, by thinning instead of terminus retreat. As a result, the surface area change and terminus moraine position are poor indicators of glacier change for a debris-covered glacier. For example, in the Mont Blanc massif, the mostly clean-ice Mer de Glace retreated 2400 m since the 1820s LIA maximum while, over the same period, the debris-covered Miage Glacier retreated only 300 m (Deline 2005). Furthermore, this pattern of surface lowering ultimately causes a reduction in the downglacier surface slope, which reduces the driving stress. This causes progressive stagnation (Bolch *et al.* 2008b; Quincey *et al.* 2009) unless increased water availability induces widespread basal sliding of the glacier tongue (Pieczonka *et al.* 2018). Low-angled and stagnating glacier tongues featuring hummocky relief with large terminal moraines means that glacier meltwater cannot be efficiently evacuated through or from the glacier system. The glacier's hydrological network thus transitions from a moderately efficient, linked system to a discontinuous and inefficient network (Benn *et al.* 2017), with consequences for glacier lake formation and associated hazards (Benn *et al.* 2012).

### Development of glacial lakes and implications for hazards

Many debris-covered glaciers have developed proglacial lakes over the past several decades (Fig. 3) (Basnett *et al.* 2013; Racoviteanu *et al.* 2015; Nie *et al.* 2017; Shukla *et al.* 2018). Patterns of thinning and stagnation associated with many debris-covered glaciers suggest the further development of numerous additional glacial lakes is likely to continue over the next few decades (Quincey *et al.* 2007). Large lake formation and increased hazard potential is commonly associated with climate change and glacier recession (e.g. Zheng *et al.* 2021), but other analyses suggest no clear link to climate change (Harrison *et al.* 2018), so the subject remains controversial. A first step towards estimating the consequences of lake growth for hazards is to understand where new lakes will emerge, how large they will become, and how lake levels will change in relation to the surrounding landscape elements. It is important to recognize that lake expansion by itself is not the main criterion that renders a lake hazardous, and that lake elevation changes may be even more important than areal changes.

The current fundamental theory of debris cover evolution and lake development is heavily based on a few well-documented examples, notably in the Everest area of Nepal (Benn *et al.* 2012). During a period of sustained negative mass balance, a debris-covered glacier with large impounding moraines in this region (Fig. 4, ELA1) is expected to first undergo an upward expansion of the debris cover in response to a rise in the equilibrium line altitude (ELA), following which it will undergo a period of downwasting, stagnation and supraglacial pond formation (Fig. 4, ELA2) before fully stagnating at the terminus and forming a terminal lake into which the glacier terminus calves (Fig. 4, ELA3). Each of these stages is linked to several factors, notably specific mass balance and hydrological conditions. In the first stage, the rate of debris cover expansion is not solely related to the rise in ELA, but also conditioned by pre-existing debris content and changing supply rates. In the second stage, surface downwasting of the hummocky surface, coupled with inefficient meltwater evacuation leads to storage of water in perched lakes. It has been established by field studies and remote sensing techniques that the formation of supraglacial lakes is coupled to sustained negative glacier mass balance, substantial historical surface lowering and glacier stagnation towards the snout. Supraglacial lakes tend to occur primarily where the surface slope is less than 2° (Reynolds 2000; Quincey *et al.* 2007; Sakai and Fujita 2010; Linsbauer *et al.* 2016; Pandit and Ramsankaran 2020). The third stage (Fig. 4, ELA3) is marked by the coalescence of



**Fig. 4.** Stages in the evolution of the surface and equilibrium line altitude of a Himalayan debris-covered glacier. Adapted from Benn *et al.* (2012). Credit: Gareth Evans.

supraglacial lakes to form a large ice-contact lake at the englacial water table; during this regime, glacier mass losses at the terminus are strongly governed by calving and water-driven ablation processes within this ice-contact lake. This process has been documented in detail for two sites in the Nepal Himalaya: Imja Tsho (Watanabe *et al.* 1995, 2009) and Tsho Rolpa (Reynolds 1999; Sakai *et al.* 2000a). Numerous proto-lake systems have been identified at other glacier termini using remote sensing, e.g. Ngozumpa Glacier (Thompson *et al.* 2012). While the processes controlling pond expansion are well studied (e.g. Mertes *et al.* 2017), the rates of surface pond expansion and coalescence are not well understood and can change over time (e.g. Thompson *et al.* 2016). Further studies to determine if existing lakes contain buried subaqueous ice would be helpful in constraining lake deepening and basin volume over time. Such studies can be based on comparison of contemporary glacier lake bathymetry with historical ice thickness, in conjunction with studies of the sedimentation rates within lakes.

A further understanding of the glacier overdeepenings and slope conditions that may favour the formation of lakes and therefore may impose some controls on maximum lake volume requires accurate knowledge of debris-covered ice thickness. Consensus estimates of global ice thickness (e.g. Farinotti *et al.* 2017, 2021) developed within the framework of the Working Group on Glacier Ice Thickness Estimation of the International Association of Cryospheric Sciences (ITMIX) (<http://cryosphericciences.org/>) are a valuable addition, but their appropriateness for debris-covered glaciers is unclear.

## Strategies for assessing the hazard potential of a glacial lake

### Key concepts and terminology

Having described the debris-covered glacier landsystem and its key components, in this section we turn our attention to the concept of hazard associated with these glaciers. We focus on glaciers that have receded from their terminal moraines and where moraine-dammed glacio-terminal lakes are created as they recede, because this is the projected end-member state for many debris-covered glaciers under future climate warming. If these moraine-dammed lakes drain rapidly because of dam failure or over-topping, a GLOF can occur, with potentially damaging consequences for downstream populations and infrastructure. Developing a robust framework for describing and assessing the potential glacial hazards associated with moraine-dammed lakes is therefore an important and societally relevant issue.

First of all, any discussion of hazard assessment requires a clear understanding of the terminology used. A common area of confusion is over usage of terms such as ‘hazard’, ‘risk’ and ‘vulnerability’, or ‘hazard assessment’ and ‘risk management’. *Hazards* are defined as potentially damaging physical events or

phenomena which may cause the loss of life or injury, property damage, social and economic disruption, or environmental degradation. *Vulnerability* refers to a set of conditions and processes resulting from physical, social, economic and environmental factors, which increase the susceptibility of a community to the impact of hazard. *Risk* implies the probability of harmful consequences or expected loss (of lives, people injured, property, livelihoods, economic activity disrupted or environment damaged) resulting from actions between natural or human-induced hazards and vulnerable/capable conditions. ‘*Resilience*’ is defined as the capacity of a system, community or society to resist or to change in order that it may obtain an acceptable level in functioning and structure, and ‘*capacity*’ as the way people and organizations use existing resources to achieve various beneficial ends during unusual, abnormal and adverse conditions of a disaster event or process. Conventionally, ‘*risk*’ is expressed as  $\text{risk} = \text{hazards} \times \text{vulnerability/capacity}$  (United Nations 2002); thus, ‘*risk management*’ implies ways in which the hazard might be mitigated as well as increasing the resilience of an affected community. On the other hand, ‘*hazard assessment*’ focuses on the initial physical processes involved with the hazardous situation, such as the triggering and development of a breach in a moraine dam. In the following section we specifically focus on ways to assess the hazard potential of a glacier lake system.

### Glacier lake hazard assessment methods at multiple scales

When assessing the hazard of a glacial lake, besides a solid understanding of the glacial landsystem as detailed earlier, it is fundamental to understand the components of a glacial lake system at the catchment scale and how each of those components behaves in response to the triggering of a GLOF. To fully assess the potential of a glacial lake hazard, it is there important to evaluate lake-specific factors, processes and dynamics of lakes at different stages of glacier lake development in relation to potential triggers in the surrounding landscape. As noted earlier, just because a glacial lake may contain a large volume of water, this does not necessarily make it inherently hazardous. Other factors within the glacier ‘system’ such as the range of landforms, possible mass movement processes, and other influencing factors from within the glacier system environment and the surrounding mountain flanks from the top of the headwall to the lowest terminal moraine dam need to be thoroughly evaluated. This requires a holistic overview of the glacial system in order to identify key components that, if present, may trigger one or more processes that might lead to the formation of a GLOF. The goal is to identify key components that may trigger one or more potentially cascading processes that might lead to a GLOF event.

When assessing the glacier lake hazard potential, two important issues exist: (a) how to assess the hazard across a region in a consistent and meaningful way and (b) how to rank them in terms of the severity of the hazard (Reynolds 2014) in a systematic,

quantitative manner. In quantifying GLOF hazard, remote sensing techniques have been used to develop Tier 1 (first-pass) assessments over large areas (tens to hundreds of square kilometres) (Kääb *et al.* 2005; Quincey *et al.* 2005). Such first-pass automated assessment schemes have been developed for the Tibetan Plateau (Allen *et al.* 2019), the Indian Himalayas (Dubey and Goyal 2020), the Andes (Frey *et al.* 2018; Kougkoulos *et al.* 2018) and the European Alps (Huggel *et al.* 2004). For Tier 2 local assessments of specific glacial lake systems, very high-resolution imagery (< 1 m spatial resolution) and associated DEMs have been used for small areas (e.g. 25–100 km<sup>2</sup> or more); drones have been used to produce very high-resolution imagery and photogrammetry for this purpose (Westoby *et al.* 2012; Fugazza *et al.* 2018; Wilson *et al.* 2019). The UAV and terrestrial structure from motion (SfM) photogrammetry techniques bridge the gap between the difficult field campaigns and the coarse satellite data, and emerged in the last decade as a promising opportunity for estimating hazard potential and hazard management strategies. The results from such analyses can be used to complement or support field campaigns that include, for example, detailed geomorphological, geophysical, topographical and engineering geological surveying and mapping (Hambrey *et al.* 2008). However, a better integration of Tier 1 and Tier 2 assessments is currently needed to assess hazard potentials at multi-scales.

**The requirement for a standardized lake ranking scheme**


Despite technical guidelines on the assessment of glacier and permafrost hazards in mountain regions published by the International Association of Cryospheric Sciences and International Permafrost Association Standing Group on Glacier and Permafrost Hazards (Allen *et al.* 2017), a standard lake hazard assessment scheme does not exist. The existing glacial lake ranking schemes (e.g. Quincey *et al.* 2007; Bolch *et al.* 2008a; Wang *et al.* 2011; Worni *et al.* 2013; Iribarren Anacona *et al.* 2014; Rounce *et al.* 2016; Aggarwal *et al.* 2017; Kougkoulos *et al.* 2018; Dubey and Goyal 2020; Pandit and Ramsankaran 2020) all differ based on the parameters used, the weight assigned to each parameter and the source of the data used (field/remote sensing/a priori knowledge) (Emmer and Vilimek 2013; Rounce *et al.* 2016). Furthermore, existing schemes do not always parameterize key GLOF processes.

Consequently, there is interest in developing a standard, objective unified ranking scheme on the basis of new remote sensing data. Such a scheme would ideally be decision-based, constructed on multi-criteria and using state-of-the-art techniques such as machine learning. Furthermore, such a scheme needs to quantify both observable conditioning and triggering factors related to GLOF formation rather than on subjective criteria or derived parameters, such as lake volume. There are four threshold factors that can be used to categorize any given glacial lake system, but which on their own do not designate the existence of any hazard (Table 4). These have been designed to be used especially as a Tier 1 preliminary screening/hazard ranking tool. However, for a hazard to exist, there must be potential for a trigger event to occur that can lead to a possible GLOF. The key factors affecting the likelihood of a GLOF include: (a) minimal moraine freeboard above the lake level with narrow dam width, rendering the dam vulnerable to overtopping; (b) evidence of avalanches from valley sides and backwall, and/or from hanging glaciers directly into the lake that might induce either a seiche or avalanche push wave; (c) evidence of seepage and/or piping through the moraine dam; (d) evidence of degradation of an ice core within the terminal moraine dam that might cause progressive collapse (RGSL 2003).

For both threshold and trigger parameters, scale factors can be used to weigh how important or significant any factor is. In general, to derive a hazard score, each threshold parameter (Table 4) is ranked using one value from each of the weighting columns and

**Table 4.** Trigger potential and threshold parameters for glacial hazard assessment (modified from RGSL 2003; Reynolds 2014)

Parameter affecting hazard Score	Parameter affecting hazard Score				
	0	2	10	30	50
<b>Threshold factors</b>					
1 Effective volume of lake water available for flood	N/A	Low	Moderate	Large	Very large
2 Height of freeboard relative to lake level	No dam	Very high	High	Moderate	Low
3 Width/height ratio of terminal moraine dam	$w \gg h$	$w > h$	$w \approx h$	$w < h$	$w \ll h$
4 Gradient of distal moraine dam	Flat-5°	5–10°	10–25°	25–40°	>40°
<b>Triggering factors</b>					
5 Height of glacier ice cliff and calving potential	No cliff	Low	Moderate	High	Very high
6 Ice/rock avalanche into lake	Open basin	Low	Moderate	High	Very high
7 Thermokarst degradation of ice within terminal moraine	No ice	Low	Moderate	High	Very high
8 Buoyancy of submerged stagnant ice based on possible ice volume	No ice	Low	Moderate	High	Very high



**Fig. 5.** Example plot of the output of a hazard ranking scheme shown from the analysis of 41 glacial lakes in the Pumqu catchment in Tibet.

summed; similarly, a trigger parameter score (Table 4) is similarly derived. This enables a hazard score to be derived using the weightings for both threshold and trigger parameters. For example, to account for relative differences in lake volume, which is often a derived value based on lake area, measured areas are used. The two scores for the threshold and trigger parameters are used as  $(x, y)$  coordinates to plot on a hazard ranking graph (see Fig. 5).

### *Towards an integrated geohazard assessment*

In addition to glacier and lake processes occurring upstream, a full hazard assessment scheme needs to include the impact on the populations downstream, and a full socio-economic assessment (Carey *et al.* 2012). In recent years there have been advances in both GLOF modelling and integration of models with robust assessments of glacial hazards and their societal impacts. For example, losses incurred from hydropower schemes following GLOFs has led the international hydropower sector to build greater resilience to climate change impacts (RGSL 2015; Reynolds 2018). The complexity of such damaging events in triggering mechanisms and in the changing processes as they propagate downstream calls for catchment-wide assessments of such geohazards. The challenge is that modelling the GLOF impact downstream requires sophisticated flow modelling which implies a number of assumptions about the characteristics of the material, lake volume, peak discharge, sediment load, channel roughness, etc. (Fig. 6) which are difficult to measure (Iribaren Anacona *et al.* 2015). In the last decade, multiple studies have tested and deployed a variety of modelling tools to perform numerical simulations of GLOFs downstream and to simulate different type of flows (Westoby *et al.* 2015). Numerical modelling approaches that coupled glacial lake impact, dam breach and flood processes are reviewed in Worni *et al.* (2014). One of the shortcomings of current models is that flow characteristics are complex and commonly develop as a cascade of physical processes as the flow propagates downstream. This poses the need for modelling multi-phase GLOF process cascades (e.g. Schneider *et al.* 2014; Worni *et al.* 2014; Mergili *et al.* 2020). Furthermore, the extreme flows are difficult to measure for calibration purposes, which entails a large degree of uncertainty (Worni *et al.* 2014).


Given the large uncertainties associated with the GLOF process chain in terms of timing, location and intensity of triggers (Schneider *et al.* 2014; Allen *et al.* 2017), one of the key remaining

challenges is how best to communicate the changing nature of hazards (and implications for GLOF model uncertainty) to communities/stakeholders. Finally, one aspect of hazard and risk assessment that is now well established in the private sector but less so in the academic world is exposure to legal responsibility and the consequential liability arising from making statements about risk that could have outcomes affecting asset values.

### **Remaining challenges and limitations**

Even with the substantial progress on mapping of the debris cover and associated lakes, there remain significant challenges to be addressed in terms of approaching the landsystem using a holistic approach in view of developing a standard hazard ranking scheme. Here we summarize the remaining limitations and gaps in our knowledge of the system.

- **Mapping of debris-covered glaciers** often relies on expert knowledge, which is subjective and often subject to disagreement, especially when independent, ground truth data are not available. There is no standardized mapping for debris-covered glaciers, and existing methods are generally 'semi-automated' because they involve some manual correction (Racoviteanu *et al.* 2009; Herreid and Pellicciotti 2020). While providing important information, available global or automated methods are only suited to mapping debris within assumed glacier extents (Scherler *et al.* 2018). Some of the new methods for debris cover mapping have the potential to automate the mapping process of debris-covered glacier tongues, but these need further and robust testing. There remain significant gaps in high-quality debris cover outlines in some glacierized regions, and retrieval of most key debris properties from remote sensing is at a very early stage;
- **Within the debris-covered glacier landsystem, debris sourcing and evacuation** is a key gap in knowledge; understanding of erosion rates varies regionally, but most erosion rates are millennial-scale values. Thus, further advances need to be made to assess contemporary and recent erosion and debris supply rates within debris-covered glacier landsystems. This might include the use of fine-resolution imagery, derivation of debris supply from avalanche cones,



**Fig. 6.** Elements of a hazardous moraine-dammed glacial lake showing the key stages of a glacier lake outburst flood: (1) propagation of displacement or seiche waves in the lake, and/or piping through the dam; (2) breach initiation and breach formation; (3) propagation of resultant flood wave(s) down-valley. Key triggers are labelled A to F: (A) glacier calving; (B) icefall from hanging glaciers; (C) rock/ice/snow avalanches; (D) dam settlement and/or piping; (E) ice-cored moraine degradation; (F) rapid input of water from supra-, en-, or subglacial (including subaqueous) sources. Conditioning factors are labelled a to d: increased lake volume, low dam width to height ratio, ice degradation, minimal freeboard (modified from Richardson and Reynolds 2000; Westoby *et al.* 2014).

and other creative analysis or numerical modelling approaches (e.g. Banerjee and Wani 2018; Scherler and Egholm 2020). It is key to study both singular, large debris supply events (e.g. Berthier and Brun 2019), and smaller events of debris supply using holistic efforts. Furthermore, debris flux out of the system is so far only crudely represented despite being a key property governing glacier development over time, and it may be valuable to identify the determinants of whether or not a glacier forms a large latero-terminal moraine;

- **The issue of scale in remote sensing remains an important challenge.** While significant progress has been made in monitoring debris cover surface properties using both field methods and remote sensing, these are often applied at different spatial scales (local to regional), making it difficult to transfer the observations from one scale to the other. Detailed field studies offer insight into specific processes (e.g. ice ablation), but they are often site-specific; remote sensing studies, on the other hand, can be applied at multi-scales, but face limitations due to spatial and temporal resolution, i.e. lack of high-resolution thermal data or surface velocities.
- **The implications of increased debris supply remain unclear, for example:**
  - How do glacier thermal and dynamic regimes respond to the increased debris resulting from glacier thinning and upwards migration of debris? What are the implications of the increased debris for basal sliding, glacier thinning and stagnation, ice thickness and deformation?
  - What are the typical glacial structures associated with increased debris supply, and what are the consequences of these structures downglacier? How will these glaciers respond to changes in terms of hydrology, ice deformation and surface debris?
  - What are the consequences of different levels of debris sequestration on glacial landscape evolution and geomorphology? How will subglacial erosion, moraine building, lake development and sedimentation change through time?
  - How do permafrost and debris-covered/rock glaciers interact with the glacier(s) and debris/mass fluxes through the system?
  - How does the increased debris supply influence the formation of proglacial and supraglacial lakes impounded

by lateral or terminal moraines and by supra-glacial debris deposits? How does this affect the probability of glacier-related hazards (lake outbursts floods and debris slides) under a transient climate?

- **The rates at which the system transitions to different states** are unclear, and proxies to project them forward in time are needed to accommodate them in glacier model projections. For example, more work is needed in order to make projections of rated of debris cover expansion/thickening from an initial state of unknown debris load in the system, and with uncertain debris inputs/outputs. There is a need for improved models (i) to reasonably account for the meltwater production role of transient features such as ice cliffs, (ii) to project glacier downwasting and surface slope evolution to the threshold of supraglacial pond development and (iii) to parameterize rates of supraglacial pond expansion to allow the likely timing of pond coalescence to be estimated.
- **The development and testing of a standardized, integrated lake hazard ranking scheme** remains a challenge. This requires better parameterization of key GLOF processes in the glacier lake system, and the ability to capture the multi-phase GLOF process cascade.

In order to address these questions and to complete the picture of processes associated with the debris-cover glacier system, improved datasets are still needed, i.e. meteorological data from weather stations installed at high altitudes (e.g. Matthews *et al.* 2020), monitoring of the rates and controls on rockwall debris supply, gauging of water and sediment amounts discharged within turbid glacier streams (e.g. Heckmann *et al.* 2016), spatially distributed measurements of ice thickness from new technologies such as airborne ice-sounding radar suited to debris-covered glaciers (e.g. Pritchard *et al.* 2020) and debris thickness distributions (e.g. Nicholson *et al.* 2018) with which to optimized models of debris thickness, studies of the sediment discharge to the glacier foreland high-resolution regular-repeat imagery for selected debris-covered glacier landsystems with differing characteristics. There is a need for more studies that link climate, mass balance, ice dynamics, topographic evolution and hydrology to quantify how hazard potential emerges from interactions between these processes. Finally, there is a need to bridge spatial scales both in terms of

connecting processes and resolving them in remotely sensed data. For example, some satellite imagery cannot resolve metre-scale features, even though features such as ice cliffs at this scale may collectively be significant to runoff generation, and the melt processes operating locally at ice cliffs need to be integrated into a glacier scale representation of the ablation regime (Ferguson and Vieli 2021).

## Conclusions and outlook

This paper stems from a workshop supported by the Geological Society of London in 2019, that brought together researchers with a shared interest in debris-covered glaciers and related hazards with a broad range of experience and activities in approaching these landscape systems. As such, this perspectives paper draws together key insights, state-of-the-art and consensus research priorities from the exchanges fostered by the workshop. While the key state of the knowledge has been described in the preceding sections, to conclude we wish to draw out a small number of key messages.

When considering debris-covered glaciers, we argue that it is vital to adopt a landsystems approach that includes the flux of both solid and liquid water and sediment within catchments as well as estimates of how these processes influence and are influenced by glacier behaviour over time. Despite key developments and advances in the use of satellite remote sensing to estimate these processes, there remain gaps in the validation of these tools using field-based measurements as these remain scarce. There remains much work to be done to develop robust tools to upscale the knowledge gained from small process studies to a landsystems scale so that it can be integrated in satellite monitoring and numerical models of larger spatio-temporal scale glacier and landscape development.

Debris-covered glaciers are projected to increase in number proportionately as mountain glaciers diminish, but the specific trajectories of glacier development are elusive due to the complex coupling of non-stationary processes and feedbacks within the debris-covered glacier system. Critically, some glaciers form large impounding latero-terminal moraines that drive local hydrological processes and which have implications for glacier hazards, while others do not, and we lack a clear method of discriminating which pathway a given glacier or glacierized region will follow.

Finally, we suggest that consideration of cascading hazards within the wider landsystem is critical for developing meaningful glacial lake hazard assessment. There is a need to address this issue due to communication failures in the past, so a better interaction between the debris-covered glacier community and the geomorphological and climate science communities is needed for this perspective framework to be successful.

**Acknowledgements** We thank the Geological Society for their support in hosting the ‘Debris-covered Glaciers and Related Lakes: Understanding the Challenges’ workshop held in London in September 2019. We thank all the speakers in the workshop: T. Bolch, J. Carrivick, E. Miles, D. Quincey, J. Reynolds, M. Westoby and S. Watson for their insights during the workshop discussions, and all workshop participants for their efforts and enthusiasm.

**Author contributions** **AER**: conceptualization (lead), investigation (equal), writing – original draft (lead), writing – review & editing (equal); **LN**: conceptualization (equal), writing – original draft (equal), writing – review & editing (supporting); **NFG**: project administration (lead), supervision (lead), writing – review & editing (supporting); **EM**: conceptualization (supporting), methodology (supporting), writing – original draft (equal), writing – review & editing (equal); **SH**: writing – review & editing (supporting); **JMR**: writing – review & editing (supporting)

**Funding** AR’s research was supported by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 663830 and SCoRE Cymru. JR was supported by Reynolds

International Ltd. LN was supported by Austrian Science Fund (FWF) Grant P28521.

**Competing interests** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Data availability** Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

*Scientific editing by Philip Hughes*

## References

- Aggarwal, S., Rai, S.C., Thakur, P.K. and Emmer, A. 2017. Inventory and recently increasing GLOF susceptibility of glacial lakes in Sikkim, Eastern Himalaya. *Geomorphology*, **295**, 39–54, <https://doi.org/10.1016/j.geomorph.2017.06.014>
- Alifu, H., Tateishi, R. and Johnson, B. 2015. A new band ratio technique for mapping debris-covered glaciers using Landsat imagery and a digital elevation model. *International Journal of Remote Sensing*, **36**, 2063–2075, <https://doi.org/10.1080/2150704X.2015.1034886>
- Allen, S., Frey, H. and Huggel, C. 2017. Assessment of Glacier and Permafrost Hazards in Mountain Regions. *Technical Guidance Document*. <https://doi.org/10.13140/RG.2.2.26332.90245>
- Allen, S.K., Zhang, G., Wang, W., Yao, T. and Bolch, T. 2019. Potentially dangerous glacial lakes across the Tibetan Plateau revealed using a large-scale automated assessment approach. *Sci. Bull.*, **64**, 435–445, <https://doi.org/10.1016/j.scib.2019.03.011>
- Anderson, R.S. 2000. A model of ablation-dominated medial moraines and the generation of debris-mantled glacier snouts. *Journal of Glaciology*, **46**, 459–469, <https://doi.org/10.3189/172756500781833025>
- Anderson, L.S. and Anderson, R.S. 2016. Modeling debris-covered glaciers: response to steady debris deposition. *The Cryosphere*, **10**, 1105–1124, <https://doi.org/10.5194/tc-10-1105-2016>
- Anderson, L. and Anderson, R. 2018. Debris thickness patterns on debris-covered glaciers. *Geomorphology*, **311**, <https://doi.org/10.1016/j.geomorph.2018.03.014>
- Anderson, B. and Mackintosh, A. 2012. Controls on mass balance sensitivity of maritime glaciers in the Southern Alps, New Zealand: The role of debris cover. *Journal of Geophysical Research: Earth Surface*, **117**, <https://doi.org/10.1029/2011JF002064>
- Anderson, K., Fawcett, D., Cugulliere, A., Benford, S., Jones, D. and Leng, R. 2020. Vegetation expansion in the subnival Hindu Kush Himalaya. *Global Change Biology*, **26**, 1608–1625, <https://doi.org/10.1111/gcb.14919>
- Anderson, L.S., Armstrong, W.H., Anderson, R.S. and Buri, P. 2021. Debris cover and the thinning of Kennicott Glacier, Alaska: in situ measurements, automated ice cliff delineation and distributed melt estimates. *The Cryosphere*, **15**, 265–282, <https://doi.org/10.5194/tc-15-265-2021>
- Ballantyne, C.K. 2002. Paraglacial geomorphology. *Quaternary Science Reviews*, **21**, 1935–2017, [https://doi.org/10.1016/S0277-3791\(02\)00005-7](https://doi.org/10.1016/S0277-3791(02)00005-7)
- Banerjee, A. and Shankar, R. 2013. On the response of Himalayan glaciers to climate change. *Journal of Glaciology*, **59**, 480–490, <https://doi.org/10.3189/2013JoG12J130>
- Banerjee, A. and Wani, B.A. 2018. Exponentially decreasing erosion rates protect the high-elevation crests of the Himalaya. *Earth and Planetary Science Letters*, **497**, 22–28, <https://doi.org/10.1016/j.epsl.2018.06.001>
- Bartlett, O.T., Ng, F.S.L. and Rowan, A.V. 2021. Morphology and evolution of supraglacial hummocks on debris-covered Himalayan glaciers. *Earth Surface Processes and Landforms*, **46**, 525–539, <https://doi.org/10.1002/esp.5043>
- Basnett, S., Kulkarni, A. and Bolch, T. 2013. The influence of debris cover and glacial lakes on the recession of glaciers in Sikkim Himalaya, India. *Journal of Glaciology*, **59**, 1035–1046, <https://doi.org/10.3189/2013JoG12J184>
- Benn, D.I. and Evans, D.J.A. 1998. *Glaciers and glaciations*. John Wiley & Sons, Inc., New York.
- Benn, D.I. and Lehmkuhl, F. 2000. Mass balance and equilibrium-line altitudes of glaciers in high-mountain environments. *Quaternary International*, **65–66**, 15–29, [https://doi.org/10.1016/S1040-6182\(99\)00034-8](https://doi.org/10.1016/S1040-6182(99)00034-8)
- Benn, D.I. and Owen, L.A. 2002. Himalayan glacial sedimentary environments: a framework for reconstructing and dating the former extent of glaciers in high mountains. *Quaternary Int.*, **97–98**, 3–25, [https://doi.org/10.1016/S1040-6182\(02\)00048-4](https://doi.org/10.1016/S1040-6182(02)00048-4)
- Benn, D.I., Wiseman, S. and Hands, K.A. 2001. Growth and drainage of supraglacial lakes on debris-mantled Ngozumpa Glacier, Khumbu Himal, Nepal. *Journal of Glaciology*, **47**, 626–638, <https://doi.org/10.3189/172756501781831729>
- Benn, D., Kirkbride, M.P., Owen, L. and Brazier, V. 2004. Glaciated valley landsystems. In: Evans, D.A.J. (ed.) *Glacial landsystems*, Arnold, 372–406.
- Benn, D.I., Bolch, T., Aaa, A.A., et al. 2012. Response of debris-covered glaciers in the Mount Everest region to recent warming, and implications for outburst flood hazards. *Earth Science Reviews*, **114**, 156–174, <https://doi.org/10.1016/j.earscirev.2012.03.008>

- Benn, D., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L. 2017. Structure and evolution of the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss. *The Cryosphere*, **11**, 2247–2264, <https://doi.org/10.5194/tc-11-2247-2017>
- Berthier, E. and Brun, F. 2019. Karakoram geodetic glacier mass balances between 2008 and 2016: persistence of the anomaly and influence of a large rock avalanche on Siachen Glacier. *Journal of Glaciology*, **65**, 494–507, <https://doi.org/10.1017/jog.2019.32>
- Berthier, E., Raup, B. and Scambos, T. 2003. New velocity map and mass-balance estimate of Mertz Glacier, East Antarctica, derived from Landsat sequential imagery. *Journal of Glaciology*, **49**, 503–511, <https://doi.org/10.3189/172756503781830377>
- Berthier, E., Vadon, H., Aaa, A.A., *et al.* 2005. Surface motion of mountain glaciers derived from satellite optical imagery. *Remote Sensing of Environment*, **95**, 14–28, <https://doi.org/10.1016/j.rse.2004.11.005>
- Berthier, E., Schiefer, E., Clarke, G., Menounos, B. and Rémy, F. 2010. Contribution of Alaskan glaciers to sea-level rise derived from satellite imagery. *Nature Geoscience*, **3**, 92–95, <https://doi.org/10.1038/ngeo737>
- Bhambri, R., Bolch, T. and Chaujar, R.K. 2011. Mapping of Debris-covered Glaciers in the Garhwal Himalayas using ASTER DEMs and Thermal Data. *International Journal of Remote Sensing*, **32**, 8095–8119, <https://doi.org/10.1080/01431161.2010.532821>
- Bishop, M.P., Bonk, R., Kamp, U. and Schroder, J.F., J. 2001. Terrain analysis and data modeling for alpine glacier mapping. *Polar Geogr.*, **25**, 182–201, <https://doi.org/10.1080/10889370109377712>
- Bisset, R.R., Dehecq, A., Goldberg, D.N., Huss, M., Bingham, R.G. and Goumelen, N. 2020. Reversed Surface-Mass-Balance Gradients on Himalayan Debris-Covered Glaciers Inferred from Remote Sensing. *Remote Sensing*, **12**, <https://doi.org/10.3390/rs12101563>
- Bolch, T., Buchroithner, M.F., Kunert, A. and Kamp, U. 2007. Automated delineation of debris-covered glaciers based on ASTER data. Geoinformation in Europe (Proc. of 27th EARSeL Symposium, 04–07 June 2007), Bozen, Italy, 403–410.
- Bolch, T., M.F. Buchroithner, J. Peters, M. Baessler and Bajracharya, S. 2008a. Identification of glacier motion and potentially dangerous glacial lakes in the Mt. Everest region/Nepal using spaceborne imagery. *Nat. Hazards Earth Syst. Sci.*, **8**, 1329–1340, <https://doi.org/10.5194/nhess-8-1329-2008>
- Bolch, T., Buchroithner, M.F., Pieczonka, T. and Kunert, A. 2008b. Planimetric and Volumetric Glacier Changes in the Khumbu Himalaya since 1962 Using Corona, Landsat TM and ASTER Data. *J. Glaciol.*, **54**, 592–600, <https://doi.org/10.3189/002214308786570782>
- Bolch, T., Pieczonka, T. and Benn, D.I. 2011. Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery. *The Cryosphere*, **5**, 349–358, <https://doi.org/10.5194/tc-5-349-2011>
- Bolch, T., Kulkarni, A., Aaa, A.A., *et al.* 2012. The State and Fate of Himalayan Glaciers. *Science*, **336**, 310, <https://doi.org/10.1126/science.1215828>
- Boxall, K., Willis, I., Giese, A. and Liu, Q. 2021. Quantifying Patterns of Supraglacial Debris Thickness and Their Glaciological Controls in High Mountain Asia. *Frontiers in Earth Science*, **9**, 504, <https://doi.org/10.3389/feart.2021.657440>
- Brun, F., Buri, P., Aaa, A.A., *et al.* 2016. Quantifying volume loss from ice cliffs on debris-covered glaciers using high-resolution terrestrial and aerial photogrammetry. *Journal of Glaciology*, **62**, 684–695, <https://doi.org/10.1017/jog.2016.54>
- Brun, F., Wagnon, P., Aaa, A.A., *et al.* 2018. Ice cliff contribution to the tongue-wide ablation of Changri Nup Glacier, Nepal, central Himalaya. *The Cryosphere*, **12**, 3439–3457, <https://doi.org/10.5194/tc-12-3439-2018>
- Brun, F., Wagnon, P., Berthier, E., Jomelli, V., Maharjan, S.B., Shrestha, F. and Kraaijenbrink, P.D.A. 2019. Heterogeneous Influence of Glacier Morphology on the Mass Balance Variability in High Mountain Asia. *Journal of Geophysical Research: Earth Surface*, **124**, 1331–1345, <https://doi.org/10.1029/2018JF004838>
- Buri, P. and Pellicciotti, F. 2018. Aspect controls the survival of ice cliffs on debris-covered glaciers. *Proceedings of the National Academy of Sciences*, **115**, 4369, <https://doi.org/10.1073/pnas.1713892115>
- Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P. and Pellicciotti, F. 2016. A physically based 3-D model of ice cliff evolution over debris-covered glaciers. *Journal of Geophysical Research: Earth Surface*, **121**, 2471–2493, <https://doi.org/10.1002/2016JF004039>
- Buri, P., Miles, E.S., Steiner, J.F., Ragetti, S. and Pellicciotti, F. 2021. Supraglacial Ice Cliffs Can Substantially Increase the Mass Loss of Debris-Covered Glaciers. *Geophysical Research Letters*, **48**, e2020GL092150, <https://doi.org/10.1029/2020GL092150>
- Carey, M., Huggel, C., Bury, J., Portocarrero, C. and Haeblerli, W. 2012. An integrated socio-environmental framework for glacier hazard management and climate change adaptation: lessons from Lake 513, Cordillera Blanca, Peru. *Climatic Change*, **112**, 733–767, <https://doi.org/10.1007/s10584-011-0249-8>
- Carrivick, J.L. and Tweed, F.S. 2013. Proglacial lakes: character, behaviour and geological importance. *Quaternary Science Reviews*, **78**, 34–52, <https://doi.org/10.1016/j.quascirev.2013.07.028>
- Casey, K. and Kääh, A. 2012. Estimation of Supraglacial Dust and Debris Geochemical Composition via Satellite Reflectance and Emissivity. *Remote Sensing*, **4**, 2554–2575, <https://doi.org/10.3390/rs4092554>
- Casey, K.A., Kääh, A. and Benn, D.I. 2012. Geochemical characterization of supraglacial debris via in situ and optical remote sensing methods: a case study in Khumbu Himalaya, Nepal. *The Cryosphere*, **6**, 85–100, <https://doi.org/10.5194/tc-6-85-2012>
- Chen, W., Fukui, H., Doko, T. and Gu, X. 2013. Improvement of glacial lakes detection under shadow environment using ASTER data in Himalayas, Nepal. *Chinese Geographical Science*, **23**, 216–226, <https://doi.org/10.1007/s11769-012-0584-3>
- Chen, F., Zhang, M., Guo, H., Allen, S., Kargel, J.S., Haritashya, U.K. and Watson, C.S. 2020. Annual 30-meter Dataset for Glacial Lakes in High Mountain Asia from 2008 to 2017. *Earth Syst. Sci. Data Discuss.*, **2020**, 1–29, <https://doi.org/10.5194/essd-2020-57>
- Cogley, J.G., Kargel, J.S., Kaser, G. and van der Vee, C.J. 2010. Tracking the Source of Glacier Misinformation. *Science*, **327**, 522, <https://doi.org/10.1126/science.327.5965.522-a>
- Collier, E., Nicholson, L.L., Brock, B.W., Maussion, F., Essery, R. and Bush, A.B.G. 2014. Representing moisture fluxes and phase changes in glacier debris cover using a reservoir approach. *The Cryosphere*, **8**, 1429–1444, <https://doi.org/10.5194/tc-8-1429-2014>
- Cook, S. and Quincey, D. 2015. Estimating the volume of Alpine glacial lakes. *Earth Surface Dynamics Discussions*, **3**, 909–940, <https://doi.org/10.5194/esurf-d-3-909-2015>
- Dehecq, A., Goumelen, N. and Trouve, E. 2015. Deriving large-scale glacier velocities from a complete satellite archive: Application to the Pamir-Karakoram-Himalaya. *Remote Sensing of Environment*, **162**, 55–66, <https://doi.org/10.1016/j.rse.2015.01.031>
- Dehecq, A., Goumelen, N., Aaa, A.A., *et al.* 2019. Twenty-first century glacier slowdown driven by mass loss in High Mountain Asia. *Nature Geoscience*, **12**, 22–27, <https://doi.org/10.1038/s41561-018-0271-9>
- Deline, P. 2005. Change in surface debris cover on Mont Blanc massif glaciers after the ‘Little Ice Age’ termination. *The Holocene*, **15**, 302–309, <https://doi.org/10.1191/0959683605hl809rr>
- Deline, P. 2009. Interactions between rock avalanches and glaciers in the Mont Blanc massif during the late Holocene. *Quaternary Science Reviews*, **28**, 1070–1083, <https://doi.org/10.1016/j.quascirev.2008.09.025>
- Deline, P., Hewitt, K., Reznichenko, N. and Shugar, D. 2015. Rock Avalanches onto Glaciers. In: Shroder, J.F. and Davies, T. (eds) *Landslide Hazards, Risks and Disasters*, Academic Press, Boston, 263–319, <https://doi.org/10.1016/B978-0-12-396452-6.00009-4>
- Detert, M. and Weitbrecht, V. 2020. Determining image-based grain size distribution with suboptimal conditioned photos. 1045–1052, <https://doi.org/10.1201/b22619-146>
- Dubey, S. and Goyal, M.K. 2020. Glacial Lake Outburst Flood Hazard, Downstream Impact, and Risk Over the Indian Himalayas. *Water Resources Research*, **56**, e2019WR026533, <https://doi.org/10.1029/2019WR026533>
- Emmer, A. and Vilimek, V. 2013. Review Article: Lake and breach hazard assessment for moraine-dammed lakes: an example from the Cordillera Blanca (Peru). *Natural Hazards and Earth System Sciences*, **13**, 1551–1565, <https://doi.org/10.5194/nhess-13-1551-2013>
- Evatt, G.W., Mayer, C., Mallinson, A.M.Y., Abrahams, I.D., Heil, M. and Nicholson, L. 2017. The secret life of ice sails. *Journal of Glaciology*, **63**, 1049–1062, <https://doi.org/10.1017/jog.2017.72>
- Farinotti, D., Brinkerhoff, D.J., Aaa, A.A., *et al.* 2017. How accurate are estimates of glacier ice thickness? Results from ITMIX, the Ice Thickness Models Intercomparison eXperiment. *The Cryosphere*, **11**, 949–970, <https://doi.org/10.5194/tc-11-949-2017>
- Farinotti, D., Brinkerhoff, D., Aaa, A.A., *et al.* 2021. Results from the Ice Thickness Models Intercomparison eXperiment Phase 2 (ITMIX2). *Frontiers in Earth Science*, **8**, <https://doi.org/10.3389/feart.2020.571923>
- Ferguson, J. and Vieli, A. 2020. Modelling steady states and the transient response of debris-covered glaciers. *The Cryosphere Discuss.*, **2020**, 1–31, <https://doi.org/10.5194/tc-2020-228>
- Ferguson, J.C. and Vieli, A. 2021. Modelling steady states and the transient response of debris-covered glaciers. *The Cryosphere*, **15**, 3377–3399, <https://doi.org/10.5194/tc-15-3377-2021>
- Fickert, T., Friend, D., Grüninger, F., Molnia, B. and Richter, M. 2007. Did Debris-Covered Glaciers Serve as Pleistocene Refugia for Plants? A New Hypothesis Derived from Observations of Recent Plant Growth on Glacier Surfaces. *Arctic, Antarctic, and Alpine Research*, **39**, 245–257, [https://doi.org/10.1657/1523-0430\(2007\)39\[245:DDGSAP\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2007)39[245:DDGSAP]2.0.CO;2)
- Foster, L.A., Brock, B.W., Cutler, M.E.J. and Diotri, F. 2012. A physically based method for estimating supraglacial debris thickness from thermal band remote-sensing data. *Journal of Glaciology*, **58**, 677–690, <https://doi.org/10.3189/2012JoG11J194>
- Frey, H., Paul, F. and Strozzi, T. 2012. Compilation of a glacier inventory for the western Himalayas from satellite data: methods, challenges, and results. *Remote Sensing of Environment*, **124**, 832–843, <https://doi.org/10.1016/j.rse.2012.06.020>
- Frey, H., Huggel, C., Chisolm, R.E., Baer, P., McArdell, B., Cochachin, A. and Portocarrero, C. 2018. Multi-Source Glacial Lake Outburst Flood Hazard Assessment and Mapping for Huaraz, Cordillera Blanca, Peru. *Frontiers in Earth Science*, **6**, <https://doi.org/10.3389/feart.2018.00210>
- Fugazza, D., Scaioni, M., Aaa, A.A., *et al.* 2018. Combination of UAV and terrestrial photogrammetry to assess rapid glacier evolution and map glacier hazards. *Nat. Hazards Earth Syst. Sci.*, **18**, 1055–1071, <https://doi.org/10.5194/nhess-18-1055-2018>
- Fujii, Y. 1977. Field Experiment on Glacier Ablation under a Layer of Debris Cover Glaciological Expedition of Nepal, Contribution No. 33. *Journal of the*

- Japanese Society of Snow and Ice, **39**, 20–21, [https://doi.org/10.5331/seppy.39.Special\\_20](https://doi.org/10.5331/seppy.39.Special_20)
- Fyffe, C.L., Brock, B.W., Aa, A.A., et al. 2019b. Do debris-covered glaciers demonstrate distinctive hydrological behaviour compared to clean glaciers? *Journal of Hydrology*, **570**, 584–597, <https://doi.org/10.1016/j.jhydrol.2018.12.069>
- Fyffe, C.L., Brock, B.W., Kirkbride, M.P., Black, A.R., Smiraglia, C. and Diolaiuti, G. 2019a. The impact of supraglacial debris on proglacial runoff and water chemistry. *Journal of Hydrology*, **576**, 41–57, <https://doi.org/10.1016/j.jhydrol.2019.06.023>
- Fyffe, C.L., Woodget, A.S., Kirkbride, M.P., Deline, P., Westoby, M.J. and Brock, B.W. 2020. Processes at the margins of supraglacial debris cover: Quantifying dirty ice ablation and debris redistribution. *Earth Surface Processes and Landforms*, **45**, 2272–2290, <https://doi.org/10.1002/esp.4879>
- Gardelle, J., Arnaud, Y. and Berthier, E. 2011. Contrasted evolution of glacial lakes along the Hindu Kush Himalaya mountain range between 1990 and 2009. *Global and Planetary Change*, **75**, <https://doi.org/10.1016/j.gloplacha.2010.10.003>
- Gardelle, J., Berthier, E., Arnaud, Y. and Kääb, A. 2013. Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999–2011. *The Cryosphere*, **7**, 1263–1286, <https://doi.org/10.5194/tc-7-1263-2013>
- Gardner, A.S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., van den Broeke, M. and Nilsson, J. 2018. Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years. *The Cryosphere*, **12**, 521–547, <https://doi.org/10.5194/tc-12-521-2018>
- Giese, A., Boone, A., Wagon, P. and Hawley, R. 2020. Incorporating moisture content in surface energy balance modeling of a debris-covered glacier. *The Cryosphere*, **14**, 1555–1577, <https://doi.org/10.5194/tc-14-1555-2020>
- GLAMOS 2020. *Swiss Glacier Inventory 2016, release 2020*.
- Greuell, W. and Oerlemans, J. 2004. Narrowband-to-broadband albedo conversion for glacier ice and snow: equations based on modeling and ranges of validity of the equations. *Remote Sensing of Environment*, **89**, 95–105, <https://doi.org/10.1016/j.rse.2003.10.010>
- Gulley, J. and Benn, D.I. 2007. Structural control of englacial drainage systems in Himalayan debris-covered glaciers. *Journal of Glaciology*, **53**, 399–412, <https://doi.org/10.3189/002214307783258378>
- Gulley, J.D., Benn, D.I., Müller, D. and Luckman, A. 2009a. A cut-and-closure origin for englacial conduits in uncrevassed regions of polythermal glaciers. *Journal of Glaciology*, **55**, 66–80, <https://doi.org/10.3189/002214309788608930>
- Gulley, J.D., Benn, D.I., Sreaton, E. and Martin, J. 2009b. Mechanisms of englacial conduit formation and their implications for subglacial recharge. *Quaternary Science Reviews*, **28**, 1984–1999, <https://doi.org/10.1016/j.quascirev.2009.04.002>
- Hagg, W., Mayer, C., Lambrecht, A. and Helm, A. 2008. Sub-debris melt rates on Southern Inylchek Glacier, central Tian Shan. *Geografiska Annaler: Series A, Physical Geography*, **90**, 55–63, <https://doi.org/10.1111/j.1468-0459.2008.00333.x>
- Hambrey, M.J., Quincey, D.J., Glasser, N.F., Reynolds, J.M., Richardson, S.J. and Clemmens, S. 2008. Sedimentological, geomorphological and dynamic context of debris-mantled glaciers, Mount Everest (Sagarmatha) region, Nepal. *Quaternary Science Reviews*, **27**, 2361–2389, <https://doi.org/10.1016/j.quascirev.2008.08.010>
- Han, H., Wang, J., Junfeng, W. and Liu, S. 2010. Backwasting rate on debris-covered Koxkar glacier, Tuomuer mountain, China. *Journal of Glaciology*, **56**, 287–296, <https://doi.org/10.3189/002214310791968430>
- Harithashya, U.K., Pleasants, M.S. and Copland, L. 2015. Assessment of the evolution in velocity of two debris-covered valley glaciers in nepal and new zealand. *Geografiska Annaler: Series A, Physical Geography*, **97**, 737–751, <https://doi.org/10.1111/geoa.12112>
- Harithashya, U.K., Kargel, J.S., Aa, A.A., et al. 2018. Evolution and Controls of Large Glacial Lakes in the Nepal Himalaya. *Remote Sensing*, **10**, <https://doi.org/10.3390/rs10050798>
- Harrison, S. 2009. Climate sensitivity: implications for the response of geomorphological systems to future climate change. *Geological Society, London, Special Publications*, **320**, 257, <https://doi.org/10.1144/SP320.16>
- Harrison, S., Kargel, J.S., Aa, A.A., et al. 2018. Climate change and the global pattern of moraine-dammed glacial lake outburst floods. *The Cryosphere*, **12**, 1195–1209, <https://doi.org/10.5194/tc-12-1195-2018>
- Heckmann, T., McColl, S. and Morche, D. (eds) 2016. The Geomorphology, Dynamism, and Significance of Proglacial Environments in the 21st Century. *Earth Surface Processes and Landforms*, **41**, 271–276., <https://doi.org/10.1002/esp.3858>
- Heimsath, A. and McGlynn, R. 2008. Quantifying periglacial erosion in the Nepal high Himalaya. *Geomorphology*, **97**, 5–23, <https://doi.org/10.1016/j.geomorph.2007.02.046>
- Herreid, S. and Pellicciotti, F. 2018. Automated detection of ice cliffs within supraglacial debris cover. *The Cryosphere*, **12**, 1811–1829, <https://doi.org/10.5194/tc-12-1811-2018>
- Herreid, S. and Pellicciotti, F. 2020. The state of rock debris covering Earth's glaciers. *Nature Geoscience*, **13**, 621–627, <https://doi.org/10.1038/s41561-020-0615-0>
- Hewitt, K. 2009. Rock avalanches that travel onto glaciers and related developments, Karakoram Himalaya, Inner Asia. *Geomorphology*, **103**, 66–79, <https://doi.org/10.1016/j.geomorph.2007.10.017>
- Hewitt, K. and Shroder, J. 1993. *Himalaya to the Sea: Geology, Geomorphology, and the Quaternary*. Routledge, London, UK.
- Hock, R. 2003. Temperature index melt modelling in mountain areas. *J Hydrol*, **282**, 104–115, [https://doi.org/10.1016/S0022-1694\(03\)00257-9](https://doi.org/10.1016/S0022-1694(03)00257-9)
- Holobacă, I.-H., Tielidze, L.G., et al. 2021. Multi-sensor remote sensing to map glacier debris cover in the Greater Caucasus, Georgia. *Journal of Glaciology*, **67**, 685–696, <https://doi.org/10.1017/jog.2021.47>
- Huang, L., Li, Z., Tian, B.S., Han, H.D., Liu, Y.Q., Zhou, J.M. and Chen, Q. 2017. Estimation of supraglacial debris thickness using a novel target decomposition on L-band polarimetric SAR images in the Tianshan Mountains. *Journal of Geophysical Research: Earth Surface*, **122**, 925–940, <https://doi.org/10.1002/2016JF004102>
- Huggel, C., Haeblerli, W., Kääb, A., Bieri, D. and Richardson, S. 2004. An assessment procedure for glacial hazards in the Swiss Alps. *Canadian Geotechnical Journal*, **41**, 1068–1083, <https://doi.org/10.1139/t04-053>
- Hugonnet, R., McNabb, R., Aa, A.A., et al. 2021. Accelerated global glacier mass loss in the early twenty-first century. *Nature*, **592**, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>
- Huo, D., Bishop, M.P. and Bush, A.B.G. 2021. Understanding Complex Debris-Covered Glaciers: Concepts, Issues, and Research Directions. *Frontiers in Earth Science*, **9**, 358.
- Huss, M. 2011. Present and future contribution of glacier storage change to runoff from macroscale drainage basins in Europe. *Water Resources Research*, **47**, <https://doi.org/10.1029/2010WR010299>
- Immerzeel, W.W., van Beek, L.P.H., Konz, M., Shrestha, A.B. and Bierkens, M.F.P. 2012. Hydrological response to climate change in a glacierized catchment in the Himalayas. *Climatic Change*, **110**, 721–736, <https://doi.org/10.1007/s10584-011-0143-4>
- Iribarren Anacona, P., Norton, K.P. and Mackintosh, A. 2014. Moraine-dammed lake failures in Patagonia and assessment of outburst susceptibility in the Baker Basin. *Natural Hazards and Earth System Sciences*, **14**, 3243–3259, <https://doi.org/10.5194/nhess-14-3243-2014>
- Iribarren Anacona, P., Mackintosh, A. and Norton, K.P. 2015. Hazardous processes and events from glacier and permafrost areas: lessons from the Chilean and Argentinean Andes. *Earth Surf. Procc Land.*, **40**, 2–21, <https://doi.org/10.1002/esp.3524>
- Irvine-Fynn, T.D.L., Porter, P.R., Aa, A.A., et al. 2017. Supraglacial ponds regulate runoff from Himalayan debris-covered glaciers. *Geophysical Research Letters*, **44**, 11894–11904, <https://doi.org/10.1002/2017GL075398>
- Iwata, S., Aoki, T., Kadota, T., Seko, K. and Yamaguchi, S. 2000. Morphological evolution of the debris cover on Khumbu Glacier, Nepal, between 1978 and 1995. In: Nakawo, M., Raymond, C.F. and Fountain, A. (eds) *Debris-Covered Glaciers*. 3–11, IAHS.
- Jones, D.B., Harrison, S. and Anderson, K. 2019. Mountain glacier-to-rock glacier transition. *Global and Planetary Change*, **181**, 102999, <https://doi.org/10.1016/j.gloplacha.2019.102999>
- Juen, M., Mayer, C., Lambrecht, A., Wirbel, A. and Kueppers, U. 2013. Thermal properties of a supraglacial debris layer with respect to lithology and grain size. *Geografiska Annaler: Series A, Physical Geography*, **95**, 197–209, <https://doi.org/10.1111/geoa.12011>
- Juen, M., Mayer, C., Lambrecht, A., Han, H. and Liu, S. 2014. Impact of varying debris cover thickness on ablation: a case study for Koxkar Glacier in the Tien Shan. *The Cryosphere*, **8**, 377–386, <https://doi.org/10.5194/tc-8-377-2014>
- Kääb, A. 2005. Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow velocities in the Bhutan Himalaya. *Remote Sensing of Environment*, **94**, 463–474, <https://doi.org/10.1016/j.rse.2004.11.003>
- Kääb, A., Huggel, C., Aa, A.A., et al. 2005. Remote sensing of glacier- and permafrost-related hazards in high mountains: an overview. *Natural Hazards and Earth System Sciences*, **5**, 527–554, <https://doi.org/10.5194/nhess-5-527-2005>
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y. 2012. Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas. *Nature*, **488**, 495–498, <https://doi.org/10.1038/nature11324>
- Kamp, U., Byrne, M. and Bolch, T. 2011. Glacier fluctuations between 1975 and 2008 in the Greater Himalaya Range of Zaskar, southern Ladakh. *Journal of Mountain Sciences*, **8**, 374–389, <https://doi.org/10.1007/s11629-011-2007-9>
- Kargel, J.S., Bush, A.B.G., Aa, A.A., et al. 2014. A world of changing glaciers: Summary and climatic context. In: Kargel, J.S., Leonard, G.J., Bishop, M.P., Kääb, A. and Raup, B.H. (eds) *Global Land Ice Measurements from Space*. Springer Berlin Heidelberg, Berlin, Heidelberg, 781–840, [https://doi.org/10.1007/978-3-540-79818-7\\_33](https://doi.org/10.1007/978-3-540-79818-7_33)
- Kaser, G., Grosshauser, M. and Marzeion, B. 2010. Contribution potential of glaciers to water availability in different climate regimes. *Proceedings of the National Academy of Sciences*, **107**, 20223–20227, <https://doi.org/10.1073/pnas.1008162107>
- Kayastha, R.B., Takeuchi, Y., Nakawo, M. and Ageta, Y. 2000. Practical prediction of ice melting beneath various thickness of debris cover on Khumbu Glacier, Nepal, using a positive degree-day factor. In: Raymond, C.F., Nakawo, M., Fountain, A. (eds) *Debris-Covered Glaciers*. IAHS, Wallingford, UK, 71–81.
- Khan, A.A., Jamil, A., Hussain, D., Taj, M., Jabeen, G. and Malik, M.K. 2020. Machine-Learning Algorithms for Mapping Debris-Covered Glaciers: The Hunza Basin Case Study. *IEEE Access*, **8**, 12725–12734, <https://doi.org/10.1109/ACCESS.2020.2965768>
- King, O., Bhattacharya, A., Bhabri, R. and Bolch, T. 2019. Glacial lakes exacerbate Himalayan glacier mass loss. *Scientific Reports*, **9**, 18145, <https://doi.org/10.1038/s41598-019-53733-x>

- King, O., Turner, A.G.D., Quincey, D.J. and Carrivick, J.L. 2020a. Morphometric evolution of Everest region debris-covered glaciers. *Geomorphology*, **371**, 107422, <https://doi.org/10.1016/j.geomorph.2020.107422>
- King, O., Bhattacharya, A., Ghuffar, S., Tait, A., Guilford, S., Elmore, A.C. and Bolch, T. 2020b. Six Decades of Glacier Mass Changes around Mt. Everest Are Revealed by Historical and Contemporary Images. *One Earth*, **3**, 608–620, <https://doi.org/10.1016/j.oneear.2020.10.019>
- Kirkbride, M. 1989. About the concepts of continuum and age. *Boreas*, **18**, 87–88, <https://doi.org/10.1111/j.1502-3885.1989.tb00376.x>
- Kirkbride, M.P. 2011. Debris-Covered Glaciers. In: Singh, V.P., Singh, P. and Haritashya, U.K. (eds) *Encyclopedia of Snow, Ice and Glaciers*. Springer Netherlands, Dordrecht, 180–182, [https://doi.org/10.1007/978-90-481-2642-2\\_622](https://doi.org/10.1007/978-90-481-2642-2_622)
- Kirkbride, M.P. and Deline, P. 2013. The formation of supraglacial debris covers by primary dispersal from transverse englacial debris bands. *Earth Surface Processes and Landforms*, **38**, 1779–1792, <https://doi.org/10.1002/esp.3416>
- Knap, W.H., Reijmer, C.H. and Oerlemans, J. 1999. Narrowband to broadband conversion of Landsat TM glacier albedos. *International Journal of Remote Sensing*, **20**, 2091–2110, <https://doi.org/10.1080/014311699212362>
- Kneib, M., Miles, E.S., Aaa, A.A., *et al.* 2020. Mapping ice cliffs on debris-covered glaciers using multispectral satellite images. *Remote Sensing of Environment*, **112201**, <https://doi.org/10.1016/j.rse.2020.112201>
- Kneib, M., Miles, E.S., Buri, P., Molnar, P., McCarthy, M., Fugger, S. and Pellicciotti, F. 2021. Interannual Dynamics of Ice Cliff Populations on Debris-Covered Glaciers From Remote Sensing Observations and Stochastic Modeling. *Journal of Geophysical Research: Earth Surface*, **126**, e2021JF006179, <https://doi.org/10.1029/2021JF006179>
- Knight, J. and Harrison, S. 2014. Mountain Glacial and Paraglacial Environments under Global Climate Change: Lessons from the Past, Future Directions and Policy Implications. *Geografiska Annal Phys Geogr*, **96**, 245–264, <https://doi.org/10.1111/geoa.12051>
- Knight, J. and Harrison, S. 2018. Transience in cascading paraglacial systems. *Land Degradation & Development*, **29**, 1991–2001, <https://doi.org/10.1002/ldr.2994>
- Knight, J., Harrison, S. and Jones, D.B. 2019. Rock glaciers and the geomorphological evolution of deglaciating mountains. *Geomorphology*, **324**, 14–24, <https://doi.org/10.1016/j.geomorph.2018.09.020>
- Komori, J. 2008. Recent expansions of glacial lakes in the Bhutan Himalayas. *Quaternary International*, **184**, 177–186, <https://doi.org/10.1016/j.quaint.2007.09.012>
- Korup, O. and Tweed, F. 2007. Ice, moraine, and landslide dams in mountainous terrain. *Quaternary Science Reviews*, **26**, 3406–3422, <https://doi.org/10.1016/j.quascirev.2007.10.012>
- Koukoulos, I., Cook, S., *et al.* 2018. Use of multi-criteria decision analysis to identify potentially dangerous glacial lakes. *Science of The Total Environment*, **621**, 1453–1466, <https://doi.org/10.1016/j.scitotenv.2017.10.083>
- Kraaijenbrink, P.D.A., Shea, J.M., Pellicciotti, F., Jong, S.M.D. and Immerzeel, W.W. 2016. Object-based analysis of unmanned aerial vehicle imagery to map and characterise surface features on a debris-covered glacier. *Remote Sensing of Environment*, **186**, 581–595, <https://doi.org/10.1016/j.rse.2016.09.013>
- Kraaijenbrink, P.D.A., Bierkens, M.F.P., Lutz, A.F. and Immerzeel, W.W. 2017. Impact of a global temperature rise of 1.5 degrees Celsius on Asia's glaciers. *Nature*, **549**, 257–260, <https://doi.org/10.1038/nature23878>
- Kraaijenbrink, P.D.A., Shea, J.M., Litt, M., Steiner, J.F., Treichler, D., Koch, I. and Immerzeel, W.W. 2018. Mapping Surface Temperatures on a Debris-Covered Glacier With an Unmanned Aerial Vehicle. *Frontiers in Earth Science*, **6**, <https://doi.org/10.3389/feart.2018.00064>
- Kumar, V., Venkataramana, G. and Høgdal, K.A. 2011. Glacier surface velocity estimation using SAR interferometry technique applying ascending and descending passes in Himalayas. *International Journal of Applied Earth Observation and Geoinformation*, **13**, 545–551, <https://doi.org/10.1016/j.jag.2011.02.004>
- Leclercq, P.W., Oerlemans, J. and Cogley, J.G. 2011. Estimating the Glacier Contribution to Sea-Level Rise for the Period 1800–2005. *Surveys in Geophysics*, **32**, 519, <https://doi.org/10.1007/s10712-011-9121-7>
- Leprince, S., Ayoub, F., Klingler, Y. and Avouac, J. 2007. Co-Registration of Optically Sensed Images and Correlation (COSI-Corr): an operational methodology for ground deformation measurements. IEEE International Geoscience and Remote Sensing Symposium, 23–28 July 2007, Barcelona, Spain, 1943–1946.
- Liang, S. 2001. Narrowband to broadband conversions of land surface albedo I: Algorithms. *Remote Sensing of Environment*, **76**, 213–238, [https://doi.org/10.1016/S0034-4257\(00\)00205-4](https://doi.org/10.1016/S0034-4257(00)00205-4)
- Linsbauer, A., Frey, H., Haerberli, W., Machguth, H., Azam, M.F. and Allen, S. 2016. Modelling glacier-bed overdeepenings and possible future lakes for the glaciers in the Himalaya—Karakoram region. *Annals of Glaciology*, **57**, 119–130, <https://doi.org/10.3189/2016AoG71A627>
- Linsbauer, A., Hodel, E., Huss, M., Bauder, A., Fischer, M., Weidmann, Y. and Bärtschi, H. 2020. The new Swiss Glacier Inventory SGI2020: From a topographic to a glaciological dataset. presented at the EGU General Assembly 2020, 4–8 May 2020, online.
- Lipp, S., Vijay, S. and Braun, M. 2018. Automatic delineation of debris-covered glaciers using InSAR coherence derived from X-, C- and L-band radar data: a case study of Yazgyl Glacier. *Journal of Glaciology*, **64**, 811–821, <https://doi.org/10.1017/jog.2018.70>
- Liu, Q., Mayer, C. and Liu, S. 2015. Distribution and interannual variability of supraglacial lakes on debris-covered glaciers in the Khan Tengri-Tumour Mountains, Central Asia. *Environmental Research Letters*, **10**: 014011–014010, <https://doi.org/10.1088/1748-9326/10/1/014011>
- Mancini, D. and Lane, S.N. 2020. Changes in sediment connectivity following glacial debuitressing in an Alpine valley system. *Geomorphology*, **352**, 106987, <https://doi.org/10.1016/j.geomorph.2019.106987>
- Marston, R.A. 1983. Supraglacial Stream Dynamics on the Juneau Icefield. *Annals of the Association of American Geographers*, **73**, 597–608, <https://doi.org/10.1111/j.1467-8306.1983.tb01861.x>
- Marzeion, B., Jarosch, A.H. and Hofer, M. 2012. Past and future sea-level change from the surface mass balance of glaciers. *The Cryosphere*, **6**, 1295–1322, <https://doi.org/10.5194/tc-6-1295-2012>
- Marzeion, B., Cogley, J.G., Richter, K. and Parkes, D. 2014. Attribution of global glacier mass loss to anthropogenic and natural causes. *Science*, **345**, 919, <https://doi.org/10.1126/science.1254702>
- Matta, E., Giardino, C., Boggero, A. and Bresciani, M. 2017. Use of Satellite and In Situ Reflectance Data for Lake Water Color Characterization in the Everest Himalayan Region. *Mountain Research and Development*, **37**, 16–23, <https://doi.org/10.1659/MRD-JOURNAL-D-15-00052.1>
- Matthews, T., Perry, L.B., Aaa, A.A., *et al.* 2020. Going to Extremes: Installing the World's Highest Weather Stations on Mount Everest. *Bulletin of the American Meteorological Society*, **101**, E1870–E1890, <https://doi.org/10.1175/BAMS-D-19-0198.1>
- Mattson, L.E., Gardner, J.S. and Young, G.J. 1993. Ablation on debris covered glaciers: an example from the Rakhiot Glacier, Panjab, Himalaya. In: Young, G.J. (ed.) *Snow and Glacier Hydrology*. Proceedings of the International Symposium, 16–21 November 1992, Kathmandu, Nepal. IAHS Publication.
- Maurer, J.M., Schaefer, J.M., Rupper, S. and Corley, A. 2019. Acceleration of ice loss across the Himalayas over the past 40 years. *Science Advances*, **5**, eaav7266, <https://doi.org/10.1126/sciadv.aav7266>
- Mayer, C., Lambrecht, A., Belò, M., Smiraglia, C. and Diolaiuti, G. 2006. Glaciological characteristics of the ablation zone of Baltoro glacier, Karakoram, Pakistan. *Annals of Glaciology*, **43**, 123–131, <https://doi.org/10.3189/172756406781812087>
- McCarthy, M., Pritchard, H., Willis, I.A.N. and King, E. 2017. Ground-penetrating radar measurements of debris thickness on Lirung Glacier, Nepal. *Journal of Glaciology*, **63**, 543–555, <https://doi.org/10.1017/jog.2017.18>
- Mergili, M., Pudasaini, S.P., Emmer, A., Fischer, J.T., Cochachin, A. and Frey, H. 2020. Reconstruction of the 1941 GLOF process chain at Lake Palcacocha (Cordillera Blanca, Peru). *Hydrol. Earth Syst. Sci.*, **24**, 93–114, <https://doi.org/10.5194/hess-24-93-2020>
- Mertes, J., Thompson, S., Booth, A., Gullely, D.J. and Benn, D. 2016. A conceptual model of supraglacial lake formation on debris-covered glaciers based on GPR facies analysis: GPR facies analysis of Spillway Lake, Ngozumpa Glacier, Nepal. *Earth Surface Processes and Landforms*, **42**, 903–914, <https://doi.org/10.1002/esp.4068>
- Mertes, J.R., Thompson, S.S., Booth, A.D., Gullely, J.D. and Benn, D.I. 2017. A conceptual model of supra-glacial lake formation on debris-covered glaciers based on GPR facies analysis. *Earth Surface Processes and Landforms*, **42**, 903–914, <https://doi.org/10.1002/esp.4068>
- Mihalcea, C., Mayer, C., Diolaiuti, G., Lambrecht, A., Smiraglia, C. and Tartari, G. 2006. Ice ablation and meteorological conditions on the debris-covered area of Baltoro glacier, Karakoram, Pakistan. *Annals of Glaciology*, **43**, 292–300, <https://doi.org/10.3189/172756406781812104>
- Mihalcea, C., Mayer, C., Aaa, A.A., *et al.* 2008. Spatial distribution of debris thickness and melting from remote-sensing and meteorological data, at debris-covered Baltoro glacier, Karakoram, Pakistan. *Annals of Glaciology*, **48**, 49–57, <https://doi.org/10.3189/172756408784700680>
- Miles, E.S., Pellicciotti, F., Willis, I.C., Steiner, J.F., Buri, P. and Arnold, N.S. 2016. Refined energy-balance modelling of a supraglacial pond, Langtang Khola, Nepal. *Annals of Glaciology*, **57**, 29–40, <https://doi.org/10.3189/2016AoG71A421>
- Miles, E.S., I.C. Willis, N.S. Arnold, Steiner, J. and Pellicciotti, F. 2017a. Spatial, seasonal and interannual variability of supraglacial ponds in the Langtang Valley of Nepal, 1999–2013. *Journal of Glaciology*, **63**, 88–105, <https://doi.org/10.1017/jog.2016.120>
- Miles, E.S., Steiner, J., Willis, I., Buri, P., Immerzeel, W.W., Chesnokova, A. and Pellicciotti, F. 2017b. Pond Dynamics and Supraglacial-Englacial Connectivity on Debris-Covered Lirung Glacier, Nepal. *Frontiers in Earth Science*, **5**, <https://doi.org/10.3389/feart.2017.00069>
- Miles, E., Watson, C.S., Aaa, A.A., *et al.* 2018a. Glacial and geomorphic effects of a supraglacial lake drainage and outburst event, Everest region, Nepal Himalaya. *The Cryosphere*, **12**, 3891–3905, <https://doi.org/10.5194/tc-12-3891-2018>
- Miles, E.S., Willis, I., Buri, P., Steiner, J.F., Arnold, N.S. and Pellicciotti, F. 2018b. Surface Pond Energy Absorption Across Four Himalayan Glaciers Accounts for 1/8 of Total Catchment Ice Loss. *Geophysical Research Letters*, **45**, 464–473, <https://doi.org/10.1029/2018GL079678>
- Miles, K.E., Hubbard, B., Quincey, D.J., Miles, E.S., Irvine-Fynn, T.D.L. and Rowan, A.V. 2019. Surface and subsurface hydrology of debris-covered Khumbu Glacier, Nepal, revealed by dye tracing. *Earth and Planetary Science Letters*, **513**, 176–186, <https://doi.org/10.1016/j.epsl.2019.02.020>
- Miles, K.E., Hubbard, B., Irvine-Fynn, T.D.L., Miles, E.S., Quincey, D.J. and Rowan, A.V. 2020. Hydrology of debris-covered glaciers in High Mountain

- Asia. *Earth-Science Reviews*, **207**, 103212, <https://doi.org/10.1016/j.earscirev.2020.103212>
- Millan, R., Mouginot, J. *et al.* 2019. Mapping Surface Flow Velocity of Glaciers at Regional Scale Using a Multiple Sensors Approach. *Remote Sensing*, **11**, <https://doi.org/10.3390/rs11212498>
- Mölg, N., Bolch, T., Walter, A. and Vieli, A. 2019. Unravelling the evolution of Zmuttgletscher and its debris cover since the end of the Little Ice Age. *The Cryosphere*, **13**, 1889–1909, <https://doi.org/10.5194/tc-13-1889-2019>
- Mölg, N., Ferguson, J., Bolch, T. and Vieli, A. 2020. On the influence of debris cover on glacier morphology: How high-relief structures evolve from smooth surfaces. *Geomorphology*, **357**, 107092, <https://doi.org/10.1016/j.geomorph.2020.107092>
- Monnier, S. and Kinnard, C. 2017. Pluri-decadal (1955–2014) evolution of glacier–rock glacier transitional landforms in the central Andes of Chile (30–33°S). *Earth Surf. Dynam.*, **5**, 493–509, <https://doi.org/10.5194/esurf-5-493-2017>
- Mool, P.K., Bajracharya, S.R., Joshi, S.P., Sakya, K. and Baidya, A. 2002. *Inventory of Glaciers, Glacial Lakes and GLOF Monitoring [...] Nepal*. International Center for Integrated Mountain Development, Nepal.
- Moore, P.L. 2018. Stability of supraglacial debris. *Earth Surface Processes and Landforms*, **43**, 285–297, <https://doi.org/10.1002/esp.4244>
- Naegeli, K., Damm, A., Huss, M., Wulf, H., Schaepman, M. and Hoelzle, M. 2017. Cross-Comparison of Albedo Products for Glacier Surfaces Derived from Airborne and Satellite (Sentinel-2 and Landsat 8) Optical Data. *Remote Sensing*, **9**, <https://doi.org/10.3390/rs9020110>
- Nagai, H., Fujita, K., Nuimura, T. and Sakai, A. 2013. Southwest-facing slopes control the formation of debris-covered glaciers in the Bhutan Himalaya. *The Cryosphere*, **7**, 1303–1314, <https://doi.org/10.5194/tc-7-1303-2013>
- Nakawo, M. and Young, G.J. 1982. Estimate of glacier ablation under a debris layer from surface temperature and meteorological variables. *Journal of Glaciology*, **28**, 29–34, <https://doi.org/10.1017/S002214300001176X>
- Nakawo, M., Iwata, S., Watanabe, O. and Yoshida, M. 1986. Processes which Distribute Supraglacial Debris on the Khumbu Glacier, Nepal Himalaya. *Annals of Glaciology*, **8**, 129–131, <https://doi.org/10.3189/S0260305500001294>
- Narama, C., Daiyrov, M., Tadono, T., Yamamoto, M., Käab, A., Morita, R. and Ukita, J. 2017. Seasonal drainage of supraglacial lakes on debris-covered glaciers in the Tien Shan Mountains, Central Asia. *Geomorphology*, **286**, 133–142, <https://doi.org/10.1016/j.geomorph.2017.03.002>
- Nicholson, L. and Benn, D.I. 2006. Calculating ice melt beneath a debris layer using meteorological data. *Journal of Glaciology*, **52**, 463–470, <https://doi.org/10.3189/172756506781828584>
- Nicholson, L. and Benn, D.I. 2012. Properties of natural supraglacial debris in relation to modelling sub-debris ice ablation. *Earth Surface Processes and Landforms*, **38**, 490–501, <https://doi.org/10.1002/esp.3299>
- Nicholson, L. and Mertes, J. 2017. Thickness estimation of supraglacial debris above ice cliff exposures using a high-resolution digital surface model derived from terrestrial photography. *Journal of Glaciology*, **63**, 989–998, <https://doi.org/10.1017/jog.2017.68>
- Nicholson, L.I., McCarthy, M., Pritchard, H.D. and Willis, I. 2018. Supraglacial debris thickness variability: impact on ablation and relation to terrain properties. *The Cryosphere*, **12**, 3719–3734, <https://doi.org/10.5194/tc-12-3719-2018>
- Nicholson, L.I., Wirbel, A., Mayer, C. and Lambrecht, A. in press. The challenge of non-stationary feedbacks in the response of debris-covered glaciers to climate forcing. *Frontiers in Earth Science*, <https://doi.org/10.3389/feart.2021.662695>
- Nie, Y., Sheng, Y., Liu, Q., Liu, L., Liu, S., Zhang, Y. and Song, C. 2017. A regional-scale assessment of Himalayan glacial lake changes using satellite observations from 1990 to 2015. *Remote Sensing of Environment*, **189**, 1–13, <https://doi.org/10.1016/j.rse.2016.11.008>
- Noh, M.-J. and Howat, I.M. 2015. Automated stereo-photogrammetric DEM generation at high latitudes: Surface Extraction with TIN-based Search-space Minimization (SETSM) validation and demonstration over glaciated regions. *GIScience & Remote Sensing*, **52**, 198–217, <https://doi.org/10.1080/15481603.2015.1008621>
- Nuimura, T., Fujita, K., Yamaguchi, S. and Sharma, R.R. 2012. Elevation changes of glaciers revealed by multitemporal digital elevation models calibrated by GPS survey in the Khumbu region, Nepal Himalaya, 1992–2008. *Journal of Glaciology*, **58**, 648–656, <https://doi.org/10.3189/2012JoG11J061>
- Østrem, G. 1959. Ice Melting under a Thin Layer of Moraine, and the Existence of Ice Cores in Moraine Ridges. *Geografiska Annaler*, **41**, 228–230, <https://doi.org/10.1080/20014422.1959.11907953>
- Owen, L.A., Derbyshire, E. and Scott, C.H. 2003. Contemporary sediment production and transfer in high-altitude glaciers. *Sedimentary Geology*, **155**, 13–36, [https://doi.org/10.1016/S0037-0738\(02\)00156-2](https://doi.org/10.1016/S0037-0738(02)00156-2)
- Panday, P., Bulley, H., Haritashya, U. and Ghimire, B. 2011. Supraglacial Lake Classification in the Everest Region of Nepal Himalaya. In: Thakur, J.K., *et al.* (eds) *Geospatial Techniques for Managing Environmental Resources*. Capital Publishing Company, 86–99, [https://doi.org/10.1007/978-94-007-1858-6\\_6](https://doi.org/10.1007/978-94-007-1858-6_6)
- Pandit, A. and Ramsankaran, R. 2020. Identification of Potential Sites for Future Lake Formation and Expansion of Existing Lakes in Glaciers of Chandra Basin, Western Himalayas, India. *Frontiers in Earth Science*, **8**, 382, <https://doi.org/10.3389/feart.2020.500116>
- Pant, S.R. and Reynolds, J.M. 2000. Application of electrical imaging techniques for the investigation of natural dams: an example from the Thulagi Glacier Lake, Nepal. *Journal of Nepal Geological Society*, **22**, <https://doi.org/10.3126/jngs.v22i0.32348>
- Paul, F., Huggel, C. and Käab, A. 2004. Combining satellite multispectral image data and a digital elevation model for mapping debris-covered glaciers. *Rem Sens Environ*, **89**, 510–518, <https://doi.org/10.1016/j.rse.2003.11.007>
- Paul, F., Kaab, A. and Haerberli, W. 2007. Recent glacier changes in the Alps observed by satellite: Consequences for future monitoring strategies. *Global and Planetary Change*, **56**, 111–122, <https://doi.org/10.1016/j.gloplacha.2006.07.007>
- Paul, F., Barrand, N.E., Aaa, A.A., *et al.* 2013. On the accuracy of glacier outlines derived from remote-sensing data. *Annals of Glaciology*, **54**, 171–182, <https://doi.org/10.3189/2013AoG63A296>
- Pellicciotti, F., Stephan, C., Miles, E., Herreid, S., Immerzeel, W.W. and Bolch, T. 2015. Mass-balance changes of the debris-covered glaciers in the Langtang Himal, Nepal, from 1974 to 1999. *Journal of Glaciology*, **61**, 373–386, <https://doi.org/10.3189/2015JoG13J237>
- Pfeffer, W.T., Arendt, A.A., Aaa, A.A., *et al.* 2014. The Randolph Glacier Inventory: a globally complete inventory of glaciers. *Journal of Glaciology*, **60**, <https://doi.org/10.3189/2014JoG131176>
- Pieczonka, T., Bolch, T., Kröhnert, M., Peters, J. and Liu, S. 2018. Glacier branch lines and glacier ice thickness estimation for debris-covered glaciers in the Central Tien Shan. *Journal of Glaciology*, **64**, 835–849, <https://doi.org/10.1017/jog.2018.75>
- Pritchard, H.D., King, E.C., Goodger, D.J., McCarthy, M., Mayer, C. and Kayastha, R. 2020. Towards Bedmap Himalayas: development of an airborne ice-sounding radar for glacier thickness surveys in High-Mountain Asia. *Annals of Glaciology*, **61**, 35–45, <https://doi.org/10.1017/aog.2020.29>
- Quincey, D.J., Lucas, R.M., Richardson, S.D., Glasser, N.F., Hambrey, M.J. and Reynolds, J.M. 2005. Optical remote sensing techniques in high-mountain environments: application to glacial hazards. *Progress in Physical Geography: Earth and Environment*, **29**, 475–505, <https://doi.org/10.1191/0309133305pp456ra>
- Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J. and Glasser, N.F. 2007. Early recognition of glacial lake hazards in the Himalaya using remote sensing datasets. *Global and Planetary Change*, **56**, 137–152, <https://doi.org/10.1016/j.gloplacha.2006.07.013>
- Quincey, D.J., Luckman, A. and Benn, D. 2009. Quantification of Everest region glacier velocities between 1992 and 2002, using satellite radar interferometry and feature tracking. *Journal of Glaciology*, **55**, 596–606, <https://doi.org/10.3189/002214309789470987>
- Racoviteanu, A.E. and Williams, M.W. 2012. Decision tree and texture analysis for mapping debris-covered glaciers: a case study from Kangchenjunga, eastern Himalaya. *Remote Sensing Special Issue*, **4**, 3078–3109, <https://doi.org/10.3390/rs4103078>
- Racoviteanu, A., Arnaud, Y. and Williams, M. 2008. Decadal changes in glacier parameters in Cordillera Blanca, Peru derived from remote sensing. *Journal of Glaciology*, **54**, 499–510, <https://doi.org/10.3189/002214308785836922>
- Racoviteanu, A., Paul, F., Raup, B., Khalsa, S.J.S. and Armstrong, R. 2009. Challenges and recommendations in mapping of glacier parameters from space: results of the 2008 Global Land Ice Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA. *Annals of Glaciology*, **50**, <https://doi.org/10.3189/172756410790595804>
- Racoviteanu, A.E., Arnaud, Y., Williams, M.W. and Manley, W.F. 2015. Spatial patterns in glacier characteristics and area changes from 1962 to 2006 in the Kanchenjunga; Sikkim area, eastern Himalaya. *The Cryosphere*, **9**, 505–523, <https://doi.org/10.5194/tc-9-505-2015>
- Racoviteanu, A., Nicholson, L. and Glasser, N. 2021. Surface composition of debris covered glaciers across the Himalaya using linear spectral unmixing of Landsat 8 data. *The Cryosphere*, **15**, 4557–4588, <https://doi.org/10.5194/tc-15-4557-2021>
- Ragetli, S., Pellicciotti, F., Aaa, A.A., *et al.* 2015. Unraveling the hydrology of a Himalayan catchment through integration of high resolution in situ data and remote sensing with an advanced simulation model. *Advances in Water Resources*, **78**, 94–111, <https://doi.org/10.1016/j.advwatres.2015.01.013>
- Rana, B., Shrestha, A.B., Reynolds, J.M., Aryal, R., Pokhrel, A.P. and Budhathoki, K.P. 2000. Hazard assessment of the Tsho Rolpa Glacier Lake and ongoing remediation measures. *Journal of the Nepal Geological Society*, **22**, 563–570, <https://doi.org/10.3126/jngs.v22i0.32432>
- Ranzi, R., Grossi, G., Iacovelli, L. and Taschner, S. 2004. Use of multispectral ASTER images for mapping debris-covered glaciers within the GLIMS project. IGARSS 2004. 2004 IEEE International Geoscience and Remote Sensing Symposium, 20–24 Sept. 2004, Anchorage, AK, USA, 1144–1147.
- Rashid, I., Majeed, U., Jan, A. and Glasser, N.F. 2020. The January 2018 to September 2019 surge of Shisper Glacier, Pakistan, detected from remote sensing observations. *Geomorphology*, **351**, 106957, <https://doi.org/10.1016/j.geomorph.2019.106957>
- Reid, T.D. and Brock, B.W. 2010. An energy-balance model for debris-covered glaciers including heat conduction through the debris layer. *Journal of Glaciology*, **56**, 903–916, <https://doi.org/10.3189/002214310794457218>
- Reid, T.D. and Brock, B.W. 2014. Assessing ice-cliff backwasting and its contribution to total ablation of debris-covered Miage glacier, Mont Blanc massif, Italy. *Journal of Glaciology*, **60**, 3–13, <https://doi.org/10.3189/2014JoG13J045>
- Reynolds, J.M. 1999. Glacial hazard assessment at Tsho Rolpa, Rolwaling, Central Nepal. *Quarterly Journal of Engineering Geology and Hydrogeology*, **32**, 209–214, <https://doi.org/10.1144/GSL.QJEG.1999.032.P3.01>

- Reynolds, J.M. 2000. On the formation of supraglacial lakes on debris-covered glaciers. In: Nakawo, M., Raymond, C.F. and Fountain, A. (eds) *Debris-Covered Glaciers*. IAHS, Wallingford, 153–161.
- Reynolds, J.M. 2006. Role of geophysics in glacial hazard assessment. *First Break*, **24**, 61–66, <https://doi.org/10.3997/1365-2397.24.8.27068>
- Reynolds, J.M. 2014. Assessing glacial hazards for hydropower development in the Himalayas, Hindu Kush and Karakoram. *The International Journal on Hydropower & Dams*, **21**, 60–65.
- Reynolds, J.M. 2018. Integrated Geohazard Assessments in high mountain environments: examples from the Hindu Kush-Karakoram-Himalayan Region. presented at the ASIA 2018, 13-15 March 2018, Da Nang, Vietnam.
- Reznichenko, N., Davies, T., Shulmeister, J. and McSaveney, M. 2010. Effects of debris on ice-surface melting rates: An experimental study. *Journal of Glaciology*, **56**, <https://doi.org/10.3189/002214310792447725>
- Reznichenko, N.V., Davies, T.R.H. and Alexander, D.J. 2011. Effects of rock avalanches on glacier behaviour and moraine formation. *Geomorphology*, **132**, 327–338, <https://doi.org/10.1016/j.geomorph.2011.05.019>
- RGI Consortium. 2017. *Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0: Technical Report, Global Land Ice Measurements from Space, Colorado, USA*. Digital Media.
- RGSL. 2003. Development of glacial hazard and risk minimisation protocols in rural environments: Guidelines for the management of glacial hazards and risks. *Technical report R7816.142*, Reynolds Geo-Sciences Ltd, Mold, UK, 62 pp. <https://www.reynolds-international.co.uk/services/glacial-and-geological-hazards/dfid/> [last accessed 16 December 2021]
- RGSL. 2015. On the need to integrate Disaster Risk Management within the hydropower sector. *Technical report R7816.142*, Reynolds Geo-Sciences Ltd, Mold, UK.
- Richardson, S. and Reynolds, J. 2000. An overview of glacial hazards in the Himalayas. *Quaternary International*, **65–66**, 31–47, [https://doi.org/10.1016/S1040-6182\(99\)00035-X](https://doi.org/10.1016/S1040-6182(99)00035-X)
- Robson, B.A., Nuth, C., Dahl, S.O., Hölbling, D., Strozzi, T. and Nielsen, P.R. 2015. Automated classification of debris-covered glaciers combining optical, SAR and topographic data in an object-based environment. *Remote Sensing of Environment*, **170**, 372–387, <https://doi.org/10.1016/j.rse.2015.10.001>
- Roe, G.H., Baker, M. B. and Herla, F. 2017. Centennial glacier retreat as categorical evidence of regional climate change. *Nature Geoscience*, **10**, 95–99, <https://doi.org/10.1038/ngeo2863>
- Rounce, D.R. and McKinney, D.C. 2014. Debris thickness of glaciers in the Everest area (Nepal Himalaya) derived from satellite imagery using a nonlinear energy balance model. *The Cryosphere*, **8**, 1317–1329, <https://doi.org/10.5194/tc-8-1317-2014>
- Rounce, D.R., McKinney, D.C., Lala, J.M., Byers, A.C. and Watson, C.S. 2016. A new remote hazard and risk assessment framework for glacial lakes in the Nepal Himalaya. *Hydrol. Earth Syst. Sci.*, **20**, 3455–3475, <https://doi.org/10.5194/hess-20-3455-2016>
- Rounce, D.R., King, O., McCarthy, M., Shean, D.E. and Salerno, F. 2018. Quantifying Debris Thickness of Debris-Covered Glaciers in the Everest Region of Nepal Through Inversion of a Subdebris Melt Model. *Journal of Geophysical Research: Earth Surface*, **123**, 1094–1115, <https://doi.org/10.1029/2017JF004395>
- Rounce, D.R., Hock, R., *et al.* 2021. Distributed Global Debris Thickness Estimates Reveal Debris Significantly Impacts Glacier Mass Balance. *Geophysical Research Letters*, **48**, e2020GL091311, <https://doi.org/10.1029/2020GL091311>
- Rowan, A.V., Egholm, D.L., Quincey, D.J. and Glasser, N.F. 2015. Modelling the feedbacks between mass balance, ice flow and debris transport to predict the response to climate change of debris-covered glaciers in the Himalaya. *Earth and Planetary Science Letters*, **430**, 427–438, <https://doi.org/10.1016/j.epsl.2015.09.004>
- Rowan, A.V., Quincey, D.J., Aaa, A.A., *et al.* 2017. *The sustainability of water resources in High Mountain Asia in the context of recent and future glacier change*. In: Pant, N.C., Ravindra, R., Srivastava, D. and Thompson, L.G. (eds) *The Himalayan Cryosphere: Past and Present*, Geological Society, London, Special Publications, **462**, 189–204, <https://doi.org/10.1144/SP462.12>
- Sakai, A. 2012. Glacial lakes in the Himalayas: a review on formation and expansion processes. *Global Environmental Research*, **16**, 23–30.
- Sakai, A. and Fujita, K. 2010. Correspondence: Formation conditions of supraglacial lakes on debris covered glaciers in the Himalaya. *Journal of Glaciology*, **56**, 177–181, <https://doi.org/10.3189/002214310791190785>
- Sakai, A. and Fujita, K. 2017. Contrasting glacier responses to recent climate change in high-mountain Asia. *Scientific Reports*, **7**, 13717–13717, <https://doi.org/10.1038/s41598-017-14256-5>
- Sakai, A., Chikita, K. and Yamada, T. 2000a. Expansion of a moraine-dammed glacial lake, Tsho Rolpa, in Rolwaling Himal, Nepal Himalaya. *Limnology and Oceanography*, **45**, 1401–1408, <https://doi.org/10.4319/lo.2000.45.6.1401>
- Sakai, A., Takeuchi, N., Fujita, K. and Nakawo, M. 2000b. Role of supraglacial ponds in the ablation process of a debris-covered glacier in the Nepal Himalayas. *International Association of Hydrological Sciences*, **264**, 119–130.
- Sakai, A., Nakawo, M. and Fujita, K. 2002. Distribution Characteristics and Energy Balance of Ice Cliffs on Debris-Covered Glaciers, Nepal Himalaya. *Arctic Antarctic and Alpine Research*, **34**, 12–19, <https://doi.org/10.1080/15230430.2002.12003463>
- Salerno, F., Thakuri, S., D'Agata, C., Smiraglia, C., Manfredi, E.C., Viviano, G. and Tartari, G. 2012. Glacial lake distribution in the Mount Everest region: Uncertainty of measurement and conditions of formation. *Global and Planetary Change*, **92–93**, 30–39, <https://doi.org/10.1016/j.gloplacha.2012.04.001>
- Salerno, F., Thakuri, S., Tartari, G., Nuimura, T., Sunako, S., Sakai, A. and Fujita, K. 2017. Debris-covered glacier anomaly? Morphological factors controlling changes in the mass balance, surface area, terminus position, and snow line altitude of Himalayan glaciers. *Earth and Planetary Science Letters*, **471**, 19–31, <https://doi.org/10.1016/j.epsl.2017.04.039>
- Schauwecker, S., Rohrer, M., Aaa, A.A., *et al.* 2015. Remotely sensed debris thickness mapping of Bara Shigri Glacier, Indian Himalaya. *Journal of Glaciology*, **61**, 675–688, <https://doi.org/10.3189/2015JG14J102>
- Scherler, D. and Egholm, D.L. 2020. Production and Transport of Supraglacial Debris: Insights From Cosmogenic <sup>10</sup>Be and Numerical Modeling, Chhota Shigri Glacier, Indian Himalaya. *Journal of Geophysical Research: Earth Surface*, **125**, e2020JF005586, <https://doi.org/10.1029/2020JF005586>
- Scherler, D., Leprince, S. and Strecker, M.R. 2008. Glacier-surface velocities in alpine terrain from optical satellite imagery—Accuracy improvement and quality assessment. *Remote Sensing of Environment*, **112**, 3806–3819, <https://doi.org/10.1016/j.rse.2008.05.018>
- Scherler, D., Wulf, H. and Gorelick, N. 2018. Global Assessment of Supraglacial Debris-Cover Extents. *Geophysical Research Letters*, **45**, 11798–11805, <https://doi.org/10.1029/2018GL080158>
- Schneider, D., Huggel, C., Cochachin, A., Guillén, S. and García, J. 2014. Mapping hazards from glacier lake outburst floods based on modelling of process cascades at Lake 513, Carhuaz, Peru. *Adv. Geosci.*, **35**, 145–155, <https://doi.org/10.5194/adgeo-35-145-2014>
- Seong, Y.B., Owen, L., Aaa, A.A., *et al.* 2009. Rates of basin-wide rockwall retreat in the K2 region of the Central Karakoram defined by terrestrial cosmogenic nuclide Be-10. *Physics Research Publications*, **107**.
- Shannon, S., Smith, R., Aaa, A.A., *et al.* 2019. Global glacier volume projections under high-end climate change scenarios. *The Cryosphere*, **13**, 325–350, <https://doi.org/10.5194/tc-13-325-2019>
- Shean, D.E. 2017. *HMA 8-meter DEM Mosaics Derived from Optical Imagery, v1 NSIDC Distributed Active Archive Center*. NSIDC DAAC.
- Shugar, D.H., Burr, A., Aaa, A.A., *et al.* 2020. Rapid worldwide growth of glacial lakes since 1990. *Nature Climate Change*, **10**, 939–945, <https://doi.org/10.1038/s41558-020-0855-4>
- Shukla, A., Arora, M.K. and Gupta, R.P. 2010a. Synergistic approach for mapping debris-covered glaciers using optical-thermal remote sensing data with inputs from geomorphometric parameters. *Remote Sensing of Environment*, **114**, 1378–1387, <https://doi.org/10.1016/j.rse.2010.01.015>
- Shukla, A., Gupta, R. and Arora, M. 2010b. Delineation of debris-covered glacier boundaries using optical and thermal remote sensing data. *Remote Sensing Letters*, **1**, 11–17, <https://doi.org/10.1080/01431160903159316>
- Shukla, A., Garg, P.K. and Srivastava, S. 2018. Evolution of Glacial and High-Altitude Lakes in the Sikkim, Eastern Himalaya Over the Past Four Decades (1975–2017). *Front. Env. Science*, **6**, 81, <https://doi.org/10.3389/fenvs.2018.00081>
- Solomina, O., Bradley, R., Aaa, A.A., *et al.* 2016. Glacier fluctuations during the past 2000 years. *Quaternary Science Reviews*, **149**, 161–190, <https://doi.org/10.1016/j.quascirev.2016.04.008>
- Stefaniak, A.M., Robson, B.A., Cook, S.J., Clutterbuck, B., Midgley, N.G. and Labadz, J.C. 2021. Mass balance and surface evolution of the debris-covered Miage Glacier, 1990–2018. *Geomorphology*, **373**, 107474, <https://doi.org/10.1016/j.geomorph.2020.107474>
- Steiner, J.F. and Pellicciotti, F. 2016. Variability of air temperature over a debris-covered glacier in the Nepalese Himalaya. *Annals of Glaciology*, **57**, 295–307, <https://doi.org/10.3189/2016AoG71A066>
- Steiner, J., Pellicciotti, F., Buri, P., Miles, E., Immerzeel, W.W. and Reid, T. 2015. Modelling ice-cliff backwasting on a debris-covered glacier in the Nepalese Himalaya. *Journal of Glaciology*, **61**, 889–907, <https://doi.org/10.3189/2015JG14J194>
- Steiner, J.F., Buri, P., Miles, E.S., Ragetti, S. and Pellicciotti, F. 2019. Supraglacial ice cliffs and ponds on debris-covered glaciers: spatio-temporal distribution and characteristics. *Journal of Glaciology*, **65**, 617–632, <https://doi.org/10.1017/jog.2019.40>
- Steiner, J.F., Kraaijenbrink, P.D.A. and Immerzeel, W.W. 2021. Distributed Melt on a Debris-Covered Glacier: Field Observations and Melt Modeling on the Lirung Glacier in the Himalaya. *Frontiers in Earth Science*, **9**, 567, <https://doi.org/10.3389/feart.2021.678375>
- Stewart, R.L., Westoby, M., Pellicciotti, F., Rowan, A., Swift, D., Brock, B. and Woodward, J. 2021. Using climate reanalysis data in conjunction with multi-temporal satellite imagery to derive supraglacial debris thickness changes from energy-balance modelling. *Journal of Glaciology*, **67**, 366–384, <https://doi.org/10.1017/jog.2020.111>
- Stokes, C.R., Popovnin, V., Aleynikov, A., Gurney, S.D. and Shahgedanova, M. 2007. Recent glacier retreat in the Caucasus Mountains, Russia, and associated increase in supraglacial debris cover and supra-/proglacial lake development. *Annals of Glaciology*, **46**, 195–203, <https://doi.org/10.3189/172756407782871468>
- Strozzi, T., Paul, F. and Kaab, A. 2010. Glacier Mapping with ALOS PALSAR Data within the ESA GlobGlacier Project. In: Lacoste-Francis, H. (ed.) *ESA Living Planet Symposium, 2010/12/1, Bergen, Norway*, 114.

- Strozzi, T., Wiesmann, A., Käab, A., Joshi, S. and Mool, P. 2012. Glacial lake mapping with very high resolution satellite SAR data. *Natural Hazards and Earth System Sciences*, **12**, 2487–2498, <https://doi.org/10.5194/nhess-12-2487-2012>
- Suzuki, R., Fujita, K. and Ageta, Y. 2007. Spatial distribution of thermal properties on debris-covered glaciers in the Himalayas derived from ASTER data. *Bulletin of Glaciological Research*, **24**, 13–22.
- Tampucci, D., Citterio, C., Gobbi, M. and Caccianiga, M. 2016. Vegetation outlines of a debris-covered glacier descending below the treeline. *Plant Sociology*, **53**, 43–52, <https://doi.org/10.7338/pls201653103>
- Taschner, S. and Ranzi, R. 2002. Landsat-TM and ASTER data for monitoring a debris covered glacier in the Italian Alps within the GLIMS project. Proceedings IGARSS 2002, Toronto, ON, Canada, **4**, 1044–1046, <https://doi.org/10.1109/IGARSS.2002.1025770>
- Thompson, S.S., Benn, D.I., Dennis, K. and Luckman, A. 2012. A rapidly growing moraine-dammed glacial lake on Ngozumpa Glacier, Nepal. *Geomorphology*, **145**, 1–11, <https://doi.org/10.1016/j.geomorph.2011.08.015>
- Thompson, S., Benn, D., Mertes, J. and Luckman, A. 2016. Stagnation and mass loss on a Himalayan debris-covered glacier: Processes, patterns and rates. *Journal of Glaciology*, **62**, 1–19, <https://doi.org/10.1017/jog.2016.37>
- Thompson, S., Kulesha, B., Douglas I. Benn and Mertes, J.R. 2017. Anatomy of terminal moraine segments and implied lake stability on Ngozumpa Glacier, Nepal, from electrical resistivity tomography (ERT). *Scientific Reports*, **7**, 46766, <https://doi.org/10.1038/srep46766>
- United Nations 2004. *Living with Risk: a global review of disaster reduction initiatives*. United Nations Publications, New York and Geneva, [https://www.unisdr.org/files/657\\_lwrl.pdf](https://www.unisdr.org/files/657_lwrl.pdf)
- van Woerkom, T., Steiner, J.F., Kraaijenbrink, P.D.A., Miles, E.S. and Immerzeel, W.W. 2019. Sediment supply from lateral moraines to a debris-covered glacier in the Himalaya. *Earth Surface Dynamics*, **7**, 411–427, <https://doi.org/10.5194/esurf-7-411-2019>
- Vincent, C., Wagon, P., Aaa, A.A., et al. 2016. Reduced melt on debris-covered glaciers: investigations from Changri Nup Glacier, Nepal. *The Cryosphere*, **10**, 1845–1858, <https://doi.org/10.5194/tc-10-1845-2016>
- Vuichard, D. and Zimmermann, M. 1987. The 1985 Catastrophic Drainage of a Moraine-Dammed Lake, Khumbu Himal, Nepal: Cause and Consequences. *Mountain Research and Development*, **7**, 91, <https://doi.org/10.2307/3673305>
- Wang, W., Yao, T., Gao, Y., Yang, X. and Kattel, D.B. 2011. A First-order Method to Identify Potentially Dangerous Glacial Lakes in a Region of the Southeastern Tibetan Plateau. *Mountain Research and Development*, **31**, 122–130, <https://doi.org/10.1659/MRD-JOURNAL-D-10-00059.1>
- Wang, W., Xiang, Y., Gao, Y., Lu, A. and Yao, T. 2014. Rapid expansion of glacial lakes caused by climate and glacier retreat in the Central Himalayas. *Hydrological Processes*, **29**, <https://doi.org/10.1002/hyp.10199>
- Wang, X., Guo, X., Aaa, A.A., et al. 2020. Glacial lake inventory of high-mountain Asia in 1990 and 2018 derived from Landsat images. *Earth Syst. Sci. Data*, **12**, 2169–2182, <https://doi.org/10.5194/essd-12-2169-2020>
- Wangchuk, S. and Bolch, T. 2020. Mapping of glacial lakes using Sentinel-1 and Sentinel-2 data and a random forest classifier: Strengths and challenges. *Science of Remote Sensing*, **2**, 100008, <https://doi.org/10.1016/j.srs.2020.100008>
- Watanabe, T., Kameyama, S. and Sato, T. 1995. Imja Glacier Dead-Ice Melt Rates and Changes in a Supra-Glacial Lake, 1989–1994, Khumbu Himal, Nepal: Danger of Lake Drainage. *Mountain Research and Development*, **15**, 293–300, <https://doi.org/10.2307/3673805>
- Watanabe, T., Lamsal, D. and Ives, J.D. 2009. Evaluating the growth characteristics of a glacial lake and its degree of danger of outburst flooding: Imja Glacier, Khumbu Himal, Nepal. *Norsk Geografisk Tidsskrift-Norwegian Journal of Geography*, **63**, 255–267, <https://doi.org/10.1080/00291950903368367>
- Watson, C.S., Quincey, D.J., Carrivick, J.L. and Smith, M.W. 2016. The dynamics of supraglacial ponds in the Everest region, central Himalaya. *Global and Planetary Change*, **142**, 14–27, <https://doi.org/10.1016/j.gloplacha.2016.04.008>
- Watson, C.S., Quincey, D.J., Carrivick, J.L. and Smith, M.W. 2017. Ice cliff dynamics in the Everest region of the Central Himalaya. *Geomorph.*, **278**, 238–251, <https://doi.org/10.1016/j.geomorph.2016.11.017>
- Watson, C.S., King, O., Miles, E.S. and Quincey, D.J. 2018a. Optimising NDWI supraglacial pond classification on Himalayan debris-covered glaciers. *Remote Sensing of Environment*, **217**, 414–425, <https://doi.org/10.1016/j.rse.2018.08.020>
- Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., Rowan, A.V. and Richardson, R. 2018b. Heterogeneous water storage and thermal regime of supraglacial ponds on debris-covered glaciers. *Earth Surface Processes and Landforms*, **43**, 229–241, <https://doi.org/10.1002/esp.4236>
- Wessel, B., Huber, M., Wohlfart, C., Marschalk, U., Kosmann, D. and Roth, A. 2018. Accuracy assessment of the global TanDEM-X Digital Elevation Model with GPS data. *ISPRS Journal of Photogrammetry and Remote Sensing*, **139**, 171–182, <https://doi.org/10.1016/j.isprsjprs.2018.02.017>
- Wessels, R.L., Kargel, J.S. and Kieffer, H.H. 2002. ASTER measurement of supraglacial lakes in the Mount Everest region of the Himalaya. *Annals of Glaciology*, **34**, 399–408, <https://doi.org/10.3189/172756402781817545>
- Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J. and Reynolds, J.M. 2012. ‘Structure-from-Motion’ photogrammetry: A low-cost, effective tool for geoscience applications. *Geomorphology*, **179**, 300–314, <https://doi.org/10.1016/j.geomorph.2012.08.021>
- Westoby, M.J., Glasser, N.F., Brasington, J., Hambrey, M.J., Quincey, D.J. and Reynolds, J.M. 2014. Modelling outburst floods from moraine-dammed glacial lakes. *Earth-Science Reviews*, **134**, 137–159, <https://doi.org/10.1016/j.earscirev.2014.03.009>
- Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J., Reynolds, J.M., Hassan, M.A.A.M. and Lowe, A. 2015. Numerical modelling of glacial lake outburst floods using physically based dam-breach models. *Earth Surface Dynamics*, **3**, 171–199, <https://doi.org/10.5194/esurf-3-171-2015>
- Westoby, M.J., Rounce, D.R., Shaw, T.E., Fyffe, C.L., Moore, P.L., Stewart, R.L. and Brock, B.W. 2020. Geomorphological evolution of a debris-covered glacier surface. *Earth Surface Processes and Landforms*, **45**, 3431–3448, <https://doi.org/10.1002/esp.4973>
- Whalley, W.B. and Martin, H.E. 1992. Rock glaciers: II models and mechanisms. *Progress in Physical Geography: Earth and Environment*, **16**, 127–186, <https://doi.org/10.1177/030913339201600201>
- Wilson, R., Glasser, N.F., Reynolds, J.M., Harrison, S., Anaconda, P.I., Schaefer, M. and Shannon, S. 2018. Glacial lakes of the Central and Patagonian Andes. *Global and Planetary Change*, **162**, 275–291, <https://doi.org/10.1016/j.gloplacha.2018.01.004>
- Wilson, R., Harrison, S., Aaa, A.A., et al. 2019. The 2015 Chileno Valley glacial lake outburst flood, Patagonia. *Geomorphology*, **332**, 51–65, <https://doi.org/10.1016/j.geomorph.2019.01.015>
- Wirbel, A., Jarosch, A.H. and Nicholson, L. 2018. Modelling debris transport within glaciers by advection in a full-Stokes ice flow model. *The Cryosphere*, **12**, 189–204, <https://doi.org/10.5194/tc-12-189-2018>
- Worni, R., Huggel, C. and Stoffel, M. 2013. Glacial lakes in the Indian Himalayas – From an area-wide glacial lake inventory to on-site and modeling based risk assessment of critical glacial lakes. *Science of The Total Environment*, **468–469**, S71–S84, <https://doi.org/10.1016/j.scitotenv.2012.11.043>
- Worni, R., Huggel, C., Clague, J. and Schaub, Y. 2014. Coupling glacial lake impact, dam breach, and flood processes: A modeling perspective. *Geomorphology*, **224**, 161–176, <https://doi.org/10.1016/j.geomorph.2014.06.031>
- Xie, F., Liu, S., Aaa, A.A., et al. 2020a. Upward Expansion of Supra-Glacial Debris Cover in the Hunza Valley, Karakoram, during 1990–2019. *Frontiers in Earth Science*, **8**, <https://doi.org/10.3389/feart.2020.00308>
- Xie, Z., Haritashya, U.K., Asari, V.K., Young, B.W., Bishop, M.P. and Kargel, J.S. 2020b. GlacierNet: A Deep-Learning Approach for Debris-Covered Glacier Mapping. *IEEE Access*, **8**, 83495–83510, <https://doi.org/10.1109/ACCESS.2020.2991187>
- Xu, X., Asawa, T. and Kobayashi, H. 2020. Narrow-to-Broadband Conversion for Albedo Estimation on Urban Surfaces by UAV-Based Multispectral Camera. *Remote Sensing*, **12**, <https://doi.org/10.3390/rs12142214>
- Zemp, M., Frey, H., Aaa, A.A., et al. 2015. Historically unprecedented global glacier decline in the early 21st century. *Journal of Glaciology*, **61**, 745–762, <https://doi.org/10.3189/2015JG15J017>
- Zhang, Y., Fujita, K., Liu, S., Liu, Q. and Nuimura, T. 2011. Distribution of debris thickness and its effect on ice melt at Hailuoguo glacier, southeastern Tibetan Plateau, using in situ surveys and ASTER imagery. *Journal of Glaciology*, **57**, 1147–1157, <https://doi.org/10.3189/002214311798843331>
- Zhang, G., Yao, T., Xie, H., Wang, W. and Yang, W. 2015. An inventory of glacial lakes in the Third Pole region and their changes in response to global warming. *Global and Planetary Change*, **131**, 148–157, <https://doi.org/10.1016/j.gloplacha.2015.05.013>
- Zhang, M.-M., Chen, F. and Tian, B.-S. 2018. An automated method for glacial lake mapping in High Mountain Asia using Landsat 8 imagery. *Journal of Mountain Science*, **15**, 13–24, <https://doi.org/10.1007/s11629-017-4518-5>
- Zhang, J.J., Menenti, M.L. and Hu, G. 2019. Glacier Facies Mapping Using a Machine-Learning Algorithm: The Parlung Zangbo Basin Case Study. *Remote Sensing*, **11**, 452, <https://doi.org/10.3390/rs11040452>
- Zhang, B., Liu, G., Aaa, A.A., et al. 2021. Monitoring Dynamic Evolution of the Glacial Lakes by Using Time Series of Sentinel-1A SAR Images. *Remote Sensing*, **13**, <https://doi.org/10.3390/rs13071313>
- Zhao, H., Chen, F. and Zhang, M. 2018. A Systematic Extraction Approach for Mapping Glacial Lakes in High Mountain Regions of Asia. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, **11**, 2788–2799, <https://doi.org/10.1109/JSTARS.2018.2846551>
- Zheng, G., Allen, S.K., Aaa, A.A., et al. 2021. Increasing risk of glacial lake outburst floods from future Third Pole deglaciation. *Nature Climate Change*, **11**, 411–417, <https://doi.org/10.1038/s41558-021-01028-3>