



HAL
open science

Allophanes, a significant soil pool of silicon for plants

Sophie Cornu, Jean-Dominique Meunier, Céline Ratie, Frédéric Ouedraogo, Yves Lucas, Patricia Merdy, Doris Barboni, Camille Delvigne, Daniel Borschneck, Olivier Grauby, et al.

► To cite this version:

Sophie Cornu, Jean-Dominique Meunier, Céline Ratie, Frédéric Ouedraogo, Yves Lucas, et al.. Allophanes, a significant soil pool of silicon for plants. *Geoderma*, 2022, 412, pp.115722. <10.1016/j.geoderma.2022.115722>. <hal-04018105v2>

HAL Id: hal-04018105

<https://hal.science/hal-04018105v2>

Submitted on 7 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

Allophanes, a significant soil pool of silicon for plants

Sophie Cornu ^{a,*}, Jean-Dominique Meunier ^a, Céline Ratie ^b, Frédéric Ouedraogo ^a, Yves Lucas ^c, Patricia Merdy ^c, Doris Barboni ^a, Camille Delvigne ^a, Daniel Borschneck ^a, Olivier Grauby ^d, Catherine Keller ^a

^aAix-Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, 13545 Aix en Provence Cedex 4, France

^bINRAE, InfoSol, 45075 Orléans, France

^cUniversité de Toulon, Aix-Marseille Univ, CNRS, IM2NP, 83041 Toulon Cedex 9, France

^dAix-Marseille Univ, CNRS – UMR 7325 CINaM, campus de Luminy, Case 913 13288 Marseille Cedex 9, France

Keywords: Land use change Clay Phytolith Temperate ecosystems Bioavailable Si Pedogenesis

DOI: <https://doi.org/10.1016/j.geoderma.2022.115722>

Corresponding author. E-mail address: sophie.cornu@inrae.fr (S. Cornu).

1. Introduction

The variability of Si content in soils is large, from <1 % wt in Histosols to greater than 45 % wt in some Podzols (Sommer et al., 2006). Si is included in a large number of soil solid phases of various origins: (1) minerals derived from the parent material as quartz, feldspar, phyllosilicates (e.g. muscovite, biotite) etc.; (2) solid phases resulting from the transformation by weathering and pedogenesis of the primary minerals into clay minerals (kaolinite, smectite, illite among others), crystalline and amorphous material, including poorly crystalline Fe oxides and short-range ordered (SRO) aluminosilicates (e.g. allophanes and imogolites); (3) amorphous silica particles (BSi) formed by Si biocycling and synthesized by plants (phytoliths), animals (sponge spicules) or protists (testate amoeba shells, diatom frustules) (Sommer et al., 2006; Puppe et al., 2015). While soils offer a wide range of Si sources, plants absorb Si only from the soil solution as Si(OH)₄. Therefore, deciphering the source of soluble Si (DSi) is challenging because of the complex biogeochemical interactions involving Si in soils (Schaller et al., 2021). The conversion of forest to cropland was shown to lead to several Si pool changes: (1) a decrease in the phytolith pool (Conley et al., 2008; Struyf et al., 2010; Guntzer et al., 2012; Vandevenne et al., 2015), in the easily soluble Si pool (Clymans et al., 2011) and the weathering of silicate minerals (Struyf et al., 2010) and (2) an increase in the pedogenic Si fraction i.e. clays (Vandevenne et al., 2015). In addition, the growing crops such as rice, sugar cane, or wheat (Hodson et al., 2005; Guntzer et al., 2012; Katz, 2015) that are among the most important Si accumulators (1 % or more Si by dry weight) are predicted to change the global Si biogeochemistry in the future by increasing the reservoir of BSi on earth (Carey et al., 2016).

Various methods have been used to estimate DSi and bioavailable Si (Liang et al., 2015). Their results suggest that bioavailable Si concentration is both negatively correlated with total Si content, reflecting the predominance of low solubility minerals such as quartz in the soils (Yanai et al., 2016; Landr'e et al., 2020) and positively correlated with soil properties such as weatherable aluminosilicates (Li et al., 2020a), phytolith content (Li et al., 2020b; de Tombeur et al., 2020), pH (Klotzbücher et al., 2018; Meunier et al., 2018; Miles et al., 2014; Caubet et al., 2020), soil organic matter, iron oxides (Yanai et al., 2016) and the clay (<2 µm) fraction contents (Liang et al., 2015; Yanai et al., 2016; Caubet et al., 2020).

In temperate soils, despite the observed decrease of soil phytolith content under agriculture, higher DSi (estimated by the CaCl₂ method referred to herein as Si_{CaCl2}) concentration was observed under agriculture compared to forest developed on sedimentary rocks (Vandevenne et al., 2015; Caubet et al., 2020). This was related to a higher pH observed in agricultural soils compared to forest soils (Caubet et al., 2020) pointing out two mechanisms: greater phytoliths dissolution, or greater adsorption onto clay minerals at higher pH. Cornu et al. (2012) also observed that the liming of agricultural soils in Europe induced a change in the nature of the clay minerals, and hence, potentially in the relative proportion of the different Si pools. Consequently, despite their lower solubility compared to phytoliths (Frayse et al., 2009), clays may play an important role in maintaining high values of Si_{CaCl2}, somehow cancelling out the phytolith pool decrease (Vandevenne et al., 2015; Caubet et al., 2020).

To identify the mechanisms controlling Si_{CaCl2} in temperate soils and to understand the impact of land use on these mechanisms in different soil types, we designed a paired-site approach in which we sampled adjacent soils differing only in their land use while other pedological factors (topography, parent material, climate, and age) were the same. Seven paired sites were selected with calcaric and non-calcaric soils, developed on sedimentary parent materials (limestone and loess). Beside the Si_{CaCl2} pool, we measured the pools of Si associated to crystalline and amorphous material, including poorly crystalline Fe oxides and SRO-aluminosilicates, and biogenic Si particles. Only the topsoil horizons (0–0.3 m) were analysed as they were considered the most impacted by land use change.

2. Material and methods

2.1. Identification of the paired sites and sampling strategy

Paired sites are not easy to locate as they have to differ only by their land use, all other pedological factors (parent material, climate, topography, and time) being kept constant between the two sites (Filippi et al., 2016). We chose soils on sedimentary materials as land use impacts Si_{CaCl2} on these soils, and calcaric and non-calcaric soils as their behaviour with respect to Si_{CaCl2} differs (Caubet et al., 2020). In France, most cultivated soils with these characteristics are developed either on limestone or on decarbonised loess. We also

chose to study soils with different degrees of pedogenetic evolution: calcareic and hypereutric Cambisols for limestone; Luvisols and Albeluvisol for loess (WRB, 2006). To identify the paired sites, we used the French Soil Quality Network (RMQS) of 2111 georeferenced sites in France spread over a regular grid (16*16 km) for which soil type, parent material and land use are provided (Jolivet et al., 2006). All the RMQS sites with the appropriate parent material and soil type either cultivated or under forest, were extracted from the database. For each selected RMQS site, an adjacent site (i.e. with the same climate), with the same topographical position (therefore expected to be on the same type of soil and parent material) and under the complementary land use (either forest or cultivation) was identified on the basis of: (i) the maps of the French National Institute of Geographic and Forest Information (IGN, 1:25000); (ii) Cassini's maps, the oldest available maps with records of land use (Cassini, 1750), both available on Geoportail (<https://www.geoportail.gouv.fr/>); (iii) aerial photographs taken at different times during the 20th century. Only paired sites for which the land use change occurred more than 100 years ago were kept. In most cases, the forested areas have been preserved from cultivation because they belonged to the state or to castle parks. Nineteen pairs of adjacent plots were identified. Another paired site of Albeluvisol (WRB, 2006) on already well-characterised loess was added (Montagne et al., 2016). The 20 sites were in the northern half of France (Table SI1), where the cultivation of wheat is common. Most of them included cereals in their rotation (Table SI1). Forests were mixed deciduous forests except for site 62 that had a conifer forest (Table SI1). All the selected paired sites were auger-sampled and described in the spring of 2015. In the cultivated area, one sample was taken in the upper horizon which is homogenized by ploughing (usually 0–25 cm). In the nearby forested area two samples were taken (usually 0–5 and 5–25 cm). On the basis of the soil morphology, eighteen out of the 20 paired sites were kept and analysed for rare earth element (REE) concentrations and particle size distribution to detect potential changes in parent materials. Analyses were performed at the Laboratory of Soil Analysis (Arras, France). Seven paired sites were finally kept (Table SI1), as their REE and grain size skeleton patterns did not differ between land uses considering the analytical uncertainties (see supplementary information, SI, part 3 for more details) and thus could be considered as having the same parent material. Bulk density measurements were performed using the volumetric cylinder method in cultivated areas and the excavation method in forested areas because the root density prevented the use of the cylinder. When bulk density could not be measured (e.g., at site 1166), it was estimated using the pedotransfer function proposed by Alexander (1980), which relates bulk density to soil organic carbon (SOC) content. To compare the forested and cultivated sites, composite values (0–25 cm) were calculated for the forested sites considering the thickness of the two sampled horizons (0–5 cm and 5–25 cm) and their respective bulk densities.

2.2. Samples characterisation

2.2.1. Pedological analysis Air dried samples were analysed for pedological characteristics at the Soil Analysis Laboratory of INRAE (Arras, France). The particle-size distribution was obtained by wet sieving and the pipette method (NF X 31-107). Total SOC concentration was measured by dry combustion (NF ISO 10694), CEC (cation exchange capacity) by the cobaltihexamine chlorine method (NF X 31-130), and pH in water with a 1:5 soil:water ratio (NF X 31-107). The particle size fraction contents are hereafter denominated < 2 µm, 2–20 µm, 20–50 µm, 50–200 µm, 200–2000 µm. The < 2 µm fraction CEC (CEC<2µm) was calculated following equation (1).

$$CEC_{<2\mu m} = ((CEC_{SOC} * SOC) / < 2\mu m) * 1000 \quad (1)$$

where SOC and < 2 µm are the contents in SOC and in the < 2 µm fraction, respectively. We used 200 cmol+ kg⁻¹ as an estimate of CEC_{SOC} (Baize, 2000) of both the ploughed horizon of the cultivated site and for the 5–25 cm horizon of the forested plots and 100 cmol+ kg⁻¹ as an estimate of CEC_{SOC} of the 0–5 cm horizon of the forested plots that contains a greater fraction of particulate organic matter with a lower CEC (Balesdent, pers. comm., 2018).

Total major elements concentrations in soil, including Si (Si_{tot}), were also analysed at the Soil Analysis Laboratory of INRAE (Arras, France) by ICP-MS analysis after HF tri-acid dissolution (Al_{tot}, Fe_{tot}, Na_{tot}, Ca_{tot}, Mg_{tot}, K_{tot}, Mn_{tot}) or alkaline fusion (Si_{tot}, NF ISO 14869-2).

2.2.2. Chemical extraction of different soil Si pools

We used three types of soft chemical extraction in parallel to access different soil Si pools:

(1) Soluble and bioavailable Si was extracted using the CaCl₂ method (Si_{CaCl2}; Haysom and Chapman, 1975; Narayanaswamy and Prakash, 2010). The soils were mixed with a 0.01 mol L⁻¹ CaCl₂ solution with a solid/solution ratio of 10, in two replicates. The suspension was stirred for 16 h and filtered at 0.45 µm.

(2) Poorly crystalline Fe oxides and short-range ordered (SRO) aluminosilicates such as allophanes and imogolites were extracted by the oxalate method (Si_o, Tamm, 1922; Wada, 1989; Harsh, 2005). It is assumed that this method does not dissolve amorphous silica particles (i.e., phytoliths) (Kodama and Ross, 1991; Saccone et al., 2007). The Si_o pool thus represents Si associated with amorphous co-precipitated Fe and Al compounds. Allophane content was estimated using Harsh's (2005) equation (2):

$$\% \text{ allophane} = Si_o / (0.067 Al_o / Si_o + 0.27) \quad (2)$$

(3) Iron oxides were extracted by the dithionite-citrate-bicarbonate (DCB) method (Si_d, Mehra and Jackson, 1960), which also dissolves the poorly crystalline fraction and may dissolve phytoliths (Georgiadis et al., 2013).

Si extracted by these three methods and Fe and Al extracted by the latter two were measured using ICP-OES (Inductively Coupled Plasma Optical Emission Spectroscopy). Both Si_o and Si_d were extracted and measured at the Soil Analysis Laboratory of INRAE (Arras, France).

The different soil Si forms were then estimated considering that (i) the DSi equals the Si_{CaCl2}; (ii) Si in SRO minerals (allophanes, imogolites and poorly crystalline Fe oxides) equals (Si_o - Si_{CaCl2}); (iii) Si associated to crystalline Fe oxides or to the part of phytoliths dissolved by the extractant equals (Si_d - Si_o); and (iv) Si in crystalline silicates plus part of the phytoliths equals (Si_{tot} - Si_d). We have to keep in mind that all these extractions are operationally defined and may extract part of soil solid phases other than the targeted ones as already extensively discussed in the literature (e.g. Georgiadis et al., 2013).

Error bars were estimated considering 5 % uncertainty on the chemical and bulk density measurements.

2.2.3. Phytolith extraction and characterisation

Soil phytoliths were quantified, identified and their weathering state was evaluated.

BSi particles were extracted from the soils using wet digestion and heavy liquid separation (Jouquet et al., 2020; see SI part 2 for more details). No replicates have been done. Extracts, which happened to include phytoliths as well as low density particles such as micro-charcoals and amorphous silica skeletons of chrysophytes, diatoms, sponges, etc., were dried and weighed before being mounted on microscope slides to evaluate the real proportion of phytoliths versus non-phytolith particles in the extracts and, therefore, provide an estimation of the phytolith contents.

We considered that phytoliths contained 42 % of Si (Alexandre et al., 1997) and could thus estimate the soil phytolith Si pool based on the phytolith abundances in the soils.

Since cultivated plants produce morphotypes of phytoliths different from forest species (trees, shrubs, ferns, forbs), phytolith taxonomic identification was performed according to the International Committee for Phytolith Taxonomy (2019) and counting was carried out.

Lastly, the analysis of the phytolith surface is a good indicator of their weathering state (Sommer et al., 2006; Jouquet et al., 2020), we measured the proportion of corroded vs non-corroded phytoliths (for more details, see SI part 2).

2.2.4. Mineralogical characterisation of the < 2 μm fraction

Dried soils were dispersed in 200 to 300 mL of ultrapure water by rotative shaking for 22 h using plastic bottles and 20 agate balls (\varnothing 5 mm) (Balesdent et al., 1991). The mass of dried soil was chosen to obtain ca. 4 g of < 2 μm fraction. The suspension thus obtained was further dispersed by addition of 2 mL of Na-hexametaphosphate at 150 g L⁻¹, mechanical shaking, and 5-min ultrasonic dispersion. The < 2 μm fraction was then sampled according to the Stokes law and the settled soil particles resuspended for a new extraction step. The process was repeated until more than 80 % of the total fraction was recovered (that is more than 3.2 g of < 2 μm fraction out of 4 g). An aliquot of the < 2 μm suspension was dried to determine the < 2 μm suspension concentration. Oriented slides were made by deposition on glass slides of 10 mg of < 2 μm particles spiked with 10 % of boehmite added as a reference for X-ray diffraction (XRD) studies. The glass slides were then analysed with a Panalytical X'Pert Pro X-ray diffractometer equipped with a cobalt anti-cathode ($\lambda = 1.79 \text{ \AA}$) at 40 kV and 40 mA, with three treatments following Robert and Tessier (1974): air-dried (AD) and ethylene glycol (EG) solvation and heating at 490 °C for 4 h.

Peak areas of the main mineral phases and boehmite were measured using the Fityk software (Wojdyr, 2010). The peak area ratio with respect to boehmite was used to compare samples. Transmission electron microscope (TEM, JEOL JEM 2011) coupled to an energy-dispersive X-ray spectrometer (EDX, X-Flash Silicon Drift Detector 5030, Bruker) analysis was performed on the < 2 μm fraction of sample BS3F1 (Table SI1) for checking the presence of allophanes.

2.3. Data analysis

Pedological characteristics and major element concentrations of the seven studied paired sites were analysed by principal component analysis (PCA). To identify different groups of soils from a pedological and chemical point of view, a hierarchical classification was performed on the first two factors of the PCA.

Soil characteristics of the paired sites were considered as different when the difference for a given soil characteristic exceeded the analytical uncertainties. As only one measurement by site was performed, no statistical tests could be performed to further test the statistical significance of this difference.

Significance of Pearson correlations between the SiCa12 concentrations and different soil parameters was estimated based on the p-values considering a threshold level of 5 %. All the statistical analyses were performed using the XLSTAT software for windows (Addinsoft).

3. Results

3.1. Seven paired sites with contrasted pedological characteristics and total concentrations of major elements

Pedological characteristics and total contents of major elements in the topsoils of the seven paired sites are reported in Tables SI1 and SI2. They were computed by PCA. The first two components explained 72 % of the dataset variance with 60 % for the first component (Fig. 1). The first component was correlated positively to Cat_{tot}, the < 2 μm fraction, Fe_{tot}, Alt_{tot}, Mg_{tot} and the SOC contents, as well to pH and the CEC, and negatively to Si_{tot}, Na_{tot}, silt (2–20 μm and 20–50 μm fractions) and fine sand (50–200 μm fraction) contents. The second component (12 % of the variance) was correlated positively to the 2–20 μm fraction, and negatively to the 50–200 μm fraction (Fig. 1a). The PCA discriminated three groups of soils confirmed by a hierarchical clustering (Fig. 1b). The first group corresponds to calcareous Cambisols and is positively correlated with the first PCA component, which means high pH, < 2 μm fraction, Fe_{tot} and SOC contents. The second group corresponds to soils developed on loess (Luvisols and Albeluvisols) and is negatively correlated to the first PCA component, which means high 20–50 μm fraction and Si_{tot} contents, low pH, Fe_{tot} and SOC contents. The third group corresponds to hypereutric Cambisols and is intermediary between the other two groups, showing more similarities with the soils developed on loess than with the calcareous Cambisols.

Although the PCA did not highlight the impact of land use, pH and Si content in SRO were higher and SOC values lower under cultivation than under forest (Fig. 2). The CEC of the < 2 μm fraction did not differ with land use in Cambisols developed on limestone but was higher under cultivation in Luvisols and Albeluvisols developed on loess (Fig. 2).

3.2. Phytoliths of grasses were dominant in the two land uses

The relative abundances of the main phytolith types grouped according to plant types and confidence intervals are given in Table 1. On average, the biogenic silica extracts contained 89 % of phytoliths (range 51–99 %), the remaining fraction consisting of 3 % of other biogenic silica particles (range 0–8 %) and of 9 % charcoal (range 0–48 %); all particles had roughly the same size. For each soil, the mass of phytoliths corrected by phytolith relative abundance (in %) (Table 1) indicates that phytolith pools in soils ranged from < 0.01 to 0.616 g 100 g⁻¹ corresponding to 90 and up to 3978 kg Si ha⁻¹. For comparison, 800 kg Si ha⁻¹ was found in a rice field from Camargue, France (Desplanques et al., 2006), and values ranging from 450 to 1400 kg Si ha⁻¹ in French forest soils (Bartoli, 1983), using a similar procedure (heavy liquid extraction).




Fig. 1. Principal component analysis (PCA) of the pedological characteristics and major element concentrations for the seven paired sites studied. a- Circle of correlations; b- space of the individuals (F states for forest and C for cultivation). Different symbols correspond to the different groups defined by a hierarchical classification performed on the first two factors of the PCA. Green diamonds encompass the calcaric Cambisols, blue circles the hypereutric Cambisols and red triangles correspond to soils developed on loess parent materials. Forest sites correspond to composite values (0–25 cm) calculated considering the thickness of the two sampled horizons (0–5 cm and 5–25 cm), their concentrations in the different elements and their respective bulk densities.

The phytolith contents were very low in calcaric Cambisols and Albeuvisols (Fig. 2i), for which there was no clear differences due to land use. For Luvisols, we observed clearly higher phytolith contents under forest (Fig. 2i). For the hypereutric Cambisols, a clear difference in the amount of phytoliths between land uses was also observed, although not always in the same direction: for the BS2 site more phytoliths were recorded under forest, while for the 1166 site more phytoliths were observed under cultivation (Fig. 2i).

Grass phytoliths (silica short cells and various silicified epidermal tissues) represented >64 % in all forest and cultivated soil samples, except in forest samples 755-F1, 755-F2, and 1166-F1 (25 % only) (Fig. 3a). Phytoliths assigned to forest species (see Tables in SI part 2 for details) only represented 1–20 % of the total phytoliths sum, except in the forest soil sample 1166-F1 (45 %) (Fig. 3b).

The proportions of ELONGATE DENDRITIC phytoliths, mainly produced by grass inflorescences, tended to be higher in the cultivated soils of the hypereutric Cambisols (Fig. 3c). Again, in the other soils their abundance was too low to discuss any differences between land uses (Fig. 3c). These results showed that in the forest soils, except for the forest site 1166, the forest-indicator phytoliths exhibited a very low abundance relative to grass phytoliths (Fig. 3), suggesting that wild grasses were the main producers of phytoliths in forest understorey. The proportion of corroded phytoliths (ELONGATE and BLOCKY) was not different between forest and cultivated soils, except for samples from sites 1166 and BS2 where it was higher in the forest soil and in the cultivated soil respectively (Table 1).

3.3. Mineralogy of the < 2 µm fraction in the topsoils of the seven paired sites

The mineralogical composition of the < 2 µm fractions was rather similar for all forest soils (Fig. 4). All the fractions contained quartz (peak at 3.34 Å in all treatments), kaolinite (peak at 7.13 Å in both AD and EG treatments), illite (peak at 10 Å in all treatments), chlorite (peak at 14 Å in all treatments), poorly crystalline smectites and/or interlayered illite-smectite minerals (large bulgearound 12–11 Å for Na-exchanged samples shifting around 14 Å with glycol with no well-marked peak, more or less marked depending on the considered sample). Soils on loess clearly contained more quartz in their < 2 µm fraction than soils on limestone as shown by their higher quartz:boehmite peak ratio (Table SI3). Cultivation appeared to slightly increase the amount of smectite and interlayered minerals and the amount of illite and decrease the amount of chlorite in soils on loess and in hypereutric Cambisols (Fig. 4 b to d, and Table SI3). In calcaric Cambisols, however, cultivation seemed to result in a decrease of all clay minerals, possibly due to the non-dissolution of the carbonate fraction that acts as clay cement, with no clear change in proportion between the different types of clay minerals (Fig. 4a). Transmission electron microscope (TEM) images showed the presence of discrete spherical particles with external diameter of 6 nm (Fig. 5). These spherical particles appear in the form of a disc with a white central part interpreted as being the projection of a hollow sphere.

3.4. Si pools in the topsoils of the seven paired sites assessed by chemical extractions

Whatever the soil type, more than 99 % of Si_{tot} was contained in crystalline silicates (Si_{tot} - Si_d) (Fig. 2j). The other Si pools (Si in SRO minerals (Si_o - Si_{CaCl2}), Si associated to crystalline oxides (Si_d - Si_o) and Si in phytoliths) represented <0.3 % of the total soil Si. Si in SRO minerals was the most abundant of these pools in calcaric Cambisols, especially when cropped (Fig. 2g), while Si in phytoliths was the most abundant in Luvisols (Fig. 2i). Si associated to crystalline Fe oxides (Si_d - Si_o) represented also a non-negligible part of the non-silicate Si pools, notably under forest (Fig. 2h).




Fig. 2. $< 2 \mu\text{m}$ fraction (a), pH (b), SOC (c), CEC of the $< 2 \mu\text{m}$ fraction (d) and the Fe in iron oxides (Fe_o ; e), and different Si forms (f: soluble and bioavailable Si ($\text{Si}_{\text{CaCl}_2}$); g: Si in SRO minerals (Si_s - $\text{Si}_{\text{CaCl}_2}$); h: Si in crystalline Fe oxides (Si_c - Si_o); i: Si in phytoliths; j: Si in crystalline silicate minerals (Si_{sil} - Si_o)) in cultivated (in grey) and forest (in black) soils of the paired sites studied. Forested sites correspond to composite values (0–25 cm) calculated considering the thickness of the two sampled horizons (0–5 cm and 5–25 cm), their concentrations in the different elements and their respective bulk densities. Error bars correspond to the analytical uncertainties.

Part of this pool may originate from amorphous silica partly dissolved by the DCB reagent (Georgiadis et al., 2013). However, the pool associated to crystalline oxides (Si_c - Si_o) was clearly lower than the Si content in phytoliths, indicating that the phytoliths were not entirely dissolved by DCB.

The $\text{Si}_{\text{CaCl}_2}$ pool represented a negligible proportion of the Si_{tot} (Table 1) and was larger in soils developed on limestone than in soils developed on loess (Fig. 2f). Cultivation increased both the $\text{Si}_{\text{CaCl}_2}$ pool, especially in soils on loess, and the SRO-Si pool, especially in soils on limestone parent material (Fig. 2g). Cultivation, however, did not seem to modify the amount of Si associated to well-crystalline iron oxides and included in crystalline silicates (Fig. 2h and j).

Table 1

The Si pools: Si_{tot} , smooth Si extracted concentrations (Si_o , Si_d and Si_{CaCl2}) and phytolith weight corrected from counting (see text); Si_{phyt} equals phytoliths $\times 0.42$ (see text). (*) The percentage of corroded phytoliths was evaluated on elongated and blocky morphotypes using the microscopic characteristics presented in figure S12.

Soil	Land use	Vegetation/ crop rotation	Sample	Si_{tot}			Phytoliths	Corroded phytoliths	
				$g\ 100\ g^{-1}$	$mg\ kg^{-1}$	$g\ 100\ g^{-1}$			
Calcaric Cambisol	Agric.	N.D.	701C1	26	0.19	0.20	24.6 \pm 0.1	0.030	nd
	Forest	Beech and other deciduous trees	701F1	28	0.075	0.14	21.2 \pm 0.5	0.011	nd
	Agric.	N.D.	755C1	28	0.17	0.21	30 \pm 2	0.002	nd
	Forest	N.D.	755F1	24	0.12	0.19	35 \pm 5	0.028	nd
			755F2	25	0.12	0.17	25.2 \pm 0.3	0.021	nd
Hypereutric Cambisol	Agric.	Cereals	1166C1	34	0.19	0.26	34 \pm 1	0.102	33 \pm 1
	Forest	Maple, hornbeam, oak	1166F1	31	0.072	0.14	37 \pm 1	0.006	60 \pm 3
			1166F2	32	0.063	0.15	24 \pm 1	0.023	nd
	Agric.	Wheat/barley	B62C1	34	0.14	0.24	48.8 \pm 0.4	0.064	62 \pm 6
	Forest	Pine, various deciduous trees	B62F1	33	0.046	0.16	40 \pm 3	0.091	32 \pm 2
			B62F2	34	0.046	0.14	39.6 \pm 0.3	0.239	50 \pm 2
Luvisol	Agric.	Cereals	340C1	39	0.061	0.13	31.4 \pm 0.7	0.246	nd
	Forest	Deciduous trees	340F1	37	0.028	0.1	7.2 \pm 0.9	0.066	nd
			340F2	40	0.046	0.12	9.6 \pm 0.5	0.616	nd
	Agric.	N.D.	62C1	36	0.066	0.14	22.8 \pm 0.3	0.292	25 \pm 1
	Forest	Conifer	62F1	36	0.031	0.14	9.3 \pm 0.3	0.646	37 \pm 1
			62F2	38	0.039	0.11	14 \pm 2	0.351	52 \pm 8
Albeluvisol	Agric.	Wheat, maize, rape	B63C1	40	0.063	0.147	24.5 \pm 0.3	0.009	36 \pm 6
	Forest	Oak, holly, hazel tree	B63F1	41	0.018	0.099	5.9 \pm 0.4	0.046	44 \pm 3
			B63F2	42	0.037	0.125	6.9 \pm 0.3	0.029	56 \pm 6

4. Discussion

4.1. Allophanes are responsible for the Si_{CaCl2} concentration in forest soils

When considering the 21 soil samples of the seven paired sites, the correlations between Si_{CaCl2} and other soil parameters were in a good agreement with previous studies for the following parameters:

1) positive correlations between Si_{CaCl2} and the $< 2\ \mu m$ fraction and its composition (CEC $<2\mu m$, Al_{tot} , Fe_{tot}), pH, CEC (Table 2), as already observed by Caubet et al. (2020), but also between Si_{CaCl2} and Si_o and Si_d (that contains Si_o see section 2.2.2). Such positive correlations of Si_{CaCl2} with Si_o and Si_d were already observed for many soil types by Yanai et al. (2016).

2) a negative correlation of Si_{CaCl2} with Si_{tot} (Table 2), which can be explained by the predominance of low solubility minerals such as quartz in the total Si pool (Yanai et al., 2016; Landr'e et al., 2020).

However, contrary to previous studies made for very different soils, Si_{CaCl2} was not positively correlated with iron oxide (estimated by Fe_a) contents (a sandy soil with very low content of Si_{CaCl2} ; de Tombeur et al., 2020) or the amount of phytoliths (estimated by Na_2CO_3 for Podzol and Chernozem, Li et al., 2020a) (Table 2). Li et al. (2020b) demonstrated that the extraction by Na_2CO_3 gave very different results from the extraction of phytoliths through heavy liquid separation. In the studied soils, the phytolith abundances estimated through heavy liquid separation were negatively correlated to the pH of forest soils ($r = -0.826$, p -value = 0.022). This result agrees with an increasing solubility of phytoliths with increasing pH (Frayssse et al., 2009). As a result, the amount of phytoliths in the calcaric Cambisol and in the hypereutric Cambisol 1166, that had a pH value close to 7 (Table S11), was low. It was also the case for the Albeluvisol despite a soil pH value of 5.2 (Table S11). In this soil, reducing conditions occurring in winter and resulting in a transient high pH (Cornu et al., 2018) may be responsible for an increase in the phytolith dissolution over that period. For the Luvisols and the BS2 hypereutric Cambisol, the amount of phytoliths in forest increased as the pH decreased. The absence of correlation between Si_{CaCl2} and phytoliths suggests that this pool is however not the main pool controlling the Si_{CaCl2} in temperate soils and that another pool of pedogenic origin must exist as already evoked by Vandevenne et al. (2015) and Keller et al. (2021).

Si_o was very strongly correlated with Al_o (Fig. 6a), excluding the cultivated calcaric Cambisol 701 that was very rich in SOC compared to the other cultivated soils (Fig. 2c and discussion below). Contrarily Si_o was not significantly correlated with Fe_o in cultivated plots ($r = 0.6$, p -value = 0.2) or in forests ($r = 0.5$, p -value = 0.3). In addition, the Al_o/Si_o ratio observed for both forest and cultivated soils was close to 2, a typical value for allophanes (Harsh, 2005; Wada, 1989). Hence, it is likely that the Si_o pool consisted in SRO aluminosilicates. SRO aluminosilicates are found in most soils, including those developed on loess and sedimentary rocks as in this study, insofar as their DSI are between 0.1 and 4 mmol L $^{-1}$ (Harsh, 2005). DSI values close to 0.1 mmol L $^{-1}$ were indeed obtained for the soil solution of the BS3 soil (Cornu et al., 2018). In addition, the spherical particles observed by TEM (Fig. 5) are commonly interpreted as allophanes (Kitagawa, 1971; Henmi and Wada, 1976; Parfitt, 1990).





Fig. 3. Relative abundance of (a) grass phytoliths (including RONDEL, CREMATE, ELOBATE, POLYLOBATE, and CROSS silica short cells, silicified epidermal tissues, FLABELLATE BULLIFORM cells, and ELONGATE DENDRITIC), (b) forest indicator phytoliths (including SPHEROIDS, sclereids and other silica bodies observed in woody dicotyledons, two BLOCKY FACETED bodies typical of conifers, and a very long and large ELONGATE phytolith typical of some ferns (see details in S1 Table-Phyto). (c) Relative abundance of ELONGATE DENDRITIC phytoliths typical of grass inflorescences. Bars represent 95% confidence intervals. Percentages were calculated based on (a, b) total sum of particles counted and (c) on total phytolith sum.

Our dataset shows also a strong correlation between the $< 2 \mu\text{m}$ fraction and the allophane contents (Fig. 6b), suggesting that allophanes may either be a product of clay mineral weathering or be part of the clay mineral formation, notably of the finest clays, the smectites and/or interlayered clay minerals that were shown to be present in all the soils considered (Fig. 4). Li et al. (2020a) also observed the formation of allophanes in a pot experiment as the result of weathering. When the Al_0/Si_0 ratio is above 2, which is the case in most forest soils (Table 1 and SI2), the Al_0 content in excess is present as Al-OC complexes (Harsh, 2005). These Al-OC complexes partially preclude the formation of short range ordered aluminosilicates (Harsh, 2005). SiCaCl_2 was well correlated to allophane content (Fig. 7a), excluding the BS2 site for which part of the Si_0 extracted was not contained in allophanes contrarily to all the other sites (as calculated by eq. (2)).

Therefore, allophanes are assumed to control a fraction of SiCaCl_2 in soils. This agrees well with the high SiCaCl_2 content observed in Andosols, which contain a large amount of imogolites and allophanes (Caubet et al., 2020). Caubet et al. (2020) concluded that SiCaCl_2 was controlled by both sorption and dissolution processes, which are strongly controlled by pH for soils with $< 2 \mu\text{m}$ fractions both larger than 50 g kg^{-1} and lower than 325 g kg^{-1} . For Indian soils this relation is not linear but polynomial, with a stabilisation of the SiCaCl_2 concentrations for pH higher than 7 (Meunier et al., 2018). We used a comparable relationship for the soils considered here excluding the BS2 and 62 forest soils (Fig. 7b). The pH value of the 62 forest plot lower than 4.5 suggests the beginning of podzolisation (Caner et al., 2010; Sauer et al., 2008; Fekiacova et al., 2017). The conifer vegetation observed on this plot (Table SI1) is compatible with the onset of such a process. This low pH also corresponds to the upper limit of the Al/Fe oxide buffer range (Stützer, 1998).

Podzolisation induces dissolution of the clay minerals, which may be responsible for the higher SiCaCl_2 concentration observed with respect to the regression model presented in Fig. 7b. The above findings have the following consequences on the terrestrial Si cycle described by Struyf et al. (2010). In all soil types, 99 % of Si was contained in crystalline silicates. However, in soils developed on loess almost 60 % of the crystalline silicates consisted of quartz (e.g., Montagne et al., 2008 for BS3), a poorly weatherable mineral. In the calcaric Cambisols, the total Si content in crystalline silicate was much lower than in soils developed on loess and was mainly contained in clay minerals and some quartz and feldspar. Despite this mineralogical difference, we concluded that the main weatherable mineral compartment contributing to SiCaCl_2 consists in the clay minerals (Fig. 8). This compartment was almost twofold larger in the calcaric Cambisols than in the soils on loess (Fig. 2a). The clay compartment controlled allophanes (Fig. 6b) which in turns controlled the SiCaCl_2 as discussed above.

Allophanes are an easily available source of DSi as shown by dissolution rates much higher than those of phytoliths at pH lower than 9 (Ralston et al., 2021; Fraysse et al., 2009). Consequently, phytoliths did not appear to be the first source of Si in the considered soils, in agreement with Keller et al. (2021), who found that the clay minerals can compete with phytoliths as a main source of plant Si when the proportion of clay is high.



D patterns of the $< 2 \mu\text{m}</math> fraction on oriented slides of the cultivated (in grey) and forest (in orange) soils: a- for the calcaric Cambisol 701; b- the c Cambisol 1166; c- the Luvisol 340; and d- the Albeluvisol.$

Reflecting the pH-dependent dissolution rate of phytoliths (Frayssé et al., 2009), the phytolith pool was lower in calcaric Cambisols (pH around 7, Table S11) than in soils developed on loess (pH around 5, Table S11). Consequently, the higher $\text{Si}_{\text{CaCl}_2}$ in calcaric Cambisols compared to soils on loess could be explained by the higher amount of allophanes in calcaric Cambisol, but also, to a lesser extent, by a higher dissolution of the phytolith pool.

4.2. Impact of land use on the Si biogeochemical cycle and its various mineralogical pools

As expected, cultivation increased the soil pH (Fig. 2b) and decreased the SOC content (Fig. 2c) in all soils. For soils developed on loess, the pH was two units higher under cultivation than under forest because of liming. For calcaric Cambisols, the pH was one unit higher under cultivation than under forest because of the acidifying capacity of the forest (Fig. 2b, Table S11). We confirm here that cultivation increases the amount of $\text{Si}_{\text{CaCl}_2}$ for soils developed on sediments (Caubet et al., 2020; Vandevenne et al., 2015; Klotzbücher et al., 2018).

The change in pH and SOC induced by cultivation may act on the allophanes and phytoliths pools and therefore modify the terrestrial Si cycle. Based on the literature, a lower phytolith content was expected in cultivated soils compared to topsoils under forest (Guntzer et al., 2012; Vandevenne et al., 2015). This was clearly observed for Luvisols, and to a lesser extent for the BS2 hypereutric Cambisol (Fig. 2i). This decrease in phytoliths may be explained by both a larger dissolution of phytoliths under cultivation due to the pH increase and phytolith exportation due to harvesting. The relative abundance of phytoliths (Fig. 3) suggests that wild grasses were the main producers of phytoliths in forest undergrowth. This is in agreement with temperate trees being poor Si accumulators (Bremond et al., 2004).

Cultivation had no clear impact on the phytolith pool in the calcaric Cambisol or in the Albeluvisols, where the amounts of extracted phytoliths were very low. The proportion of corroded phytoliths (ELONGATE and BLOCKY) was higher in the forest 1166 hypereutric Cambisol (Table 1), that also exhibits a lower amount of phytoliths than under cultivation. Differences in surface areas between grass and forest phytoliths might explain the observed proportions with a higher rate of dissolution of forest phytoliths (Wilding and Drees, 1974).

However, this hypothesis is not supported by the laboratory measurements (Frayssé et al., 2009). Crop management through the restitution strategy of inedible plant parts and the crop rotation system may also lead to site fluctuation in exportation, maybe explaining the exception of the 1166 site. Indeed, crop straw recycling in agricultural soils have proven to be an efficient strategy for limiting desilication (Puppe et al., 2021).

By decreasing SOC (Fig. 2c), cultivation promotes the formation of allophanes that are twice as abundant under cultivation as under forest in the calcaric Cambisols and soils developed on loess (Fig. 2c and 5b). We observed Al_0/Si_0 ratios higher than 2 in forest but close to 2 in cultivated soils (Table 1), evidencing a decrease in the Al-OC complexes that prevent allophane formation under forest as discussed above. For the cultivated soils with a pH close to 8, pH may be less favourable to Si adsorption on allophanes that are at

zero point of charges (Harsh, 2005). The high Si_{CaCl2} values may then be explained by Si adsorbed on carbonate surfaces (Caubet et al., 2020).




Fig. 5. TEM image of a few discrete spherical particles interpreted as allophanes (Okagawa, 1971; Hanmi and Wada, 1976; Parfitt, 1990) at the edges of clay particles (see in the circles) for the cultivated BS3 site. Spherical particles have a diameter of 6 nm and are composed of a disc with a white central part interpreted as being the projection of a hollow sphere. It was not possible to observe them individually and they are systematically associated with clay particles.

Table 2
Pearson correlation coefficients (r) and associated p -values of Si_{CaCl2} with the main soil variables analysed for the 21 samples of the seven soil pairs (2 samples per forest and one per crop plot). Coefficients in bold are significant at a 5 % level of confidence.

Variables	< 2 μm	2-20 μm	20-50 μm	50-200 μm	200-2000 μm	SOC	pH		
r	0.498	0.424	0.410	0.297	0.105	0.267	0.614		
p -value	0.025	0.055	0.065	0.192	0.651	0.241	0.005		
Variables	Si_{CaCl2}	Al_{CaCl2}	F_{CaCl2}	Si_{d}	Al_{d}	F_{d}	Si_{tot}	Al_{tot}	F_{tot}
r	0.526	0.261	0.185	0.684	0.261	0.373	0.479	0.452	0.485
p -value	0.014	0.253	0.420	0.001	0.254	0.006	0.028	0.039	0.026
Variables	phytoliths	CEC	CEC _{CaCl2}						
r	0.360	0.602	0.824						
p -value	0.100	0.004	<0.0001						




Fig. 6. Impact of land use on the allophane amount in the paired-sites: a- Al_0 as a function of Si_0 ; b- amount in allophanes as a function of the < 2 μm concentration. Allophane content was calculated using Harsh's (2005) equation: % allophane = $Si_0 / (-0.067 Al_0 / Si_0 + 0.27)$. Forest soils are reported in black and the cultivated soils in grey. Pearson correlation coefficient and p -values provided on the figures were calculated excluding the samples identified by their name (for figure a only). Forested sites correspond to composite values (0-25 cm) calculated considering the thickness of the two sampled horizons (0-5 cm and 5-25 cm), their concentrations in the different elements and their respective bulk densities.

Struyf et al. (2010) observe that cultivation decreases the exportation of DSi at the watershed outlet. They interpret this decrease in exportation by a reduction in the silicate mineral weathering. On the contrary, our results suggest that this decrease might be due to a higher retention of Si in the soils by the increase of allophane formation under cultivation (Fig. 8).




Fig. 7. Soluble Si ($\text{Si}_{\text{CaCl}_2}$) as a function of the allophane amount (a) and pH (b) in forest soils in black and cultivated soils in grey. Allophane content was calculated using Harsh's (2005) equation: $\% \text{ allophane} = \text{Si}_e / (-0.067 \text{ Al}_e / \text{Si}_e + 0.27)$. The grey rectangle corresponds both to the point of zero charge of the allophanes and to the equilibrium with calcite. In both cases, the regression line was performed on all the soils that are not identified. Forest sites correspond to composite values (0–25 cm) calculated considering the thickness of the two sampled horizons (0–5 cm and 5–25 cm), their concentrations in the different elements and their respective bulk densities.




Fig. 8. Adaptation of the Struyf et al. (2010) conceptual model for the impact of land use change from mature forests to long-term cultivated areas on the main pools that control soluble Si in soils developed on loess (left-hand side) and calcareous soils (right-hand side). Boxes represent the main Si pools with the size of the boxes representing the relative size of the stock. Arrows represent fluxes with a thickness depending on their relative importance. Dashed arrows represent irrelevant fluxes. Ph, Si in phytoliths; DSi, soluble Si estimated by $\text{Si}_{\text{CaCl}_2}$; Allo, Si in allophanes.

5. Conclusion

In temperate soils, phytoliths may not be the main Si pool contributing to bioavailable Si in contrast to tropical soils that are strongly depleted in Si and have kaolinite as dominant clay mineral. Temperate soils indeed exhibit a more complex clay mineral assemblage (smectite, illite, kaolinite and interlayered minerals). Allophanes - either precursors or dissolution products of these clay minerals - appear as the main Si pool contributing to DSi, estimated by CaCl_2 extraction. This explains the correlation between $\text{Si}_{\text{CaCl}_2}$ and the clay fraction reported in the literature. Cultivation seems to impact soil Si pools and DSi in two ways. First, cultivation increases pH by liming in the case of non-calcareous soils and by suppressing forest acidification. The increase in pH enhances Si sorption and phytolith dissolution. Secondly, cultivation decreases soil organic carbon, promoting the formation of allophanes. At pH lower than 4.5, other processes occur, such as podzolisation, modifying the relationship between DSi and pH. So far, there has been a lot of emphasis put on phytoliths as a source of DSi. Our work highlights the importance of allophanes as the missing link between the soil solution and the clay minerals that could explain why clay minerals may be so important, despite lower solubility rates than phytoliths.

Declaration of Competing Interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was performed in the framework of the French ANR BioSiSol (ANR-14-CE01-0002) project. The authors are grateful to the CEES team (10.15454/VCZ6BS) (INFOSOL, INRAE, Orléans) for their support in site selection, sampling and bulk density measurements and to B. Angeletti for ICP measurements at CEREGE.

Appendix A. Supplementary data Supplementary data to this article can be found online at <https://doi.org/10.1016/j.geoderma.2022.115722>.

References

- Alexander, E.B., 1980. Bulk Densities of California Soils in Relation to Other Soil Properties. *Soil Sci. Soc. Am. J.* 44 (4), 689–692. <https://doi.org/10.2136/sssaj1980.03615995004400040005x>.
- Alexandre, A., Meunier, J.-D., Colin, F., Koud, J.-M., 1997. Plant impact on the biogeochemical cycle of silicon and related weathering processes. *Geochim. et Cosmochim. Acta* 61 (3), 677–682.

- Baize, D., 2000. Guide des analyses en pédologie, Quae. ed. INRA Editions, Paris (France).
- Balesdent, J., P'etraud, J.P., Feller, C., 1991. Effet des ultrasons sur la distribution granulométrique des matières organiques des sols. *Science du sol* 29 (2), 95–106.
- Bartoli, F., 1983. The biogeochemical cycle of silicon in two temperate forest ecosystems. In: R. Hallberg (ed.) "Environmental Biogeochemistry", Ecol. Bull., Stockholm, 35, 469-476.
- Bremont, L., Alexandre, A., V'ela, E., Guiot, J., 2004. Advantages and disadvantages of phytolith analysis for the reconstruction of Mediterranean vegetation: an assessment based on modern phytolith, pollen and botanical data (Luberon, France). *Rev. Palaeobot. Palynol.* 129 (4), 213–228. <https://doi.org/10.1016/j.revpalbo.2004.02.002>.
- Carey, J.C., Fulweiler, R.W., DeGabriel, J., 2016. Human appropriation of biogenic silicon—the increasing role of agriculture. *Funct. Ecol.* 30 (8), 1331–1339.
- Cassini, C.F., 1750. Composite: Carte de France. Carte de France. Levee par ordre du Roy. Retrieved from. <http://www.davidrumsey.com/xmaps10000.html>.
- Caner, L., Joussein, E., Salvador-Blanes, S., Hubert, F., Schlicht, J.-F., Duigou, N., 2010. Shorttime clay-mineral evolution in a soil chronosequence in Ol'eron Island (France). *J. Plant Nutr. Soil Sci.* 173 (4), 591–600.
- Caubet, M., Cornu, S., Saby, N.P.A., Meunier, J.D., 2020. Agriculture increases the bioavailability of silicon, a beneficial element for crop, in temperate soils. *Sci. Rep.* 10, 19999. <https://doi.org/10.1038/s41598-020-77059-1>.
- Conley, D.J., Likens, G.E., Buso, D.C., Saccone, L., Bailey, S.W., Johnson, C.E., 2008. Deforestation causes increased dissolved silicate losses in Hubbard Brook Experimental Forest. *Glob. Change Biol.* 14, 1–7.
- Cornu, S., Montagne, D., Hubert, F., Barr'e, P., Caner, L., 2012. Evidence of short-term clay evolution in soils under human impact. *C. R. Geosci.* 344 (11-12), 747–757.
- Cornu, S., Samou'elian, A., Ayzac, A., Montagne, D., 2018. Soluble and colloidal translocation of Al, Fe, Si and Mn in an artificially drained French Retisol. *Geoderma* 330, 193–203. <https://doi.org/10.1016/j.geoderma.2018.05.032>.
- Clymans, W., Struyf, E., Govers, G., Vandevenne, F., Conley D.J., 2011. Anthropogenic impact on amorphous silica pools in temperate soils. *Biogeosciences*, 8, 2281–2293, 2011.25.
- Desplanques, V., Cary, L., Mouret, J.-C., Trolard, F., Bourri'e, G., Grauby, O., Meunier, J.-D., 2006. Silicon transfers in a rice field in Camargue (France). *J. Geochem. Explor.* 88 (1-3), 190–193.
- de Tombeur, F., Turner, B.L., Lalibert'e, E., Lambers, H., Mahy, G., Faucon, M.-P., Zemunik, G., Cornelis, J.-T., 2020. Plants sustain the terrestrial silicon cycle during ecosystem retrogression. *Science* 369 (6508), 1245–1248.
- Fekiacova, Z., Vermeire, M.L., Bechon, L., Cornelis, J.T., Cornu, S., 2017. Can Fe isotope fractionations trace the pedogenetic mechanisms involved in podzolization? *Geoderma* 296, 38–46. <https://doi.org/10.1016/j.geoderma.2017.02.020>.
- Filippi P., Minasny B., Cattle S.R., Bishop T.F.A., 2016. Chapter Four - Monitoring and Modeling Soil Change: The Influence of Human Activity and Climatic Shifts on Aspects of Soil Spatiotemporally. In *Advances in Agronomy*, eds Sparks D.L., 139, 153-214, 10.1016/bs.agron.2016.06.001.
- Fraysse, F., Pokrovsky, O.S., Schott, J., Meunier, J.D., 2009. Surface chemistry and reactivity of plant phytoliths in aqueous solutions. *Chem. Geol.* 258 (3–4), 197–206. <https://doi.org/10.1016/j.chemgeo.2008.10.003>.
- Georgiadis, A., Sauer, D., Herrmann, L., Breuer, J., Zarei, M., Stahr, K., 2013. Development of a method for sequential Si extraction from soils. *Geoderma* 209-210, 251–261.
- Guntzer, F., Keller, C., Poulton, P.R., McGrath, S.P., Meunier, J.-D., 2012. Long-term removal of wheat straw decreases soil amorphous silica at Broadbalk, Rothamsted. *Plant Soil* 352 (1-2), 173–184.
- Harsh, J., 2005. In: *Encyclopedia of Soils in the Environment*. Elsevier, pp. 64–71. <https://doi.org/10.1016/B0-12-348530-4/00207-1>.
- Haysom, M.B.C., Chapman, L.S., 1975. Some aspects of the calcium silicate trials at Mackay. *Proc. Qld. Soc. Sugar Cane Technol.* 42, 117–122.
- Henmi, T., Wada, K., 1976. Morphology and composition of allophane. *Am. Mineral.* 61, 379–390.
- Hodson, M. J., White, P. J., Mead, A., Broadley, M. R., 2005. Phylogenetic variation in the silicon composition of plants. *Annals of Botany* 96, 1027–1046.
- International Committee for Phytolith Taxonomy (ICPT), Neumann, K., Str'omberg, C.A. E., Ball, T., Albert, R.M., Vrydaghs, L., Cummings, L.S., 2019. International Code for Phytolith Nomenclature (ICPN) 2.0. *Annals of Botany*. 10.1093/aob/mcz064.
- Jolivet, C., Boulonne, L., Rati'e, C., 2006. Manuel du Réseau de Mesures de la Qualité des Sols.
- Jouquet, P., Jamoteau, F., Majumdar, S., Podwojewski, P., Nagabovanalli, P., Caner, L., Barboni, D., Meunier, J.D., 2020. The distribution of silicon in soils influenced by termite bioturbation in South Indian forest soils. *Geoderma*. <https://doi.org/10.1016/j.geoderma.2020.114362>.
- Katz, O., 2015. Silica phytoliths in angiosperms: Phylogeny and early evolutionary history. *New Phytol.* 208 (3), 642–646.
- Keller, C., Rizwan, M., Meunier, J.D., 2021. Are clay minerals a significant source of Si for crops? A comparison of amorphous silica and the roles of the minerals type and pH. *Soil Silicon*. 10.1007/s12633-020-00877-5.
- Kitagawa, Y., 1971. The "unit particle" of allophane. *Am. Mineral.* 56, 465–475.
- Klotzbücher, T., Klotzbücher, A., Kaiser, K., Merbach, I., Mikutta, R., 2018. Impact of agriculture practices on plant-available silicon. *Geoderma* 331, 15–17.
- Kodama, H., Ross, G.J., 1991. Tiron dissolution method used to remove and characterize inorganic components in soils. *Soil Sci. Soc. Am. J.* 55, 1180–1187.
- Landr'e, A., Cornu, S., Meunier, J.-D., Guerin, A., Arrouays, D., Caubet, M., Rati'e, C., Saby, N.P.A., 2020. Do climate and land use affect the pool of total silicon concentration? A digital soil mapping approach of French topsoils. *Geoderma* 364, 114175. <https://doi.org/10.1016/j.geoderma.2020.114175>.
- Li, Z., Cornelis, J.T., Vander Linden, C., Van Ranst, E., Delvaux, B., 2020a. Neoformed aluminosilicates and phytogenic silica are competitive sinks in the silicon soil-plant cycle. *Geoderma* 114308. Li, Z., de Tombeur, F., Linden, C.V., Cornelis, J.-T., Delvaux, B., 2020b. Soil microaggregates store phytoliths in a sandy loam. *Geoderma* 360, 114037.
- Liang, Y., Nikolic, M., B'elanger, R., Gong, H., Song, A., 2015. Silicon in Agriculture. (Springer Netherlands). 10.1007/978-94-017-9978-2.
- Mehra, O.P., Jackson, M.L., 1960. Iron Oxide Removal from Soils and Clays by a Dithionite-Citrate System Buffered with Sodium Bicarbonate. *Clays Clay Miner.* 7, 317–327.
- Meunier, J.-D., Sandhya, K., Prakash, N.B., Borschneck, D., Dussouillez, P., 2018. pH as a proxy for estimating plant-available Si? A case study in rice fields in Karnataka (South India). *Plant Soil* 432 (1-2), 143–155.
- Miles, N., Manson, A.D., Rhodes, R., van Antwerpen, R., Weigel, A., 2014. Extractable Silicon in Soils of the South African Sugar Industry and Relationships with Crop Uptake. *Commun. Soil Sci. Plant Anal.* 45 (22), 2949–2958.
- Montagne, D., Cousin, I., Cornu, S., 2016. Changes in the pathway and the intensity of albic material genesis: Role of agricultural practices. *Geoderma* 268, 156–164.
- Montagne, D., Cornu, S., Le Forestier, L., Hardy, M., Josi'ere, O., Caner, L., Cousin, I., 2008. Impact of drainage on soil-forming mechanisms in a French Albelvisol: input of mineralogical data in mass-balance modelling. *Geoderma* 145 (3-4), 426–438.
- Narayanaswamy, C., Prakash, N.B., 2010. Evaluation of selected extractants for plant-available Silicon in Rice Soils of Southern India. *Commun. Soil Sci. Plant Anal.* 41 (8), 977–989. <https://doi.org/10.1080/00103621003646063>.
- Parfitt, R.L., 1990. Allophane in new Zealand - a review. *Aust. J. Soil Res.* 28, 343–360.
- Puppe, D., Ehrmann, O., Kaczorek,

- D., Wanner, M., Sommer, M., 2015. The protozoic Si pool in temperate forest ecosystems—quantification, abiotic controls and interactions with earthworms. *Geoderma* 243-244, 196–204.
- Puppe, D., Kaczorek, D., Schaller, J., Barkusky, D., Sommer, M., 2021. Crop straw recycling prevents anthropogenic desilication of agricultural soil-plant systems in the temperate zone—Results from a long-term field experiment in NE Germany. *Geoderma* 403, 115187. <https://doi.org/10.1016/j.geoderma.2021.115187>.
- Ralston, S.J., Hausrath, E.M., Tschauer, Oliver, Rampe, Elizabeth, Peretyazhko, T.S., Christoffersen, ROY, Defelice, Chris, Lee, Hyejeong. 2021. Dissolution rates of allophane with variable Fe contents: implications for aqueous alteration and the preservation of X-Ray amorphous materials on Mars. *Clays Clay Miner.* 69 (2), 263–288.
- Robert, M., Tessier, D., 1974. Méthode de préparation des argiles des sols pour des études minéralogiques. *Ann. Agron.* 25, 859–882.
- Saccone, L., Conley, D.J., Koning, E., Sauer, D., Sommer, M., Kaczorek, D., Blecker, S.W., Kelly, E.F., 2007. Assessing the extraction and quantification of amorphous silica in soils of forest and grassland ecosystems. *Eur. J. Soil Sci.* 58 (6), 1446–1459.
- Sauer, D., Schüllli-Maurer, I., Sperstad, R., Sørensen, R., Stahr, K., 2008. Podzol development with time in sandy beach deposits in southern Norway. *J. Plant Nutr.* 171 (4), 483–497.
- Schaller, J., Puppe, D., Kaczorek, D., Ellerbrock, R., Sommer, M., 2021. Silicon Cycling in Soils Revisited. *Plants* 10, 295. <https://doi.org/10.3390/plants10020295>.
- Sommer, M., Kaczorek, D., Kuzyakov, Y., Breuer, J., 2006. Silicon pools and fluxes in soils and landscapes—a review. *J. Plant Nutr. Soil Sci.* 169 (3), 310–329. <https://doi.org/10.1002/jpln.200521981>.
- Struyf, E., Smis, A., Van Damme, S., Garnier, J., Govers, G., Van Wesemael, B., Conley, D. J., Batelaan, O., Frot, E., Clymans, W., Vandevenne, F., Lancelot, C., Goos, P., Meire, P., 2010. Historical land use change has lowered terrestrial silica mobilization. *Nat. Commun.* 1 (1) <https://doi.org/10.1038/ncomms1128>.
- Stützer, A., 1998. Early stages of podzolisation in young aeolian sediments, western Jutland. *Catena* 32 (2), 115–129. Tamm, O., 1922. Eine Method zur Bestimmung der anorganischen Komponenten des Golkomplex in Boden. *Medd. Statens skogforsöksanst.* 19, 385–404.
- Vandevenne, F.I., Barão, L., Ronchi, B., Govers, G., Meire, P., Kelly, E.F., Struyf, E., 2015. Silicon pools in human impacted soils of temperate zones. *Global Biogeochem. Cycles* 29 (9), 1439–1450. <https://doi.org/10.1002/2014GB005049>.
- WRB, 2006. World reference base for soil resources. *World Soil Resources Reports*, 103. FAO, Rome.
- Wada, K., 1989. Allophane and imogolite, in: *Minerals in Soil Environments*, Dixon, J. B., Weed, S. B., & Wada, K. eds. Soil Science Society of America, Madison, USA, 1051–1087. 10.2136/sssabookser1.2ed.c21.
- Wilding, L.P., Drees, L.R., 1974. Contributions of Forest Opal and Associated Crystalline Phases to Fine Silt and Clay Fractions of Soils. *Clays Clay Miner.* 22, 295–306. <https://doi.org/10.1346/CCMN.1974.0220311>.
- Wojdyr, M., 2010. Fityk: a general-purpose peak fitting program. *J. Appl. Cryst.* 43 (5), 1126–1128.
- Yanai, J., Taniguchi, H., Nakao, A., 2016. Evaluation of available silicon content and its determining factors of agricultural soils in Japan. *Soil Sci. Plant Nutr.* 62 (5-6), 511–518.