

The periodic motion of a disk freely falling in a tube Patricia Ern, Nicolas Brosse

▶ To cite this version:

Patricia Ern, Nicolas Brosse. The periodic motion of a disk freely falling in a tube. 9th Euromech Fluid Mechanics Conference (EFMC9), Sep 2012, Rome, Italy. pp.0. hal-04019436

HAL Id: hal-04019436 https://hal.science/hal-04019436

Submitted on 8 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <u>http://oatao.univ-toulouse.fr/</u> Eprints ID: 10391

Official URL: http://www.euromech.org/conferences/EFMC/EFMC9

To cite this version:

Ern, Patricia and Brosse, Nicolas *The periodic motion of a disk freely falling in a tube*. (2013) In: 9th Euromech Fluid Mechanics Conference (EFMC9), 09 September 2012 - 13 September 2012 (Rome, Italy).

The periodic motion of a disk freely falling in a tube

P. Ern^{*}, N. Brosse^{*}

This study is devoted to the rectilinear and periodic paths of an axisymmetric solid body (short-length cylinder and disk of diameter d, thickness h and aspect ratio $3 \leq \chi = d/h \leq 10$) falling in a vertical tube of diameter D. Three-dimensional trajectography was used to record the body motion (see figure 1). We investigated the influence of the confinement ratio (S = d/D) on the characteristics of the body motion, for different aspect ratios and Reynolds numbers (80 < Re < 320), and a density ratio between the fluid and the body close to unity. The critical Reynolds number for the onset of the periodic motion decreases with S in the case of thin bodies $(\chi = 10)$, whereas it appears unaffected by S for thicker bodies ($\chi = 3$ and 6). For all aspect ratios, the mean fall velocity of the body decreases when S increases. The characteristics of the oscillatory motion are also strongly modified by the confinement ratio. A thick body ($\chi = 3$) tends to stabilize and to go back to a rectilinear path when the confinement ratio increases, while a thin body ($\chi = 10$) displays oscillations of growing amplitude with S until it touches the tube (at about S = 0.5). However, the amplitudes of the oscillatory motion of bodies with a given aspect ratio gather on a unique curve for all S (including the case with no confinement, S = 0), when plotted as a function of the relative distance of the Reynolds numbers of the bodies to the threshold of path instability for each confinement ratio S.

^{*}Institut de Mécanique des Fluides de Toulouse, Université de Toulouse and CNRS, Allée du Prof. C. Soula, 31400 Toulouse, France




Figure 1: Three-dimensional view of the path of a body of aspect ratio $\chi = 3$ falling in a tube with S = 0.32 and Re = 245. The grid represents the wall of the tube.