



HAL
open science

Cancer Among Adolescents and Young Adults Between 2000 and 2016 in France: Incidence and Improved Survival

Thomas Raze, Brigitte Lacour, Anne Cowppli-Bony, Patricia Delafosse, Michel Velten, Brigitte Trétarre, Gautier Defossez, Karima Hammas, Anne-Sophie Woronoff, Olivier Ganry, et al.

► To cite this version:

Thomas Raze, Brigitte Lacour, Anne Cowppli-Bony, Patricia Delafosse, Michel Velten, et al.. Cancer Among Adolescents and Young Adults Between 2000 and 2016 in France: Incidence and Improved Survival. *Journal of Adolescent and Young Adult Oncology*, 2021, 10 (1), pp.29-45. 10.1089/jayao.2020.0017 . hal-04020187

HAL Id: hal-04020187

<https://hal.science/hal-04020187v1>

Submitted on 8 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**CANCER AMONG ADOLESCENTS AND YOUNG ADULTS BETWEEN 2000 AND 2016 IN FRANCE:
INCIDENCE AND IMPROVED SURVIVAL**

Thomas Raze, MD,¹, Brigitte Lacour, MD,^{1,2}, Anne Cowppli-Bony, MD,³, Patricia Delafosse, MD,⁴, Michel Velten, MD, PhD,⁵, Brigitte Trétarre, MD,⁶, Gautier Defossez, MD,⁷, Karima Hammas, MD,⁸, Anne-Sophie Woronoff, MD,⁹, Olivier Ganry, MD, PhD,¹⁰, Sandrine Plouvier, MD,¹¹, Gaëlle Coureau, MD,¹², Anne-Valérie Guizard, MD,¹³, Simona Bara, MD,¹⁴, Alain Monnereau, MD, PhD,¹⁵, Laetitia Daubisse-Marliac, MD,¹⁶, Xavier Troussard, MD, PhD,¹⁷, Tania D'Almeida, MD,¹⁸, Isabelle Baldi, MD, PhD,¹⁹, Véronique Bouvier, MD,²⁰, Nicolas Boissel, MD, PhD,²¹, Jacqueline Clavel, MD, PhD,^{2,22}, Emmanuel Desandes, MD,^{1,2}

¹Registre National des cancers de l'Enfant, Registre National des Tumeurs Solides de l'Enfant, CHRU Nancy, Vandœuvre-lès-Nancy, France

²Centre de Recherche en Epidémiologie et en Statistique Sorbonne-Paris Cité (CRESS), UMR 1153, INSERM, Université Paris-Descartes, Paris, France

³Registre des tumeurs de Loire-Atlantique et Vendée, CHU de Nantes, SIRIC-ILIAD, Nantes, France

⁴Registre Général des Cancers de l'Isère, Meylan, France

⁵Registre des cancers du Bas-Rhin, Laboratoire d'épidémiologie et de santé publique, Université de Strasbourg, Inserm UMR-S1113/Centre Paul Strauss, Service de santé publique, Strasbourg, France

⁶Registre des Tumeurs de l'Hérault, Montpellier, France

⁷Registre des Cancers de Poitou-Charentes, CHU Poitiers, France

⁸Registre des cancers du Haut-Rhin, ARER68, Groupe hospitalier de la région de Mulhouse et Sud-Alsace (GHRMSA), Mulhouse, France

⁹Registre des tumeurs du Doubs, EA3181, CHRU Besançon, France

- ¹⁰Registre Général du Cancer de la Somme, CHU Amiens, France
- ¹¹Registre général des cancers de Lille et de sa région, GCS-C2RC, Lille, France
- ¹²Université Bordeaux, Epicene, centre Inserm U1219/CHU de Bordeaux, service d'information médicale, Bordeaux, France
- ¹³Registre Général des Tumeurs du Calvados, Caen, France
- ¹⁴Registre des Cancers de la Manche, Cherbourg-Octeville, France
- ¹⁵Institut Bergonié, registre des hémopathies malignes de la Gironde/Université de Bordeaux, Inserm U1219, équipe EPICENE, ISPED, Bordeaux, France
- ¹⁶Institut Claudius Regaud, IUCT-O, Registre des cancers du Tarn, CHU Toulouse/ UMR 1027 Université de Toulouse, UPS, Inserm, Toulouse, France
- ¹⁷Registre des Hémopathies Malignes de Basse Normandie, Caen, France
- ¹⁸Registre général des cancers en Région Limousin, CHU de Limoges, France
- ¹⁹Equipe EPICENE, INSERM U1219, Université de Bordeaux/Service Santé Travail Environnement, CHU Bordeaux, France
- ²⁰Registre des Tumeurs Digestives du Calvados, Caen, France
- ²¹Unité d'Hématologie Adolescents et Jeunes Adultes, Hôpital Saint-Louis, Assistance Publique des Hôpitaux de Paris/EA-3518, Institut de Recherche Saint-Louis, Université de Paris, France
- ²²Registre National des Hémopathies Malignes de l'Enfant, Villejuif, France

Corresponding author

Emmanuel Desandes, MD, Registre National des cancers de l'Enfant, Registre National des Tumeurs Solides de l'Enfant, Université de Lorraine, Faculté de médecine, 9 Avenue de la Forêt de Haye, BP 20199, 54505 Vandœuvre-lès-Nancy cedex, France (tel: +33(0)372746328, fax: +33(0)372746332, Email: emmanuel.desandes@univ-lorraine.fr)

Funding

The current work is funding by the French Public Health Agency « Santé Publique France » (project number: 18DMNA53-0).

Keywords

Adolescents, Young Adults, Cancer, Incidence, Survival, Trends

Running Head

Cancers in French AYA, 2000-2016

Count

Text word count: 2,901

Abstract word count: 250

Tables: 5

Figures: 1

Number of references : 41

1 **CANCER AMONG ADOLESCENTS AND YOUNG ADULTS BETWEEN 2000 AND 2016 IN FRANCE:**
2 **INCIDENCE AND IMPROVED SURVIVAL**

3 **ABSTRACT**

4 **Purpose:** This study was undertaken to determine cancer survival and describe the spectrum
5 of cancers diagnosed among French adolescent and young adult (AYA) population.

6 **Methods:** All cases of cancer diagnosed in 15-24 years, recorded by all French population-
7 based registries (18% of the French population), over the 2000-2016 period, were included.

8 Age-standardized incidences rates (ASR), conventional annual percentage change (cAPC) of
9 incidence over time, and 5-year overall survival (5yOS) were calculated.

10 **Results:** We analyzed 2,734 cancer diagnoses in adolescents and 4,199 in young adults.

11 Overall incidence rates were 231.9/10⁶ in 15-19 year olds and 354.0/10⁶ in 20-24 year olds.

12 The most frequently diagnosed cancers in male AYA were malignant gonadal germ-cell
13 tumors (GCT), Hodgkin lymphoma (HL), and malignant melanoma, and were HL, thyroid

14 carcinoma, and malignant melanoma in females. Cancer incidence was stable over time with a
15 cAPC of 0.8% (p=0.72). For all cancers combined, 5yOS was 86.6% (95%CI: 85.8-87.4),

16 greater than 85% for HL, non-Hodgkin lymphomas (NHL), GCT, thyroid carcinomas, and
17 malignant melanomas, and around 60% and lower for osteosarcomas, Ewing tumors, hepatic

18 carcinomas, and rhabdomyosarcomas. The 5yOS has significantly improved from 2000-2007
19 to 2008-2015 for all cancers pooled, with a substantial gain of 4% for 15-19 year-olds, and

20 3% for 20-24 year-olds.

21 **Conclusion:** Notwithstanding the encouraging results for some cancers, and overall,
22 persistent poorer survivals in AYA were shown compared to children for acute lymphoblastic

23 leukemia, osteosarcoma, Ewing tumor, rhabdomyosarcoma and malignant hepatic tumors.

24 These disparities require further investigation to identify and address the causes of these
25 inferior outcomes.

26

27 **Introduction**

28 Incidence and survival within the adolescents and young adults (AYA) population can be
29 misleading when assessed across all tumor streams. The latest French report (2000-2008) ¹
30 described an overall incidence rate (IR) 219.4 per million adolescents (age 15-19 years) and
31 293.1 per million young adults (age 20-24 years). For all cancers combined, substantial
32 survival gains were achieved, with an increase of 5-year overall survival (OS) from 62.0%
33 (95%CI: 57.5-66.5) in the 1978-1982 period to 81.8% (95%CI: 79.5-83.9) in the 2000-2004
34 period. An improvement in survival was found for leukemia, lymphoma, and central nervous
35 system (CNS) tumors in subgroup analysis. Conversely, no increase in survival over time was
36 observed for malignant bone tumors.^{1,2} AYA had significantly worse 5-year OS, during the
37 2000s, than children (0-14 years) for acute lymphoblastic leukemia (ALL, 62.8% vs. 89.8%),
38 non-Hodgkin lymphomas (NHL, 78.9% vs. 88.3%), CNS tumors (67.3% vs. 71.9%), and
39 bone sarcomas (50.0% vs. 74.4%).^{1,3} To improve cancer outcomes for AYA, various
40 initiatives have been implemented in France since 2011,⁴ including promoting collaboration
41 between pediatric and adult oncologists, developing French national policies for managing
42 AYA with cancer, and setting up specific treatment units. Nevertheless it is unclear whether
43 the adoption of pediatric-based protocols in adult centers has abolished disparities in AYA
44 outcomes between pediatric and adult centers and improved outcomes at a population-level.
45 Using a population-based cohort of AYA in France, the objectives were to provide updated
46 statistics covering incidence and survival and to describe incidence and survival trends of
47 these patients from 2000 to 2016.

48

49 **Methods**

50 *Source of data*

51 The registration of cancers in AYA is ensured by eighteen general and specialized
52 population-based registries of the Francim network, covering 18% of the French population.
53 Medico-administrative sources (health institution claims databases of public or private health
54 institutions used to obtain hospital patient information coded on discharge, and health
55 insurance schemes databases) and medical sources (multidisciplinary team meetings
56 databases in oncology, histology reports databases) were used to identify cases. Vital status
57 was obtained by active search in medical records and by matching the French Cancer
58 Registries files and the national population register “Repertoire National d’Identification des
59 Personnes Physiques” sorted by date and place of birth to obtain the mention of death and the
60 date of death. The cut-off date was defined as June 30, 2018.

61

62 *Population*

63 Data were taken from the common database of all Francim network registries, and included
64 gender, age at diagnosis, and diagnostic information. In line with the recommendations of the
65 European Network of Cancer Registries⁵ and the third version of the International
66 Classification of Childhood Cancer (ICCC-3),⁶ eligibility criteria were: diagnosis of cancer,
67 for CNS localization according to the first revision of the third edition of the International
68 Classification of Disease for Oncology (ICDO-3),⁷ benign and borderline tumors were also
69 considered; from 15 to 24 years of age at diagnosis; and period of diagnosis between January
70 1, 2000 and December 31, 2016. Cutaneous carcinomas were excluded.

71

72 *Statistical Analyses*

73 All data were retrieved, tabulated by tumor type and categorized using the ICCC-3 levels: 12
74 main diagnostic groups, 47 diagnostic subgroups, and 2-11 divisions of selected subgroups.⁶
75 The males/females (M/F) ratio was the ratio of number of cases in males to that in females.

76 The IR were calculated for the 17-year period, 2000-2016, and based on person-years derived
77 from the annual estimates of population by age and sex provided by the Institute of Statistics
78 and Economic Studies. Crude and age-specific IR were calculated and age-standardized rates
79 (ASR) were estimated by the direct method using the world population for the age 15–24
80 years.⁸ Methods based on Poisson distribution were used to specify the 95% confidence
81 interval (95%CI) for the IR.⁹ Time trend in the incidence was estimated by the conventional
82 annual percent change (cAPC) calculated as the slope of the linear regression used to model
83 the natural logarithm of the IR as a function of the calendar year.¹⁰ Survival analysis was
84 performed on a dataset containing all cases diagnosed between January 1, 2000 and December
85 31, 2015 and followed-up until June 30, 2018. The probability of OS was estimated using the
86 Kaplan–Meier method.¹¹ The endpoint of interest was death from any cause for OS, with the
87 date of diagnosis acting as the time origin. Survival curves between groups were compared
88 using the log-rank test and the trend in survival over time periods was estimated using the log-
89 rank test for trend.^{12,13} Cox models were used to calculate the hazard ratios of death (HRs)
90 and their 95%CI.¹⁴ Statistical tests were two-sided with a significant level of 0.05. All
91 statistical analyses were performed using the SAS version 9.4 (SAS Institute Inc., Cary, North
92 Carolina, USA, 2013).

93

94 **Results**

95 *Patient characteristics*

96 Table 1 shows the number of cancers in AYA, diagnosed from 2000 to 2016, with main data
97 quality indicators by French administrative areas where a French Cancer Registry is
98 established. Most cancers were microscopically verified (98.6%) and only 2% had
99 unspecified morphology. A total of 2,734 15-19-year-old adolescents (39.4%) and 4,199 20-
100 24-year-old young adults (60.6%) were included. Table 2 shows the distribution of tumors by

101 age group and diagnostic group and the M/F ratios for each tumor type. The most common
102 diagnoses in 15-19 year olds were lymphomas (27.7%): Hodgkin lymphoma (HL) 20.4%,
103 NHL 6.0%. In young adults (20-24 years), the most frequent tumors were malignant epithelial
104 tumors (33.8%): thyroid carcinomas (13.5%), malignant melanomas (10.2%). In 15-24-year-
105 olds, approximately 30.7% of cancers were tumors of adulthood, especially thyroid
106 carcinomas 11.6% and malignant melanomas 8.5%. Embryonal malignancies, including
107 Wilms tumor, hepatoblastoma, neuroblastoma, and medulloblastoma, were rare accounting
108 for less than 1% of cases. For all cancers combined, the M/F ratios were 1.0 for adolescents,
109 for young adults, and for both. In male AYA, the most common cancers were malignant
110 germ-cell tumors (GCT), malignant bone tumors, soft-tissue sarcomas, and leukemia, with
111 M/F ratio of 5.0, 1.5, 1.4, and 1.4 respectively. Conversely, malignant melanomas and thyroid
112 carcinomas were more frequent in female AYA (M/F ratio 0.6 and 0.2, respectively).

113

114 *Incidence*

115 Table 2 shows the specific IR and ASR from 2000 to 2016 for the main tumor types by age
116 group, computed with both genders. The ASR for the 15-24-year-old population was 289.4
117 per million (95%CI: 279.8-299.2). The incidence (rate per million) was lowest in 15-19
118 year olds: 232.2 (males) and 231.7 (females). This incidence increased to 351.4 (males) and
119 372.2 (females) in 20-24 year olds. The highest ASR in males aged from 15 to 24 years was
120 seen for malignant gonadal GCT (ASR = 89.2, 95%CI: 82.8-96.1). HL was the second most
121 frequent tumor type in males (ASR = 51.7, 95%CI: 46.8-56.9), followed by malignant
122 melanoma (ASR = 21.7, 95%CI: 18.5-25.2). In AYA females, HL represented the most
123 frequent cancer type with an ASR of 58.6(95%CI: 52.6-65.2), thyroid carcinoma the second
124 (ASR=54.1, 95%CI: 48.4-60.2), and malignant melanoma the third (ASR=30.7 95%CI: 26.6-
125 35.4). Cancer incidence in AYA for both genders was stable over time with a cAPC of 0.8%

126 (95%CI: -3.5%;+5.3%, p=0.72, Fig.1). No variation in incidence was observed by age group,
127 gender, or each ICCC-3 diagnostic group.

128

129 *Overall Survival*

130 Table 3 describes the 1-, 2-, and 5-year OS among AYA with cancer from 2000 to 2015 by
131 ICCC-3 diagnostic groups and subgroups (n=6,480). With 2.9% lost to follow-up (Table 1),
132 the median follow-up time in the cohort was 103 months (min-max: 0-222 months). At the
133 end time, 983 AYA had died with a median time after diagnosis of 21 months (min-max: 0-
134 194 months). For all cancers combined, OS was 95.3% (95%CI: 94.8-95.8) at 1 year, 91.3%
135 (95%CI: 90.6-92.0) at 2 years, and 86.6% (95%CI: 85.8-87.4) at 5 years. The 5-year OS
136 greater than 85% were observed for HL, NHL, GCT, thyroid carcinomas, cervix uteri
137 carcinomas, and malignant melanomas whereas 5-year OS was around 60% and lower for
138 osteosarcomas, Ewing tumors, hepatic carcinomas, and rhabdomyosarcomas. The 5-year OS
139 were close to 65-80% for ALL, acute myeloid leukemia (AML), CNS tumors, and malignant
140 renal tumors.

141 A comparison of 5-year OS in the studied periods (2000-2007 and 2008-2015) with those in
142 earlier time periods (1972-1982, 1983-1987, 1988-1992, and 1993-1997 data from reference²)
143 indicated that substantial survival gains were achieved for all tumors pooled (Table 4), with
144 an increase from 62.0% (95%CI:57.5-66.5) in the 1978-82 period to 88.0% (95%CI:86.9-
145 89.0) in the 2008-15 period. Since 1997, survival improvement was especially seen in
146 leukemia and CNS tumors. For malignant bone tumors, the increase in survival was only
147 observed during the last analyzed period (2008-15). Since 1993, survival was stable for soft-
148 tissue sarcomas.

149 The 5-year OS has significantly improved from 2000-2007 to 2008-2015 for all cancers
150 pooled, with a substantial gain of about 4% for 15-19 year-olds, and 3% for 20-24 year-olds

151 (Table 5). Increases in 5-year OS concerned most of diagnostic groups and were especially
152 apparent for osteosarcomas in both age groups ($p<0.01$), and carcinomas in 20-24 year-olds
153 ($p=0.04$). Conversely, a significant decrease over time was observed for astrocytomas in 20-
154 24 year-olds ($p=0.01$), due to differences in percentage of pilocytic astrocytoma and
155 glioblastoma, respectively 51% vs. 12% during the 2000-2007 period, and 24% vs. 45%
156 during the 2008-2015 period.

157 During the 2000-2007 period, there were no statistically significant differences in 5-year OS
158 between adolescents (15-19 years) and young adults (20-24 years) whatever the main
159 diagnostic group (Table 5). However, during the latest studied period, young adults (20-24
160 years) diagnosed with ALL, astrocytoma, or Ewing's sarcoma had significantly a higher risk
161 of dying from the disease compared to adolescents (15-19 years): HR=2.0 (IC95%: 1.1-3.5),
162 HR=2.2 (IC95%: 1.2-3.9), and HR=2.3 (IC95%: 1.1-4.8), respectively. The histologic
163 distribution of pilocytic astrocytoma and glioblastoma were significantly different between
164 adolescents and young adults: 61% and 12% in 15- to 19-year-olds, vs. 24% and 45% in 20-
165 to 24-year-olds ($p<0.01$).

166

167 **Discussion**

168 Considering the emergence of interest in oncology in the AYA group, we examined incidence
169 and survival of 15-24 year olds in order to better define the burden of the cancer for this age
170 group. The two major findings in the present study were that cancer incidence in AYA
171 remained stable during the 2000-2016 period, and the 5-year OS combined has significantly
172 improved over time, especially for osteosarcoma. Another major finding of our study was
173 that, during the latest studied period, adolescents had significantly better 5-year OS than
174 young adults for ALL, Ewing's sarcomas, and astrocytomas.

175 The major strength of this study is its homogenization of compulsory registration based to
176 various sources of notification of 18 population-based cancer registries providing a nonbiased
177 and valid view of the situation in France during the studied period, compared to clinical series
178 which report better survival than population-based studies. The coverage is 18% of the whole
179 country. However, these registries have provided the estimates of national rates for decades
180 and they are the gold standard referred to by all descriptive studies. Roughly 2,300 new cases
181 of cancers in AYA are currently diagnosed each year in France (900 in adolescents and 1,400
182 in young adults).

183 Several studies have described cancer incidence and survival in AYA, but few of them were
184 based on the same period and the same age ranges.^{10,15-21} Our incidence rates in 15-19 year
185 olds were comparable to those reported in other countries.^{10,15-18} In 20-24 year olds, highest
186 incidence was observed in the United States (US): 386 per million.¹⁰ Marked differences
187 between France and the US occurred for thyroid carcinoma (IR per million: 78 versus 122 for
188 young females, no difference for males) and for melanoma (IR per million: 47 versus 57 for
189 young white females, no difference for males). In the US, the incidence of thyroid carcinoma
190 in AYA population increased by 4.0% per year from 1988 to 2006, 8.6% per year from 2006
191 to 2009, and 1.4% from 2009 to 2016, likely a consequence of ultrasound screening.²² For
192 melanoma, this suggests that American white female young adults may share high exposures
193 that are associated with malignant melanoma. Jones et al²³ have reported that, during 1999-
194 2009, the percentage of white American students who never or rarely wore sunscreen when
195 outside on a sunny day for >1 hour increased from 57.5% to 69.4% (with an increase most
196 pronounced among white female students). We reported a non-significant increase of +0.8 in
197 incidence for all malignancies over time in 15- to 24-year olds. Vast differences between
198 countries were observed in the incidence trend for all cancers combined. Gupta et al²⁴ have
199 reported that the incidence of AYA (15-39 years) cancer rose over the 1998-2012 period in 23

200 countries, decreased in two countries, and remained stable in the other 16. Striking increases
201 were noted for thyroid and testicular in 33 and 22 countries respectively, including France.
202 The 5-year observed survivals in this study were similar to those reported for AYA in the US
203 (2009-2015)¹⁰, in Australia (2000-2014)¹⁸, in the United-Kingdom (UK: 2005-2010)¹⁹, and in
204 27 European countries participating in EUROCORE-5 (2000-2007)²¹. However marked
205 differences between France and the UK occurred for NHL (88% vs. 79%), bone sarcomas
206 (65% vs. 56%), soft-tissue sarcomas (67% vs. 61%), and GCT (93% vs. 82%). Compared to
207 EUROCORE-5 study²¹, we reported higher survival for ALL, CNS tumors, Ewing's sarcoma,
208 and soft-tissue sarcomas especially in 15-19 years. These survival disparities may be due to
209 differences in tumor classification used (ICDO-2 vs. ICDO-3), in tumor selection (inclusion
210 or not of benign/borderline CNS tumors), in distribution of cases by stages at diagnosis, by
211 histological or biological subtypes, or by grades of neoplasm, and in treatment strategies by
212 age over time.

213 Compared to results of the latest French reports among AYA,^{1,2} we found an improvement in
214 survival from 1978 to 2015, to achieve a 5-year OS of 88.0% (95%CI: 86.9-89.0) for all
215 cancers combined during the 2008-2015 period. Increases in survival over time have been
216 observed for all diagnostic groups, except for soft-tissue sarcomas with a 5-year OS around
217 70% since the mid-90s. Compared to children and older adults, AYA with soft-tissue
218 sarcomas have not experienced an improvement in survival.^{25,26} The reasons for this disparity
219 remain unclear, but following issues have been mentioned: delays in diagnosis, trial
220 availability and participation, aspects of the organization of care, national centralization of
221 sarcoma care, international consortia, and factors related to tumor biology.²⁷ In our series,
222 comparison of 5-year OS of osteosarcoma AYA treated during 2000 to 2007 and 2008 to
223 2015 revealed a significant improvement of 14% in adolescents and 40% in young adults.
224 One of the most crucial factors contributing to the increase in survival is the setting up of the

225 French OS2006/sarcome-09 study using methotrexate-etoposide-ifosfamide based regimen,
226 randomized to receive or not high doses of zoledronic acid adapted from those given for
227 pediatric osteosarcoma patients.^{28,29}

228 During the latest studied period, 5-year OS was worse in young adults than in adolescents and
229 in children³ for ALL (63.2%, 79.3%, and 89.8%, respectively), for astrocytoma (49.6%,
230 74.8%, and 87.0%, respectively), and Ewing tumor (38.1%, 70.4%, and 73.8%, respectively).

231 Interest has recently been paid to AYA with ALL, particularly because all reports so far
232 published indicate that these patients have a better outcome when treated with pediatric rather
233 than adult therapeutic protocols.³⁰ In France since 2003, adolescents with ALL have been
234 treated in or according to pediatric trials (either the FRALLE-2000 or EORTC-58951 trials)³¹
235 and young adults markedly benefited from a pediatric approach by the Group for Research on
236 Adult Acute Lymphoblastic Leukemia (GRAALL).^{32,33} This may explain the difference in
237 survival between adolescents and young adults. Although the inclusion of asparaginase in
238 pediatric-inspired chemotherapy regimens has a positive impact on survival in AYA patients,
239 factors accounting for differences in outcome compared to children include incidence of
240 adverse events³⁴ and heterogeneity in ALL biology.³⁵ Differences in survival between
241 children, adolescents and young adults with astrocytoma result of different distribution of
242 gliomas with varying prognosis depending on tumor grade. Low-grade astrocytomas (WHO
243 grades I and II including especially pilocytic astrocytomas) are the most frequently
244 encountered brain tumors in early childhood and adolescence, while high-grade astrocytomas
245 (WHO grades III and IV, such as anaplastic astrocytomas and glioblastomas) tend to occur at
246 increasing rates with increasing age.³⁶ Several studies have analyzed potential correlations
247 between Ewing sarcoma patients' ages and prognosis, but results have been inconsistent, and
248 it remains unproven whether age in the AYA range by itself truly confers inferior survival.³⁷

249 For this reason, European cooperative group trials have included patients up to the age of 50

250 years.³⁸ Moreover, in the EURO-EWING99 clinical trial, it has been observed that some
251 toxicities (neutropenia, thrombocytopenia and stomatitis) may even decrease with age.³⁸ It
252 remains to be determined whether less toxicity represents less treatment administered to older
253 patients, or whether age distribution of biologic factors and Ewing-like tumor entities, treated
254 as Ewing sarcoma and having different clinical behavior, can be held responsible.³⁹ Even
255 though the survival between adolescents and young adults seemed to be similar for the other
256 cancers, these AYA patients had persistent worse 5-year overall survival than children for
257 osteosarcoma, rhabdomyosarcoma, and malignant hepatic tumors.^{1,3} Several factors are
258 believed to have influenced the lack of survival improvement noted for AYA: access to
259 clinical trials, biologic characteristics of these cancers, access to health-care professionals and
260 services, and specific treatment strategies.^{4,40,41}

261

262 **Conclusion**

263 We have presented updated data representing a comprehensive analysis of cancer incidence
264 and survival in AYA in France during the 2000-2016 period. Notwithstanding the
265 encouraging results for some cancers, and overall, persistent poorer survivals in AYA were
266 shown compared to children for ALL, osteosarcoma, Ewing tumor, rhabdomyosarcoma and
267 malignant hepatic tumors. These survival disparities observed require further investigation to
268 identify and address the causes of these inferior outcomes.

269

270 **Author Disclosure Statements**

271 No competing financial interests exist.

272

273 **Funding Information**

274 The current work is funding by the French Public Health Agency « Santé Publique France »
275 (project number: 18DMNA53-0).

276

277 **Acknowledgments**

278 The authors thank all those who contributed toward the recording of cancer data: the
279 laboratories and departments of anatomy, cytology and pathology, the departments of medical
280 information of the public and private hospitals, and the French health insurance. The authors
281 thank the French Public Health Agency «Santé Publique France» for financial support.

282

283 **References**

- 284 1. Desandes E, Lacour B, Belot A, et al. Cancer Incidence and Survival in Adolescents
285 and Young Adults in France, 2000-2008. *Pediatr Hematol Oncol.* 2013;30(4):291-306.
- 286 2. Desandes E, Lacour B, Belot A, et al. Cancer incidence and survival among
287 adolescents and young adults in France (1978-1997). *Bull Cancer.* 2007;94(4):331-7.
- 288 3. Lacour B, Goujon S, Guissou S, et al. Childhood cancer survival in France, 2000–
289 2008. *Eur J Cancer Prev.* 2014;23(5):449-57.
- 290 4. Desandes E, Brugieres L, Molinie F, et al. Adolescent and young adult oncology
291 patients in France: Heterogeneity in pathways of care. *Pediatr Blood Cancer.*
292 2018;65(9):e27235.
- 293 5. Tycsynski J, Demaret E, Parkin D. Standards and Guidelines for Cancer Registration
294 Network in Europe. The ENCR recommendations. Lyon, IARC Technical Publication n°40,
295 2003, pp 97.
- 296 6. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of
297 Childhood Cancer, third edition. *Cancer.* 2005;103(7):1457-67.

- 298 7. Frizt A, Percy C, Jack A, et al. International Classification of Diseases for Oncology.
299 Third edition. First Revision. Geneva, World Health Organization, 2013.
- 300 8. Bray F. Chapter 8. Age-standardization, in Parkin D, Whelan SL, Ferlay J, et al (eds):
301 Cancer Incidence in Five Continents. Lyon, IARC Scientific Publications n° 156, 2002, pp
302 87-9.
- 303 9. Dobson AJ, Kuulasmaa K, Eberle E, Scherer J. Confidence intervals for weighted
304 sums of Poisson parameters. *Stat Med.* 1991;10(3):457-62.
- 305 10. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-
306 2016, National Cancer Institute. Bethesda, MD, https://seer.cancer.gov/csr/1975_2016/,
307 accessed on November 8, 2019.
- 308 11. Kaplan E, Meier P. Non-parametric estimation from incomplete observation. *J Am*
309 *Stat Assoc.* 1958;53:457-81.
- 310 12. Mantel N. Evaluation of survival date and two new rank order statistics arising in its
311 consideration. *Cancer Chemother Rep.* 1966;50:163-70.
- 312 13. Peto R, Pike MC, Armitage P, et al. Design and analysis of randomized clinical trials
313 requiring prolonged observation of each patient. II. analysis and examples. *Br J Cancer.*
314 1977;35(1):1-39.
- 315 14. Cox DR, Oakes D. Analysis of survival data. New-York, Chapman and Hall, 1984, pp
316 212.
- 317 15. Carreira H, Antunes L, Castro C, et al. Cancer incidence and survival (1997-2006)
318 among adolescents and young adults in the north of Portugal. *Pediatr Hematol Oncol.*
319 2012;29(7):663-76.
- 320 16. Furlong W, Rae C, Greenberg ML, Barr RD. Surveillance and survival among
321 adolescents and young adults with cancer in Ontario, Canada. *Int J Cancer.*
322 2012;131(11):2660-7.

- 323 17. van Gaal JC, Bastiaannet E, Schaapveld M, et al. Cancer in adolescents and young
324 adults in north Netherlands (1989-2003): increased incidence, stable survival and high
325 incidence of second primary tumours. *Ann Oncol.* 2009;20(2):365-73.
- 326 18. Australian Institute of Health and Welfare 2018. Cancer in adolescents and young
327 adults in Australia. Cat. no. CAN 110, Canberra: AIHW, <https://www.aihw.gov.au>, accessed
328 on April 1, 2020.
- 329 19. Stark D, Bowen D, Dunwoodie E, et al. Survival patterns in teenagers and young
330 adults with cancer in the United Kingdom: Comparisons with younger and older age groups.
331 *Eur J Cancer.* 2015;51(17):2643-54.
- 332 20. Gondos A, Hiripi E, Hollecsek B, et al. Survival among adolescents and young adults
333 with cancer in Germany and the United States: an international comparison. *Int J Cancer.*
334 2013;133(9):2207-15.
- 335 21. Trama A, Botta L, Foschi R, et al. Survival of European adolescents and young adults
336 diagnosed with cancer in 2000-07: population-based data from EURO CARE-5. *Lancet Oncol.*
337 2016;17(7):896-906.
- 338 22. Pereira M, Williams VL, Hallanger Johnson J, Valderrabano P. Thyroid Cancer
339 Incidence Trends in the United States: Association with Changes in Professional Guideline
340 Recommendations. *Thyroid.* 2020;doi: 10.1089/thy.2019.
- 341 23. Jones SE, Saraiya M, Miyamoto J, Berkowitz Z. Trends in sunscreen use among U.S.
342 high school students: 1999-2009. *J Adolesc Health.* 2012;50(3):304-7.
- 343 24. Gupta S, Harper A, Ruan Y, et al. International trends in the incidence of cancer
344 among adolescents and young adults. *J Natl Cancer Inst.* 2020;doi: 10.1093/jnci/djaa007
345 [Epub ahead of print].

- 346 25. Keegan TH, Ries LA, Barr RD, et al. Comparison of cancer survival trends in the
347 United States of adolescents and young adults with those in children and older adults. *Cancer*.
348 2016;122(7):1009-16.
- 349 26. Papworth KE, Arroyo VM, Styring E, et al. Soft-tissue sarcoma in adolescents and
350 young adults compared with older adults: A report among 5000 patients from the
351 Scandinavian Sarcoma Group Central Register. *Cancer*. 2019;125(20):3595-602.
- 352 27. van der Graaf WTA, Orbach D, Judson IR, Ferrari A. Soft tissue sarcomas in
353 adolescents and young adults: a comparison with their paediatric and adult counterparts.
354 *Lancet Oncol*. 2017;18(3):e166-75.
- 355 28. Battaglia S, Dumoucel S, Chesneau J, et al. Impact of oncopediatric dosing regimen of
356 zoledronic acid on bone growth: preclinical studies and case report of an osteosarcoma
357 pediatric patient. *J Bone Miner Res*. 2011;26(10):2439-51.
- 358 29. Gaspar N, Océan BV, Pacquement H, et al. Results of methotrexate-etoposide-
359 ifosfamide based regimen (M-EI) in osteosarcoma patients included in the French
360 OS2006/sarcome-09 study. *Eur J Cancer*. 2018;8857-66.
- 361 30. Boissel N, Sender LS. Best Practices in Adolescent and Young Adult Patients with
362 Acute Lymphoblastic Leukemia: A Focus on Asparaginase. *J Adolesc Young Adult Oncol*.
363 2015;4(3):118-28.
- 364 31. Garand R, Beldjord K, Cavé H, et al. Flow cytometry and IG/TCR quantitative PCR
365 for minimal residual disease quantitation in acute lymphoblastic leukemia: a French
366 multicenter prospective study on behalf of the FRALLE, EORTC and GRAALL. *Leukemia*.
367 2013;27(2):370-6.
- 368 32. Huguet F, Leguay T, Raffoux E, et al. Pediatric-inspired therapy in adults with
369 Philadelphia chromosome-negative acute lymphoblastic leukemia: the GRAALL-2003 study.
370 *J Clin Oncol*. 2009;27(6):911-8.

- 371 33. Desjonquères A, Chevallier P, Thomas X, et al. Acute lymphoblastic leukemia
372 relapsing after first-line pediatric-inspired therapy: a retrospective GRAALL study. *Blood*
373 *Cancer J.* 2016;9(12):e504.
- 374 34. Plourde PV, Jeha S, Hijiya N, et al. Safety profile of asparaginase Erwinia
375 chrysanthemi in a large compassionate-use trial. *Pediatr Blood Cancer.* 2014;61(7):1232-8.
- 376 35. Roberts KG, Mullighan CG. Genomics in acute lymphoblastic leukaemia: insights and
377 treatment implications. *Nat Rev Clin Oncol.* 2015;12(6):344-57.
- 378 36. Diwanji TP, Engelman A, Snider JW, Mohindra P. Epidemiology, diagnosis, and
379 optimal management of glioma in adolescents and young adults. *Adolesc Health Med Ther.*
380 2017;899-113.
- 381 37. Wilhelm M, Dirksen U, Bielack SS, et al. ENCCA WP17-WP7 consensus paper on
382 teenagers and young adults (TYA) with bone sarcomas. *Ann Oncol.* 2014;25(8):1500-5.
- 383 38. Juergens C, Weston C, Lewis I, et al. Safety assessment of intensive induction with
384 vincristine, ifosfamide, doxorubicin, and etoposide (VIDE) in the treatment of Ewing tumors
385 in the EURO-E.W.I.N.G. 99 clinical trial. *Pediatr Blood Cancer.* 2006;47(1):22-9.
- 386 39. Gaspar N, Hawkins DS, Dirksen U, et al. Ewing Sarcoma: Current Management and
387 Future Approaches Through Collaboration. *J Clin Oncol.* 2015;33(27):3036-46.
- 388 40. Desandes E, Brugieres L, Laurence V, et al. Survival of adolescents with cancer
389 treated at pediatric versus adult oncology treatment centers in France. *Pediatr Blood Cancer.*
390 2017;64(5):1-8.
- 391 41. Desandes E, Bonnay S, Berger C, et al. Pathways of care for adolescent patients with
392 cancer in France from 2006 to 2007. *Pediatr Blood Cancer.* 2012;58(6):924-9.
- 393
- 394

395 TABLE 1. CANCER CASES IN ADOLESCENTS AND YOUNG ADULTS DIAGNOSED IN 2000–2016, IN
396 19 FRENCH ADMINISTRATIVE AREAS, WITH DATA QUALITY INDICATORS

397 Abbreviations: ICCC-3, third version of the International Classification of Childhood Cancer.

398

399 TABLE 2. NUMBER OF TUMORS, SEX RATIO, AND CANCER IRs PER MILLION IN AYA IN FRANCE
400 FROM 2000 TO 2016 DETAILED BY DIAGNOSTIC GROUPS AND SUBGROUPS

401 Abbreviations: AYA, adolescents and young adults; ICCC-3, third version of the International
402 Classification of Childhood Cancer; GCT, Germ-Cell tumors; cPNET, central primitive
403 neuroectodermal tumors; pPNET, peripheral primitive neuroectodermal tumors; M/F ratio,
404 Male/Female ratio; IR, Incidence rates per million; ASR, age-standardized incidence rates per
405 million; 95%CI, 95% confidence interval.

406

407 **FIG. 1.** Time trend in age-standardized incidence rates estimated by the conventional annual
408 percent change, in AYA (15–24 years) in France between 2000 and 2016

409 Abbreviations: cAPC, conventional annual percent change; ASR, age-standardized incidence
410 rates per million; AYA, adolescents and young adults.

411

412 TABLE 3. ONE-, TWO-, AND FIVE-YEAR OVERALL SURVIVAL AMONG AYA WITH CANCER
413 FROM 2000 TO 2015 BY ICCC-3 DIAGNOSTIC GROUPS

414 Abbreviations: AYA, adolescents and young adults; ICCC-3, third version of the International
415 Classification of Childhood Cancer; GCT, Germ-Cell tumors; cPNET, central primitive
416 neuroectodermal tumors; pPNET, peripheral primitive neuroectodermal tumors; OS, Overall
417 Survival; 95%CI, 95% confidence interval.

418

419 TABLE 4. TRENDS IN FIVE-YEAR OVERALL SURVIVAL AMONG ADOLESCENTS AND YOUNG
420 ADULTS WITH CANCER FROM 1978 TO 2015 BY ICCC-3 DIAGNOSTIC GROUPS
421 Abbreviations: ICCC-3: third version of the International Classification of Childhood Cancer;
422 OS: Overall Survival; 95%CI: 95% confidence interval
423 * Data from reference ²
424 # Only malignant CNS tumors were included (exclusion of benign or borderline tumors)
425 ^a p<0.05 Comparison between diagnostic periods (2000-2007 vs 2008-2015)

426

427 TABLE 5. TRENDS IN FIVE-YEAR OVERALL SURVIVAL AMONG ADOLESCENTS AND YOUNG
428 ADULTS WITH CANCER FROM 2000-2007 TO 2008-2015 BY ICCC-3 DIAGNOSTIC GROUPS
429 Abbreviations: ICCC-3, third version of the International Classification of Childhood Cancer;
430 5yOS, Five-year Overall Survival; 95%CI, 95% confidence interval.
431 ^a p<0.05 Comparison between diagnostic periods (2000-2007 vs 2008-2015)
432 ^b p<0.05 Comparison between age groups (15-19 years vs 20-24 years)

TABLE 1. CANCER CASES IN ADOLESCENTS AND YOUNG ADULTS DIAGNOSED IN 2000–2016, IN 19 FRENCH ADMINISTRATIVE AREAS, WITH DATA

QUALITY INDICATORS

French Administrative Areas	Registered period	Incidence analysis (2000-2016)			Survival analysis (2000-2015)	
		Patient included N	Microscopically Verified N (%)	Unspecified cases according to ICCC-3 N (%)	Patient included N	Lost to follow-up N (%)
14-Calvados	2000-2015	396	391 (98.7)	6 (1.5)	392	5 (1.3)
16-Charente	2008-2016	123	120 (97.6)	1 (0.8)	110	5 (4.5)
17-Charente/Maritime	2008-2016	189	187 (98.9)	2 (1.1)	164	19 (11.6)
25-Doubs	2000-2016	336	326 (97.0)	10 (3.0)	320	0 (0.0)
33-Gironde	2005,2008-2016	594	579 (97.5)	7 (1.2)	532	18 (3.4)
34-Hérault	2000-2016	632	624 (98.7)	12 (1.9)	573	25 (4.4)
38-Isère	2000-2016	816	810 (99.3)	7 (0.9)	775	17 (2.2)
44-Loire Atlantique	2000-2016	836	830 (99.3)	10 (1.2)	780	22 (2.8)
50-Manche	2000-2016	245	242 (98.8)	4 (1.6)	234	14 (6.0)
59-Lille Area	2008-2016	315	313 (99.4)	4 (1.3)	284	4 (1.4)
67- Bas Rhin	2000-2014	644	631 (98.0)	10 (1.6)	642	13 (2.0)
68- Haut Rhin	2000-2016	457	454 (99.3)	1 (0.2)	429	19 (4.4)
79-Deux-Sèvres	2008-2016	124	122 (98.4)	3 (2.4)	108	2 (1.9)
80-Somme	2000-2016	325	320 (98.5)	7 (2.2)	309	9 (2.9)
81-Tarn	2000-2016	173	171 (98.8)	1 (0.6)	157	3 (1.9)
85-Vendée	2000-2016	429	427 (99.5)	3 (0.7)	404	7 (1.7)
86-Vienne	2008-2016	151	146 (96.7)	2 (1.3)	133	3 (2.3)
87-Haute Vienne	2009-2016	99	97 (98.0)	2 (2.0)	89	2 (2.2)
90- Territoire de Belfort	2007-2016	49	48 (98.0)	1 (2.0)	45	1 (2.2)
Total	2000-2016	6933	6838 (98.6)	93 (1.3)	6480	188 (2.9)

Abbreviations: ICCC-3, third version of the International Classification of Childhood Cancer.

TABLE 2. NUMBER OF TUMORS, SEX RATIO, AND CANCER IRS PER MILLION IN AYA IN FRANCE FROM 2000 TO 2016 DETAILED BY DIAGNOSTIC

GROUPS AND SUBGROUPS

Diagnostic groups and subgroups (ICCC-3)	Adolescents (15–19 years)			Young adults (20–24 years)			AYA (15–24 years)		
	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	ASR [95%CI]
I. Leukemias, myeloproliferative/dysplastic diseases	365 (13.4)	1.5	31.0 [27.9-34.3]	318 (7.6)	1.2	26.8 [23.9-29.9]	683 (9.9)	1.4	29.0 [26.0-32.2]
Ia. Lymphoblastic leukemias	196 (7.2)	2.1	16.6 [14.4-19.1]	111 (2.6)	2.2	9.4 [7.7-11.3]	307 (4.4)	2.1	13.2 [11.2-15.4]
Ia1. Precursor cell leukemias	195 (7.1)	2.1	16.5 [14.3-19]	100 (2.4)	2.2	8.4 [6.9-10.3]	295 (4.3)	2.1	12.7 [10.8-14.9]
Ia2. Mature B-cell leukemias	1 (0.0)	0.0	0.1 [0.0-0.5]	8 (0.2)	3.0	0.7 [0.3-1.3]	9 (0.1)	2.0	0.4 [0.1-0.9]
Ia3. Mature T-cell and NK cell leukemias	0 (0.0)	-	0.0 [0.1-0.3]	3 (0.1)	0.5	0.3 [0.0-0.7]	3 (0.0)	0.5	0.1 [0.1-0.5]
Ib. Acute myeloid leukemias	102 (3.7)	1.1	8.7 [7.1-10.5]	87 (2.1)	0.9	7.3 [5.9-9.0]	189 (2.7)	1.0	8.0 [6.5-9.8]
Ic. Chronic myeloproliferative diseases	42 (1.5)	1.0	3.6 [2.6-4.8]	96 (2.3)	1.0	8.1 [6.6-9.9]	138 (2.0)	1.0	5.7 [4.4-7.2]
Id. Myelodysplastic diseases	16 (0.6)	0.8	1.4 [0.8-2.2]	19 (0.5)	0.7	1.6 [1.0-2.5]	35 (0.5)	0.8	1.5 [0.9-2.3]
Ie. Unspecified and other specified leukemias	9 (0.3)	3.5	0.8 [0.3-1.5]	5 (0.1)	0.3	0.4 [0.1-1.0]	14 (0.2)	1.3	0.6 [0.2-1.2]
II. Lymphomas and lymphoreticular neoplasms	758 (27.7)	0.9	64.3 [59.8-69.1]	956 (22.8)	1.0	80.6 [75.6-85.9]	1714 (24.7)	0.9	72.0 [67.2-77.0]
IIa. Hodgkin disease	557 (20.4)	0.8	47.3 [43.4-51.3]	703 (16.7)	0.9	59.3 [55-63.8]	1260 (18.2)	0.8	52.9 [48.8-57.2]
IIb. Non-Hodgkin lymphomas	164 (6.0)	1.0	13.9 [11.9-16.2]	227 (5.4)	1.2	19.1 [16.7-21.8]	391 (5.6)	1.1	16.4 [14.2-18.8]
IIb1. Precursor-cell lymphomas	12 (0.4)	5.0	1.0 [0.5-1.8]	14 (0.3)	2.5	1.2 [0.6-2.0]	26 (0.4)	3.3	1.1 [0.6-1.9]
IIb2. Mature B-cell lymphomas	101 (3.7)	0.9	8.6 [7.0-10.4]	160 (3.8)	1.3	13.5 [11.5-15.8]	261 (3.8)	1.1	10.9 [9.1-12.9]
IIb3. Mature T-cell and NK-cell lymphomas	47 (1.7)	1.0	4.0 [2.9-5.3]	46 (1.1)	0.8	3.9 [2.8-5.2]	93 (1.3)	0.9	3.9 [2.9-5.2]
IIb4. Unspecified Non-Hodgkin lymphomas	4 (0.1)	0.0	0.3 [0.1-0.9]	7 (0.2)	0.8	0.6 [0.2-1.2]	11 (0.2)	0.4	0.5 [0.2-1.0]
IIc. Burkitt lymphoma	23 (0.8)	3.6	2.0 [1.2-2.9]	15 (0.4)	4.0	1.3 [0.7-2.1]	38 (0.5)	3.8	1.6 [1.0-2.5]
IId. Miscellaneous lymphoreticular neoplasms	13 (0.5)	3.3	1.1 [0.6-1.9]	8 (0.2)	1.0	0.7 [0.3-1.3]	21 (0.3)	2.0	0.9 [0.4-1.6]
IIe. Unspecified lymphoma	1 (0.0)	0.0	0.1 [0.0-0.5]	3 (0.1)	0.0	0.3 [0.0-0.7]	4 (0.1)	0.0	0.2 [0.0-0.6]

TABLE 2. (CONTINUED)

Diagnostic groups and subgroups (ICCC-3)	Adolescents (15–19 years)			Young adults (20–24 years)			AYA (15–24 years)		
	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	ASR [95%CI]
III. Central nervous system tumors	332 (12.1)	1.1	28.2 [25.2-31.4]	411 (9.8)	1.3	34.6 [31.4-38.2]	743 (10.7)	1.2	31.2 [28.1-34.6]
IIIa. Ependymomas and choroid plexus tumor	31 (1.1)	1.6	2.6 [1.8-3.7]	25 (0.6)	1.5	2.1 [1.4-3.1]	56 (0.8)	1.5	2.4 [1.6-3.4]
IIIa1. Ependymomas	26 (1.0)	1.9	2.2 [1.4-3.2]	23 (0.5)	1.3	1.9 [1.2-2.9]	49 (0.7)	1.6	2.1 [1.3-3.1]
IIIa2. Choroid plexus tumor	5 (0.2)	0.7	0.4 [0.1-1.0]	2 (0.0)	-	0.2 [0.0-0.6]	7 (0.1)	1.3	0.3 [0.1-0.8]
IIIb. Astrocytomas	106 (3.9)	1.3	9.0 [7.4-10.9]	122 (2.9)	1.7	10.3 [8.5-12.3]	228 (3.3)	1.5	9.6 [7.9-11.5]
IIIc. Intracranial and intraspinal embryonal tumors	26 (1.0)	2.3	2.2 [1.4-3.2]	25 (0.6)	5.3	2.1 [1.4-3.1]	51 (0.7)	3.3	2.2 [1.4-3.2]
IIIc1. Medulloblastomas	18 (0.7)	3.5	1.5 [0.9-2.4]	19 (0.5)	8.5	1.6 [1.0-2.5]	37 (0.5)	5.2	1.6 [0.9-2.5]
IIIc2. ePNET	5 (0.2)	0.7	0.4 [0.1-1.0]	5 (0.1)	4.0	0.4 [0.1-1.0]	10 (0.1)	1.5	0.4 [0.1-1.0]
IIIc3. Medulloepithelioma	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	0.0	0.1 [0.0-0.5]	2 (0.0)	0.0	0.1 [0.0-0.5]
IIIc4. Atypical teratoid/rhabdoid tumor	2 (0.1)	-	0.2 [0.0-0.6]	0 (0.0)	-	0.0 [0.1-0.3]	2 (0.0)	-	0.1 [0.0-0.5]
IIId. Other gliomas	43 (1.6)	0.7	3.6 [2.6-4.9]	101 (2.4)	1.2	8.5 [6.9-10.3]	144 (2.1)	1.1	5.9 [4.7-7.5]
IIId1. Oligodendrogliomas	20 (0.7)	0.8	1.7 [1.0-2.6]	58 (1.4)	1.2	4.9 [3.7-6.3]	78 (1.1)	1.1	3.2 [2.3-4.4]
IIId2. Mixed and unspecified gliomas	22 (0.8)	0.7	1.9 [1.2-2.8]	42 (1)	1.2	3.5 [2.6-4.8]	64 (0.9)	1.0	2.7 [1.8-3.8]
IIId3. Neuroepithelial glial tumors	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	-	0.1 [0.0-0.5]	2 (0.0)	1.0	0.1 [0.0-0.5]
IIIe. Other specified intracranial and intraspinal tumors	115 (4.2)	0.9	9.8 [8.1-11.7]	121 (2.9)	0.8	10.2 [8.5-12.2]	236 (3.4)	0.9	10.0 [8.2-11.9]
IIIe1. Pituitary adenomas and carcinomas	6 (0.2)	2.0	0.5 [0.2-1.1]	4 (0.1)	0.3	0.3 [0.1-0.9]	10 (0.1)	1.0	0.4 [0.1-1.0]
IIIe2. Tumors of sellar region	10 (0.4)	0.7	0.8 [0.4-1.6]	20 (0.5)	1.2	1.7 [1.0-2.6]	30 (0.4)	1.0	1.2 [0.7-2.1]
IIIe3. Pineal parenchymal tumors	2 (0.1)	1.0	0.2 [0.0-0.6]	1 (0.0)	0.0	0.1 [0.0-0.5]	3 (0.0)	0.5	0.1 [0.0-0.6]
IIIe4. Neuronal and mixed neuronal-glial tumors	68 (2.5)	0.9	5.8 [4.5-7.3]	51 (1.2)	1.0	4.3 [3.2-5.7]	119 (1.7)	0.9	5.1 [3.9-6.5]
IIIe5. Meningiomas	29 (1.1)	0.9	2.5 [1.6-3.5]	45 (1.1)	0.7	3.8 [2.8-5.1]	74 (1.1)	0.8	3.1 [2.2-4.3]
IIIff. Unspecified intracranial and intraspinal neoplasms	11 (0.4)	0.8	0.9 [0.5-1.7]	17 (0.4)	1.4	1.4 [0.8-2.3]	28 (0.4)	1.2	1.2 [0.6-2.0]
IV. Neuroblastoma	13 (0.5)	1.6	1.1 [0.6-1.9]	12 (0.3)	0.3	1.0 [0.5-1.8]	25 (0.4)	0.8	1.1 [0.6-1.8]
IVa. Neuroblastoma and ganglioneuroblastoma	6 (0.2)	2.0	0.5 [0.2-1.1]	3 (0.1)	0.5	0.3 [0.0-0.7]	9 (0.1)	1.3	0.4 [0.1-0.9]
IVb. Other peripheral nervous cell tumors	7 (0.3)	1.3	0.6 [0.2-1.2]	9 (0.2)	0.3	0.8 [0.3-1.4]	16 (0.2)	0.6	0.7 [0.3-1.3]
VI. Renal tumors	20 (0.7)	0.7	1.7 [1.0-2.6]	42 (1)	1.2	3.5 [2.6-4.8]	62 (0.9)	1.0	2.6 [1.7-3.6]
VIa. Nephroblastoma, other nonepithelial renal tumors	6 (0.2)	0.0	0.5 [0.2-1.1]	7 (0.2)	1.3	0.6 [0.2-1.2]	13 (0.2)	0.4	0.5 [0.2-1.2]
VIa1. Nephroblastoma	4 (0.1)	0.0	0.3 [0.1-0.9]	6 (0.1)	2.0	0.5 [0.2-1.1]	10 (0.1)	0.7	0.4 [0.1-1.0]
VIa2. Rhabdoid renal tumors	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	0.0	0.1 [0.0-0.5]	2 (0.0)	0.0	0.1 [0.0-0.5]
VIa4. pPNET of kidney	1 (0.0)	0.0	0.1 [0.0-0.5]	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	0.0	0.0 [0.0-0.4]
VIb. Renal carcinoma	14 (0.5)	1.3	1.2 [0.6-2.0]	34 (0.8)	1.3	2.9 [2.0-4.0]	48 (0.7)	1.3	2.0 [1.3-2.9]
VIc. Unspecified malignant renal tumors	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	0.0	0.0 [0.0-0.4]

TABLE 2. (CONTINUED)

Diagnostic groups and subgroups (ICCC-3)	Adolescents (15–19 years)			Young adults (20–24 years)			AYA (15–24 years)		
	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	ASR [95%CI]
VII. Hepatic tumors	11 (0.4)	0.8	0.9 [0.5-1.7]	20 (0.5)	1.9	1.7 [1.0-2.6]	31 (0.4)	1.4	1.3 [0.7-2.1]
VIIa. Hepatoblastoma	1 (0.0)	-	0.1 [0.0-0.5]	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.0 [0.0-0.4]
VIIb. Hepatic carcinoma	9 (0.3)	0.5	0.8 [0.3-1.5]	20 (0.5)	1.9	1.7 [1.0-2.6]	29 (0.4)	1.2	1.2 [0.7-2.0]
VIIc. Unspecified malignant hepatic tumors	1 (0.0)	-	0.1 [0.0-0.5]	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.0 [0.0-0.4]
VIII. Malignant bone tumors	217 (7.9)	1.5	18.4 [16.0-21.0]	122 (2.9)	1.6	10.3 [8.5-12.3]	339 (4.9)	1.5	14.6 [12.5-16.9]
VIIIa. Osteosarcomas	107 (3.9)	1.7	9.1 [7.4-11.0]	48 (1.1)	1.4	4.0 [3.0-5.4]	155 (2.2)	1.6	6.7 [5.3-8.3]
VIIIb. Chondrosarcomas	17 (0.6)	0.9	1.4 [0.8-2.3]	32 (0.8)	0.9	2.7 [1.8-3.8]	49 (0.7)	0.9	2.0 [1.3-3.0]
VIIIc. Ewing tumor and related sarcomas of bone	83 (3.0)	1.5	7.0 [5.6-8.7]	36 (0.9)	3.5	3.0 [2.1-4.2]	119 (1.7)	1.9	5.2 [4.0-6.6]
VIII d. Other specified malignant bone tumors	8 (0.3)	0.3	0.7 [0.3-1.3]	6 (0.1)	2.0	0.5 [0.2-1.1]	14 (0.2)	0.8	0.6 [0.2-1.2]
VIII d2. Malignant chordomas	6 (0.2)	0.0	0.5 [0.2-1.1]	2 (0.0)	-	0.2 [0.0-0.6]	8 (0.1)	0.3	0.3 [0.1-0.9]
VIII d3. Odontogenic malignant tumors	1 (0.0)	-	0.1 [0.0-0.5]	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.0 [0.0-0.4]
VIII d4. Miscellaneous malignant bone tumors	1 (0.0)	-	0.1 [0.0-0.5]	4 (0.1)	1.0	0.3 [0.1-0.9]	5 (0.1)	1.5	0.2 [0.0-0.7]
VIII e. Unspecified malignant bone tumors	2 (0.1)	-	0.2 [0.0-0.6]	0 (0.0)	-	0.0 [0.1-0.3]	2 (0.0)	-	0.1 [0.0-0.5]
IX. Soft tissue and other extrasosseous sarcomas	134 (4.9)	1.4	11.4 [9.5-13.5]	165 (3.9)	1.4	13.9 [11.9-16.2]	299 (4.3)	1.4	12.6 [10.6-14.8]
IXa. Rhabdomyosarcomas	33 (1.2)	1.5	2.8 [1.9-3.9]	23 (0.5)	2.8	1.9 [1.2-2.9]	56 (0.8)	1.9	2.4 [1.6-3.5]
IXb. Fibrosarcomas, and other fibrous neoplasms	26 (1.0)	0.9	2.2 [1.4-3.2]	26 (0.6)	0.9	2.2 [1.4-3.2]	52 (0.8)	0.9	2.2 [1.4-3.2]
IXb1. Fibroblastic and myofibroblastic tumors	17 (0.6)	1.4	1.4 [0.8-2.3]	13 (0.3)	0.6	1.1 [0.6-1.9]	30 (0.4)	1.0	1.3 [0.7-2.1]
IXb2. Nerve sheath tumors	9 (0.3)	0.3	0.8 [0.3-1.5]	13 (0.3)	1.2	1.1 [0.6-1.9]	22 (0.3)	0.7	0.9 [0.5-1.7]
IXc. Kaposi sarcoma	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	0.0	0.0 [0.0-0.4]
IXd. Other specified soft tissue sarcomas	62 (2.3)	1.6	5.3 [4.0-6.7]	87 (2.1)	1.4	7.3 [5.9-9.0]	149 (2.1)	1.4	6.2 [4.9-7.8]
IXd1-2. Ewing tumors and pPNET of soft tissue	16 (0.6)	1.7	1.4 [0.8-2.2]	11 (0.3)	1.8	0.9 [0.5-1.7]	27 (0.4)	1.7	1.2 [0.6-2.0]
IXd3. Extrarenal rhabdoid tumor	0 (0.0)	-	0.0 [0.1-0.3]	3 (0.1)	0.5	0.3 [0.0-0.7]	3 (0.0)	0.5	0.1 [0.1-0.5]
IXd4. Liposarcomas	8 (0.3)	0.6	0.7 [0.3-1.3]	14 (0.3)	1.3	1.2 [0.6-2.0]	22 (0.3)	1.0	0.9 [0.5-1.6]
IXd5. Fibrohistiocytic tumors	2 (0.1)	1.0	0.2 [0.0-0.6]	5 (0.1)	0.7	0.4 [0.1-1.0]	7 (0.1)	0.8	0.3 [0.1-0.8]
IXd6. Meitomyosarcomas	6 (0.2)	-	0.5 [0.2-1.1]	10 (0.2)	1.0	0.8 [0.4-1.6]	16 (0.2)	2.2	0.7 [0.3-1.3]
IXd7. Synovial sarcomas	19 (0.7)	1.4	1.6 [1.0-2.5]	23 (0.5)	1.9	1.9 [1.2-2.9]	42 (0.6)	1.6	1.8 [1.1-2.7]
IXd8. Blood vessel tumors	4 (0.1)	0.3	0.3 [0.1-0.9]	5 (0.1)	0.3	0.4 [0.1-1.0]	9 (0.1)	0.3	0.4 [0.1-0.9]
IXd9. Osseous and chondromatous neoplasms	1 (0.0)	-	0.1 [0.0-0.5]	5 (0.1)	0.7	0.4 [0.1-1.0]	6 (0.1)	1.0	0.2 [0.1-0.7]
IXd10. Alveolar soft parts sarcoma	1 (0.0)	0.0	0.1 [0.0-0.5]	4 (0.1)	-	0.3 [0.1-0.9]	5 (0.1)	4.0	0.2 [0.0-0.7]
IXd11. Miscellaneous soft tissue sarcomas	5 (0.2)	-	0.4 [0.1-1.0]	7 (0.2)	2.5	0.6 [0.2-1.2]	12 (0.2)	5.0	0.5 [0.2-1.1]
IXe. Unspecified soft tissue sarcomas	13 (0.5)	2.3	1.1 [0.6-1.9]	28 (0.7)	1.5	2.4 [1.6-3.4]	41 (0.6)	1.7	1.7 [1.0-2.6]

TABLE 2. (CONTINUED)

Diagnostic groups and subgroups (ICCC-3)	Adolescents (15–19 years)			Young adults (20–24 years)			AYA (15–24 years)		
	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	ASR [95%CI]
X. Germ-cell tumors, and other neoplasms of gonads	292 (10.7)	2.7	24.8 [22.0-27.8]	721 (17.2)	7.0	60.8 [56.4-65.4]	1013 (14.6)	5.0	41.7 [38.2-45.5]
Xa. Intracranial and intraspinal germ cell tumors	31 (1.1)	6.8	2.6 [1.8-3.7]	16 (0.4)	7.0	1.3 [0.8-2.2]	47 (0.7)	6.8	2.0 [1.3-3.0]
Xa1. germinomas	20 (0.7)	19.0	1.7 [1-2.6,0]	9 (0.2)	-	0.8 [0.3-1.4]	29 (0.4)	28.0	1.3 [0.7-2.1]
Xa2. teratomas	3 (0.1)	0.5	0.3 [0.0-0.8]	5 (0.1)	1.5	0.4 [0.1-1.0]	8 (0.1)	1.0	0.3 [0.1-0.9]
Xa3. embryonal carcinomas	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.1 [0.0-0.5]	1 (0.0)	-	0.0 [0.0-0.4]
Xa4. yolk sac tumor	1 (0.0)	-	0.1 [0.0-0.5]	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.0 [0.0-0.6]
Xa5. choriocarcinomas	3 (0.1)	2.0	0.3 [0.0-0.8]	1 (0.0)	-	0.1 [0.0-0.5]	4 (0.1)	3.0	0.2 [0.0-0.6]
Xa6. tumors of mixed forms	4 (0.1)	-	0.3 [0.1-0.9]	0 (0.0)	-	0.0 [0.1-0.3]	4 (0.1)	-	0.2 [0.1-0.6]
Xb. Malignant extracranial and extragonadal GCT	14 (0.5)	2.5	1.2 [0.6-2.0]	35 (0.8)	2.5	3.0 [2.1-4.1]	49 (0.7)	2.5	2.0 [1.3-3.0]
Xb1. germinomas	8 (0.3)	1.7	0.7 [0.3-1.3]	10 (0.2)	9.0	0.8 [0.4-1.6]	18 (0.3)	3.5	0.8 [0.3-1.4]
Xb2. teratomas	3 (0.1)	-	0.3 [0.0-0.8]	3 (0.1)	2.0	0.3 [0.0-0.7]	6 (0.1)	5.0	0.3 [0.0-0.8]
Xb3. Embryonal carcinomas	0 (0.0)	-	0.0 [0.1-0.3]	1 (0.0)	-	0.1 [0.0-0.5]	1 (0.0)	-	0.0 [0.0-0.4]
Xb4. Yolk sac tumor	0 (0.0)	-	0.0 [0.1-0.3]	7 (0.2)	-	0.6 [0.2-1.2]	7 (0.1)	-	0.3 [0.2-0.8]
Xb5. Choriocarcinomas	1 (0.0)	0.0	0.1 [0.0-0.5]	10 (0.2)	0.4	0.8 [0.4-1.6]	11 (0.2)	0.4	0.4 [0.2-1.0]
Xb6. Other and unspecified malignant mixed GCT	2 (0.1)	-	0.2 [0.0-0.6]	4 (0.1)	3.0	0.3 [0.1-0.9]	6 (0.1)	5.0	0.2 [0.0-0.7]
Xc. Malignant gonadal germ cell tumors	224 (8.2)	3.5	19.0 [16.6-21.7]	626 (14.9)	15.1	52.8 [48.7-57.1]	850 (12.3)	8.6	34.9 [31.7-38.3]
Xc1. germinomas	48 (1.8)	1.2	4.1 [3.0-5.4]	189 (4.5)	9.5	15.9 [13.7-18.4]	237 (3.4)	4.9	9.7 [8.1-11.5]
Xc2. teratomas	49 (1.8)	2.3	4.2 [3.1-5.5]	106 (2.5)	8.6	8.9 [7.3-10.8]	155 (2.2)	5.0	6.4 [5.1-8.0]
Xc3. embryonal carcinomas	37 (1.4)	36.0	3.1 [2.2-4.3]	133 (3.2)	132.0	11.2 [9.4-13.3]	170 (2.5)	84.0	6.9 [5.6-8.5]
Xc4. yolk sac tumor	12 (0.4)	1.4	1.0 [0.5-1.8]	20 (0.5)	2.3	1.7 [1.0-2.6]	32 (0.5)	1.9	1.3 [0.8-2.2]
Xc5. choriocarcinomas	7 (0.3)	2.5	0.6 [0.2-1.2]	9 (0.2)	-	0.8 [0.3-1.4]	16 (0.2)	7.0	0.7 [0.3-1.3]
Xc6. tumors of mixed forms	71 (2.6)	13.2	6.0 [4.7-7.6]	169 (4.0)	55.3	14.2 [12.2-16.6]	240 (3.5)	29.0	9.9 [8.2-11.8]
Xd. Gonadal carcinoma	17 (0.6)	0.0	1.4 [0.8-2.3]	38 (0.9)	0.0	3.2 [2.3-4.4]	55 (0.8)	0.0	2.3 [1.5-3.3]
Xe. Other and unspecified malignant gonadal tumors	6 (0.2)	0.5	0.5 [0.2-1.1]	6 (0.1)	2.0	0.5 [0.2-1.1]	12 (0.2)	1.0	0.5 [0.2-1.1]

TABLE 2. (CONTINUED)

Diagnostic groups and subgroups (ICCC-3)	Adolescents (15–19 years)			Young adults (20–24 years)			AYA (15–24 years)		
	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	IR [95%CI]	n (%)	M/F ratio	ASR [95%CI]
XI. Epithelial neoplasms and malignant melanomas	580 (21.2)	0.5	49.2 [45.3-53.4]	1419 (33.8)	0.4	119.6 [113.5-126]	1999 (28.8)	0.4	82.3 [77.4-87.6]
XIa. Adenocarcinomas	4 (0.1)	1.0	0.3 [0.1-0.9]	10 (0.2)	0.4	0.8 [0.4-1.6]	14 (0.2)	0.6	0.6 [0.2-1.2]
XIb. Thyroid carcinomas	241 (8.8)	0.3	20.4 [17.9-23.2]	566 (13.5)	0.2	47.7 [43.9-51.8]	807 (11.6)	0.2	33.3 [30.1-36.7]
XIc. Nasopharyngeal carcinomas	21 (0.8)	3.2	1.8 [1.1-2.7]	14 (0.3)	1.0	1.2 [0.6-2.0]	35 (0.5)	1.9	1.5 [0.9-2.4]
XId. Malignant melanomas	156 (5.7)	0.7	13.2 [11.2-15.5]	430 (10.2)	0.5	36.3 [32.9-39.8]	586 (8.5)	0.6	24.1 [21.4-26.9]
XIf. Other and unspecified carcinomas	158 (5.8)	0.6	13.4 [11.4-15.7]	399 (9.5)	0.4	33.6 [30.4-37.1]	557 (8)	0.4	22.9 [20.3-25.8]
XIf1. Carcinomas of salivary glands	21 (0.8)	0.4	1.8 [1.1-2.7]	22 (0.5)	0.7	1.9 [1.2-2.8]	43 (0.6)	0.5	1.8 [1.1-2.8]
XIf2. Carcinomas of colon and rectum	16 (0.6)	0.3	1.4 [0.8-2.2]	49 (1.2)	0.5	4.1 [3.1-5.5]	65 (0.9)	0.5	2.7 [1.8-3.7]
XIf3. Carcinomas of appendix	67 (2.5)	0.7	5.7 [4.4-7.2]	55 (1.3)	0.4	4.6 [3.5-6.0]	122 (1.8)	0.6	5.2 [4.0-6.7]
XIf4. Carcinomas of lung	13 (0.5)	1.2	1.1 [0.6-1.9]	41 (1.0)	1.0	3.5 [2.5-4.7]	54 (0.8)	1.0	2.2 [1.5-3.2]
XIf5. Carcinomas of thymus	5 (0.2)	0.7	0.4 [0.1-1.0]	6 (0.1)	1.0	0.5 [0.2-1.1]	11 (0.2)	0.8	0.5 [0.2-1.0]
XIf6. Carcinomas of breast	4 (0.1)	0.0	0.3 [0.1-0.9]	90 (2.1)	0.0	7.6 [6.1-9.3]	94 (1.4)	0.0	3.8 [2.9-4.9]
XIf7. Carcinomas of cervix uteri	1 (0.0)	0.0	0.1 [0.0-0.5]	40 (1)	0.0	3.4 [2.4-4.6]	41 (0.6)	0.0	1.6 [1.1-2.4]
XIf8. Carcinomas of bladder	1 (0.0)	-	0.1 [0.0-0.5]	1 (0.0)	-	0.1 [0.0-0.5]	2 (0.0)	-	0.1 [0.0-0.5]
XIf9. Carcinomas of eye	0 (0.0)	-	0.0 [0.1-0.3]	2 (0.0)	1.0	0.2 [0.0-0.6]	2 (0.0)	1.0	0.1 [0.1-0.5]
XIf10. Carcinomas of other specified sites	29 (1.1)	0.7	2.5 [1.6-3.5]	82 (2.0)	0.6	6.9 [5.5-8.6]	111 (1.6)	0.6	4.6 [3.5-5.9]
XIf11. Carcinomas of unspecified sites	1 (0)	-	0.1 [0.0-0.5]	11 (0.3)	2.7	0.9 [0.5-1.7]	12 (0.2)	3.0	0.5 [0.2-1.0]
XII. Other and unspecified malignant neoplasms	12 (0.4)	3.0	1.0 [0.5-1.8]	13 (0.3)	1.2	1.1 [0.6-1.9]	25 (0.4)	1.8	1.1 [0.6-1.8]
XIIa. Other specified malignant tumors	9 (0.3)	3.5	0.8 [0.3-1.5]	6 (0.1)	1.0	0.5 [0.2-1.1]	15 (0.2)	2.0	0.6 [0.3-1.3]
XIIa1. Gastrointestinal stromal tumor	5 (0.2)	4.0	0.4 [0.1-1.0]	2 (0.0)	1.0	0.2 [0.0-0.6]	7 (0.1)	2.5	0.3 [0.1-0.8]
XIIa4. Other complex mixed and stromal neoplasms	1 (0.0)	0.0	0.1 [0.0-0.5]	1 (0.0)	0.0	0.1 [0.0-0.5]	2 (0.0)	0.0	0.1 [0.0-0.5]
XIIa5. Mesothelioma	3 (0.1)	-	0.3 [0.0-0.8]	3 (0.1)	2.0	0.3 [0.0-0.7]	6 (0.1)	5.0	0.3 [0.0-0.8]
XIIb. Other unspecified malignant tumors	3 (0.1)	2.0	0.3 [0.0-0.8]	7 (0.2)	1.3	0.6 [0.2-1.2]	10 (0.1)	1.5	0.4 [0.1-1.0]
Total	2734 (100.0)	1.0	231.9 [223.3-240.8]	4199 (100.0)	1.0	354.0 [343.4-364.9]	6933 (100.0)	1.0	289.4 [279.8-299.2]

Abbreviations: AYA, adolescents and young adults; ICC-3, third version of the International Classification of Childhood Cancer; GCT, Germ-Cell tumors; cPNET, central primitive neuroectodermal tumors; pPNET, peripheral primitive neuroectodermal tumors; M/F ratio, Male/Female ratio; IR, Incidence rates per million; ASR, age-standardized incidence rates per million; 95%CI, 95% confidence interval.

TABLE 3. ONE-, TWO-, AND FIVE-YEAR OVERALL SURVIVAL AMONG AYA WITH CANCER FROM 2000 TO 2015 BY ICCC-3 DIAGNOSTIC GROUPS

ICCC-3 diagnostic groups and subgroups	n	1-year OS % [95%CI]	2-year OS % [95%CI]	5-year OS % [95%CI]
I. Leukemias, myeloproliferative/dysplastic diseases	645	88.4 [85.6-90.6]	80.4 [77.1-83.3]	73.4 [69.7-76.6]
Ia. Lymphoblastic leukemias	292	87.3 [82.9-90.6]	78.7 [73.5-83.0]	69.6 [63.9-74.6]
Ia1. Precursor cell leukemias	280	87.8 [83.4-91.2]	79.2 [74.0-83.5]	69.8 [63.9-74.9]
Ia2. Mature B-cell leukemias	9	77.8 [36.5-93.9]	66.7 [28.2-87.8]	66.7 [28.2-87.8]
Ia3. Mature T-cell and NK cell leukemias	3	66.7 [5.4-94.5]	66.7 [5.4-94.5]	66.7 [5.4-94.5]
Ib. Acute myeloid leukemias	176	83 [76.5-87.8]	72.2 [64.9-78.2]	64.4 [56.8-71.0]
Ic. Chronic myeloproliferative diseases	130	100.0 [-]	99.2 [94.7-99.9]	96.8 [91.7-98.8]
Id. Myelodysplastic diseases	34	91.2 [75.1-97.1]	82.1 [64.3-91.5]	75.6 [57.0-87.0]
Ie. Unspecified and other specified leukemias	14	64.3 [34.3-83.3]	42.9 [17.7-66.0]	42.9 [17.7-66.0]
II. Lymphomas and reticuloendothelial neoplasms	1594	98.1 [97.3-98.7]	96.0 [94.9-96.8]	94.1 [92.8-95.1]
IIa. Hodgkin disease	1170	99.7 [99.1-99.9]	98.0 [97.0-98.7]	96.5 [95.3-97.4]
IIb. Non-Hodgkin lymphomas	365	94.5 [91.5-96.4]	91.1 [87.7-93.6]	87.6 [83.7-90.6]
IIb1. Precursor-cell lymphomas	25	84 [62.8-93.7]	76.0 [54.2-88.4]	68.0 [46.1-82.5]
IIb2. Mature B-cell lymphomas	239	95.3 [91.7-97.4]	92.4 [88.1-95.1]	88.7 [83.8-92.2]
IIb3. Mature T-cell and NK-cell lymphomas	91	95.5 [88.5-98.3]	93.3 [85.7-96.9]	91.0 [82.8-95.4]
IIb4. Unspecified Non-Hodgkin lymphomas	10	90.0 [47.3-98.5]	80.0 [40.9-94.6]	80.0 [40.9-94.6]
IIc. Burkitt lymphoma	38	94.7 [80.6-98.7]	89.5 [74.3-95.9]	89.5 [74.3-95.9]
IId. Miscellaneous lymphoreticular neoplasms	17	88.2 [60.6-96.9]	82.4 [54.7-93.9]	82.4 [54.7-93.9]
IIe. Unspecified lymphoma	4	50.0 [5.8-84.5]	50.0 [5.8-84.5]	50.0 [5.8-84.5]
III. Central nervous system tumors	695	92.7 [90.5-94.4]	86.4 [83.6-88.7]	78.1 [74.7-81]
IIIa. Ependymomas and choroid plexus tumor	52	98 [86.9-99.7]	96.1 [85.2-99.0]	89.2 [76.0-95.4]
IIIa1. Ependymomas	45	100.0 [-]	97.7 [84.9-99.7]	90.0 [75.4-96.1]
IIIa2. Choroid plexus tumor	7	85.7 [33.4-97.9]	85.7 [33.4-97.9]	85.7 [33.4-97.9]
IIIb. Astrocytomas	213	90.9 [86.2-94.1]	78.0 [71.7-83.0]	66.7 [59.8-72.8]
IIIc. Intracranial and intraspinal embryonal tumors	47	89.4 [76.3-95.4]	85.1 [71.3-92.6]	71.1 [55.3-82.1]
IIIc1. Medulloblastomas	34	97.1 [80.9-99.6]	91.2 [75.1-97.1]	81.7 [63.6-91.4]
IIIc2. cPNET	9	66.7 [28.2-87.8]	66.7 [28.2-87.8]	20.0 [1.2-55.7]
IIIc3. Medulloepithelioma	2	100.0 [-]	100.0 [-]	100.0 [-]
IIIc4. Atypical teratoid/rhabdoid tumor	2	50.0 [0.6-91.0]	50.0 [0.6-91.0]	50.0 [0.6-91.0]
IIId. Other gliomas	141	85.6 [78.6-90.5]	77.7 [69.8-83.8]	63.7 [55.0-71.1]
IIId1. Oligodendrogliomas	78	85.5 [75.4-91.7]	78.9 [68.0-86.5]	68.4 [56.7-77.6]
IIId2. Mixed and unspecified gliomas	61	85.2 [73.6-92.0]	77.0 [64.3-85.7]	59.7 [46.1-71.0]
IIId3. Neuroepithelial glial tumors	2	100.0 [-]	50.0 [0.6-91.0]	50.0 [0.6-91.0]
IIIe. Other intracranial and intraspinal tumors	217	98.1 [95.1-99.3]	97.2 [93.9-98.7]	96.1 [92.4-98]
IIIe1. Pituitary adenomas and carcinomas	10	100.0 [-]	100.0 [-]	100 [100-100]
IIIe2. Tumors of sellar region	29	100.0 [-]	100.0 [-]	95.5 [71.9-99.3]
IIIe3. Pineal parenchymal tumors	3	100.0 [-]	100.0 [-]	100.0 [-]
IIIe4. Neuronal and mixed neuronal-glial tumors	112	98.2 [93.0-99.5]	97.3 [91.8-99.1]	96.3 [90.4-98.6]
IIIe5. Meningiomas	63	96.8 [87.8-99.2]	95.2 [85.8-98.4]	95.2 [85.8-98.4]
IIIe6. Unspecified neoplasms	27	92.3 [72.6-98.0]	92.3 [72.6-98.0]	88.5 [68.4-96.1]
IV. Neuroblastoma	24	87.5 [66.1-95.8]	83.3 [61.5-93.4]	79.2 [57.0-90.8]
IVa. Neuroblastoma and ganglioneuroblastoma	9	66.7 [28.2-87.8]	66.7 [28.2-87.8]	55.6 [20.4-80.5]
IVb. Other peripheral nervous cell tumors	15	100 -	93.3 [61.3-99.0]	93.3 [61.3-99.0]

Table 3. (CONTINUED)

ICCC-3 diagnostic groups and subgroups	n	1-year OS % [95%CI]	2-year OS % [95%CI]	5-year OS % [95%CI]
VI. Renal tumors	55	90.7 [79.2-96.0]	88.9 [76.9-94.8]	79.1 [65.4-87.9]
VIa. Nephroblastoma and other nonepithelial tumors	11	81.8 [44.7-95.1]	81.8 [44.7-95.1]	71.6 [35.0-89.9]
VIa1. Nephroblastoma	8	100.0 [-]	100.0 [-]	85.7 [33.4-97.9]
VIa2. Rhabdoid renal tumors	2	0.0 [-]	0.0 [-]	0.0 [-]
VIa4. pPNET of kidney	1	100.0 [-]	100.0 [-]	100.0 [-]
VIb. Renal carcinoma	43	92.9 [79.5-97.6]	90.5 [76.6-96.3]	80.5 [64.6-89.7]
VIc. Unspecified malignant renal tumors	1	100.0 [-]	100.0 [-]	100.0 [-]
VII. Hepatic tumors	31	71.0 [51.6-83.7]	58.1 [39.0-73.1]	33.4 [17.3-50.4]
VIIa. Hepatoblastoma	1	100.0 [-]	100.0 [-]	100.0 [-]
VIIb. Hepatic carcinoma	29	69.0 [48.8-82.5]	55.2 [35.6-71.0]	28.4 [13.0-46.0]
VIIc. Unspecified malignant hepatic tumors	1	100.0 [-]	100.0 [-]	100.0 [-]
VIII. Malignant bone tumors	313	94.2 [90.9-96.3]	82.2 [77.5-86.1]	65.1 [59.4-70.3]
VIIIa. Osteosarcomas	145	93.8 [88.3-96.7]	82.6 [75.3-87.9]	60.3 [51.5-67.9]
VIIIb. Chondrosarcomas	42	100.0 [-]	92.7 [79.0-97.6]	90.0 [75.6-96.1]
VIIIc. Ewing tumor and related sarcomas of bone	113	92.9 [86.3-96.4]	77.0 [68.1-83.7]	61.1 [51.3-69.5]
VIId. Other specified malignant bone tumors	12	90.9 [50.8-98.7]	90.9 [50.8-98.7]	72.7 [37.1-90.3]
VIId2. Malignant chordomas	6	80.0 [20.4-96.9]	80.0 [20.4-96.9]	40.0 [5.2-75.3]
VIId3. Odontogenic malignant tumors	1	100.0 [-]	100.0 [-]	100.0 [-]
VIId4. Miscellaneous malignant bone tumors	5	100.0 [-]	100.0 [-]	100.0 [-]
VIIIe. Unspecified malignant bone tumors	1	100.0 [-]	100.0 [-]	100.0 [-]
IX. Soft tissue and other extraosseous sarcomas	288	90.5 [86.5-93.4]	79.9 [74.8-84.2]	66.8 [60.9-72.1]
IXa. Rhabdomyosarcomas	52	84.6 [71.6-92.0]	61.5 [47.0-73.2]	48.1 [34.1-60.8]
IXb. Fibrosarcomas, and other fibrous neoplasms	50	89.8 [77.2-95.6]	85.7 [72.4-92.9]	79.1 [64.6-88.2]
IXb1. Fibroblastic and myofibroblastic tumors	29	100.0 [-]	96.4 [77.2-99.5]	89.0 [69.7-96.3]
IXb2. Nerve sheath tumors	21	76.2 [51.9-89.3]	71.4 [47.2-86.0]	66.7 [42.5-82.5]
IXc. Kaposi sarcoma	1	100.0 [-]	100.0 [-]	100.0 [-]
IXd. Other specified soft tissue sarcomas	144	93.6 [88.1-96.6]	84.4 [77.3-89.5]	70.6 [62.1-77.6]
IXd1-2. Ewing tumors and pPNET of soft tissue	27	92.6 [73.5-98.1]	74.1 [53.2-86.7]	58.3 [37.4-74.4]
IXd3. Extrarenal rhabdoid tumor	3	33.3 [0.9-77.4]	0.0 [-]	0.0 [-]
IXd4. Liposarcomas	22	100.0 [-]	100.0 [-]	95.2 [70.7-99.3]
IXd5. Fibrohistiocytic tumors	7	100.0 [-]	100.0 [-]	100.0 [-]
IXd6. Meioyosarcomas	16	100.0 [-]	100.0 [-]	87.5 [58.6-96.7]
IXd7. Synovial sarcomas	39	97.4 [82.8-99.6]	89.3 [73.8-95.8]	65.7 [47.3-78.9]
IXd8. Blood vessel tumors	8	75 [31.5-93.1]	62.5 [22.9-86.1]	50.0 [15.2-77.5]
IXd9. Osseous and chondromatous neoplasms	6	100.0 [-]	100.0 [-]	100.0 [-]
IXd10. Alveolar soft parts sarcoma	4	100.0 [-]	100.0 [-]	100.0 [-]
IXd11. Miscellaneous soft tissue sarcomas	12	81.8 [44.7-95.1]	54.5 [22.9-78.0]	36.4 [11.2-62.7]
IXe. Unspecified soft tissue sarcomas	41	87.8 [73.2-94.7]	80.5 [64.8-89.7]	62.6 [45.7-75.6]

Table 3. (CONTINUED)

ICCC-3 diagnostic groups and subgroups	n	1-year OS % [95%CI]	2-year OS % [95%CI]	5-year OS % [95%CI]
X. Germ-cell tumors, and other neoplasms of gonads	948	97.0 [95.7-97.9]	94.5 [92.9-95.8]	92.9 [91-94.4]
Xa. Intracranial and intraspinal germ cell tumors	42	97.6 [84.3-99.7]	88.1 [73.7-94.9]	83.3 [68.2-91.7]
Xa1. germinomas	26	100.0 [-]	92.3 [72.6-98.0]	88.5 [68.4-96.1]
Xa2. teratomas	7	100.0 [-]	100.0 [-]	100.0 [-]
Xa4. yolk sac tumor	1	100.0 [-]	0.0 [-]	0.0 [-]
Xa5. choriocarcinomas	4	75.0 [12.8-96.1]	75.0 [12.8-96.1]	75.0 [12.8-96.1]
Xa6. tumors of mixed forms	4	100.0 [-]	75.0 [12.8-96.1]	50.0 [5.8-84.5]
Xb. Malignant extracranial and extragonadal GCT	47	85.1 [71.3-92.6]	72.3 [57.2-82.9]	69.4 [53.8-80.7]
Xb1. germinomas	17	88.2 [60.6-96.9]	88.2 [60.6-96.9]	88.2 [60.6-96.9]
Xb2. teratomas	6	66.7 [19.5-90.4]	33.3 [4.6-67.6]	33.3 [4.6-67.6]
Xb3. Embryonal carcinomas	1	100.0 [-]	100.0 [-]	100.0 [-]
Xb4. Yolk sac tumor	6	100.0 [-]	50.0 [11.1-80.4]	50.0 [11.1-80.4]
Xb5. Choriocarcinomas	11	81.8 [44.7-95.1]	72.7 [37.1-90.3]	72.7 [37.1-90.3]
Xb6. Other and unspecified malignant mixed GCT	6	83.3 [27.3-97.5]	83.3 [27.3-97.5]	62.5 [14.2-89.3]
Xc. Malignant gonadal germ cell tumors	798	98.2 [97.0-98.9]	96.9 [95.5-97.9]	95.7 [94.0-96.9]
Xc1. germinomas	220	99.5 [96.7-99.9]	99.1 [96.3-99.8]	99.1 [96.3-99.8]
Xc2. teratomas	148	99.3 [95.2-99.9]	96.6 [91.9-98.6]	93.5 [87.9-96.6]
Xc3. embryonal carcinomas	159	100.0 [-]	99.4 [95.5-99.9]	98.0 [94.1-99.4]
Xc4. yolk sac tumor	29	96.4 [77.2-99.5]	96.4 [77.2-99.5]	96.4 [77.2-99.5]
Xc5. choriocarcinomas	16	87.1 [57.3-96.6]	73.7 [44.1-89.2]	73.7 [44.1-89.2]
Xc6. tumors of mixed forms	226	96.0 [92.4-97.9]	95.1 [91.3-97.2]	93.7 [89.6-96.2]
Xd. Gonadal carcinoma	50	88.0 [75.2-94.4]	82.0 [68.3-90.2]	77.9 [63.6-87.1]
Xe. Other and unspecified malignant gonadal tumors	11	100.0 [-]	100.0 [-]	100.0 [-]
XI. Epithelial neoplasms and malignant melanomas	1883	97.1 [96.2-97.8]	95.3 [94.2-96.1]	92.5 [91.2-93.7]
XIa. Adrenocortical carcinomas	14	78.6 [47.2-92.5]	57.1 [28.4-78.0]	57.1 [28.4-78.0]
XIb. Thyroid carcinomas	746	99.7 [98.9-99.9]	99.7 [98.9-99.9]	99.4 [98.5-99.8]
XIc. Nasopharyngeal carcinomas	33	90.9 [74.4-97.0]	84.8 [67.4-93.4]	77.8 [58.7-88.8]
XId. Malignant melanomas	560	98.9 [97.6-99.5]	97.8 [96.2-98.7]	95.3 [93.1-96.8]
XIf. Other and unspecified carcinomas	531	92.3 [89.7-94.3]	87.9 [84.8-90.4]	81.7 [78.1-84.8]
XIf1. Carcinomas of salivary glands	40	100.0 [-]	100.0 [-]	100.0 [-]
XIf2. Carcinomas of colon and rectum	64	92.2 [82.2-96.7]	85.9 [74.7-92.4]	67.9 [54.7-78.0]
XIf3. Carcinomas of appendix	109	100.0 [-]	100.0 [-]	100.0 [-]
XIf4. Carcinomas of lung	54	90.6 [78.8-96.0]	88.6 [76.4-94.7]	82.7 [69.4-90.6]
XIf5. Carcinomas of thymus	11	100.0 [-]	90.9 [50.8-98.7]	81.8 [44.7-95.1]
XIf6. Carcinomas of breast	91	94.5 [87.3-97.7]	91.2 [83.2-95.5]	82.7 [73.0-89.2]
XIf7. Carcinomas of cervix uteri	40	97.5 [83.5-99.6]	87.5 [72.5-94.6]	87.5 [72.5-94.6]
XIf8. Carcinomas of bladder	2	50.0 [0.6-91.0]	50.0 [0.6-91.0]	50.0 [0.6-91.0]
XIf9. Carcinomas of eye	2	100.0 [-]	100.0 [-]	100.0 [-]
XIf10. Carcinomas of other specified sites	107	84.1 [75.7-89.8]	75.7 [66.4-82.8]	70.7 [61.0-78.4]
XIf11. Carcinomas of unspecified sites	11	40.9 [12.7-67.9]	30.7 [7.3-58.6]	0.0 [-]
XII. Other and unspecified malignant neoplasms	24	74.6 [51.9-87.7]	70.2 [47.4-84.5]	65.5 [42.6-81.1]
XIIa. Other specified malignant tumors	14	92.9 [59.1-99.0]	92.9 [59.1-99.0]	85.1 [52.3-96.1]
XIIa1. Gastrointestinal stromal tumor	6	100.0 [-]	100.0 [-]	100.0 [-]
XIIa4. Other complex mixed and stromal neoplasms	2	50.0 [0.6-91.0]	50.0 [0.6-91.0]	50.0 [0.6-91.0]
XIIa5. Mesothelioma	6	100.0 [-]	100.0 [-]	83.3 [27.3-97.5]
XIIb. Other unspecified malignant tumors	10	46.7 [15.0-73.7]	35.0 [8.5-64.0]	35.0 [8.5-64.0]
Total	6480	95.3 [94.8-95.8]	91.3 [90.6-92.0]	86.6 [85.8-87.4]

Abbreviations: AYA, adolescents and young adults; ICC-3, third version of the International Classification of Childhood Cancer; GCT, Germ-Cell tumors; cPNET, central primitive neuroectodermal tumors; pPNET, peripheral primitive neuroectodermal tumors; OS, Overall Survival; 95%CI, 95% confidence interval.

TABLE 4. TRENDS IN FIVE-YEAR OVERALL SURVIVAL AMONG ADOLESCENTS AND YOUNG ADULTS WITH CANCER FROM 1978 TO 2015 BY

ICCC-3 DIAGNOSTIC GROUPS

ICCC-3 diagnostic groups	5-year OS % (95%CI)					
	1978-82*	1983-87*	1988-92*	1993-97*	2000-07	2008-15
I. Leukemia	28.6 (16.1-41.1)	25.3 (13.9-36.7)	30.3 (20.3-40.3)	49.5 (38.9-60.1)	69.5 (63.5-74.7)	76.2 (71.5-80.2)
II. Lymphomas	72.0 (63.8-80.2)	77.9 (71.4-84.4)	86 (80.3-91.7)	87.6 (82.9-92.3)	93.4 (91.6-95.5)	94.2 (92.5-95.5)
III. Central Nervous System tumors	47.4 (32.5-62.3)#	61.7 (46.4-77.0)#	63.3 (51.3-75.3)#	60.1 (47.9-72.3)#	77.6 (72.1-82.2)	78.3 (73.9-82.1)
VIII. Malignant bone tumors	59.3 (40.7-77.9)	48.4 (30.8-66.0)	58.6 (43.5-73.7)	60.3 (47.2-73.4)	58.2 (48.9-66.4)	70.0 (62.7-76.2) ^a
IX. Soft tissue and other extraosseous sarcomas	56.1 (36.3-75.9)	50.7 (33.8-67.6)	49.2 (35.7-62.7)	73.1 (60.0-86.2)	65.6 (56.6-73.2)	67.5 (59.3-74.5)
X. Germ-cell tumors	73.2 (63.4-83.0)	78 (69.4-86.6)	88.5 (83.2-93.8)	91.1 (86.6-95.6)	91.2 (87.9-93.6)	94.3 (91.9-96.0) ^a
XI. epithelial neoplasms and melanomas	75.9 (66.7-85.1)	74.6 (66.6-82.6)	85.7 (80.4-91.0)	89.9 (86.2-93.6)	90.9 (88.5-92.9)	93.5 (91.8-94.7) ^a
All tumors	62.0 (57.5-66.5)	66.0 (61.9-70.1)	73.2 (70.1-76.3)	80.2 (77.7-82.8)	81.8 (79.5-83.9)	88.0 (86.9-89.0) ^a

Abbreviations: ICCC-3: third version of the International Classification of Childhood Cancer; OS: Overall Survival; 95%CI: 95% confidence interval

* Data from reference ²

Only malignant CNS tumors were included (exclusion of benign or borderline tumors)

^a p<0.05 Comparison between diagnostic periods (2000-2007 vs 2008-2015)

TABLE 5. TRENDS IN FIVE-YEAR OVERALL SURVIVAL AMONG ADOLESCENTS AND YOUNG ADULTS WITH CANCER FROM 2000-2007 TO 2008-2015 BY

ICCC-3 DIAGNOSTIC GROUPS

ICCC-3 diagnostic groups and subgroups	Adolescents (15-19 years)						Young adults (20-24 years)					
	2000-2007			2008-2015			2000-2007			2008-2015		
	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]
I. Leukemias, myeloproliferative/myelodysplastic diseases												
Ia. Lymphoblastic leukemias	83	69.6 [58.4-78.3]	100	79.3 [69.8-86.2] ^b	39	59.0 [42.0-72.5]	36	63.9 [46.1-77.2]	70	63.2 [50.4-73.5] ^b	46	71.4 [55.9-82.3]
Ib. Acute myeloid leukemias	39	59.0 [42.0-72.5]	55	62.4 [47.7-74.0]	35	94.3 [61.3-99.0]	35	94.3 [79.0-98.5]	54	98.1 [87.6-99.7]	54	98.1 [87.6-99.7]
Ic. Chronic myeloproliferative diseases	2	50.0 [0.6-91.0]	14	77.9 [45.9-92.3]	9	77.8 [36.5-93.9]	9	77.8 [36.5-93.9]	9	76.2 [33.2-93.5]	9	76.2 [33.2-93.5]
Id. Myelodysplastic / myeloproliferative diseases	3	33.3 [0.9-77.4]	6	50.0 [11.1-80.4]	3	33.3 [0.9-77.4]	3	33.3 [0.9-77.4]	2	50.0 [0.6-91.0]	2	50.0 [0.6-91.0]
Ie. Unspecified and other specified leukemias	286	94.4 [91.0-96.5]	409	95.0 [92.4-96.8]	335	93.4 [90.2-95.6]	335	93.4 [90.2-95.6]	564	93.6 [91.2-95.4]	564	93.6 [91.2-95.4]
II. Lymphomas and reticuloendothelial neoplasms												
IIa. Hodgkin disease	210	97.6 [94.3-99.0]	300	98.0 [95.5-99.1]	241	95.8 [92.4-97.7]	241	95.8 [92.4-97.7]	419	95.5 [93.0-97.2]	419	95.5 [93.0-97.2]
IIb. Non-Hodgkin lymphomas	66	86.2 [75.1-92.5]	85	86.8 [77.5-92.5]	82	87.8 [78.5-93.2]	82	87.8 [78.5-93.2]	132	88.2 [80.8-92.9]	132	88.2 [80.8-92.9]
IIc. Burkitt lymphoma	7	100.0 [-]	16	81.3 [52.5-93.5]	8	100.0 [-]	8	100.0 [-]	7	85.7 [33.4-97.9]	7	85.7 [33.4-97.9]
IId. Miscellaneous lymphoreticular neoplasms	3	33.3 [0.9-77.4]	7	100.0 [-]	3	66.7 [5.4-94.5]	3	66.7 [5.4-94.5]	4	100.0 [-]	4	100.0 [-]
IIe. Unspecified lymphoma	-	-	-	-	1	0.0 [-]	1	0.0 [-]	2	50.0 [0.6-91.0]	2	50.0 [0.6-91.0]
III. Central nervous system tumors												
IIIa. Ependymomas and choroid plexus tumor	119	80.3 [71.9-86.5]	187	81.3 [74.7-86.4]	151	75.4 [67.6-81.6]	151	75.4 [67.6-81.6]	238	76.0 [69.8-81.1]	238	76.0 [69.8-81.1]
IIIb. Astrocytomas	6	83.3 [27.3-97.5]	22	85.7 [53.9-96.2]	6	83.3 [27.3-97.5]	6	83.3 [27.3-97.5]	18	94.4 [66.6-99.2]	18	94.4 [66.6-99.2]
IIIc. Intracranial and intraspinal embryonal tumors	37	77.8 [60.4-88.2]	61	74.8 [61.6-84.0] ^b	43	73.7 [57.5-84.5] ^a	43	73.7 [57.5-84.5] ^a	72	49.6 [37.0-61.0] ^{a,b}	72	49.6 [37.0-61.0] ^{a,b}
IIId. Other gliomas	9	66.7 [28.2-87.8]	14	85.7 [53.9-96.2]	10	50.0 [18.4-75.3]	10	50.0 [18.4-75.3]	14	76.6 [43.3-91.9]	14	76.6 [43.3-91.9]
IIIe. Other specified intracranial and intraspinal neoplasms	16	43.8 [19.8-65.6]	24	47.8 [26.8-66.1]	50	67.3 [52.3-78.6]	50	67.3 [52.3-78.6]	51	73.6 [58.7-83.9]	51	73.6 [58.7-83.9]
IIIf. Unspecified intracranial and intraspinal neoplasms	45	97.7 [84.9-99.7]	61	96.7 [87.3-99.2]	38	94.4 [79.5-98.6]	38	94.4 [79.5-98.6]	73	95.3 [85.9-98.5]	73	95.3 [85.9-98.5]
IV. Neuroblastoma, other peripheral nervous cell tumors												
IVa. Neuroblastoma and ganglioneuroblastoma	6	83.3 [27.3-97.5]	5	100.0 [-]	5	80.0 [20.4-96.9]	5	80.0 [20.4-96.9]	11	90.0 [47.3-98.5]	11	90.0 [47.3-98.5]
IVb. Other peripheral nervous cell tumors	5	60.0 [12.6-88.2]	7	100.0 [-]	6	66.7 [19.5-90.4]	6	66.7 [19.5-90.4]	6	83.3 [27.3-97.5]	6	83.3 [27.3-97.5]
VI. Renal tumors												
VIa. Neuroblastoma and other nonepithelial renal tumors	2	0.0 [-]	4	100.0 [-]	1	100.0 [-]	1	100.0 [-]	2	50.0 [0.6-91.0]	2	50.0 [0.6-91.0]
VIb. Other peripheral nervous cell tumors	3	100.0 [-]	3	100.0 [-]	5	80.0 [20.4-96.9]	5	80.0 [20.4-96.9]	4	100.0 [-]	4	100.0 [-]
VII. Renal tumors												
VIIa. Nephroblastoma and other nonepithelial renal tumors	-	-	-	-	1	100.0 [-]	1	100.0 [-]	4	37.5 [1.1-80.8]	4	37.5 [1.1-80.8]
VIIb. Renal carcinoma	6	80.0 [20.4-96.9]	5	80.0 [20.4-96.9]	11	81.8 [44.7-95.1]	11	81.8 [44.7-95.1]	21	80.0 [54.9-92.0]	21	80.0 [54.9-92.0]
VIII. Hepatic tumors												
VIIIa. Hepatic carcinoma	6	33.3 [4.6-67.6]	5	60.0 [12.6-88.2]	6	16.7 [0.8-51.7]	6	16.7 [0.8-51.7]	14	35.7 [13.0-59.4]	14	35.7 [13.0-59.4]
VIIIb. Hepatic carcinoma	5	20.0 [0.8-58.2]	4	50.0 [5.8-84.5]	6	16.7 [0.8-51.7]	6	16.7 [0.8-51.7]	14	35.7 [13.0-59.4]	14	35.7 [13.0-59.4]

TABLE 5. (CONTINUED)

	15-19 years				20-24 years			
	2000-2007	2008-2015	2000-2007	2008-2015	2000-2007	2008-2015	2000-2007	2008-2015
	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]	n	5yOS [95%CI]
VIII. Malignant bone tumors	82	60.5 [49.0-70.2]^a	120	72.1 [62.8-79.4]^a	41	53.7 [37.4-67.4]	70	66.5 [53.7-76.6]
VIIIa. Osteosarcomas	46	55.6 [40-68.6] ^a	56	69.8 [55.3-80.4] ^a	14	28.6 [8.8-52.4] ^b	29	68.4 [47.9-82.2] ^b
VIIIb. Chondrosarcomas	8	75.0 [31.5-93.1]	7	100.0 [-]	8	100.0 [-]	19	87.7 [58.8-96.8]
VIIIc. Ewing tumor and related sarcomas of bone	25	64.0 [42.2-79.4]	53	70.4 [55.6-81.1] ^b	17	52.9 [27.6-73.0]	18	38.1 [16.6-59.5] ^b
VIII d. Other specified malignant bone tumors	3	66.7 [5.4-94.5]	3	66.7 [5.4-94.5]	2	50.0 [0.6-91.0]	4	100.0 [-]
IX. Soft tissue and other extrasosseous sarcomas	52	70.6 [56.0-81.1]	76	67.7 [55.3-77.4]	74	62.2 [50.1-72.1]	86	67.4 [55.7-76.6]
IXa. Rhabdomyosarcomas	12	41.7 [15.2-66.5]	17	58.8 [32.5-77.8]	13	38.5 [14.1-62.8]	10	50.0 [18.4-75.3]
IXb. Fibrosarcomas, and other fibrous neoplasms	10	77.8 [36.5-93.9]	16	80.8 [51.4-93.4]	13	76.9 [44.2-91.9]	11	81.8 [44.7-95.1]
IXd. Other specified soft tissue sarcomas	24	79.2 [57.0-90.8]	36	66.8 [47.6-80.3]	37	70.3 [52.8-82.3]	47	67.9 [50.8-80.2]
IXe. Unspecified soft tissue sarcomas	6	83.3 [27.3-97.5]	7	68.6 [21.3-91.2]	10	40.0 [12.3-67.0]	18	65.8 [39.1-83.0]
X. Germ-cell tumors, and other neoplasms of gonads	112	88.0 [80.2-92.8]	156	93.4 [88.1-96.4]	293	92.4 [88.7-94.9]	387	94.6 [91.8-96.5]
Xa. Intracranial and intraspinal germ cell tumors	12	75.0 [40.8-91.2]	17	82.4 [54.7-93.9]	5	80.0 [20.4-96.9]	8	100.0 [-]
Xb. Malignant extracranial and extragonadal germ cell tumors	7	71.4 [25.8-92.0]	6	66.7 [19.5-90.4]	12	66.7 [33.7-86.0]	22	72.7 [49.1-86.7]
Xc. Malignant gonadal germ cell tumors	88	92.9 [84.9-96.8]	118	96.4 [90.7-98.6]	255	94.4 [90.7-96.6]	337	97.2 [94.7-98.5]
Xd. Gonadal carcinoma	5	60.0 [12.6-88.2]	10	90.0 [47.3-98.5]	16	81.3 [52.5-93.5]	19	73.3 [47.2-87.9]
Xe. Other and unspecified malignant gonadal tumors	-	-	-	-	5	100.0 [-]	1	100.0 [-]
XI. Other malignant epithelial neoplasms and melanomas	207	92.1 [87.5-95.1]	342	93.7 [90.3-95.9]	463	90.4 [87.3-92.8]^a	871	93.4 [91.4-94.9]^a
XIa. Adrenocortical carcinomas	2	0.0 [-]	2	0.0 [-]	2	50.0 [0.6-91.0]	8	87.5 [38.7-98.1]
XIb. Thyroid carcinomas	94	100.0 [-]	136	100.0 [-]	166	99.4 [95.8-99.9]	350	99.1 [97.3-99.7]
XIc. Nasopharyngeal carcinomas	5	60.0 [12.6-88.2]	15	78.0 [45.5-92.5]	4	75.0 [12.8-96.1]	9	87.5 [38.7-98.1]
XId. Malignant melanomas	61	94.9 [85.1-98.3]	86	95.9 [87.8-98.7]	168	93.4 [88.4-96.3]	245	96.4 [93.0-98.2]
XIf. Other and unspecified carcinomas	45	79.8 [64.8-89.0]	103	87.3 [78.6-92.6]	123	75.4 [66.7-82.1]	260	83.2 [78.0-87.3]
XII. Other and unspecified malignant neoplasms	7	71.4 [25.8-92.0]	5	80.0 [20.4-96.9]	7	51.4 [11.8-81.3]	5	60.0 [12.6-88.2]
XIIa. Other specified malignant tumors	5	100.0 [-]	4	100.0 [-]	1	0.0 [-]	4	75.0 [12.8-96.1]
XIIb. Other unspecified malignant tumors	2	0.0 [-]	1	0.0 [-]	6	62.5 [14.2-89.3]	1	0.0 [-]
Total	1028	83.3 [80.8-85.5]^a	1511	87.0 [85.2-88.7]^a	1509	85.5 [83.6-87.2]^a	2432	88.6 [87.2-89.8]^a

Abbreviations: ICCC-3, third version of the International Classification of Childhood Cancer; 5yOS, Five-year Overall Survival; 95%CI, 95% confidence interval.

^a p<0.05 Comparison between diagnostic periods (2000-2007 vs 2008-2015)^b p<0.05 Comparison between age groups (15-19 years vs 20-24 years)

