



HAL
open science

Search for Effective Antimycotic Agents against Microsporium Gypseum from 61 Ethno Medicinal Plants of Hyderabad, Karnataka Region, Karnataka, India

Shivakumar Singh, Vidyasagar Gm

► To cite this version:

Shivakumar Singh, Vidyasagar Gm. Search for Effective Antimycotic Agents against Microsporium Gypseum from 61 Ethno Medicinal Plants of Hyderabad, Karnataka Region, Karnataka, India. Journal of Bioanalysis & Biomedicine, 2014, 06 (5), pp.33-35. 10.4172/1948-593x.1000106 . hal-04020783

HAL Id: hal-04020783

<https://hal.science/hal-04020783>

Submitted on 9 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Search for Effective Antimycotic Agents against *Microsporium Gypseum* from 61 Ethno Medicinal Plants of Hyderabad, Karnataka Region, Karnataka, India

Shivakumar Singh P and Vidyasagar GM*

Medicinal Plants and Microbiology Research Laboratory, Department of Post- Graduate Studies and Research in Botany, Gulbarga University, Gulbarga-585 106, Karnataka, India

Abstract

The 61 ethno medicinal plants of Hyderabad Karnataka region belonging to 33 different families used in skin diseases were screened for their antidermatophytic properties. Screening was carried out at 5 and 2.5 mg/ml concentrations of pet ether, chloroform, ethylacetate, methanol and aqueous extracts of each plant by agar well diffusion technique against *Microsporium gypseum*. Out of 61 plants, 05 (*Ceasalpinia bonducella*, *Coccinia indica*, *Corchorus olerarius*, *Lawsonia inermis* and *Tridax procumbens*) showed very effective antidermatophytic activity in ethyl acetate, chloroform and in aqueous extracts, effective activity observed in 11 plants (*Achyranthes aspera*, *Allium sativum*, *Celosia argentea*, *Citrus medica*, *Curcuma longa*, *Emblica officinalis*, *Gymnosporia montana*, *Lycopersicon esculentum*, *Milletia pinnata*, *Ricinus communis*, *Zingiber officinale*) in different extracts, whereas 38 plants showed moderate activity, 07 plants (*Euphorbia tirucalli*, *Lantana camara*, *Mentha viridis*, *Tinospora cordifolia* and *Tridax procumbens*) showed weak activity. The minimum inhibitory concentrations of 05 very effective plants were determined. The potential minimum inhibitory concentrations of 0.15 mg/ml conc. were detected from *C. indica*. This study provides scientific base for the isolation and purification of antidermatophytic compound(s) from ethno medicinal plants.

Keywords: Traditional medicinal plants; *Microsporium gypseum*; Antifungal screening

Introduction

Medicinal plants have been used in traditional treatment of skin diseases in worldwide [1], India [2] and Karnataka [3]. Herbal medicines are the basis of treatment and cure for various diseases in traditional methods practiced, such as Ayurveda, Unani and Siddha [4]. The rich availability and easy access to these medicinal plant resources have made them almost inevitable in the healthcare practices especially for those residing in the forest, hilly, rural and remote areas. Mycotic infections are the most common cause of skin infection in tropical developing countries. The incidence of dermatophytosis raised dramatically in the past one decade. Humid weather, over population and poor hygiene are the ideal conditions for the growth of dermatophytes [5]. These dermatophytes invade skin, hair and nail and cause dermatophytosis. Though these dermatophytes respond to treatment with conventional antifungal agents, the disease had a tendency to reoccur in the same area or other ones [4]. The most frequent fungal pathogens include the dermatophytes *Microsporium sp.*, *Trichophyton sp.*, and *Epidermophyton sp.* [6]. In the past decade, the incidence of dermatophytosis has risen dramatically. The humid weather, over population and poor hygienic conditions are ideally suited for the growth of dermatophytes and these factors are more important in country like India. Among the infectious diseases, diseases caused by fungal infections account for a larger proportion of health problems in humans [7]. Therefore, the present report focused on the antidermatophytic activity of petroleum ether, chloroform, ethylacetate, methanol and aqueous extracts of 61 plant parts against common dermatophytic fungi *Microsporium gypseum*.

Materials and Methods

Plant materials

Plant materials were collected from various localities of Hyderabad Karnataka region and Identified with the help of Gulbarga district flora

[8] the voucher specimens deposited in herbarium centre, Department of Botany, Gulbarga University, Karnataka, India. The collected plant materials were initially rinsed with distilled water to remove soil and other contaminants and dried on paper towel in laboratory at 37 ± 2°C for week.

Preparation of the plant extracts

The selected plant materials after shade drying were ground in a grinding machine in the laboratory. 25 g of shade dried powder was weighed and extracted successively with petroleum ether, chloroform, ethyl acetate, and methanol and aqueous in soxhlet extractor for 48 h. The extracts were concentrated under reduced pressure and preserved in refrigerator in airtight bottles for further use.

Extract stock solution: Dissolved 50 mg of crude in 10 ml DMF with glass beads, vortex to homogenize and a two-fold serial dilution was prepared. As a precaution not to miss trace amounts of antifungals for screening, a relatively high concentration of 2.5 to 5 mg/ml of each extract was prepared for bioassays.

Microbial culture and growth conditions

Test microorganism *Microsporium gypseum* used in the present study was obtained from M.R. medical college, Gulbarga, Karnataka,

*Corresponding author: Vidyasagar GM, Professor, Department of Botany, Gulbarga University, Gulbarga, Karnataka, India, Tel: 094-492-588-12; E-mail: gmvidyasagar@rediffmail.com

Received July 23, 2014; Accepted October 06, 2014; Published October 10, 2014

Citation: Lenjisa JL, Fereja TH (2014) A Retrospective Analysis of Prescribing Practices through WHO Prescribing Indicators at Four Selected Hospitals of West Ethiopia. J Bioanal Biomed 6: 033-036. doi:10.4172/1948-593X.1000106

Copyright: © 2014 Lenjisa JL, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

India. The Culture of *M. gypseum* grown on Sabouraud dextrose broth (HiMedia) at 28°C for 48 h and it was maintained on agar slants at 4°C.

Inoculum preparation

Stock inoculum of *M. gypseum* strain was prepared from 10-day cultures in Potato Dextrose Agar at 28°C to induce sporulation. Fungal colonies were covered with 5 mL of sterile saline solution (NaCl 0.85% w/v), the surface gently scraped with a sterile loop and the resultant mixture of fungal units was then transferred to a sterile tube. The turbidity of the final inoculum was standardized according to a McFarland scale 0.5 tubes and adjusted to a fungal population of 106 Colony Former Units (CFU). The confirmation of inoculum quantification was done by plating 0.01 mL of inoculum suspension in Sabouraud Dextrose Agar (SDA). The dishes were incubated at 28°C and examined daily for the presence of fungal colonies which were counted as soon as growth became visible [8,9].

Agar-well diffusion method

The assay was conducted by agar well diffusion method [10]. About 15 to 20 ml of potato dextrose agar medium was poured in the sterilized petri dishes and allowed to solidify. Fungal lawn was prepared using 5 days old culture strains. The fungal strains were suspended in a saline solution (0.85% NaCl) and adjusted to a turbidity of 0.5 Mac Farland standards (108 CFU/ml). 1 ml of fungal strain was spread over the medium using a sterilized glass spreader. Using flamed sterile borer, wells of 4 mm diameter were punctured in the culture medium and required concentrations of serially diluted extract (2.5, 5 mg/ml) was added to the 20 µl to each wells. The plates thus prepared were left for diffusion of extracts into media for one hour in the refrigerator and then incubated at 30°C. After incubation for 48 h, the plates were observed for zone of inhibition. Diameter zone of inhibition was measured and expressed in millimeters. Dimethyl formamide (DMF) was used as a negative control. The experiments were conducted in triplicates. The obtained results classified into three groups on the basis of zone of inhibition i.e., very effective (above 12 mm), effective (10-11 mm), moderate (7-9 mm), weak (No zone of inhibition).

Minimum Inhibitory Concentration

One ml of sterile liquid Sabouraud medium was added to 08 sterile capped tubes, 1 ml of each solvent extracts suspension was added to tube 1 [11]. The contents were mixed and 1 ml was transferred to tube 2. This serial dilution was repeated through to tube six and 1 ml was discarded from tube 6. 50 µl of inoculum was added to tubes 1-8 and the contents were mixed. Medium control (no inoculum and no drug) and inoculum control (no drug) tubes were prepared. The final concentrations of each plant solvent extracts ranged from 0.5 mg/ml to 0.15 mg/ml. The tubes were incubated at 30°C for 96 h. The fungal growth in each tube was evaluated visually depending up on the turbidity in the tubes. MIC was defined as the drug concentration at which the turbidity of the medium was the same as the medium control.

Statistical analysis


All the experiments were conducted in triplicate unless stated otherwise and statistical analysis of the data was performed by Analysis Of Variance (ANOVA), using STATISTICA 5.5 (Stat Soft Inc, Tulsa, Oklahoma, USA) software. A probability value of difference $p \sim 0.05$ was considered to denote a statistically significance All data were presented as mean values \pm Standard Deviation (SD).

Results and Discussion

The 61 ethno medicinal plants of Hyderabad Karnataka region belonging to 33 different families used in skin diseases were screened for their antidermatophytic properties against *Microsporum gypseum*. The screening was carried out at 5 and 2.5 mg/ml concentrations of pet ether, chloroform, ethylacetate, methanol and aqueous extracts of each plant by agar well diffusion technique the obtained results were given in (Table 1). Out of 61 plants, 05 (*Ceasalpinia bonducella*, *Coccinia indica*, *Corchorus olerius*, *Lawsonia inermis* and *Tridax procumbens*) showed very effective antidermatophytic activity in ethyl acetate, chloroform and in aqueous extracts. The effective activity observed in 11 plants (*Achyranthes aspera*, *Allium sativum*, *Celosia argentea*, *Citrus medica*, *Curcuma longa*, *Embllica officinalis*, *Gymnosporia montana*, *Lycopersicon esculentum*, *Milletia pinnata*, *Ricinus communis*, *Zingiber officinale*) in different extracts, whereas 38 plants showed moderate activity, 07 plants (*Euphorbia tirucalli*, *Lantana camara*, *Mentha viridis*, *Tinospora cordifolia* and *Tridax procumbens*) showed weak activity. The negative control (DMF) was not showed activity, while the standard drug, Ketoconazole significantly inhibited (28.66 ± 1.15 to 12.33 ± 1.52 mm) the growth of the test dermatophyte.

The minimum inhibitory concentrations of very effective 05 plants were determined. The 0.15 mg/ml of minimum inhibitory concentration was recorded with *C. indica* followed by 0.31 mg/ml conc. observed with three plants extracts i.e., *C. bonducella*, *C. olerius*, *L. inermis*. Whereas the weak 1.25 mg/ml minimum inhibitory concentration was observed with *T. procumbens* (Figure 1).

The response of *M. gypseum* to treatment with various plants extracts varied from solvent extract to extract. The ethyl acetate seed extract of *C. bonducella* was showed effective activity against test dermatophyte. This was supported by the previous work of Kavitha Sagar and Vidyasagar GM. Where they used ethyl acetate leaves extracts of *C. bonducella* [12]. Farukh and Iqbal reported antimicrobial activity using seed extract of *C. bonducella* [13]. In the present report *Lawsonia inermis* leaves showed very effective activity in chloroform extract it was correlated with past reports of Bhatnagar et al. [14], Misra and Dixit [15]. Whereas in another report the effective activity showed in ethanolic leaves extract against *T. rubrum*, *T. mentagrophytes* [16]. The ethanolic extract of the whole plant of *Lawsonia inermis* showed antifungal activity against *Trichophyton mentagrophytes* and *Microsporum canis* [17]. In the present study, the ethylacetate seed extract of *Corchorus olerius* showed an effective activity. Though no report is available on the antidermatophytic activity of *Corchorus*



Sl. No.	Name of the Plant	Part used	Zone of Inhibition in different solvent extracts (mm)										C		S	
			P		C		E		M		A		DMF	Ketoconazole		
			1	2	1	2	1	2	1	2	1	2				
01	<i>Achyranthesaspera</i> L.	L	07.66 ± 1.15	06.20 ± 1.00	11.00 ± 0.00	06.33 ± 1.52	07.00 ± 0.00	05.33 ± 1.52	05.33 ± 1.52	07.00 ± 0.00	04.50 ± 0.00	NA	NA	17.66 ± 0.57		
02	<i>Aeglemarmelos</i> (L.)	L	07.00 ± 0.00	04.00 ± 1.00	06.33 ± 1.52	04.33 ± 1.52	10.33 ± 1.52	05.33 ± 1.52	05.33 ± 1.52	NA	06.33 ± 1.52	NA	NA	18.33 ± 1.52		
03	<i>Allium cepa</i> Linn.	B	05.00 ± 0.00	04.66 ± 1.15	04.66 ± 1.15	04.33 ± 1.52	04.66 ± 1.15	NA	10.00 ± 0.00	05.00 ± 0.00	06.00 ± 1.00	05.00 ± 0.00	NA	12.33 ± 1.52		
04	<i>Allium sativum</i> L.	B	07.33 ± 0.57	06.66 ± 0.57	11.33 ± 0.57	06.66 ± 0.57	07.00 ± 0.00	05.66 ± 0.57	05.00 ± 0.00	07.00 ± 0.00	NA	NA	NA	17.33 ± 1.52		
05	<i>Aloe vera</i> L.	L	08.66 ± 1.15	04.00 ± 0.00	05.33 ± 0.57	04.66 ± 0.57	07.00 ± 0.00	07.33 ± 0.57	06.66 ± 0.57	06.66 ± 1.15	05.33 ± 0.57	NA	NA	18.33 ± 1.52		
06	<i>Amaranthuspinosus</i> L.	L	07.00 ± 0.00	05.00 ± 1.00	05.33 ± 1.52	05.66 ± 1.52	05.00 ± 0.00	05.00 ± 1.00	07.33 ± 1.15	04.66 ± 1.57	05.00 ± 1.00	NA	NA	18.33 ± 1.52		
07	<i>Annonareticulata</i> L.	L	05.66 ± 0.57	04.00 ± 0.00	05.00 ± 1.00	04.33 ± 0.57	08.66 ± 0.57	05.00 ± 0.00	07.00 ± 0.00	04.66 ± 0.57	05.33 ± 1.15	04.00 ± 0.00	NA	21.66 ± 1.15		
08	<i>Annonasquamosa</i> L.	L	07.00 ± 0.00	05.00 ± 0.00	06.66 ± 1.52	06.33 ± 1.15	07.00 ± 0.00	06.66 ± 0.57	07.00 ± 0.00	06.33 ± 0.57	07.66 ± 1.57	NA	NA	20.33 ± 1.52		
09	<i>Argemonemexicana</i> L.	L	05.66 ± 1.52	04.00 ± 0.00	0.66 ± 0.57	04.66 ± 1.52	05.33 ± 1.52	04.33 ± 0.57	05.66 ± 1.52	04.00 ± 0.00	04.66 ± 0.57	NA	NA	15.66 ± 0.57		
10	<i>Azadirachtaindica</i> A. Juss.	L	05.00 ± 0.00	06.66 ± 0.57	05.33 ± 0.57	07.00 ± 0.00	08.66 ± 0.57	05.33 ± 1.52	07.66 ± 0.57	10.00 ± 0.00	04.33 ± 0.57	NA	NA	24.00 ± 1.00		
11	<i>Bergerakoentigi</i> L.	L	05.66 ± 0.57	04.00 ± 0.00	07.00 ± 0.00	05.33 ± 0.57	10.33 ± 1.52	06.66 ± 1.52	06.33 ± 1.52	05.33 ± 1.52	07.66 ± 1.52	05.66 ± 1.57	NA	18.00 ± 1.00		
12	<i>Buteamonosperma</i> (Lam.) Taub.	L	06.66 ± 1.52	05.00 ± 0.00	07.66 ± 1.52	06.66 ± 1.52	08.33 ± 1.52	05.00 ± 0.00	08.66 ± 1.52	08.66 ± 1.52	04.33 ± 0.57	NA	NA	19.66 ± 0.57		
13	<i>Cajanuscajan</i> (L.) Mill.	L	07.66 ± 1.57	05.00 ± 1.00	06.00 ± 1.00	05.00 ± 1.00	06.00 ± 1.00	05.66 ± 1.57	06.66 ± 1.57	05.00 ± 1.00	NA	NA	NA	22.33 ± 1.52		
14	<i>Calotropisgigantea</i> L.	L	06.66 ± 1.57	06.00 ± 1.00	07.33 ± 0.57	06.66 ± 1.57	06.66 ± 1.57	06.33 ± 0.57	06.33 ± 0.57	06.66 ± 1.57	05.66 ± 1.57	NA	NA	19.33 ± 1.52		
15	<i>Carica papaya</i> L.	L	05.00 ± 0.00	05.00 ± 0.00	07.00 ± 1.00	05.00 ± 0.00	07.66 ± 0.57	04.33 ± 0.57	07.00 ± 0.00	05.00 ± 1.00	08.00 ± 0.00	05.66 ± 1.57	NA	16.33 ± 1.52		
16	<i>Ceasalpinibonducella</i> (L.) Flem.	S	13.66 ± 1.57	08.33 ± 1.15	13.00 ± 0.00	07.33 ± 1.15	12.33 ± 1.15	07.33 ± 1.15	10.66 ± 1.57	07.33 ± 1.15	05.00 ± 0.00	NA	NA	23.66 ± 0.57		
17	<i>Celosia argentea</i> L.	S	05.33 ± 1.52	05.33 ± 1.15	06.33 ± 1.52	05.66 ± 0.57	07.33 ± 1.52	05.33 ± 0.57	11.33 ± 0.57	04.33 ± 1.52	05.33 ± 0.57	NA	NA	18.33 ± 1.52		
18	<i>Citrus medica</i> L.	L	08.00 ± 0.00	05.33 ± 0.57	10.00 ± 0.00	08.66 ± 1.52	11.66 ± 0.57	06.00 ± 0.00	10.33 ± 0.57	07.66 ± 1.52	04.33 ± 1.15	NA	NA	15.00 ± 0.00		
19	<i>Coccinia indica</i> Wt. & Arn.	L	10.00 ± 0.00	05.66 ± 0.57	07.33 ± 1.15	05.33 ± 0.57	10.33 ± 1.15	06.66 ± 1.52	07.33 ± 0.57	06.33 ± 1.15	19.33 ± 1.15	07.66 ± 1.57	NA	18.00 ± 0.00		
20	<i>Corchorusoleterius</i> L.	S	07.00 ± 0.00	NA	19.66 ± 0.57	05.33 ± 1.15	21.00 ± 0.00	05.33 ± 0.57	08.66 ± 1.52	05.00 ± 0.00	11.33 ± 1.15	05.66 ± 1.57	NA	18.00 ± 0.00		
21	<i>Coriandratsativum</i> L.	A	05.33 ± 1.15	04.00 ± 0.00	07.33 ± 0.57	04.33 ± 1.15	09.00 ± 0.00	05.33 ± 0.57	07.33 ± 1.15	04.33 ± 1.15	06.66 ± 1.52	NA	NA	24.66 ± 1.15		
22	<i>Cryptolepisbuchananii</i> Roem & Schult.	A	07.00 ± 0.00	05.66 ± 1.52	07.66 ± 1.52	07.00 ± 0.00	07.33 ± 0.57	05.66 ± 1.52	07.00 ± 0.00	06.01 ± 0.00	05.33 ± 0.57	NA	NA	19.00 ± 0.00		
23	<i>Curcuma longa</i> Linn.	R	18.66 ± 1.52	09.33 ± 0.57	11.33 ± 1.15	08.33 ± 1.15	07.33 ± 0.57	06.00 ± 0.00	09.33 ± 1.15	05.66 ± 1.52	07.01 ± 0.00	05.66 ± 1.57	NA	18.00 ± 0.00		
24	<i>Dalbergiasisso</i> Roxb.	L	05.33 ± 1.52	05.66 ± 0.57	06.66 ± 1.57	05.66 ± 1.52	07.33 ± 1.52	05.66 ± 0.57	07.66 ± 0.57	05.66 ± 1.52	06.66 ± 1.52	NA	NA	17.66 ± 1.15		
25	<i>Daturametel</i> L.	L	05.66 ± 1.57	05.66 ± 0.57	07.66 ± 1.52	05.66 ± 0.57	06.66 ± 1.52	04.66 ± 1.57	06.66 ± 1.52	05.66 ± 0.57	06.33 ± 1.52	NA	NA	17.33 ± 1.52		
26	<i>Emblicaoofficinalis</i> Gaertn.	L	06.33 ± 1.52	05.33 ± 1.52	10.00 ± 0.00	05.33 ± 1.52	13.66 ± 1.57	06.33 ± 1.52	07.33 ± 1.15	05.33 ± 1.52	04.66 ± 0.57	NA	NA	20.00 ± 0.00		
27	<i>Euphorbia tirucalli</i> L.	L	NA	NA	04.66 ± 0.57	04.00 ± 0.00	05.66 ± 1.52	04.33 ± 1.52	06.33 ± 1.52	05.66 ± 0.57	NA	NA	NA	18.66 ± 0.57		
28	<i>Ficusracemosa</i> L.	L	05.66 ± 0.57	05.66 ± 1.57	05.33 ± 1.52	05.66 ± 0.57	09.66 ± 0.57	07.33 ± 1.15	07.33 ± 1.52	04.66 ± 1.52	07.33 ± 1.15	05.33 ± 1.52	NA	20.00 ± 1.00		
29	<i>Gymnosporiamontana</i> (Roth) Benth	L	06.66 ± 1.57	05.66 ± 1.57	08.33 ± 1.15	05.66 ± 0.57	11.33 ± 1.15	05.00 ± 0.00	11.00 ± 1.00	06.00 ± 0.00	06.66 ± 1.57	NA	NA	16.00 ± 0.00		
30	<i>Hibiscus rosa-sinensis</i> L.	F	07.33 ± 1.52	05.33 ± 1.52	06.33 ± 1.52	04.66 ± 0.57	06.00 ± 1.00	05.33 ± 1.15	08.00 ± 0.00	04.00 ± 1.00	04.00 ± 1.00	NA	NA	19.00 ± 0.00		
31	<i>Hyptissuoveolens</i> (L.) Poit.	L	06.66 ± 1.52	05.66 ± 1.57	07.00 ± 0.00	05.33 ± 1.15	07.66 ± 1.52	06.33 ± 1.15	06.66 ± 0.57	05.33 ± 1.52	04.66 ± 0.57	NA	NA	19.00 ± 1.00		
32	<i>Ixoracoccinea</i> L.	F	05.00 ± 0.00	04.33 ± 1.15	06.33 ± 1.15	05.33 ± 0.57	06.33 ± 1.15	05.00 ± 0.00	07.33 ± 1.15	05.66 ± 1.52	05.66 ± 1.52	NA	NA	16.00 ± 0.00		
33	<i>Jatrophaglandulifera</i> Roxb.	L	05.33 ± 1.52	04.33 ± 1.15	07.66 ± 1.52	05.00 ± 0.00	06.66 ± 0.57	05.33 ± 1.15	07.66 ± 1.57	05.66 ± 1.52	06.33 ± 0.57	NA	NA	19.00 ± 0.00		
34	<i>Lantana camara</i> L.	L	NA	NA	05.66 ± 1.52	05.66 ± 1.52	NA	NA	05.66 ± 1.52	NA	NA	NA	NA	19.66 ± 1.15		
35	<i>Lawsoniainermis</i> Linn.	L	04.00 ± 0.00	04.33 ± 1.15	13.33 ± 1.52	06.33 ± 0.57	12.66 ± 1.52	05.33 ± 1.15	09.66 ± 1.57	05.33 ± 1.15	05.33 ± 0.57	NA	NA	24.33 ± 1.52		
36	<i>Lycopersiconesculentum</i> L.	L	08.66 ± 1.52	05.33 ± 1.52	09.33 ± 1.15	05.00 ± 0.00	06.3 ± 1.15	05.66 ± 1.52	09.33 ± 0.57	04.3 ± 1.15	10.3 ± 0.57	05.66 ± 1.57	NA	17.66 ± 0.57		
37	<i>Mangifera indica</i> Linn.	L	05.33 ± 1.52	05.33 ± 1.15	07.66 ± 1.57	06.00 ± 0.00	06.33 ± 1.52	05.66 ± 1.57	06.33 ± 0.57	04.00 ± 0.00	05.3 ± 1.15	NA	NA	19.00 ± 1.00		
38	<i>Menthavididis</i> L.	A	NA	04.66 ± 1.52	04.00 ± 1.00	04.66 ± 1.52	04.33 ± 0.57	04.00 ± 1.00	04.33 ± 0.57	04.6 ± 1.57	NA	NA	NA	18.50 ± 0.00		
39	<i>Milletiapinnata</i> (L.) Panigrahi	L	07.00 ± 1.00	09.33 ± 1.52	10.00 ± 1.00	10.00 ± 1.00	08.66 ± 1.52	10.00 ± 1.00	08.66 ± 1.52	09.00 ± 1.00	05.66 ± 1.52	NA	NA	18.33 ± 1.52		
40	<i>Momordicacharantia</i> L.	L	07.33 ± 1.52	06.33 ± 1.52	07.00 ± 1.00	06.33 ± 0.57	06.33 ± 0.57	05.00 ± 1.00	06.66 ± 0.57	05.33 ± 1.52	NA	NA	NA	18.66 ± 0.57		
41	<i>Neriumdorum</i> Solander.	L	05.33 ± 1.52	04.66 ± 1.57	04.00 ± 1.00	04.66 ± 0.57	07.33 ± 1.15	06.00 ± 1.00	06.33 ± 1.15	06.33 ± 1.52	NA	NA	NA	21.33 ± 1.52		
42	<i>Ocimum sanctum</i> L.	A	06.00 ± 1.00	04.33 ± 1.52	07.33 ± 0.57	NA	05.66 ± 1.52	04.33 ± 1.15	05.66 ± 0.57	04.00 ± 1.00	04.66 ± 0.57	NA	NA	20.66 ± 0.57		
43	<i>Piper nigrum</i> L.	S	04.00 ± 1.00	07.33 ± 0.57	NA	05.00 ± 1.00	06.66 ± 1.57	05.33 ± 1.52	08.00 ± 1.00	05.33 ± 0.57	04.66 ± 1.57	NA	NA	16.33 ± 1.52		
44	<i>Plumbagozeylanica</i> L.	L	07.33 ± 0.57	04.66 ± 0.57	06.00 ± 1.00	05.33 ± 0.57	08.33 ± 1.52	04.00 ± 0.00	08.33 ± 0.57	05.33 ± 1.52	08.66 ± 1.57	05.66 ± 1.57	NA	22.00 ± 1.00		
45	<i>Ricinuscommunis</i> L.	S	07.66 ± 0.57	05.66 ± 1.57	06.66 ± 0.57	05.66 ± 1.52	10.66 ± 0.57	05.33 ± 1.52	08.66 ± 1.57	05.66 ± 0.57	04.33 ± 1.52	NA	NA	18.66 ± 1.15		
46	<i>Santalum album</i> L.	L	04.66 ± 1.57	04.33 ± 1.15	04.66 ± 0.57	06.33 ± 1.52	09.66 ± 0.57	05.66 ± 1.52	07.33 ± 1.52	04.66 ± 0.57	05.66 ± 1.57	NA	NA	17.00 ± 0.00		
47	<i>Sennaauriculata</i> (L.) Roxb.	F	06.00 ± 0.00	04.66 ± 0.57	07.66 ± 1.52	05.33 ± 1.15	07.66 ± 1.52	05.33 ± 1.52	08.33 ± 1.15	05.66 ± 1.57	05.66 ± 0.57	NA	NA	21.00 ± 0.00		
48	<i>Sennatoral</i> L.	L	06.33 ± 0.57	05.66 ± 1.52	06.66 ± 1.57	05.66 ± 0.57	09.33 ± 0.57	05.66 ± 1.57	08.66 ± 1.57	06.66 ± 1.52	05.33 ± 1.52	NA	NA	16.66 ± 0.57		
49	<i>Solanumnigrum</i> L.	L	06.66 ± 0.57	05.00 ± 0.00	08.66 ± 0.57	07.00 ± 0.00	08.33 ± 1.15	07.66 ± 0.57	08.66 ± 1.52	08.66 ± 1.57	08.33 ± 1.15	06.33 ± 1.15	NA	21.66 ± 1.15		
50	<i>Sterculiafoetida</i> L.	S	10.33 ± 0.57	05.33 ± 1.52	08.33 ± 1.15	06.66 ± 1.57	07.66 ± 1.57	06.66 ± 0.57	09.33 ± 1.52	06.66 ± 0.57	07.00 ± 0.00	NA	NA	21.00 ± 0.00		
51	<i>Semecarpusanacardium</i> L.	B	07.33 ± 1.52	05.66 ± 0.57	07.66 ± 1.57	05.66 ± 1.52	06.66 ± 0.57	05.66 ± 1.52	06.33 ± 1.52	05.66 ± 0.57	NA	NA	NA	17.66 ± 1.15		
52	<i>Tamarindus indica</i> Linn.	L	NA	04.66 ± 0.57	06.66 ± 1.57	04.66 ± 0.57	08.33 ± 1.52	07.66 ± 1.57	08.33 ± 1.52	06.00 ± 0.00	06.33 ± 1.52	NA	NA	21.00 ± 0.00		
53	<i>Tectonagrandidis</i> L.	L	06.66 ± 1.57	05.66 ± 1.57	08.66 ± 0.57	06.66 ± 1.57	07.00 ± 0.00	04.33 ± 1.52	09.00 ± 1.00	06.66 ± 0.57	05.66 ± 1.57	NA	NA	18.66 ± 0.57		
54	<i>Tinosporacordifolia</i> (Willd.) J. Hook & Thoms.	L	NA	NA	08.66 ± 1.57	05.33 ± 1.15	08.00 ± 0.00	05.33 ± 1.52	07.66 ± 0.57	06.33 ± 1.15	05.33 ± 0.57	NA	NA	05.33 ± 1.52		
55	<i>Tephrosiapurpurea</i> (L.) Pers.	L	05.33 ± 1.52	05.66 ± 0.57	06.33 ± 0.57	05.33 ± 0.57	08.33 ± 1.52	05.66 ± 0.57	09.00 ± 0.00	04.33 ± 0.57	04.66 ± 1.57	NA	NA	17.66 ± 1.15		
56	<i>Thevetianerifolia</i> Juss.	L	05.33 ± 0.57	04.00 ± 0.00	07.00 ± 0.00	04.66 ± 0.57	08.66 ± 1.57	05.33 ± 0.57	06.33 ± 1.52	04.00 ± 0.00	05.00 ± 0.00	NA	NA	22.00 ± 0.00		
57	<i>Tribulusterestril</i> L.	A	06.66 ± 1.57	04.66 ± 1.57	09.00 ± 0.00	05.66 ± 1.57	07.33 ± 1.15	05.66 ± 0.57	07.33 ± 1.52	05.66 ± 1.57	05.33 ± 1.52	NA	NA	20.00 ± 0.00		
58	<i>Tridaxprocumbens</i> Linn.	A	07.66 ± 0.57	14.66 ± 1.57	12.33 ± 1.52	16.66 ± 0.57	NA	11.66 ± 1.57	06.33 ± 0.57	08.00 ± 0.00	NA	NA	NA	21.66 ± 1.15		
59	<i>Vitexnegundo</i> L.	L	05.33 ± 1.52	05.00 ± 0.00	05.66 ± 0.57	05.00 ± 0.00	09.33 ± 1.52	07.00 ± 0.00	07.33 ± 0.57	04.00 ± 1.00	07.00 ± 0.00	0				

oleterius. In this study at lower concentration of *Allium sativum* aqueous extract has not showed activity whereas the report of Sowjanya NC, and Manohara C Chary revealed the aqueous extracts of *Allium sativum* and *Ocimum sanctum* at 10% concentration were more pronounced antifungal properties against *Microsporum gypseum* [18].

In the present report the best MIC was obtained on *M. gypseum* with remarkable activity for ethylacetate leaf extract of *C. indica*. This is supported by the work of Narasimha et al., [19] where they reported effective antibacterial activity of floral petal extracts of *C. Indica*.

Conclusion

Of the antimycotic activity of 61 ethno medicinal plants (05 very effective + 11 effective) showed maximum activity against *M. gypseum*. This may be attributed to the various phytochemical constituents present in the crude extracts. The purified components may have even more potency with respect to inhibition of dermatophytes. The work carried was a basic approach to find out the effective antimycotic agents from 61 medicinal plants. Further works on the types of phytoconstituents and purification of individual group (s) of bioactive compounds (s) can reveal the exact potential of the plants to inhibit skin pathogenic microbes.

Acknowledgement

The authors are thankful to UGC for extending financial support and Gulbarga University for providing lab facilities.

References

1. Adeleye A, Ogunniyi AA, Omonigbehin EA (2011) Antimicrobial activity of some local herbs on common skin pathogens. Bioscience Research Communications 15: 231-236.
2. Kotoky J, Das PN (2008) Medicinal Plants used for liver diseases in some parts of Kamrup district of Assam, a North Eastern State of India. Fitoterapia 79: 384-387.
3. Shiva kumar Singh P, Vidyasagar GM (2013) Ethno medicinal plants used in the treatment of skin diseases in Hyderabad Karnataka region, Karnataka, India. Asian Pacific Journal of Tropical Biomedicine 3: 882-886.
4. Sukhdev (1997) Ethanotherapeutics and modern drug development. The Potential Aurvedha Curr Sci 73: 909-928.
5. Di Santo R (2010) Natural products as antifungal agents against clinically relevant pathogens. Nat Prod Rep 27: 1084-1098.
6. Simaljakova M, Skutilova (1995) E. Bratisl-Lek-Listy 96: 122-126.
7. Seetharam YN, Kotresh K, Upalaonkar SB (2000) Flora of Gulbarga district (Gulbarga University, Gulbarga).
8. Santos DA, Barros MES, Hamdan JS (2006) Establishing a method of inoculum preparation for susceptibility testing of *Trichophytonrubrum* and *Trichophytonmentagrophytes*. J Clin Microbiol 44: 98-101.
9. Hadacek F, Greger H (2000) Testing of antifungal natural products: methodologies, comparability of results and assay choice. Phytochem Anal 11: 137-147.
10. Magaldi S, Mata-Essayag S, Hartung de Capriles C, Perez C, Colella MT, et al. (2004) Well diffusion for antifungal susceptibility testing. International Journal of Infectious Diseases 8: 39-45.
11. National Committee for Clinical Laboratory Standards (NCCLS) (1997). Approved Standard M2-A6(5th edition) NCCLS: Wayne, PA.
12. Kavitha Sagar, Vidyasagar GM (2013) Anti-dermatophytic activity of some traditionally used medicinal plants of North Karnataka Region. J App Pharm Sci 3: 077-083.
13. Ayneshi Y, Emami M, Shidfar MR, Moghadam M, Lordbacheh P, et al. (1999) Broad-spectrum antibacterial Screening of Iranian Plants of antifungal activity Part 1. DARU. 10: 78-89.
14. Bhatnagar SS, Santapau H, Desa JDH, Maniar AC, Ghadially NC, et al. (1961) Biological activity of Indian Medicinal plants Part I. Antibacterial, Antitubercular and Antifungal action. Ind J Med Res 49: 799-805.
15. Misra SS, Dixit SN (1979) Antifungal activity of leaf-extracts of some higher plants. Acta Bot Indica 7: 147-150.
16. Vijayanthimala J, Rajendra Prasad J, Anandi C, Pugalendi KV (2004) Antidermatophytic activity of some Indian medicinal plants. Journal of Natural Remedies 26: 26-31.
17. Bhakuni DS, Dhar ML, Dhar MM, Dhawan, BN, Gupta B, Srimal RC. Indian J. Exp. Biol 1971; 9: 91.
18. Sowjanya NC, Manohara Chary C (2012) Effect of plant extracts on the growth of *Microsporumgypseum*. Journal of Phytology 4: 41-44.
19. Narasimaha G, Sridevi A, Vijayalakshmi M (2011) Antibacterial activity of some higher plant floral petals. NPAIJ 7: 168-170.