



HAL
open science

The contribution of weathering of the main Alpine rivers on the global carbon cycle

Marco Donnini, Jean-Luc Probst, Anne Probst, Francesco Frondini, Ivan
Marchesini, Fausto Guzzetti

► To cite this version:

Marco Donnini, Jean-Luc Probst, Anne Probst, Francesco Frondini, Ivan Marchesini, et al.. The contribution of weathering of the main Alpine rivers on the global carbon cycle. EGU General Assembly 2013, Apr 2013, Vienne, Austria. pp.0. hal-04021127

HAL Id: hal-04021127

<https://hal.science/hal-04021127>

Submitted on 9 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 8939

To cite this version: Donnini, Marco and Probst, Jean-Luc and Probst, Anne and Frondini, Francesco and Marchesini, Ivan and Guzzetti, Fausto *The contribution of weathering of the main Alpine rivers on the global carbon cycle.* (2013) In: EGU General Assembly 2013, 07-12 Apr 2013, Vienna, Austria.

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

The contribution of weathering of the main Alpine rivers on the global carbon cycle

Marco Donnini (1,4), Jean-Luc Probst (2,3), Anne Probst (2,3), Francesco Frondini (1), Ivan Marchesini (4), and Fausto Guzzetti (4)

(1) Università degli Studi di Perugia. Dipartimento di Scienze della Terra. Perugia, Italy, (4) Consiglio Nazionale delle Ricerche, Istituto di Ricerca per la Protezione Idrogeologica, Perugia, Italy, (2) University of Toulouse ; INPT, UPS ; Laboratoire Ecologie Fonctionnelle et Environnement (EcoLab), ENSAT, Castanet Tolosan, France, (3) Centre National de la Recherche Scientifique (CNRS), EcoLab, ENSAT Castanet Tolosan, France

The contribution of weathering of the main Alpine rivers on the global carbon cycle

Abstract

On geological time-scales the carbon fluxes from the solid Earth to the atmosphere mainly result from volcanism and metamorphic-decarbonation processes, whereas the carbon fluxes from atmosphere to solid Earth mainly depend on weathering of silicates and carbonates, biogenic precipitation and removal of CaCO_3 in the oceans and volcanic gases – seawater interactions. Quantifying each contribution is critical. In this work, we estimate the atmospheric CO_2 uptake by weathering in the Alps, using results of the study of the dissolved loads transported by 33 main Alpine rivers. The chemical composition of river water in unpolluted areas is a good indicator of surface weathering processes (Garrels and Mackenzie, 1971; Drever, 1982; Meybeck, 1984; Tardy, 1986; Berner and Berner, 1987; Probst et al., 1994). The dissolved load of streams originates from atmospheric input, pollution, evaporite dissolution, and weathering of carbonate and silicate rocks, and the application of mass balance calculations allows quantification of the different contributions. In this work, we applied the MEGA (Major Element Geochemical Approach) geochemical code (Amiotte Suchet, 1995; Amiotte Suchet and Probst, 1996) to the chemical compositions of the selected rivers in order to quantify the atmospheric CO_2 consumed by weathering in Alpine region. The drainage basins of the main Alpine rivers were sampled near the basin outlets during dry and flood seasons. The application of the MEGA geochemical consisted in several steps. First, we subtracted the rain contribution in river waters knowing the X/Cl ($X = \text{Na}, \text{K}, \text{Mg}, \text{Ca}$) ratios of the rain. Next, we considered that all $(\text{Na}+\text{K})$ came from silicate weathering. The average molar ratio $R_{\text{sil}} = (\text{Na}+\text{K})/(\text{Ca}+\text{Mg})$ for rivers draining silicate terrains was estimated from unpolluted French stream waters draining small monolithological basins (Meybeck, 1986; 1987). For the purpose, we prepared a simplified geo-lithological map of Alps according to the lithological classification of Meybeck (1986, 1987). Then for each basin we computed R_{sil} weighted average considering the surface and the mean precipitation for the surface area of each lithology. Lastly, we estimated the $(\text{Ca}+\text{Mg})$ originating from carbonate weathering as the remaining cations after silicate correction. Depending on time-scales of the phenomena (shorter than about 1 million year i.e. correlated to the short term carbon cycle, or longer than about 1 million years i.e. correlated to the long-term carbon cycle), we considered different equations for the quantification of the atmospheric CO_2 consumed by weathering (Huh, 2010). The results show the net predominance of carbonate weathering on fixing atmospheric CO_2 and that, considering the long-term carbon cycle, the amount of atmospheric CO_2 uptake by weathering is about one order of magnitude lower than considering the short-term carbon cycle. Moreover, considering the short-term carbon cycle, the mean CO_2 consumed by Alpine basins is of the same order of magnitude of the mean CO_2 consumed by weathering by the 60 largest rivers of the world estimated by Gaillardet et al. (1999).

References

- Amiotte-Suchet, P. "Cycle Du Carbone, Érosion Chimique Des Continents Et Transfert Vers Les Océans." *Sci. Géol. Mém. Strasbourg* 97 (1995): 156.
- Amiotte-Suchet, P., and J.-L. Probst. "Origins of dissolved inorganic carbon in the Garonne river waters: seasonal and interannual variations." *Sci. Géologiques Bull. Strasbourg* 49, no. 1-4 (1996): 101-126.
- Berner, E.K., and R.A. Berner. *The Global Water Cycle. Geochemistry and Environment*. Prentice Hall. Englewood Cliffs, NJ, 1987.
- Drever, J.L. *The Geochemistry of Natural Waters*. Prentice Hall, 1982.
- Gaillardet, J., B. Dupré, P. Louvat, and C.J. Allègre. "Global Silicate Weathering and CO₂ Consumption Rates Deduced from the Chemistry of Large Rivers." *Chemical Geology* 159 (1999): 3-30.
- Garrels, R.M., and F.T. Mackenzie. *Evolution of Sedimentary Rocks*. New York: W.W. Norton and, 1971.
- Huh, Y. "Estimation of Atmospheric CO₂ Uptake by Silicate Weathering in the Himalayas and the Tibetan Plateau: a Review of Existing Fluvial Geochemical Data." In *Monsoon Evolution and Tectonics-Climate Linkage in Asia*, 129-151. Geological Society of London, Special Publications. 342. London, 2010.
- Meybeck, M. "Composition Chimique Naturelle Des Ruisseaux Non Pollués En France." *Bulletin De La Société Géologique* 39 (1986): 3-77.
- Meybeck, M. "Global Chemical Weathering of Surficial Rocks Estimated from River Dissolved Load." *Am. J. Sci* 287 (1987): 401-428.
- Probst J.L., Mortatti J. And Tardy Y. 1994- Carbon river fluxes and global weathering CO₂ consumption in the Congo and Amazon river basins. *Applied Geochemistry*, 9, p 1-13
- Tardy, Y. *Le Cycle De l'Eau. Climats, Paléoclimates Et Géochimie Globale*. Masson. Paris, 1986.