



HAL
open science

New insights on Cu origin and fate from combined chemical extraction and Cu isotopic composition: Application to Cu transfers in a Mediterranean vineyard catchment

Désirée El Azzi, Jérôme Viers, Maritxu Guisresse, Anne Probst, Dominique Aubert, Jocelyne Caparros, François Charles, Katell Guizien, Jean-Luc Probst

► **To cite this version:**

Désirée El Azzi, Jérôme Viers, Maritxu Guisresse, Anne Probst, Dominique Aubert, et al.. New insights on Cu origin and fate from combined chemical extraction and Cu isotopic composition: Application to Cu transfers in a Mediterranean vineyard catchment. Joint European Stable Users Group Meeting 2012 (JESIUM), Sep 2012, Leipzig, Germany. pp.90-90. hal-04021322

HAL Id: hal-04021322

<https://hal.science/hal-04021322v1>

Submitted on 9 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 6627

To cite this version: El Azzi, Désirée and Viers, Jérôme and Guiresse, Maritxu and Probst, Anne and Aubert, Dominique and Caparros , Jocelyne and Charles , François and Guizien, Katell and Probst, Jean-Luc *New insights on Cu origin and fate from combined chemical extraction and ^{65}Cu isotopic composition: Application to Cu transfers in a Mediterranean vineyard catchment*. In: Joint European Stable Users Group Meeting 2012 (JESIUM), 2-7 September 2012, Leipzig - Germany. (Unpublished)

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

New insights on Cu origin and fate from combined chemical extraction and $\delta^{65}\text{Cu}$ isotopic composition: Application to Cu transfers in a Mediterranean vineyard catchment

D. EL AZZI^{1,2}, J. VIERS^{3,4}, M. GUIRESSE^{1,2}, A. PROBST^{1,2}, D. AUBERT^{5,6}, J. CAPARROS^{7,8}, F. CHARLES^{7,8}, K. GUIZIE^{7,8} & J. L. PROBST^{1,2,*}

¹Université de Toulouse, INPT, UPS, Laboratoire Ecologie Fonctionnelle et Environnement (ECOLAB), ENSAT, Castanet Tolosan, France, jean-luc.probst@ensat.fr

²CNRS, ECOLAB; ENSAT, Castanet Tolosan, France

³Université de Toulouse; UPS; Géosciences Environnement Toulouse (GET), Toulouse, France

⁴CNRS, IRD, CNES; GET, Toulouse, France

⁵Université de Perpignan Via Domitia, Centre de Formation et de Recherche sur les Environnements Méditerranéens (CEFREM), Perpignan, France

⁶CNRS, CEFREM, UMR 5110, Perpignan, France.

⁷CNRS, FRE 3350, LECOB, Observatoire Océanologique, Banyuls/mer, France

⁸UPMC Université Paris 6, FRE 3350, LECOB, Observatoire Océanologique, Banyuls/mer, France

Repeated use of Cu based fungicides (Bordeaux mixture: $\text{Ca}(\text{OH})_2 + \text{CuSO}_4$) to control vine downy has led to significant increase of Cu in vineyard soils. In Mediterranean catchments, brief and intense flood events can multiply stream discharge by up to 10 and are responsible for important soil leaching and therefore for high fluxes of Cu exported at the outlet of the catchment. In order to assess the origin and fate of Cu measured in the Baillaury catchment (South of France), combined sequential chemical extraction (SCE) and isotopic Cu compositions ($\delta^{65}\text{Cu}$) approaches are proposed in this study. A cultivated vineyard soil (CV) and an abandoned one (AV) were sampled at different depths. River bed sediments (BS), suspended particulate matter (SPM) and river water were also collected during the flash flood event of February 2009. In the samples, Cu partitioning between residual and non residual fractions (dissolved with water, exchangeable, acido soluble, manganese oxides, amorphous and crystalline iron oxides, organic matter) was studied using a 7-steps SCE procedure. Data showed a more important residual fraction in AV soil (> 60%) than in CV soil, riverine BS and SPM (>25%). By normalizing Cu concentrations to those of the local bedrock using Fe, Sc and Ti as normalizers, enrichment factors (EF) were calculated revealing moderate to significant enrichments (EF > 2). Anthropogenic contributions in all samples were between 50 and 85 % attesting the impact of human activities on Cu content currently detected in the catchment. In non residual fractions of the samples, Cu showed the most important affinity for amorphous and crystalline Fe oxides (25-75%) and to a lesser extent for organic matter (5-10%). $\delta^{65}\text{Cu}$ of bulk samples were similar to the bedrock and detailed isotopic investigations on SCE fractions of superficial CV and AV soils and SPM were therefore powerful to discriminate Cu origin and distribution. Copper in residual fractions of soils, BS and SPM is of natural origin because it has $\delta^{65}\text{Cu}$ close to the local bedrock (0.07‰). In the surface horizon of the vineyard soil, copper has an isotopic ratio close to that ($\delta^{65}\text{Cu} = -0.34$) of Cu fungicide. ^{65}Cu is shown to have a preferential adsorption onto iron oxides. Cu transported in solution ($\delta^{65}\text{Cu} = 0.31$ ‰) seemed to be associated with the dissolved fraction ($\delta^{65}\text{Cu} = 0.26$ ‰) of SPM without the occurrence of any fractionation. Cu extracted from the organic matter fraction has negative isotopic values ($\delta^{65}\text{Cu}$: - 0.37‰ and -0.20‰ respectively in the cultivated soil and in the river suspended matters) showing that Cu from fungicide is bound to organic matter without or with slight isotopic fractionation. As a conclusion, combining a chemical sequential extraction with the isotopic measurements is a powerful tool to get a clearer insight of Cu distribution and mobility in the environment.