



HAL
open science

Pore-scale modelling of biofilm activity in the underground storage of hydrogen

Anozie Ebigbo, Fabrice Golfier, Michel Quintard

► To cite this version:

Anozie Ebigbo, Fabrice Golfier, Michel Quintard. Pore-scale modelling of biofilm activity in the underground storage of hydrogen. *Interpore 2012*, May 2012, West Lafayette, United States. pp.0. <hal-04023499>

HAL Id: hal-04023499

<https://hal.science/hal-04023499v1>

Submitted on 10 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 8801

To cite this version :

Ebigbo, Anozie and Golfier, Fabrice and Quintard, Michel *Pore-scale modelling of biofilm activity in the underground storage of hydrogen*. (2012) In: Interpore 2012, 14-16 May, Purdue University, West Lafayette, USA. (Unpublished)

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes.diff.inp-toulouse.fr

Pore-scale modelling of biofilm activity in the underground storage of hydrogen.

A. Ebigbo¹, F. Golfier¹, and M. Quintard^{2,3}

¹*Laboratoire Environnement, Géomécanique et Ouvrages (LAEGO), Université de Lorraine, Nancy, France.*

²*Université de Toulouse; INPT, UPS; IMFT (Institut de Mécanique des Fluides de Toulouse), France*

³*CNRS; IMFT; Toulouse, France*

ABSTRACT: The storage of hydrogen in the subsurface to compensate fluctuations in energy demand and supply is considered an important part of future energy strategies. It has been observed that, within the period of storage, there is a partial conversion of hydrogen in the presence of carbon dioxide to methane. This has been attributed to the activity of micro-organisms (archaea and bacteria) indigenous to the storage site.

The talk will look at pore-scale phenomena including the interplay of different microbes (methanogens, acetogens, and acetotrophs) within a biofilm at the gas-water interface, the growth and decay of the multi-species biofilm, and the diffusion, consumption, and production of the dissolved gases. A numerical model with interface tracking, based on a volume-of-fluid method, is proposed for investigating these effects.

The aim of the study is the description and quantification of the dominant processes which determine the amount of biomass such a reservoir can support and the rate at which the micro-organisms produce methane as a contribution to explaining the observed field-scale phenomenon.