



HAL
open science

Electron dissociative attachment to ArH^+ , HD^+ , N_2^+ and CO_2

A Abdoulanziz, E Djuissi, C Argentin, Y Moulane, Arnaud Bultel, J Zs Mezei, J Tennyson, K Chakrabarti, V Laporta, I Schneider

► To cite this version:

A Abdoulanziz, E Djuissi, C Argentin, Y Moulane, Arnaud Bultel, et al.. Electron dissociative attachment to ArH^+ , HD^+ , N_2^+ and CO_2 . *Journal of Physics: Conference Series*, 2020, 31st International Conference on Photonic, Electronic and Atomic Collisions (ICPEAC XXXI) 23-30 July 2019, Deauville, France, 1412 (17), pp.172005. 10.1088/1742-6596/1412/17/172005 . hal-04024922

HAL Id: hal-04024922

<https://hal.science/hal-04024922>

Submitted on 9 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PAPER • OPEN ACCESS

Electron dissociative attachment to ArH^+ , HD^+ , N_2^+ and CO_2

To cite this article: A Abdoulanziz *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 172005

View the [article online](#) for updates and enhancements.

You may also like

- [The Cosmic-Ray Ionization Rate in the Galactic Disk, as Determined from Observations of Molecular Ions](#)
David A. Neufeld and Mark G. Wolfire
- [NEW ACCURATE MEASUREMENT OF \$^{36}\text{ArH}^+\$ AND \$^{38}\text{ArH}^+\$ RO-VIBRATIONAL TRANSITIONS BY HIGH RESOLUTION IR ABSORPTION SPECTROSCOPY](#)
M. Cueto, J. Cernicharo, M. J. Barlow et al.
- [Exploring the Possibility of Identifying Hydride and Hydroxyl Cations of Noble Gas Species in the Crab Nebula Filament](#)
Ankan Das, Milan Sil, Bratati Bhat et al.

PRIME
PACIFIC RIM MEETING
ON ELECTROCHEMICAL
AND SOLID STATE SCIENCE

HONOLULU, HI
Oct 6-11, 2024

Abstract submission deadline:
April 12, 2024

Learn more and submit!

Joint Meeting of
The Electrochemical Society
•
The Electrochemical Society of Japan
•
Korea Electrochemical Society

Electron dissociative attachment to ArH^+ , HD^+ , N_2^+ and CO_2

A Abdoulanziz^{1*}, E Djuissi¹, C Argentin¹, Y Moulane², A Bultel³, J Zs Mezei⁴,
J Tennyson⁵, K Chakrabarti⁶, V Laporta^{1,7} and I Schneider^{1,8†}

¹LOMC UMR CNRS 6294, Université du Havre, Le Havre, 76058, France

²Space Sciences, Technologies & Astrophysics Research Institute, University of Liege, Liege, Belgium

³CORIA, UMR CNRS 6614, Université de Rouen, Saint Etienne du Rouvray, 76801, France

⁴Institute for Nuclear Research, Hungarian Academy of Sciences, Debrecen, H-4001, Hungary

⁵Dept. of Physics and Astronomy, University College London, London WC1E 6BT, UK




⁶Dept. Of Mathematics, Scottish Church College 1&3 Urquhart Sq., Kolkata 700 006, India

⁷P.Las.M.I. lab, Nanotec, CNR, 70126 Bari, Italy

⁸Laboratoire Aimé-Cotton CNRS-UPR-3321, Université Paris-Sud, 91405 Orsay, France

Synopsis We will present our recent results in the study of electron-induced reactivity of ionized and neutral molecules. The Configuration Interaction method and the Multichannel Quantum Defect Theory have been employed in order to obtain cross sections for collisions of electrons with ArH^+ , HD^+ , N_2 and CO_2 .


Using the Multichannel Quantum Defect Theory (MQDT) [1-6], we computed the cross sections and the rate coefficients for the dissociative recombination (DR) (1.1) and the related competitive processes - vibrational excitation or de-excitation (VE or VdE) (1.2) ($v_f^+ > v_i^+$ or $v_f^+ < v_i^+$) and dissociative excitation (DE) (1.3):



AB standing for ArH [7], HD and N_2 [6]. Here v_i^+ and v_f^+ stand for the initial and final vibrational quantum number of the target ion.

These processes occur in the interstellar media, in (re)entry, fusion edge and laboratory plasmas.

The configuration interaction method [8] has been alternatively used for exploring the dissociative electron attachment (1.4) and dissociative excitation (1.5) of CO_2 whose removal is crucial for depollution:



The study of the reactional dynamics relies on the calculation of the relevant potential energy curves and of the corresponding mutual interactions. In these calculations one bond of the target (OC-O) has been stretched only, in order to obtain CO in its electronic ground state.

* E-mail: abdillah.abdoulanziz@univ-lehavre.fr

† E-mail: ioan.schneider@univ-lehavre.fr




Figure 1. Reactive collisions of HD^+ in its ground vibrational state with electrons.

References

- [1] Giusti A 1980 *J. Phys. B* **13** 3867
- [2] Guberman S L *et al* 1991 *The Journal of Chemical Physics* **95** 2602
- [3] Chakrabarti K *et al* 2013 *Phys. Rev. A*, **87** 022702
- [4] Motapon O *et al* 2014 *Phys. Rev. A* **90** 012706
- [5] Epée Epée M *et al* 2015 *MNRAS* **455** 276
- [6] Little D A *et al* 2014, *J. Phys. B* **47** 105204
- [7] Abdoulanziz A *et al* 2018 *MNRAS* **479** 2415
- [8] Bardsley J N 1968 *J. Phys. B: At. Mol. Phys.* **1** 349