



Distinct roles of forward and backward alpha-band waves in spatial visual attention

Andrea Alamia, Lucie Terral, Malo Renaud d'Ambra, Rufin Vanrullen

► To cite this version:

Andrea Alamia, Lucie Terral, Malo Renaud d'Ambra, Rufin Vanrullen. Distinct roles of forward and backward alpha-band waves in spatial visual attention. *eLife*, 2023, 12, 10.7554/eLife.85035 . hal-04025214

HAL Id: hal-04025214

<https://hal.science/hal-04025214>

Submitted on 12 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Distinct roles of forward and backward alpha-band waves**
2 **in spatial visual attention**

3 Andrea Alamia^{1,2}, Lucie Terral¹, Malo Renaud D’Ambra¹ and Rufin VanRullen^{1,2}

4 ¹ Cerco, CNRS Université de Toulouse, Toulouse 31052 (France)

5 ²Artificial and Natural Intelligence Toulouse Institute (ANITI), Toulouse (France)

6 andrea.alamia@cnrs.fr

7 Running title: *“Alpha-band traveling waves in visual attention”*

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33 **Abstract (237/250)**

34 Previous research has associated alpha-band [8-12Hz] oscillations with inhibitory functions:
35 for instance, several studies showed that visual attention increases alpha-band power in the
36 hemisphere ipsilateral to the attended location. However, other studies demonstrated that
37 alpha oscillations positively correlate with visual perception, hinting at different processes
38 underlying their dynamics. Here, using an approach based on traveling waves, we
39 demonstrate that there are two functionally distinct alpha-band oscillations propagating in
40 different directions. We analyzed EEG recordings from three datasets of human participants
41 performing a covert visual attention task (one new dataset with N=16, two previously
42 published datasets with N=16 and N=31). Participants were instructed to detect a brief target
43 by covertly attending to the screen's left or right side. Our analysis reveals two distinct
44 processes: allocating attention to one hemifield increases top-down alpha-band waves
45 propagating from frontal to occipital regions ipsilateral to the attended location, both with or
46 without visual stimulation. These top-down oscillatory waves correlate positively with alpha-
47 band power in frontal and occipital regions. Yet, different alpha-band waves propagate from
48 occipital to frontal regions and contralateral to the attended location. Crucially, these forward
49 waves were present only during visual stimulation, suggesting a separate mechanism related
50 to visual processing. Together, these results reveal two distinct processes reflected by
51 different propagation directions, demonstrating the importance of considering oscillations as
52 traveling waves when characterizing their functional role.

53 **Keywords:** Traveling waves, covert visual attention, alpha-band oscillations, EEG recordings.

54

55

56

57

58

59

60

61

62

63 **Introduction**

64 Brain oscillations are related to several cognitive functions, as they orchestrate neuronal
65 activity at distinct temporal and spatial scales (Buzsáki and Draguhn, 2004; Buzsáki, 2009).
66 Alpha-band oscillations [8-12Hz] are the most prevailing rhythms in electrophysiological
67 (EEG) recordings, spreading through most cortical regions.

68 Several studies investigated their functional role in various cognitive processes (Palva and
69 Palva, 2007, 2011), providing mixed results. On the one hand, some studies showed that
70 alpha-band oscillations might filter sensory information, regulating excitation and inhibition
71 of sensory-specific brain regions (Jensen and Mazaheri, 2010; Mathewson et al., 2011;
72 Klimesch, 2012; Sadaghiani and Kleinschmidt, 2016). Accordingly, researchers interpreted
73 alpha oscillations as a top-down mechanism involved in inhibitory control and timing of
74 cortical processing (Klimesch et al., 2007), as well as modulating cortical excitability (Jensen
75 and Mazaheri, 2010; Mathewson et al., 2011). Experimental studies corroborated this
76 hypothesis, demonstrating how the phase of alpha-band oscillation affects visual perception
77 (Busch et al., 2009; Fakche et al., 2022; but see also Ruzzoli et al., 2019). Another highly
78 replicated result regarding the inhibitory role of alpha oscillations consists in the hemispheric
79 modulation in occipital regions associated with visual attention, having an increase of power
80 ipsilateral to the attended hemifield, and a corresponding decrease contralaterally (Worden et
81 al., 2000; Sauseng et al., 2005; Kelly et al., 2006; Thut et al., 2006; Händel et al., 2011). On
82 the other hand, other experimental studies have related alpha-band oscillations in occipital and
83 parietal regions to perceptual processing and visual memory (Bonnefond and Jensen, 2012;
84 VanRullen, 2016; Pang et al., 2020; Luo et al., 2021). For example, reverse-correlation
85 techniques reveal that the visual system reverberates sensory information in the alpha-band
86 for as long as one second, in what has been dubbed ‘perceptual echoes’ (Vanrullen and
87 MacDonald, 2012). Importantly, these echoes are a clear signature of sensory processing as
88 they reflect the input’s precise time course, are modulated by attention and have been
89 dissociated from inhibitory alpha power modulation (Vanrullen and MacDonald, 2012;
90 VanRullen, 2016; Brüers and VanRullen, 2018; Schwenk et al., 2020).

91 Altogether, these experimental evidences support distinct and contradictory conclusions about
92 alpha-band oscillation’s functional role(s), which remains an open debate. Here, we address
93 this question from a different perspective that interprets alpha-band oscillations as traveling
94 waves (Muller et al., 2018; Alamia and VanRullen, 2019), thus considering their spatial

component, and their propagation direction. Considering the case of visual attention, we tested the hypothesis that two functionally distinct alpha-band oscillations propagate along the frontal-occipital line in opposite directions. This compelling hypothesis about the different functional roles of alpha-band traveling waves derives from our previous studies (Alamia and VanRullen, 2019; Pang et al., 2020), in which we showed how visual perception modulates alpha waves, i.e. forward waves during visual stimulation, backward waves when the stimulus was off. In addition, this hypothesis is in line with previous studies suggesting that distinct alpha-band oscillations are related to specific cognitive processes (Gulbinaite et al., 2017; Deng et al., 2019; Schuhmann et al., 2019; Sokoliuk et al., 2019; Kasten et al., 2020). In this study, we analyzed three datasets, two publicly available (Foster et al. 2017 and Feldmann-Wustefeld et al. 2019, see below), and one collected specifically for this study. In all dataset, participants attended either to the left or the right hemifield, while keeping central fixation. Our results confirmed the hemispheric modulation of alpha-band oscillations in posterior regions (Worden et al., 2000; Sauseng et al., 2005; Kelly et al., 2006; Thut et al., 2006; Händel et al., 2011) and revealed two distinct alpha-band traveling waves propagating in opposite directions. First, visual attention increases top-down alpha-band waves propagating from frontal to occipital regions ipsilateral to the attended location, and such waves correlate positively with alpha power in frontal and occipital regions. Moreover, our analysis demonstrates that visual attention also modulates contralateral forward waves, i.e., waves propagating from occipital to frontal areas. Importantly, the attentional modulation of forward waves is crucially dependent on sustained sensory processing, as this modulation disappears in the absence of visual stimulation. In contrast, alpha-band top-down waves are present and modulated by visual attention irrespective of the presence or absence of concurrent sensory stimulation. These results demonstrate two distinct alpha-band oscillatory waves propagating in opposite directions, seemingly underlying different cognitive processes. The well-known lateralization effect observed in alpha-band can be interpreted as top-down traveling waves, and it's most likely related to inhibitory processes, in line with previous studies (Jensen and Mazaheri, 2010; Händel et al., 2011). However, different alpha-band oscillations propagate in a forward direction and are directly related to sensory processing, reconciling previous evidence linking alpha-band oscillations with visual processing (Vanrullen and MacDonald, 2012; Lozano-Soldevilla and VanRullen, 2019).

Results

Travelling waves' spectral profile. The goal of the study was to investigate how visual attention modulates alpha-band travelling waves in the hemisphere contra- and ipsilateral to the attended location. To test this, we considered 11 lines of electrodes running from occipital to frontal regions (figure 1C), 5 for each hemisphere and one midline. It is important to note that the spatial resolution of these lines is not critical for our analysis, as we do not expect significant differences within each hemisphere. However, before testing how visual attention modulates travelling waves, we explored the amount of waves propagating forward (FW) and backward (BW) as a function of their temporal frequency (see figure 5 and methods for a detailed description of the analysis). Figure 1D shows the spectral profile of FW and BW waves in the midline (along the Oz-Fz axis) and the contra- and ipsilateral lines: confirming previous experimental studies (Alamia and VanRullen, 2019; Pang et al., 2020), we found that alpha-band oscillatory waves propagate in both directions during visual stimulation, whereas theta (4-7Hz) and high-beta/gamma (24-45Hz) bands propagate mostly bottom-up from occipital to frontal regions, and low-beta (13-23Hz) waves flow in the top-down direction. Interestingly, this pattern of results confirms previous studies using different methods, in which higher frequency bands (i.e., high-beta/gamma) have been associated with forward processing, whereas low-beta and alpha frequencies have been related to top-down processing (Bastos et al., 2012, 2015; van Kerkoerle et al., 2014; Michalareas et al., 2016; but see also Schneider et al., 2021).




Figure 1 – Experimental design and waves’ spectral profile. A) Each trial lasted 5 seconds, in which two flickering stimuli were presented to both hemifield. Participants were instructed to attend either the left or the right hemifield, as indicated by a central cue. In some trials, a target or a distractor appeared for 100ms as a square either in the attended or unattended location. B) The target and distractor luminance changed over trials due to the QUEST algorithm, which kept participants’ performance around 80%. C) We quantified travelling waves along 11 electrodes lines, running along the anterior-posterior axis. These lines were located in the contralateral or the ipsilateral hemisphere to the attended location. D) The amount of waves in dB computed for forward (in blue) and backward (in red) waves in the midline (central subplots, thinner lines represent standard errors of the mean) and in the ipsi- and contralateral hemisphere (left and right panels, respectively). These waves were computed on trials without target or distractors. Positive (negative) values reflect more (less) waves than the chance level (as quantified by the surrogate distribution), whereas values around 0 indicate no difference between the real and the null distribution. E) Simulated data providing a schematic representation of forward and backward waves in the time domain in a given line of electrodes (from more frontal E1 to more occipital E7). A positive and a negative phase shift characterized forward and backward waves, respectively.

Attending to visual stimuli modulates traveling waves. In this analysis, we investigated how covert visual attention influences the traveling wave pattern. We focused on trials where neither a target nor a distractor was presented. First, we quantified the amount of traveling waves in the contra- and ipsi- lateral hemispheres to the attentional allocation. As shown in figure 2 (left column), we found a strong lateralization effect revealing an increase

170 (respectively, decrease) of contralateral (ipsilateral) forward waves in the alpha band, and the
 171 opposite pattern in waves propagating in the opposite direction. These results were confirmed
 172 by a Bayesian ANOVA, considering as factors DIRECTION (FW or BW), LINES (distance
 173 from the midline), and LATERALITY (contra vs ipsi). The results revealed strong evidence in
 174 favor of the interaction between DIRECTION and LATERALITY factors ($BF_{10}=31.230$,
 175 estimated error~1%, $\eta^2=0.08$ as estimated from a classical ANOVA), whereas all other
 176 factors and their interactions revealed evidence in favor of the absence of an effect
 177 ($BF_{s10}<0.3$). We also found no significant effect in the other frequency bands (as shown in
 178 figure 1D, namely theta, low and high beta), hence we focused the following analyses on
 179 alpha-band oscillatory waves. These results demonstrate that the direction of alpha-band
 180 oscillatory traveling waves shows a laterality effect during a task involving covert selective
 181 attention.




Figure 2 – Traveling waves block analysis. Each column in the figure represents a different EEG dataset involving experiments with visual stimulation (left and middle column) and without visual stimulation (right column). In the upper panels, the net amount of forward (blue) and backward (red) waves is represented along different lines of electrodes, normalized to the midline. The left and central panels reveal an increase (decrease) of forward (backward) waves contralateral to the attended location when participants attended to visual stimulation. The right column shows that when participants attended an empty screen (data from Foster et al. 2017), only backward waves were modulated by visual attention, and no effect was observed in the forward waves without visual stimulation. Error bars represent standard errors of the mean. The middle row shows schematic representations of the screen during the tasks: the central panel illustrates the task from Feldmann-Wustefeld et al. 2019, where D and T stand for Distractor and Target, respectively. In the task from Foster et al. 2017, the screen was empty, as the eight circles were not displayed during the task but here illustrate the stimulus positions (Foster et al., 2017) (Foster et al., 2017). The lower panels represent the lines of electrodes in all datasets.

Backward waves correlate with alpha-band power. Previous studies investigating the role of alpha-band oscillations in visual attention reported a lateralization effect in the spectral power of alpha-band oscillations (Worden et al., 2000; Sauseng et al., 2005; Kelly et al., 2006; Thut et al., 2006; Jensen and Mazaheri, 2010; Händel et al., 2011). One may then wonder about the relationship between alpha power and traveling waves. To address this question, we investigated whether the oscillatory activity we observe propagating through the cortex relates to the spectral power in either occipital or frontal regions. We computed the averaged alpha-band power in frontal and occipital areas, contra- and ipsi- laterally to the target presentation, considering the same electrodes used for quantifying the traveling waves (see figure 3A). Interestingly, we found a significant positive correlation between alpha-band power in both occipital and frontal regions and backward waves, but not with forward waves (figure 3A – and Table 1). Next, we considered the lateralization effect in the alpha-band, as shown in figure 3B (topographic plot in the right panel) and well-replicated in previous studies (Sauseng et al., 2005; Thut et al., 2006; Händel et al., 2011). We wondered whether we could observe a correlation between such lateralization, defined as the difference between alpha-band power when attention is allocated to one side of the screen and to the other side, and the effect we reported in the travelling waves (figure 2). As shown in figure 3B (left panels), our results demonstrate a lack of correlation for both backward and forward waves in both frontal and occipital regions (all $|r| < 0.1$, $BF_{10} \sim 0.3$).

Pearson r (BF_{10})		FW		BW	
		CONTRA	IPSI	CONTRA	IPSI
○	CONTRA	-0.297 (0.549)	-0.350 (0.697)	0.720 (28.519)	0.698 (19.503)

	IPSI	-0.305 (0.566)	-0.342 (0.669)	0.786 (116.990)	0.746 (47.512)
FRONT	CONTRA	-0.222 (0.422)	-0.252 (0.465)	0.772 (84.225)	0.712 (24.645)
	IPSI	-0.327 (0.625)	-0.354 (0.710)	0.747 (48.448)	0.705 (21.841)

Table 1 – Correlation with alpha-band power. The table reports the Pearson correlation coefficient and the Bayes Factor (BF_{10} supporting the alternative hypothesis, that is the presence of a correlation) between frontal and occipital electrodes and forward (FW) and backward (BW) waves, in both contralateral and ipsilateral hemisphere. All correlations were computed on trials when neither a target nor a distractor was displayed.

To further investigate the relation between alpha-band travelling waves and alpha power, we performed the same analysis focusing on the correlation within each participant. In particular, we correlated trial-by-trial forward and backward waves with alpha-band power for each subject, obtaining correlation coefficients ‘r’ and their respective p-values. As in the previous analysis, we correlated forward and backward waves with frontal and occipital electrodes in both contro- and ipsilateral hemispheres. We applied the Fisher method (Fisher, 1992, see Methods for details) to combine all subjects' p-values in every conditions. Overall, we found a significant effect of all combined p-values ($p < 0.0001$), except in the lateralization condition (contra- minus ipsilateral hemisphere), similar to our previous analysis. Additionally, we tested for a consistent positive or negative distribution of the correlation coefficients. As shown in figure 3C, the results support a significant correlation between backward waves and alpha-power in the hemisphere contralateral to the attended location ($BF_{10}=10.7$ and $BF_{10}=7.4$ for occipital and frontal regions, respectively; all other BF_{10} were between 1 and 2, providing inconclusive evidence). Interestingly, this analysis also revealed a small but consistent effect in the correlation between lateralization effects, as we reported a consistently positive correlation in the contra- minus ipsilateral difference between forward waves and alpha power ($BF_{10} \sim 5$ for both frontal and occipital electrodes). However, it's important to notice that the combined p-values obtained using the Fisher method did not reach the significance threshold in the lateralization condition, reducing the relevance of this specific result.




Figure 3 - Correlation with alpha-band power. A) Panel A reveals a correlation between backward waves and alpha power (static, standing power, i.e. measured via wavelets transform), in both frontal and occipital areas, in both hemispheres. We did not observe such correlation with forward waves. The plot to the right reveals the topographic distribution of alpha-power when participants attended to the right hemifield (we included the ‘left’ condition by flipping the electrodes symmetrically to the midline). The white dots indicate the electrodes used for the correlation. B) The plots to the left show the correlation between the laterality effect in the alpha power and in the waves (laterality measured as the mean difference between contra and ipsilateral hemispheres for both alpha power and the waves –

for the waves we computed the difference using lines of electrodes symmetrical to the (midline). We did not observe any correlation in neither forward or backward waves, with neither frontal or occipital alpha power. The topography to the right reveal a lateralization effect in the alpha power (attention to the left minus attention to the right), confirming the presence of alpha power lateralization, in line with previous studies (Sauseng et al., 2005; Thut et al., 2006; Händel et al., 2011). C) Panel C shows the trial-by-trial correlation coefficients averaged across participants for different conditions (as indicated in the x-axis). Confirming the results in panel A, we found a positive correlation across participants between backward waves and alpha-power, specifically in the contralateral hemisphere. We also observed a positive global effect of the laterality condition across participants in the forward waves, even though the combined p-values for the trial-by-trial correlation did not reach the significant threshold.

Covert attention modulates forward waves only with visual stimuli. To confirm our previous results, we replicated the same traveling waves analysis on two publicly available EEG datasets in which participants performed similar attentional tasks (experiment 1 of Foster et al., 2017 and experiment 1 of Feldmann-Wüstefeld and Vogel, 2019). In the first experiment from the Feldmann-Wüstefeld and Vogel dataset, participants were instructed to perform a visual working memory task in which, while keeping a central fixation, they had to memorize a set of items while ignoring a group of distracting stimuli. We focused our analysis on those trials in which the visual items to remember were placed either to the right or the left side of the screen, while the distractors were either in the upper or lower part of the screen (we pulled together the trials with either 2 or 4 distractors, as this factor was irrelevant for our analysis). The stimuli were shown for 200ms, and we computed the amount of forward and backward waves in the 500ms following stimulus onset. As shown in figure 2 (central column), the analysis confirmed our previous results, demonstrating a strong interaction between the factors DIRECTION and LATERALITY ($BF_{10}=667$, error~2%; independently, the factors DIRECTION and LATERALITY had $BF_{10}=0.2$ and $BF_{10}=0.4$, respectively). These results confirmed that spatial attention modulates both forward and backward waves in the presence of visual stimulation. Next, we analyzed another publicly available dataset from Foster et al., 2017. In the first experiment of Foster's study, participants completed a spatial cueing task, requiring them to identify a digit among distractor letters. After a central cue was displayed for 250ms, participants attended one of eight locations for 1000ms before the onset of the target and distractors. As in our design, participants allocated attention to different locations to the left or right of the screen while keeping central fixation. However, unlike in our and in Feldmann-Wüstefeld's study, no stimulus was displayed while participants were attending one of the possible locations. Here, we assessed the amount of waves in the 1000ms before the onset of the stimulus during attention allocation, when no visual stimuli were

shown on the screen. Remarkably, as shown in figure 2 (right panel), our analysis demonstrated an effect of the lateralization (LATERALITY: $BF_{10}=3.571$, error~1%), revealing more waves contralateral to the attended location, but inconclusive results regarding the interaction between DIRECTION and LATERALITY ($BF_{10}=2.056$, error~1%). However, using a classical ANOVA (i.e., without modeling the slope of the random terms), the interaction between DIRECTION and LATERALITY proved significant ($F(1,16)=9.81$, $p=0.003$, $\eta^2=0.13$). In addition, when testing LATERALITY separately for forward and backward waves, we observed an effect in the backward waves ($BF_{10}=3.497$, error<0.01%) but not in the forward waves ($BF_{10}=0.231$, error<0.01%, supporting evidence in favor of the absence of an effect). In addition, as analyzed in our dataset, we tested the correlation between backward waves and alpha-band power in occipital (electrodes: PO3, PO4) and frontal (electrodes: F3, F4) regions. We found moderate evidence of a positive correlation between contra- and ipsi- lateral backward waves, and occipital (all Pearson's $r\sim 0.4$, all $BF_{S_{10}}\sim 3$) but inconclusive evidence in the frontal areas (all Pearson's $r\sim 0.3$, all $BF_{S_{10}}\sim 2$). These results supported those from our dataset, despite having a smaller amount of electrodes' lines, and potentially reduced statistical power (see figure 2, lower panels). All in all, we could confirm our previous conclusion that covert visual attention modulates top-down oscillatory waves, showing this effect even in the absence of visual stimulation. In addition, we surmised that the lateralization effect we reported in the forward waves in our dataset (absent in the Foster dataset) is related to the steady visual stimulation during the attentional allocation, in line with our previous results demonstrating that oscillatory bottom-up waves reflect sensory processing (Alamia and VanRullen, 2019; Pang et al., 2020).

Both detected targets and distractors elicit FW waves, but not missed targets. In our previous analysis, based on a subset of trials in which neither a target nor a distractor occurred, we demonstrated that sustained attention to one hemifield generates oscillatory alpha-band waves propagating forward in the contralateral hemisphere and backward in the ipsilateral one. We now assess whether the occurrence of a specific event, such as the onset of a target or a distractor stimulus, could induce the generation of transient oscillatory waves. For this reason, we replicated the same analysis on those trials including either a target or a distractor (on average, each participant performed 146.25 trials in each condition), to quantify the amount of waves locked to the onset of these events.




Figure 4 – Event analysis. A) The figure shows the amount of forward (in blue) and backward (in red) contralateral waves around the onset of the target/distractor (left) or hit and missed targets (right panel). Error bars are standard error of the mean. We found an interaction effect when we analyzed the hit versus missed target. B) The 2D maps represent the amount of waves in the 11 lines of electrodes (x-axis) and around the onset time (y-axis) for forward and backward waves, and for hits and missed targets separately. The opposite pattern for hits vs. misses, already visible before the target onset, suggests that missed targets are due to a failure of attentional allocation rather than sensory processing; and consequently, that proper attentional allocation is characterized by contralateral forward waves and ipsilateral backward waves.

The upper panels of figure 4A reveal the amount of forward and backward waves contralateral to the stimulus. Note that the targets and distractors appeared in the attended and unattended locations, respectively. A Bayesian ANOVA reveals no difference between targets and distractors (EVENT: $BF_{10}=0.206$, error~1%), or their interaction (DIRECTION x EVENT: $BF_{10}=0.423$, error~5%), as shown in the top-right panel of figure 4. This result reveals that both target and distractor elicit forward waves propagating contralateral to the hemifield where they occur. Next, we investigated whether the waves in the hemisphere contralateral to the attended hemifield correlate with the participant's performance in detecting the target (a QUEST algorithm kept the accuracy throughout the experiment around 80%). Remarkably,

we found an effect concerning the ‘hit’ and ‘miss’ target, as revealed by a significant interaction of the DIRECTION and EVENT factors (DIRECTION x EVENT: $BF_{10}=4.085$, error~2%), as shown in the bottom-right panel of figure 4A. Interestingly, figure 4B reveals the amount of waves 400ms before and after the onset of the stimulus, showing how a missed target is related to a decrease (increase) in forward (backward) waves contralateral (ipsilateral) to the attended location, possibly due to attentional fluctuations during each trial.

Discussion

Previous studies demonstrated that selective attention modulates alpha-band oscillations in occipital and parietal regions (Worden et al., 2000; Sauseng et al., 2005; Kelly et al., 2006; Thut et al., 2006; Händel et al., 2011), supposedly indicating their involvement in top-down, inhibitory functions. Here, we took a novel perspective on these results by interpreting oscillations as traveling waves (Muller et al., 2018), thus considering their spatial component on top of the temporal one. Our results revealed two distinct alpha-band waves propagating in opposite directions: attention modulates oscillations traveling from occipital to frontal regions only in the presence of visual stimulation, thus relating forward waves to visual processing (Lozano-Soldevilla and VanRullen, 2019; Pang et al., 2020); whereas oscillations propagating in the opposite, top-down direction were modulated by attention irrespective of the presence or absence of concurrent visual stimulation; as in standard studies of alpha power lateralization (Worden et al., 2000; Sauseng et al., 2005; Kelly et al., 2006; Thut et al., 2006; Händel et al., 2011), this attentional modulation involved both an decrease of alpha waves contralateral to the attended location, and an ipsilateral increase.

In line with previous studies (Gulbinaite et al., 2017; Deng et al., 2019; Schuhmann et al., 2019; Sokoliuk et al., 2019; Kasten et al., 2020), our results support the thesis that distinct alpha-band oscillations are involved in separate cognitive processes. A recent study from Sokoliuk and colleagues (Sokoliuk et al., 2019) demonstrated two different sources of alpha-band oscillations during a multisensory task: one, located in visual areas, reflects the “spotlight of attention” and decreases linearly with increasing attention, whereas another one indicates attentional efforts and occurs in parietal regions. Gulbinaite and colleagues also demonstrated that parietal, but not occipital alpha-band oscillations are responsible for the oscillatory reverberation causing the ‘triple-flash’ illusion (Gulbinaite et al., 2017). Similarly, another study (Kasten et al., 2020) disentangled two primary sources of alpha oscillations, revealing a differential effect of tACS stimulation on endogenous but not exogenous attention. The authors interpreted their results as evidence supporting the hypothesis that alpha-band

oscillations play a causal role in top-down but not bottom-up attention (Schuhmann et al., 2019; Kasten et al., 2020). Our results are consistent with these findings, including the spatial dimension in analyzing and interpreting alpha-band oscillations. Additionally, we also found a significant correlation between backward waves and occipital and frontal alpha-band power, consistently with Kasten's study (Kasten et al., 2020) and the inhibitory role of alpha-band oscillations. Our findings support the hypothesis that top-down processes, as reflected by backward waves, drive the well-documented hemispheric asymmetries reported in the literature (Händel et al., 2011; Klimesch, 2012; Waldhauser et al., 2012; Peylo et al., 2021). All in all, previous studies and our results pave the way for a more comprehensive understanding of the role of alpha-oscillations in cognition.

One may wonder whether alpha-band oscillations during attention relate to target enhancement or distractor suppression (Schneider et al., 2021a). In the first case, the desynchronization of alpha activity would favor the sensory processing in the hemisphere contralateral to the target, whereas in the second case, alpha synchronization would suppress the processing of the distractor (Kelly et al., 2006; Noonan et al., 2018; Peylo et al., 2021). Our findings don't address this question directly but provide another element to the picture, suggesting the intriguing hypothesis that target enhancement is not reflected in the alpha power decrease but rather in an increase in the contralateral alpha-band waves processing the target and propagating forward. Our results thus support the hypothesis that alpha waves are involved in both distractor suppression (via ipsilateral top-down inhibitory feedback) and target enhancement (via contralateral bottom-up alpha-band waves). Future studies will precisely characterize the anatomical pathways of the distinct alpha-band oscillations, possibly involving cortical dynamics in the ventral and dorsal streams (Capilla et al., 2014) or cortico-thalamic connections (Lopes da Silva et al., 1980; Halgren et al., 2019).

Concerning the anatomical pathway of waves' propagation, our analysis based on EEG recordings prevents us from clearly determining whether the observed waves propagate through the cortex or whether more localized dipoles generate such widespread observations at the sensor level. A previous source-analysis study on different visual-task datasets (Lozano-Soldevilla and VanRullen, 2019) leaves both possibilities open. However, recent simulations on perceptual echo data (related to bottom-up, sensory waves, Zhigalov and Jensen, 2022) suggest that two dipoles in occipital and parietal regions could be responsible for the generation of the waves propagating in the occipital-to-frontal direction. Supposing this conclusion generalizes to raw EEG data and not only perceptual echoes (Vanrullen and MacDonald, 2012), one could speculate that visual attention modulates dipoles selectively in

each hemisphere. However, one may wonder whether similar dipoles are also responsible for generating top-down waves in frontal regions or whether other mechanisms are involved in generating alpha-band backward waves.

Our previous work proposed an alternative cause for the generation of cortical waves (Alamia and VanRullen, 2019). We demonstrated that a simple multi-level hierarchical model based on Predictive Coding (PC) principles and implementing biologically plausible constraints (temporal delays between brain areas and neural time constants) gives rise to oscillatory traveling waves propagating both forward and backward. This model is also consistent with the 2-dipoles hypothesis (Zhigalov and Jensen, 2022), considering the interaction between the parietal and occipital areas (i.e., a model of 2 hierarchical levels). However, dipoles in parietal regions are unlikely to explain the observed pattern of top-down waves, suggesting that more frontal areas may be involved in generating the feedback. This hypothesis is in line with the PC framework, in which top-down connections have an inhibitory function, suppressing the activity predicted by higher-level regions (Huang and Rao, 2011). Interestingly, Spratling proposed a simple reformulation of the terms in the PC equations that could describe it as a model of biased competition in visual attention, thus corroborating the interpretation of our finding within the PC framework (Spratling, 2008, 2012).

In conclusion, our study demonstrated the existence of two functionally distinct alpha-band traveling waves propagating in opposite directions and modulated by visual attention. Top-down waves prevail in the hemisphere ipsilateral to the attended location and are related to inhibitory functions, whereas forward waves reflect ongoing visual processes in the contralateral hemisphere.

Methods

Participants. EEG data were recorded from 16 volunteers (aged 20–32 years old, four women, three left-handed). All subjects reported normal or corrected-to-normal vision, and they had no history of epileptic seizures or photosensitivity. All participants gave written informed consent before starting the experiment, following the Declaration of Helsinki. This study adheres to the guidelines for research at the “Centre de Recherche Cerveau et Cognition,” and the protocol was approved by the local ethical committee “Comité de protection des Personnes Sud Méditerranée 1” (ethics approval number N° 2019-A02641-56). Furthermore, we included EEG recordings from two additional publicly available datasets

investigating distinct scientific questions and using different analyses than our study. In the first one, 31 participants performed a visual working memory task involving spatial attention. The data were published in a previous study (Feldmann-Wüstefeld and Vogel, 2019, data available online at <https://osf.io/a65xz/>). In the second dataset, 16 participants performed a task involving covert spatial attention. These data were published in another study (Foster et al., 2017, data available online at <https://osf.io/m64ue>). The number of participants in our dataset was estimated based on a power analysis of previous studies investigating travelling waves in vision (Luo et al., 2021) and to match the number of participant in the third dataset (Foster et al., 2017). Our dataset is also available online at <https://osf.io/pn784/>.

Experimental procedure. The following describes the experimental procedure to collect the data never published before. After setting the EEG device and placing the electrodes, participants performed the task in a dim and quiet room. The experiment was composed of 10 blocks of 60 trials each. During each trial (described in figure 1A), two flickering luminance disks were displayed for 5 seconds, 9° to the left and right from the center of the screen. The flicker had a frequency of 160Hz, and the intensities were pooled from a uniform distribution. We chose to apply flickering stimulation to keep participants engaged in the task and avoid attentional captures due to sudden target/distractor onset and offset (see below). Before each block, participants were instructed to allocate attention to either the right or the left stimulus while focusing on a central arrow located at the center of the screen. The arrow pointed to the attended location and served as a visual reminder throughout the block. In half of the trials, a target or a distractor flashed 100ms inside the attended or non-attended stimulus (see figure 1A). Their onset could occur any time after the first 500ms of the trial. Both target and distractor were squares whose luminance was a percentage of the disk's luminance (i.e., when at 100%, targets/distractors were indiscernible from the disk, as they have the same luminance). A QUEST algorithm (Watson and Pelli, 1983) modulated such percentage to keep participants' performance around 80% throughout the experiment (see fig. 1B). In the other half of the trials, either the target followed the distractor's onset, or neither the target nor the distractor was presented (in sum, four possible trials occurred with the same frequency: either only a target, or only a distractor, or a target preceded by a distractor, or neither of them). Participants reported whether they saw a target only at each trial's end to prevent motor activity from confounding the EEG signals. All stimuli were generated in MATLAB using the Psychtoolbox (Brainard, 1997).

We included two additional datasets in this study. In both studies, participants performed a visual attention task while keeping their fixation in the center of the screen.

Regarding the Feldmann-Wüstefeld and Vogel, 2019 study, participants were asked to memorize the colors of two stimuli while ignoring a set of distractors stimuli. We analyzed uniquely those trials in which the visual stimuli were presented to the left or right side of the screen, while the distractors were placed above or below the fixation cross. After 500ms of the fixation cross, two colored ‘target’ stimuli were presented for 200ms. Participants were asked to memorize these stimuli, and a new ‘probe’ stimulus was shown after an additional second. Participants reported whether the probe matched the target stimuli or not. We analyzed the traveling waves in the 500ms following the target stimulus onset.

Participants performed a spatial attention task in the second dataset from Foster et al. 2017. First, the fixation cross cued participants to covertly attend one of eight possible spatial positions uniformly distributed around the center of the screen. After one second, a digit was displayed either in the cued location or in any other one. The remaining locations were filled with letters. Participants were instructed to report the displayed digit. We analyzed the waves the second before the stimuli onset when participants were attending to the locations cued to the left or right side of the screen (we discarded trials in which participants attended locations above or below the fixation cross). For additional details about both experimental procedures, we refer the reader to Foster et al., 2017 and Feldmann-Wüstefeld and Vogel, 2019.

EEG recording and Preprocessing. Throughout the experiment, we recorded EEG signals using a 64-channel active BioSemi EEG system (1024Hz sampling rate), with three additional ocular electrodes. The preprocessing consisted of down-sampling the data to 256Hz, followed by a high-pass (>1Hz) and a notch (47-53Hz) filter. Data were then average-referenced and segmented from 500ms before trial onset to its end. We performed the preprocessing in EEGLab (Delorme and Makeig, 2004). Importantly, we carefully discarded from all analyses all trials in which the EOG signals revealed eye movements.

Traveling wave analysis. As in our previous studies (Alamia et al., 2020; Pang et al., 2020), we quantified traveling waves' propagation along eleven lines of seven electrodes, running from occipital to frontal regions. As shown in figure 1C, we considered one midline (Oz, POz, Pz, CPz, Cz, FCz, Fz) and five lines spreading through the left and right hemispheres, symmetrically to the midline. The electrodes' choice overlapped and covered a large portion of each hemisphere. For each set of seven electrodes, we created 2D maps by sliding a 500ms time window over the EEG signals (having a 250ms overlap) and computing 2D-FFT representations of each map (figure 5B). Notably, the power in the lower and upper quadrants quantifies the amount of waves propagating forward (FW - from occipital to frontal electrodes) and backward (BW - from frontal to occipital), respectively (see figure 5A). Next,

we performed the same procedure after shuffling the electrodes to obtain a baseline with the same spectral power but without information about the amount of FW/BW waves (note in figure 1D that the surrogate distribution accounts for the typical $\frac{1}{f}$ power trend, as well as the alpha peak). Lastly, for each frequency in the range [2-45Hz], we extracted the maximum values in the 2D-FFT spectra in both the real (FW, BW) and the shuffled data (FW_{ss}, BW_{ss}), obtaining the waves' amount in decibel [dB] as:

$$FW \text{ waves [dB]} = 10 * \log_{10} \left(\frac{FW}{FW_{ss}} \right); \quad BW \text{ waves [dB]} = 10 * \log_{10} \left(\frac{BW}{BW_{ss}} \right).$$

This value quantifies the total waves compared to the null distribution, thus being informative when contrasted against zero. Importantly, this waves analysis is limited to the surface level, and does not directly inform about the underlying sources. It is also necessary to keep in mind issues related to long-range connections and distortions due to scalp interference (Nunez, 1974; Alexander et al., 2019).




Figure 5 – Waves analysis. A) The 2D-Fast Fourier Transform (2D-FFT) decomposes an image (e.g. a space-time representation of an EEG signal) into its spectral components. The upper part shows the decomposition of a 2D sinusoid propagating along the vertical or horizontal axis of the image. The corresponding peaks are found on the axis in the spectral domain, and their position depends on the frequency of the oscillations. The lower part of the figure show how the spectra change when the oscillations propagate with a backward- or forward- like pattern. Importantly, the spectral peaks rotate in two of the four quadrants depending on the direction, providing a reliable measure of forward or backward waves in the image. B) Schematic of the waves' quantification method. After defining time windows over each electrode line, we computed 2D Fourier transformation to quantify the amount of

forward (in blue) and backward (in red) waves. From the upper and lower quadrant of the 2D-FFT spectra, we consider the maximum value over spatial frequencies, providing a 1D spectrum of forward and backward waves in the temporal frequency domain. The same procedure after shuffling the electrodes' order provides a surrogate measure, used as a baseline. Notably, such surrogate distribution captures the 1/f trend and the alpha-band peak, accounting for these factors in the final waves' quantification.

Block analysis. This analysis quantified the amount of traveling waves across all trials when neither a target nor a distractor appeared. Participants paid attention to either one or the other side of the screen, thus defining a contralateral and an ipsilateral hemisphere (see figure 1C). First, we averaged the forward and backward waves separately (in dB, see above) for each map computed along the 11 lines of electrodes within the 5-second trial (five lines for each hemisphere and the midline). Next, we averaged the results between trials, thus obtaining five contra- and five ipsi- lateral spectra per subject for both FW and BW waves. Although our hypothesis focuses on alpha-band oscillations, we also assessed the amount of waves in other frequency bands. Accordingly, from each spectrum, we computed the average per frequency band defined as θ (4-7Hz), α (8-12Hz), low β (13-24Hz) and high β/γ (25-45Hz). Besides following the frequency band definition found in the literature, such division reflects the waves' profile observed in the midline (fig.1D). Then, we normalized each pair of symmetric lines (e.g., L1) by subtracting their mean value separately for each frequency band (i.e., $\frac{L_{i, \text{contra}} + L_{i, \text{ipsi}}}{2}$). This normalization allows to remove power differences across lines (e.g., L1, L2, etc.) and to compare the effects between hemispheres. Lastly, we tested an ANOVA considering as factors DIRECTION (either FW or BW), LINE (a value from 1 to 5 to define the distance from the midline), LATERALITY (contra vs ipsi lateral), and all their interactions. We considered SUBJECTS as the random factor in the model. All models in this study relied on Bayesian statistics (see below for details). We performed the same analysis on the EEG recordings from the second dataset (16 participants performing a similar attentional task, see Foster et al., 2017, data available online at <https://osf.io/m64ue>). However, given the available data, we were able to consider only one electrodes' line per hemisphere, using the sensors O1-PO3-P3-C3-F3 and O2-PO4-P4-C4-F4 (see figure 2, lower left panel).

Waves and power correlation. This analysis assessed the correlation between FW and BW waves computed in the block analysis, with occipital and frontal alpha power. First, we estimated the mean alpha-power in contra- and ipsilateral electrodes in both frontal and parieto-occipital regions, using the same electrodes involved in the waves' analysis (see figure 3B). We computed power spectra using wavelet transform (1-45Hz in log-space frequency steps with 1-20 cycles) for all trials when neither a target nor a distractor appeared. We then

correlated the mean alpha power in both frontal and posterior regions with alpha-band forward and backward waves between subjects in both contra and ipsilateral hemispheres. We reported Bayes Factor and Pearson's coefficients. Additionally, we computed trial-by-trial correlations between waves and alpha power for all participants. First, we tested the correlation coefficient against zero in all conditions. Then, we obtained a combined p-value per condition using the log/lin regress Fisher method (Fisher, 1992), as shown in (Zoefel et al., 2019). Specifically, we computed the T value of a chi-square distribution with $2*N$ degrees of freedom from the p_i values of the N participants as:

$$T = -2 * \sum_{i=1}^N \ln(p_i)$$

Event analysis. In this analysis, we first investigated how the onset of a target or a distractor modulates the amount of both forward and backward waves, then whether a missed or correctly identified target elicits different patterns of waves. In both cases, we performed the same procedure as in the *block analysis*: we computed forward and backward waves separately for each line of electrodes obtaining five contra- and five ipsilateral spectra per subject. First, we computed the waves 500ms before and 500ms after the target or distractor's onset, and we normalized each pair of symmetric lines as in the *block analysis* (see above). Then, we tested two separate ANOVAs considering in the first analysis the factor EVENT (either a target or a distractor occurred on the screen), and in the second the factor CORRECT (either a hit or a missed target) in the second analysis. We included DIRECTION (either FW or BW) in both models as a fixed factor and SUBJECTS as the random term.

Statistical analyses. We computed Bayes Factors (BF) in all statistical analyses, measured as the ratio between the models testing the alternative against the null hypothesis. All BFs follow this indication throughout the paper and are denoted as BF_{10} . Conventionally, large BFs provide substantial ($BF > \sim 3$) or strong ($BF > \sim 10$) evidence in favor of the alternative hypothesis, whereas low BF ($BF < \sim 0.333$) suggests a lack of effect (Smith, 2001; Masson, 2011). We performed all analyses in JASP (Love et al., 2015), considering default uniform prior distributions. The code to analyze the travelling waves is freely available at the following link: <https://github.com/artipago/Travelling-waves-EEG-2.0> (Alamia, 2023).

Acknowledgments

This work was funded by an ANR (OSCI-DEEP grant ANR-19-NEUC-0004) and ANITI (Artificial and Natural Intelligence Toulouse Institute) Research Chair (grant ANR-19-PI3A-0004) to R.V. A.A. has received funding from the European Research Council (ERC) under

the European Union's Horizon 2020 research and innovation program (grant agreement No 101075930)

References

- Alamia A. (2023). Travelling-waves-EEG-2.0. <https://github.com/artipago/Travelling-waves-EEG-2.0>.
- Alamia A, Timmermann C, Nutt DJ, Vanrullen R, Carhart-Harris RL (2020) DMT alters cortical travelling waves. *Elife* 9:1–16.
- Alamia A, VanRullen R (2019) Alpha oscillations and traveling waves: Signatures of predictive coding? *PLOS Biol* 17:e3000487.
- Alexander DM, Ball T, Schulze-Bonhage A, Van Leeuwen C (2019) Large-scale cortical travelling waves predict localized future cortical signals. *PLoS Comput Biol* 15.
- Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ (2012) Canonical microcircuits for predictive coding. *Neuron* 76:695–711 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3777738&tool=pmcentrez&rendertype=abstract> [Accessed May 23, 2014].
- Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, DeWeerd P, Kennedy H, Fries P (2015) Visual areas exert feedforward and feedback influences through distinct frequency channels. *Neuron* 85:390–401.
- Bonnefond M, Jensen O (2012) Alpha oscillations serve to protect working memory maintenance against anticipated distracters. *Curr Biol* 22:1969–1974.
- Brainard DH (1997) The Psychophysics Toolbox. *Spat Vis* 10:433–436 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9176952>.
- Brüers S, VanRullen R (2018) Alpha power modulates perception independently of endogenous factors. *Front Neurosci* 12.
- Busch NA, Dubois J, VanRullen R (2009) The phase of ongoing EEG oscillations predicts visual perception. *J Neurosci* 29:7869–7876.
- Buzsáki G (2009) *Rhythms of the Brain*. Oxford University Press.
- Buzsáki G, Draguhn A (2004) Neuronal oscillations in cortical networks. *Science* (80-) 304:1926–1929.
- Capilla A, Schoffelen JM, Paterson G, Thut G, Gross J (2014) Dissociated α -band modulations in the dorsal and ventral visual pathways in visuospatial attention and perception. *Cereb Cortex*.
- Delorme A, Makeig S (2004) EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods* 134:9–21.
- Deng Y, Reinhart RMG, Choi I, Shinn-Cunningham B (2019) Causal links between parietal alpha activity and spatial auditory attention. *Elife*.
- Fakche C, Vanrullen R, Marque P, Dugué L (2022) α Phase-Amplitude Tradeoffs Predict Visual Perception. *eNeuro* 9.
- Feldmann-Wüstefeld T, Vogel EK (2019) Neural Evidence for the Contribution of Active Suppression During Working Memory Filtering. *Cereb Cortex*.
- Fisher RA (1992) *Statistical Methods for Research Workers*.

- Foster JJ, Sutterer DW, Serences JT, Vogel EK, Awh E (2017) Alpha-Band Oscillations Enable Spatially and Temporally Resolved Tracking of Covert Spatial Attention. *Psychol Sci* 28:929–941.
- Gulbinaite R, İlhan B, Vanrullen R (2017) The triple-flash illusion reveals a driving role of alpha-band reverberations in visual perception. *J Neurosci* 37:7219–7230.
- Halgren M, Ulbert I, Bastuji H, Fabó D, Eross L, Rey M, Devinsky O, Doyle WK, Mak-McCully R, Halgren E, Wittner L, Chauvel P, Heit G, Eskandar E, Mandell A, Cash SS (2019) The generation and propagation of the human alpha rhythm. *Proc Natl Acad Sci U S A* 116:23772–23782.
- Händel BF, Haarmeier T, Jensen O (2011) Alpha oscillations correlate with the successful inhibition of unattended stimuli. *J Cogn Neurosci* 23:2494–2502.
- Huang Y, Rao RPN (2011) Predictive coding. *Wiley Interdiscip Rev Cogn Sci* 2:580–593 Available at: <http://doi.wiley.com/10.1002/wcs.142> [Accessed May 24, 2014].
- Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Front Hum Neurosci* 4 Available at: <http://journal.frontiersin.org/article/10.3389/fnhum.2010.00186/abstract>.
- Kasten FH, Wendeln T, Stecher HI, Herrmann CS (2020) Hemisphere-specific, differential effects of lateralized, occipital–parietal α - versus γ -tACS on endogenous but not exogenous visual-spatial attention. *Sci Rep* 10.
- Kelly SP, Lalor EC, Reilly RB, Foxe JJ (2006) Increases in alpha oscillatory power reflect an active retinotopic mechanism for distracter suppression during sustained visuospatial attention. *J Neurophysiol* 95:3844–3851.
- Klimesch W (2012) Alpha-band oscillations, attention, and controlled access to stored information. *Trends Cogn Sci* 16:606–617.
- Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Res Rev* 53:63–88.
- Lopes da Silva FH, Vos JE, Mooibroek J, van Rotterdam A (1980) Relative contributions of intracortical and thalamo-cortical processes in the generation of alpha rhythms, revealed by partial coherence analysis. *Electroencephalogr Clin Neurophysiol* 50:449–456.
- Love J, Selker R, Verhagen J, Marsman M, Gronau QF, Jamil T, Smira M, Epskamp S, Wild A, Ly A, Matzke D, Wagenmakers E-J, Morey RD, Rouder JN (2015) Software to sharpen your stats. *APS Obs* 28:27–29.
- Lozano-Soldevilla D, VanRullen R (2019) The Hidden Spatial Dimension of Alpha: 10-Hz Perceptual Echoes Propagate as Periodic Traveling Waves in the Human Brain. *Cell Rep* 26:374–380.e4 Available at: <https://doi.org/10.1016/j.celrep.2018.12.058>.
- Luo C, VanRullen R, Alamia A (2021) Conscious perception and perceptual echoes: a binocular rivalry study. *Neurosci Conscious* 2021.
- Masson MEJ (2011) A tutorial on a practical Bayesian alternative to null-hypothesis significance testing. :679–690.
- Mathewson KE, Lleras A, Beck DM, Fabiani M, Ro T, Gratton G (2011) Pulsed out of awareness: EEG alpha oscillations represent a pulsed-inhibition of ongoing cortical processing. *Front Psychol* 2.
- Michalareas G, Vezoli J, van Pelt S, Schoffelen JM, Kennedy H, Fries P (2016) Alpha-Beta and Gamma Rhythms Subserve Feedback and Feedforward Influences among Human Visual Cortical Areas. *Neuron* 89:384–397.

681 Muller L, Chavane F, Reynolds J, Sejnowski TJ (2018) Cortical travelling waves: Mechanisms and
682 computational principles. *Nat Rev Neurosci* 19:255–268.

683 Noonan MAP, Crittenden BM, Jensen O, Stokes MG (2018) Selective inhibition of distracting input. *Behav*
684 *Brain Res.*

685 Nunez PL (1974) The brain wave equation: a model for the EEG. *Math Biosci* 21:279–297.

686 Palva S, Palva JM (2007) New vistas for α -frequency band oscillations. *Trends Neurosci* 30:150–158.

687 Palva S, Palva JM (2011) Functional roles of alpha-band phase synchronization in local and large-scale cortical
688 networks. *Front Psychol* 2.

689 Pang Z, Alamia A, Vanrullen R (2020) Turning the stimulus on and off changes the direction of α traveling
690 waves. *eNeuro* 7:1–11.

691 Peylo C, Hilla Y, Sauseng P (2021) Cause or consequence? Alpha oscillations in visuospatial attention. *Trends*
692 *Neurosci* 44:705–713.

693 Ruzzoli M, Torralba M, Moris Fernández L, Soto-Faraco S (2019) The relevance of alpha phase in human
694 perception. *Cortex* 120:249–268.

695 Sadaghiani S, Kleinschmidt A (2016) Brain Networks and α -Oscillations: Structural and Functional Foundations
696 of Cognitive Control. *Trends Cogn Sci.*

697 Sauseng P, Klimesch W, Stadler W, Schabus M, Doppelmayr M, Hanslmayr S, Gruber WR, Birbaumer N (2005)
698 A shift of visual spatial attention is selectively associated with human EEG alpha activity. *Eur J Neurosci*
699 22:2917–2926.

700 Schneider D, Herbst SK, Klatt LI, Wöstmann M (2021a) Target enhancement or distractor suppression?
701 Functionally distinct alpha oscillations form the basis of attention. *Eur J Neurosci.*

702 Schneider M, Broggin AC, Dann B, Tzanou A, Uran C, Sheshadri S, Scherberger H, Vinck M (2021b) A
703 mechanism for inter-areal coherence through communication based on connectivity and oscillatory power.
704 *Neuron* 109:4050–4067.e12.

705 Schuhmann T, Kemmerer SK, Duecker F, de Graaf TA, Oever S ten, de Weerd P, Sack AT (2019) Left parietal
706 tACS at alpha frequency induces a shift of visuospatial attention. *PLoS One* 14.

707 Schwenk JCB, VanRullen R, Bremmer F (2020) Dynamics of visual perceptual echoes following short-term
708 visual deprivation. *Cereb Cortex Commun.*

709 Smith JMB and AFM (2001) Bayesian Theory. *Meas Sci Technol* 12:221–222.

710 Sokoliuk R, Mayhew SD, Aquino KM, Wilson R, Brookes MJ, Francis ST, Hanslmayr S, Mullinger KJ (2019)
711 Two spatially distinct posterior alpha sources fulfill different functional roles in attention. *J Neurosci*
712 39:7183–7194.


713 Spratling MW (2008) Predictive coding as a model of biased competition in visual attention. *Vision Res*
714 48:1391–1408.


715 Spratling MW (2012) Predictive coding as a model of the V1 saliency map hypothesis. *Neural Networks* 26:7–
716 28.




717 Thut G, Nietzel A, Brandt SA, Pascual-Leone A (2006) α -Band electroencephalographic activity over occipital
718 cortex indexes visuospatial attention bias and predicts visual target detection. *J Neurosci* 26:9494–9502.

719 van Kerkoerle T, Self MW, Dagnino B, Gariel-Mathis M-A, Poort J, van der Togt C, Roelfsema PR (2014)
720 Alpha and gamma oscillations characterize feedback and feedforward processing in monkey visual cortex.


721 Proc Natl Acad Sci 111:14332–14341 Available at:
 722 <http://www.pnas.org/lookup/doi/10.1073/pnas.1402773111>.
 723 VanRullen R (2016) Perceptual Cycles. Trends Cogn Sci 20:723–735.
 724 Vanrullen R, MacDonald JSP (2012) Perceptual echoes at 10 Hz in the human brain. Curr Biol 22:995–999.
 725 Waldhauser GT, Johansson M, Hanslmayr S (2012) Alpha/Beta Oscillations Indicate Inhibition of Interfering
 726 Visual Memories. J Neurosci 32:1953–1961.
 727 Watson AB, Pelli DG (1983) Quest: A Bayesian adaptive psychometric method. Percept Psychophys 33:113–
 728 120.
 729 Worden MS, Foxe JJ, Wang N, Simpson G V. (2000) Anticipatory biasing of visuospatial attention indexed by
 730 retinotopically specific alpha-band electroencephalography increases over occipital cortex. J Neurosci 20.
 731 Zhigalov A, Jensen O (2022) Travelling waves observed in MEG data can be explained by two discrete sources.
 732 bioarxiv Available at: <https://www.biorxiv.org/content/10.1101/2022.09.28.509870v2.abstract>.
 733 Zoefel B, Davis MH, Valente G, Riecke L (2019) How to test for phasic modulation of neural and behavioural
 734 responses. Neuroimage.
 735







A FORWARD AND BACKWARD WAVES' AT STIMULUS ONSET IN CONTRALATERAL ELECTRODES



B

FORWARD

