



HAL
open science

Lithium isotope geochemistry in the Barton Peninsula, King George Island, Antarctica

Jong-Sik Ryu, Nathalie Vigier, Hye-Bin Choi, Hyoun Soo Lim, Jeonghoon Lee

► To cite this version:

Jong-Sik Ryu, Nathalie Vigier, Hye-Bin Choi, Hyoun Soo Lim, Jeonghoon Lee. Lithium isotope geochemistry in the Barton Peninsula, King George Island, Antarctica. *Goldschmidt 2021*, Jul 2021, Virtual, France. <10.7185/gold2021.4010>. <hal-04028194>

HAL Id: hal-04028194

<https://hal.science/hal-04028194v1>

Submitted on 30 Jun 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Lithium isotope geochemistry in the Barton Peninsula, King George Island, Antarctica

JONG-SIK RYU¹, NATHALIE VIGIER², HYE-BIN CHOI³,
HYOUN SOO LIM⁴ AND JEONGHOON LEE⁵

¹Pukyong National University

²LOV - CNRS - Sorbonne Université

³Korea Basic Science Institute

⁴Pusan National University

⁵Ewha Womans University

Presenting Author: jongsikryu@gmail.com

Lithium (Li) has two stable isotopes, ⁶Li and ⁷Li, whose large relative mass difference is responsible for significant isotopic fractionation during physico-chemical processes. Thus Li isotopes are a good tracer of continental chemical weathering. Although physical erosion is dominant in the Polar regions due to cold climate and limited river systems, increasing global surface temperature may enhance chemical weathering, with possible consequences on carbon cycle and nutriment flux to the ocean. Fine fractions of soils have been forming during the last 6000 yr since the last deglaciation under warmer and more humid climate than other parts of Antarctica. Here, we examined elemental and Li isotope geochemistry of meltwaters, suspended sediments, soils and bedrocks in the Barton Peninsula, King George Island, Antarctica. Li concentrations range from 8.7 nM to 23.3 μM in water, from 0.01 ppm to 1.43 ppm in suspended sediment, from 9.56 ppm to 36.9 ppm in soil, and from 0.42 ppm to 28.3 ppm in bedrock. δ⁷Li values are also variable, ranging from +16.4 to +41.1‰ in water, from -0.4 to +13.4‰ in suspended sediment, from -2.5 to +6.9‰ in soil, and from -1.8 to +11.7‰ in bedrock. Correlation between elemental and Li isotope geochemistry reveals that both congruent and incongruent dissolution controls δ⁷Li values of water, rather than seasalt inputs from atmosphere or ice melting. Furthermore, PHREEQC modeling indicates that Fe-oxyhydroxides are oversaturated in some meltwater, explaining their high δ⁷Li values that resulted from preferential uptake of ⁶Li. Likewise, δ⁷Li values of suspended sediment indicate are mostly caused by modern weathering. These results confirm that increasing global surface temperature enhances modern chemical weathering in Antarctica, which therefore is expected to be stronger with time.