

Celebrating the 50th anniversary of CAIs discovery by Mireille Christophe Michel-Lévy

Matthieu Gounelle

▶ To cite this version:

Matthieu Gounelle. Celebrating the 50th anniversary of CAIs discovery by Mireille Christophe Michel-Lévy. Meteoritics and Planetary Science, 2018, 53 (11), pp.2427-2429. 10.1111/maps.13127. hal-04028403

HAL Id: hal-04028403

https://hal.science/hal-04028403

Submitted on 14 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MAPS 13127-3011 WILEY Dispatch: 29.5.18

Journal Code Manuscript No. WILEY No. of pages: 3



Meteoritics & Planetary Science 1–3 (2018) doi: 10.1111/maps.13127

Report

Celebrating the 50th anniversary of CAIs discovery by Mireille Christophe Michel-Lévy

Matthieu GOUNELLE D*

Muséum National d'Histoire Naturelle *Corresponding author. E-mail: gounelle@mnhn.fr

(Received 09 April 2018; revision accepted 14 May 2018)

Abstract–The scope of the present short report is to honour the discovery of CAIs in 1968 and her discoverer, the French mineralogist, Mireille Christophe Michel-Lévy.

In spring 1968, a paper describing an unusual chondrule in the Vigarano meteorite by Mireille Christophe Michel–Lévy appeared in the Bulletin de la Société Française de Minéralogie et de Cristallographie (Christophe Michel-Lévy 1968). Written in French, it was submitted October 30, 1967, and despite the name "chondrule," it offered the first description and interpretation of a calcium-aluminum-rich inclusion (CAI). CAIs are now known to be the first solids to have formed in our solar system (e.g., Connelly et al. 2017). The scope of the present short report was to honor the discovery and the discoverer.

In 1968, Mireille Christophe was in her 40s. She was a CNRS researcher at the Laboratoire de Minéralogie-Cristallographie, part of the Université de la Sorbonne. After an early career dedicated to mineral synthesis, she had decided to devote herself to the study of meteorites, taking advantage of the large meteorite collection gathered at the nearby Muséum by Alfred Lacroix (Lacroix 1927). Before her study of Vigarano, she had published a couple of papers on meteorites (e.g., Cristophe Michel-Lévy and Curien 1965). At that time, far fewer individuals worked on meteorites. As an example, 117 people attended the Meteoritical Society Meeting at Washington in 1966, while 539 were present in Berlin 50 years later. Meteorites were also less numerous. There were roughly 2020 known specimens in world's collections in 1966 (Hey 1966), while now researchers can count on more than 57800 samples to develop their research. The CV chondrite group, which had been defined by Wiik (1956), included eight specimens. Only five of them (the falls) had been described with some detail (Mason 1963). Vigarano, which had fallen in 1910 in Northern Italy, had been examined only by a handful of authors (Rosati 1910a,b; Mason 1963).


CE: Malarvizhi

PE: Nishanthan P

The organization of the discovery paper sounds mildly familiar. In the introduction, Madame Christophe reports that she has worked on a thin section from the Muséum national d'histoire naturelle in Paris (Fig. 1). She discusses previous work and mentions that spinel has been reported previously by Mason (1963). Given the "extraordinary" nature of the chondrule she describes, she gives no real justification of her work. In addition to optical microscope, she uses two techniques, X-ray diffraction and electron microprobe, that were at the time quite "modern" and required some specific experience. At that time, it was mainly used for elementary profiles (Fig. 1), and quantitative analyses became accurate only a few years later.

In that exceptional millimeter-sized chondrule (Fig. 1), Madame Christophe reports to have found numerous titaniferous spinel grains whose abundance decreases outwards. Spinel grains are enclosed in melilite whose akermanite content (40–70%) is determined thanks to the electron microprobe. Some areas richer in titanium are interpreted to be rutile or perovskite. Finally, she observes a reaction rim made of spinel and other minerals too fine-grained to be identified. In addition to CAIs, she had discovered the so-called Wark-Lovering rims (Wark and Lovering 1977).

In the discussion, Madame Christophe notes that such a calcium and aluminum-rich object had never been seen before in a chondritic meteorite. She clearly states it is "a primary silico-alumino-calcic condensate." From its





M. Gounelle




Fig. 1. a) Thin section used by Mireille Christophe. The large, presumably type B, CAI she discovered is on the right of the sample. The label reads "Vigarano – gros chondre à mélilite-spinelle". b) Optical micrograph of the CAI originally published in the 1968 paper. c) Electron microprobe profile published in the 1968 paper.

composition, she calculates that the extraordinary chondrule crystallized between 1400 and 1450 °C. She observes that these temperatures are of the same order than those calculated by Larimer (1967) for vapor condensation from a gas. Finally, she remarks that this object has reacted with the matrix, not during metamorphism, but when it was still hot.

The importance of the discovery is considerable. CAIs are probably among the most studied rocks on Earth, of interest for both cosmochemists and astronomers (e.g., Reipurth et al. 2007). An immense deal of effort has been invested in measuring their ages, which defines that of the solar system (Manhès et al. 1988; Connelly et al. 2017). The discovery is all the more remarkable as CAIs are far less abundant and visible in Vigarano compared to Allende that was to fall the following year (in 1969), and where CAIs were rediscovered (Marvin et al. 1970).

Remarkably enough, most of what would make later on the importance of CAIs (early high-temperature condensates made in a specific environment and transported in the chondrite-forming region) had been clearly identified by Madame Christophe in 1967. Formation mechanisms and transport modes of CAIs are now a subject of important research (Nittler and Ciesla

2016 for a review). Given the importance of the discovery, it is a bit sad that this key paper is not more quoted¹. I hope that the present note will contribute to a wider reconnaissance of the key contribution Mireille Christophe made to Meteoritics and Astrophysics.

Editorial Handling—Dr. A. J. Timothy Jul

REFERENCES

Christophe Michel-Lévy M. 1968. Un chondre exceptionnel dans la météorite de Vigarano. *Bulletin de la Société française de Minéralogie et de Cristallographie* 91:212–214.

Connelly J. N., Bollard J., and Bizzarro M. 2017. Pb-Pb chronometry and the early solar system. *Geochimica et Cosmochimica Acta* 201:345–363.

Cristophe Michel-Lévy M. and Curien H. 1965. Étude à microsonde électronique d'un chondre d'olivine et d'un fragment riche en cristobalite de la météorite de Nadiabondi. Bulletin de la Société française de Minéralogie et de Cristallographie 88:122–125.

Lacroix A. 1927. Les météorites tombées en France et dans ses colonies et conservées au Muséum National d'Histoire Naturelle, avec remarques sur la classification des

¹Possibly because it is published in French, in a relatively confidential journal, this paper is not referenced by ADS.

CAI discovery

- météorites. Bulletin de Museum National d'Histoire Naturelle 33:411–455.
- Larimer J. W. 1967. Chemical fractionations in meteorites I. Condensation of the elements. Geochimica et Cosmochimica Acta 31:1215–1238.
- Manhès G., Göpel C., and Allègre C. J. 1988. Systématique U-Pb dans les inclusions réfractaires d'Allende: le plus vieux matériau solaire.1988. *Journées de Planétologie* 323–326.
- Marvin U. B., Wood J. A., and Dickey J. S. 1970. Ca-Al rich phases in the Allende meteorite. *Earth and Planetary Science Letters* 7:346–350.
- Mason B. 1963. The carbonaceous chondrites. *Space Science Reviews* 1:621–646.
- Nittler L. R. and Ciesla F. 2016. Astrophysics with extraterrestrial materials. *Annual Review of Astronomy and Astrophysics* 54:53–93.

- Reipurth B., Jewitt D., and Keil K., ed. 2007. *Protostars and planets V.* Tucson, Arizona: The University of Arizona Press. 951 p.
- Rosati A. 1910a. Studio microscopico della meteorite caduta a Vigarano Pieve, presso Ferrara, nel gennaio 1910. *Atti R. Academia Lincei* 19 (1° semestre):841–846.
- Rosati A. 1910b. Studio microscopico di una seconda meteorite trovata a Vigarano Pieve, presso Ferrara, nel febbraio 1910. Atti R. Academia Lincei 19 (2° semestre):25–27.
- Wark D. A. and Lovering J. F. 1977. Marker events in the early solar system: evidence from rims on Ca–Al-rich inclusions in carbonaceous chondrites. Proceedings, 8th Lunar Science Conference. pp. 95–112.
- Wiik H. B. 1956. The chemical composition of some stony meteorites. *Geochimica et Cosmochimica Acta* 9:279–289.