

Mixture of segmentation for heterogeneous functional data

Vincent Brault, Emilie Devijver, Charlotte Laclau

To cite this version:

Vincent Brault, Emilie Devijver, Charlotte Laclau. Mixture of segmentation for heterogeneous functional data. 2023. hal-04036482v1

HAL Id: hal-04036482 <https://hal.science/hal-04036482v1>

Preprint submitted on 19 Mar 2023 (v1), last revised 19 Oct 2024 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

[Electronic Journal of Statistics](https://imstat.org/journals-and-publications/electronic-journal-of-statistics/) ISSN: 1935-7524

Mixture of segmentation for heterogeneous functional data

Vincent Brault1 **, Émilie Devijver**2 **and Charlotte Laclau**3

¹*Univ. Grenoble Alpes, CNRS, Grenoble INP*∗ *, LJK, 38000 Grenoble, France, e-mail:* vincent.brault@univ-grenoble-alpes.fr

²*CNRS, Univ. Grenoble Alpes, Grenoble INP*[∗] *, LIG, 38000 Grenoble, France, e-mail:* emilie.devivjer@univ-grenoble-alpes.fr

> ³*Télécom Paris, Institut Polytechnique de Paris, e-mail:* charlotte.laclau@telecom-paris.fr

Abstract: In this paper we consider functional data with heterogeneity in time and in population. We propose a mixture model with segmentation of time to represent this heterogeneity while keeping the functional structure. Maximum likelihood estimator is considered, proved to be identifiable and consistent. In practice, an EM algorithm is used, combined with dynamic programming for the maximization step, to approximate the maximum likelihood estimator. The method is illustrated on a simulated dataset, and used on a real dataset of electricity consumption.

Keywords and phrases: Mixture model, Segmentation, Functional Data, Consistency.

Contents

∗ Institute of Engineering Univ. Grenoble Alpes

/Mixture of segmentation 2

1. Introduction

Functional Data Analysis (FDA) deals with the theory and the exploration of data observed over a finite discrete grid and expressed as curves (or mathematical functions) varying over some continuum such as time. This type of data is commonly encountered in many fields, including economy (Bugni et al. (2009)), computational biology (Giacofci et al. (2013)) or environmental sciences (Bouveyron et al. (2021a)), to name a few. For an in-depth review of techniques and applications, we refer the interested readers to the books of Ferraty and Vieu (2006) and Ramsay and Silverman (2002, 2005). In many of these applications, such as electricity load, used for illustration here, we observe multiple curves corresponding to several individuals over a given time interval. As a result, one can expect a high heterogeneity of the data, both at the level of the studied individuals, that may correspond to different behavior or consumer profiles, but also on the time dimension where changes of power consumption regimes are likely to occur over the course of one year for instance. To consider a parametric model, homogeneous data is required, both at population and time levels. In this paper, we propose to split the considered heterogeneous data into homogeneous clusters of individual curves, each of them being segmented over time into homogeneous regimes. To this end, we consider a mixture of segmentation over the projection of the curves onto some functional basis. Figure 1 serves to illustrate this objective. The top-row represents the initial functional data consisting of 100 individuals (curves) observed over a period of 50 days. Following rows allow to visualize on the one hand the decomposition of the population into clusters (here 3 clusters - red yellow, purple) and on the other hand, within each cluster the segmentation obtained on the time dimension. Note that, in our case we allow each cluster to have a different segmentation, leading to a more flexible model. In this example, we visualize the segmentation on the three dimensions, denoted by *r*, of the projection on a wavelet basis.

Model-based clustering approaches for functional data have been extensively studied in the literature (James and Sugar (2003); Liu and Yang (2009); Bouveyron and Jacques (2011); Jacques and Preda (2013, 2014); Devijver (2017)). For the particular case of heterogeneous data that interests us in this article, one

Fig 1*. Motivation example. The top-row represents the initial functional data consisting of 100 individuals (curves) observed every half-hour over a period of 50 days. Following rows allow to visualize 1/ the decomposition of the population into clusters (here 3 clusters - red yellow, purple), 2/ within each cluster the segmentation obtained on the time dimension, 3/ the projection on a wavelet basis at several level r.*

can broadly differentiate between methods that perform simultaneously clustering and segmentation (Alon et al. (2003); Hébrail et al. (2010); Samé et al. (2011); Samé and Govaert (2012); Chamroukhi (2016)) and co-clustering based methods (Bouveyron et al. (2017, 2021a,b); Galvani et al. (2021)). We provide further details on these two families of approaches hereafter and position our contributions with respect to the existing state of the art.

Clustering and segmentation Samé et al. (2011) proposed to deal with heterogeneous time series by integrating the notion of change of regimes within a mixture of hidden logistic process regressions. The model is considering two latent variables, one for the mixture component and one for the segmentation. Model selection is done through an adapted BIC criterion. However, while attempting to consider changes of regime, this approach fails to account for the ordering of observations, a key feature when dealing with functional data. Samé and Govaert (2012) extended this model for online segmentation of time series. In an effort to account for these potential changes of regimes, another family of mixture models, namely the mixture of piecewise regression, has been proposed. Hébrail et al. (2010) first define this notion of piecewise regression to analyze temporal data, by proposing a distance-based model that simultaneously performs clustering on the set of functional observations (through a Kmeans-like algorithm) and segmentation (in the form of piecewise constant function summarizing) within each of the obtained cluster. This work was further generalized to a more flexible probabilistic framework by Chamroukhi (2016), who designed a model based on a mixture of piecewise regression densities. The piecewise regression is modeled by a segmentation of polynomial functions, as a generalization of spline basis where knots have to be fixed. However, this sets a particular form within each segment.

Model-based co-clustering for FDA Bouveyron et al. (2017) proposed a co-clustering model to analyze multivariate functional data. They apply this model to analyze electricity consumption curves, and found that due to the nature of the temporal data, the clustering over timepoints is in fact close to a segmentation over time. Bouveyron et al. (2021a) extend this method to multivariate time series (with several time series for each observation and each timepoint), using a sparse representation over principal components. In Bouveyron et al. (2021b), authors extend this co-clustering approach using a shape invariant model, allowing for translation in time, and translation and scaling in mean. Galvani et al. (2021) propose another bi-clustering algorithm for functional data while considering a potential misalignment through translation. While co-clustering based approach have proven efficient in this context, the clustering obtained on the time dimension do not account for the ordering of the observation.

Contributions and organization of the paper Our contribution is threefold, we propose (1) a method to study multivariate functional data, decomposing the population into homogeneous clusters and the time into homogeneous segments, where we ensure coherence on the time order; (2) we then focus on the theoretical study of the model (identifiability) and the estimation of the parameters (consistency), which is completely missing from all aforementioned related articles; (3) finally, we study a real-world electricity consumption data set to illustrate the benefits of our method.

The paper is organized as follows. In Section 2 the modeling framework is introduced together with the necessary notations. The identifiability of the model is obtained. Details about the estimation procedure are provided in Section 3. The maximum likelihood estimator is proposed, approximated by an EM algorithm. The maximization step is solved by a dynamical programming. The consistency of the estimator is provided. The finite-sample performance of the proposed estimation method is investigated in Section 4. The methodology is finally used to analyse electricity consumption in Section 5. The paper concludes by some discussion in Section 6. The code is publicly available at <https://github.com/laclauc/MixtSegmentation>. All proofs are given in the Appendix.

2. The model and its identifiability

Suppose one observes multivariate individual curves $X_1(t), \ldots, X_n(t)$ on discrete time points t_1, \ldots, t_d . First, we introduce the various elements of the modeling framework, and provide the identifiability of the model. The proof of identifiability can be found in Appendix A.

2.1. A multivariate functional model with segments in time and clusters in population

We observe multivariate individual curves $(X_{ih}(t_i))_{1 \leq i \leq n, 1 \leq j \leq d, 1 \leq h \leq H}$ of dimension *H* over *d* timepoints and within a population of size *n*. The heterogeneous population is studied through a mixture model of *K* clusters, encoded indifferently in its binary form, $z_{ik} = 1$ if and only if the curve *i* belongs to the cluster *k*, and its vector form, $z_i = k$ if and only if the curve *i* belongs to the cluster *k*, for $1 \leq k \leq K$ and $1 \leq i \leq n$. Each observation belongs to the cluster $k \in \{1, ..., K\}$ with probability $\pi_k \in [0, 1]$. The heterogeneity in time is represented through $L_k + 1$ segments $(I_{k\ell})_{0 \leq \ell \leq L_K}$: if $z_{ik} = 1$ and $j \in I_{k\ell}$, encoded by $w_{i\ell} = 1$,

$$
X_{ih}(t_j) = f_{k\ell h}(t_j) + \eta_{ijh} \tag{1}
$$

with η_{ijh} corresponds to some random noise, more details being given in Section 2.2. Usually in segmentation, we assume that the signal is constant. Here, we would like to emphasize some coherence in time, but not necessarily through a strong assumption as constant. Then, we propose to decompose our signal into several time periods that are meaningful in practice (in hours, in days, in weeks depending on the application), and to have the same function *fkℓh* within the considered interval, through the same segment.

The modeling assumption is equivalent to a main function *fkℓh* for the *h*th component, for individuals belonging to the cluster *k*, and for a timepoint in the *ℓ*th segment. This means that within a segment and a cluster, there is a random variation (seen as a noise) independent and identically distributed over each component of the multivariate curve.

2.2. Projection onto a functional basis and matrix-variate model

We denote $(\mathbf{y}_{ij}) \in \mathbb{R}^p$ the coefficient decomposition vectors of the component $j \in \{1, \ldots, d\}$ onto the functional basis, and the individual $i \in \{1, \ldots, n\}$, and the orthonormal characterization leads to, for the level *M*,

$$
(\mathbf{X}_i(t_j))_{1\leq j\leq d}=\Pi \mathbf{y}_{ij};
$$

where Π is a matrix defined by the functional basis of size *M*. We consider the wavelet coefficient dataset $(Y_i)_{1 \leq i \leq n} = (y_i)_{1 \leq i \leq n} \in (\mathbb{R}^{d \times p})^n$, which defines *n* observations whose probability distribution is modeled by the following finite matrix-variate Gaussian mixture of segmentation model. As mentioned previously, the heterogeneous population is represented by *K* clusters. For the cluster $k \in \{1, \ldots, K\}$, the heterogeneity in time is described by $L_k + 1$ segments, defined by L_k break-points $T_{k0} < T_{k1} < \cdots < T_{kL_k} < T_{k,L_k+1}$. Then, for an observation $i \in \{1, \ldots, n\}$ in the cluster $k \in \{1, \ldots, K\}$, for $\ell \in \{0, \ldots, L_k\}$ such that $j \in \{T_{k,\ell} + 1, T_{k,\ell} + 2, \ldots, T_{k,\ell+1}\}$, we have:

$$
[\mathbf{Y}_i]_j \, |(z_{ik} = 1, W_{j\ell} = 1) = \boldsymbol{\mu}_{k\ell} + \varepsilon_{ij} \tag{2}
$$

with $\varepsilon_{ij} \sim \mathcal{N}_p(0, \Sigma_{k\ell})$ where $\Sigma_{k\ell}$ is diagonal with the values $(\sigma_{k\ell r})_{1 \leq r \leq p}$.

2.3. Identifiability of the model

In this section, we first establish the identifiability of the multivariate model $(2).$

Theorem 1 (Identifiability of (2))**.** *Assume that:*

(ID.1) For every $k \in \{1, \ldots, K\}$ and $\ell \in \{0, \ldots, L_k\}$, there exists at least one $r \in \{1, \ldots, p\}$ *such that* $\sigma_{k\ell r} \neq \sigma_{k,\ell+1,r}$ *or* $\mu_{k\ell r} \neq \mu_{k,\ell+1,r}$.

(ID.2) We have:

$$
p \ge \max_{k \in \{1, ..., K\}} L_k + 1.
$$

(ID.3) If there exists $k \neq k'$ *such that* $L_k = L_{k'}$ *then:*

- *there exists* $\ell \in \{0, \ldots, L_k\}$ *such that* $T_{k\ell} \neq T_{k',\ell}$ *,*
- *or there exists* $l \in \{0, \ldots, L_k\}$ *and* $r \in \{1, \ldots, p\}$ *such that:*

 $\sigma_{k\ell r} \neq \sigma_{k',\ell,r}$ *or* $\mu_{k\ell r} \neq \mu_{k',\ell,r}$.

(ID.4) For every $k \in \{1, \ldots, K\}$, $\pi_k > 0$.

Under these assumptions, the model (2) *is identifiable.*

The breakpoints models of each cluster are identifiable by the assumption (ID.1) and (ID.2). The assumption (ID.3) allows to differentiate the clusters.

Mixture models are known to be identifiable up to a label switching: two partitions can be the same while the cluster labels being reversed. In this model, a natural order is to choose the labeling of each cluster such that

$$
k \leq k' \Leftrightarrow L_k \leq L_{k'}.
$$

This alleviates the problem of label switching; and it can be completely removed when the $(L_k)_{1\leq k\leq K}$ are all different.

3. Estimation

In this paper, we assume that *K* the number of clusters is known, as well as the number of segment within each cluster $(L_k)_{1\leq k\leq K}$.

3.1. Maximum Likelihood Estimation

Using the model (2) , under identifiability, by noting T the set of the break points and $\boldsymbol{\theta} = ((\mu_{k\ell r}, \sigma_{k\ell r})_{1 \leq k \leq K, 0 \leq \ell \leq L_K, 1 \leq r \leq p}, (\pi_k)_{1 \leq k \leq K})$ the set of parameters, we obtain the following likelihood:

lik
$$
(Y; K, T, \theta) = \prod_{i=1}^{n} \sum_{k=1}^{K} \pi_k \prod_{\ell=0}^{L_k} \prod_{j=T_{k\ell}+1}^{T_{k,\ell+1}} \prod_{r=1}^{p} \left[\frac{1}{\sqrt{2\pi\sigma_{k\ell r}}} e^{-\frac{1}{2\sigma_{k\ell r}} (Y_{ijr} - \mu_{k\ell r})^2} \right].
$$

The mixture model leads to the product over individuals $i \in \{1, \ldots, n\}$ and the sum over the clusters $k \in \{1, ..., K\}$ while the segmentation is related to the product over each segment $\ell \in \{0, \ldots, L_k\}$ and timepoints indexed by $j \in \{T_{k\ell}+1,\ldots,T_{k,\ell+1}\},$ for the cluster $k \in \{1,\ldots,K\}.$ In addition to the parameters K, T and θ , we search to estimate the partition z .

We denote $\hat{\theta}$ the maximum likelihood estimator.

3.2. EM algorithm

Considering a mixture model, we use the *Expectation Maximisation* (EM) algorithm (Dempster, Laird and Rubin (1977)) to estimate the parameters. The principle is, for the step (*c*), to fix a parameter $\boldsymbol{\theta}^{(c)}$ and break points **T**, and to maximize the following function:

$$
\left(\boldsymbol{\theta}, \boldsymbol{T}\right) \mapsto Q\left(\boldsymbol{\theta}, \boldsymbol{T}\left|\boldsymbol{\theta}^{(c)}, \boldsymbol{T}^{(c)}\right.\right) = \mathbb{E}_{\boldsymbol{z} \big| \boldsymbol{Y}; \boldsymbol{\theta}^{(c)}, \boldsymbol{T}^{(c)}}\left[\log p\left(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}\right)\right]
$$

where $p(Y, z; \theta, T)$ is the joint distribution of Y and z for fixed parameters θ and break points T . To do so, we alternate between two steps:

E-step: For $i \in \{1, \ldots, n\}$, $k \in \{1, \ldots, K\}$ compute the values $s_{ik}^{(c)}$ defined by:

$$
s_{ik}^{(c)} = \mathbb{P}\left(z_{ik} = 1 \middle| \boldsymbol{Y}; \boldsymbol{\theta}^{(c)}, \boldsymbol{T}^{(c)}\right);
$$

M-step: Maximization with respect to *θ* and *T* the function $(\boldsymbol{\theta}, \boldsymbol{T}) \mapsto Q\left(\boldsymbol{\theta}, \boldsymbol{T} \Big| \boldsymbol{\theta}^{(c)}, \boldsymbol{T}^{(c)}\right)$.

For $i \in \{1, \ldots, n\}, k \in \{1, \ldots, K\}$, the computation of $s_{ik}^{(c)}$ is explicit using the following proposition.

Proposition 1 (E-Step). *For every* $i \in \{1, \ldots, n\}$ and $k \in \{1, \ldots, K\}$ *we have:*

$$
s_{ik}^{(c)} = \frac{\pi_k^{(c)} \prod_{\ell=0}^{L_k^{(c)}} \prod_{j=T_{k\ell}^{(c)}+1}^{T_{k,\ell+1}^{(c)}} \prod_{r=1}^p \left[\frac{1}{\sqrt{2\pi \sigma_{k\ell r}^{(c)}}} e^{-\frac{1}{2\sigma_{k\ell r}^{(c)}} \left(Y_{i\ell r} - \mu_{k\ell r}^{(c)}\right)^2} \right]}{\sum_{k'=1}^K \pi_{k'}^{(c)} \prod_{\ell=0}^{L_{k'}^{(c)}} \prod_{j=T_{k'\ell}^{(c)}+1}^{T_{k',\ell+1}^{(c)}} \prod_{r=1}^p \left[\frac{1}{\sqrt{2\pi \sigma_{k'\ell r}^{(c)}}} e^{-\frac{1}{2\sigma_{k'\ell r}^{(c)}} \left(Y_{i\ell r} - \mu_{k'\ell r}^{(c)}\right)^2} \right]}.
$$

For the maximization, the problem is more difficult due to the unknown segmentation over time. The next proposition explicit the formulae for the proportions $(\pi_k)_{1\leq k\leq K}$.

Proposition 2 (Proportion in the M-Step). For every $k \in \{1, ..., K\}$ we have:

$$
\pi_k^{(c)} = \frac{s_{+k}^{(c)}}{n} = \frac{\sum_{i=1}^n s_{ik}^{(c)}}{n}.
$$

The other parameters $(\mu_{k\ell r}, \sigma_{k\ell r})_{1 \leq k \leq K, 1 \leq \ell \leq L_K, 1 \leq r \leq p}$ are given using the dynamic programming ((see for example Bellman and Kalaba, 1957; Kay, 1993)). We start by observing that

$$
\max_{(\theta, T)} Q\left(\theta, T \middle| \theta^{(c)}, T^{(c)}\right) = -\frac{1}{2} \sum_{k=1}^{K} \left[\min_{T_k \in \mathcal{T}} \sum_{\ell=0}^{L_k} \Delta_k^{(c)}\left(T_{k\ell}; T_{k,\ell+1}\right) \right] - \frac{ndp}{2} \log\left(2\pi\right) + \sum_{k=1}^{K} s_{+k}^{(c)} \log \pi_k,
$$
\n(3)

where $\mathcal{T} = \{0 = T_{k0} < T_{k1} < \cdots < T_{k,L_k+1} = d+1\}$ with for every $k \in \mathcal{T}$ $\{1, \ldots, K\}, 0 \le t_1 < t_2 \le d,$

$$
\Delta_k^{(c)}(t_1; t_2) = \min_{\substack{\mu \in \mathbb{R}^p, \\ \sigma \in (\mathbb{R}_+^*)^{p}}}\sum_{r=1}^p \left[s_{+k}^{(c)}(t_2 - t_1) \log \left(\sigma_r\right) + \frac{1}{\sigma_r} \sum_{j=t_1+1}^{t_2} \sum_{i=1}^n s_{ik}^{(c)} \left(Y_{ijr} - \mu_r\right)^2 \right].
$$
 (4)

This optimization problem is explicitly solved in the following proposition, for each cluster independently.

Proposition 3 (Form of $\Delta_k^{(c)}$) *k*^(c)). For all $k \in \{1, ..., K\}$ and $0 \le t_1 < t_2 \le d$, *we have:*

$$
\Delta_k^{(c)}(t_1; t_2) = \sum_{r=1}^p \left[s_{+k}^{(c)}(t_2 - t_1) \log(\hat{\sigma}_{kt_1r}) + \frac{1}{\hat{\sigma}_{kt_1r}} \sum_{j=t_1+1}^{t_2} \sum_{i=1}^n s_{ik}^{(c)} (Y_{ijr} - \hat{\mu}_{kt_1r})^2 \right]
$$

with

$$
\overline{Y}_{i,t_1:t_2,r} = \frac{1}{t_2 - t_1} \sum_{j=t_1+1}^{t_2} Y_{ijr}
$$
\n
$$
\widehat{\mu}_{kt_1r} = \sum_{i=1}^n \frac{s_{ik}^{(c)}}{s_{+k}^{(c)}} \overline{Y}_{i,t_1:t_2,r}
$$
\n
$$
\widehat{\sigma}_{kt_1r} = \sum_{i=1}^n \frac{s_{ik}^{(c)}}{s_{+k}^{(c)}} \frac{1}{t_2 - t_1} \sum_{j=t_1+1}^{t_2} (Y_{ijr} - \overline{Y}_{i,t_1:t_2,r})^2.
$$

To optimize the computation time, we suggest to first compute all the means $\overline{Y}_{i,t_1:t_2,r}^{(c)}$ for $t_2 > t_1$. In particular, in the case of σ depends only on the cluster *k*, we improve the complexity.

Proposition 4 (Complexity)**.** *The complexity of the EM algorithm with Nalgo iterations is* $\mathcal{O}\left[pnd^2 \max\left(d, KN_{algo}\right)\right]$.

Proof. The values $\overline{Y}_{i,t_1:t_2,r}$ are computed for each *i*, each *r* and each pair t_1 \lt t_2 : as the computation is a mean, the final complexity is $\mathcal{O}(pnd^3)$. For each iteration in the algorithm, the complexity of E-step is $\mathcal{O}(Kpnd)$ and, thanks the storage of the $\overline{Y}_{i,t_1:t_2,r}$, the complexity of the computation of each matrix $\Delta_k^{(c)}$ $\binom{c}{k}$ is only $\mathcal{O}(npd^2)$. Finally, the update of the estimation of μ and σ is $\mathcal{O}(Knpd)$. The combination of all these complexities gives the result.

Remark that the two most time-consuming steps are the computation of the means $\overline{Y}_{i,t_1:t_2,r}$ and the computations of $\Delta_k^{(c)}$ $k^(c)$ at each iteration; these two steps can be easily parallelized. Moreover, the table of averages can be stored for later use.

3.3. Consistency

In this section, we prove that the model introduced in Equation (2) is consistent. To simplify notations, we consider univariate functional data or projection of the observed functions onto a 1-dimensional basis, such that $p = 1$ in this section, but the conclusion would be the same. To simplify the notations, we set $\sigma_{k\ell} = 1$, but the results can be extended as well to any variance.

First, we assume that the parameter space is bounded.

Assumption C.1. *There exists* $M > 0$ *such that for all* $k \in \{1, \ldots, K\}$ *and* $l \in \{0, \ldots, L_k\},\$

$$
\mu_{k\ell} \in [-M;M].
$$

We assume that there are enough observations in each segment:

Assumption C.2. *There exists* $\tau_{\min} > 0$ *such that for all* $k \in \{1, \ldots, K\}$ *and* $l \in \{0, \ldots, L_k\},\$

$$
T_{k,\ell+1} - T_{k\ell} > \tau_{\min} d.
$$

Let $\mathcal{T}_{\tau_{\min}}$ the set of breakpoints satisfying Assumption (C.2).

We need an assumption about the rate of convergence.

Assumption C.3. *We assume that* $\log(n)/d \longrightarrow_{n,d \to +\infty} 0$ *.*

We also want to distinguish between clusters. To do so, we introduce the notion of equivalent clusters and the related symmetry.

Definition 1 (Equivalent clusters)**.** *Two partitions z [⋆] and z with K clusters are* equivalent, *denoted* $z^* \sim z$ *, if there exists a permutation* $\sigma \in \mathfrak{S}(\{1, \ldots, K\})$ *such that for all* $i \in \{1, ..., n\}$ *and* $k \in \{1, ..., K\}$ *,* $z_{i\sigma(k)} = z_{ik}^{\star}$ *.*

By similarity, we denote $\boldsymbol{\theta}', \boldsymbol{T}' \sim \boldsymbol{\theta}, \boldsymbol{T}$ *if there exists a permutation that leads to the same parameters.*

We can thus define a distance between two partitions.

Definition 2 (Distance for partitions). We define the distance $d_{0,\sim}$ between *two partitions* z, z^* *with* K *clusters by*

$$
d_{0,\sim}(z^*,z) = \min_{\sigma \in \mathfrak{S}(\{1,\ldots,K\})} \sum_{k=1}^K \sum_{i=1}^n z_{ik}^* z_{i\sigma(k)}.
$$
 (5)

For a partition z^* and a radius $r > 0$, and Z is the set of all potential partitions, let $\mathcal{B}(z^*; r)$ the ball

$$
\mathcal{B}(\boldsymbol{z}^{\star};r)=\left\{ \boldsymbol{z}\in\mathcal{Z}\left|d_{0,\sim}(\boldsymbol{z},\boldsymbol{z}^{\star})\leq rn\right. \right\} .
$$

For the consistency of the estimator, we need to distinguish between close clusters, assuming something stronger than Assumption (ID.3).

Assumption ID.3.s. If there exists $k \neq k'$ such that $L_k = L_{k'}$ then we as*sume that there exists at least τ*min*d coordinates j such that the distribution of* $Y_{ij} | z_{ik}^* = 1$ *is different from the distribution of* $Y_{ij} | z_{ik}^* = 1$ *.*

This is needed when the models within each segment are the same, and only the segments are different.

We also need an assumption stronger than $(ID.4)$ about the number of curves in each cluster:

Assumption ID.4.s. *There exists a constant* $c > 0$ *such that for every* $k \in$ $\{1, \ldots, K\}, \pi_k > c.$

Variances are supposed to be equal whatever the cluster, the segment and the dimension, and to have the value $\sigma_{kjr} = 1$. Extension to any variance is straightforward but derivations of formula are more technical.

Theorem 2. Let **Y** be a matrix of a $n \times T$ observations of the model (2) *with true parameter* $\boldsymbol{\theta}^*, \boldsymbol{T}^*$ *where the number of clusters K and the number of segments* $(L_k)_{1 \leq k \leq K}$ *are known. We assume (ID.1),* (ID.3.s), (ID.4.s), (C.1)*,* (C.2) and (C.3). Then, for every parameter $\theta \in \Theta$ and $T \in \mathcal{T}_{\tau_{\min}}$,

$$
\frac{p(\boldsymbol{Y}; \boldsymbol{\theta}, \boldsymbol{T})}{p(\boldsymbol{Y}; \boldsymbol{\theta}^\star, \boldsymbol{T}^\star)} = \max_{(\boldsymbol{\theta}', \boldsymbol{T}') \sim (\boldsymbol{\theta}, \boldsymbol{T})} \frac{p\left(\boldsymbol{Y}, \boldsymbol{z}^\star; \boldsymbol{\theta}', \boldsymbol{T}'\right)}{p\left(\boldsymbol{Y}, \boldsymbol{z}^\star; \boldsymbol{\theta}^\star, \boldsymbol{T}^\star\right)}\left[1 + o_P(1)\right] + o_P(1)
$$

where o_P *are uniform over* Θ *and* $\mathcal{T}_{\tau_{\min}}$ *, and* $(\theta', T') \sim (\theta, T)$ *means for any parameter up to the label switching.*

Sketch of proof. We will prove that the complete likelihood with a bad clustering becomes small asymptotically with respect to the complete likelihood associated to the true partition. To do so, we decompose the probability with respect to potential partitions.

$$
p(\boldsymbol{Y}; \boldsymbol{\theta}, \boldsymbol{T}) = \sum_{\boldsymbol{z} \in \mathcal{Z}} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T})
$$

=
$$
\sum_{\boldsymbol{z} \sim \boldsymbol{z}^*} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) + \sum_{\boldsymbol{z} \in \mathcal{B}(\boldsymbol{z}^*; c)} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) + \sum_{\boldsymbol{z} \notin \mathcal{B}(\boldsymbol{z}^*; c)} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}).
$$

Each term is controlled by the following propositions, that are proved in Appendix B.

Proposition 5 (Equivalent partitions)**.** *Under Assumptions (ID.1) and (ID.3), we have for all* $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ *and* $\boldsymbol{T} \in \mathcal{T}_{\tau_{\min}}$ *:*

$$
\sum_{\boldsymbol{z}\sim\boldsymbol{z}^{\star}}\frac{p(\boldsymbol{Y},\boldsymbol{z};\boldsymbol{\theta},\boldsymbol{T})}{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}\right)}=\max_{(\boldsymbol{\theta}',\boldsymbol{T}')\sim(\boldsymbol{\theta},\boldsymbol{T})}\frac{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}',\boldsymbol{T}'\right)}{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}\right)}\left[1+o_{P}(1)\right]
$$

where o_P *is uniform on* Θ *.*

Proposition 6 (Partitions close to the true one)**.** *Under Assumptions (ID.3.s), (ID.4.s), (C.1), (C.2) and (C.3), we have that for all* $\tilde{c} < c/4$ *:*

$$
\sup_{(\bm{\theta},\bm{T})\in\bm{\Theta}\times\mathcal{T}_{\tau_{\min}}}\sum_{\bm{z}\in\mathcal{B}(\bm{z}^\star;\tilde{c})}p\left(\bm{Y},\bm{z};\bm{\theta},\bm{T}\right)=o_P\left[p\left(\bm{Y},\bm{z}^\star;\bm{\theta}^\star,\bm{T}^\star\right)\right].
$$

Proposition 7 (Partitions far from the true one)**.** *Under Assumptions (C.1) and (ID.4.s), asymptotically in n and d, if there exists a sequence of radius Rnd converging to 0 such that* $R_{nd} > \max\left(\sqrt{\log K/d}, \frac{4Diam(\Theta)K^2}{\sqrt{nd}\delta(\theta^*)}, \frac{4\log(\frac{1}{c})}{d\delta(\theta^*)}\right)$ *, then for all* $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ *and all* $\boldsymbol{T} \in \mathcal{T}_{\tau_{\min}}$ *:*

$$
\sup_{(\theta,\mathbf{T})\in\Theta\times\mathcal{T}_{\tau_{\min}}}\sum_{\mathbf{z}\notin\mathcal{B}(\mathbf{z}^{\star};R_{nd})}p(\mathbf{Y},\mathbf{z};\theta,\mathbf{T})=p(\mathbf{Y},\mathbf{z}^{\star};\theta^{\star},\mathbf{T}^{\star})o_{P}(1)
$$
(6)

with probability $1 - \exp(-\varepsilon_n^2 nd)$ *where* $\varepsilon_{nd} = \min\left(\frac{\delta(\theta^*)R_{nd}}{16}, 1/\right)$ √ $\overline{2}$.

Then, we get that

$$
p(\boldsymbol{Y}; \boldsymbol{\theta}, \boldsymbol{T}) = \sum_{\boldsymbol{z} \sim \boldsymbol{z}^{\star}} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) + p(\boldsymbol{Y} | \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{\star}, \boldsymbol{T}^{\star}) o_{P}(1)
$$

which gives the result.

Corollary 1. Let **Y** be a matrix of a $n \times T$ observations of the model (2) *with true parameter* $\boldsymbol{\theta}^*, \boldsymbol{T}^*$ *where the number of clusters K and the number of segments* $(L_k)_{1\leq k \leq K}$ *are known. We assume (ID.1),* (ID.3.s), (ID.4.s), (C.1)*,* (C.2) *and* (C.3)*. Then,*

$$
\left(\widehat{\pmb{\theta}},\widehat{\pmb{T}}\right)\underset{n,d\rightarrow+\infty}{\overset{\mathbb{P}}{\rightarrow}}\left(\pmb{\theta}^{\star},\pmb{T}^{\star}\right).
$$

From Theorem 2, we have that the likelihood focuses on the true partition of the data, hence the maximum likelihood estimator is asymptotically close to the complete maximum likelihood, given the true partitions. In this particular case, since the partitions are known, our problem boils down to a standard segmentation problem.

4. Simulation study

We first provide empirical evaluation of our model on univariate generated data.

4.1. Experimental Protocol

Data generation process We simulate data based on the following generative process. For a given number of multivariate observations $n \in \{100, 1000\}$ with $H = 32$ and a number of days $d \in \{50, 100\}$, we start by generating a mixture model with $K = 3$ clusters with equal proportions $(\pi_k = \frac{1}{3} \text{ for } k \in \{1, 2, 3\}).$ For each $k \in \{1, \ldots, K\}$, we take:

$$
f_{k\ell}(t_j) = (-1)^k \cdot \alpha \cdot \cos\left(\frac{2\pi t_j}{1+\ell}\right),\,
$$

where α guides the amplitude of the generated curves, $\ell \in \{0, \ldots, L_k\}$ is the index of the current segment, and $j \in \{1, \ldots, T\}$ is the index of the timepoint. As a result α plays a dual role: the smaller α is, the harder it is to both differentiate between the clusters and to detect the break-points. In the sequel, we consider different settings by varying the values of $\alpha \in \{0.1, 0.2, 1\}$. Finally, we fix the variance $\sigma_{k\ell} = 1$ for all cluster $k \in \{1, \ldots, 3\}$ and segment $\ell \in \{1, \ldots, L_k\}.$ Figure 2 illustrates all settings for *K* = 3 and fixed *n* and *d*.

Projection onto a wavelet basis The discretization of each component of the p -dimensionial curve is projected onto a wavelet basis¹, that represents localized features of functions in a sparse way (Mallat (1999)). In our paper, the Discrete Wavelet Transform (DWT) is performed using a computationally fast pyramid algorithm (Mallat (1989); Misiti et al. (2004)). We use both scaling functions to construct approximations of the function of interest, and the wavelet functions serve to provide the details not captured by successive approximations.

Evaluation Metrics In order to evaluate the quality of our output, we consider different metrics of evaluation. For the clustering part, we compute the Adjusted Rand Index (ARI) (Hubert and Arabie (1985)) and the Normalized Classification Error (NCE) (Robert, Vasseur and Brault (2021)).

The ARI is a measure of agreement between two partitions defined by

$$
\text{ARI}(\mathbf{z}^*, \hat{\mathbf{z}}) = \frac{\sum_{k,k'} \binom{n_{k,k'}2}{2} \left[\sum_k \binom{n_k}{2} \sum_{k'} \binom{\hat{n}_{k'}}{2} / \binom{n}{2} \right]}{\frac{1}{2} \left[\sum_k \binom{n_k}{2} + \sum_{k'} \binom{\hat{n}_{k'}}{2} \right] - \left[\sum_k \binom{n_k}{2} \sum_{k'} \binom{\hat{n}_{k'}}{2} \right] / \binom{n}{2}},
$$

where n_k denotes the number of observations contained in the k -th cluster described by z^* , \hat{n}'_k is the number of observations in the estimated *k*-th cluster described by \hat{z} and $n_{kk'}$ denotes the number of observations that are in the intersection between ground-truth cluster k and the estimated cluster k' . The

 1 Note that any functional basis might be used to project the observed functions.

ARI lies between 0 and 1, with 1 indicating a perfect agreement between the two partitions and 0 that the two partitions are random.

The NCE is defined by

$$
\text{NCE}(\boldsymbol{z}^{\star}, \hat{\boldsymbol{z}}) = 1 - \frac{d_{0,\sim}(\boldsymbol{z}^{\star}, \hat{\boldsymbol{z}})}{n/K},
$$

where the distance $d_{0,\sim}$ between two partitions z, z^* is given by Eq. (5). The NCE lies in [0, 1], with 0 indicating a perfect estimation of z^* .

Regarding the quality of the segmentation part, we compute the Hausdorff distance (Brault et al. (2018a)), defined by

$$
d_{\text{Haus}}(\mathbf{T}^*; \hat{\mathbf{T}}) = \max_{k \in \{1,2,3\}} \max_{1 \leq \ell \leq L_k} \left\{ \frac{|T^*_{k\ell} - \hat{T}_{k\ell}|}{d} \right\}.
$$

Note that $d_{\text{Haus}} \in [0,1]$, where 0 indicates a perfect matching between the ground-truth and the estimated break-points.

Finally, we propose to evaluate the quality of the parameter estimation with respect to the real values, by taking the difference between the ground-truth parameters (obtained by using the projection of *fkℓ*, without additional noise) and the estimated ones.

FIG 2. Illustration of the different settings for $K = 3$ with $(\alpha = 0.1$ (left column), 0.2 (middle *column)* and 1 *(right column). The number of breakpoints is* $L_1 = 1, L_2 = 2, L_3 = 3$.

/Mixture of segmentation 14 ${\it Table~1}$

4.2. Results

ARI and NCE results are summarized in Table 1. We observe that our method behave as expected. As α increases (i.e. the task becomes easier), ARI and NCE increases and decreases, respectively. We obtain a perfect clustering when $\alpha = 1$. In addition, we recover the results of theorem 2: one can see that as the number of observations (*d*) and the number of individuals (*n*) increases, the classification gets better (contrary to the one dimensional mixture model where the proportion of errors tends toward a limit value even if the number of individuals keeps increasing).

The same trend can is observed for the segmentation results presented in Figure 3. We observe that the quality of the segmentation part (and hence the breakpoints localization) is correlated with the clustering performance aforementioned.

Finally, Figure 4 shows the performance of our approach on the estimation of the model parameters. First, we recover the consistency results stated in corollary 1. We also observe that the parameters from cluster 1 are better estimated than the ones from cluster 3. This comes as no surprise as the number of breakpoints is less for cluster 1, resulting in more observations to perform the parameters estimation.

5. Real Data Analysis

We propose to apply our model to analyze electricity consumption using the Enedis Open Data Set². We focus on the year 2020 (52 weeks), corresponding to the outburst of the COVID-19 pandemic. We built 984 observations by crossreferencing information on the type of contract subscribed to, the customer profile and the region of France. Out of these 984 observations, we removed those with missing values to obtain a final population of 889 individuals. The

²available at <https://data.enedis.fr/pages/accueil/?id=init>

Fig 3*. Boxplots of the Hausdorff distance following the number of curves (rows), observations (columns) and difficulties (colors).*

curves are observed in kW every 30 minutes. We have chosen to analyze the curves by considering the week as a time unit of interest, hence we project those curves onto the Haar basis with $p = 42$ ($d = 52$ weeks).

Remark We first ran our approach on such data. The result of the model selection (see next paragraph) resulted in 2 clusters. After having analyzed these results, we found that the model isolated all profiles associated with public lighting contracts, which are not subject to a notion of energy consumption behavior in the sense that interests us. For this reason, we chose to discard these observations in the following, which give us a final population of 791 observations.

5.1. Model Selection

When dealing with real data, we have no access to the *true* number of clusters or the number of breakpoints. We therefore propose to adapt a model selection strategy proposed in Zhang and Siegmund (2007) relying on the Bayesian information criterion (BIC) (Schwarz (1978)). We obtain the following criterion for

FIG 4. Boxplots of the difference $\hat{\mu}_{k\ell}$ and $\mu_{k\ell}^*$ following the number of curves (rows), observations (columns), difficulties (x-axis of each graphic) and cluster (colors).

our model:

$$
\text{BIC}(K, \mathbf{L}) = \max_{\theta, T} \text{lik}(\mathbf{Y}; K, T, \theta) - \underbrace{\frac{K-1}{2} \log(n)}_{\text{for } \pi}
$$
\n
$$
-\frac{1}{2} \sum_{k=1}^{K} \left[3p(L_k+1) \log(ndp) + \sum_{\ell=0}^{L_k} \log \left(\frac{\widehat{T}_{k,\ell+1} - \widehat{T}_{k\ell}}{d} np \right) \right]
$$
\n
$$
-\frac{K}{2} \log(ndp).
$$
\nfor θ

The general form of the penalty in this criterion allow us to account for the specificities of all parameters.

Since an exhaustive exploration of the number of clusters and breakpoints is not possible, we adapt the bikm1 strategy proposed by Robert (2021). Given a reference configuration (the current state of the model), we proceed as follows: */Mixture of segmentation* 17

Fig 5*. Distribution of the observations across clusters (horizontal dashed lines) and localization of the break points within each cluster (vertical dashed lines). Vertical dot points areas correspond to the start and end weeks of the lockdown periods.*

TABLE 2 *Contingency table between clusters and Enedis profiles.*

		ENT/PRO RES1, 3 or 4 Other RES	
w		36	-82

- **Backward Search**: we remove a cluster (K possible options). For each of these options, we make 10 random initializations as well as a random distribution of the observations of the deleted cluster in the remaining ones.
- **Forward Search** : we add a cluster with 1 breakpoint and make 10 random initializations.
- **Number of breakpoints**: we proceed with the same principles (backward and forward searches) for the number of breakpoints (with *K* fixed).

In the end, we obtain $K = 3$ clusters and the number of breakpoints within each cluster is given by $L_1 = 1$, $L_2 = 2$ and $L_3 = 6$. Figure 5 shows the distribution of our observations within clusters and the locations of the different breakpoints. We observe that cluster 3 only contains one observation and hence overfit on the number of breakpoints. On the basis of the information at our disposal (type of contract, profile and geographic area), we have not been able to draw interesting conclusions for this particular observation³. Aside from that particular point, the obtained clusters are essentially shaped to distinguish the profiles built by Enedis (see Table 2). We believe that choosing a less penalizing BIC criterion could reveal less discriminating effects than the profile, such as a regional effect for instance.

³We are currently investigating this point with the help of Enedis

5.2. Results Discussion

Let us begin by recalling the two periods of lockdown observed in France in 2020: the first one lasted from March 17^{th} until May 11^{th} , while the second one started on October 30*th* and ended on December 15*th*. For cluster 1, which is mainly composed of enterprises and professionals, we observe a unique breakpoint that matches with the beginning of the first lockdown. We note that for this cluster no other breakpoints are observed. We can make two assumptions regarding this point: (1) it indicates that the regime change operated by these consumers did not return to *normal* $\frac{4}{3}$ after the beginning of the crisis; (2) the return to this socalled *normal* regime did not occur at the same moment for all individuals. For the second cluster, corresponding mainly to the Residential profiles, we observe two breakpoints, each appearing during the lockdown period (mid April and end of November). Once again, two hypothesis are in order: (1) a delayed effect of each lockdown or (2) the impact of outdoor temperatures, particularly high in France between mid-April and the end of November. Finally, for the last cluster, the fact that it only contains one observation does not allow us to draw significant conclusions. We can only note that most of the breakpoints matches breakpoints from both cluster 1 and 2. Additional results are provided in Figures 6, 7 and 8 in the Supplementary.

6. Conclusion

In this paper, we define a novel model to analyze multivariate functional data by performing clustering and segmentation simultaneously. We derive an EM algorithm where the maximization step is carried out by dynamic programming. From a theoretical point of view, we establish the identifiability and the consistency of the proposed model. We apply this model on synthetic data to control the behavior and validate our theoretical statements. We also demonstrate the usefulness of our model on real electricity consumption data by focusing on the year 2020 corresponding to the outbreak of the COVID pandemic.

This work can be further extended in different directions. On the estimation part, three developments can be considered. To speed up the parameter estimation, we could adapt the pruned dynamic programming algorithm proposed by Rigaill (2015) and recently improved by Maidstone et al. (2017), who proposed to prune the set of candidate change-points rather than looking at all possible cases. In order to circumvent this speed problem, a second alternative could be to replace the current maximization step with a group-lasso procedure for the parameter estimation. For instance Brault, Chiquet and Lévy-Leduc (2017) proposed to transform the problem into an equivalent estimate of a linear regression whose parameter of interest would be sparse and thus to use the LASSO (Least Absolute Shrinkage and Selection Operator) procedures. The main challenge of these methods being the regularization parameter of the LASSO method and the multiplication of the number of estimated breakpoints. Moreover, in relation

⁴here normal refers to prior to the crisis

to the real observed data, a non parametric extension based on rank statistics (see Brault et al. (2018b)) can be studied.Finally, regarding the model selection criterion proposed in the experimental part, a theoretical study of the latter would allow us to define the most appropriate form for the penalty. Indeed, the first results appear to indicate that the proposed version is too penalizing, not allowing to highlight fine-grained information.

Appendix A: Identifiability

To prove Theorem 1, we need the following lemma:

Lemma 1 (Identifiability for the breakpoints model)**.** *We define a L-breakpoints model of p-dimensional spherical Gaussian of length d* with the parameters $0 =$ $T_0 < T_1 < \cdots < T_L < T_{L+1} = d$, $\mu \in \mathcal{M}_{(L+1) \times p}(\mathbb{R})$ and $\sigma \in (\mathbb{R}^+_*)^{L+1}$ with the $\emph{following likelihood for every $\boldsymbol{x}\in\mathbb{R}^{d\times p}$:}$

$$
p(\boldsymbol{x};\boldsymbol{T},\boldsymbol{\mu},\boldsymbol{\sigma})=\prod_{\ell=0}^{L}\prod_{j=T_{\ell}+1}^{T_{\ell+1}}\prod_{r=1}^{p}f\left(x_{jr};\mu_{lr},\sigma_{\ell r}\right)
$$

where f is the density of an univariate Gaussian distribution. We assume that: (ID.a) For every $\ell \in \{0, \ldots, L\}$ *, there exists* $r \in \{1, \ldots, p\}$ *such that:*

$$
\sigma_{\ell r} \neq \sigma_{\ell+1,r} \text{ or } \mu_{\ell r} \neq \mu_{\ell+1,r}.
$$

(ID.b) We have $d \geq L + 1$ *.*

Under the assumptions (ID.a) and (ID.b), the model is identifiable.

Proof. Let two parameters (L, T, μ, σ) and (L', T', μ', σ') satisfying the assumptions (ID.a) and (ID.b), and \boldsymbol{X} and \boldsymbol{X}' the random matrices depending of each parameter. We assume that X and X' have the same distribution. To prove that the parameters are equal, we use the characteristic function defined for every $\boldsymbol{\xi} \in \mathbb{R}^{d \times p}$ by:

$$
\Phi_{\boldsymbol{X}}\left(\boldsymbol{\xi}\right)=\mathbb{E}\left[e^{\left\langle i\boldsymbol{\xi},\boldsymbol{X}\right\rangle }\right]
$$

where $\langle \cdot, \cdot \rangle$ is the scalar product and *i* is the imaginary number, satisfying $i^2 =$

−1. In our case, we have, for every $\boldsymbol{\xi} \in \mathbb{R}^{d \times p}$:

$$
\Phi_{\mathbf{X}}\left(\boldsymbol{\xi}\right) = \mathbb{E}\left[e^{\langle i\boldsymbol{\xi},\mathbf{X}\rangle}\right] = \prod_{j=1}^{d} \mathbb{E}\left[e^{\langle i\boldsymbol{\xi}_{j},\mathbf{X}_{j}\rangle}\right]
$$
\n
$$
= \prod_{\ell=0}^{L} \prod_{j=T_{\ell}+1}^{T_{\ell+1}} \prod_{r=1}^{p} \mathbb{E}\left[e^{\langle i\boldsymbol{\xi}_{j},\mathbf{X}_{j}\rangle}\right] = \prod_{\ell=0}^{L} \prod_{j=T_{\ell}+1}^{T_{\ell+1}} \prod_{r=1}^{p} \left(e^{i\xi_{jr}\mu_{\ell r} - \frac{\sigma_{\ell r}^{2}\xi_{jr}^{2}}{2}}\right)
$$
\n
$$
= \exp\left[\sum_{\ell=0}^{L} \sum_{j=T_{\ell}+1}^{T_{\ell+1}} \sum_{r=1}^{p} \left(i\xi_{jr}\mu_{\ell r} - \frac{\sigma_{\ell r}^{2}\xi_{jr}^{2}}{2}\right)\right]
$$
\n
$$
= \exp\left[i\sum_{\ell=0}^{L} \sum_{r=1}^{p} \mu_{\ell r} \sum_{j=T_{\ell}+1}^{T_{\ell+1}} \xi_{jr} - \sum_{\ell=0}^{L} \sum_{r=1}^{p} \frac{\sigma_{\ell r}^{2}}{2} \sum_{j=T_{\ell}+1}^{T_{\ell+1}} \xi_{jr}^{2}\right].
$$

Let *w* be the vector of $\{0, \ldots, L\}^d$ with $w_j = \ell$ if and only if the $T_{\ell} + 1 \leq j \leq$ *Tℓ*+1. Then,

$$
\Phi_{\mathbf{X}}\left(\xi\right) = \exp\left[i\sum_{j=1}^{d}\sum_{r=1}^{p}\xi_{jr}\mu_{w_jr} - \frac{1}{2}\sum_{j=1}^{d}\sum_{r=1}^{p}\xi_{jr}^{2}\sigma_{w_jr}^{2}\right].
$$

The distribution of X and X' are equal if and only if the characteristic functions are equal: for all $\xi \in \mathbb{R}^{d \times p}$, we have

$$
\Phi_{\mathbf{X}}(\boldsymbol{\xi}) = \Phi_{\mathbf{X}'}(\boldsymbol{\xi})
$$
\n
$$
\Leftrightarrow i \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr} \mu_{w_{j}r} - \frac{1}{2} \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr}^{2} \sigma_{w_{j},r}^{2} =
$$
\n
$$
i \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr} \mu_{w'_{j}r}' - \frac{1}{2} \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr}^{2} \sigma_{w'_{j},r}'^{2}
$$
\n
$$
\Leftrightarrow i \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr} (\mu_{w_{j}r} - \mu_{w'_{j}r}') - \frac{1}{2} \sum_{j=1}^{d} \sum_{r=1}^{p} \xi_{jr}^{2} (\sigma_{w_{j}r}^{2} - \sigma_{w'_{j}r}') = 0.
$$

A polynomial is null if and only all coefficients are null:

$$
\text{for every } j \in \{1, \dots, d\}, \begin{cases} \text{for every } r \in \{1, \dots, p\}, \ \mu_{w_j r} = \mu'_{w'_j r} \\ \text{and } \sigma^2_{w_j r} = \sigma'_{w'_j}^2. \end{cases}
$$

By the assumption (ID.b), we know that there is at least one observation by segment (by definition of T and T') then, by definition also, $w_1 = 0 = w'_1$ which implies that

for every
$$
r \in \{1, ..., p\}
$$
, $\mu_{0,r} = \mu'_{0,r}$ and $\sigma^2_{0,r} = {\sigma'}^2_{0,r}$.

Then, if we assume that $T_1 \neq T'_1$ (for example, $T'_1 > T_1$), we observe that $w_{T_1+1} = 0$ and $w'_{T_1+1} = 1$ but we know that:

for every
$$
r \in \{1, ..., p\}
$$
, $\mu_{w_{T_1+1},r} = \mu'_{w'_{T_1+1},r}$ and $\sigma^2_{w_{T_1+1}} = {\sigma'}^2_{w'_{T_1+1}}$
\n \Leftrightarrow for every $r \in \{1, ..., p\}$, $\mu_{0,r} = \mu'_{1r}$ and $\sigma^2_{0,r} = {\sigma'}^2_{1r}$
\n \Leftrightarrow for every $r \in \{1, ..., p\}$, $\mu'_{0,r} = \mu'_{1r}$ and ${\sigma'}^2_{0,r} = {\sigma'}^2_{1r}$

by the previous results. This affirmation contradicts the assumption (ID.b), therefore $T_1 = T'_1$. We then continue with $w_{T_1+1} = 1 = w'_{T_1+1}$ and by an identical reasoning, we show that

for every
$$
r \in \{1, ..., p\}
$$
, $\mu_{1r} = \mu'_{1r}$ and $\sigma_{1r}^2 = {\sigma'}_{1r}^2$

and so on until the segment $[T_{\min L,L'} + 1; T_{\min L,L'+1}]$. If $L = L'$, the proof is finished since we prove that all the parameters are identical. If $L \neq L'$, for example $L > L'$, we observe that $w_{T_{L'+1}+1} = L + 1$ (since by the previous reasoning $T_L = T'_L$ and $T_{L'} < T_{L'+1} < d$ and $w'_{T_{L'+1}+1} = L' = L$ then, by the same reasoning, this implies:

for every
$$
r \in \{1, ..., p\}
$$
, $\mu_{w_{T_{L'+1}+1},r} = \mu'_{w'_{T_{L'+1}+1},r}$
\nand $\sigma^2_{w_{T_{L'+1}+1},r} = \sigma'^2_{w'_{T_{L'+1}+1},r}$
\n \Leftrightarrow for every $r \in \{1, ..., p\}$, $\mu_{L'+1,r} = \mu'_{L',r}$ and $\sigma^2_{L'+1,r} = \sigma'^2_{L',r}$
\n \Leftrightarrow for every $r \in \{1, ..., p\}$, $\mu_{L'+1,r} = \mu_{L',r}$ and $\sigma^2_{L'+1,r} = \sigma^2_{L',r}$

which again contradicts the assumption (ID.b). Then, $L = L'$ and, the parameters being identical, the model is identifiable. \Box

To prove Theorem 1, we start by observing that the assumptions (ID.1) and (ID.3) imply the assumptions (ID.a) and (ID.b) of Lemma 1 and, with the assumptions (ID.2), we have that the distribution of each cluster is unique. As the image of the distribution functions by any isomorphism defined on the vector subspace generated by the set of distribution functions is a free family in the arrival space, then model is identifiable (see Droesbeke, Saporta and Thomas-Agnan (2013)).

Appendix B: Consistency

B.1. Notations and first results

Let z^* be the true partition, and z any partition.

We denote $\mathbb{R}_K(z^*, z)$ the matrix of size K about contingency of intersection of the two clusterings: for all $(k, k') \in \{1, \ldots, K\}^2$,

$$
\mathbb{R}_K(z^\star, z)_{k,k'} = \sum_{i=1}^n z_{ik}^\star z_{i,k'}.
$$
 (7)

Particularly, remark that marginals give the contingency table:

$$
\forall k \in \{1, ..., K\}, \mathbb{R}_K (z^*, z)_{k,+} = z^*_{+,k} \tag{8}
$$

$$
\forall k' \in \{1, \dots, K\}, \mathbb{R}_K(z^*, z)_{+, k'} = z^*_{k', +} .
$$
 (9)

Also, if the two partitions are equal up to label switching, then $\mathbb{R}_K (z^*, z)$ is diagonal up to a permutation of rows and columns.

Similarly, we denote $\mathbb{N}_{(L_k+1)\times (L_{k'}+1)}^{(k,k')}$ (T^{\star}, T) the same matrix for the segments: for all $(k, k') \in \{1 \ldots, K\}^2$, for all $(\ell, \ell') \in \{1 \ldots, L_k\} \times \{1 \ldots, L_{k'}\},$

$$
\mathbb{N}_{(L_k+1)\times(L_{k'}+1)}^{(k,k')}(\mathbf{T}^{\star},\mathbf{T})_{\ell,\ell'} = \left| D^{\star}_{\ell}^{k} \bigcap D_{\ell'}^{k'} \right| \tag{10}
$$

where

$$
D_{\ell'}^{k'} = \{j \in \{1, ..., d\} | T_{k'\ell'} + 1 \le j \le T_{k',\ell'+1} \},
$$

$$
D^{\star}_{\ell'}^{k'} = \{j \in \{1, ..., d\} | T_{k'\ell'}^{\star} + 1 \le j \le T_{k',\ell'+1}^{\star} \}.
$$

Remark that

$$
\sum_{\ell_1=1}^{L_{k_1}} \mathbb{N}_{(L_{k_1}+1)\times(L_k+1)}^{(k_1,k)} (\mathbf{T}^{\star}, \mathbf{T})_{\ell_1, \ell} = \left| D^{\star}_{\ell} \right|.
$$
 (11)

For every (θ, T, z) , we denote

$$
F_{n}\left(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z}\right)=\log\frac{p(\boldsymbol{Y}|\boldsymbol{z};\boldsymbol{\theta},\boldsymbol{T})}{p(\boldsymbol{Y}|\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star})},
$$

and its expectation

$$
g_{n}(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z})=\mathbb{E}_{\boldsymbol{Y}|\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}}\left[F_{n}(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z})\right].
$$

In the following, we study the Kullback-Leibler divergence between two Gaussian distributions with variance 1. As the distributions only depend on the respective means, we denote $KL(\mu, \mu')$ this divergence. Remark that

$$
KL(\mu, \mu') = \frac{1}{2}(\mu - \mu')^2.
$$
 (12)

Proposition 8 (Distribution of $F_n(\theta, T, z)$). For all $\theta, T \in \mathcal{T}_{\tau_{\min}}$ and z *, we get*

$$
F_n(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}) \;\;\sim\;\; \mathcal{N}\left(g_n\left(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}\right), g_n\left(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}\right)\right).
$$

$$
g_n(\theta, T, z) = - \sum_{k=1}^K \sum_{k'=1}^K \sum_{\ell=1}^{L_k} \sum_{\ell'=1}^{L_{k'}} KL(\mu_{k\ell}^*, \mu_{k'\ell'}) \, R_K(z^*, z)_{k,k'} N_{(L_k+1)\times (L_{k'}+1)}^{(k,k')} (T^*, T)_{\ell,\ell'}.
$$

The proof stands in Section B.5.

We are also particularly interested in the maximum of g_n

$$
\widetilde{\Lambda}_n(\boldsymbol{z},\boldsymbol{T})\;\;=\;\;\max_{\boldsymbol{\theta}\in\Theta}g_n\left(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z}\right)
$$

Proposition 9. *For all z, T ,*

$$
\Lambda_{n}(z, T)
$$
\n
$$
= -\frac{1}{2} \sum_{k=1}^{K} \sum_{\ell=1}^{L_{k}} \frac{1}{\left| D^{*}{}_{\ell}^{k} \right| z_{+,k}^{*}} \sum_{k_{1}=1}^{K} \sum_{\ell_{1}=1}^{L_{k_{1}}} \mathbb{R}_{K} (z^{*}, z)_{k_{1},k} \mathbb{N}_{(L_{k_{1}}+1) \times (L_{k}+1)}^{(k_{1},k)} (T^{*}, T)_{\ell_{1},\ell}
$$
\n
$$
\times \sum_{k_{2}=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K} (z^{*}, z)_{k_{2},k} \mathbb{N}_{(L_{k_{2}}+1) \times (L_{k}+1)}^{(k_{2},k)} (T^{*}, T)_{\ell_{2},\ell} KL(\mu_{k_{1},\ell_{1}}^{*}, \mu_{k_{2},\ell_{2}}^{*})
$$

The proof stands in Section B.5.

Then, we need a measure of the minimal difference between two different parameters, computed through the Kullback-Leibler divergence.

Definition 3 (Minimal Kullback-Leibler divergence). Let $\delta(\theta^*)$ be the minimal *nonzero Kullback-Leibler divergence:*

$$
\delta(\boldsymbol{\theta}^{\star}) = \min_{\substack{k_1, \ell_1, k_2, \ell_2 \\ \mu^{\star}_{k_1, \ell_1} \neq \mu^{\star}_{k_2, \ell_2}}} KL\left(\mu^{\star}_{k_1, \ell_1}, \mu^{\star}_{k_2, \ell_2}\right) > 0.
$$

B.2. Proof of Proposition 5: equivalent partitions

Let $\sigma \in \mathfrak{S}(\{1,\ldots,K\})$ be a permutation. We say that (π, μ, T) has a *symmetry* for σ if we have, for all $k \in \{1, \ldots, K\}$:

$$
\pi_{\sigma(k)} = \pi_k, \forall \ell \in \{1, \ldots, L_k\}, \mu_{\sigma(k), \ell} = \mu_{k\ell} \text{ and } T_{\sigma(k), \ell} = T_{k\ell}.
$$

We denote $Sym(\theta, T)$ the set of permutations such that (θ, T) has a symmetry. Remark that under Assumption (ID.3), we have

$$
\#\mathrm{Sym}\left(\boldsymbol{\theta}, \boldsymbol{T}\right) = 1,
$$

which makes the next computations easier than in Brault et al. (2020), where we directly get, in our particular case, $\{\sigma \in \mathfrak{S}_{\mathbf{z},\mathbf{z}^*}\} = \{(\boldsymbol{\theta}', \mathbf{T}') \sim (\boldsymbol{\theta}, \mathbf{T})\}.$

Let σ be a permutation such that for all $i \in \{1, \ldots, n\}$ and $k \in \{1, \ldots, K\}$, we have

$$
z_{i\sigma(k)} = z_{ik}^{\star},
$$

 \mathfrak{S}_{z,z^*} the set of all possible permutations and we denote $z^{(\sigma)} := (z_{i\sigma(k)})_{ik}$. We have:

$$
p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) = p(\boldsymbol{Y}, \boldsymbol{z}^{\star(\sigma)}; \boldsymbol{\theta}, \boldsymbol{T}) = p(\boldsymbol{Y}, \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{(\sigma)}, \boldsymbol{T}^{(\sigma)})
$$

If $\sigma \in \text{Sym}(\theta)$, it leads to $p(Y, z; \theta, T) = p(Y, z^*; \theta, T)$. By summing, we get

$$
\sum_{z \sim z^*} p(Y, z; \theta, T) = \sum_{\sigma \in \mathfrak{S}_{z,z^*}} p(Y, z^{*(\sigma)}; \theta, T)
$$

=
$$
\sum_{\sigma \in \mathfrak{S}_{z,z^*}} p(Y, z^*; \theta^{(\sigma)}, T^{(\sigma)})
$$

=
$$
\sum_{(\theta', T') \sim (\theta, T)} p(Y, z^*; \theta', T').
$$

However, the function $\theta \mapsto p(Y, z^*, \theta, T)$ is unimodal and maximal for the maximum of the complete likelihood. As the estimator is consistent as soon as we have the true partition, under Assumption (ID.1), we have $p(Y, z^*; \theta, T) =$ $\mathcal{O}_P[p(Y, z^*, \theta^*, T^*)]$ when θ is in a neighborhood of θ^* and $p(Y, z^*, \theta, T)$ $\rho_P[p(Y, z^*, \theta^*, T^*)]$ elsewhere. If θ is close to θ^* , the set of equivalent θ' but not symmetric are far and we get:

$$
p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}',\boldsymbol{T}'\right)=o_{P}\left[p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}\right)\right].
$$

Then,

$$
\sum_{(\boldsymbol{\theta}',\boldsymbol{T}')\sim(\boldsymbol{\theta},\boldsymbol{T})}\frac{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}',\boldsymbol{T}'\right)}{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}\right)}=\max_{(\boldsymbol{\theta}',\boldsymbol{T}')\sim(\boldsymbol{\theta},\boldsymbol{T})}\frac{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}',\boldsymbol{T}'\right)}{p\left(\boldsymbol{Y},\boldsymbol{z}^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star}\right)}\left[1+o_{P}(1)\right].
$$

B.3. Proof of Proposition 6: partitions that are close

Lemma 2. *Assume Ass. (ID.4.s)* with parameter $c > 0$. Let the event

$$
\Omega_1(c) = \left\{ \boldsymbol{z}^{\star} \in \mathcal{Z} \middle| \forall k \in \{1,\ldots,K\}, \, z_{+,k}^{\star} \geq nc/2 \right\},\,
$$

where Z *is the set of all possible partitions. Then,*

$$
\mathbb{P}_{\theta^{\star}}\left(\overline{\Omega_1(c)}\right) \leq Ke^{-\frac{nc^2}{2}}.
$$

Proof. First, remark that for all $k \in \{1, ..., K\}$, $Z_{+,k}^{\star} \sim \mathcal{B}(\pi_k^{\star})$ with $\pi_k^{\star} > c$ according to Assumption (ID.4.s). Then, using Hoeffding inequality,

$$
\mathbb{P}_{\theta^*} \left(\overline{\Omega_1(c)} \right) = \mathbb{P}_{\theta^*} \left(\bigcup_{k=1}^K \{ Z_{+,k}^* < nc/2 \} \right) \le \sum_{k=1}^K \mathbb{P}_{\theta^*} \left(Z_{+,k}^* < nc/2 \right) \\
\le \sum_{k=1}^K \mathbb{P}_{\theta^*} \left(Z_{+,k}^* < n\pi_k^* / 2 \right) \le \sum_{k=1}^K \mathbb{P}_{\theta^*} \left(Z_{+,k}^* - n\pi_k^* < -n\pi_k^* / 2 \right) \\
\le \sum_{k=1}^K \exp \left[-\frac{2(-n\pi_k^* / 2)^2}{\sum_{i=1}^n (1-0)^2} \right] \le \sum_{k=1}^K \exp \left[-\frac{2n^2 \pi_k^{*2}}{4n} \right] \\
\le \sum_{k=1}^K \exp \left[-\frac{n\pi_k^{*2}}{2} \right] \le \sum_{k=1}^K \exp \left[-\frac{nc^2}{2} \right] \le K \exp \left[-\frac{nc^2}{2} \right].
$$

In the next lemma, we split the balls into slices.

Lemma 3 (Upper bounding of $F_n(\theta, T, z)$). *Under Assumptions (ID.3.s)*, *(C.1)* and *(C.2)*, for all $r \geq 1/n$, for all $\theta \in \Theta$ and $z \in \mathcal{Z}$ such that $d_{0,\sim}(z, z^*) = rn$ *, we have:*

$$
F_n(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}) \le -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^*)}{2}rn\left[1 + o_P(1)\right]. \tag{13}
$$

Proof. First, we remark that for all θ , $T \in \mathcal{T}_{\tau_{\min}}$ and z , we have:

$$
F_n(\theta, T, z) \leq F_n(\theta, T, z) - g_n(\theta, T, z) + \hat{\Lambda}_n(z)
$$

\n
$$
\leq F_n(\theta, T, z) - g_n(\theta, T, z) - \frac{d\tau_{\min}\delta(\theta^*)}{2} d_{0,\sim}(z, z^*)
$$

\n
$$
\leq F_n(\theta, T, z) - g_n(\theta, T, z) - \frac{d\tau_{\min}\delta(\theta^*)}{2}rn.
$$

By Lemma 8, $F_n(\theta, T, z) - g_n(\theta, T, z)$ is a centered Gaussian. We also know that

$$
(\mu_{k\ell} - \mu_{k'\ell'})^2 \leq (\text{Diam}\Theta)^2
$$

$$
\sum_{\ell=1}^{L_k} \sum_{\ell'=1}^{L_{k'}} \mathbb{N}_{(L_k+1)\times(L_{k'}+1)}^{(k,k')}
$$

$$
\sum_{k=1}^K \sum_{\ell'=1}^K \mathbb{R}_K (\mathbf{z}^*, \mathbf{z})_{k,k'} \leq n,
$$

then using Lemma 6, we get for all $t > 0$:

$$
\mathbb{P}\left(F_n\left(\theta,\boldsymbol{T},\boldsymbol{z}\right)-g_n\left(\theta,\boldsymbol{T},\boldsymbol{z}\right)\geq t\right)\leq \exp\left\{-\frac{t^2}{2\mathrm{Diam}(\boldsymbol{\Theta})^2nd}\right\},\,
$$

then for all partition such that $d_{0,\sim}(z, z^*) = rn$, we have:

$$
\mathbb{P}\left(F_n\left(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}\right) - g_n\left(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z}\right) \geq \frac{dd_{0,\sim}(\boldsymbol{z}, \boldsymbol{z}^{\star})\tau_{\min}\delta\left(\boldsymbol{\theta}^{\star}\right)}{2}\right)
$$
\n
$$
\leq \exp\left\{-\frac{\frac{d^2\tau_{\min}^2d_{0,\sim}(\boldsymbol{z}, \boldsymbol{z}^{\star})^2\delta\left(\boldsymbol{\theta}^{\star}\right)^2}{2\mathrm{Diam}(\boldsymbol{\Theta})^2nd}\right\}
$$
\n
$$
\leq \exp\left\{-\frac{\delta\left(\boldsymbol{\theta}^{\star}\right)^2d\tau_{\min}d_{0,\sim}(\boldsymbol{z}, \boldsymbol{z}^{\star})r}{8\mathrm{Diam}(\boldsymbol{\Theta})^2}\right\} \underset{n, d \to +\infty}{\longrightarrow} 0,
$$

where *d*0*,*[∼] depends on *n*. Then,

$$
F_n(\theta, T, z) \leq o_P \left[\frac{d\tau_{\min} d_{0,\sim}(z, z^*) \delta(\theta^*)}{2} \right] - \frac{d\tau_{\min} \delta(\theta^*)}{2} r n
$$

$$
\leq -\frac{d\tau_{\min} \delta(\theta^*)}{2} r n \left[1 + o_P(1) \right]
$$

 \Box

Lemma 4 (Having different partitions)**.** *For all c >* 0*, considering the event* $\Omega_1(c)$ *, we have for all* $\tilde{c} \leq c/4$ *and* $\boldsymbol{z} \in \mathcal{B}(\boldsymbol{z}^*, \tilde{c})$ *:*

$$
\frac{p(z;\boldsymbol{\theta},\boldsymbol{T})}{p(z^{\star};\boldsymbol{\theta}^{\star},\boldsymbol{T}^{\star})} \leq \mathcal{O}_P\left(1\right) e^{M_{c/4} \|z-z^{\star}\|_0}.
$$
\n(14)

Proof. Considering $\Omega_1(c)$: for all $k \in \{1, ..., K\}$, we have $z_{+,k}^* \geq nc/2$. As $z \in \mathcal{B}(z^*, \tilde{c})$ with $\tilde{c} \leq c/4$ then for all $k \in \{1, \ldots, K\}$, we have $z_{+,k} \geq nc/4$.

For a partition *z*, let $\hat{\pi}(z)$ be the maximum of $\pi \mapsto p(z;\pi)$: for all $k \in$ $\{1, \ldots, K\},\$

$$
\widehat{\boldsymbol{\pi}}\left(\boldsymbol{z}\right)_k = \frac{z_{+,k}}{n}.
$$

Then, we have

$$
\frac{p(z;\theta,T)}{p(z^\star;\theta^\star,T^\star)} = \frac{p(z;\pi)}{p(z^\star;\pi^\star)} \leq \frac{p(z;\widehat{\pi}(z))}{p(z^\star;\widehat{\pi}(z^\star))} \times \frac{p(z;\widehat{\pi}(z^\star))}{p(z^\star;\pi^\star)},
$$

by definition of $\hat{\pi}(z)$. Using Brault et al. (2020)[Lemma D.2] we get:

$$
\log \left[\frac{p(z; \hat{\pi}(z))}{p(z^{\star}; \hat{\pi}(z^{\star}))} \right] = \log p(z; \hat{\pi}(z)) - \log p(z^{\star}; \hat{\pi}(z^{\star}))
$$

$$
= \sum_{i=1}^{n} \sum_{k=1}^{K} z_{ik} \log \hat{\pi}(z)_{k} - \sum_{i=1}^{n} \sum_{k=1}^{K} z_{ik}^{\star} \log \hat{\pi}(z^{\star})_{k}
$$

$$
= n \sum_{k=1}^{K} [\hat{\pi}(z)_{k} \log \hat{\pi}(z)_{k} - \hat{\pi}(z^{\star})_{k} \log \hat{\pi}(z^{\star})_{k}].
$$

Let $H(\pi) = \sum_{k=1}^{K} \pi_k \log \pi_k$. This function is differentiable and we can use the mean value theorem to the function $x \mapsto -x \log x$ with derivative $x \mapsto \log x + 1$. Then, for all $k \in \{1, ..., K\}$, there exists $\kappa_k \in]\pi_k^{\star}; \pi_k[$ such that

$$
\frac{\pi_k \log \pi_k - \pi_k^* \log \pi_k^*}{\pi_k - \pi_k^*} = -\log \kappa_k - 1
$$

$$
\Rightarrow \left| \frac{\pi_k \log \pi_k - \pi_k^* \log \pi_k^*}{\pi_k - \pi_k^*} \right| = \left| -\log \kappa_k - 1 \right|.
$$

However, as π and π^* are regular enough, we know that $\kappa_k \in]c/4; 1 - c/4[$ then there exists a constant $M_{c/4}$ such that

$$
\left|\frac{\pi_k \log \pi_k - \pi_k^* \log \pi_k^*}{ \pi_k - \pi_k^*}\right| \le \frac{M_{c/4}}{2}.
$$

By summing, we get

$$
|H(\pi) - H(\pi^*)| = \left| \sum_{k=1}^K \pi_k \log \pi_k - \sum_{k=1}^K \pi_k^* \log \pi_k^* \right|
$$

$$
\leq \sum_{k=1}^K \frac{M_{c/4}}{2} |\pi_k - \pi_k^*| \leq \frac{M_{c/4}}{2} ||\pi - \pi^*||_1.
$$

Finally, we have

$$
\begin{aligned}\n&\left|\log\left[\frac{p(z;\hat{\pi}(z))}{p(z^{\star};\hat{\pi}(z^{\star}))}\right]\right| \\
&= n\left|H(\hat{\pi}(z))-H(\hat{\pi}(z^{\star}))\right| \le \frac{nM_{c/4}}{2} \left\|\hat{\pi}(z)-\hat{\pi}(z^{\star})\right\|_{1} \\
&\le \frac{nM_{c/4}}{2} \sum_{k=1}^{K} \left|\frac{z_{+,k}}{n} - \frac{z_{+,k}^{\star}}{n}\right| \\
&\le \frac{M_{c/4}}{2} \sum_{i=1}^{n} \sum_{k=1}^{K} |z_{ik} - z_{ik}^{\star}| \le \frac{M_{c/4}}{2} \sum_{i=1}^{n} \sum_{k=1}^{K} \mathbb{1}_{\{z_{ik} \neq z_{ik}^{\star}\}} \\
&\le M_{c/4} \left\|z-z^{\star}\right\|_{0}.\n\end{aligned}
$$

Indeed, if i does not belong to the true partition, there are two nonzero terms. On the other side, by the law of large numbers, we know that $\hat{\pi}(z^*) = (z^*)$ and $\hat{\pi}(\bar{z}^*)$ is regular, we have: $\mathcal{O}_P(\pi^*)$ and as π^* is regular, we have:

$$
\frac{p(z;\hat{\pi}(z^*))}{p(z^*,\pi^*)} = \mathcal{O}_P(1).
$$
\n(15)

Proof of Proposition 6. Considering $\tilde{c} < c/4$, $\theta \in \Theta$ and $T \in \mathcal{T}_{\tau_{\min}}$, considering the event $\Omega_1(c)$:

$$
\sum_{z \in \mathcal{B}(z^*,\tilde{c})} p(Y,z;\theta,T) \n= \sum_{z \in \mathcal{B}(z^*,\tilde{c})} p(Y|z;\theta,T) p(z;\theta,T) \frac{p(Y|z^*,\theta^*,T^*)}{p(Y|z^*,\theta^*,T^*)} \n= \sum_{z \in \mathcal{B}(z^*,\tilde{c})} \frac{p(Y,z^*,\theta^*,T^*)}{p(z^*,\theta^*,T^*)} p(z;\theta,T) \frac{p(Y|z;\theta,T)}{p(Y|z^*,\theta^*,T^*)} \n= p(Y,z^*,\theta^*,T^*) \sum_{z \in \mathcal{B}(z^*,\tilde{c})} \frac{p(z;\theta,T)}{p(z^*,\theta^*,T^*)} e^{F_n(\theta,T,z)} \n= p(Y,z^*,\theta^*,T^*) \sum_{z \in \mathcal{B}(z^*,\tilde{c})} \mathcal{O}_P(1) e^{M_{c/4}||z-z^*||_0} e^{-\frac{d\tau_{\min}\delta(\theta^*)}{2}||z-z^*||_0[1+o_P(1)]}
$$

using Eqs. (13) and (14) . Then,

$$
\sum_{\mathbf{z}\in\mathcal{B}(\mathbf{z}^*,\tilde{c})} p(\mathbf{Y},\mathbf{z};\boldsymbol{\theta},\mathbf{T})
$$
\n
$$
\leq p(\mathbf{Y},\mathbf{z}^*,\boldsymbol{\theta}^*,\mathbf{T}^*) \mathcal{O}_P(1) \sum_{R=1}^{[\tilde{c}n]} {n \choose R} K^R e^{M_{c/4}R - \frac{d\tau_{\min}\delta(\boldsymbol{\theta}^*)}{2}R[1+o_P(1)]}
$$

because there are at most $\binom{n}{R} K^R$ possible partitions at distance R , and then

$$
\sum_{z \in \mathcal{B}(z^*,\tilde{c})} p(Y,z;\theta,T) \n\leq p(Y,z^*,\theta^*,T^*) \mathcal{O}_P(1) \sum_{R=1}^{[\tilde{c}n]} {n \choose R} \left(e^{\log K + M_{c/4} - \frac{d\tau_{\min}\delta(\theta^*)}{2} [1 + o_P(1)]} \right)^R \n\leq p(Y,z^*,\theta^*,T^*) \mathcal{O}_P(1) \left(1 + e^{\log K + M_{c/4} - \frac{d\tau_{\min}\delta(\theta^*)}{2} [1 + o_P(1)]} \right)^n \n\leq p(Y,z^*,\theta^*,T^*) \mathcal{O}_P(1) \exp \left\{ n \log \left[1 + e^{\log K + M_{c/4} - \frac{d\tau_{\min}\delta(\theta^*)}{2} [1 + o_P(1)]} \right] \right\} \n\leq p(Y,z^*,\theta^*,T^*) \mathcal{O}_P(1) \exp \left\{ n e^{\log K + M_{c/4} - \frac{d\tau_{\min}\delta(\theta^*)}{2} [1 + o_P(1)]} [1 + o(1)] \right\} \n\leq p(Y,z^*;\theta^*,T^*) \mathcal{O}_P(1) \n\times \exp \left\{ e^{-d \left\{ \frac{\log(n)}{d} + \frac{\log K + M_{c/4}}{d} - \frac{\delta(\theta^*)}{2} [1 + o_P(1)] \right\}} [1 + o(1)] \right\} \n\leq p(Y,z^*;\theta^*,T^*) o_P(1) \text{ using Assumption (C.3).}
$$

As this is true for all $\theta \in \Theta$ and all $T \in \mathcal{T}_{\tau_{\min}}$, this is also true for the maximum. \Box

B.4. Proof of Proposition 7: partitions that are far

Proposition 10 (Separability). *Assume (ID.3.s) and (C.2). There exists* $R > 0$ *and a constant B*(*R*) *such that*

$$
\max_{\mathbf{T}} \widetilde{\Lambda}_n(\mathbf{z}, \mathbf{T}) \le -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^*)\max_k L_k}{2} d_{0,\sim}(\mathbf{z}, \mathbf{z}^*)
$$
 for $\mathbf{z} \in \mathcal{B}(\mathbf{z}^*; R)$;
\n
$$
\max_{\mathbf{T}} \widetilde{\Lambda}_n(\mathbf{z}, \mathbf{T}) \le -B(R)dn
$$
 for $\mathbf{z} \notin \mathcal{B}(\mathbf{z}^*; R)$.

Proof. If $z \notin B(z^*; R)$, because the two partitions z and z^* are far from each other (at least a radius $R > 0$), there exists a constant $B(R)$ such that the second inequality holds.

Else, let assume that $z \in \mathcal{B}(z^*, R)$. From Assumptions (ID.3.s) and (C.2), for all $k \neq k'$, there exists at least $\tau_{\text{min}}d$ columns such that the Kullback-Leibler divergence is strictly positive, then

$$
\sum_{\ell_2=1}^{L_{k_2}} \mathbb{N}_{(L_{k_2}+1)\times(L_{k+1})}^{(k_2,k)}(T^{\star},T)_{\ell_2,\ell} \operatorname{KL}\left(\mu^{\star}_{k_1,\ell_1},\mu^{\star}_{k_2,\ell_2}\right) \geq d\tau_{\min}\delta(\boldsymbol{\theta}^{\star}).
$$

Then, according to Proposition 9, we get

$$
\widetilde{\Lambda}_{n}(z, T) \n\leq -\frac{d\tau_{\min}\delta(\theta^{\star})}{2} \sum_{k=1}^{K} \sum_{\ell=1}^{L_{k}} \frac{1}{\left| D^{\star}_{\ell} \right| z_{+,k}^{\star}} \sum_{k=1}^{K} \mathbb{R}_{K} (z^{\star}, z)_{k_{1},k} \n\times \sum_{\ell_{1}=1}^{L_{k_{1}}} \mathbb{N}_{(L_{k_{1}}+1)\times(L_{k}+1)}^{(k_{1},k)} (T^{\star}, T)_{\ell_{1},\ell} \n\times \left[\sum_{k_{2}=1}^{K} \mathbb{R}_{K} (z^{\star}, z)_{k_{2},k} - \mathbb{R}_{K} (z^{\star}, z)_{k_{1},k} \right] \n\times -\frac{d\tau_{\min}\delta(\theta^{\star})}{2} \sum_{k=1}^{K} \sum_{\ell=1}^{L_{k}} \frac{1}{z_{+,k}^{\star}} \sum_{k_{1}=1}^{K} \mathbb{R}_{K} (z^{\star}, z)_{k_{1},k} \n\times \left[\sum_{k_{2}=1}^{K} \mathbb{R}_{K} (z^{\star}, z)_{k_{2},k} - \mathbb{R}_{K} (z^{\star}, z)_{k_{1},k} \right],
$$

using Eq. (11) in the last inequality. Then, using Eq. (9), and the fact that $\mathbb{R}_K \left(\boldsymbol{z}^\star, \boldsymbol{z} \right)_{k_1,k} \leq \boldsymbol{z}^\star_{+,k},$

$$
\Lambda_n(z,T)
$$
\n
$$
\leq -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^{\star})\max_k L_k}{2} \sum_{k=1}^K \frac{1}{z_{+,k}^{\star}} \sum_{k=1}^K \mathbb{R}_K (z^{\star}, z)_{k_1,k}
$$
\n
$$
\times \left[\sum_{k_2=1}^K \mathbb{R}_K (z^{\star}, z)_{k_2,k} - \mathbb{R}_K (z^{\star}, z)_{k_1,k} \right]
$$
\n
$$
\leq -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^{\star})\max_k L_k}{2} \left[n - \sum_{k=1}^K \sum_{k_1=1}^K \frac{\mathbb{R}_K (z^{\star}, z)_{k_1,k}^2}{z_{+,k}^{\star}} \right]
$$
\n
$$
\leq -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^{\star})\max_k L_k}{2} \left[n - \sum_{k=1}^K \sum_{k_1=1}^K \mathbb{R}_K (z^{\star}, z)_{k_1,k} \right]
$$
\n
$$
\leq -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^{\star})\max_k L_k}{2} d_{0,\sim}(z, z^{\star}).
$$

Lemma 5 (Large deviation). *Under assumption (C.1), and for all* $\varepsilon_{nd} < 1$ / √ 2*, we get:*

$$
\mathbb{P}\left(\sup_{\theta,\mathbf{T},\mathbf{z}}\left[F_n\left(\theta,\mathbf{T},\mathbf{z}\right)-\widetilde{\Lambda}_n(\mathbf{z},\mathbf{T})\right]\geq Diam(\Theta)\sqrt{nd}K^2+4\varepsilon_{nd}Diam(\Theta)nd\right)\n\leq K^n\exp\left(-\varepsilon_{nd}^2nd\right).
$$

Proof. By definition of $F_n(\theta, T, z)$ and $g_n(\theta, T, z)$:

$$
F_n(\theta, T, z) - \Lambda_n(z, T) \leq F_n(\theta, T, z) - g_n(\theta, T, z)
$$

\n
$$
\leq - \sum_{i=1}^n \sum_{k=1}^K \sum_{k'=1}^K z_{ik}^* z_{i,k'} \sum_{\ell=1}^{L_k} \sum_{\ell'=1}^{L_{k'}} \times \sum_{\ell=1}^{L_{k'}} \sum_{\ell'=1}^{L_{k'}} \left(Y_{ij} - \mathbb{E}_{Y|z^*, \theta^*, T^*}[Y_{ij}]\right) (\mu_{k\ell}^* - \mu_{k'\ell'})
$$

\n
$$
\leq \sup_{\substack{\Gamma \in \mathbb{R}^{K \times K} \\ \|\Gamma\|_{\infty} \leq \text{Diam}(\Theta)}} \sum_{k=1}^K \sum_{k'=1}^K \sum_{\ell=1}^{L_k} \sum_{\ell'=1}^L \Gamma_{k,k'} \sum_{i=1}^n z_{ik}^* z_{i,k'} \sum_{j \in D^*_{\ell}^k \cap D_{\ell'}^{k'}} (\mu_{k\ell}^* - Y_{ij})
$$

\n
$$
\leq \sup_{\substack{\Gamma \in \mathbb{R}^{K \times K} \\ \|\Gamma\|_{\infty} \leq \text{Diam}(\Theta)}} \sum_{k=1}^K \sum_{k'=1}^K \Gamma_{k,k'} \sum_{i=1}^n z_{ik}^* z_{i,k'} \sum_{\ell=1}^{L_k} \sum_{j \in D^*_{\ell}^k} (\mu_{k\ell}^* - Y_{ij}).
$$

Let

$$
W_{k,k'} = \sum_{i=1}^n z_{ik}^{\star} z_{i,k'} \sum_{\ell=1}^{L_k} \sum_{j \in D^{\star}_\ell^k} \left(\mu_{k\ell}^{\star} - Y_{ij} \right) \sim \mathcal{N}\left(0, d \mathbb{R}_K \left(z^{\star}, z\right)_{k',k}\right).
$$

Then, using Brault et al. (2020)[Proposition C.4], for $\mathbf{\Gamma} \in \mathbb{R}^{K \times K}$ such that $\|\Gamma\|_{\infty} \leq \text{Diam}(\Theta),$

$$
\mathbb{E}_{\theta^*}\left[\sum_{k=1}^K \sum_{k'=1}^K \Gamma_{k,k'} W_{k,k'}\right] \leq \text{Diam}(\Theta) \sqrt{n} \, dK^2.
$$

Then, for all nonnegative sequence $(\varepsilon_{nd})_{(n,d)\in(\mathbb{N}^*)}$ satisfying $4\varepsilon_{nd}$ Diam $(\boldsymbol{\Theta})nd \leq \frac{8\text{Diam}(\boldsymbol{\Theta})^2nd}{2\sqrt{2}\text{Diam}(\boldsymbol{\Theta})}$ $\frac{\text{3Diam}(\Theta)^2 nd}{2\sqrt{2}\text{Diam}(\Theta)} = 2\sqrt{2}\text{Diam}(\Theta)nd$, using Lemma 6:

$$
\mathbb{P}\left(\sum_{k=1}^K \sum_{k'=1}^K \Gamma_{k,k'} W_{k,k'} - \mathbb{E}_{\theta^{\star}} \left[\sum_{k=1}^K \sum_{k'=1}^K \Gamma_{k,k'} W_{k,k'}\right] \ge 4\varepsilon_{nd} \text{Diam}(\Theta)nd\right) \n\le \exp\left[-\frac{(4\varepsilon_{nd} \text{Diam}(\Theta)nd)^2}{16 \text{Diam}(\Theta)^2 nd}\right] \le \exp\left(-\varepsilon_{nd}^2 nd\right).
$$

Then,

$$
\mathbb{P}\left(F_n\left(\theta,\mathbf{T},\mathbf{z}\right)-\widetilde{\Lambda}_n(\mathbf{z},\mathbf{T})\geq \text{Diam}(\Theta)\sqrt{nd}K^2+4\varepsilon_{nd}\text{Diam}(\Theta)nd\right)\\
\leq \exp\left(-\varepsilon_{nd}^2nd\right).
$$

This inequality holds for all (θ, T) and z , then we get:

$$
\mathbb{P}\left(\sup_{\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z}}\left[F_n(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z})-\widetilde{\Lambda}_n(\boldsymbol{z},\boldsymbol{T})\right]\ge
$$

\n
$$
\text{Diam}(\boldsymbol{\Theta})\sqrt{nd}K^2+4\varepsilon_{nd}\text{Diam}(\boldsymbol{\Theta})nd\right)
$$

\n
$$
\leq \sum_{\boldsymbol{z}\in\mathcal{Z}}\mathbb{P}\left(\sup_{\boldsymbol{\theta},\boldsymbol{T}}\left[F_n(\boldsymbol{\theta},\boldsymbol{T},\boldsymbol{z})-\widetilde{\Lambda}_n(\boldsymbol{z},\boldsymbol{T})\right]\ge
$$

\n
$$
\text{Diam}(\boldsymbol{\Theta})\sqrt{nd}K^2+4\varepsilon_{nd}\text{Diam}(\boldsymbol{\Theta})nd\right)
$$

\n
$$
\leq \sum_{\boldsymbol{z}\in\mathcal{Z}}\exp\left(-\varepsilon_{nd}^2nd\right)\leq K^n\exp\left(-\varepsilon_{nd}^2nd\right).
$$

 \Box

Let $R > 0$. Then, according to Proposition 10,

$$
\widetilde{\Lambda}_n(\mathbf{z}, \mathbf{T}) \le -B(C)nd \text{ if } \mathbf{z} \notin \mathcal{B}(\mathbf{z}^*; R)
$$
\n
$$
\widetilde{\Lambda}_n(\mathbf{z}, \mathbf{T}) \le -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^*)}{2}d_{0,\sim}(\mathbf{z}, \mathbf{z}^*)
$$
\n
$$
\le -\frac{d\tau_{\min}\delta(\boldsymbol{\theta}^*)}{2}nR_{nd} \text{ if } \mathbf{z} \in \mathcal{B}(\mathbf{z}^*; R) \setminus \mathcal{B}(\mathbf{z}^*; R_{nd}).
$$

Using Lemma 5, with $\varepsilon_{nd} = \min\left(\frac{\delta(\theta^*)\tau_{\min}R_{nd}}{16}, 1/\right)$ √ $\overline{2}$, we get with probability $1 - K^n \exp(-\varepsilon_{nd}^2 nd)$,

$$
F_n(\theta, T, z) - \tilde{\Lambda}_n(z, T) + \tilde{\Lambda}_n(z, T)
$$

\n
$$
\leq F_n(\theta, T, z) - \tilde{\Lambda}_n(z, T) - \frac{d\tau_{\min}\delta(\theta^*)}{2} nR_{nd}
$$

\n
$$
\leq \text{Diam}(\Theta) \sqrt{n d} K^2 + 4\varepsilon_{nd} \text{Diam}(\Theta) nd - \frac{d\tau_{\min}\delta(\theta^*)}{2} nR_{nd}
$$

\n
$$
\leq \text{Diam}(\Theta) \sqrt{n d} K^2 + 4 \frac{\delta(\theta^*) \tau_{\min} R_{nd}}{16} \text{Diam}(\Theta) nd - \frac{d\tau_{\min}\delta(\theta^*)}{2} nR_{nd}
$$

\n
$$
\leq \text{Diam}(\Theta) \sqrt{n d} K^2 + \frac{d\tau_{\min}\delta(\theta^*)}{4} nR_{nd} - \frac{d\tau_{\min}\delta(\theta^*)}{2} nR_{nd}
$$

\n
$$
\leq \text{Diam}(\Theta) \sqrt{n d} K^2 - \frac{d\tau_{\min}\delta(\theta^*)}{4} nR_{nd}
$$

\n
$$
\leq -d nR_{nd} \frac{\delta(\theta^*) \tau_{\min}}{4} \left[1 - \frac{4 \text{Diam}(\Theta) K^2}{\sqrt{n d} R_{nd} \tau_{\min}\delta(\theta^*)} \right]
$$

\n
$$
\leq -d nR_{nd} \frac{\tau_{\min}\delta(\theta^*)}{4}
$$

for *n* and *d* large enough. We also know that

$$
p(\mathbf{z}^{\star}; \boldsymbol{\theta}^{\star}, \mathbf{T}^{\star}) \leq e^{-\log \prod_{i=1}^{n} \prod_{k=1}^{K} \pi_{k}^{z_{ik}^{\star}}} \leq e^{-\sum_{i=1}^{n} \sum_{k=1}^{K} z_{ik}^{\star} \log \pi_{k}}
$$

\n
$$
\leq e^{-\sum_{i=1}^{n} \sum_{k=1}^{K} z_{ik}^{\star} \log c} \text{ by Assumption (C.1)},
$$

\n
$$
\leq e^{-\sum_{i=1}^{n} \log c} \leq e^{n \log \frac{1}{c}}.
$$

Then, it leads to

$$
\sum_{z \notin \mathcal{B}(\boldsymbol{z}^{\star}; R_{nd})} p(\boldsymbol{Y}, \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) = \sum_{z \notin \mathcal{B}(\boldsymbol{z}^{\star}; R_{nd})} p(\boldsymbol{Y} | \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) p(\boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T})
$$
\n
$$
= p(\boldsymbol{Y} | \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{\star}, \boldsymbol{T}^{\star}) \sum_{z \notin \mathcal{B}(\boldsymbol{z}^{\star}; R_{nd})} p(\boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) \frac{p(\boldsymbol{Y} | \boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T})}{p(\boldsymbol{Y} | \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{\star}, \boldsymbol{T}^{\star})}
$$
\n
$$
= p(\boldsymbol{Y} | \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{\star}, \boldsymbol{T}^{\star}) \sum_{z \notin \mathcal{B}(\boldsymbol{z}^{\star}; R_{nd})} p(\boldsymbol{z}; \boldsymbol{\theta}, \boldsymbol{T}) e^{F_n(\boldsymbol{\theta}, \boldsymbol{T}, \boldsymbol{z})}
$$
\n
$$
\leq p(\boldsymbol{Y}, \boldsymbol{z}^{\star}; \boldsymbol{\theta}^{\star}, \boldsymbol{T}^{\star}) e^{-n \left[dR_{nd} \frac{\delta(\boldsymbol{\theta}^{\star}) \tau_{\min}}{4} - \log \frac{1}{c} \right]}.
$$

Then, with probability $1 - K^n \exp(-\varepsilon_n^2 nd)$, under Assumption (C.1), we get:

$$
\sum_{z \notin \mathcal{B}(\mathbf{z}^\star; R_{nd})} p(\mathbf{Y}, \mathbf{z}; \boldsymbol{\theta}, \mathbf{T}) = p(\mathbf{Y}, \mathbf{z}^\star; \boldsymbol{\theta}^\star, \mathbf{T}^\star) o_P(1)
$$

for all $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ and $\boldsymbol{T} \in \mathcal{T}_{\tau_{\min}}$.

B.5. Tools and details

Lemma 6 (Chernoff's lemma). Let $Z \sim \mathcal{N}(0, \sigma^2)$. Then, for all $t > 0$:

$$
\mathbb{P}\left(Z \ge t\right) \le e^{-\frac{t^2}{2\sigma^2}}.
$$

Proof of Proposition 8. By definition,

$$
F_n(\theta, T, z) = - \sum_{i=1}^n \sum_{k=1}^K \sum_{k'=1}^K z_{ik}^* z_{i,k'} \sum_{\ell=1}^{L_k} \sum_{\ell'=1}^{L_{k'}} \sum_{j \in D^*_{\ell}^k \cap D_{\ell'}^{k'}} \log \left[\frac{\varphi(Y_{ij}; \mu_{k\ell}^*)}{\varphi(Y_{ij}; \mu_{k'\ell'})} \right].
$$

The computation gives:

$$
\log\left[\frac{\varphi(Y_{ij};\mu_{k\ell}^*)}{\varphi(Y_{ij};\mu_{k'\ell'})}\right] = \log\frac{\frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}(Y_{ij}-\mu_{k\ell})^2}}{\frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}(Y_{ij}-\mu_{k'\ell'})^2}}
$$

$$
= -\frac{1}{2}(Y_{ij}-\mu_{k\ell}^*)^2 + \frac{1}{2}(Y_{ij}-\mu_{k'\ell'})^2
$$

$$
= Y_{ij}(\mu_{k\ell}^* - \mu_{k'\ell'}) - \frac{1}{2}(\mu_{k\ell}^*^2 - \mu_{k'\ell'}^2)
$$
(16)

which leads to

$$
F_{n}(\theta, T, z) = -\sum_{i=1}^{n} \sum_{k=1}^{K} \sum_{k'=1}^{K} z_{ik}^{*} z_{i,k'} \sum_{\ell=1}^{L_{k}} \sum_{\ell'=1}^{L_{k'}} \sum_{j \in D^{*k}_{\ell}} Y_{ij} (\mu_{k\ell}^{*} - \mu_{k'\ell'})
$$

$$
-\frac{1}{2} \sum_{k=1}^{K} \sum_{k'=1}^{K} \sum_{\ell'=1}^{L_{k}} \sum_{\ell'=1}^{L_{k'}} \mathbb{R}_{K} (z^{*}, z)_{k,k'} \mathbb{N}_{(L_{k}+1) \times (L_{k'}+1)}^{(k,k')} (T^{*}, T)_{\ell,\ell'} (\mu_{k\ell}^{*2} - \mu_{k'\ell'}^{2})
$$

We recognize the linear combination of independent Gaussian variable, then the distribution of $F_n(\theta, T, z)$ is also Gaussian. The computation of the expectation and the variance are straightforward. \Box

Proof of Proposition 9. The function $\theta \mapsto g_n(\theta, T, z)$ is maximal for

$$
\widehat{\mu}_{k'\ell'} = \frac{\sum_{k=1}^{K} \sum_{\ell=1}^{L_k} \mathbb{R}_K(\boldsymbol{z}^\star, \boldsymbol{z})_{k,k'} \mathbb{N}_{(L_k+1)\times(L_{k'}+1)}^{(k,k')}(\boldsymbol{T}^\star, \boldsymbol{T})_{\ell,\ell'} \mu_{k\ell}^\star}{\sum_{k=1}^{K} \sum_{\ell=1}^{L_k} \mathbb{R}_K(\boldsymbol{z}^\star, \boldsymbol{z})_{k,k'} \mathbb{N}_{(L_k+1)\times(L_{k'}+1)}^{(k,k')}(\boldsymbol{T}^\star, \boldsymbol{T})_{\ell,\ell'}}.
$$

Indeed, the Kullback-Leibler divergence is equal to

KL
$$
(\mu, \mu') = \frac{1}{2} (\mu - \mu')^2
$$
,

and the maximum is get by differentiating in each value. Then,

$$
\mu_{k_{1},\ell_{1}}^{\star} - \widehat{\mu}_{k\ell} = \mu_{k_{1},\ell_{1}}^{\star} - \frac{\sum_{k_{2}=1}^{K}\sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K}\left(\boldsymbol{z}^{\star},\boldsymbol{z}\right)_{k_{2},k} \mathbb{N}_{\left(L_{k_{2}}+1\right)\times\left(L_{k}+1\right)}^{(k_{2},k)} \left(\boldsymbol{T}^{\star},\boldsymbol{T}\right)_{\ell_{2},\ell} \mu_{k_{2},\ell_{2}}^{\star}}{\sum_{k_{2}=1}^{K}\sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{E}_{k_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K}\left(\boldsymbol{z}^{\star},\boldsymbol{z}\right)_{k_{2},k} \mathbb{N}_{\left(L_{k_{2}}+1\right)\times\left(L_{k}+1\right)}^{(k_{2},k)} \left(\boldsymbol{T}^{\star},\boldsymbol{T}\right)_{\ell_{2},\ell}}{\left(\mu_{k_{1},\ell_{1}}^{\star} - \mu_{k_{2},\ell_{2}}^{\star}\right)} = \frac{\sum_{k_{2}=1}^{K}\sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K}\left(\boldsymbol{z}^{\star},\boldsymbol{z}\right)_{k_{2},k} \mathbb{N}_{\left(L_{k_{2}}+1\right)\times\left(L_{k}+1\right)}^{(k_{2},k)} \left(\boldsymbol{T}^{\star},\boldsymbol{T}\right)_{\ell_{2},\ell}}{\sum_{k_{2}=1}^{K}\sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K}\left(\boldsymbol{z}^{\star},\boldsymbol{z}\right)_{k_{2},k} \mathbb{N}_{\left(L_{k_{2}}+1\right)\times\left(L_{k}+1\right)}^{(k_{2},k)} \left(\boldsymbol{T}^{\star},\boldsymbol{T}\right)_{\ell_{2},\ell}}.
$$

.

However, if ones want to take the square of the previous formulae, we get

$$
\left(\mu_{k_1,\ell_1}^{\star} - \mu_{k_2,\ell_2}^{\star}\right)\left(\mu_{k_1,\ell_1}^{\star} - \mu_{k_2',\ell_2'}^{\star}\right) = \mu_{k_1,\ell_1}^{\star^2} - \mu_{k_1,\ell_1}^{\star}\mu_{k_2,\ell_2}^{\star} - \mu_{k_2',\ell_2'}^{\star}\left(\mu_{k_1,\ell_1}^{\star} - \mu_{k_2,\ell_2}^{\star}\right),
$$

then it leads to

$$
(\mu_{k_{1},\ell_{1}}^{*} - \widehat{\mu}_{k\ell})^{2} = \frac{\sum_{k_{2}=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \sum_{\ell_{2}=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \sum_{\ell_{2}=1}^{L_{k_{2}}} \sum_{\ell_{2}=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \sum_{\ell_{2
$$

When considering $\Lambda_n(z, T)$, we are summing with respect to k_1, ℓ_1 as well, and the second term becomes 0. Then, using the explicit form of F_n given in the proof of Proposition 8, it leads to

$$
\Lambda_{n}(z, T) = -\frac{1}{2} \sum_{k=1}^{K} \sum_{\ell=1}^{L_{k}} \frac{\sum_{k_{1}=1}^{K} \sum_{\ell_{1}=1}^{L_{k_{1}}} \sum_{k_{2}=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K} (z^{\star}, z)_{k_{1},k} \mathbb{R}_{K} (z^{\star}, z)_{k_{2},k}}{\sum_{k=1}^{K} \sum_{\ell=1}^{K} \sum_{\ell_{2}=1}^{L_{k_{2}}} \mathbb{R}_{K} (z^{\star}, z)_{k_{2},k} \mathbb{N}_{(L_{k_{2}}+1)\times(L_{k}+1)}^{(k_{2},k)} (T^{\star}, T)_{\ell_{2},\ell}} \times \mathbb{N}_{(L_{k_{1}}+1)\times(L_{k}+1)}^{(k_{1},k)} (T^{\star}, T)_{\ell_{1},\ell} \mathbb{N}_{(L_{k_{2}}+1)\times(L_{k}+1)}^{(k_{2},k)} (T^{\star}, T)_{\ell_{2},\ell} \frac{1}{2} (\mu_{k_{1},\ell_{1}}^{\star} - \mu_{k_{2},\ell_{2}}^{\star})^{2}.
$$

Then, using 8, 11 and 12, we finally get the desired formulae.

\Box

Acknowledgments

 \sim

This work has been partially supported by MIAI@Grenoble Alpes (ANR-19- P3IA-0003). All the computations presented in this paper were performed using the GRICAD infrastructure (<https://gricad.univ-grenoble-alpes.fr>), which is supported by Grenoble research communities.

Supplementary Material

Fig 6*. Evolution of the coefficients grouped per day (rows) and time slots (columns) for the cluster 1. For each graphics, the evolution week per week of each curve is represented in color, the vertical black line corresponds at the estimated break-point, the horizontal black dashed line at the mean and the horizontal black dotted lines at the confident intervals. The vertical blue dashed lines correspond at the begin and the end of the both lock-down.*

Additional figures for Enedis data analysis

In this part, three additional figures (6, 7 and 8) for the section 5 are presented.

References

- ALON, J., SCLAROFF, S., KOLLIOS, G. and PAVLOVIC, V. (2003). Discovering clusters in motion time-series data. In *2003 IEEE Computer Society Conference on Computer Vision and Pattern Recognition, 2003. Proceedings.*
- Bellman, R. and Kalaba, R. (1957). Dynamic programming and statistical communication theory. *Proceedings of the National Academy of Sciences of the United States of America* **43** 749.
- Bouveyron, C. and Jacques, J. (2011). Model-based Clustering of Time Series in Group-specific Functional Subspaces. *Advances in Data Analysis and Classification* 281-300.
- Bouveyron, C., Bozzi, L., Jacques, J. and Jollois, F.-X. (2017). The Functional Latent Block Model for the Co-Clustering of Electricity Consump-

/Mixture of segmentation 36

Fig 7*. Evolution of the coefficients grouped per day (rows) and time slots (columns) for the cluster 2. For each graphics, the evolution week per week of each curve is represented in color, the vertical black lines correspond at the estimated break-points, the horizontal black dashed line at the mean and the horizontal black dotted lines at the confident intervals. The vertical blue dashed lines correspond at the begin and the end of the both lock-down.*

tion Curves. *Journal of the Royal Statistical Society: Series C Applied Statistics*.

- Bouveyron, C., Jacques, J., Schmutz, A., Simoes, F. and Bottini, S. (2021a). Co-Clustering of Multivariate Functional Data for the Analysis of Air Pollution in the South of France. *Annals of Applied Statistics*.
- Bouveyron, C., Casa, A., Erosheva, E. and Menardi, G. (2021b). Coclustering of Time-Dependent Data via the Shape Invariant Model. *Journal of Classification*.
- BRAULT, V., CHIQUET, J. and LÉVY-LEDUC, C. (2017). Efficient block boundaries estimation in block-wise constant matrices: An application to HiC data. *Electron. J. Statist.* **11** 1570-1599.
- BRAULT, V., OUADAH, S., SANSONNET, L. and LÉVY-LEDUC, C. (2018a). Nonparametric multiple change-point estimation for analyzing large Hi-C data matrices. *J. Multivar. Anal.* **165** 143–165.
- BRAULT, V., OUADAH, S., SANSONNET, L. and LÉVY-LEDUC, C. (2018b). Nonparametric multiple change-point estimation for analyzing large Hi-C data matrices. *Journal of Multivariate Analysis* **165** 143–165.
- BRAULT, V., KERIBIN, C., MARIADASSOU, M. et al. (2020). Consistency and

Fig 8*. Evolution of the coefficients grouped per day (rows) and time slots (columns) for the cluster 3. For each graphics, the evolution week per week of each curve is represented in color, the vertical black lines correspond at the estimated break-points, the horizontal black dashed line at the mean and the horizontal black dotted lines at the confident intervals. The vertical blue dashed lines correspond at the begin and the end of the both lock-down.*

asymptotic normality of Latent Block Model estimators. *Electronic journal of statistics* **14** 1234–1268.

- BUGNI, F. A., HALL, P., HOROWITZ, J. L. and NEUMANN, G. R. (2009). Goodness-of-fit tests for functional data. *The Econometrics Journal* **12** S1– S18.
- Chamroukhi, F. (2016). Piecewise Regression Mixture for Simultaneous Functional Data Clustering and Optimal Segmentation. *Journal of Classification* **33** 374-411.
- DEMPSTER, A. P., LAIRD, N. M. and RUBIN, D. B. (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)* **39** 1–22.
- Devijver, E. (2017). Model-Based Regression Clustering for High-Dimensional Data: Application to Functional Data. *Advances in Data Analysis and Classification* **11** 243–279.
- Droesbeke, J.-J., Saporta, G. and Thomas-Agnan, C. (2013). *Modèles à variables latentes et modèles de mélange*. Editions TECHNIP.
- FERRATY, F. and VIEU, P. (2006). *Nonparametric Functional Data Analysis. Theory and Practice*. *Springer Series in Statistics*. Springer.
- GALVANI, M., TORTI, A., MENAFOGLIO, A. and VANTINI, S. (2021). FunCC: A new bi-clustering algorithm for functional data with misalignment. *Computational Statistics & Data Analysis* **160** 107219.
- GIACOFCI, M., LAMBERT-LACROIX, S., MAROT, G. and PICARD, F. (2013). Wavelet-Based Clustering for Mixed-Effects Functional Models in High Dimension. *Biometrics* **69** 31–40.
- Hubert, L. and Arabie, P. (1985). Comparing partitions. *Journal of classification* **1** 193–218.
- Hébrail, G., Hugueney, B., Lechevallier, Y. and Rossi, F. (2010). Exploratory analysis of functional data via clustering and optimal segmentation. *Neurocomputing* **73** 1125-1141. Advances in Computational Intelligence and Learning.
- JACQUES, J. and PREDA, C. (2013). Funclust: A curves clustering method using functional random variables density approximation. *Neurocomputing* **112** 164-171. Advances in artificial neural networks, machine learning, and computational intelligence.
- JACQUES, J. and PREDA, C. (2014). Functional Data Clustering: A Survey. *Advances in Data Analysis and Classification* **8** 231-255.
- JAMES, G. M. and SUGAR, C. A. (2003). Clustering for Sparsely Sampled Functional Data. *Journal of the American Statistical Association* **98** 397-408.
- Kay, S. M. (1993). *Fundamentals of statistical signal processing*. Prentice Hall PTR.
- Liu, X. and Yang, M. C. K. (2009). Simultaneous curve registration and clustering for functional data. *Computational Statistics & Data Analysis* **53** 1361-1376.
- MAIDSTONE, R., HOCKING, T., RIGAILL, G. and FEARNHEAD, P. (2017). On optimal multiple changepoint algorithms for large data. *Statistics and computing* **27** 519–533.
- Mallat, S. (1989). A theory for multiresolution signal decomposition: the wavelet representation. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **11** 674–693.
- Mallat, S. (1999). *A wavelet tour of signal processing.* Academic Press.
- Misiti, M., Misiti, Y., Oppenheim, G. and Poggi, J.-M. (2004). Matlab Wavelet Toolbox User's Guide. Version 3. The Mathworks, Inc., Natick, MA.
- Ramsay, J. and Silverman, B. (2002). *Applied Functional Data Analysis. Methods and Case Studies*. Springer.
- Ramsay, J. O. and Silverman, B. W. (2005). *Functional Data Analysis*. Springer.
- Rigaill, G. (2015). A pruned dynamic programming algorithm to recover the best segmentations with 1 to *K*max change-points. *Journal de la Société Française de Statistique* **156** 180–205.
- ROBERT, V. (2021). bikm1: Co-Clustering Adjusted Rand Index and Bikm1 Procedure for Contingency and Binary Data-Sets R package version 1.1.0.
- ROBERT, V., VASSEUR, Y. and BRAULT, V. (2021). Comparing High-Dimensional Partitions with the Co-Clustering Adjusted Rand Index. *J. Classif.* **38** 158–186.
- SAMÉ, A. and GOVAERT, G. (2012). Online Time Series Segmentation Using Temporal Mixture Models and Bayesian Model Selection. In *2012 11th International Conference on Machine Learning and Applications* **1** 602-605.
- SAMÉ, A., CHAMROUKHI, F., GOVAERT, G. and AKNIN, P. (2011). Modelbased clustering and segmentation of time series with changes in regime. *Advances in Data Analysis and Classification* **5** 301-321.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics* **6** 461–464.
- ZHANG, N. R. and SIEGMUND, D. O. (2007). A modified Bayes information criterion with applications to the analysis of comparative genomic hybridization data. *Biometrics* **63** 22–32.