



**HAL**  
open science

## Scavenger receptor class B type I genetic variants associated with disease severity in chronic hepatitis C virus infection

Victoria L Arandhara, Charles Patrick McClure, Alexander W Tarr, Sally Chappell, Kevin Morgan, Thomas F Baumert, William L Irving, Jonathan K Ball

### ► To cite this version:

Victoria L Arandhara, Charles Patrick McClure, Alexander W Tarr, Sally Chappell, Kevin Morgan, et al.. Scavenger receptor class B type I genetic variants associated with disease severity in chronic hepatitis C virus infection. *Journal of Medical Virology*, 2023, 95 (1), 10.1002/jmv.28331 . hal-04036854

**HAL Id: hal-04036854**

**<https://hal.science/hal-04036854>**

Submitted on 20 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Precise. Reliable.  
Sleek and  
Squeaky Clean.

Ensure excellence with PHCbi brand products.



**PHCbi**

Life Science Innovator Since 1966

# Scavenger receptor class B type I genetic variants associated with disease severity in chronic hepatitis C virus infection

Victoria L. Arandhara<sup>1</sup> | Charles Patrick McClure<sup>1,2,3</sup>  | Alexander W. Tarr<sup>1,2,3</sup> | Sally Chappell<sup>1,†</sup> | Kevin Morgan<sup>1</sup> | Thomas F. Baumert<sup>4,5</sup> | William L. Irving<sup>1,2,3</sup> | Jonathan K. Ball<sup>1,2,3</sup>

<sup>1</sup>School of Life Sciences, The University of Nottingham, Queen's Medical Centre, Nottingham, UK

<sup>2</sup>Wolfson Centre for Global Virus Research, The University of Nottingham, Queen's Medical Centre, Nottingham, UK

<sup>3</sup>NIHR Nottingham Biomedical Research Centre at the Nottingham University Hospitals NHS Trust and University of Nottingham, Nottingham, UK

<sup>4</sup>Institut de Recherche sur les Maladies Virales et Hépatiques, Université de Strasbourg, Inserm, Strasbourg, France

<sup>5</sup>IHU Strasbourg, Pôle hépato-digestif, Hôpitaux Universitaires de Strasbourg, Strasbourg, France

## Correspondence

Jonathan K. Ball, School of Life Sciences, The University of Nottingham, Queen's Medical Centre, Nottingham NG7 2UH, UK.  
Email: [jonathan.ball@nottingham.ac.uk](mailto:jonathan.ball@nottingham.ac.uk)

## Funding information

European Commission, Grant/Award Numbers: QLK2-CT-2001-01120, MRTN-CT-2006-035599, HEALTH-F4-2012-305600; NIHR Nottingham Biomedical Research Centre, Grant/Award Number: IS-BRC-1215-20003; European Research Council, Grant/Award Number: ERC-AdG-2020-FIBCAN#101021417; Medical Research Council, Grant/Award Number: MR/S009434/1; IdEx Unistra, Grant/Award Number: ANR-10-IDEX-0002; SFRI-STRATUS, Grant/Award Number: ANR20-SFRI-0012; Special Trustees for Nottingham University Hospital, Grant/Award Number: STR64/03/N; University of Nottingham Research Committee

## Abstract

Analysis of host genetic polymorphisms is an increasingly important tool for understanding and predicting pathogenesis and treatment response of viral diseases. The gene locus of scavenger receptor class B type I (SR-BI), encoding a cell entry factor and receptor for hepatitis C virus (HCV), contains several genetic polymorphisms. We applied a probe extension assay to determine the frequency of six single nucleotide polymorphisms (SNPs) within the SR-BI gene locus in 374 individuals with history of HCV infection. In addition, SR-BI messenger RNA (mRNA) levels were analyzed in liver biopsy specimens of chronically infected HCV subjects. The rs5888 variant allele T was present at a higher frequency in subjects with advanced fibrosis ( $\chi^2$ ,  $p = 0.016$ ) and after adjusting for age, duration of infection and alcohol intake as confounding factors. Haplotype analysis of SNP frequencies showed that a haplotype consisting of rs61932577 variant allele C and rs5888 variant allele T was associated with an increased risk of advanced liver fibrosis (defined by an Ishak score 4–6) (adjusted odds ratio 2.81; 95% confidence interval 1.06–7.46.  $p = 0.038$ ). Carriers of the rs5888 variant allele T displayed reduced SR-BI mRNA expression in liver biopsy specimens. In conclusion the rs5888 polymorphism variant is associated with decreased SR-BI expression and an increased risk of development of advanced fibrosis in chronic HCV infection. These findings provide further evidence for a role of SR-BI in HCV pathogenesis and provides a genetic marker for prediction of those infected individuals at greater risk of developing severe disease.

†Died. June 16, 2017.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Journal of Medical Virology* published by Wiley Periodicals LLC.

## KEYWORDS

HCV, liver disease, rs5888, scavenger receptor class B type 1, single nucleotide polymorphisms, SR-BI

## 1 | INTRODUCTION

The consequences of infection with hepatitis C virus (HCV) differ considerably between individuals and the spectrum of liver disease in chronic HCV infection can vary widely. Host factors such as age, gender, duration of infection, alcohol consumption, and route of transmission may influence disease progression.<sup>1–5</sup> There is also evidence that host genetic factors influence the course of infection and development of liver disease,<sup>6,7</sup> the most noticeable being the influence of genetic variation within the interferon lambda locus on infection and treatment outcome (reviewed in).<sup>8,9</sup> Additional polymorphisms within genes encoding cell surface proteins, including low density lipoprotein receptor (LDL-R)<sup>10</sup> and apolipoprotein E<sup>11</sup> are associated with severity of fibrosis, viral clearance, overall inflammation or treatment response.<sup>10</sup> The scavenger receptor class B type I (SR-BI), which is primarily expressed in the liver and in steroidogenic tissues,<sup>12–15</sup> serves as one of several host surface molecules involved in HCV cell entry.<sup>16</sup> Its exact role in entry has not been fully elucidated, but direct binding to the E2 envelope protein is thought to facilitate subsequent binding to the major receptor, CD81,<sup>17–19</sup> and SR-BI-specific monoclonal antibodies inhibit HCV infection in cell-based and animal models.<sup>20,21</sup>

SR-BI plays an important role in regulating plasma HDL levels and is a major determinant of HDL cholesterol levels.<sup>22–26</sup> In selective lipid uptake, HDL binds to the cells but only the lipid part of the particle enters the cells via an unknown mechanism that does not require endocytosis of the lipoprotein particle.<sup>27</sup> The SR-BI gene locus contains several genetic polymorphisms<sup>23</sup> and there is evidence that these are associated with plasma lipid profiles and the development of atherosclerosis and coronary artery disease.<sup>6,23,28–40</sup> Previous studies have shown that some common SR-BI polymorphisms are associated with differential HCV treatment outcome and viral load.<sup>41,42</sup> In addition, naturally occurring, but very rare SR-BI coding mutations (specifically S112F and T175A) have reduced HCV entry function.<sup>42–44</sup> Despite the association between SR-BI polymorphisms and key facets of HCV replication and biology, their potential impact on disease severity has not been defined. To address this shortfall, we used archived DNA extracts obtained from the extensively characterized Trent HCV cohort<sup>45</sup> to investigate the prevalence of relatively high frequency polymorphisms in individuals with severe versus mild HCV-induced liver disease. We also defined the relationship between SR-BI polymorphisms and SR-BI messenger RNA (mRNA) expression in liver biopsies from chronically infected HCV subjects using real-time polymerase chain reaction (PCR).

## 2 | PATIENTS AND METHODS

### 2.1 | Human subjects

Archived samples from patients with a history of HCV infection were obtained from the Trent HCV study cohort, which was established in 1991 with the aim of studying the epidemiology and natural history of HCV infection.<sup>1,4,46</sup> All patients referred to one of the participating clinics were invited to enroll in the cohort study with informed consent following approval by the local ethics committee. HCV antibody positive subjects (third-generation enzyme-linked immunosorbent assays, Ortho; Sanofi) were divided, based on historic diagnostic testing protocols, into those who were negative (acute resolved infection) or positive (chronic infection) for HCV RNA (Amplicor reverse-transcription PCR assay; Roche). Diagnostic liver biopsy specimens were scored using the Ishak modified histological activity index<sup>47</sup> and grouped into severe liver disease, defined by the presence of advanced fibrosis (Ishak fibrosis stage 4–6), and mild liver disease (Ishak fibrosis stage 0–3). Following the Trent Hepatitis C Cohort Study assessment and management protocol<sup>1</sup> patients are offered repeat liver biopsies at 2-year intervals. Subsequently several patients in this study had undergone more than one liver biopsy. For this study, the highest fibrosis score achieved was recorded irrespective of whether this was on a first or subsequent biopsy. Risk factors for acquisition of the virus were categorized as injecting drug use, blood product transfusion, other known risk factor (e.g., tattoo and body piercing) and no known risk factor. The duration of infection at the time of biopsy was estimated assuming that the first exposure to risk was the date of infection. At each clinical visit the average weekly alcohol intake since the previous visit was recorded. Clinical, ApoE and interferon lambda genotype data for some of these patients have previously been described.<sup>4,11,48</sup>

### 2.2 | Genotyping of SR-BI polymorphisms

To assess the possible relationship between SR-BI polymorphisms and the outcome of infection with HCV, four single nucleotide polymorphisms (SNPs) present within the gene locus were selected, based on previous evidence of an association with either lipid metabolism and/or HCV outcomes.<sup>23,33,36,39,41,42,49</sup> In addition, two control SNPs, where no associations with disease had been reported, were included in the analysis (Table 1). DNA was extracted from 200 µl of either whole blood or plasma using Qiagen DNA blood mini kit according to the manufacturer's protocol. Oligonucleotide primer pairs (Table 2) were derived

**TABLE 1** SR-BI SNPs under study

SNP	Alleles	Highest minor allele frequency <sup>a</sup>	cDNA position	Amino acid	Phenotype	Reference
rs4238001	G > A	6%	4	G2S	Increased MI risk; Decreased HDL/Elevated LDL (in males); Decreased SR-BI protein expression	23,36,39,49
rs5889	C > T	11%	502	G167G	Elevated HDL; Decreased coronary artery disease;	70
rs5892	C > T	4%	905	F301F	-	
rs5888	C > T	32%	1050	A350A	Decreased LDL/Increased BMI (in women);	23,37,40
rs61932577	C > T	14%	-	Intron	-	
rs2293440	T > C	24%	-	Intron	Altered HDL levels	71

<sup>a</sup>Minor allele frequency as recorded in the 1000 Genomes Project Phase 3, Ensembl database.<sup>46</sup>

**TABLE 2** Characteristics of patients grouped by HCV status and Ishak fibrosis stage

	HCV RNA		HCV/Ishak Fibrosis	
	Negative (resolved) n = 107	Positive (chronic) n = 267	Mild (0–3) n = 160	Severe (4–6) n = 92
Gender				
Male	67 (63%)	183 (69%) <sup>a</sup>	111 (69%)	63 (69%) <sup>a</sup>
Female	38 (35%)	81 (30%)	49 (31%)	26 (28%)
Unknown	2 (2%)	3 (1%)	0 (0%)	3 (3%)
Risk factor				
Injecting drug user	40 (37%)	151 (56%)	106 (66%)	36 (39%)
Blood product transfusion	4 (4%)	45 (17%)	27 (17%)	15 (16%)
Other known risk factor	4 (4%)	15 (6%)	6 (4%)	9 (10%)
No known risk factor	1 (1%)	33 (12%)	14 (9%)	17 (19%)
No risk factor data	58 (54%) <sup>b</sup>	23 (9%)	7 (4%)	15 (16%)
Mean (SD) age at Biopsy (year)			39 (10)	50 (11) <sup>c</sup>
Mean (SD) duration of Infection (year)			17 (8)	23 (10) <sup>c</sup>
Median weekly alcohol intake (range)			4 (0–65)	0 (0–60) <sup>d</sup>

<sup>a</sup>NS ( $\chi^2$  test);

<sup>b</sup>Risk factor data was not routinely collected from HCV RNA negative patients;

<sup>c</sup> $p < 0.001$  (unpaired *t*-test);

<sup>d</sup> $p = 0.001$  (Mann–Whitney *U* test).

from a previous study<sup>23</sup> or mapped to human contiguous sequences (Genbank Accession AC126309 and AC073593) flanking each SNP of interest. Primers were designed to amplify a 200–300 nucleotide region surrounding the SNP of interest. Aliquots of DNA (1  $\mu$ l) were used as templates in PCR reactions containing 0.2  $\mu$ M of each primer pair, 200  $\mu$ M dinucleotide triphosphates (dNTPs) and 0.625 U Hot Star Taq (Qiagen,) in 1  $\times$  PCR buffer in a 25  $\mu$ l reaction volume. The amplification cycle parameters were an initial denaturation step of 94°C for 10 min followed by 35 cycles of denaturation (94°C, 50 s), annealing (55°C, 30 s) and extension (72°C, 1 min) and an extended elongation step (72°C, 5 min). PCR products were purified by

treatment with shrimp alkaline phosphatase (SAP and GE Healthcare) and exonuclease I (Exo I, New England Biolabs) to remove unincorporated dNTPs and primers.

Allelic composition was determined using the commercially available ABI PRISM Multiplex SNaPshot Reaction kit (ThermoFisher). SNaPshot primers (Supporting Information: Table 1) were designed adjacent to the SNP of interest and 5' polyA tails added to produce primers of differing lengths to allow discrimination of individual SNPs in a multiplex reaction. The SNaPshot reaction was prepared according to the manufacturer's protocol. Samples were resolved using an Applied Biosystems Genetic analyzer according to the manufacturer's protocol.



## 2.3 | Sequencing of PCR products

Each allelic variant was confirmed by direct sequencing of PCR products that were used in the SNP analysis using the ABI Prism Dye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems) according to the manufacturer's protocol. Samples were selected for sequencing at the first detection instance of each SNP genotype.

## 2.4 | RNA extraction from liver biopsies

Percutaneous needle liver biopsies were obtained from subjects attending the liver clinic at Lincoln City Hospital between 1999 and 2003, consented to join the HCV Trent Study cohort.<sup>1,4</sup> Biopsies were snap frozen in liquid nitrogen and nucleic acids using TRIzol reagent (Invitrogen) according to the manufacturer's guidelines. 1 ml TRIzol was added to the frozen liver biopsies and the tissue homogenized using a Ribolyser (Bio-Rad) at speed 6 for 40 s, repeating until the biopsy was homogenized, before nucleic acid extraction. Reverse transcription was performed using ThermoScript RT-PCR System (ThermoFisher) according to the manufacturer's guidelines using random hexamers for priming.

## 2.5 | Quantitative real-time PCR

Investigation of mRNA copy numbers in liver biopsy specimens was performed using Brilliant SYBR Green Quantitative PCR Master Mix (ThermoFisher) according to the manufacturer's instructions. HPRT was amplified as a reference gene using the following primers: HPRTA, 5' GACCAGTCAACAGGGGACAT 3' and HPRTB 5' CGACCTTGACCATCTTTGGA 3'. SR-BI mRNA was amplified using primers SR-BIF 5' GAAACTGCAGCTGAGCCTCT 3' and SR-BIR 5' ATTCTCTTGGCTCCGGATT 3'. A standard curve of cycle threshold ( $C_t$ ) plotted against a standard of known copy number was used to interpolate the copy number of unknown samples using the  $C_t$  for each sample. Relative copy numbers of the reference gene HPRT and SR-BI were calculated for each unknown sample for comparison.

## 2.6 | Statistical analyses

SPSS version 12.0.1 was used for statistical analyses. Variables were compared using the unpaired *t*-test for two independent groups for normally distributed variables and the Mann-Whitney *U* test for non-normally distributed variables. SR-BI genotype allele frequencies were estimated by allele counting and deviations from Hardy-Weinberg equilibrium were assessed for the whole cohort. The statistical significance of associations between genotype and total allele frequency distribution for each SNP between patient groups was assessed using  $\chi^2$ -tests. Logistic regression analysis was performed to adjust for confounding factors. A *p* value of <0.05

was considered statistically significant. Haplotype analysis and linkage disequilibrium (LD) analysis was performed using the COCAPHASE software utility of the UNPHASED suite of programs for association analysis for multilocus haplotypes (<https://gaow.github.io/genetic-analysis-software/u/unphased/>)<sup>50</sup> and using the Ensembl Genome Browser.<sup>46</sup> To compare the SR-BI mRNA levels in liver biopsy samples the number of copies of SR-BI per HPRT mRNA copy was calculated by dividing the number of copies of SR-BI by the number of copies of the reference gene HPRT. The copies of SR-BI per HPRT were compared between genotypes for rs4238001, rs61932577 and rs5888 and the significance of differences between the median values determined using Kruskal-Wallis test for three or more groups and the Mann-Whitney *U* tests for two groups. Analyses were performed using GraphPad Prism version 9 for Windows (GraphPad Software). To assess inter- and intra-assay variation the coefficient of variation (CV) of  $C_t$  values was calculated as (standard deviation/mean) x 100.<sup>51</sup> The CV for HPRT  $C_t$  values was less than 3.21% for standards (range 0.81% to 3.21%) and less than 7.7% for unknown samples (range 0%–7.7%). For SR-BI the CV of the  $C_t$  values was less than 11.24% for standards (range 0.95%–11.24%) and less than 8.94% for unknown samples (range 0.37%–8.94%). CVs for intra-assay variation from two intra-assay repeats were below 7% for both HPRT and SR-BI unknown samples and less than 2% for standard  $C_t$  values.

## 3 | RESULTS

### 3.1 | Patient characteristics

A total of 374 subjects with a history of HCV infection were studied. Of these, 107 (28.6%) had acute resolved infection and 267 (71.4%) had chronic infection. Histological assessment of liver biopsies was available for 252 chronically infected subjects of which 92 (36.5%) had severe liver disease (Ishak fibrosis stage 4–6) and 160 (63.5%) had mild liver disease (Ishak fibrosis stage 0–3). Characteristics of patients grouped by HCV status and Ishak fibrosis stage are given in Table 2. Several host factors may influence the degree of liver fibrosis in chronic infection with HCV including gender, alcohol intake and duration of infection.<sup>3</sup> No associations were found between gender and HCV status ( $\chi^2$  test, *p* = 0.33) or severity of liver disease ( $\chi^2$  test, *p* = 0.89). The duration of infection was estimated for 187 subjects from whom risk factors were known and was estimated from the first exposure to that risk factor. The mean estimated duration of infection and mean age at the time of biopsy procedure were higher in the subjects with severe liver disease (unpaired *t*-tests, *p* < 0.001). The median average weekly alcohol intake (units) since the previous clinical visit was higher in the group of subjects with mild liver disease compared to those with severe liver disease (*n* = 252, Mann-Whitney *U*, *p* = 0.001). The cohort was infected with genotypes 1–5, with genotypes 1 and 3 the most prevalent in both groups (Data not shown). Genotype prevalence did not differ significantly between the groups ( $\chi^2$  test, *p* = 0.122).

### 3.2 | Genotyping and distribution of SR-BI polymorphisms

To determine possible associations with SR-BI polymorphisms and the outcome of HCV infection, the allele frequencies at six polymorphic sites were assessed. Four of the SNPs were in exon sequences and two in introns (Table 1). Allele counting revealed that the frequency of minor allelic variants defined by rs5889, rs2293440, and rs5892 were 0.003, 0.011, and 0.003 respectively. Since the frequencies of these three SNPs were so low in our cohort they were excluded from further analyses. The frequencies of the minor allelic variant for each of the remaining SNPs within the whole cohort were 0.111, 0.099, and 0.473 for rs4238001, rs61932577, and rs5888, respectively. All three SNPs were in Hardy–Weinberg equilibrium. Allele frequencies did not differ statistically by gender.

Table 3 shows the frequencies of the SR-BI SNP genotypes and alleles grouped by HCV RNA status and Ishak fibrosis stage for chronically infected HCV subjects. LD analysis indicated that rs4238001 was not linked to either rs61932577 ( $D' = 0.075$ ,  $r^2 = 0.005$ ) or rs5888 ( $D' = 0.053$ ,  $r^2 = 0.0004$ ). Rs61932577 was found to be linked to rs5888 using the  $D'$  parameter of LD

( $D' = 0.766$ ). However, since the allele frequencies of rs61932577 and rs5888 differ widely the  $r^2$  parameter has more desirable statistical properties,<sup>52</sup> and this value does not indicate linkage between the two SNPs ( $r^2 = 0.059$ ).

For subjects with a history of HCV infection, there were no differences observed in the frequencies of rs4238001 and rs61932577 variants between subjects who had cleared the virus versus subjects with chronic infection or between subjects with severe versus mild liver disease. The rs5888 variant allele T was higher in HCV RNA positive subjects (48.1%) compared to those who had cleared the virus (40.6%) but this difference was not statistically significant ( $\chi^2$  test,  $p = 0.06$ ). Genotype frequencies for the rs5888 SNP differed between chronically infected HCV subjects with severe and mild liver disease ( $\chi^2$  test,  $p = 0.039$ ) and the variant allele T was found at a higher frequency in subjects with severe liver disease compared to those with mild liver disease ( $\chi^2$  test,  $p = 0.016$ ). Since the frequency of rs5888 SNP heterozygotes was similar between the two groups of chronically infected subjects, analysis of homozygous genotypes only was performed. The frequency of homozygous genotype TT was higher in subjects with severe liver disease compared to subjects with mild liver disease ( $\chi^2$  test,  $p = 0.014$ ).

**TABLE 3** SR-BI SNP genotypes and allele frequencies

SRB1 genotype		HCV RNA		HCV Ishak fibrosis Severe <i>n</i> = 92	HCV Ishak fibrosis Mild <i>n</i> = 160
		Positive <i>n</i> = 267	Negative <i>n</i> = 107		
rs4238001	GG	211 (79.0)	85 (79.4)	70 (76.1)	130 (81.3)
	GA	52 (19.5)	20 (18.7)	22 (23.9)	27 (16.9)
	AA	4 (1.5)	2 (1.9)	0 (0.0)	3 (1.9)
rs61932577	CC	217 (81.3)	80 (74.8)	74 (80.4)	129 (80.6)
	CT	48 (18.0)	27 (25.2)	17 (18.5)	30 (18.8)
	TT	2 (0.7)	0 (0.0)	1 (1.1)	1 (0.6)
rs5888	CC	70 (26.2)	37 (34.6)	19 (20.7)	48 (30.0) <sup>a,b</sup>
	CT	137 (51.3)	53 (49.5)	45 (48.9)	84 (52.9)
	TT	60 (22.5)	17 (15.9)	28 (30.4)	28 (17.5)
SRB1 alleles		<i>N</i> = 534	<i>N</i> = 214	<i>N</i> = 184	<i>N</i> = 320
rs4238001	G	474 (88.8)	190 (88.8)	162 (88.0)	287 (89.7)
	A	60 (11.2)	24 (11.2)	22 (12.0)	33 (10.3)
rs61932577	C	482 (90.3)	187 (87.4)	165 (89.7)	288 (90.0)
	T	52 (9.7)	27 (12.6)	19 (10.3)	32 (10.0)
rs5888	C	277 (51.9)	127 (59.4)	83 (45.1)	180 (56.3) <sup>c</sup>
	T	257 (48.1)	87 (40.6)	101 (54.9)	140 (43.7)

Abbreviations: *n*, number of subjects; *N*, number of alleles; percentages are given in brackets.

<sup>a</sup>Genotype versus disease severity  $p = 0.039$  ( $\chi^2$  test);

<sup>b</sup>Homozygous genotype versus disease severity  $p = 0.014$  ( $\chi^2$  test);

<sup>c</sup>Allele frequency versus disease severity  $p = 0.016$  ( $\chi^2$  test).

Logistic regression analysis was performed to adjust for gender, age at biopsy, estimated duration of infection, and alcohol intake as confounding factors. Since the duration of infection was estimated from the date of exposure to risk, this excluded 56 patients with missing risk factor data from the analysis. Unadjusted and adjusted odds ratios are given in Table 4. There was a significant association between the severity of liver disease and the rs5888 genotype TT, compared to genotype CC, after adjusting for the confounding factors with an adjusted odds ratio of 3.32 (95% confidence interval (CI) 1.05–10.53,  $p = 0.041$ ).

Haplotype analysis was performed to increase the statistical power of single locus analyses. All combinations of haplotypes of rs4238001, rs61932577 and rs5888 SNPs were analyzed. Haplotypes occurring at a frequency of 0.10 or more in both severe and mild HCV subjects are shown in Table 5. Haplotype rs4238001/rs61932577/rs5888 G/C/C showed the most significant association with disease severity and was present at higher frequencies in subjects with mild liver disease compared to severe liver disease ( $p = 0.009$ ). When the rs4238001 SNP was excluded from the haplotype, the frequency of rs61932577/rs5888 haplotype C/C was higher in patients with mild liver disease ( $p = 0.012$ ) and haplotype C/T was found more frequently in subjects with severe liver disease ( $p = 0.015$ ). rs4238001/rs5888 haplotype G/C was observed at a higher frequency in subjects with mild liver disease compared to severe liver disease ( $p = 0.018$ ). Subjects who are heterozygous at multiple SNP loci cannot be unequivocally assigned a haplotype therefore subjects homozygous for SNP haplotypes were compared between the severe and mild liver disease groups using logistic regression. Since haplotypes containing the rs5888 SNP were consistently associated with disease severity only haplotypes containing rs5888 were compared. Unadjusted and adjusted odds ratios and CIs are shown in Table 6. After adjusting for gender, age at biopsy, estimated duration of infection and alcohol intake, subjects who were homozygous for both rs61932577 C and rs5888 T showed the most significant association with severity of liver disease compared to other haplotypes with an adjusted odds ratio of 2.81 (95% CI 1.06–7.46,  $p = 0.038$ ).

Finally, since previous studies have reported gender-specific effects of SR-BI polymorphisms,<sup>31,32</sup> analysis of the cohort split by gender was undertaken. When the data obtained for the male patients was analyzed separately, a stronger association of the rs5888 polymorphism with disease severity was observed compared to the analysis for the full cohort

( $n = 174$ ,  $\chi^2$  test,  $p = 0.011$ ). The rs5888 allele T was found at a higher frequency in male subjects with severe liver disease (55%) compared to those with mild liver disease (40%) ( $\chi^2$  test,  $p = 0.006$ ). By contrast, analysis of the association of the rs5888 allele on disease severity in females did not reveal any significant associations, with T allele frequencies of 54% and 53% in those with severe or mild disease, respectively ( $n = 75$ ,  $\chi^2$  test,  $p = 0.927$ ). Using logistic regression analysis to account for age at biopsy, estimated duration of infection and alcohol intake as confounding factors, carriage of rs5888 genotype TT was associated with an increased risk of severe disease compared to genotype CC in men, with an odds ratio of 4.38 (95% CI 1.05–18.27,  $p = 0.043$ ). Similarly, haplotype analysis showed that the association of the haplotype G/C/C with disease severity was greater for the male data ( $p = 0.004$ ) (Supporting Information: Table 2) than for the whole cohort.

### 3.3 | SR-BI polymorphisms and haplotypes are associated with differing SR-BI mRNA expression

The SR-BI polymorphism associated with disease severity was a synonymous nucleotide substitution. To assess whether this polymorphism was exerting an effect through altered gene

**TABLE 5** Haplotype frequencies subjects grouped by HCV liver disease severity

SNP Haplotype			Frequency	
rs4238001	rs61932577	rs5888	Severe	Mild
G	C	C	0.31	0.43**
		C	0.35	0.48**
	T	0.55	0.42*	
G		C	0.40	0.50*
G	C	T	0.48	0.39*
G		T	0.49	0.40*
G	C		0.79	0.81

Note: haplotypes present at a frequency of at least 0.10 in both severe and mild liver disease subjects are shown. Haplotypes are listed in order of decreasing association.

\*\* $p < 0.01$ ; \* $p \leq 0.05$ .

rs5888 Genotype	Unadjusted				Adjusted <sup>a</sup>			
	OR	95% CI		$p$	OR	95% CI		$p$
		Lower	Upper			Lower	Upper	
CC	1.00				1.00			
CT	1.35	0.71	2.57	0.356	1.52	0.60	3.93	0.384
TT	2.53	1.20	5.33	0.015	3.32	1.05	10.53	0.041

Note: exon 8 genotype frequencies are from 132 patients with mild and 46 subjects with severe liver disease.

<sup>a</sup>Adjusted for gender, age at biopsy, estimated duration of infection and alcohol intake.

**TABLE 4** Logistic regression analysis of rs5888 genotype frequencies and the severity of liver disease




**TABLE 6** Logistic regression analysis of frequency of homozygous haplotypes and the severity of liver disease

Haplotype <sup>a</sup>				Unadjusted				Adjusted <sup>b</sup>			
rs4238001	rs61932577	rs5888	n	OR	95% CI		p	OR	95% CI		p
					Lower	Upper			Lower	Upper	
G	C	C	38	0.58	0.27	1.25	0.16	0.64	0.21	2.02	0.45
		C	47	0.54	0.26	1.09	<b>0.09</b>	0.46	0.16	1.37	0.17
		T	54	2.26	1.22	4.16	<b>0.009</b>	2.81	1.06	7.46	<b>0.038</b>
G		C	51	0.60	0.30	1.18	0.14	0.74	0.29	1.86	0.52
G	C	T	40	2.20	1.10	4.35	<b>0.024</b>	2.37	0.78	7.20	0.13
G		T	42	1.96	1.00	3.82	<b>0.049</b>	1.91	0.65	5.60	0.24

Abbreviations: CI, confidence interval; n, number of subjects with homozygous haplotype; OR, odd ratio.

<sup>a</sup>Homozygous haplotype versus all other haplotypes present;

<sup>b</sup>Adjusted for gender, age at biopsy, estimated duration of infection and alcohol.



**FIGURE 1** mRNA copies of SR-BI per HPRT for SR-BI variant rs5888 genotypes and haplotype. Log scale expression of SR-BI per HPRT for rs5888 genotypes CC, CT and CC (A) and rs61932577/rs5888 homozygous haplotypes; C/T and C/C (B). Lines represent the median of the data set and data are representative of at least 2 repeats for each data point. \* $p < 0.05$ , \*\* $p < 0.01$ . Statistical comparisons performed were the Kruskal–Wallis ANOVA test with Dunn's multiple comparisons tests (A) or the Mann–Whitney test (B). ANOVA, analysis of variance.

transcription SR-BI RNA expression levels were determined in freshly obtained liver biopsies.  $C_t$  values were plotted against starting copy number to construct a standard curve from which to interpolate copy numbers of unknown samples. Using the standard curves of SR-BI and HPRT the number of copies of SR-BI and HPRT in the liver biopsy samples was determined and SR-BI per HPRT RNA copy number plotted according to the SNP genotype and haplotype (Figure 1). Analysis of the median copies of SR-BI per HPRT between rs5888 genotypes showed that rs5888 genotype CC was associated with twofold more SR-BI expression compared to genotype TT ( $n = 17$ , Dunn's post-test following Kruskal–Wallis test,  $p = 0.048$ ). Since a SR-BI homozygous haplotype consisting of rs61932577 C and rs5888 T was previously found to be associated with increased liver disease severity in chronically infected HCV subjects, analysis of SR-BI mRNA levels grouped by the same homozygous haplotype was performed. Carriers of the homozygous intron 5/exon8 haplotype C/C had a twofold higher expression of copies of SR-BI per HPRT compared with C/T ( $n = 15$ , Mann–Whitney  $U$  test,  $p = 0.008$ ).

## 4 | DISCUSSION

The clinical outcome of HCV infection is variable and remains poorly understood. Susceptibility to infection or disease progression may be influenced by the host genetic background, and polymorphisms within several genes have been associated with differing disease progression.<sup>6,10,11,53–55</sup> Genetic polymorphisms within the SR-BI gene locus have been associated with differing plasma lipid profiles and risk of atherosclerosis and coronary artery disease,<sup>23,28,30–32,35,38,56,57</sup> however the impact of SR-BI allelic variation on outcome of HCV infection is still only partially understood. Since SR-BI is a putative receptor for HCV entry,<sup>17</sup> we performed the present study to define possible associations between SR-BI polymorphisms with HCV disease. Genotyping was performed using a probe extension assay, which has previously been shown to produce comparable data to other genotyping methods.<sup>58,59</sup> Genotyping errors were further minimized by direct sequencing of a subset of PCR products used in the multiplex genotyping analysis. The minor allelic variants of rs5889, rs2293440 and rs5892 were rare in our

population. Whilst this is at odds with reported global minor variant allele frequencies, this discrepancy can be explained by the ethnic composition of our cohort. For example, the global minor allele frequency (MAF) for rs5889 is 11%, whereas the frequency in our cohort was only 0.3%, although this latter value accords with the MAF frequency reported in for the European Caucasian 1000 genome project (phase 3) data set reported in the Ensembl database.<sup>46</sup> Total allele frequencies for the SNPs in rs4238001, rs61932577, and rs5888 were similar to those reported elsewhere.<sup>23</sup> Previous studies have reported linkage between the rs61932577 and rs5888 alleles.<sup>23,32</sup> A similar analysis of our data performed using the  $D'$  parameter also indicated linkage of these two alleles. However, the  $r^2$  value has more desirable properties.<sup>60,61</sup> A  $r^2$  value of 1 has a stricter interpretation compared to  $D'$ , as  $r^2$  can only equal 1 when the marker loci have identical allele frequencies and every occurrence of an allele at each of the markers perfectly predicts the allele at the other locus.  $D'$  can equal 1 even when the allele frequencies vary widely.<sup>52</sup> Reanalysis of our data using the more stringent  $r^2$  value failed to show linkage between any of the alleles under study.

Our main finding was that the SR-BI rs5888 polymorphism is associated with severity of liver fibrosis in chronic HCV infection. Since advanced liver fibrosis is a key factor for hepatocellular carcinoma and liver disease outcome,<sup>62</sup> this finding is likely clinically relevant. We chose the Ishak score since Ishak fibrosis stage has been shown to predict clinical outcomes, need for liver transplantation, and liver-related death in patients with chronic hepatitis C.<sup>63</sup> Even in a background of confounding factors, such as age, alcohol consumption and duration of infection,<sup>1,2,4,5</sup> the rs5888 SNP allelic variant was associated with increased fibrosis. The effect of the rs5888 polymorphism was more pronounced when the data obtained for males was analyzed separately, but analysis of the female data set was not feasible due to low numbers. Analysis of a larger cohort of male and female HCV infected individuals would be necessary to elucidate the role of gender in defining the effects of this polymorphism in disease progression. Similarly, haplotype analyses using COCAPHASE for case-control data<sup>50</sup> showed that inheritance of a haplotype consisting of rs61932577 allele C and rs5888 allele T was significantly associated with increased risk of severe liver disease development.


To determine whether the rs5888 could be exerting an effect at the transcriptional level, SR-BI mRNA was quantified in liver biopsy samples of chronically infected HCV subjects. The rs5888 minor variant homozygous genotype was associated with reduced SR-BI mRNA levels. One previous study of the impact of SR-BI variation on HCV<sup>42</sup> reported reduced liver-associated SR-BI protein-expression for the rs5888 TT genotype (together with an intronic variant, rs3782287, which we did not include in our study). Whilst these differences did not reach statistical significance, their sample size was relatively small ( $n = 40$ ). The same paper highlighted that both the rs5888 CT heterozygote and minor homozygous TT genotypes were associated with significantly increased HCV viral load compared with the CC genotype. Unfortunately, viral load data for our historic cohort was not available, so we were unable to assess this relationship.

The association between rs5888 minor variant with decreased SR-BI mRNA transcripts and increased liver disease severity was even more pronounced when considered in the context of the rs5888/rs61932577 haplotype. The study by Westhaus and colleagues<sup>42</sup> also reported a similar increased association between reduced viral load when assessing rs5888 in the context of a haplotype containing a different intron 5 variant rs3782287. (This variant alone was significantly associated with increased viral load, and a trend toward lower liver SR-BI protein expression.) Our combined data are comparable, and both show strong associations of SR-BI variants and haplotypes with key outcomes of HCV infection as well as SR-BI expression.

The mechanism by which reduced SR-BI mRNA and protein expression might lead to increased HCV viral load and disease severity is unclear. Downregulation of SR-BI expression on Huh-7 cells by RNA silencing reduces HCVpp infectivity,<sup>19</sup> therefore in vitro evidence would suggest decreased mRNA levels should result in lower HCV infectivity. However, such an effect would be less marked for HCV variants that are less dependent on SR-BI for direct cell-to-cell spread.<sup>64</sup> In addition, SR-BI mRNA levels have been found to correlate with HCV RNA levels in the liver.<sup>65</sup> However, viraemia can be present even in the absence of liver damage<sup>66</sup>; therefore the correlation of SR-BI mRNA levels with HCV RNA levels may not necessarily be important in the severity of liver disease. Reduced SR-BI transcription is associated with an increase in levels of HDL<sup>22,29,67</sup> and the HDL has recently been shown to significantly enhance HCVpp entry through interaction with SR-BI.<sup>68</sup> Therefore, decreased SR-BI transcription may lead to increased HDL levels, ultimately resulting in increased HCV infectivity. In this scenario, levels of receptor expression are not as important as the availability of HDL to potentiate the uptake of viral particles by hepatocytes.

An alternative hypothesis is that levels of plasma lipoproteins per se are more important in determining the severity of liver disease than the direct interaction of the virus with SR-BI. Patients with chronic liver diseases often have impaired lipid metabolism.<sup>69</sup> Cicognani et al.<sup>69</sup> showed that patients with cirrhosis had reduced LDL, HDL and total cholesterol serum levels compared to subjects with coronary artery disease and normal controls. In these patients a decrease in LDL was related to increasing liver disease severity. The rs5888 T variant of SR-BI has also been associated with reduced LDL-C levels in large populations of healthy subjects.<sup>31,32</sup> The rs5888 polymorphism may therefore be a marker for altered levels of serum lipoproteins, augmenting an already existing consequence of severe liver disease, thereby resulting in more severe liver disease. Analysis of serum lipoprotein levels in the cohort studied here, together with cohorts of individuals with non-HCV liver disease would be necessary to determine whether this polymorphism may be exerting an effect via alterations in lipoprotein levels. Furthermore, analysis of SR-BI protein levels, relative to mRNA levels will be necessary to determine whether the rs5888 SNP variant influences receptor expression at the cell surface.

The allele frequency for the rs5888 minor variant in our HCV cohort was 46%. Our cohort consists of predominantly white British people, and this MAF is similar to that reported in Phase 3 of the



**FIGURE 2** Frequency of rs5888 alleles in different populations. Data was obtained from the latest Ensembl release of SR-B1 variant data derived from Phase 3 of the 1000 Genomes project.<sup>46</sup> The relative frequency of each allele is represented in pie charts for individuals living in each geographical location.

1000 Genomes Project for British people living in Scotland and England (47%), and identical to that reported in European data set.<sup>46</sup> However, it is important that additional studies investigating the potential association between this, and other SR-B1 variants are carried out in additional populations with different ethnic mixes; especially considering the major differences in MAF observed in the 1000 Genome Project data set (Figure 2).

In conclusion the rs5888 polymorphism variant is associated with an increased risk of severe liver disease in chronic HCV infection. Chronically infected HCV patients who are carriers of the rs5888 polymorphism were found to have reduced SR-B1 mRNA levels in liver biopsies. The mechanism whereby reduced SR-B1 mRNA and protein expression in the liver might lead to increased viral load and severity of liver disease will need to be elucidated. An assessment of any association with rs5888 (and its haplotypes) in other chronic liver diseases will provide at least some insight into the relative contribution of viral factors or lipid metabolism in disease pathogenesis. The data presented here support the involvement of SR-B1 in HCV infection and show that SR-B1 polymorphisms are important genetic predictors of liver disease progression in HCV infection.

#### AUTHOR CONTRIBUTIONS

*Conceptualization:* Jonathan K. Ball. *Data curation:* Victoria L. Arandhara. *Formal Analysis:* Victoria L. Arandhara, Sally Chappell, Kevin Morgan, and Jonathan K. Ball. *Funding acquisition:* Jonathan K. Ball, William L. Irving, and Thomas F. Baumert. *Investigation:* Victoria L. Arandhara, Sally Chappell, and C. Patrick McClure. *Methodology:* Victoria L. Arandhara, C. Patrick McClure, and Alexander W. Tarr. *Resources:* William L. Irving. *Supervision:* Jonathan K. Ball and C.

Patrick McClure. *Writing – original draft:* Victoria L. Arandhara and Jonathan K. Ball. *Writing – review and editing:* All authors.

#### ACKNOWLEDGMENTS

This work was co-funded by the NIHR Nottingham Biomedical Research Center, through contracts from the European Union (QLK2-CT-2001-01120, MRTN-CT-2006-035599, HEALTH-F4-2012-305600) and by grants from the Special Trustees for Nottingham University Hospital (STR 64/03/N), Medical Research Council (MR/S009434/1) and the University of Nottingham Research Committee. VLA was supported by a postgraduate studentship from the Medical Research Council. TFB acknowledges funding by ERC-AdG-2020-FIBCAN #101021417, IdEx Unistra (ANR-10-IDEX-0002) and by SFRI-STRAT'US project (ANR 20-SFRI-0012) under the framework of the French Investments for the Future Program. The authors would also like to acknowledge Sue Bainbridge, Prof Patrick Tighe and Dr Olivia Belbin for technical advice and Prof Carol Coupland for advice on statistical analyses.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### ORCID

Charles Patrick McClure  <http://orcid.org/0000-0002-1710-1049>

## REFERENCES

- Mohsen AH, Group THS. The epidemiology of hepatitis C in a UK health regional population of 5.12 million. *Gut*. 2001;48(5):707-713.
- Poynard T, Bedossa P, Opolon P. Natural history of liver fibrosis progression in patients with chronic hepatitis C. *Lancet*. 1997;349(9055):825-832.
- Datz C, Cramp M, Haas T, et al. The natural course of hepatitis C virus infection 18 years after an epidemic outbreak of non-A, non-B hepatitis in a plasmapheresis centre. *Gut*. 1999;44(4):563-567.
- Ryder SD, Irving WL, Jones DA, et al. Progression of hepatic fibrosis in patients with hepatitis C: a prospective repeat liver biopsy study. *Gut*. 2004;53(3):451-455.
- Wiley TE, McCarthy M, Breidi L, McCarthy M, Layden TJ. Impact of alcohol on the histological and clinical progression of hepatitis C infection. *Hepatology*. 1998;28(3):805-809.
- Battaller R. Genetic polymorphisms and the progression of liver fibrosis: a critical appraisal. *Hepatology*. 2003;37(3):493-503.
- McCaughan GW. Fibrosis progression in chronic hepatitis C virus infection. *Gut*. 2004;53(3):318-321.
- Møhlenberg M, O'Brien TR, Hartmann R. The role of IFNL4 in liver inflammation and progression of fibrosis. *Genes Immun*. 2022;23(3-4):111-117.
- O'Brien TR, Yang HI, Groover S, Jeng WJ. Genetic factors that affect spontaneous clearance of hepatitis C or B virus, response to treatment, and disease progression. *Gastroenterology*. 2019;156(2):400-417.
- Hennig BJW, Hellier S, Frodsham AJ, et al. Association of low-density lipoprotein receptor polymorphisms and outcome of hepatitis C infection. *Genes Immun*. 2002;3(6):359-367.
- Wozniak MA, Itzhaki RF, Faragher EB, James MW, Ryder SD, Irving WL. Apolipoprotein E-ε4 protects against severe liver disease caused by hepatitis C virus: apolipoprotein E-ε4 protects against severe liver disease caused by hepatitis C virus. *Hepatology*. 2002;36(2):456-463.
- Murao K, Terpstra V, Green SR, Kondratenko N, Steinberg D, Quehenberger O. Characterization of CLA-1, a human homologue of rodent scavenger receptor BI, as a receptor for high density lipoprotein and apoptotic thymocytes. *J Biol Chem*. 1997;272(28):17551-17557.
- Cao G, Garcia CK, Wyne KL, Schultz RA, Parker KL, Hobbs HH. Structure and localization of the human gene encoding SR-BI/CLA-1. *J Biol Chem*. 1997;272(52):33068-33076.
- Calvo D, Gómez-Coronado D, Lasunción MA, Vega MA. CLA-1 is an 85-kD plasma membrane glycoprotein that acts as a high-affinity receptor for both native (HDL, LDL, and VLDL) and modified (OxLDL and AcLDL) lipoproteins. *Arterioscler Thromb Vasc Biol*. 1997;17(11):2341-2349.
- Malerød L, Juvet L, Gjøen T, Berg T. The expression of scavenger receptor class B, type I (SR-BI) and caveolin-1 in parenchymal and nonparenchymal liver cells. *Cell Tissue Res*. 2002;307(2):173-180.
- Popescu CI, Dubuisson J. Role of lipid metabolism in hepatitis C virus assembly and entry. *Biol Cell*. 2012;102(1):63-74.
- Scarselli E, Ansuini H, Cerino R, et al. The human scavenger receptor class B type I is a novel candidate receptor for the hepatitis C virus. *EMBO J*. 2002;21(19):5017-5025.
- Bartosch B, Vitelli A, Granier C, et al. Cell entry of hepatitis C virus requires a set of co-receptors that include the CD81 tetraspanin and the SR-B1 scavenger receptor. *J Biol Chem*. 2003;278(43):41624-41630.
- Lavillette D, Tarr AW, Voisset C, et al. Characterization of host-range and cell entry properties of the major genotypes and subtypes of hepatitis C virus. *Hepatology*. 2005;41(2):265-274.
- Zeisel MB, Koutsoudakis G, Schnober EK, et al. Scavenger receptor class B type I is a key host factor for hepatitis C virus infection required for an entry step closely linked to CD81. *Hepatology*. 2007;46(6):1722-1731.
- Vercauteren K, Van Den Eede N, Mesalam AA, et al. Successful anti-scavenger receptor class B type I (SR-BI) monoclonal antibody therapy in humanized mice after challenge with HCV variants within viroresistance to SR-BI-targeting agents. *Hepatology*. 2014;60(5):1508-1518.
- Varban ML, Rinninger F, Wang N, et al. Targeted mutation reveals a central role for SR-BI in hepatic selective uptake of high density lipoprotein cholesterol. *Proc Natl Acad Sci U.S.A.* 1998;95(8):4619-4624.
- Acton S, Osgood D, Donoghue M, et al. Association of polymorphisms at the SR-BI gene locus with plasma lipid levels and body mass index in a white population. *Arterioscler Thromb Vasc Biol*. 1999;19(7):1734-1743.
- Acton S, Rigotti A, Landschulz KT, Xu S, Hobbs HH, Krieger M. Identification of scavenger receptor SR-BI as a high density lipoprotein receptor. *Science*. 1996;271(5248):518-520.
- Gu X, Trigatti B, Xu S, Acton S, Babitt J, Krieger M. The efficient cellular uptake of high density lipoprotein lipids via scavenger receptor class B type I requires not only receptor-mediated surface binding but also receptor-specific lipid transfer mediated by its extracellular domain. *J Biol Chem*. 1998;273(41):26338-26348.
- Wadsack C, Hirschmugl B, Hammer A, et al. Scavenger receptor class B, type I on non-malignant and malignant human epithelial cells mediates cholesterol ester-uptake from high density lipoproteins. *Int J Biochem Cell Biol*. 2003;35(4):441-454.
- Nieland TJF, Ehrlich M, Krieger M, Kirchhausen T. Endocytosis is not required for the selective lipid uptake mediated by murine SR-BI. *Biochim Biophys Acta*. 2005;1734(1):44-51.
- Hong SH, Kim YR, Yoon YM, Min WK, Chun SI, Kim JQ. Association between HaeIII polymorphism of scavenger receptor class B type I gene and plasma HDL-cholesterol concentration. *Ann Clin Biochem*. 2002;39(Pt 5):478-481.
- Hsu LA, Ko YL, Wu S, et al. Association between a novel 11-base pair deletion mutation in the promoter region of the scavenger receptor class B type I gene and plasma HDL cholesterol levels in Taiwanese Chinese. *Arterioscler Thromb Vasc Biol*. 2003;23(10):1869-1874.
- McCarthy JJ, et al. Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease. *J Med Genet*. 2003;40(6):453-458.
- Morabia A, Ross BM, Costanza MC, et al. Population-based study of SR-BI genetic variation and lipid profile. *Atherosclerosis*. 2004;175(1):159-168.
- Osgood D, Corella D, Demissie S, et al. Genetic variation at the scavenger receptor class B type I gene locus determines plasma lipoprotein concentrations and particle size and interacts with type 2 diabetes: the framingham study. *J Clin Endocrinol Metab*. 2003;88(6):2869-2879.
- Richard E, von Muhlen D, Barrett-Connor E, Alcaraz J, Davis R, McCarthy JJ. Modification of the effects of estrogen therapy on HDL cholesterol levels by polymorphisms of the HDL-C receptor, SR-BI: the rancho bernardo study. *Atherosclerosis*. 2005;180(2):255-262.
- Rodríguez-Esparragón F, Rodríguez-Pérez C, Hernández-Trujillo Y, et al. Allelic variants of the human scavenger receptor class B type 1 and paraoxonase 1 on coronary heart disease: genotype-phenotype correlations. *Arterioscler Thromb Vasc Biol*. 2005;25(4):854-860.
- Tai E, Adiconis X, Ordovas J, et al. Polymorphisms at the SRBI locus are associated with lipoprotein levels in subjects with heterozygous familial hypercholesterolemia. *Clin Genet*. 2003;63(1):53-58.
- Çaykara B, Tokat B, Coşkunpınar E, et al. Investigation of scavenger receptor class B type I gene variants in patients with coronary heart disease with a history of early myocardial infarction. *Turk Kardiyol Dern Ars*. 2021;49(8):641-653.



37. Goodarzynejad H, Boroumand M, Behmanesh M, Ziaee S, Jalali A. The rs5888 single nucleotide polymorphism in scavenger receptor class B type 1 (SCARB1) gene and the risk of premature coronary artery disease: a case-control study. *Lipids Health Dis.* 2016;15:7. doi:10.1186/s12944-016-0176-9
38. Pérez-Martínez P, Ordovás JM, López-Miranda J, et al. Polymorphism exon 1 variant at the locus of the scavenger receptor class B type I gene: influence on plasma LDL cholesterol in healthy subjects during the consumption of diets with different fat contents. *Am J Clin Nutr.* 2003;77(4):809-813.
39. West M, Greason E, Kolmakova A, et al. Scavenger receptor class B type I protein as an independent predictor of high-density lipoprotein cholesterol levels in subjects with hyperalphalipoproteinemia. *J Clin Endocrinol Metab.* 2009;94(4):1451-1457.
40. Wu D-F, Yin RX, Cao XL, et al. Scavenger receptor class B type 1 gene rs5888 single nucleotide polymorphism and the risk of coronary artery disease and ischemic stroke: a case-control study. *Int J Med Sci.* 2013;10:1771-1777. doi:10.7150/ijms.7044
41. Hsu CS, Hsu SJ, Liu WL, Chen DS, Kao JH. Association of SCARB1 gene polymorphisms with virological response in chronic hepatitis C patients receiving pegylated interferon plus ribavirin therapy. *Sci Rep.* 2016;6:32303.
42. Westhaus S, Deest M, Nguyen ATX, et al. Scavenger receptor class B member 1 (SCARB1) variants modulate hepatitis C virus replication cycle and viral load. *J Hepatol.* 2017;67(2):237-245.
43. Yamamoto S, Fukuhara T, Ono C, et al. Lipoprotein receptors redundantly participate in entry of hepatitis C virus. *PLoS Pathog.* 2016;12(5):e1005610.
44. Gao R, Gao W, Xu G, Xu J, Ren H. Single amino acid mutation of SR-BI decreases infectivity of hepatitis C virus derived from cell culture in a cell culture model. *World J Gastroenterol.* 2017;23(28):5158-5166.
45. Irving WL, Neal KR, Underwood JCE, Simmonds PN, James V. Chronic hepatitis in United Kingdom blood donors infected with hepatitis C virus. *BMJ.* 1994;308(6930):695-696.
46. Cunningham F, Allen JE, Allen J, et al. Ensembl 2022. *Nucleic Acids Res.* 2021;50(D1):D988-D995.
47. Ishak K, Baptista A, Bianchi L, et al. Histological grading and staging of chronic hepatitis. *J Hepatol.* 1995;22(6):696-699.
48. Neal KR, et al. Excess mortality rates in a cohort of patients infected with the hepatitis C virus: a prospective study. *Gut.* 2007;56(8):1098-1104.
49. Constantineau J, Greason E, West M, et al. A synonymous variant in scavenger receptor, class B, type I gene is associated with lower SR-BI protein expression and function. *Atherosclerosis.* 2010;210(1):177-182.
50. Dudbridge F. Pedigree disequilibrium tests for multilocus haplotypes. *Genet Epidemiol.* 2003;25(2):115-121.
51. Bustin S. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol.* 2000;25(2):169-193.
52. Zondervan KT, Cardon LR. The complex interplay among factors that influence allelic association. *Nat Rev Genet.* 2004;5(2):89-100.
53. Goyal A, Kazim SN, Sakhuja P, Malhotra V, Arora N, Sarin SK. Association of TNF- $\alpha$  polymorphism with disease severity among patients infected with hepatitis C virus. *J Med Virol.* 2004;72(1):60-65.
54. Mangia A, Santoro R, Piattelli M, et al. IL-10 haplotypes as possible predictors of spontaneous clearance of HCV infection. *Cytokine.* 2004;25(3):103-109.
55. Tokushige K, Tsuchiya N, Hasegawa K, et al. Influence of TNF gene polymorphism and HLA-DRB1 haplotype in Japanese patients with chronic liver disease caused by HCV. *Am J Gastroenterol.* 2003;98(1):160-166.
56. Di Lello FA, Caruz A, Rallon NI, et al. Effects of the genetic pattern defined by low-density lipoprotein receptor and IL28B genotypes on the outcome of hepatitis C virus infection. *Eur J Clin Microbiol Infect Dis.* 2013;32(11):1427-1435.
57. Caruz A, Neukam K, Rivero-Juárez A, et al. Association of low-density lipoprotein receptor genotypes with hepatitis C viral load. *Genes Immun.* 2014;15(1):16-24.
58. Turner D, Choudhury F, Reynard M, Railton D, Navarrete C. Typing of multiple single nucleotide polymorphisms in cytokine and receptor genes using SNaPshot. *Hum Immunol.* 2002;63(6):508-513.
59. Yamaguchi N, Kobayashi K, Yasuda T, et al. Screening of SLC25A13 mutations in early and late onset patients with citrin deficiency and in the Japanese population: identification of two novel mutations and establishment of multiple DNA diagnosis methods for nine mutations. *Hum Mutat.* 2002;19(2):122-130.
60. Carlson CS, Eberle MA, Kruglyak L, Nickerson DA. Mapping complex disease loci in whole-genome association studies. *Nature.* 2004;429(6990):446-452.
61. Weiss KM, Clark AG. Linkage disequilibrium and the mapping of complex human traits. *TIG.* 2002;18(1):19-24.
62. Roehlen N, Crouchet E, Baumert TF. Liver fibrosis: mechanistic concepts and therapeutic perspectives. *Cells.* 2020;9(4):875.
63. Everhart JE, Wright EC, Goodman ZD, et al. Prognostic value of Ishak fibrosis stage: findings from the hepatitis C antiviral long-term treatment against cirrhosis trial. *Hepatology.* 2010;51(2):585-594.
64. Catanese MT, Loureiro J, Jones CT, Dorner M, von Hahn T, Rice CM. Different requirements for scavenger receptor class B type I in hepatitis C virus Cell-Free versus Cell-to-Cell transmission. *J Virol.* 2013;87(15):8282-8293.
65. Ikura Y, Ohsawa M, Ueda M. The amount of hepatitis C viral RNA is correlated to the expression level of scavenger receptor class B type 1 in patients' livers. *J Clin Virol.* 2005;33(4):345-346.
66. Alberti A, et al. Prevalence of liver disease in a population of asymptomatic persons with hepatitis C virus infection. *Ann Intern Med.* 2002;137(12):961-964.
67. Rigotti A, Trigatti BL, Penman M, Rayburn H, Herz J, Krieger M. A targeted mutation in the murine gene encoding the high density lipoprotein (HDL) receptor scavenger receptor class B type I reveals its key role in HDL metabolism. *Proc Natl Acad Sci U.S.A.* 1997;94(23):12610-12615.
68. Voisset C, et al. High density lipoproteins facilitate hepatitis C virus entry through the scavenger receptor class B type I. *J Biol Chem.* 2005;280(9):7793-7809.
69. Cicognani C, et al. Serum lipid and lipoprotein patterns in patients with liver cirrhosis and chronic active hepatitis. *Arch Intern Med.* 1997;157(7):792-796.
70. Carlquist JF, McKinney JT, Horne BD, et al. Common variants in 6 lipid-related genes discovered by high-resolution DNA melting analysis and their association with plasma lipids. *J Clin Exp Cardiol.* 2011;2(138). doi:10.4172/2155-9880.1000138
71. Niemsiri V, Wang X, Pirim D, et al. Genetic contribution of SCARB1 variants to lipid traits in African Blacks: a candidate gene association study. *BMC medical genetics.* 2015;16:106. doi:10.1186/s12881-015-0250-6

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Arandhara VL, McClure CP, Tarr AW, et al. Scavenger receptor class B type I genetic variants associated with disease severity in chronic hepatitis C virus infection. *J Med Virol.* 2022;e28331. doi:10.1002/jmv.28331