



HAL
open science

Mercury dynamic in a mountainous catchment polluted by chlor-alkali activity (Alsace, France)

Christophe Hissler, Jean-Luc Probst

► **To cite this version:**

Christophe Hissler, Jean-Luc Probst. Mercury dynamic in a mountainous catchment polluted by chlor-alkali activity (Alsace, France). The 15th Annual Goldschmidt Conference: A voyage of discovery, May 2005, Moscow, United States. pp.A707-A707. hal-04037423

HAL Id: hal-04037423

<https://hal.science/hal-04037423>

Submitted on 20 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mercury dynamic in a mountainous catchment polluted by chlor-alkali activity (Alsace, France)

C. HISSLER ^{1,2} AND J.L. PROBST ^{1,2}

¹Institut National Polytechnique, ENSAT, AEE Laboratory, Castanet Tolosan, France (jean-luc.probst@ensat.fr)

²CNRS, Lab. des Mécanismes de Transfert en Géologie, Toulouse, France (christophe-hissler@hotmail.fr)

Total atmospheric mercury contamination was measured using CV-AFS in soils, river bottom sediments, river waters and bryophytes (*Hypnum cupressiform*) samples collected in a French mountainous catchment upstream from a chlor-alkali industrial activity. Natural and atmospheric/anthropogenic contributions to total Hg contents in soils and stream sediments were assessed using Hg concentrations in the deepest soil mineral horizon and in the stream bottom sediments. Soil Hg stock and mean annual particulate and dissolved Hg river fluxes coming from the upper part of the basin are estimated for the first time in this catchment.

The natural background level of Hg content deriving from rock weathering was estimated to 32 ± 9 ng g⁻¹ in the deepest soil layers. The soils appear to be Hg contaminated in two ways: atmospheric deposition and Hg-organic matter complexes leaching through the soil profiles. Hg concentrations in upper stream sediments varied between 108 and 639 ng g⁻¹. The Hg enrichment factor (EF_{Hg}) which could be calculated by normalization to Sc, allow to estimate the major contribution (63 to 95%) of the atmospheric inputs, even in the upper part of the basin. This contribution may be attributed to diffuse regional atmospheric deposition of Hg and is mainly due to the geographic location of this chemical plant. The suspended matter was the main phase of Hg fluvial transportation in the different tributaries, with concentrations close to those of river bottom sediments. The Hg flux due to the weathering of the upper basin ($0,58 \cdot 10^{-3}$ mg m⁻² y⁻¹) was negligible in comparison with the yearly Hg discharges in the Thur River due to the anthropogenic activities. The calculation of the Hg residence time indicates that Hg is trapped in the soils of the upper catchment for a long time in such a polluted hydrosystem.