

Seven-year monitoring of mercury in wet precipitation and atmosphere at the Amsterdam Island GMOS station

Antonella Tassone, Olivier Magand, Attilio Naccarato, Maria Martino, Domenico Amico, Francesca Sprovieri, Hippolyte Leuridan, Yann Bertrand, Michel Ramonet, Nicola Pirrone, et al.

► To cite this version:

Antonella Tassone, Olivier Magand, Attilio Naccarato, Maria Martino, Domenico Amico, et al.. Sevenyear monitoring of mercury in wet precipitation and atmosphere at the Amsterdam Island GMOS station. Heliyon, 2023, 9 (3), pp.e14608. 10.1016/j.heliyon.2023.e14608 hal-04037485v1

HAL Id: hal-04037485 https://hal.science/hal-04037485v1

Submitted on 20 Mar 2023 (v1), last revised 24 Mar 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Seven-year monitoring of mercury in wet precipitation and atmosphere at the Amsterdam Island GMOS station

Antonella Tassone, Olivier Magand, Attilio Naccarato, Maria Martino, Domenico Amico, Francesca Sprovieri, Hippolyte Leuridan, Yann Bertrand, Michel Ramonet, Nicola Pirrone, Aurelien Dommergue

PII: S2405-8440(23)01815-7

DOI: https://doi.org/10.1016/j.heliyon.2023.e14608

Reference: HLY 14608

To appear in: HELIYON

Received Date: 20 December 2022

Revised Date: 5 March 2023

Accepted Date: 13 March 2023

Please cite this article as: , Seven-year monitoring of mercury in wet precipitation and atmosphere at the Amsterdam Island GMOS station, *HELIYON* (2023), doi: https://doi.org/10.1016/j.heliyon.2023.e14608.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.



CRedit Authorship contribution statement

Antonella Tassone: Conceptualization of the study, preparation of material for sampling, laboratory Hg analysis, data analysis, original manuscript writing; Olivier Magand: Implementation of Hg measurements at TAAF-IPEV Amsterdam Island district, observations monitoring management, field campaign, data treatment qualification and validation, IPEV GMOStral-1028 and Hg dataset AERIS (GMOS-FR national database https://gmos.aeris-data.fr/) program managing, funding acquisition, original manuscript writing; Attilio Naccarato: Conceptualization of the study, supervision and coordination of the work, original manuscript writing, funding acquisition and project managing; Maria Martino: Preparation of material for sampling, laboratory Hg analysis, data analysis; Domenico Amico: Preparation of material for sampling, laboratory Hg analysis, data analysis; Francesca Sprovieri: Manuscript review and editing, funding acquisition and project managing, GMOS network coordination; Hippolyte Leuridan: ICOS station management at Amsterdam Island; Yann Bertrand: IPEV GMOStral-1028 field campaign on THg in precipitation; THg collector system maintenance; Michel Ramonet: ICOS station management at Amsterdam Island; Nicola Pirrone: Funding acquisition and project managing, GMOS network coordination; Aurelien Dommergue: Implementation of Hg measurements at TAAF-IPEV Amsterdam Island district, funding acquisition. All co-authors contributed to data interpretation, reviewing, and editing the manuscript.

JunalPre



And a series during dur



2013 – 2019 monitoring of GEM and total Hg in wet deposition

Journal Prosproof

1	Seven-year monitoring of mercury in wet precipitation and atmosphere at the Amsterdam
2	Island GMOS station
3	
4	Antonella Tassone ^{1*} , Olivier Magand ² , Attilio Naccarato ^{1,3} , Maria Martino ¹ , Domenico Amico ¹ ,
5	Francesca Sprovieri ¹ , Hippolyte Leuridan ⁴ , Yann Bertrand ² , Michel Ramonet ⁴ , Nicola Pirrone ¹
6	and Aurelien Dommergue ²
7	
8	
9	
10	1 CNR-Institute of Atmospheric Pollution Research (CNR-IIA), Rende, Italy
11	2 Institut des Géosciences de l'Environnement, Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP,
12	38400 Grenoble, France
13	3 Department of Chemistry and Chemical Technologies, University of Calabria, Via P. Bucci,
14	Rende, Italy
15	4 Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL (CEA-CNRS-UVSQ),
16	Université Paris-Saclay, 91191 Gif-sur-Yvette, France
17	
18	
19	
20	*Corresponding author:
21	Antonella Tassone
22	CNR-Institute of Atmospheric Pollution Research, Rende, Italy
23	e-mail: antonella.tassone@iia.cnr.it
24	

26 Abstract

Mercury (Hg) fate and transport research requires more effort to obtain a deep knowledge of its
biogeochemical cycle, particularly in the Southern Hemisphere and Tropics that are still missing
of distributed monitoring sites.

Continuous monitoring of atmospheric Hg concentrations and trend worldwide is relevant for the 30 effectiveness evaluation of the Minamata Convention on Mercury (MCM) actions. In this context, 31 Gaseous Elemental Mercury (GEM) and total mercury (THg) in precipitations were monitored 32 from 2013 to 2019 at the Amsterdam Island Observatory (AMS - 37°48'S, 77°34'E) to provide 33 insights into the Hg pathway in the remote southern Indian Ocean, also considering ancillary 34 dataset of Rn-222, CO₂, CO₂, and CH₄. GEM average concentration was 1.06 ± 0.07 ng m⁻³, with 35 a slight increase during the austral winter due to both higher wind speed over the surface ocean 36 and contributions from southern Africa. In wet depositions, THg average concentration was 2.39 37 \pm 1.17 ng L⁻¹, whereas the annual flux averaged 2.04 \pm 0.80 µg m⁻² yr⁻¹. In general, both GEM and 38 Volume-Weighted Mean Concentration (VWMC) of THg did not show an increasing/decreasing 39 trend over the seven-year period, suggesting a substantial lack of evolution about emission of Hg 40 reaching AMS. 41

Air masses Cluster Analysis and Potential Source Contribution Function showed that oceanic
 evasion was the main Hg contributor at AMS, while further contributions were attributable to long range transport events from southern Africa, particularly when the occurrence of El Niño increased
 the frequency of wildfires.

- 46
- 47

48 Keywords: Wet deposition flux; Southern Hemisphere; Atmospheric pollution; Source
49 identification; Atmospheric transport; Mercury measurements

51 **1. Introduction**

Due to its toxicity and great impact on human health and the environment, mercury (Hg) has 52 increasingly been investigated by the scientific community [1,2] with the overall aim to reduce 53 its emissions in each environmental compartment [3]. The Minamata Convention on Mercury 54 (MCM) is perhaps the most important international regulatory framework aimed at protecting 55 human health and the environment from anthropogenic emissions of Hg and its compounds [4] 56 and the effectiveness evaluation of its actions has highlighted the need to increase the number of 57 ground-based monitoring sites worldwide, particularly in areas of scarce data coverage and to 58 provide comparable Hg data both in the background and in impacted areas [5]. This complex 59 framework draws attention to the need for multiple analytical methods and the exploitation of 60 new analytical strategies for comprehensive investigations, which should also include remote 61 areas [6–9]. 62

Although several Hg monitoring programs and networks have been established over the years,
[10–13] the monitoring stations that provide Hg data both in air and wet depositions in the open
ocean and remote areas are still limited in number [8,10,14–24] particularly in the Southern
Hemisphere.

In this context, the station on Amsterdam Island (AMS) has been set up in the framework of the Global Mercury Observation System (GMOS) Hg global network (<u>http://www.gmos.eu/</u>), with the support of the French Polar Institute (IPEV). The unique geographic location of this site has enabled the study of background data on major air pollutants in the Southern Hemisphere due to the absence of local anthropogenic sources and its distance from maritime traffic routes [15,17,25–29].

The present manuscript is an updated study on the temporal variability of Gaseous Elemental Mercury (GEM) and total Hg (THg) in precipitations from 2013 to 2019 at AMS providing new insights into the Hg pathway in the Southern Hemisphere. The investigation of seasonal and interannual variability, combined with the analysis of CO₂, CO, CH₄, and radon-222 (Rn-222) data,

\sim	111			5	rn	\sim	
U	uu				ιU	U.	

- with the support of Potential Source Contribution Function (PSCF) and Cluster Analysis (CA),
 permitted to assess the evolutionary tendencies of Hg at the Southern Hemisphere midlatitudes
 and pinpoint the potential sources associated with long-range transport and El Niño Southern
 Oscillation (ENSO).
- 81 **2.** Materials and methods
- 82 **2.1 Sampling site**

Amsterdam Island (37.7983° S, 77.5378° E) is a small remote volcanic island of 55 km² in the southern Indian Ocean, 3200 km from Australia, 3400 km from Madagascar, 4200 km from South Africa, and 3300 km from Antarctica. The atmospheric observatory of Pointe Benedicte is northwest to the island at an altitude of 55 m above sea level, and due to its strategic position that ensures minimal local interference, the station is of particular interest to study Hg dynamics and pathway under unique background conditions.

Within the GMOS global network, it has been classified as a master site [11] and it is also part of
other atmospheric monitoring programs such as the Global Atmosphere Watch World
Meteorological Organization (GAW WMO), the French national monitoring system Integrated
Carbon Observation System ICOS-France-Atmosphere for the long-term observation of
greenhouse gases (GHGs), and the Global Observation System for Mercury (GOS⁴M) GEO
Flagship [10,16,17]. A more detailed description of the site is given in the Supporting Information
(Text S1).

96

2.2 Meteorological variables and CO₂, CO, CH₄, and Rn-222 measurements

97 Meteorological data – air temperature, rainfall amount, wind speed and direction – were provided 98 by the local meteorological station. The dataset of GHGs included CO₂, CO, and CH₄ from March 99 23, 2013, to December 22, 2019. A temporal gap exists in the CO measurements, which started 100 from December 2, 2015. CO₂, CO, and CH₄ were monitored using Picarro gas analyzers, which 101 used cavity ring-down spectroscopy to measure the gas concentration every 30 seconds with a 102 precision (1 sigma confidence) for hourly measurements of 10 ppb, 1 ppb, and 0.3 ppb,

respectively, in compliance with the specifications of the manufacturer. Calibration was performed
 monthly. The raw data were manually checked and validated by ICOS staff [30], before being
 hourly averaged.

106 Radon-222 measurements, covering the same period considered for GHGs, were performed using 107 an alpha spectrometer which required monthly calibration and ensured a minimum detected 108 activity of about 10 mBq m⁻³ with an uncertainty of about 40%.

This set of ancillary data proved useful to track continental air over the ocean [31]. Indeed, Rn-109 222 concentration at AMS exceeding the threshold of 80 mBq m⁻³ (i.e., radonic storms), indicated 110 that the air mass over the island recently passed over the mainland, usually southern Africa [17,32]. 111 The time series of all gases were analyzed on an hourly, seasonal, and annual basis, with seasonal 112 boundaries defined as follows: austral winter from July to September, austral summer from 113 December to February, intermediate season 1 (IS1) from March to June, and intermediate season 114 2 (IS2) from October to November. This classification revealed useful since the meteorological 115 conditions in June are more similar to that in the preceding months than in July-August, as also 116 the case of other sites in the Southern Hemisphere, and more strictly in Africa [33]. 117

118 119

2.3 Mercury determination in the atmosphere and wet deposition

2.3.1 **GEM** measurements

GEM was continuously measured from March 23, 2013, to December 22, 2019, using Tekran analyzer 2537A/B models (Tekran Inc., Toronto, Canada) based on amalgamation onto dual gold cartridge design, thermal desorption, and quantification by cold vapor atomic fluorescence spectroscopy (CV-AFS). Calibration was performed using both the internal permeation sources (every 69 h) and manual injections of saturated Hg vapor (every year). To ensure comparability of the Hg data, GEM measurements were carried out according to GMOS Standard Operating Procedures (SOPs) [34].

127

2.3.2 Rainwater sampling and total mercury quantification

Wet deposition samples were continuously collected from March 24, 2013, using an automatic 128 wet-only NSA-171 Eigenbrodt collector which comprises a glass funnel connected to a 1-L 129 fluorinated high-density polyethylene (FLPE) bottle. The entire sampling apparatus was 130 thoroughly acid-cleaned, according to the procedures reported by Tassone et al. [35], based on the 131 SOPs developed within the GMOS network [36]. Each deployment lasted 6-45 days, depending 132 on the season (wet or dry) and according to exceptional periods of intense rainfall events. Analyses 133 of wet deposition samples were performed at the CNR-IIA laboratories following the US-EPA 134 method 1631E guidance reported in the literature [35]. Quality assurance/quality control (QA/QC) 135 procedures were routinely implemented, including the analysis of duplicate samples, accuracy test 136 using certified reference material (NIST1641), matrix spikes, and assessment of Hg contamination 137 through the analysis of the system, transport, reagent and field blanks (more details in Text S2). 138

139

2.4 Data treatment

Raw GEM time series were quality-controlled according to GMOS SOPs, using a dedicated software developed at the Institute of Environmental Geosciences (<u>https://doi.org/10.25326/168</u> available at https://gmos.aeris-data.fr, last access: 10 February 2023) [37]. Specifically, the QA/QC procedure involved comparing the raw data with fixed criteria, concerning both the calibration and the readout steps of airborne measurements [34].

THg values were derived according to UNI 15853:2010 and converted into volume-weighted mean concentration (VWMC) values, while seasonal and annual Hg wet deposition fluxes (F_w) were calculated as reported in Sprovieri at al. [10] by using the seasonal/yearly VWMCs and cumulative rain depths (P^i , mm), according to Eq. (1):

149

$$F_{w} = VWMC_{Hg} \sum_{i=1}^{n} P^{i} / 1000$$
(1)

The statistical significance of temporal variability of each variable was determined using linear regression with the method of Pearson's correlation coefficient and the differences in annual and seasonal concentration values were investigated using Mann-Whitney and ANOVA tests.

The seasonal Mann-Kendall trend test was applied to monthly mean data of each variable, after verification of their significant seasonality using the Kruskal-Wallis test, to detect trends while Sen's slope estimator was applied to assess their magnitude, using the Theil-Sen method [38].

The investigation of the impact of El Niño on Amsterdam Island was conducted using crosscorrelation to seek lag times between the Southern Oscillation Index (SOI), as a tracer of El Niño occurrence, and GEM and CO concentrations. All data analyses were performed using the R software.

160

2.5 Backward trajectories and source identification techniques

The interpretation of the surveyed time series was carried out using the U.S. NOAA's HYbrid 161 Single-Particle Lagrangian Trajectory 162 Integrated (HYSPLIT) model 163 (http://ready.arl.noaa.gov/HYSPLIT.php, last access: May 3, 2022), which permitted to obtain information on the path of air masses to the receptor site. For this purpose, the READY website 164 was used to compute 5-day back-trajectories, at 3-hour intervals, at 50 m above ground level in 165 the domain 70°-0° S and 0°-120° E [33,39,40]. Considering the remoteness of the island, 5-day 166 167 trajectories may be deemed short for tracing possible Hg sources reaching AMS. However, longer trajectories calculated for the entire 7-year monitoring period would have entailed a higher 168 computational cost. We therefore calculated 10-day trajectories only for few cases, in order to 169 support our findings overcoming computational limitations. More specifically, 10-day trajectories 170 were used to facilitate the interpretation of particularly high or low values of GEM and Rn as well 171 172 as to run the CA. The GDAS 1°x1° from the Global Data Assimilation System was selected as the meteorological dataset [41]. 173

In addition, CA was applied to group air masses according to their origin through the computation of the proportion of time that a particular back-trajectory was in a specific wind sector (e.g., S, SW, W). CA was performed using Ward's Minimum-Variance method which, starting with groups containing single trajectories, iteratively merges pairs of them so as to obtain the minimum increase in the sum of squared distances between the members and the centroids of the

corresponding groups, summed over the resulting groups [42-44]. The centroid of the newly-179 formed group, i.e., our mean trajectory, is recomputed each time using the data from the individual 180 groups previously separated, before calculating the squared distances [45]. In our study, an angle 181 distance matrix was used to assess the similarity between the trajectories to be merged. The optimal 182 number of clusters was chosen following the total spatial variance (TSV) metric in the Elbow 183 method [45,46], selecting a variance criterion of 20%. This approach relies on calculating the sum 184 of the spatial variance of all the clusters, until TSV increases significantly, thus indicating that the 185 merged clusters are not very similar. This increase in TSV denotes the completion of the clustering 186 process [47], and in our case, it resulted in four clusters as the optimal number. Furthermore, a 187 hybrid receptor model, namely the Potential Source Contribution Function, was used to locate 188 potential emission sources around the receptor site [48,49]. This technique assumes that air masses 189 that intercept emission sources in the grid cell (i, j) can get enriched in the pollutants emitted by 190 that source and transport them to the receptor site. The PSCF is defined by Eq. (2): 191

$$PSCF = \frac{m_{ij}}{n_{ij}}$$
(2)

where n_{ij} is the number of times that the trajectories passed through the cell and m_{ij} is the number of times that a source concentration was high (>90th percentile) when the trajectories passed through the cell (i, j). The cell with the higher PSCF value thereby characterizes the most likely source location.

197 **3. Results and discussion**

192

198

3.1 Variability of CO₂, CO, CH₄, Rn, and meteorological conditions

The summary of the annual meteorological variables and atmospheric gases is listed in Table 1. The monthly median air temperature ranged from 11 °C in winter to 19 °C in summer (Fig. S1) while winds mainly blew from NNW to WSW, without significant differences in direction among the seasons (Fig. S1), except for a higher frequency of N-NW air masses during winter. The most abundant precipitations were measured during IS1 except for 2013, when the maximum was

The measured CO₂ (n=56976, 89.5% of available hourly data) showed a statistically significant increase over the investigated period, from 390.5 ppm in 2013 to 412.8 ppm in 2019 (Fig. S2), with an average rate of 2.46 ppm year⁻¹ (p<0.05), thus confirming the growing trend already found in previous studies [26].. Daily CO₂ concentration showed little variability among the seasons, with a slight decrease between 9 a.m. and 6 p.m., suggesting a weak influence from the photosynthetic activity of local vegetation (Fig. S2).

The hourly CO values (n=34264, 88% of available data) ranged from 30 ppb to a maximum of 65 212 213 ppb (Fig. S3), with an upward significant trend (p<0.05), as confirmed by the positive Sen's slope value, which was 6.8e-05 ppb year⁻¹. However, it is worth noticing that this increasing tendency 214 was very small, thus not physically meaningful. On a seasonal basis, a clear distribution was 215 observed, with the lowest values being obtained in February and March (Fig. S3), due to removal 216 phenomena favored by the high presence of radical OH during IS1. Conversely, these removal 217 processes were weak during winter and the CO concentration increased significantly, with maxima 218 in October [50]. In addition, between June and October, this pattern was probably influenced by 219 220 the long-range transport of CO released during the biomass burning events in southern Africa [51]. 221 The hourly CH₄ values (Fig. S4) showed a statistically significant increase from 1747 ppb to 1841 ppb with an average value of 1794 ± 21 ppb (n=56964, 89.5% of available hourly data) and a 222 growth rate of 8.5 ppb year⁻¹ (p < 0.05). The lowest values were recorded in January-April (Fig. S4) 223 224 and were mainly due to the reaction with OH radicals, as was also observed for CO. Indeed, CH4 and CO showed a significant correlation (r=0.57, p<0.05) over the covered period, suggesting that 225 common mechanisms have driven their pattern on a seasonal scale [50,52,53]. 226

The 2-hour Rn-222 activity was on average 35.36 ± 32.24 mBq m⁻³ (n=26988, 92.3% of available data) with a maximum of 398.80 mBq m⁻³ (Fig. S5). The Mann-Kendall test revealed a significant decrease that was confirmed by the Sen's slope value (-4.1e-04 mBq m⁻³ year⁻¹). However, this

tendency was quite small, therefore not physically meaningful, as in the case of CO. Annual Rn 230 activities (Fig. S5) ranged between 24.93 ± 26.78 mBg m⁻³ (in 2016) and 48.10 ± 19.4 mBg m⁻³ 231 (in 2013), as typical for marine air masses which usually range between 30-50 mBq m⁻³ [27,54,55]. 232 Radonic storms from southern Africa were mainly detected during winter (Fig. S5), which is the 233 season with the greatest contribution of the long-range transport [17]. Their continental origin was 234 confirmed by the correlation between Rn and CO, which improved when considering only the 235 biomass burning season (i.e., austral winter) rather than the entire period, with a 10-fold increase 236 in the correlation coefficient, from 0.02 to 0.20 (p<0.05). Besides, by looking at the relationship 237 between Rn and wind direction (Fig. S5), it can be noticed a lower frequency of radonic storms 238 239 transported up to AMS during the second half of the investigated period. Although there was substantial uniformity in the direction of Rn origin (WNW), these radonic storms decreased their 240 frequency from W over time, and in 2019 they were recorded only from the northernmost (NW) 241 sectors. 242

243

244

3.2.1 GEM measurements

3.2 Hg in the atmosphere and wet deposition

The individual mean hourly GEM values from March 2013 to December 2019 (Fig. 2) ranged 245 from 0.70 to 1.58 ng m⁻³, resulting in a mean concentration of 1.06 ± 0.07 ng m⁻³ (n=45018, 70.7%) 246 of available data), which is comparable with the concentrations (0.9-1.0 ng m⁻³) usually found at 247 background sites of the Southern Hemisphere [11]. The observed GEM was also consistent with 248 previous measurements at AMS, which reported concentrations of 1.03 ng m⁻³ [17] and 1.05 ng 249 m⁻³ [16], as well as at the Cape Point station in southern Africa, where the annual mean GEM was 250 1.09 ng m⁻³ [16]. However, concentrations at AMS were slightly higher than in other subequatorial 251 sites, such as the EMMA station in Argentina, where the average concentration was 0.9 ng m^{-3} 252 [22], and the Cape Grim station in Tasmania, where concentrations of 0.9-1.0 ng m⁻³ were found 253 [56,57]. On a seasonal basis (Fig. 2), we recorded a slightly but significantly higher Hg 254 concentration in winter than in summer $(1.08 \pm 0.05 \text{ vs.} 1.06 \pm 0.07 \text{ ng m}^{-3}, \text{p}<0.05)$. This seasonal 255

variability may be ascribed to the impact of the Intertropical Convergence Zone (ITCZ) shift 256 during the austral winter, which caused changes in the wind speed on the sea surface [58], thus 257 favoring Hg emission. Indeed, ITCZ is a belt of air masses from both the Northern and Southern 258 hemispheres mixing together which crosses the equator from south to north around April-May 259 (IS1) and reaches its northernmost point around June-July (winter), and then moves south again 260 around August-September [58,59]. Therefore, the relocation of the ITCZ over the Indian Ocean 261 was likely a vector of air masses from the north to AMS that reasonably influenced the recorded 262 GEM concentrations. In addition, a non-negligible contribution to the observed GEM seasonal 263 variability was the long-range transport from southern Africa during the biomass burning season, 264 that was confirmed by the significant positive correlation between GEM and CO only in winter 265 (r=0.28, p<0.05), contrary to the findings regarding the correlations between GEM and CO₂ or 266 CH4, which were in all cases non-significant. 267

268

3.2.2 Total mercury in rainwater

The annual-averaged VWMC of THg in precipitation samples (n=27) was 2.39 ± 1.17 ng L⁻¹ and 269 varied in the range from 1.16 ng L⁻¹ to 4.67 ng L⁻¹, which were obtained in 2015 and 2018, 270 respectively, when total rainfall was 916 mm and 855 mm (Table 1 and Fig. 1). On both a seasonal 271 272 and annual basis, the VWMC showed no significant increase or decrease over time, suggesting 273 that no substantial changes in the concentration of atmospheric oxidized Hg species during the investigated period were observed. Since a significant source of these species at AMS is biomass 274 burning in southern Africa [16], it can be hypothesized that these emissions did not change during 275 276 the last decade.

Even focusing only on the winter seasons, when African air masses are detectable at AMS, we did
not observe a clear upward/downward tendency of VWMC, rather it was mostly fluctuating (Fig.
1). Except for 2016, the lowest values were observed during the rainy season, i.e., from May to
October, due to the movement of cold fronts from the west. Conversely, higher values were mainly
found in IS2 (Fig. 1), except for 2014 and 2017 when a peak was recorded during summer. In

particular, during IS2 2018 we observed a VWMC value of 10.93 ng L⁻¹, associated a rainfall 282 amount (199 mm) that was not notably different from the typically measured values. The analysis 283 of the wind roses combined with backward trajectories during IS2 2018 did not provide us with 284 immediate answers to this event. Within the same synoptic pattern, more intense winds from the 285 W and WNW sectors in IS2 than other seasons were recorded (Fig. S6), suggesting the potential 286 contribution of long-range transport from Africa. Therefore, we hypothesize during this season 287 some unusual emission phenomena occurred in Africa, whose oxidized Hg was then scavenged by 288 the wet deposition reaching AMS [60,61]. This hypothesis is supported by previous measurements 289 at AMS, which showed that the highest concentrations of particle-bound Hg were observed during 290 291 episodes of stronger winds from the NW sector [17].

The effect of washout of atmospheric oxidized Hg species from wet deposition was studied on a 292 seasonal basis by correlating VWMC with total rainfall (n=27, r=-0.21, p>0.05) (Fig. 3), which 293 resulted in a negative but not sufficient coefficient. The correlation was low because the occurrence 294 was observed by considering seasonal values from samples that averaged out their respective 295 contributions. Similarly, a non-significant correlation was found between GEM and wet deposition 296 volume (n=27, r=0.11, p>0.05), suggesting rainfall events did not affect GEM concentration (Fig. 297 3). As regards the VWMC and the corresponding seasonal GEM concentrations (Fig. 3), the 298 299 correlation study revealed a not significant linear relationship between the two forms (n=27, r=0.05, p>0.05), which was probably due to their different temporal variability. Indeed, GEM 300 showed a quite stationary pattern compared to the more variable VWMC throughout the covered 301 302 period.

The values of VWMC in precipitation samples and their respective precipitation volumes resulted in an average wet deposition flux of $2.04 \pm 0.80 \ \mu g \ m^{-2} \ year^{-1}$, and therefore slightly higher than the value of $1.95 \ \mu g \ m^{-2} \ year^{-1}$ reported in a previous study [10]. This finding was impacted by the higher value of Hg in wet deposition observed in 2018. In fact, during 2018 the highest wet

deposition flux was attained (3.31 μ g m⁻² year⁻¹), whereas the lowest value (1.06 μ g m⁻² year⁻¹) resulted in 2015. On a seasonal scale, the maximum flux varied year-to-year between IS1 and IS2. A positive significant correlation (n=27, r=0.48, p<0.05) was observed between the wet deposition flux and associated rainfall (Fig. 3), indicating that wet Hg deposition flux increased with increasing rainfall, as also reported in other studies showing that rainfall had a greater influence on the variation of wet deposition fluxes [62].

The average THg (whence the HgII forms) wet deposition flux is lower than the GEOS-Chem 313 model results that produced wet Hg deposition flux around 5-6 μ g m⁻² year⁻¹ [63.64] in the areas 314 surrounding AMS. Compared with other models, such as GLEMOS, GEM-MACH-Hg, and 315 ECHMERIT, our experimental result was also lower, as can be seen from the spatial distribution 316 maps reported in Travnikov et al. [65], from which it can be inferred that the flux value in 2013 317 ranged in 2.9-4.4, 2.9-7.3, and 4.4-7.3 µg m⁻² year⁻¹, respectively. This model-to-measurement gap 318 outlines the importance of monitoring wet deposition at AMS to gather experimental results that 319 contribute to the study of the latitudinal gradient of HgII wet deposition over the oceans, hence 320 improving the understanding of Hg inputs to different ocean regions. 321

322

3.3 Influence of the El Niño Southern Oscillation (ENSO)

The El Niño Southern Oscillation (ENSO) is a cyclical meteorological event that occurs approximately every 3-7 years and causes variation of rainfall amount with consequent impact on emission phenomena, such as biomass burning due to long periods of drought [66–68]. The occurrence of El Niño has usually been detected using atmospheric indexes like SOI (Southern Oscillation Index), MEI (Multivariate ENSO Index), and ONI (Oceanic Niño Index).

From 2013-2019 (Fig. 4), ONI values reached very high levels between July 2015 and July 2016, with a peak in December-January. MEI and SOI also showed a very similar, increasing trend over the same period. However, the growth of all the indexes around 2016 was not accompanied by a sharp and direct increase in GEM, suggesting at first a limited simultaneous influence of ENSO.

On the contrary, the highest GEM concentrations were observed at the most negative SOI values

on average 6-8 months after the El Niño event (Fig. S7). This result is in agreement with the findings at Cape Point (South Africa) and Mace Head (Ireland) [67], therefore supporting the influence of El Niño on GEM variability, which can occur with the increased wildfire frequency (which also impacts CO concentration) [67] or with enhanced Hg re-emission from the ocean surface caused by the increase of the sea surface temperature [69,70]. Huang and Zhang indeed recently simulated by GEOS-Chem model a positive anomaly of atmospheric Hg concentration, lagged of 3 months, driven by Hg(0) evasion fluxes after the occurrence of El Niño [71].

In our study, we found a significant cross correlation between CO and SOI, with lag time peaking
at 4-6 months (Fig. S7) comparable to that between GEM and SOI and therefore we suggest that
ENSO-modulated biomass burning was one of the major mechanisms of GEM variability on
Amsterdam Island.

344

3.4 Backward trajectories analysis

The analysis of the 10-day backward trajectories revealed that more than 30% of the GEM peaks 345 (concentration>95th percentile) were associated with the transport of air masses coming from 346 southern Africa during the biomass burning season (Fig. S8). A similar apportionment was 347 confirmed for the high concentrations of Rn-222 (Fig. S8), whose pattern was mainly affected by 348 the emissions from the African continent. The 10-day trajectory analysis was also useful to analyze 349 350 low GEM concentrations, which were mainly linked to southerly polar air masses (Fig. S8), and low Rn values, which were also traced to air masses traveling over the ocean from the Antarctic 351 (Fig. S8), which has low local emissions of Rn, being almost entirely covered by ice [72]. 352

For each individual year, the analysis of 5-day backward trajectories reaching AMS revealed no particular differences on both annual and seasonal scales (Fig. S9), although differences could be expected on a seasonal basis at least for the summer of 2016 because of the occurrence of El Niño. The four clusters of the 10-day backward trajectories resulting from the CA were clearly associated with the marine environment, though coming from different sectors: The first cluster (C1) collected a set of trajectories coming from N-NNE; the second (C2) and fourth (C4) clusters those

coming from the SW, probably intercepting masses also from the subantarctic and Antarctic region, while the third (C3) from the NW sector collected trajectories probably intercepting masses near the African continent (Fig. S10). These results allowed us to demonstrate that air masses reaching AMS essentially traveled over the ocean, so we could assume that the origin of Hg was related to the marine environment with occasional continental sources. This outcome agrees with the literature on pollutant load at this site, which reports oceanic emissions and biomass burning as the pathways that primarily influence the occurrence of pollution events.

The results of the CA were combined with the maps obtained from the PSCF technique for GEM 366 and Rn-222, thus obtaining indications about the most likely source location of these two 367 pollutants. The map obtained for GEM over the entire 2013-2019 period showed that the highest 368 369 values of GEM were associated with clusters C1 and C3 (Fig. 5), thus supporting our hypothesis that the main contribution to GEM concentration was the emission from the subtropical ocean, 370 which contributed to depositional fluxes, as expected for marine regions [10]. The second GEM 371 source was probably of continental origin, arising from WNW air masses that crossed through 372 some areas of southern Africa, meeting the assumption of the influence of long-range transport on 373 the pollution load on Amsterdam Island. The detailed study of the PSCF maps by season did not 374 show significant differences in the PSCF probability, except for C1 where the highest values of 375 376 this probability were found during summer and IS2, two periods with higher solar radiation and temperature gradient in the air-water interface that favor the emission of elemental Hg from surface 377 water [73]. As for the lower GEM values, the PSCF confirmed the limited contribution that 378 southerly polar air masses (C4 and C2 clusters) provided to the GEM concentration at AMS. In 379 summary, our findings suggested GEM levels on Amsterdam Island are consistent with primary 380 control by evasion from subtropical marine waters and transport from the African continent. 381 Besides, considering the lack of upward/downward trend in both GEM and VWMC, we can 382 surmise that there has been no evolution in the Hg emissions intercepted at AMS from Africa. 383

A different situation was found with Rn, for which the PSCF map in the entire period showed a 384 higher probability associated with cluster C3 (Fig. S11), also providing further evidence that the 385 cluster C3 was associated with trajectories coming from southern Africa. In addition, observation 386 of the seasonal PSCF maps clarified that Rn sourced most probably from the African inland in 387 winter and was transported to AMS (Fig. S11). In support of this conclusion, previous studies of 388 air mass backward trajectories ending at AMS and Rn measurements showed that southern Africa 389 (including Madagascar) was the predominant non-marine source of trace materials transported to 390 AMS [27,74,75]. 391

4. Conclusions

Given the importance of Hg pollution control on a global scale, continuous measurements have 393 been carried out at the station on Amsterdam Island, a unique monitoring site in the Indian Ocean. 394 The GEM concentration throughout 2013-2019 was quite stationary, ranging from 0.70 to 1.58 ng 395 m⁻³, with an average value of 1.06 ± 0.07 ng m⁻³, following the typical background concentrations 396 in the Southern Hemisphere. The highest values were recorded during winter, presumably due to 397 the ITCZ variability that caused higher wind speed over the surface ocean, and the long-range 398 contribution of biomass burning from South Africa. In addition, the occurrence of the El Niño in 399 2016 led to GEM peaks which were detected at AMS with a delay of 6-8 months. 400

401 Statistical tools such as CA and PSCF showed that oceanic emission was the major driver of GEM, while Rn-222 at AMS was primarily associated with long-range transport from southern Africa. 402 In precipitation, the VWMC of THg was 2.39 ± 1.17 ng L⁻¹ on average and varied between 1.16 403 and 4.67 ng L⁻¹, which corresponded to wet deposition fluxes between 1.06 and 3.31 μ g m⁻² yr⁻¹. 404 However, the substantial lack of increasing/decreasing trend for both GEM and VWMC suggested 405 that there was no clear evolution about emission of Hg transported to AMS from Africa, hence 406 numerous efforts still need to be made for an effective long-term reduction of the concentration of 407 this pollutant to fulfill the objectives of the MCM. 408

Data availability: AMS L2 GEM data (<u>https://doi.org/10.25326/168</u>) are freely available [37] at
 <u>https://gmos.aeris-data.fr/</u> from a GMOS-FR data portal coordinated by IGE (Institut des
 Géosciences de l'Environnement, Grenoble, France; technical PI: Olivier Magand) with the
 support of the French national AERIS-SEDOO partners, data and services center for the
 atmosphere (last access: 31 January 2022).

<u>Acknowledgments:</u> We deeply thank all overwintering staff at AMS and the French Polar
Institute Paul-Emile Victor (IPEV) staff and scientists who helped with the setup and maintenance
of the experiment at AMS in the framework of the GMOStral-1028 IPEV program. AMS GEM
data were collected via instruments coordinated by the IGE-PTICHA technical platform dedicated
to atmospheric chemistry field instrumentation. The authors acknowledge the AERIS data
infrastructure for providing access to the GEM data in this study.

We thank Dr. Gianluca Bevacqua from the Language Centre of the University of Calabria andCNR-IIA for the English proofreading of the manuscript.

422 **Funding**:

The authors from CNR-IIA would like to acknowledge the contribution received from thefollowing projects:

 FET Proactive project "Towards new frontiers for distributed environmental monitoring based on an ecosystem of plant seed-like soft robots" (I-Seed). This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 101017940.

 EMPIR - EURAMET project "Metrology for traceable protocols for elemental and oxidised mercury concentrations" (SI-Hg); grant no. 19NRM03. This project (SI-Hg) has received funding from the EMPIR programme co-financed by the Participating States and from the European Union's Horizon 2020 research and innovation programme.

Amsterdam Island GEM data, accessible in GMOS-FR national database (<u>https://gmos.aeris-</u>
 <u>data.fr/</u>) have been collected with funding from European Union 7th Framework Programme

435	project Global Mercury Observation System (GMOS 2010-2015), the French Polar Institute IPEV
436	via GMOStral-1028 IPEV program since 2012, the LEFE CHAT CNRS/INSU (TOPMMODEL
437	program) and the H2020 ERA-PLANET (689443) iGOSP programme.
438	Author contribution statement:
439	
440	Antonella Tassone; Olivier Magand: Conceived and designed the experiments; Performed the
441	experiments; Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or
442	data; Wrote the paper.
443	
444	Attilio Naccarato: Conceived and designed the experiments; Analyzed and interpreted the data;
445	Contributed reagents, materials, analysis tools or data; Wrote the paper.
446	
447	Maria Martino; Domenico Amico; Hippolyte Leuridan; Michel Ramonet; Aurelien Dommergue:
448	Performed the experiments; Analyzed and interpreted the data; Contributed reagents, materials,
449	analysis tools or data; Wrote the paper.
450	
451	Francesca Sprovieri; Nicola Pirrone: Contributed reagents, materials, analysis tools or data; Wrote
452	the paper.
453	
454	Yann Bertrand: Performed the experiments; Contributed reagents, materials, analysis tools or data;
455	Wrote the paper.
456	
457	Funding statement:
458	
459	Nicola Pirrone was supported by Horizon 2020 [101017940 and 19NRM03].
460	

461	Data availability statement:
462	
463	Data associated with this study has been deposited at https://gmos.aeris-data.fr/.
464	
465	Declaration of interest's statement:
466	
467	The authors declare no conflict of interest.
468	
469	Additional information:
470	
471	Supplementary content related to this article has been published online at [URL].
472	
473	
474	

475 **References**

- 476 [1] H. Zhang, S. Wu, E.M. Leibensperger, Source-receptor relationships for atmospheric
- 477 mercury deposition in the context of global change, Atmos. Environ. 254 (2021) 118349.
- 478 https://doi.org/10.1016/j.atmosenv.2021.118349.
- 479 [2] F. Ciani, V. Rimondi, P. Costagliola, Atmospheric mercury pollution: the current
- 480 methodological framework outlined by environmental legislation, Air Qual. Atmos. Heal. 14

481 (2021) 1633–1645. https://doi.org/10.1007/s11869-021-01044-4.

- 482 [3] UNEP, Global Mercury Assessment 2018 | UNEP UN Environment Programme, (2018).
- 483 https://www.unep.org/resources/publication/global-mercury-assessment-2018 (accessed
 484 April 19, 2021).
- 485 [4] UNEP, Minamata Convention on Mercury Text and Annexes, UNEP, 2013a. (2013) 69.
- H. Selin, S.E. Keane, S. Wang, N.E. Selin, K. Davis, D. Bally, Linking science and policy to
 support the implementation of the Minamata Convention on Mercury, Ambio. 47 (2018)
 198–215. https://doi.org/10.1007/s13280-017-1003-x.
- [6] D. Amico, A. Tassone, N. Pirrone, F. Sprovieri, A. Naccarato, Recent applications and novel
 strategies for mercury determination in environmental samples using microextraction-based
- 491 approaches: A review, J. Hazard. Mater. 433 (2022) 128823.
- 492 https://doi.org/10.1016/j.jhazmat.2022.128823.
- 493 [7] A. Naccarato, A. Tassone, F. Cavaliere, R. Elliani, N. Pirrone, F. Sprovieri, A. Tagarelli, A.
- 494 Giglio, Agrochemical treatments as a source of heavy metals and rare earth elements in
- 495 agricultural soils and bioaccumulation in ground beetles, Sci. Total Environ. 749 (2020)
- 496 141438. https://doi.org/10.1016/j.scitotenv.2020.141438.
- 497 [8] A. Naccarato, A. Tassone, M. Martino, R. Elliani, F. Sprovieri, N. Pirrone, A. Tagarelli, An
- innovative green protocol for the quantification of benzothiazoles, benzotriazoles and
- benzosulfonamides in PM10 using microwave-assisted extraction coupled with solid-phase
- 500 microextraction gas chromatography tandem-mass spectrometry, Environ. Pollut. 285 (2021)

501		117487. https://doi.org/10.1016/j.envpol.2021.117487.
502	[9]	A. Naccarato, A. Tassone, M. Martino, S. Moretti, A. Macagnano, E. Zampetti, P. Papa, J.
503		Avossa, N. Pirrone, M. Nerentorp, J. Munthe, I. Wängberg, G.W. Stupple, C.P.J.J. Mitchell,
504		A.R. Martin, A. Steffen, D. Babi, E.M. Prestbo, F. Sprovieri, F. Wania, A field
505		intercomparison of three passive air samplers for gaseous mercury in ambient air, Atmos.
506		Meas. Tech. 14 (2021) 3657–3672. https://doi.org/10.5194/amt-14-3657-2021.
507	[10]	F. Sprovieri, N. Pirrone, M. Bencardino, F. D'Amore, H. Angot, C. Barbante, EG. Brunke,
508		F. Arcega-Cabrera, W. Cairns, S. Comero, M. del C. Diéguez, A. Dommergue, R.
509		Ebinghaus, X. Bin Feng, X. Fu, P.E. Garcia, B.M. Gawlik, U. Hageström, K. Hansson, M.
510		Horvat, J. Kotnik, C. Labuschagne, O. Magand, L. Martin, N. Mashyanov, T. Mkololo, J.
511		Munthe, V. Obolkin, M. Ramirez Islas, F. Sena, V. Somerset, P. Spandow, M. Vardè, C.
512		Walters, I. Wängberg, A. Weigelt, X. Yang, H. Zhang, Five-year records of mercury wet
513		deposition flux at GMOS sites in the Northern and Southern hemispheres, Atmos. Chem.
514		Phys. 17 (2017) 2689–2708. https://doi.org/10.5194/acp-17-2689-2017.
515	[11]	F. Sprovieri, N. Pirrone, M. Bencardino, F. D'Amore, F. Carbone, S. Cinnirella, V.
516		Mannarino, M. Landis, R. Ebinghaus, A. Weigelt, E.G. Brunke, C. Labuschagne, L. Martin,
517		J. Munthe, I. Wängberg, P. Artaxo, F. Morais, H. De Melo Jorge Barbosa, J. Brito, W.
518		Cairns, C. Barbante, M. Del Carmen Diéguez, P. Elizabeth Garcia, D. Aurélien, H. Angot, O.
519		Magand, H. Skov, M. Horvat, J. Kotnik, K. Alana Read, L. Mendes Neves, B. Manfred
520		Gawlik, F. Sena, N. Mashyanov, V. Obolkin, D. Wip, X. Bin Feng, H. Zhang, X. Fu, R.
521		Ramachandran, D. Cossa, J. Knoery, N. Marusczak, M. Nerentorp, C. Norstrom,
522		Atmospheric mercury concentrations observed at ground-based monitoring sites globally
523		distributed in the framework of the GMOS network, Atmos. Chem. Phys. 16 (2016) 11915-
524		11935. https://doi.org/10.5194/acp-16-11915-2016.
525	[12]	GR. Sheu, NH. Lin, Characterizations of wet mercury deposition to a remote islet

526 (Pengjiayu) in the subtropical Northwest Pacific Ocean, Atmos. Environ. 77 (2013) 474–481.

- 527 https://doi.org/10.1016/j.atmosenv.2013.05.038.
- 528 [13] Sheu, Gay, Schmeltz, Olson, Chang, Lin, Nguyen, A New Monitoring Effort for Asia: The
 529 Asia Pacific Mercury Monitoring Network (APMMN), Atmosphere (Basel). 10 (2019) 481.
 530 https://doi.org/10.3390/atmos10090481.
- [14] K.A. Pfaffhuber, T. Berg, D. Hirdman, A. Stohl, Atmospheric mercury observations from
 Antarctica: seasonal variation and source and sink region calculations, Atmos. Chem. Phys.
 12 (2012) 3241–3251. https://doi.org/10.5194/acp-12-3241-2012.
- 534 [15] F. Slemr, H. Angot, A. Dommergue, O. Magand, M. Barret, A. Weigelt, R. Ebinghaus, E.G.
- 535 Brunke, K.A. Pfaffhuber, G. Edwards, D. Howard, J. Powell, M. Keywood, F. Wang,
- Comparison of mercury concentrations measured at several sites in the Southern Hemisphere,
 Atmos. Chem. Phys. 15 (2015) 3125–3133. https://doi.org/10.5194/acp-15-3125-2015.
- 538 [16] F. Slemr, L. Martin, C. Labuschagne, T. Mkololo, H. Angot, O. Magand, A. Dommergue, P.
- 539 Garat, M. Ramonet, J. Bieser, Atmospheric mercury in the Southern Hemisphere Part 1:
- 540 Trend and inter-annual variations in atmospheric mercury at Cape Point, South Africa, in
- 541 2007–2017, and on Amsterdam Island in 2012–2017, Atmos. Chem. Phys. 20 (2020) 7683–
- 542 7692. https://doi.org/10.5194/acp-20-7683-2020.
- 543 [17] H. Angot, M. Barret, O. Magand, M. Ramonet, A. Dommergue, A 2-year record of
- atmospheric mercury species at a background Southern Hemisphere station on Amsterdam

Island, Atmos. Chem. Phys. 14 (2014) 11461–11473. https://doi.org/10.5194/acp-14-114612014.

- 547 [18] H. Angot, A. Dastoor, F. De Simone, K. Gärdfeldt, C.N. Gencarelli, I.M. Hedgecock, S.
- 548 Langer, O. Magand, M.N. Mastromonaco, C. Nordstrøm, K.A. Pfaffhuber, N. Pirrone, A.
- 549 Ryjkov, N.E. Selin, H. Skov, S. Song, F. Sprovieri, A. Steffen, K. Toyota, O. Travnikov, X.
- 550 Yang, A. Dommergue, Chemical cycling and deposition of atmospheric mercury in polar
- regions: Review of recent measurements and comparison with models, Atmos. Chem. Phys.
- 552 16 (2016) 10735–10763. https://doi.org/10.5194/ACP-16-10735-2016.

553	[19]	H. Angot, I. Dion, N. Vogel, M. Legrand, O. Magand, A. Dommergue, Multi-year record of
554		atmospheric mercury at Dumont d'Urville, East Antarctic coast: Continental outflow and
555		oceanic influences, Atmos. Chem. Phys. 16 (2016) 8265-8279. https://doi.org/10.5194/ACP-
556		16-8265-2016.

- E.-G. Brunke, C. Walters, T. Mkololo, L. Martin, C. Labuschagne, B. Silwana, F. Slemr, A. 557 [20] Weigelt, R. Ebinghaus, V. Somerset, Mercury in the atmosphere and in rainwater at Cape
- Point, South Africa, Atmos. Environ. 125 (2016) 24–32. 559
- https://doi.org/10.1016/j.atmosenv.2015.10.059. 560

- E.-G. Brunke, R. Ebinghaus, H.H. Kock, C. Labuschagne, F. Slemr, Emissions of mercury in 561 [21] southern Africa derived from long-term observations at Cape Point, South Africa, Atmos. 562
- 563 Chem. Phys. 12 (2012) 7465–7474. https://doi.org/10.5194/acp-12-7465-2012.
- M.C. Diéguez, M. Bencardino, P.E. García, F. D'Amore, J. Castagna, F. De Simone, C. Soto 564 [22] Cárdenas, S. Ribeiro Guevara, N. Pirrone, F. Sprovieri, A multi-year record of atmospheric 565
- mercury species at a background mountain station in Andean Patagonia (Argentina): 566
- Temporal trends and meteorological influence, Atmos. Environ. 214 (2019) 116819. 567
- https://doi.org/10.1016/j.atmosenv.2019.116819. 568
- A.M. Koenig, O. Magand, P. Laj, M. Andrade, I. Moreno, F. Velarde, G. Salvatierra, R. 569 [23]
- 570 Gutierrez, L. Blacutt, D. Aliaga, T. Reichler, K. Sellegri, O. Laurent, M. Ramonet, A.
- Dommergue, Seasonal patterns of atmospheric mercury in tropical South America as inferred 571
- by a continuous total gaseous mercury record at Chacaltaya station (5240 m) in Bolivia, 572
- Atmos. Chem. Phys. 21 (2021) 3447–3472. https://doi.org/10.5194/acp-21-3447-2021. 573
- M. Martino, A. Tassone, L. Angiuli, A. Naccarato, P.R. Dambruoso, F. Mazzone, L. Trizio, [24] 574
- C. Leonardi, F. Petracchini, F. Sprovieri, N. Pirrone, F. D'Amore, M. Bencardino, First 575
- atmospheric mercury measurements at a coastal site in the Apulia region: seasonal variability 576
- and source analysis, Environ. Sci. Pollut. Res. (2022) 16. https://doi.org/10.1007/s11356-577
- 022-20505-6. 578

579	[25]	J. Bieser, H. Angot, F. Slemr, L. Martin, Atmospheric mercury in the Southern Hemisphere –
580		Part 2: Source apportionment analysis at Cape Point station, South Africa, Atmos. Chem.
581		Phys. 20 (2020) 10427–10439. https://doi.org/10.5194/acp-20-10427-2020.
582	[26]	A. Gaudry, P. Monfray, G. Polian, G. Bonsang, B. Ardouin, A. Jegou, G. Lambert, Non-
583		seasonal variations of atmospheric CO 2 concentrations at Amsterdam Island, Tellus B
584		Chem. Phys. Meteorol. 43 (1991) 136–143. https://doi.org/10.3402/tellusb.v43i2.15258.
585	[27]	V. Gros, B. Bonsang, D. Martin, P.C. Novelli, V. Kazan, Carbon monoxide short term
586		measurements at Amsterdam island: estimations of biomass burning emission rates,
587		Chemosph Glob. Chang. Sci. 1 (1999) 163–172. https://doi.org/10.1016/S1465-
588		9972(99)00009-4.
589	[28]	V. Gros, B. Bonsang, R. Sarda Esteve, Atmospheric carbon monoxide 'in situ' monitoring by
590		automatic gas chromatography, Chemosph Glob. Chang. Sci. 1 (1999) 153-161.
591		https://doi.org/10.1016/S1465-9972(99)00010-0.
592	[29]	M. Jiskra, J.E. Sonke, D. Obrist, J. Bieser, R. Ebinghaus, C.L. Myhre, K.A. Pfaffhuber, I.
593		Wängberg, K. Kyllönen, D. Worthy, L.G. Martin, C. Labuschagne, T. Mkololo, M. Ramonet,
594		O. Magand, A. Dommergue, A vegetation control on seasonal variations in global
595		atmospheric mercury concentrations, Nat. Geosci. 11 (2018) 244-250.
596		https://doi.org/10.1038/s41561-018-0078-8.
597	[30]	C. Yver-Kwok, C. Philippon, P. Bergamaschi, T. Biermann, F. Calzolari, H. Chen, S. Conil,
598		P. Cristofanelli, M. Delmotte, J. Hatakka, M. Heliasz, O. Hermansen, K. Kominkova, D.
599		Kubistin, N. Kumps, O. Laurent, T. Laurila, I. Lehner, J. Levula, M. Lindauer, M. Lopez, I.
600		Mammarella, G. Manca, P. Marklund, J.M. Metzger, M. Mölder, S.M. Platt, M. Ramonet, L.
601		Rivier, B. Scheeren, M. Kumar Sha, P. Smith, M. Steinbacher, G. Vítková, S. Wyss,
602		Evaluation and optimization of ICOS atmosphere station data as part of the labeling process,
603		Atmos. Meas. Tech. 14 (2021) 89–116. https://doi.org/10.5194/AMT-14-89-2021.
604	[31]	S.D. Chambers, S. Preunkert, R. Weller, SB. Hong, R.S. Humphries, L. Tositti, H. Angot,

- 605 M. Legrand, A.G. Williams, A.D. Griffiths, J. Crawford, J. Simmons, T.J. Choi, P.B.
- 606 Krummel, S. Molloy, Z. Loh, I. Galbally, S. Wilson, O. Magand, F. Sprovieri, N. Pirrone, A.
- 607 Dommergue, Characterizing Atmospheric Transport Pathways to Antarctica and the Remote
- 608 Southern Ocean Using Radon-222, Front. Earth Sci. 6 (2018) 1–28.
- 609 https://doi.org/10.3389/feart.2018.00190.
- 610 [32] A. Gaudry, G. Polian, B. Ardouin, G. Lambert, Radon-calibrated emissions of CO 2 from
- 611 South Africa, Tellus B Chem. Phys. Meteorol. 42 (1990) 9–19.
- 612 https://doi.org/10.3402/tellusb.v42i1.15187.
- [33] L. Harrison, C. Funk, A. McNally, S. Shukla, G. Husak, Pacific sea surface temperature
- 614 linkages with Tanzania's multi-season drying trends, Int. J. Climatol. 39 (2019) 3057–3075.
- 615 https://doi.org/10.1002/joc.6003.
- 616 [34] F. D'Amore, M. Bencardino, S. Cinnirella, F. Sprovieri, N. Pirrone, Data quality through a
- 617 web-based QA/QC system: Implementation for atmospheric mercury data from the global
- 618 mercury observation system, Environ. Sci. Process. Impacts. 17 (2015) 1482–1491.
- 619 https://doi.org/10.1039/c5em00205b.
- 620 [35] A. Tassone, S. Moretti, M. Martino, N. Pirrone, F. Sprovieri, A. Naccarato, Modification of
- the EPA method 1631E for the quantification of total mercury in natural waters, MethodsX. 7
- 622 (2020) 100987. https://doi.org/10.1016/j.mex.2020.100987.
- [36] GMOS, Method for the determination of total mercury in precipitation, GMOS. (2011) 1–22.
- 624 [37] O. Magand, A. Dommergue, Continuous measurements of atmospheric mercury at
- Amsterdam Island (L2) [Dataset]. Aeris., (2021).
- 626 https://doi.org/https://doi.org/10.25326/168.
- [38] P.K. Sen, Estimates of the Regression Coefficient Based on Kendall's Tau, J. Am. Stat.
 Assoc. 63 (1968) 1379. https://doi.org/10.2307/2285891.
- 629 [39] M.J. McGill, R.J. Swap, J.E. Yorks, P.A. Selmer, S.J. Piketh, Observation and quantification
- of aerosol outflow from southern Africa using spaceborne lidar, S. Afr. J. Sci. 116 (2020)

- 631 65–70. https://doi.org/10.17159/sajs.2020/6398.
- [40] E. Baboukas, J. Sciare, N. Mihalopoulos, Spatial, Temporal and Interannual Variability of
- 633 Methanesulfonate and Non-Sea-Salt Sulfate in Rainwater in the Southern Indian Ocean
- 634 (Amsterdam, Crozet and Kerguelen Islands), J. Atmos. Chem. 48 (2004) 35–57.
- 635 https://doi.org/10.1023/B:JOCH.0000034508.13326.cd.
- 636 [41] R.R. Draxler, G.D. Rolph, HYSPLIT (HYbrid Single-Particle Lagrangian Integrated
- 637 Trajectory), NOAA Air Resour. Lab. Coll. Park. MD. (2003).
- http://www.arl.noaa.gov/ready/hysplit4.html (accessed May 3, 2022).
- [42] J.L. Moody, P.J. Samson, The influence of atmospheric transport on precipitation chemistry
- at two sites in the midwestern United States, Atmos. Environ. 23 (1989) 2117–2132.
- 641 https://doi.org/10.1016/0004-6981(89)90173-X.
- [43] J.H. Ward, Hierarchical Grouping to optimize an objective function, J. Am. Stat. Assoc. 58
 (1963) 236–244.
- [44] M.S. Landis, G.J. Keeler, Atmospheric mercury deposition to Lake Michigan during the
- Lake Michigan Mass Balance Study, Environ. Sci. Technol. (2002).
- 646 https://doi.org/10.1021/es011217b.
- [45] D.S. Wilks, Statistical Methods in the Atmospheric Sciences, Fourth, Elsevier, Oxford, 2019.
 https://doi.org/10.1016/C2017-0-03921-6.
- [46] M. Anjos, A.C. Targino, P. Krecl, G.Y. Oukawa, R.F. Braga, Analysis of the urban heat
- island under different synoptic patterns using local climate zones, Build. Environ. 185 (2020)
- 651 107268. https://doi.org/10.1016/J.BUILDENV.2020.107268.
- 652 [47] R.R. Draxler, HYSPLIT4 User 's Guide, Silver Spring, USA, 1999.
- 653 http://www.arl.noaa.gov/hysplit.html.
- [48] Z.L. Fleming, P.S. Monks, A.J. Manning, Review: Untangling the influence of air-mass
 history in interpreting observed atmospheric composition, Atmos. Res. (2012).
- [49] N. Pekney, C. Davidson, L. Zhou, P. Hopke, Application of PSCF and CPF to PMF-modeled

- 657 sources of PM2.5 in Pittsburgh, in: Aerosol Sci. Technol., 2006: pp. 952–961.
- 658 https://doi.org/10.1080/02786820500543324.
- [50] K.R. Travis, C.L. Heald, H.M. Allen, E.C. Apel, S.R. Arnold, D.R. Blake, W.H. Brune, X.
- 660 Chen, R. Commane, J.D. Crounse, B.C. Daube, G.S. Diskin, J.W. Elkins, M.J. Evans, S.R.
- Hall, E.J. Hintsa, R.S. Hornbrook, P.S. Kasibhatla, M.J. Kim, G. Luo, K. McKain, D.B.
- Millet, F.L. Moore, J. Peischl, T.B. Ryerson, T. Sherwen, A.B. Thames, K. Ullmann, X.
- 663 Wang, P.O. Wennberg, G.M. Wolfe, F. Yu, Constraining remote oxidation capacity with
- ATom observations, Atmos. Chem. Phys. 20 (2020) 7753–7781. https://doi.org/10.5194/acp-
- 66520-7753-2020.
- 666 [51] D.P. Edwards, L.K. Emmons, J.C. Gille, A. Chu, J.L. Attié, L. Giglio, S.W. Wood, J.
- 667 Haywood, M.N. Deeter, S.T. Massie, D.C. Ziskin, J.R. Drummond, Satellite-observed
- pollution from Southern Hemisphere biomass burning, J. Geophys. Res. Atmos. 111 (2006)
- 669 1–17. https://doi.org/10.1029/2005JD006655.
- [52] M.A.K. Khalil, R.A. Rasmussen, Sources, sinks, and seasonal cycles of atmospheric
- 671 methane, J. Geophys. Res. Ocean. 88 (1983) 5131–5144.
- 672 https://doi.org/10.1029/JC088iC09p05131.
- [53] M.A.K. Khalil, R.A. Rasmussen, The global cycle of carbon monoxide: Trends and mass
 balance, Chemosphere. 20 (1990) 227–242. https://doi.org/10.1016/0045-6535(90)90098-E.
- [54] J. Williams, V. Gros, B. Bonsang, V. Kazan, HO cycle in 1997 and 1998 over the southern
- Indian Ocean derived from CO, radon, and hydrocarbon measurements made at Amsterdam
- 677 Island, J. Geophys. Res. Atmos. 106 (2001) 12719–12725.
- 678 https://doi.org/10.1029/2001JD900116.
- [55] E.-G. Brunke, C. Labuschagne, B. Parker, H.E. Scheel, S. Whittlestone, Baseline air mass
- selection at Cape Point, South Africa: application of 222Rn and other filter criteria to CO2,
- 681 Atmos. Environ. 38 (2004) 5693–5702. https://doi.org/10.1016/j.atmosenv.2004.04.024.
- [56] D.S. McLagan, C.P.J. Mitchell, A. Steffen, H. Hung, C. Shin, G.W. Stupple, M.L. Olson,

- 683 W.T. Luke, P. Kelley, D. Howard, G.C. Edwards, P.F. Nelson, H. Xiao, G.R. Sheu, A.
- 684 Dreyer, H. Huang, B. Abdul Hussain, Y.D. Lei, I. Tavshunsky, F. Wania, Global evaluation
- and calibration of a passive air sampler for gaseous mercury, Atmos. Chem. Phys. 18 (2018)
- 686 5905–5919. https://doi.org/10.5194/acp-18-5905-2018.
- [57] J.A. Fisher, P.F. Nelson, Atmospheric mercury in Australia, Elem. Sci. Anthr. 8 (2020) 1–20.
 https://doi.org/10.1525/elementa.2020.070.
- [58] C. Liu, X. Liao, J. Qiu, Y. Yang, X. Feng, R.P. Allan, N. Cao, J. Long, J. Xu, Observed
 variability of intertropical convergence zone over 1998-2018, Environ. Res. Lett. 15 (2020).
- 691 https://doi.org/10.1088/1748-9326/aba033.
- [59] H. Lashkari, M. Jafari, Annual displacement and appropriate index to determine ITCZ
- position in East Africa and the Indian Ocean regions, Meteorol. Atmos. Phys. 133 (2021)
- 694 1111–1126. https://doi.org/10.1007/s00703-021-00797-y.
- [60] G. Liu, Y. Cai, N. O'Driscoll, Environmental Chemistry and Toxicology of Mercury, 2011.
 https://doi.org/10.1002/9781118146644.
- 697 [61] Y.-S. Seo, Y.-J. Han, H.-D. Choi, T.M. Holsen, S.-M. Yi, Characteristics of total mercury
- (TM) wet deposition: Scavenging of atmospheric mercury species, Atmos. Environ. 49
 (2012) 69–76. https://doi.org/10.1016/j.atmosenv.2011.12.031.
- [62] X. Fu, X. Yang, X. Lang, J. Zhou, H. Zhang, B. Yu, H. Yan, C.J. Lin, X. Feng, Atmospheric
 wet and litterfall mercury deposition at urban and rural sites in China, Atmos. Chem. Phys.

702 16 (2016) 11547–11562. https://doi.org/10.5194/acp-16-11547-2016.

- 703 [63] V. Shah, D.J. Jacob, C.P. Thackray, X. Wang, E.M. Sunderland, T.S. Dibble, A. Saiz-Lopez,
- I. Cernusák, V. Kellö, P.J. Castro, R. Wu, C. Wang, Improved mechanistic model of the
- atmospheric redox chemistry of mercury, Environ. Sci. Technol. 55 (2021) 14445–14456.
- 706 https://doi.org/10.1021/acs.est.1c03160.
- 707 [64] H.M. Horowitz, D.J. Jacob, Y. Zhang, T.S. DIbble, F. Slemr, H.M. Amos, J.A. Schmidt, E.S.
- 708 Corbitt, E.A. Marais, E.M. Sunderland, A new mechanism for atmospheric mercury redox

- chemistry: Implications for the global mercury budget, Atmos. Chem. Phys. 17 (2017) 6353–
 6371. https://doi.org/10.5194/acp-17-6353-2017.
- 711 [65] O. Travnikov, H. Angot, P. Artaxo, M. Bencardino, J. Bieser, F. D'Amore, A. Dastoor, F. De
- 512 Simone, M.C. DIéguez, A. Dommergue, R. Ebinghaus, X. Bin Feng, C.N. Gencarelli, I.M.
- Hedgecock, O. Magand, L. Martin, V. Matthias, N. Mashyanov, N. Pirrone, R.
- Ramachandran, K. Alana Read, A. Ryjkov, N.E. Selin, F. Sena, S. Song, F. Sprovieri, D.
- 715 Wip, I. Wängberg, X. Yang, Multi-model study of mercury dispersion in the atmosphere:
- Atmospheric processes and model evaluation, Atmos. Chem. Phys. 17 (2017) 5271–5295.
- 717 https://doi.org/10.5194/acp-17-5271-2017.
- 718 [66] R. V. Rohli, A.J. Vega, Climatology, Jones & Bartlett Learning, 2015.
- 719 [67] F. Slemr, C.A. Brenninkmeijer, A. Rauthe-Schöch, A. Weigelt, R. Ebinghaus, E. Brunke, L.
- 720 Martin, T.G. Spain, S. O'Doherty, El Niño–Southern Oscillation influence on tropospheric
- mercury concentrations, Geophys. Res. Lett. 43 (2016) 1766–1771.
- 722 https://doi.org/10.1002/2016GL067949.
- 723 [68] R.A. Preston-Whyte, P.D. Tyson, Atmosphere and weather of southern Africa, Oxford
- 724 University Press, 1988. https://agris.fao.org/agris-
- search/search.do?recordID=US201300685112 (accessed November 19, 2021).
- [69] F. Carbone, M.S. Landis, C.N. Gencarelli, A. Naccarato, F. Sprovieri, F. De Simone, I.M.
- Hedgecock, N. Pirrone, Sea surface temperature variation linked to elemental mercury
- concentrations measured on Mauna Loa, Geophys. Res. Lett. (2016).
- 729 https://doi.org/10.1002/2016GL069252.
- [70] L.S.P. Nguyen, K.T. Nguyen, S.M. Griffith, G.R. Sheu, M.C. Yen, S.C. Chang, N.H. Lin,
- 731 Multiscale Temporal Variations of Atmospheric Mercury Distinguished by the Hilbert–
- Huang Transform Analysis Reveals Multiple El Niño–Southern Oscillation Links, Environ.
- 733 Sci. Technol. 56 (2022) 1423–1432. https://doi.org/10.1021/acs.est.1c03819.
- [71] S. Huang, Y. Zhang, Interannual Variability of Air-Sea Exchange of Mercury in the Global

- 735 Ocean: The "seesaw Effect" in the Equatorial Pacific and Contributions to the Atmosphere,
- 736 Environ. Sci. Technol. 55 (2021) 7145–7156. https://doi.org/10.1021/acs.est.1c00691.
- 737 [72] G. Polian, G. Lambert, B. Ardouin, A. Jegou, Long-range transport of continental radon in
- subantarctic and antarctic areas, Tellus B Chem. Phys. Meteorol. 38 (1986) 178–189.
- 739 https://doi.org/10.3402/tellusb.v38i3-4.15126.
- 740 [73] C. deW. Rautenbach, I. Smith, Teleconnections between global sea-surface temperatures
- and the interannual variability of observed and model simulated rainfall over southern Africa,
- 742 J. Hydrol. 254 (2001) 1–15. https://doi.org/10.1016/S0022-1694(01)00454-1.
- 743 [74] J.M. Miller, J.L. Moody, J.M. Harris, A. Gaudry, A 10-year trajectory flow climatology for
- Amsterdam Island, 1980–1989, Atmos. Environ. Part A. Gen. Top. 27 (1993) 1909–1916.
- 745 https://doi.org/10.1016/0960-1686(93)90296-B.
- [75] M. Ramonet, J.C. Le Roulley, P. Bousquet, P. Monfray, Radon-222 Measurements During
- the Tropoz II Campaign and Comparison With a Global Atmospheric Transport Model, J.
- 748 Of Atmospheric Chem. 23 (1996) 107–136.
- 749

750 **Figure captions**

Fig. 1. The plot of the seasonal volume-weighted mean concentration (VWMC) of total mercury
in rainwater and rainfall amount in the 2013-2019 period (IS1 = M-A-M-J; austral winter = J-AS; IS2 = O-N; austral summer = D-J-F).

- Fig. 2. Temporal variability of Gaseous Elemental Mercury (GEM) from March 2013 to
 December 2019: Individual hourly mean values (left panel) and monthly median values (right
 panel). The shadings in the monthly median plot represent 25/75th and 5/95th quantile values.
- Fig. 3. Correlation plots of seasonal averages of VWMC versus total rainfall (upper left); GEM
 versus total rainfall (upper right); VWMC versus GEM (lower left); Wet deposition flux versus
 rainfall (lower right).
- 760Fig. 4. Temporal plot of the SOI, ONI, MEI indexes during the monitored period useful to assess761ENSO occurrence and trend of GEM over the same period. (For a better visualization, SOI was762sign-changed using multiplication by -1). The exceedance of the threshold of ± 0.5 °C of the ONI763for five consecutive months is used as evidence of El Niño (+0.5 °C), as well as large positive764values of -SOI and MEI. The light green box indicates the occurrence of the El Niño.
- Fig. 5. The plot of the PSCF probabilities for GEM concentrations (90th percentile) at AMS in
 the 2013-2019 period. The round black dot indicates the location of Amsterdam Island.
- 767
- 768

769	Table 1. Annual mean (± standard deviation) of the meteorological variables and monitored
770	gases, namely CO ₂ , CO, Rn-222, CH ₄ , and Gaseous Elemental Mercury (GEM) at AMS from
771	March 2013 to December 2019. Annual mean (± standard deviation) of volume-weighted mean
772	concentration (VWMC) of total mercury in rainwater and wet deposition fluxes are also reported.

773 (n.a.=not available)

	Temperature (°C)	Rainfall (mm)	CO ₂ (ppm)	CO (ppb)	Rn-222 (mBq m ⁻ ³)	CH4 (ppb)	GEM (ng m ⁻³)	VWMC (ng L ⁻¹)	Wet deposition flux (µg m ⁻² year ⁻¹)
2013	13.7 ± 2.2	833	393.4 ± 0.8	n.a.	48.1 ± 19.4	1769 ± 11	1.04 ± 0.09	2.34 ± 0.93	1.95 ± 0.15
2014	14.2 ± 2.8	864	395.2 ± 0.9	n.a.	41.7 ± 14.2	1774 ± 15	$\begin{array}{c} 1.05 \pm \\ 0.06 \end{array}$	1.80 ± 1.07	1.55 ± 0.33
2015	14.4 ± 2.8	916	397.5 ± 0.9	50.0 ± 3.3	46.8 ± 13.4	1786 ± 14	1.02 ± 0.07	1.16 ± 0.30	1.06 ± 0.13
2016	14.5 ± 2.9	967	400.7 ± 1.0	52.1 ± 6.7	24.1 ± 8.3	1792 ± 12	1.08 ± 0.06	3.04 ± 1.39	2.94 ± 0.44
2017	15.0 ± 2.9	894	402.7 ± 0.9	45.2 ± 6.8	32.0 ± 7.5	1799 ± 15	1.02 ± 0.08	2.18 ± 1.63	1.95 ± 0.31
2018	14.8 ± 3.3	855	405.2 ± 0.9	42.8 ± 8.1	31.1 ± 8.8	1810 ± 14	1.08 ± 0.05	4.67 ± 3.95	3.31 ± 0.58
2019	14.3 ± 3.0	1003	407.6 ± 1.1	50.5 ± 7.8	26.9 ± 7.1	1817 ± 13	1.08 ± 0.06	1.55 ± 0.47	1.55 ± 0.26



rainfall (mm)









Highlights

Gaseous Elemental Mercury at Amsterdam Island varied between 0.70 and 1.58 ng m⁻³ Total mercury concentration in wet deposition ranged from 1.16 to 4.67 ng L⁻¹ CO, CO₂, and CH₄ concentrations showed upward significant trends from 2013 to 2019 Gaseous Elemental Mercury was affected by El Niño-modulated biomass burning Atmospheric mercury was governed by oceanic evasion and long-range transport

political level of the second se