



HAL
open science

Association between prolonged exposure to long working hours and stroke subtypes in the CONSTANCES cohort

Marc Fadel, Grace Sembajwe, Jian Li, Annette Leclerc, Fernando Pico, Alexis Schnitzler, Yves Roquelaure, Alexis Descatha

► To cite this version:

Marc Fadel, Grace Sembajwe, Jian Li, Annette Leclerc, Fernando Pico, et al.. Association between prolonged exposure to long working hours and stroke subtypes in the CONSTANCES cohort. Occupational and Environmental Medicine, 2023, 80 (4), pp.196-201. 10.1136/oemed-2022-108656 . hal-04041101

HAL Id: hal-04041101

<https://hal.science/hal-04041101v1>

Submitted on 26 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Association between prolonged exposure to long-working hours and stroke subtypes in the CONSTANCES cohort

Marc Fadel, MD¹, Grace Sembajwe, ScD², Jian Li, PhD³, Annette Leclerc, PhD⁴, Fernando Pico, PhD^{5,6}, Alexis Schnitzler, PhD⁷, Yves Roquelaure, PhD¹, Alexis Descatha, PhD^{1,2}

1. Univ Angers, CHU Angers, Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, IRSET-ESTER, SFR ICAT, CAPTV CDC, F-49000, Angers, France
2. Department of Occupational Medicine, Epidemiology and Prevention, Donald and Barbara Zucker School of Medicine, Hofstra University Northwell Health, New York, USA
3. Department of Environmental Health Sciences, Fielding School of Public Health, School of Nursing, University of California, Los Angeles, CA, USA
4. Inserm, UMS 011, Villejuif, France
5. Neurology and stroke unit, Versailles Mignot Hospital, Le Chesnay, France
6. Versailles Saint Quentin en Yvelines Paris Saclay Univ, Versailles, France
7. PRM Department, Hôpital Lariboisière-F.Widal AP-HP, Paris, France

Correspondence: Dr Marc Fadel, Inserm, Irset - UMR_S 1085, Faculté de Santé, Département

Médecine 28 rue Roger Amsler, CS 7452, 49045 ANGERS Cedex 01, Angers, France

(marc.fadel@univ-angers.fr, Tel : +33(2)41735912)

Words count: 1876, 2 tables, 19 references

Abstract

Background: Though there is increasing evidence on the effect of long working hours (LWH) and stroke, few studies have distinguished stroke subtypes. We examined the associations between LWH and ischemic or hemorrhagic stroke after adjusting for cardiovascular risk factors.

Methods: From a national population-based cohort CONSTANCES, baseline questionnaires and initial health examinations were used to retrieve sociodemographic and cardiovascular risk factors from 2012 to 2018. LWH were defined as self-reported working time ≥ 10 hours daily for at least 50 days per year. Incident cases of stroke were collected using International Classification of Disease codes recorded in the National health data system. Associations between LWH and stroke were investigated using multinomial models adjusted for cardiovascular risk factors.

Results: Among the 160,751 participants who were free from stroke at baseline, exposure to LWH ≥ 10 years was reported by 20,723 participants, and 190 incident cases of stroke were identified, including 134 ischemic and 56 hemorrhagic. Exposure to LWH was associated with an elevated odds of ischemic stroke [OR=1.61(1.04-2.49)] and hemorrhagic stroke [OR=2.50(1.38-4.53)] in unadjusted models. In adjusted multivariable models, only the LWH association with hemorrhagic stroke remained significant [aOR=1.92(1.01-3.09)].

Conclusions: LWH were associated with stroke, though it remained significant for hemorrhagic stroke only after adjustments. Differences in direct and indirect biological pathways and lack of power in the ischemic subgroup may explain these results and further studies on the impact of mediating and effect measure modifying factors are needed. Nevertheless, policies that attenuate effects of both LWH and cardiovascular risks factor are warranted.

Keywords: stroke, work, long-working hours, epidemiology, occupational

Key messages:

What is already known about this subject?

- More than 400 000 deaths by stroke in 2016 were estimated to be caused by long-working hours.
- Few studies have differentiated the effect of long-working hours on ischemic and hemorrhagic stroke while considering strong cardiovascular risk factors.

What are the new findings?

- Prolonged exposure to long-working hours was associated with increased stroke occurrences in unadjusted models, but the association remained significant only for hemorrhagic stroke after adjustments.

How might this impact on policy or clinical practice in the foreseeable future?

- Lack of power and differences in direct and indirect biological pathways may explain this result, with a predominant mediation by modifiable behaviors for ischemic strokes.
- More studies, even underpowered ones, are needed to better understand the association between long working hours, cardiovascular risk factors and stroke.
- Better controls and interventions on both occupational exposures and lifestyle factors are needed to reduce the global burden of stroke.

INTRODUCTION

The effect of long-working hours (LWH) on cardiovascular diseases has been widely documented.(1) Systematic reviews and meta-analyses from the WHO/ILO joint estimates of disease reported sufficient evidence of harmfulness associated with working more than 55 hours per week for stroke and ischemic heart diseases.(2,3) Though the strength of the association between LWH and stroke was moderate compared to other occupational exposures, the high prevalence of LWH lead to an estimated 400 000 deaths by stroke in 2016, with a population-attributable fraction close to 7%.(1) Stroke, however, is a broad term that conflates two different types of disease with different pathophysiology: ischemic stroke is caused mainly by the clotting of a cerebral artery and hemorrhagic stroke is caused by the rupture of a cerebral artery.(4,5) The WHO/ILO systematic review did not report enough information to differentiate the effect of LWH on risk of stroke subtypes. Among the reviewed studies, some reported a higher risk of hemorrhagic stroke.(6,7) These studies, however, did not consider potentially strong confounding and effect measure modifying factors, such as diabetes and smoking, nor did they account for duration of exposure to LWH, which has been shown to influence the association between LWH and stroke.(8,9). However, there is little evidence on the confounding factors on the relation between LWH and stroke. Thus, future models assessing the effect of LWH on stroke should at least include known cardiovascular risk factors and consider behavioral risk factors depending on their relevance.

To better understand the relationship between LWH and stroke, our aim was to examine a large population cohort for associations between LWH and incident cases of ischemic and hemorrhagic stroke, while accounting for an array of cardiovascular risk factors.

METHODS

Data origin

CONSTANCES is a French population-based cohort created in 2012. Adults enrolled in the National Social Security system and aged 18 to 69 years were randomly selected. The cohort design and characteristics are detailed elsewhere.⁽¹⁰⁾ All participants underwent health screenings by medical professionals at Social Security affiliated centers and completed self-administered surveys that included “lifestyle” and “occupational exposure” questionnaires. Data from the cohort were linked to French National Health Insurance records to retrieve cases of stroke during the follow-up period (until 2018). These records are collected in the national health data system (SNDS) which evaluate all health care utilization in our country.⁽¹¹⁾ All working participants who completed the questionnaires during the data collection period (from 2012 to 2018) were included.

Variables of interest

From the self-administered questionnaires at baseline, we retrieved the following self-reported relevant variables: age (years), sex (female, male), smoking (never smoked, ever smoked <30 pack-years, ever smoked \geq 30 pack-years), occupational status using the French occupational and social classification system⁽¹²⁾ with the first digit (high-skilled white-collar jobs, Self-employed/Chief Executive Officer/Professional jobs, low-skilled white-collar jobs, blue-collar jobs), and exposure to LWH. Exposure to LWH was defined as a working time greater than 10 hours daily for at least 50 days per year. Participants also reported the number of years they were exposed. In the statistical analyses, exposure to LWH was categorized as <10 years and \geq 10 years of exposure consistent with sensitivity analyses from previous findings.^(8,9)

We retrieved data on history of stroke (all subtypes together) and age of occurrence, diagnosed diabetes (yes/no), diagnosed high blood pressure (yes/no), diagnosed dyslipidemia, hypercholesterolemia or hypertriglyceridemia (yes/no), family history of cardiovascular events (yes/ no), and body mass index (kg/m²) from the medical interview at baseline where physicians questioned participants on medical diseases and examined them.

Main outcome

Cases of stroke were retrieved via a linkage between the CONSTANCES data and the National Health Data System (SNDS), which collects ICD-10 (International Classification of Diseases) codes from all hospitalization across the country. We retrieved the principal diagnosis that allowed us to categorize stroke as hemorrhagic (I60-61-62 except I604/I605/I621) or ischemic (I63-64 except I63.6). Codes were chosen accordingly to a previous study on stroke in the National Health Data System.(13) Only the first case of stroke after study inclusion was considered. Cases of stroke that occurred before participants filled out the baseline questionnaire were not included, as well as participants with a history of stroke were excluded.

Statistical analyses

Associations between LWH and stroke were examined using crude and adjusted multinomial logistic regression models. Multivariable models were adjusted for cardiovascular confounding variables, including age, sex, BMI, high blood pressure, diabetes, dyslipidemia, familial history of cardiovascular disease and smoking, as well as occupation. We did not include alcohol in our final model because it was not significantly associated with stroke risk in the univariable analyses and because of potential risk of collinearity with other variables in the final model. Survival analyses were not chosen as the primary analyses because the focus of this study was the risk of stroke and not the time to event or when the stroke occurred, as we considered the life-time cumulative exposure. However, Cox regression models were run as sensitivity analysis (Supplemental material). All analyses were performed using R software (version 4.0.4, packages *compareGroups*, *EpiDisplay*, *survival*).

All study participants provided written informed consent prior to enrollment. The study had institutional review board approval for research on human subjects (Commission nationale de l'informatique et des libertés n° 910486, "Comité consultatif sur le traitement de l'information

en matière de recherche n°10.628). The study was approved by the Angers University Hospital ethic committee (approval #2021-186).

RESULTS

Figure 1 illustrate the sample studied from the initial population. From the years 2012 to 2018, 162,114 participants were eligible. After excluding 1,363 participants from the study for having a history of stroke, the final sample size was 160,751 participants, free from stroke at baseline. A description of the population according to stroke subtype is presented in Table 1. There were 20,763 participants (13.4%) who reported exposure to LWH for more than 10 years, and 190 incident cases of stroke (0.12%) comprising 134 ischemic and 56 hemorrhagic strokes. Self-employed/Chief Executive Officer/Professional jobs were the most exposed to LWH (Table 2), with a mean duration of exposure (5.49 years) contrary to Low-skilled white-collar jobs which had the least mean duration of exposure (1.65 years). The median for the time from last year of exposure reported to stroke event was of 7 years (mean 11.4 years), the median follow-up time was 3 years (inter quartile range (IQR) 2-5 years) and the median for time from inclusion to stroke event was of 2 years (IQR 1-3 years).

In crude models, exposure to LWH was associated with an elevated odds of ischemic stroke [OR=1.61(1.04-2.49)] and hemorrhagic stroke [OR=2.50(1.38-4.53)] (Table 3). However, in adjusted multivariable models, statistical significance waned for LWH associations with ischemic stroke [aOR=1.01(0.63-1.63)] and remained significant for hemorrhagic stroke [aOR=1.92(1.01-3.09)]. Age, sex and diagnosed diabetes were significantly associated with ischemic stroke, whereas only age and smoking were significantly associated with hemorrhagic stroke. Associations were similar when running Cox regression models (Table 4). Exposure to LWH \geq 5 years (reference <5 years) was also associated with hemorrhagic stroke [aOR=2.33 (1.27-4.27)] but not to ischemic stroke (aOR of 1.02 [95% CI, 0.66-1.58]).

DISCUSSION

Hemorrhagic stroke was associated with prolonged exposure to LWH after adjusting for cardiovascular risk factors.

These results are similar to findings in other epidemiological studies.(6) Our study further distinguishes ischemic and hemorrhagic stroke in the association with LWH, while accounting for an expansive array of cardiovascular confounders alongside number of years of LWH exposure. Both ischemic and hemorrhagic stroke had significant associations in crude bivariate analyses but our study could find an association between ischemic stroke and LWH in adjusted models. Lack of power may explain this difference as there were relatively few events. Another possible explanation is that the relation between the different risk factors and their influence on the association between LWH and stroke is different for ischemic and hemorrhagic stroke. A predominant indirect pathway may exist for ischemic stroke and thus, the effect of LWH may be mediated by modifiable behaviors. Several studies(14,15) have shown that LWH are associated with negative health behaviors that are also known risk factors for ischemic stroke.(4) Elsewhere, LWH are also described as increasing the risk of cardiac arrhythmias.(16) Stress-related mechanisms, caused by acute and chronic stress, may participate in a pro-inflammatory and hypercoagulability setting that leads to cardiovascular disease.(17,18) The effect of LWH on the risk of hemorrhagic stroke remains after adjusting for cardio-vascular risk factors, suggesting a direct pathway, or factors that possibly precipitate the occurrence of stroke. The effects of LWH on health are complex and several authors have highlighted the need of studies to better understand their effect on several unhealthy behaviors and diseases.(19,20) LWH is associated with stroke with sufficient level of evidence.(2) However, the underlying causal pathway remains unclear as well as the role of associated risks factors which can have a confounding effect or a mediating effect. This study highlights the need for exhaustive causal analyses which differentiates ischemic and hemorrhagic strokes.

The definition of LWH is unusual compared to the weekly working time hours used in many studies but some advantages. First it allows consideration of atypical working hours (long shifts) that may not fit into the usual ≥ 55 hours a week, as well as “rush period” with increase workloads for some jobs that usually have normal working hours. Second, it was possible to estimate life-time cumulative exposure to LWH. However, its main limit is that it assesses LWH as a binary variable and it differs from the usual definition of LWH in weekly hours, making direct comparison with other studies more complex, though generally, associations are similar with other studies.(2,6)

The main limitation of this study is the lack of power, especially in the case of ischemic stroke, which may be caused by the relatively small number of events. Stroke remains a rare event, especially when excluding participants with a history of stroke. However, even after full model adjustments, we found a significant association in the hemorrhagic subgroup, which had fewer events than ischemic stroke subgroup. This study together with others new studies could help clarify the effect of LWH on the risk of stroke in a future meta-analysis, which underlines the importance of individuals studies to share their results, even in the case of lack of power, we found a significant association in the hemorrhagic subgroup, which had fewer events than the ischemic stroke subgroup. In addition, self-reported exposure could be a source of recall bias, modestly influencing results, even though the baseline questionnaire was completed and retrieved independently of the outcome. Moreover, there is a possibility of potential residual confounding, in particular related to working conditions. Lastly, some potential confounder for hemorrhagic stroke could not be considered. For example, estrogen-progestin or blood pressure drugs were not available, as well as blood pressure control which may explain why high blood pressure was not significant in the multivariable analysis, in addition to the lack of power of the study. The large population-based cohort and the extensive adjustment for cardiovascular confounding variables are the main strengths of this study. The design of the study also allowed

for identifying incident cases of stroke and stroke subtypes based on medical records from a national health database.

These analyses highlight the possible differences in the association between prolonged exposure to LWH and subtypes of strokes, all limitations considered. Further studies are needed to better understand the causal pathways and possible mediators and effect measure modifiers. Reducing the global burden of stroke related to LWH, nevertheless, need better controls and interventions on both occupational exposures and lifestyle factors.

COMPETING INTEREST

None. ICMJE disclosure form completed.

FUNDING

No funding for the study (Authors are paid by their institutions). The CONSTANCES Cohort Study was supported and funded by the Caisse nationale d'assurance maladie (CNAM); it is an "Infrastructure Nationale en Biologie et Santé" and benefits from ANR (ANR-11-INBS-0002) grant funding. CONSTANCES is also partly funded by Merck Sharp & Dohme (MSD), AstraZeneca, Lundbeck and L'Oréal through Inserm-Transfert. None of these funding sources had any role in the design of the study, collection and analysis of data or decision to publish.

DATA SHARING

The data of the CONSTANCES cohort are protected by our national regulatory agency ("Commission nationale de l'informatique et des libertés", n°910486). However, the CONSTANCES cohort is "an open epidemiological laboratory" and access to study protocols and data is available on justified request.

ETHICS

All study participants provided written informed consent prior to enrollment. The study had institutional review board approval for research on human subjects (Commission nationale de l'informatique et des libertés n° 910486, “Comité consultatif sur le traitement de l’information en matière de recherche n°10.628). The study was approved by the Angers University Hospital ethic committee (approval #2021-186).

REFERENCES

1. Pega F, Náfrádi B, Momen NC, Ujita Y, Streicher KN, Prüss-Üstün AM, et al. Global, regional, and national burdens of ischemic heart disease and stroke attributable to exposure to long working hours for 194 countries, 2000-2016: A systematic analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ. Int.* . 2021 Sep;154:106595.
2. Descatha A, Sembajwe G, Pega F, Ujita Y, Baer M, Boccuni F, et al. The effect of exposure to long working hours on stroke: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ. Int.* . 2020 Sep;142:105746.
3. Li J, Pega F, Ujita Y, Brisson C, Clays E, Descatha A, et al. The effect of exposure to long working hours on ischaemic heart disease: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ. Int.* . 2020 Sep;142:105739.
4. Arboix A. Cardiovascular risk factors for acute stroke: Risk profiles in the different subtypes of ischemic stroke. *World J. Clin. Cases* . 2015 May 16;3(5):418–29.
5. Morais Filho AB de, Rego TL de H, Mendonça L de L, Almeida SS de, Nóbrega ML da, Palmieri T de O, et al. The physiopathology of spontaneous hemorrhagic stroke: a systematic review. *Rev. Neurosci.* . 2021 Aug 26;32(6):631–58.
6. Hannerz H, Albertsen K, Burr H, Nielsen ML, Garde AH, Larsen AD, et al. Long working hours and stroke among employees in the general workforce of Denmark. *Scand. J. Public Health* . 2018 May;46(3):368–74.
7. Kim BJ, Lee S-H, Ryu W-S, Kim CK, Chung J-W, Kim D, et al. Excessive work and risk of haemorrhagic stroke: a nationwide case-control study. *Int. J. Stroke Off. J. Int. Stroke Soc.* . 2013 Oct;8 Suppl A100:56–61.
8. Fadel M, Sembajwe G, Gagliardi D, Pico F, Li J, Ozguler A, et al. Association Between Reported Long Working Hours and History of Stroke in the CONSTANCES Cohort. *Stroke* . 2019;50(7):1879–82.

9. Fadel M, Li J, Sembajwe G, Gagliardi D, Pico F, Ozguler A, et al. Cumulative Exposure to Long Working Hours and Occurrence of Ischemic Heart Disease: Evidence From the CONSTANCES Cohort at Inception. *J. Am. Heart Assoc.* . 2020 Jun 16;9(12):e015753.
10. Goldberg M, Carton M, Descatha A, Leclerc A, Roquelaure Y, Santin G, et al. CONSTANCES: a general prospective population-based cohort for occupational and environmental epidemiology: cohort profile. *Occup. Environ. Med.* . 2017 Jan;74(1):66–71.
11. Tuppin P, Rudant J, Constantinou P, Gastaldi-Ménager C, Rachas A, de Roquefeuil L, et al. Value of a national administrative database to guide public decisions: From the système national d’information interrégimes de l’Assurance Maladie (SNIIRAM) to the système national des données de santé (SNDS) in France. *Rev. Epidemiol. Sante Publique* . 2017 Oct;65 Suppl 4:S149–67.
12. National Institute of Statistics and Economic Studies (INSEE). Professions et catégories socioprofessionnelles [Internet]. [cited 2022 Dec 15]. Available from: <https://www.insee.fr/fr/metadonnees/pcs2003/categorieSocioprofessionnelleAgregee/1?champRecherche=true>
13. Santé Publique France. Évolution de l’admission en soins de suite et de réadaptation des patients hospitalisés pour accident vasculaire cérébral en France, 2010-2014 [Internet]. [cited 2022 Dec 15]. Available from: <https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-cardiovasculaires-et-accident-vasculaire-cerebral/accident-vasculaire-cerebral/evolution-de-l-admission-en-soins-de-suite-et-de-readaptation-des-patients-hospitalises-pour-accident-vasculaire-cerebral-en-france-2010-2014>
14. Lee D-W, Jang T-W, Kim H-R, Kang M-Y. The relationship between working hours and lifestyle behaviors: Evidence from a population-based panel study in Korea. *J. Occup. Health* . 2021 Jan;63(1):e12280.
15. Virtanen M, Jokela M, Lallukka T, Magnusson Hanson L, Pentti J, Nyberg ST, et al. Long working hours and change in body weight: analysis of individual-participant data from 19 cohort studies. *Int. J. Obes.* 2005 . 2020 Jun;44(6):1368–75.
16. Kivimäki M, Nyberg ST, Batty GD, Kawachi I, Jokela M, Alfredsson L, et al. Long working hours as a risk factor for atrial fibrillation: a multi-cohort study. *Eur. Heart J.* . 2017 Sep 7;38(34):2621–8.
17. Virtanen M, Kivimäki M. Long Working Hours and Risk of Cardiovascular Disease. *Curr. Cardiol. Rep.* . 2018 Oct 1;20(11):123.
18. Osborne MT, Shin LM, Mehta NN, Pitman RK, Fayad ZA, Tawakol A. Disentangling the Links Between Psychosocial Stress and Cardiovascular Disease. *Circ. Cardiovasc. Imaging* . 2020 Aug;13(8):e010931.
19. Pachito DV, Pega F, Bakusic J, Boonen E, Clays E, Descatha A, et al. The effect of exposure to long working hours on alcohol consumption, risky drinking and alcohol use disorder: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ. Int.* . 2021 Jan;146:106205.

20. Rugulies R, Sørensen K, Di Tecco C, Bonafede M, Rondinone BM, Ahn S, et al. The effect of exposure to long working hours on depression: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environ. Int.* . 2021 Oct;155:106629.

Table 1: Description of the population according to stroke subtype

	<i>Total*</i> (<i>N=160751</i>)	<i>Ischemic stroke*</i> (<i>N=134</i>)	<i>Hemorrhagic stroke*</i> (<i>N=56</i>)
Age (Years)	47.7 (13.1)	58.9 (9.43)	59.7 (9.72)
Body mass index (kg/m²)	25.0 (4.48)	26.4 (3.83)	25.4 (3.71)
Sex			
Female	85303 (53.2%)	45 (33.8%)	28 (50.9%)
Male	75112 (46.8%)	88 (66.2%)	27 (49.1%)
High blood pressure diagnosed			
No	142409 (88.7%)	103 (76.9%)	41 (73.2%)
Yes	18152 (11.3%)	31 (23.1%)	15 (26.8%)
Diabetes diagnosed			
No	158466 (98.7%)	123 (91.8%)	55 (98.2%)
Yes	2095 (1.30%)	11 (8.21%)	1 (1.79%)
Familial history of cardiovascular diseases			
No	144529 (90.0%)	117 (87.3%)	50 (89.3%)
Yes	16032 (9.98%)	17 (12.7%)	6 (10.7%)
Dyslipidemia diagnosed			
No	147367 (91.8%)	105 (78.4%)	43 (76.8%)
Yes	13194 (8.22%)	29 (21.6%)	13 (23.2%)
Smoking			
Not smoker	71634 (45.6%)	53 (40.8%)	17 (30.9%)
Current/former smoker <30 pack.years	79434 (50.5%)	61 (46.9%)	27 (49.1%)
Current/former smoker ≥30pack.years	6172 (3.93%)	16 (12.3%)	11 (20.0%)
Occupation			
High-skilled white-collar jobs	43358 (30.0%)	38 (29.0%)	23 (45.1%)
Self-employed/Chief Executive Officer, Professional jobs	49236 (34.0%)	42 (32.1%)	16 (31.4%)
Low-skilled white-collar jobs	37352 (25.8%)	29 (22.1%)	8 (15.7%)
Blue-collar jobs	14723 (10.2%)	22 (16.8%)	4 (7.84%)
Long working hours exposure			
<10 years	134429 (86.6%)	101 (80.2%)	39 (72.2%)
≥10 years	20723 (13.4%)	25 (19.8%)	15 (27.8%)

*Quantitative variables are shown in mean (standard deviation) and categorical variables in number (percentage)

Table 2: Exposure to long working hours according to occupation

	<i>High-skilled white-collar jobs N=42399</i>	<i>Self-employed/Chief Executive Officer, Professional jobs N=46426</i>	<i>Low-skilled white-collar jobs N=36737</i>	<i>Blue-collar jobs N=14390</i>	<i>p.overall</i>
Long working hours (years)	2.89 (7.22)	5.49 (9.34)	1.65 (5.11)	2.91 (7.39)	<0.001
Long working hours					<0.001
<10 years	37583 (88.6%)	36142 (77.8%)	34401 (93.6%)	12767 (88.7%)	
≥10 years	4816 (11.4%)	10284 (22.2%)	2336 (6.36%)	1623 (11.3%)	

Table 3: Crude and adjusted multinomial logistic regression assessing association between LWH and stroke.

	<i>Total*</i>	<i>Stroke cases Number (%)</i>	<i>Ischemic stroke</i>		<i>Hemorrhagic stroke</i>	
			Crude Odds Ratios (95% confidence interval)	Adjusted Odds Ratios† (95% confidence interval)	Crude Odds Ratios (95% confidence interval)	Adjusted Odds Ratios† (95% confidence interval)
Age (Years)	47.7 (13.1)		1.08 (1.07-1.10)	1.08 (1.06-1.10)	1.09 (1.06-1.12)	1.08 (1.05-1.12)
Body mass index (kg/m²)	25.0 (4.48)		1.06 (1.03-1.09)	1.01 (0.96-1.05)	1.02 (0.96-1.08)	0.94 (0.88-1.02)
Sex						
<i>Female</i>	85376 (53.2%)	73 (38.8%)	ref		ref	
<i>Male</i>	75227 (46.8%)	115 (61.2%)	2.22 (1.55-3.18)	1.92 (1.26-2.91)	1.10 (0.65-1.86)	0.84 (0.45-1.56)
High blood pressure diagnosed						
<i>No</i>	142553 (88.7%)	144 (75.8%)	ref		ref	
<i>Yes</i>	18198 (11.3%)	46 (24.2%)	2.36 (1.58-3.53)	0.87 (0.54-1.41)	2.87 (1.59-5.19)	1.46 (0.73-2.91)
Diabetes diagnosed						
<i>No</i>	158644 (98.7%)	178 (93.7%)	ref		ref	
<i>Yes</i>	2107 (1.31%)	12 (6.32%)	6.76 (3.65-12.55)	3.86 (1.95-7.65)	1.38 (0.19-9.94)	0.79 (0.11-5.84)
Familial history of cardiovascular diseases						
<i>No</i>	144696 (90.0%)	167 (87.9%)	ref		ref	
<i>Yes</i>	16055 (9.99%)	23 (12.1%)	1.31 (0.79-2.18)	1.10 (0.65-1.86)	1.08 (0.46-2.52)	0.89 (0.38-2.10)
Dyslipidemia diagnosed						
<i>No</i>	147715 (91.8%)	148 (77.9%)	ref		ref	
<i>Yes</i>	13236 (8.23%)	42 (22.1%)	3.08 (2.04-4.66)	1.17 (0.72-1.88)	3.38 (1.82-6.28)	1.70 (0.84-3.41)
Smoking						
<i>Not smoker</i>	71704 (45.5%)	70 (37.8%)	ref		ref	
<i>Current/former smoker <30 pack.years</i>	79522 (50.5%)	88 (47.6%)	1.04 (0.72-1.50)	0.94 (0.64-1.39)	1.43 (0.78-2.63)	1.59 (0.82-3.08)
<i>Current/former smoker ≥30pack.years</i>	6199 (3.94%)	27 (14.6%)	3.50 (2.00-6.13)	1.70 (0.94-3.06)	7.51 (3.52-16.04)	4.58 (1.06-10.68)

Occupation						
<i>High-skilled white-collar jobs</i>	43419 (30.0%)	61 (33.5%)	ref		ref	
<i>Self-employed/Chief Executive Officer, Professional jobs</i>	49294 (34.0%)	58 (31.9%)	0.97 (0.63-1.51)	0.94 (0.59-1.51)	0.61 (0.32-1.16)	0.54 (0.28-1.06)
<i>Low-skilled white-collar jobs</i>	37389 (25.8%)	37 (20.3%)	0.89 (0.55-1.44)	1.29 (0.77-2.15)	0.40 (0.18-0.90)	0.47 (0.20-1.11)
<i>Blue-collar jobs</i>	14749 (10.2%)	26 (14.3%)	1.71 (1.01-2.88)	1.22 (0.68-2.18)	0.51 (0.18-1.48)	0.53 (0.18-1.57)
Long working hours exposure						
<i><10 years</i>	134569 (86.6%)	140 (77.8%)	ref		ref	
<i>≥10 years</i>	20763 (13.4%)	40 (22.2%)	1.61 (1.04-2.49)	1.01 (0.63-1.63)	2.50 (1.38-4.53)	1.92 (1.01-3.69)

*Quantitative variables are shown in mean (standard deviation) and categorical variables in number (percentage)

†Multivariable models are adjusted for age, body mass index, high blood pressure, diabetes, dyslipidemia, smoking and occupation.

Table 4: Crude and adjusted Cox regression models assessing association between LWH and stroke.

		<i>Ischemic stroke</i>		<i>Hemorrhagic stroke</i>	
		Crude Odds Ratios (95% confidence interval)	Adjusted Odds Ratios [†] (95% confidence interval)	Crude Odds Ratios (95% confidence interval)	Adjusted Odds Ratios [†] (95% confidence interval)
Age (Years)		1.08 (1.06-1.10)	1.08 (1.06-1.10)	1.10 (1.06-1.12)	1.08 (1.05-1.12)
Body mass index (kg/m²)		1.06 (1.03-1.09)	1.01 (0.97-1.05)	1.02 (0.96-1.08)	0.95 (0.88-1.02)
Sex					
	<i>Female</i>	ref		ref	
	<i>Male</i>	2.17 (1.51-3.11)	1.91 (1.25-2.90)	1.13 (0.66-1.93)	0.83 (0.44-1.55)
High blood pressure diagnosed					
	<i>No</i>	ref		ref	
	<i>Yes</i>	2.34 (1.56-3.50)	0.87 (0.54-1.41)	2.66 (1.44-4.88)	1.47 (0.74-2.93)
Diabetes diagnosed					
	<i>No</i>	ref		ref	
	<i>Yes</i>	5.09 (2.74-9.46)	2.97 (1.50-5.89)	1.10 (0.15-8.02)	0.64 (0.09-4.72)
Familial history of cardiovascular diseases					
	<i>No</i>	ref		ref	
	<i>Yes</i>	1.32 (0.79-2.20)	1.09 (0.65-1.86)	1.11 (0.47-2.59)	0.89 (0.38-2.11)
Dyslipidemia diagnosed					
	<i>No</i>	ref		ref	
	<i>Yes</i>	2.94 (1.95-4.45)	1.13 (0.65-1.86)	3.33 (1.78-6.21)	1.66 (0.83-3.33)
Smoking					
	<i>Not smoker</i>	ref		ref	
	<i>Current/former smoker <30 pack.years</i>	1.07 (0.74-1.56)	0.94 (0.63-1.39)	1.37 (0.74-2.52)	1.59 (0.82-3.08)
	<i>Current/former smoker ≥30pack.years</i>	3.45 (1.97-6.04)	1.68 (0.93-3.02)	7.18 (3.36-15.33)	4.54 (1.95-10.59)
Occupation					
	<i>High-skilled white-collar jobs</i>	ref		ref	
	<i>Self-employed/Chief Executive Officer, Professional jobs</i>	0.98 (0.63-1.52)	0.94 (0.59-1.50)	0.62 (0.33-1.17)	0.54 (0.28-1.05)
	<i>Low-skilled white-collar jobs</i>	0.89 (0.55-1.44)	1.29 (0.78-2.15)	0.40 (0.18-0.90)	0.47 (0.20-1.11)
	<i>Blue-collar jobs</i>	1.74 (1.03-2.94)	1.27 (0.71-2.27)	0.52 (0.18-1.50)	0.55 (0.19-1.63)
Long working hours exposure					
	<i><10 years</i>	ref		ref	
	<i>≥10 years</i>	1.61 (1.04-2.50)	1.00 (0.63-1.61)	2.57 (1.41-4.70)	1.91 (1.00-3.66)

†Multivariable models are adjusted for age, body mass index, high blood pressure, diabetes, dyslipidemia, smoking and occupation.