



HAL
open science

Influence of the healthcare pathway on the outcome of patients with infective endocarditis

Florent Arregle, Nicolas Iline, Roch Giorgi, Mary Philip, Sandrine Hubert, Frederique Gouriet, Jean Paul Casalta, Frédéric Collart, Alberto Riberi, Hélène Martel, et al.

► To cite this version:

Florent Arregle, Nicolas Iline, Roch Giorgi, Mary Philip, Sandrine Hubert, et al.. Influence of the healthcare pathway on the outcome of patients with infective endocarditis. *European Heart Journal: Acute Cardiovascular Care*, 2022, 11 (9), pp.672-681. 10.1093/ehjacc/zuac088 . hal-04043110

HAL Id: hal-04043110

<https://hal.science/hal-04043110>

Submitted on 17 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Abstract**

2 **Aims:** To determine the prognosis of patients treated for infective endocarditis (IE) according
3 to their health-care pathway. To assess how the ESC guidelines are implemented concerning
4 the performance of transesophageal echocardiography (TEE), the use of antibiotic therapy, and
5 the performance of valve surgery; and to compare the epidemiological profile of IE according
6 to the type of centers in which the patients are hospitalized.

7 **Methods and results:** In a prospective multicentric study including 22 hospitals in the South-
8 East of France, 342 patients were classified in three groups according to their health-care
9 pathway: 119 patients diagnosed and taken care entirely in a reference center or hospital with
10 cardiac surgery (Referral Center (RC) group), 111 patients diagnosed and initially taken care
11 in a non-referral center (NRC) , then referred in a center including cardiac surgery (Transferred
12 to Referral Center (TRC) group) and 112 patients totally taken care in NRC (NRC group). One-
13 year mortality was 26% (88 deaths) and was not significantly different between groups 1 and 2
14 (20% vs 21%, $p = 0.83$). Patients in NRC group had a higher mortality (37%) compared to
15 patients in RC and TRC groups ($p < 0.001$). ESC guidelines were not implemented similarly
16 depending on the health-care pathway ($p = 0.04$). Patients in NRC group were significantly older
17 ($p < 0.001$) and had more comorbidities ($p < 0.001$) than patients treated in referral centers.

18 **Conclusion:** Prognosis of patients with IE is influenced by their health-care pathway. Patients
19 treated exclusively in NRC have a worse prognosis than patients treated in referral or surgical
20 centers.

21

22 **Keywords :** endocarditis, mortality, health-care pathway, referral center

23

24

25

1 **Introduction**

2 Epidemiologic profile of infective endocarditis (IE) has changed over the past twenty
3 years,^{1 2 3} but mortality remains high^{4 5 6} despite diagnostic improvement and access to surgery
4 in up to 50% of cases.⁷ This persistent severity has justified the development of new diagnostic
5 and therapeutic strategies to improve prognosis.^{8 9} Management by a specialized
6 multidisciplinary team within a referent medical-surgical center seems to be a key factor in
7 improving patients prognosis,^{10 11} and is recommended by the ESC guidelines.^{8 9}

8 However, few studies have compared the prognosis and management of patients with
9 IE according to their health-care pathway. Indeed, three situations are usually observed: patients
10 for whom the diagnosis and management are carried out entirely in a referral center and/or
11 having a cardiac surgery department, patients secondarily referred to a referral hospital or
12 surgical center by another care center, patients for whom the entire care is provided within a
13 non-referral care center. In addition, application of European guidelines remains difficult to
14 assess in daily clinical practice outside the major referral centers. Published epidemiological
15 data only represent part of the IE, most often not taking into account patients hospitalized in
16 non-referral centers and therefore introducing a significant bias in the knowledge of this
17 pathology.¹² Finally, although the ESC guidelines suggest that patients with uncomplicated IE
18 can be managed in a non-reference center⁹, if all patients with IE¹³ or only the sickest patients
19¹⁴ should be referred to reference centers is still a matter of debate.

20 The main objective of this study was to determine the prognosis of patients treated for
21 IE according to their health-care pathway. The secondary objectives were to assess how the
22 ESC guidelines are implemented concerning the performance of transesophageal
23 echocardiography (TEE), the use of antibiotic therapy, and the performance of valve surgery,
24 and to compare the epidemiological profile of IE according to the type of centers in which the
25 patients are hospitalized.

1 **Methods**

2 **Study design and patients**

3 This was a prospective multicentric study including patients consecutively admitted for
4 suspicion of IE in 22 hospitals in South-East of France between January 2014 and June 2017.
5 These hospitals were both tertiary centers with cardiac surgery and second-level community
6 hospitals. Inclusion criteria were a diagnosis of definite IE based on Duke modified criteria ¹⁵
7 on native, prosthetic or pacemaker/implantable automatic defibrillator. Non-inclusion criteria
8 were age < 18 years and patients already included in the study and presenting with a new IE
9 diagnosis. Patients were screened in the echocardiography lab of each institution and were
10 prospectively included by local investigators.

11

12 **Patient and Public involvement**

13 All patients hospitalized for suspected IE were accepted on admission to participate in this
14 research protocol. However, they were not involved in the design, conduct, reporting or
15 dissemination plans of our research. The study complies with the Declaration of Helsinki.
16 Written consent was waived by La Timone Institutional Review Board, which gave its approval
17 for this study.

18

19 **Data collected at the time of inclusion and during hospitalization**

20 The following clinical data were collected: age, sex, previous cardiac and non-cardiac
21 diseases, Charlson index score, date of first clinical signs, body temperature, presence of cardiac
22 murmur, congestive heart failure, neurological complications, septic or cariogenic shock or
23 atrio-ventricular block on electrocardiogram at time of admission and during hospitalization.

1 Biological data were collected, including hemoglobin level, platelet count, leukocytes count,
2 C-Reactive Protein serum level, creatinine level, rheumatoid factor. Microbiological data
3 included: positive or negative blood cultures, type of microorganism and antibiogram.

4 Echocardiographic data collected were: presence of vegetation and its maximal length,
5 presence of periannular lesion (defined as the presence of abscess and/or pseudoaneurysm
6 and/or fistula), presence of valvular regurgitation, quantified according to current
7 recommendations.^{16 17}

8 Type and date of the following complications were collected: congestive heart failure,
9 peripheral embolism, neurological complications (including symptomatic and silent), mycotic
10 aneurysm, acute kidney failure (>26 μ mol/L increase of serum creatinine in 48 hours or >50%
11 increase in 7 days) , glomerulonephritis, uncontrolled infection (fever and/or positive blood
12 cultures after 7 days of appropriate antibiotic therapy and unlinked to an extracardiac cause).

13 Moreover, type, dose and length of antibiotic therapy was reported. Antibiotic therapy
14 was considered appropriate if drugs used and full duration of treatment were in accordance with
15 ESC guidelines (taking into account patients terminating antibiotics prematurely due to death).
16 Analysis of antibiotic treatment was conducted retrospectively by the Endocarditis team of the
17 investigator center (CHU La Timone, Marseille).

18 During hospitalization or follow-up the indication (according to ESC guidelines) and
19 date of surgery was also collected. Finally, surgical risk was evaluated using the Euroscore II.¹⁸

20

21 **Classification of patients according to health-care pathway**

22 Patients were classified in three groups according to their health-care pathway:

- 23 - Referral Center group (RC), including patients diagnosed and taken care entirely in a
24 reference center or hospital with cardiac surgery (4 hospitals). These centers were the
25 only surgical centers of this region of France.

- 1 - Transferred to Referral Center group (TRC), including patients diagnosed and initially
2 taken care in a non-referral center, then referred in a center including cardiac surgery
3 - Non-Referral Center group (NRC), including patients totally taken care in non-referral
4 centers (18 centers)

5

6 **Follow-up**

7 Patients had a one-year follow-up with a clinical evaluation or if not possible, phone contact.

8

9 **Primary and secondary endpoints**

10 Primary endpoint was death from all cause at one-year follow-up. Secondary endpoints
11 were:

12 - the evaluation of the implementation of European guidelines^{8 9} in the three patients groups
13 regarding: the use of TEE, the choice and duration of antibiotic therapy, and the performance
14 of valvular surgery and/or extraction of pacemaker or implantable automatic defibrillator when
15 indicated

16 - comparison of demographical and microbiological data among the three groups.

17

18 **Statistical analysis**

19 Continuous variables were described by their mean, standard deviation, minimum and
20 maximum value, 1st and 3rd quartile. Continuous variables were compared using the Mann-
21 Whitney non parametric U test or Student's t test, depending on the application conditions.
22 Categorical variables were compared using the chi-square test or the Fisher test, depending on
23 the application conditions. The event-free survival distributions were estimated with the
24 Kaplan-Meier method. The log-rank test was used to compare survival distributions for
25 categorical variables. The effect of continuous variables on the risk of an event occurring was

1 estimated and tested using a Cox model. The surgery variable, which is a time dependent
2 variable, was estimated using a Poisson approach.

3 Multivariate analyzes were tested using a Cox model. Multivariate analyzes were tested
4 using a Cox model. Inclusion of variables in the fitted model was carried out according to the
5 following criteria : result in univariate analysis with statistical significance $p < 0.20$, number of
6 missing values $< 10\%$ for this variable, effective presence in both categories of a bivariate
7 variable $> 10\%$, as well as according to their clinical relevance. The final models were selected
8 using the Akaike Information Criteria (AIC).

9 The comparison tests of group 1 to group 2, group 1 to group 3, group 1 + 2 to group 3
10 were performed in bilateral situation using Bonferroni correction and were considered
11 statistically significant for $p < 0.017$. For all other tests, they were performed bilaterally and
12 were considered statistically significant for $p < 0.05$. Statistical analysis was performed with R
13 software (version 3.6.2) and RStudio (v 1.2.5).

14

15

16

17 **Results**

18 **Patient population**

19 Between January 2014 and June 2017, a total of 342 patients with definite IE were
20 included. Among them, 119 were diagnosed and entirely managed in a reference center or
21 hospital with cardiac surgery (RC group), 111 were diagnosed and initially managed in a non-
22 referral center, then referred in a center including cardiac surgery (TRC group) and 112 were
23 totally managed in non-referral centers (NRC group). Clinical, demographical, biological, and
24 microbiological characteristics of patients are presented in **Table 1**.

25

1 *Overall population*

2 In the overall population, mean age was 68 +/- 15 years. IE were mainly community-
3 acquired (83% of cases), *S. aureus* was the most frequent bacteria involved (26% of cases).
4 Negative blood culture IE represented 17% of cases. Left-sided IE counted for 85% of all cases
5 and aortic (43%) and mitral (42%) localization were almost equally represented. Perivalvular
6 complications were observed in 16% of cases. 33% of cases were prosthetic valve IE. More
7 than half IE cases (58%) showed theoretical indication for valvular surgery or intracardiac
8 device removal but only 35% had such interventions.

9

10 *Characteristics according to health-care pathway (Table 1)*

11 Patients in NRC group were significantly older and had higher Charlson index. Sources
12 of infection did not differ between the three groups and microbiological profile of IE cases was
13 not significantly different. Echocardiographic findings showed more severe lesions in RC group
14 and TRC group: longer vegetations, more peri-valvular complications, and more severe
15 valvular regurgitation. Concerning IE complications, there was no significant difference
16 between groups in embolic events, however more silent embolisms were diagnosed in RC group
17 and TRC group. Spondylodiscitis was more frequent in NRC group.

18 Patients in RC and TRC groups had more frequent surgical indication, but surgery or
19 intracardiac device removal was performed in only 52 patients (44%) in RC group, 60
20 patients (55%) in TRC group and 6 patients (5%) in NRC group, $p < 0.001$. In RC group, 81
21 patients had an indication for intervention and 29 were not operated and in TRC group, 91
22 patients had an indication for intervention and 31 were not operated. Reasons for absence of
23 intervention were: deaths before surgery (6 patients in RC group and 2 patients in TRC group
24), patient's refusal (2 patients in RC group and 1 patient in TRC group) and decision of the
25 endocarditis team due to comorbidities (21 patients in RC group and 28 patients in TRC

1 group). Patients who benefited from surgery were significantly younger (69.4 +/- 13 years vs
2 62.4 +/- 12 years, $p=0.01$), had lower Charlson index (3.9 +/- 1.9 vs 2.8 +/- 2.7, $p=0.007$),
3 had more frequently vegetations (86 patients (73%) vs 95 patients (85%), $p=0.002$) and severe
4 valvular regurgitation (41 patients (35%) vs 68 patients (61%), $p<0.001$) but less renal failure
5 (29 patients (25%) vs 5 patients (4%), $p<0.001$) and diabetes (39 patients (33%) vs 20 patients
6 (18%), $p=0.01$).

7 Time from diagnosis to intervention did not significantly differ between RC group and
8 TRC group (mean time 30+/-32 days vs 22+/-29 days, $p=0.36$). In our cohort, mean time for
9 right-sided IE was 17+/-18 days and 25+/-28 days for left-sided IE.

10

11 **Prognosis of patients according to their health-care pathway**

12 Among the 342 patients, one year mortality was 26% (88 deaths), hospital mortality was
13 13% (44 deaths) and one year recurrence rate was 4% (13 patients). **Figure 1** shows survival
14 probability according to health-care pathway.

15 Hospital mortality was 11% (13 deaths) in RC group, 8% (9 patients) in TRC group and
16 20% (22 patients) in NRC group, $p=0.03$. One year mortality was 20% (24 deaths) in RC group,
17 21% (23 deaths) in TRC group and 37% (41 deaths) in NRC group, $p=0.003$. One year mortality
18 was not significantly different between RC and TRC groups ($p=0.83$), however it was
19 significantly higher in NRC group compared to RC and TRC groups ($p<0.001$).

20

21 **Characteristics of patients only treated in non-referral centers**

22 During the study period, 112 patients were managed only in a NRC. Among them, 28
23 patients (25%) had indication for valvular surgery or intracardiac device removal. These
24 patients had a mean age of 76+/-11 years and were at high surgical risk (mean Euroscore 17+/-
25 16%). Reasons for absence of transfer and surgery included: death before surgery for 16 patients

1 (14%), medical choice for 14 patients (13%), patient's refusal in 3 cases (3%) and high surgical
2 risk (comorbidities) for 16 patients (14%). 6 patients (5%) had intracardiac device removal
3 (pacemaker lead or catheter) in the NRC. Among these 28 patients, one year mortality was 50%
4 (14 deaths). 84 patients (75%) had indication for medical treatment of IE. One year mortality
5 among them was 29% (24 deaths).

6

7 **Prognostic factors of mortality**

8 Results of univariate analysis of prognostic factors of one year mortality are shown in
9 **Table S1. Table 2** shows prognostic factors of mortality among the 342 patients in multivariate
10 analysis. Belonging to NRC group was associated with higher risk of death (HR 2.56; 95% CI
11 1.44-4.55, $p=0.001$). Heart failure (HR 2.4; 95% CI 1.52-3.78, $p<0.001$) and peri-valvular
12 involvement (HR 3.25; 95% CI 1.86-5.68, $p<0.001$) were IE complications predictive of
13 mortality.

14

15 **Evaluation of European guidelines application according to health-care pathway**

16 **Table 3** shows the implementation of ESC guidelines in the three groups regarding use
17 of TEE, valvular surgery or intracardiac device removal and antibiotic therapy. In overall
18 population, ESC guidelines were correctly implemented in 52% of patients. TEE was
19 performed in most IE cases (92%); however it was significantly less frequently used in NRC
20 group as compared to the two other groups ($p<0.001$). Guidelines concerning surgery or
21 intracardiac device extraction were correctly implemented in 78% of cases and there was no
22 significant difference between the three groups. Antibiotic therapy was in accordance with
23 guidelines in 72% of cases and there was no significant difference between patients taken care
24 entirely or partially in RC ($p=0.66$). However, treatment was less frequently in agreement with
25 guidelines in patients in NRC group compared to the others ($p<0.001$).

1 **Discussion**

2 This prospective study carried out in 22 hospitals in the South-East of France including
3 342 IE cases divided into three groups according to their health-care pathway shows that:

4 - The prognosis of patients is influenced by their health-care pathway

5 - The number of patients managed in non-referral centers is high. They are older, have more
6 comorbidities, and less severe cardiac lesions than the others.

7 - Surgery or device removal is less frequently performed in those patients

8 - European recommendations on the management of these patients are not followed similarly
9 depending on their health-care pathway

10 - Both in-hospital and long-term mortalities are higher in non-referral centers.

11

12 **General population**

13 By including patients treated in NRC, our study has the advantage of providing global
14 epidemiological information concerning the cases of IE within the south-eastern region,
15 reflecting the management of this pathology in "real life" and not only in RC. Almost a third of
16 the patients treated for IE during the inclusion period were treated in a NRC.

17 Consistent with known data ^{19 20}, there was a predominance of IE cases in males (69%).

18 The mean age of diagnosis of IE was 68 years in our study. This age seemed more advanced in
19 comparison to other French (average age of 59 +/- 17 years in 1999 ², 62 +/- 16 years in 2008
20 ²¹) or European studies (57 +/- 16 years in 2001 ³, 59 +/- 18 years in the Euro-Endo registry ²⁰).

21 Concerning microbiological data, *S. aureus* was the predominant germ (26% of IE), which was
22 in agreement with previous studies ^{21 22}. There was a majority of left-sided IE (85%). An
23 embolic event was observed in 38% of cases. This rate was comparable to other studies that
24 describe the occurrence of an embolic event in 20 to 50% of cases ^{21 23 24}

25

1 **Mortality according to health-care pathway**

2 In our cohort, hospital mortality was 13% and one-year mortality was 26%. Few studies
3 have compared the prognosis of patients with IE according to their health-care pathway. A
4 study by Fernandez-Hidalgo et al. ²⁵, compared the hospital mortality of 223 patients treated in
5 a RC with cardiac surgery with 144 patients taken care in a NRC and secondarily referred to
6 the RC. This study did not show any significant difference in terms of hospital mortality
7 between the two groups. Likewise, an international multicenter study ¹² carried out in RC in the
8 management of IE did not show difference in hospital mortality between 1,164 patients
9 transferred from NRC and 1,596 patients treated directly within the RC. However, these studies
10 have some limitations: by considering only patients treated in RC they are not representative of
11 epidemiology of patients treated in NRC. The originality of our study is to compare three
12 groups, including a group of patients treated exclusively in NRC, these patients usually being
13 not included in previous studies. Our main findings is a significantly higher hospital (20%) and
14 one year mortality (37%) in patients treated only in NRC. Belonging to NRC group was an
15 independent risk factor in multivariate analysis of one-year mortality in the total cohort of
16 patients. This excess of mortality observed in NRC group could be explained by various factors:
17 epidemiological differences compared to other groups, recommendations on the management
18 of IE that are less implemented and a higher absence of surgery rate in patients with a theoretical
19 surgical indication.

20

21 **Implementation of European guidelines**

22 In our study, we observed a more frequent use of TEE (92% of patients) than in other
23 multicenter studies (60-75% ^{3 20}). However, within NRC group, TEE was significantly less
24 frequently performed compared to the other groups. TEE has a key role in the diagnostic
25 process of IE and in particular in the detection of peri-annular complications and should be

1 performed systematically, in accordance with current recommendations ^{8 9 26}. Thus, it is
2 possible that these complications, often requiring surgery, have not been diagnosed in some
3 patients, especially in NRC group.

4 Regarding valve surgery or the extraction of intracardiac material, there was no
5 significant difference in application of recommendations between the three groups. However,
6 we noted within each group patients with a theoretical indication for surgery for who surgery
7 was finally not performed. Within NRC group, 25% of patients presented a theoretical
8 indication and only 5% received interventional management. Conversely, in RC and TRC
9 groups, 65% of patients with theoretical surgical indication benefited from intervention.
10 Different studies have already shown the poor prognosis of patients with a theoretical indication
11 for surgery and not operated ^{20 27}, which may in part explain the higher mortality of patients in
12 NRC group. In addition, the main multicenter studies carried out over the last few years
13 highlight a surgery rate close to 50% (52% in Euro Heart Survey ³, 51% in the EuroEndo
14 register ²⁰, 48% in the ICE register ⁴). Among the patients treated exclusively in RC, the rate of
15 surgery or extraction of intracardiac material was 44%, close to previous published data. On
16 the other hand, if we considered all the patients for whom the diagnosis of IE was made in a
17 non-referral center (TRC and NRC groups), the overall rate of valve surgery or extraction of
18 intracardiac material was only 30 %. Several factors could explain this discrepancy with the
19 pre-existing studies. Indeed, most studies on IE focused on patients treated in RC with a cardiac
20 surgery department. It is possible that a referral bias may cause an overestimation of the rate of
21 cardiac surgery in this pathology, by not considering many IE of "medical" management and
22 treated exclusively in NRC. On the other hand, it is also possible that among the patients
23 diagnosed in a NRC, surgical indications were sometimes not identified, especially in NRC
24 group in which, unlike TRC group, it was not possible to re-evaluate the surgical indications
25 by a multidisciplinary team in a reference center. In a same way, a Spanish study by Lopez-

1 Dupla et al. ²⁸ in a university center without a cardiac surgery department showed that the
2 establishment of a multidisciplinary team including a cardiac surgeon was associated with an
3 increase in the rate of surgical indication identified (14.5% vs 34.5%). Finally, the European
4 guidelines concerning the antibiotic treatment were significantly less applied among patients in
5 group 3 with 35% of inappropriate antibiotic therapy. Here again, few studies concerning the
6 antibiotic treatment used in non-referral centers are available. Fernandez-Hidalgo et al. ²⁵
7 showed that among the patients diagnosed in a NRC and referred secondarily to a surgical
8 center, the antibiotic treatment initiated in the non-referral center was inadequate in 54% of
9 cases. In this study, inadequate treatment was associated with an increased risk of in-hospital
10 mortality in multivariate analysis. The goal of our study was to evaluate the implementation of
11 ESC guidelines in “real life”. It is however important to highlight that in some cases, there may
12 be logical reasons why patients do not undergo an indicated procedure such as TEE, for instance
13 when a patient is not operable, or surgery.

14

15 **Epidemiological characteristics of IE according to health-care pathway**

16 The epidemiological profile of patients in RC and TRC groups was comparable. These
17 were patients with more severe IE than in NRC group with more hemodynamic complications,
18 a higher rate of vegetations, more peri-annular complications. These elements partly explain
19 the higher surgical indication rate in these two groups. The patients in NRC group were
20 significantly older, presented more comorbidities, and a greater operative risk. Some studies
21 focusing on IE cases in the elderly have shown, including during treatment in a RC, limited
22 access to surgery and significant mortality in these patients when surgery, although indicated,
23 was not performed ^{29 30}.

24

25

1 **Limitations**

2 Some data could not be recorded, such as the reason for the transfer of patients to a
3 referral center, the time between diagnosis and the transfer of patients, and some factors who
4 could have influenced transfer decision (such as cirrhosis or poor nutritional status). Due to the
5 large number of participating non-referral centers, the reasons for transferring patients may
6 have differed from one center to another. Moreover, modified Duke criteria were used for
7 diagnosis of IE and have limited sensitivity. This study reflects the epidemiology and practices
8 used in the south-eastern region of France and cannot be generalized to all the other centers.
9 Finally, the study design was meant to compare prognosis of patients according to the health-
10 care pathway but reasons explaining the different prognosis observed could only be
11 hypothesized.

12

13 **Conclusion**

14 Prognosis of patients with IE is influenced by their health-care pathway. Patients treated
15 exclusively in NRC have a worse prognosis than patients treated in RC. These patients are often
16 fragile and their management difficult. Given the difference in mortality between patients
17 managed in referral vs non-referral centers, the current recommendations of treating patients
18 with non-complicated IE in non-referral centers should be questioned. IE is a deadly disease
19 with a better prognosis when managed in RC.

20

21 **Supplementary data**

22 Supplementary material is available online.

23

24 **Funding**

1 This study was supported by a grant from: the PHRCI (Projet Hospitalier de Recherche
2 Clinique Inter-régional 2012)

3

4 **Conflict of interest**

5 None declared

6

7 **Data availability**

8 The data underlying this article will be shared on reasonable request to the corresponding
9 author.

10

11

12

13

14 **References**

15 1. Thuny F, Grisoli D, Collart F, Habib G, Raoult D. Management of infective endocarditis:
16 challenges and perspectives. *The Lancet* 2012;379:965–975.

17 2. Hoen B. Changing Profile of Infective Endocarditis Results of a 1-Year Survey in France.
18 *JAMA* 2002;288:75.

19 3. Tornos P. Infective endocarditis in Europe: lessons from the Euro heart survey. *Heart*
20 2005;91:571–575.

21 4. Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, Karchmer AW,
22 Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones
23 P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell
24 CH, International Collaboration on Endocarditis-Prospective Cohort Study (ICE-PCS)

- 1 Investigators. Clinical presentation, etiology, and outcome of infective endocarditis in the
2 21st century: the International Collaboration on Endocarditis-Prospective Cohort Study.
3 *Arch Intern Med* 2009;169:463–473.
- 4 5. Gálvez-Acebal J, Rodríguez-Baño J, Martínez-Marcos FJ, Reguera JM, Plata A, Ruiz J,
5 Marquez M, Lomas JM, Torre-Lima J de la, Hidalgo-Tenorio C, Alarcón A de, Grupo
6 para el Estudio de las Infecciones Cardiovasculares de la Sociedad Andaluza de
7 Enfermedades Infecciosas (SAEI). Prognostic factors in left-sided endocarditis: results
8 from the Andalusian multicenter cohort. *BMC Infect Dis* 2010;10:17.
- 9 6. Thuny F, Disalvo G, Belliard O, Avierinos J-F, Pergola V, Rosenberg V, Casalta J-P,
10 Gouvernet J, Derumeaux G, Iarussi D, Ambrosi P, Calabro R, Riberi A, Collart F, Metras
11 D, Lepidi H, Raoult D, Harle J-R, Weiller P-J, Cohen A, Habib G. Risk of Embolism and
12 Death in Infective Endocarditis: Prognostic Value of Echocardiography: A Prospective
13 Multicenter Study. *Circulation* 2005;112:69–75.
- 14 7. Thuny F, Beurtheret S, Mancini J, Gariboldi V, Casalta J-P, Riberi A, Giorgi R, Gouriet
15 F, Tafanelli L, Avierinos J-F, Renard S, Collart F, Raoult D, Habib G. The timing of
16 surgery influences mortality and morbidity in adults with severe complicated infective
17 endocarditis: a propensity analysis. *Eur Heart J* 2011;32:2027–2033.
- 18 8. Endorsed by the European Society of Clinical Microbiology and Infectious Diseases
19 (ESCMID) and by the International Society of Chemotherapy (ISC) for Infection and
20 Cancer, Authors/Task Force Members, Habib G, Hoen B, Tornos P, Thuny F, Prendergast
21 B, Vilacosta I, Moreillon P, Jesus Antunes M de, Thilen U, Lekakis J, Lengyel M, Muller
22 L, Naber CK, Nihoyannopoulos P, Moritz A, Zamorano JL, ESC Committee for Practice
23 Guidelines (CPG), Vahanian A, Auricchio A, Bax J, Ceconi C, Dean V, Filippatos G,
24 Funck-Brentano C, Hobbs R, Kearney P, McDonagh T, McGregor K, et al. Guidelines on

- 1 the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): The
2 Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the
3 European Society of Cardiology (ESC). *Eur Heart J* 2009;30:2369–2413.
- 4 9. Habib G, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta J-P, Del Zotti F, Dulgheru
5 R, El Khoury G, Erba PA, Iung B, Miro JM, Mulder BJ, Plonska-Gosciniak E, Price S,
6 Roos-Hesselink J, Snygg-Martin U, Thuny F, Tornos Mas P, Vilacosta I, Zamorano JL.
7 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for
8 the Management of Infective Endocarditis of the European Society of Cardiology
9 (ESC) Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the
10 European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015;36:3075–3128.
- 11 10. Botelho-Nevers E, Thuny F, Casalta JP, Richet H, Gouriet F, Collart F, Riberi A, Habib
12 G, Raoult D. Dramatic Reduction in Infective Endocarditis–Related Mortality With a
13 Management-Based Approach. *Arch Intern Med* 2009;169:1290.
- 14 11. Chirillo F, Scotton P, Rocco F, Rigoli R, Borsatto F, Pedrocco A, De Leo A, Minniti G,
15 Polesel E, Olivari Z. Impact of a Multidisciplinary Management Strategy on the Outcome
16 of Patients With Native Valve Infective Endocarditis. *Am J Cardiol* 2013;112:1171–1176.
- 17 12. Kanafani ZA, Kanj SS, Cabell CH, Cecchi E, Oliveira Ramos A, Lejko-Zupanc T, Pappas
18 PA, Giamerellou H, Gordon D, Michelet C, Muñoz P, Pachirat O, Peterson G, Tan R-S,
19 Tattevin P, Thomas V, Wang A, Wiesbauer F, Sexton DJ. Revisiting the effect of referral
20 bias on the clinical spectrum of infective endocarditis in adults. *Eur J Clin Microbiol Infect*
21 *Dis* 2010;29:1203–1210.
- 22 13. San Román JA, Vilacosta I, López J. Comments on: ‘The infective endocarditis team:
23 recommendations from an international working group’. *Heart* 2014;100:1301.2-1302.

- 1 14. Chambers J, Sandoe J, Ray S, Prendergast B, Taggart D, Westaby S, Arden C, Grothier
2 L, Wilson J, Campbell B, Gohlke-Bärwolf C, Mestres CA, Rosenhek R, Pibarot P, Otto
3 C. The infective endocarditis team: recommendations from an international working
4 group. *Heart* 2014;100:524–527.
- 5 15. Li JS, Sexton DJ, Mick N, Nettles R, Fowler VG, Ryan T, Bashore T, Corey GR. Proposed
6 Modifications to the Duke Criteria for the Diagnosis of Infective Endocarditis. *Clin Infect*
7 *Dis* 2000;30:633–638.
- 8 16. Lancellotti P, Tribouilloy C, Hagendorff A, Popescu BA, Edvardsen T, Pierard LA,
9 Badano L, Zamorano JL, Scientific Document Committee of the European Association of
10 Cardiovascular Imaging. Recommendations for the echocardiographic assessment of
11 native valvular regurgitation: an executive summary from the European Association of
12 Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2013;14:611–644.
- 13 17. Lancellotti P, Pibarot P, Chambers J, Edvardsen T, Delgado V, Dulgheru R, Pepi M,
14 Cosyns B, Dweck MR, Garbi M, Magne J, Nieman K, Rosenhek R, Bernard A,
15 Lowenstein J, Vieira MLC, Rabischoffsky A, Vyhmeister RH, Zhou X, Zhang Y,
16 Zamorano J-L, Habib G. Recommendations for the imaging assessment of prosthetic heart
17 valves: a report from the European Association of Cardiovascular Imaging endorsed by
18 the Chinese Society of Echocardiography, the Inter-American Society of
19 Echocardiography, and the Brazilian Department of Cardiovascular Imaging. *Eur Heart J*
20 *Cardiovasc Imaging* 2016;17:589–590.
- 21 18. Nashef SAM, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, Lockowandt U.
22 EuroSCORE II. *Eur J Cardio-Thorac Surg Off J Eur Assoc Cardio-Thorac Surg*
23 2012;41:734–744; discussion 744-745.

- 1 19. Cresti A, Chiavarelli M, Scalese M, Nencioni C, Valentini S, Guerrini F, D'Aiello I, Picchi
2 A, De Sensi F, Habib G. Epidemiological and mortality trends in infective endocarditis, a
3 17-year population-based prospective study. *Cardiovasc Diagn Ther* 2017;7:27–35.
- 4 20. Habib G, Erba PA, Iung B, Donal E, Cosyns B, Laroche C, Popescu BA, Prendergast B,
5 Tornos P, Sadeghpour A, Oliver L, Vaskelyte J-J, Sow R, Axler O, Maggioni AP,
6 Lancellotti P, EURO-ENDO Investigators, Gale CP, Beleslin B, Budaj A, Chioncel O,
7 Dagres N, Danchin N, Emberson J, Erlinge D, Glikson M, Gray A, Kayikcioglu M,
8 Maggioni AP, Nagy VK, et al. Clinical presentation, aetiology and outcome of infective
9 endocarditis. Results of the ESC-EORP EURO-ENDO (European infective endocarditis)
10 registry: a prospective cohort study. *Eur Heart J* 2019;40:3222–3232.
- 11 21. Selton-Suty C, Célard M, Le Moing V, Doco-Lecompte T, Chirouze C, Iung B, Strady C,
12 Revest M, Vandenesch F, Bouvet A, Delahaye F, Alla F, Duval X, Hoen B. Preeminence
13 of *Staphylococcus aureus* in Infective Endocarditis: A 1-Year Population-Based Survey.
14 *Clin Infect Dis* 2012;54:1230–1239.
- 15 22. Fowler VG, Miro JM, Hoen B, Cabell CH, Abrutyn E, Rubinstein E, Corey GR, Spelman
16 D, Bradley SF, Barsic B, Pappas PA, Anstrom KJ, Wray D, Fortes CQ, Anguera I, Athan
17 E, Jones P, Meer JTM van der, Elliott TSJ, Levine DP, Bayer AS, ICE Investigators for
18 the. *Staphylococcus aureus* Endocarditis: A Consequence of Medical Progress. *JAMA*
19 2005;293:3012.
- 20 23. Di Salvo G, Habib G, Pergola V, Avierinos JF, Philip E, Casalta JP, Vailloud JM,
21 Derumeaux G, Gouvernet J, Ambrosi P, Lambert M, Ferracci A, Raoult D, Luccioni R.
22 Echocardiography predicts embolic events in infective endocarditis. *J Am Coll Cardiol*
23 2001;37:1069–1076.

- 1 24. Steckelberg JM. Emboli in Infective Endocarditis: The Prognostic Value of
2 Echocardiography. *Ann Intern Med* 1991;114:635.
- 3 25. Fernández- Hidalgo N, Almirante B, Tornos P, González- Alujas MT, Planes AM,
4 Larrosa MN, Sambola A, Igual A, Pahissa A. Prognosis of left- sided infective
5 endocarditis in patients transferred to a tertiary- care hospital—prospective analysis of
6 referral bias and influence of inadequate antimicrobial treatment. *Clin Microbiol Infect*
7 2011;17:769–775.
- 8 26. Habib G, (France), Badano L, (Italy), Tribouilloy C, (France), Vilacosta I, (Spain),
9 Zamorano JL, (Spain), Scientific Committee:, Galderisi M, (Italy), Voigt J-U, (Belgium),
10 Sicari R, (Italy), Document Reviewers:, Cosyns B, (Belgium), Fox K, (UK), Aakhus S,
11 (Norway), On behalf of the European Association of Echocardiography.
12 Recommendations for the practice of echocardiography in infective endocarditis. *Eur J*
13 *Echocardiogr* 2010;11:202–219.
- 14 27. Chu VH, Park LP, Athan E, Delahaye F, Freiburger T, Lamas C, Miro JM, Mudrick DW,
15 Strahilevitz J, Tribouilloy C, Durante-Mangoni E, Pericas JM, Fernández-Hidalgo N,
16 Nacinovich F, Rizk H, Krajcinovic V, Giannitsioti E, Hurley JP, Hannan MM, Wang A.
17 Association Between Surgical Indications, Operative Risk, and Clinical Outcome in
18 Infective Endocarditis: A Prospective Study From the International Collaboration on
19 Endocarditis. *Circulation* 2015;131:131–140.
- 20 28. López-Dupla M, Hernández S, Olona M, Mercé J, Lorenzo A, Tapiol J, Gómez F,
21 Santamaría J, García R, Auguet T, Richart C, Castells E, Bardají A, Vidal F. [Clinical
22 characteristics and outcome of infective endocarditis in individuals of the general
23 population managed at a teaching hospital without cardiac surgery facilities. Study of 120
24 cases]. *Rev Esp Cardiol* 2006;59:1131–1139.

- 1 29. Oliver L, Lavoute C, Giorgi R, Salaun E, Hubert S, Casalta J-P, Gouriet F, Renard S, Saby
2 L, Avierinos J-F, Maysou L-A, Riberi A, Grisoli D, Casalta A-C, Collart F, Raoult D,
3 Habib G. Infective endocarditis in octogenarians. *Heart* 2017;103:1602–1609.
- 4 30. Ramírez-Duque N, García-Cabrera E, Ivanova-Georgieva R, Noureddine M, Lomas JM,
5 Hidalgo-Tenorio C, Plata A, Gálvez-Acebal J, Ruíz-Morales J, Torre-Lima J de la,
6 Reguera JM, Martínez-Marcos FJ, Alarcón A de. Surgical treatment for infective
7 endocarditis in elderly patients. *J Infect* 2011;63:131–138.

8

9

10

11 **Figure legend**

12

13 **Figure 1 : Probability of survival according to health-care pathway**

14 A: Comparison of survival between the three groups

15 B: Comparison of survival between Referral Centers (RC) and Transferred to Referral Centers
16 (TRC) groups

17 C: Comparison of survival between RC and Non-Referral Centers (NRC) groups

18 D: Comparison of survival between groups RC and TRC together and NRC group

19

20

21

22

23

24

25

1 **Investigators and sites**

2 **La Timone Hospital Marseille:** Florent Arregle, MD, Mary Philip, MD, SanRedrine Hubert,
3 MD, Frédérique Gouriet, MD PhD, Jean-Paul Casalta, MD, Frédéric Collart, MD, Alberto
4 Riberi, MD, Héléne Martel, MD, Sébastien Renard, MD, Laurence Camoin ,MD, PhD, Anne
5 Claire Casalta , MD, Hubert Lepidi ,MD, PhD, Michel Drancourt , MD, PhD, Didier Raoult
6 ,MD, PhD, Gilbert Habib MD, PhD. **CHU Nord Marseille:** Franck Thuny. MD, Yves
7 Frances. MD, Karim Aissi. MD, Philippe Brouqui. MD, Matthieu Million. MD, **CHU de**
8 **Montpellier** Catherine Sportouch. MD, Stéphane Cade. MD, Frédéric Cransac, MD. **CHU de**
9 **Nîmes:** Patrick Messner-Pellenc. MD, Camille Soulier, MD, Jean-etienne Ricci, MD,
10 **Hôpital d’Instruction des Armées Laveran,** Marseille: Laurent Fourcade, MD, Pierre-
11 Laurent Massoure, MD, Jean-Marie Gil. MD, **Hôpital Ambroise Paré-Desbief, Marseille:**
12 Emmanuel Philip. MD, **Fondation Saint Joseph, Marseille:** Patrick Khanoyan. MD, Roger
13 Rosario, MD, Nicolas Michel, MD, **CH de Martigues:** Serge Yvorra. MD, Virginie
14 Chellini. MD, **CH d’Avignon:** Stéphanie Branger. MD, Stéphane Andrieux, MD, Saida
15 Cheggour. MD, Salah Aboukoudir, MD. **Clinique Rhône-Durance, Avignon.** Laurent
16 Meille. MD, Catherine Meulemans. MD, **CH Toulon:** Jean-Michel Tartière. MD, Danielle
17 Raufast. MD, Isabelle LeCardonnel. MD, **Hôpital d’Instruction des Armées Saint-Anne,**
18 **Toulon:** Christophe Jego. MD, Gilles Cellarier. MD, Raphael Poyet. MD, **Polyclinique Les**
19 **Fleurs, Ollioules:** Philippe Pietri. MD, Philippe Villain. MD, Jean-Paul Giorgi. MD,
20 Chirine Parsai, MD, Olivier Brissy. MD, **CH Aix-en-Provence:** Bernard Jouve. MD, Tewfik
21 Benchaa. MD, **CH d’Arles:** Salah Si Ahmed MD, François Saint-Pierre. MD, **CH de**
22 **Draguignan:** André-François Chaix. Pierrick Boyer. MD, **Clinique La Casamance,**
23 **Aubagne** Véronique Haddad. MD, **CH des Alpes du Sud, Gap-Sisteron:** Christian Tafani.
24 MD, Amar Hidoud, MD, Audrey Boudes , MD, Hakim Berguigua, MD, Fabien Devemy,
25 MD, Bernard d’Hautefeuille, MD, **CH de Manosque** Alain Rapuzzi. MD, **Clinique des**

- 1 **Franciscaines, Nîmes:** Vlad Ciobataru. MD, **Hôpital de Fréjus:** Xavier Lamit. MD,
- 2 **Hôpital de Bastia:** Aurelia Tho, MD, Thierry James, MD, Abdelkader Bensalah. MD.
- 3
- 4
- 5

1 **Abstract**

2 **Aims:** To determine the prognosis of patients treated for infective endocarditis (IE) according
3 to their health-care pathway. To assess how the ESC guidelines are implemented concerning
4 the performance of transesophageal echocardiography (TEE), the use of antibiotic therapy, and
5 the performance of valve surgery; and to compare the epidemiological profile of IE according
6 to the type of centers in which the patients are hospitalized.

7 **Methods and results:** In a prospective multicentric study including 22 hospitals in the South-
8 East of France, 342 patients were classified in three groups according to their health-care
9 pathway: 119 patients diagnosed and taken care entirely in a reference center or hospital with
10 cardiac surgery (**Referral Center (RC) group**), 111 patients diagnosed and initially taken care
11 in a non-referral center (NRC) , then referred in a center including cardiac surgery (**Transferred
12 to Referral Center (TRC) group**) and 112 patients totally taken care in **NRC (NRC) group**). One-
13 year mortality was 26% (88 deaths) and was not significantly different between groups 1 and 2
14 (20% vs 21%, $p = 0.83$). Patients in **NRC** group had a higher mortality (37%) compared to
15 patients in **RC and TRC** groups ($p < 0.001$). ESC guidelines were not implemented similarly
16 depending on the health-care pathway ($p = 0.04$). Patients in **NRC** group were significantly older
17 ($p < 0.001$) and had more comorbidities ($p < 0.001$) than patients treated in referral centers.

18 **Conclusion:** Prognosis of patients with IE is influenced by their health-care pathway. Patients
19 treated exclusively in NRC have a worse prognosis than patients treated in referral or surgical
20 centers.

21

22 **Keywords :** endocarditis, mortality, health-care pathway, referral center

23

24

25

1 **Introduction**

2 Epidemiologic profile of infective endocarditis (IE) has changed over the past twenty
3 years,^{1 2 3} but mortality remains high^{4 5 6} despite diagnostic improvement and access to surgery
4 in up to 50% of cases.⁷ This persistent severity has justified the development of new diagnostic
5 and therapeutic strategies to improve prognosis.^{8 9} Management by a specialized
6 multidisciplinary team within a referent medical-surgical center seems to be a key factor in
7 improving patients prognosis,^{10 11} and is recommended by the ESC guidelines.^{8 9}

8 However, few studies have compared the prognosis and management of patients with
9 IE according to their health-care pathway. Indeed, three situations are usually observed: patients
10 for whom the diagnosis and management are carried out entirely in a referral center and/or
11 having a cardiac surgery department, patients secondarily referred to a referral hospital or
12 surgical center by another care center, patients for whom the entire care is provided within a
13 non-referral care center. In addition, application of European guidelines remains difficult to
14 assess in daily clinical practice outside the major referral centers. Published epidemiological
15 data only represent part of the IE, most often not taking into account patients hospitalized in
16 non-referral centers and therefore introducing a significant bias in the knowledge of this
17 pathology.¹² Finally, although the ESC guidelines suggest that patients with uncomplicated IE
18 can be managed in a non-reference center⁹, if all patients with IE¹³ or only the sickest patients
19¹⁴ should be referred to reference centers is still a matter of debate.

20 The main objective of this study was to determine the prognosis of patients treated for
21 IE according to their health-care pathway. The secondary objectives were to assess how the
22 ESC guidelines are implemented concerning the performance of transesophageal
23 echocardiography (TEE), the use of antibiotic therapy, and the performance of valve surgery,
24 and to compare the epidemiological profile of IE according to the type of centers in which the
25 patients are hospitalized.

1 **Methods**

2 **Study design and patients**

3 This was a prospective multicentric study including patients consecutively admitted for
4 suspicion of IE in 22 hospitals in South-East of France between January 2014 and June 2017.
5 These hospitals were both tertiary centers with cardiac surgery and second-level community
6 hospitals. Inclusion criteria were a diagnosis of definite IE based on Duke modified criteria ¹⁵
7 on native, prosthetic or pacemaker/implantable automatic defibrillator. Non-inclusion criteria
8 were age < 18 years and patients already included in the study and presenting with a new IE
9 diagnosis. **Patients were screened in the echocardiography lab of each institution and were**
10 **prospectively included by local investigators.**

11

12 **Patient and Public involvement**

13 All patients hospitalized for suspected IE were accepted on admission to participate in this
14 research protocol. However, they were not involved in the design, conduct, reporting or
15 dissemination plans of our research. The study complies with the Declaration of Helsinki.
16 Written consent was waived by La Timone Institutional Review Board, which gave its approval
17 for this study.

18

19 **Data collected at the time of inclusion and during hospitalization**

20 The following clinical data were collected: age, sex, previous cardiac and non-cardiac
21 diseases, Charlson index score, date of first clinical signs, body temperature, presence of cardiac
22 murmur, congestive heart failure, neurological complications, septic or cariogenic shock or
23 atrio-ventricular block on electrocardiogram at time of admission and during hospitalization.

1 Biological data were collected, including hemoglobin level, platelet count, leukocytes count,
2 C-Reactive Protein serum level, creatinine level, rheumatoid factor. Microbiological data
3 included: positive or negative blood cultures, type of microorganism and antibiogram.

4 Echocardiographic data collected were: presence of vegetation and its maximal length,
5 presence of periannular lesion (defined as the presence of abscess and/or pseudoaneurysm
6 and/or fistula), presence of valvular regurgitation, quantified according to current
7 recommendations.^{16 17}

8 Type and date of the following complications were collected: congestive heart failure,
9 peripheral embolism, neurological complications (including symptomatic and silent), mycotic
10 aneurysm, acute kidney failure (>26 μ mol/L increase of serum creatinine in 48 hours or >50%
11 increase in 7 days) , glomerulonephritis, uncontrolled infection (fever and/or positive blood
12 cultures after 7 days of appropriate antibiotic therapy and unlinked to an extracardiac cause).

13 Moreover, type, dose and length of antibiotic therapy was reported. Antibiotic therapy
14 was considered appropriate if drugs used and full duration of treatment were in accordance with
15 ESC guidelines (taking into account patients terminating antibiotics prematurely due to death).
16 Analysis of antibiotic treatment was conducted retrospectively by the Endocarditis team of the
17 investigator center (CHU La Timone, Marseille).

18 During hospitalization or follow-up the indication (according to ESC guidelines) and
19 date of surgery was also collected. Finally, surgical risk was evaluated using the Euroscore II.¹⁸

20

21 **Classification of patients according to health-care pathway**

22 Patients were classified in three groups according to their health-care pathway:

23 - Referral Center group (RC), including patients diagnosed and taken care entirely in a
24 reference center or hospital with cardiac surgery (4 hospitals). These centers were the
25 only surgical centers of this region of France.

- 1 - Transferred to Referral Center group (TRC), including patients diagnosed and initially
2 taken care in a non-referral center, then referred in a center including cardiac surgery
3 - Non-Referral Center group (NRC), including patients totally taken care in non-referral
4 centers (18 centers)

5

6 **Follow-up**

7 Patients had a one-year follow-up with a clinical evaluation or if not possible, phone contact.

8

9 **Primary and secondary endpoints**

10 Primary endpoint was death from all cause at one-year follow-up. Secondary endpoints
11 were:

12 - the evaluation of the implementation of European guidelines^{8 9} in the three patients groups
13 regarding: the use of TEE, the choice and duration of antibiotic therapy, and the performance
14 of valvular surgery and/or extraction of pacemaker or implantable automatic defibrillator when
15 indicated

16 - comparison of demographical and microbiological data among the three groups.

17

18 **Statistical analysis**

19 Continuous variables were described by their mean, standard deviation, minimum and
20 maximum value, 1st and 3rd quartile. Continuous variables were compared using the Mann-
21 Whitney non parametric U test or Student's t test, depending on the application conditions.
22 Categorical variables were compared using the chi-square test or the Fisher test, depending on
23 the application conditions. The event-free survival distributions were estimated with the
24 Kaplan-Meier method. The log-rank test was used to compare survival distributions for
25 categorical variables. The effect of continuous variables on the risk of an event occurring was

1 estimated and tested using a Cox model. The surgery variable, which is a time dependent
2 variable, was estimated using a Poisson approach.

3 Multivariate analyzes were tested using a Cox model. Multivariate analyzes were tested
4 using a Cox model. Inclusion of variables in the fitted model was carried out according to the
5 following criteria : result in univariate analysis with statistical significance $p < 0.20$, number of
6 missing values $< 10\%$ for this variable, effective presence in both categories of a bivariate
7 variable $> 10\%$, as well as according to their clinical relevance. The final models were selected
8 using the Akaike Information Criteria (AIC).

9 The comparison tests of group 1 to group 2, group 1 to group 3, group 1 + 2 to group 3
10 were performed in bilateral situation using Bonferroni correction and were considered
11 statistically significant for $p < 0.017$. For all other tests, they were performed bilaterally and
12 were considered statistically significant for $p < 0.05$. Statistical analysis was performed with R
13 software (version 3.6.2) and RStudio (v 1.2.5).

14

15

16

17 **Results**

18 **Patient population**

19 Between January 2014 and June 2017, a total of 342 patients with definite IE were
20 included. Among them, 119 were diagnosed and entirely managed in a reference center or
21 hospital with cardiac surgery (RC group), 111 were diagnosed and initially managed in a non-
22 referral center, then referred in a center including cardiac surgery (TRC group) and 112 were
23 totally managed in non-referral centers (NRC group). Clinical, demographical, biological, and
24 microbiological characteristics of patients are presented in **Table 1**.

25

1 *Overall population*

2 In the overall population, mean age was 68 +/- 15 years. IE were mainly community-
3 acquired (83% of cases), *S. aureus* was the most frequent bacteria involved (26% of cases).
4 Negative blood culture IE represented 17% of cases. Left-sided IE counted for 85% of all cases
5 and aortic (43%) and mitral (42%) localization were almost equally represented. Perivalvular
6 complications were observed in 16% of cases. 33% of cases were prosthetic valve IE. More
7 than half IE cases (58%) showed theoretical indication for valvular surgery or intracardiac
8 device removal but only 35% had such interventions.

9

10 *Characteristics according to health-care pathway (Table 1)*

11 Patients in NRC group were significantly older and had higher Charlson index. Sources
12 of infection did not differ between the three groups and microbiological profile of IE cases was
13 not significantly different. Echocardiographic findings showed more severe lesions in RC group
14 and TRC group: longer vegetations, more peri-valvular complications, and more severe
15 valvular regurgitation. Concerning IE complications, there was no significant difference
16 between groups in embolic events, however more silent embolisms were diagnosed in RC group
17 and TRC group. Spondylodiscitis was more frequent in NRC group.

18 Patients in RC and TRC groups had more frequent surgical indication, but surgery or
19 intracardiac device removal was performed in only 52 patients (44%) in RC group, 60
20 patients (55%) in TRC group and 6 patients (5%) in NRC group, $p < 0.001$. In RC group, 81
21 patients had an indication for intervention and 29 were not operated and in TRC group, 91
22 patients had an indication for intervention and 31 were not operated. Reasons for absence of
23 intervention were: deaths before surgery (6 patients in RC group and 2 patients in TRC group
24), patient's refusal (2 patients in RC group and 1 patient in TRC group) and decision of the
25 endocarditis team due to comorbidities (21 patients in RC group and 28 patients in TRC

1 group). Patients who benefited from surgery were significantly younger (69.4 +/- 13 years vs
2 62.4 +/- 12 years, $p=0.01$), had lower Charlson index (3.9 +/- 1.9 vs 2.8 +/- 2.7, $p=0.007$),
3 had more frequently vegetations (86 patients (73%) vs 95 patients (85%), $p=0.002$) and severe
4 valvular regurgitation (41 patients (35%) vs 68 patients (61%), $p<0.001$) but less renal failure
5 (29 patients (25%) vs 5 patients (4%), $p<0.001$) and diabetes (39 patients (33%) vs 20 patients
6 (18%), $p=0.01$).

7 Time from diagnosis to intervention did not significantly differ between RC group and
8 TRC group (mean time 30+/-32 days vs 22+/-29 days, $p=0.36$). In our cohort, mean time for
9 right-sided IE was 17+/-18 days and 25+/-28 days for left-sided IE.

10

11 Prognosis of patients according to their health-care pathway

12 Among the 342 patients, one year mortality was 26% (88 deaths), hospital mortality was
13 13% (44 deaths) and one year recurrence rate was 4% (13 patients). **Figure 1** shows survival
14 probability according to health-care pathway.

15 Hospital mortality was 11% (13 deaths) in RC group, 8% (9 patients) in TRC group and
16 20% (22 patients) in NRC group, $p=0.03$. One year mortality was 20% (24 deaths) in RC group,
17 21% (23 deaths) in TRC group and 37% (41 deaths) in NRC group, $p=0.003$. One year mortality
18 was not significantly different between RC and TRC groups ($p=0.83$), however it was
19 significantly higher in NRC group compared to RC and TRC groups ($p<0.001$).

20

21 Characteristics of patients only treated in non-referral centers

22 During the study period, 112 patients were managed only in a NRC. Among them, 28
23 patients (25%) had indication for valvular surgery or intracardiac device removal. These
24 patients had a mean age of 76+/-11 years and were at high surgical risk (mean Euroscore 17+/-
25 16%). Reasons for absence of transfer and surgery included: death before surgery for 16 patients

1 (14%), medical choice for 14 patients (13%), patient's refusal in 3 cases (3%) and high surgical
2 risk (comorbidities) for 16 patients (14%). 6 patients (5%) had intracardiac device removal
3 (pacemaker lead or catheter) in the **NRC**. Among these 28 patients, one year mortality was 50%
4 (14 deaths). 84 patients (75%) had indication for medical treatment of IE. One year mortality
5 among them was 29% (24 deaths).

6

7 **Prognostic factors of mortality**

8 Results of univariate analysis of prognostic factors of one year mortality are shown in
9 **Table S1. Table 2** shows prognostic factors of mortality among the 342 patients in multivariate
10 analysis. Belonging to **NRC** group was associated with higher risk of death (HR 2.56; 95% CI
11 1.44-4.55, $p=0.001$). Heart failure (HR 2.4; 95% CI 1.52-3.78, $p<0.001$) and peri-valvular
12 involvement (HR 3.25; 95% CI 1.86-5.68, $p<0.001$) were IE complications predictive of
13 mortality.

14

15 **Evaluation of European guidelines application according to health-care pathway**

16 **Table 3** shows the implementation of ESC guidelines in the three groups regarding use
17 of TEE, valvular surgery or intracardiac device removal and antibiotic therapy. In overall
18 population, ESC guidelines were correctly implemented in 52% of patients. TEE was
19 performed in most IE cases (92%); however it was significantly less frequently used in **NRC**
20 group as compared to the two other groups ($p<0.001$). Guidelines concerning surgery or
21 intracardiac device extraction were correctly implemented in 78% of cases and there was no
22 significant difference between the three groups. Antibiotic therapy was in accordance with
23 guidelines in 72% of cases and there was no significant difference between patients taken care
24 entirely or partially in **RC** ($p=0.66$). However, treatment was less frequently in agreement with
25 guidelines in patients in NRC group compared to the others ($p<0.001$).

1 Discussion

2 This prospective study carried out in 22 hospitals in the South-East of France including
3 342 IE cases divided into three groups according to their health-care pathway shows that:

4 - The prognosis of patients is influenced by their health-care pathway

5 - The number of patients managed in non-referral centers is high. They are older, have more
6 comorbidities, and less severe cardiac lesions than the others.

7 - Surgery or device removal is less frequently performed in those patients

8 - European recommendations on the management of these patients are not followed similarly
9 depending on their health-care pathway

10 - Both in-hospital and long-term mortalities are higher in non-referral centers.

11

12 General population

13 By including patients treated in **NRC**, our study has the advantage of providing global
14 epidemiological information concerning the cases of IE within the south-eastern region,
15 reflecting the management of this pathology in "real life" and not only in **RC**. Almost a third of
16 the patients treated for IE during the inclusion period were treated in a **NRC**.

17 Consistent with known data ^{19 20}, there was a predominance of IE cases in males (69%).

18 The mean age of diagnosis of IE was 68 years in our study. This age seemed more advanced in
19 comparison to other French (average age of 59 +/- 17 years in 1999 ², 62 +/- 16 years in 2008
20 ²¹) or European studies (57 +/- 16 years in 2001 ³, 59 +/- 18 years in the Euro-Endo registry ²⁰).

21 Concerning microbiological data, *S. aureus* was the predominant germ (26% of IE), which was
22 in agreement with previous studies ^{21 22}. There was a majority of left-sided IE (85%). An
23 embolic event was observed in 38% of cases. This rate was comparable to other studies that
24 describe the occurrence of an embolic event in 20 to 50% of cases ^{21 23 24}

25

1 **Mortality according to health-care pathway**

2 In our cohort, hospital mortality was 13% and one-year mortality was 26%. Few studies
3 have compared the prognosis of patients with IE according to their health-care pathway. A
4 study by Fernandez-Hidalgo et al. ²⁵, compared the hospital mortality of 223 patients treated in
5 a **RC** with cardiac surgery with 144 patients taken care in a NRC and secondarily referred to
6 the **RC**. This study did not show any significant difference in terms of hospital mortality
7 between the two groups. Likewise, an international multicenter study ¹² carried out in **RC** in the
8 management of IE did not show difference in hospital mortality between 1,164 patients
9 transferred from NRC and 1,596 patients treated directly within the RC. However, these studies
10 have some limitations: by considering only patients treated in RC they are not representative of
11 epidemiology of patients treated in NRC. The originality of our study is to compare three
12 groups, including a group of patients treated exclusively in **NRC**, these patients usually being
13 not included in previous studies. Our main findings is a significantly higher hospital (20%) and
14 one year mortality (37%) in patients treated only in NRC. Belonging to **NRC** group was an
15 independent risk factor in multivariate analysis of one-year mortality in the total cohort of
16 patients. This excess of mortality observed in **NRC** group could be explained by various factors:
17 epidemiological differences compared to other groups, recommendations on the management
18 of IE that are less implemented and a higher absence of surgery rate in patients with a theoretical
19 surgical indication.

20

21 **Implementation of European guidelines**

22 In our study, we observed a more frequent use of TEE (92% of patients) than in other
23 multicenter studies (60-75% ^{3 20}). However, within **NRC group**, TEE was significantly less
24 frequently performed compared to the other groups. TEE has a key role in the diagnostic
25 process of IE and in particular in the detection of peri-annular complications and should be

1 performed systematically, in accordance with current recommendations ^{8 9 26}. Thus, it is
2 possible that these complications, often requiring surgery, have not been diagnosed in some
3 patients, especially in **NRC group**.

4 Regarding valve surgery or the extraction of intracardiac material, there was no
5 significant difference in application of recommendations between the three groups. However,
6 we noted within each group patients with a theoretical indication for surgery for who surgery
7 was finally not performed. Within **NRC** group, 25% of patients presented a theoretical
8 indication and only 5% received interventional management. Conversely, in **RC and TRC**
9 groups, 65% of patients with theoretical surgical indication benefited from intervention.
10 Different studies have already shown the poor prognosis of patients with a theoretical indication
11 for surgery and not operated ^{20 27}, which may in part explain the higher mortality of patients in
12 **NRC** group. In addition, the main multicenter studies carried out over the last few years
13 highlight a surgery rate close to 50% (52% in Euro Heart Survey ³, 51% in the EuroEndo
14 register ²⁰, 48% in the ICE register ⁴). Among the patients treated exclusively in **RC**, the rate of
15 surgery or extraction of intracardiac material was 44%, close to previous published data. On
16 the other hand, if we considered all the patients for whom the diagnosis of IE was made in a
17 non-referral center (**TRC and NRC** groups), the overall rate of valve surgery or extraction of
18 intracardiac material was only 30 %. Several factors could explain this discrepancy with the
19 pre-existing studies. Indeed, most studies on IE focused on patients treated in **RC** with a cardiac
20 surgery department. It is possible that a referral bias may cause an overestimation of the rate of
21 cardiac surgery in this pathology, by not considering many IE of "medical" management and
22 treated exclusively in **NRC**. On the other hand, it is also possible that among the patients
23 diagnosed in a **NRC**, surgical indications were sometimes not identified, especially in **NRC**
24 group in which, unlike **TRC** group, it was not possible to re-evaluate the surgical indications
25 by a multidisciplinary team in a reference center. In a same way, a Spanish study by Lopez-

1 Dupla et al. ²⁸ in a university center without a cardiac surgery department showed that the
2 establishment of a multidisciplinary team including a cardiac surgeon was associated with an
3 increase in the rate of surgical indication identified (14.5% vs 34.5%). Finally, the European
4 guidelines concerning the antibiotic treatment were significantly less applied among patients in
5 group 3 with 35% of inappropriate antibiotic therapy. Here again, few studies concerning the
6 antibiotic treatment used in non-referral centers are available. Fernandez-Hidalgo et al. ²⁵
7 showed that among the patients diagnosed in a NRC and referred secondarily to a surgical
8 center, the antibiotic treatment initiated in the non-referral center was inadequate in 54% of
9 cases. In this study, inadequate treatment was associated with an increased risk of in-hospital
10 mortality in multivariate analysis. The goal of our study was to evaluate the implementation of
11 ESC guidelines in “real life”. It is however important to highlight that in some cases, there may
12 be logical reasons why patients do not undergo an indicated procedure such as TEE, for instance
13 when a patient is not operable, or surgery.

14

15 **Epidemiological characteristics of IE according to health-care pathway**

16 The epidemiological profile of patients in RC and TRC groups was comparable. These
17 were patients with more severe IE than in NRC group with more hemodynamic complications,
18 a higher rate of vegetations, more peri-annular complications. These elements partly explain
19 the higher surgical indication rate in these two groups. The patients in NRC group were
20 significantly older, presented more comorbidities, and a greater operative risk. Some studies
21 focusing on IE cases in the elderly have shown, including during treatment in a RC, limited
22 access to surgery and significant mortality in these patients when surgery, although indicated,
23 was not performed ^{29 30}.

24

25

1 **Limitations**

2 Some data could not be recorded, such as the reason for the transfer of patients to a
3 referral center, the time between diagnosis and the transfer of patients, and some factors who
4 could have influenced transfer decision (such as cirrhosis or poor nutritional status). Due to the
5 large number of participating non-referral centers, the reasons for transferring patients may
6 have differed from one center to another. Moreover, modified Duke criteria were used for
7 diagnosis of IE and have limited sensitivity. This study reflects the epidemiology and practices
8 used in the south-eastern region of France and cannot be generalized to all the other centers.
9 Finally, the study design was meant to compare prognosis of patients according to the health-
10 care pathway but reasons explaining the different prognosis observed could only be
11 hypothesized.

12

13 **Conclusion**

14 Prognosis of patients with IE is influenced by their health-care pathway. Patients treated
15 exclusively in NRC have a worse prognosis than patients treated in RC. These patients are often
16 fragile and their management difficult. Given the difference in mortality between patients
17 managed in referral vs non-referral centers, the current recommendations of treating patients
18 with non-complicated IE in non-referral centers should be questioned. IE is a deadly disease
19 with a better prognosis when managed in RC.

20

21 **Supplementary data**

22 Supplementary material is available online.

23

24 **Funding**

1 This study was supported by a grant from: the PHRCI (Projet Hospitalier de Recherche
2 Clinique Inter-régional 2012)

3

4 **Conflict of interest**

5 None declared

6

7 **Data availability**

8 The data underlying this article will be shared on reasonable request to the corresponding
9 author.

10

11

12

13

14 **References**

15 1. Thuny F, Grisoli D, Collart F, Habib G, Raoult D. Management of infective endocarditis:
16 challenges and perspectives. *The Lancet* 2012;379:965–975.

17 2. Hoen B. Changing Profile of Infective Endocarditis Results of a 1-Year Survey in France.
18 *JAMA* 2002;288:75.

19 3. Tornos P. Infective endocarditis in Europe: lessons from the Euro heart survey. *Heart*
20 2005;91:571–575.

21 4. Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, Karchmer AW,
22 Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones
23 P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell
24 CH, International Collaboration on Endocarditis-Prospective Cohort Study (ICE-PCS)

- 1 Investigators. Clinical presentation, etiology, and outcome of infective endocarditis in the
2 21st century: the International Collaboration on Endocarditis-Pro prospective Cohort Study.
3 *Arch Intern Med* 2009;169:463–473.
- 4 5. Gálvez-Acebal J, Rodríguez-Baño J, Martínez-Marcos FJ, Reguera JM, Plata A, Ruiz J,
5 Marquez M, Lomas JM, Torre-Lima J de la, Hidalgo-Tenorio C, Alarcón A de, Grupo
6 para el Estudio de las Infecciones Cardiovasculares de la Sociedad Andaluza de
7 Enfermedades Infecciosas (SAEI). Prognostic factors in left-sided endocarditis: results
8 from the Andalusian multicenter cohort. *BMC Infect Dis* 2010;10:17.
- 9 6. Thuny F, Disalvo G, Belliard O, Avierinos J-F, Pergola V, Rosenberg V, Casalta J-P,
10 Gouvernet J, Derumeaux G, Iarussi D, Ambrosi P, Calabro R, Riberi A, Collart F, Metras
11 D, Lepidi H, Raoult D, Harle J-R, Weiller P-J, Cohen A, Habib G. Risk of Embolism and
12 Death in Infective Endocarditis: Prognostic Value of Echocardiography: A Prospective
13 Multicenter Study. *Circulation* 2005;112:69–75.
- 14 7. Thuny F, Beurtheret S, Mancini J, Gariboldi V, Casalta J-P, Riberi A, Giorgi R, Gouriet
15 F, Tafanelli L, Avierinos J-F, Renard S, Collart F, Raoult D, Habib G. The timing of
16 surgery influences mortality and morbidity in adults with severe complicated infective
17 endocarditis: a propensity analysis. *Eur Heart J* 2011;32:2027–2033.
- 18 8. Endorsed by the European Society of Clinical Microbiology and Infectious Diseases
19 (ESCMID) and by the International Society of Chemotherapy (ISC) for Infection and
20 Cancer, Authors/Task Force Members, Habib G, Hoen B, Tornos P, Thuny F, Prendergast
21 B, Vilacosta I, Moreillon P, Jesus Antunes M de, Thilen U, Lekakis J, Lengyel M, Muller
22 L, Naber CK, Nihoyannopoulos P, Moritz A, Zamorano JL, ESC Committee for Practice
23 Guidelines (CPG), Vahanian A, Auricchio A, Bax J, Ceconi C, Dean V, Filippatos G,
24 Funck-Brentano C, Hobbs R, Kearney P, McDonagh T, McGregor K, et al. Guidelines on

- 1 the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): The
2 Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the
3 European Society of Cardiology (ESC). *Eur Heart J* 2009;30:2369–2413.
- 4 9. Habib G, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta J-P, Del Zotti F, Dulgheru
5 R, El Khoury G, Erba PA, Iung B, Miro JM, Mulder BJ, Plonska-Gosciniak E, Price S,
6 Roos-Hesselink J, Snygg-Martin U, Thuny F, Tornos Mas P, Vilacosta I, Zamorano JL.
7 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for
8 the Management of Infective Endocarditis of the European Society of Cardiology
9 (ESC) Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the
10 European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015;36:3075–3128.
- 11 10. Botelho-Nevers E, Thuny F, Casalta JP, Richet H, Gouriet F, Collart F, Riberi A, Habib
12 G, Raoult D. Dramatic Reduction in Infective Endocarditis–Related Mortality With a
13 Management-Based Approach. *Arch Intern Med* 2009;169:1290.
- 14 11. Chirillo F, Scotton P, Rocco F, Rigoli R, Borsatto F, Pedrocco A, De Leo A, Minniti G,
15 Polesel E, Olivari Z. Impact of a Multidisciplinary Management Strategy on the Outcome
16 of Patients With Native Valve Infective Endocarditis. *Am J Cardiol* 2013;112:1171–1176.
- 17 12. Kanafani ZA, Kanj SS, Cabell CH, Cecchi E, Oliveira Ramos A, Lejko-Zupanc T, Pappas
18 PA, Giamerellou H, Gordon D, Michelet C, Muñoz P, Pachirat O, Peterson G, Tan R-S,
19 Tattevin P, Thomas V, Wang A, Wiesbauer F, Sexton DJ. Revisiting the effect of referral
20 bias on the clinical spectrum of infective endocarditis in adults. *Eur J Clin Microbiol Infect*
21 *Dis* 2010;29:1203–1210.
- 22 13. San Román JA, Vilacosta I, López J. Comments on: ‘The infective endocarditis team:
23 recommendations from an international working group’. *Heart* 2014;100:1301.2-1302.

- 1 14. Chambers J, Sandoe J, Ray S, Prendergast B, Taggart D, Westaby S, Arden C, Grothier
2 L, Wilson J, Campbell B, Gohlke-Bärwolf C, Mestres CA, Rosenhek R, Pibarot P, Otto
3 C. The infective endocarditis team: recommendations from an international working
4 group. *Heart* 2014;100:524–527.
- 5 15. Li JS, Sexton DJ, Mick N, Nettles R, Fowler VG, Ryan T, Bashore T, Corey GR. Proposed
6 Modifications to the Duke Criteria for the Diagnosis of Infective Endocarditis. *Clin Infect*
7 *Dis* 2000;30:633–638.
- 8 16. Lancellotti P, Tribouilloy C, Hagendorff A, Popescu BA, Edvardsen T, Pierard LA,
9 Badano L, Zamorano JL, Scientific Document Committee of the European Association of
10 Cardiovascular Imaging. Recommendations for the echocardiographic assessment of
11 native valvular regurgitation: an executive summary from the European Association of
12 Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2013;14:611–644.
- 13 17. Lancellotti P, Pibarot P, Chambers J, Edvardsen T, Delgado V, Dulgheru R, Pepi M,
14 Cosyns B, Dweck MR, Garbi M, Magne J, Nieman K, Rosenhek R, Bernard A,
15 Lowenstein J, Vieira MLC, Rabischoffsky A, Vyhmeister RH, Zhou X, Zhang Y,
16 Zamorano J-L, Habib G. Recommendations for the imaging assessment of prosthetic heart
17 valves: a report from the European Association of Cardiovascular Imaging endorsed by
18 the Chinese Society of Echocardiography, the Inter-American Society of
19 Echocardiography, and the Brazilian Department of Cardiovascular Imaging. *Eur Heart J*
20 *Cardiovasc Imaging* 2016;17:589–590.
- 21 18. Nashef SAM, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, Lockowandt U.
22 EuroSCORE II. *Eur J Cardio-Thorac Surg Off J Eur Assoc Cardio-Thorac Surg*
23 2012;41:734–744; discussion 744-745.

- 1 19. Cresti A, Chiavarelli M, Scalese M, Nencioni C, Valentini S, Guerrini F, D'Aiello I, Picchi
2 A, De Sensi F, Habib G. Epidemiological and mortality trends in infective endocarditis, a
3 17-year population-based prospective study. *Cardiovasc Diagn Ther* 2017;7:27–35.
- 4 20. Habib G, Erba PA, Iung B, Donal E, Cosyns B, Laroche C, Popescu BA, Prendergast B,
5 Tornos P, Sadeghpour A, Oliver L, Vaskelyte J-J, Sow R, Axler O, Maggioni AP,
6 Lancellotti P, EURO-ENDO Investigators, Gale CP, Beleslin B, Budaj A, Chioncel O,
7 Dagnes N, Danchin N, Emberson J, Erlinge D, Glikson M, Gray A, Kayikcioglu M,
8 Maggioni AP, Nagy VK, et al. Clinical presentation, aetiology and outcome of infective
9 endocarditis. Results of the ESC-EORP EURO-ENDO (European infective endocarditis)
10 registry: a prospective cohort study. *Eur Heart J* 2019;40:3222–3232.
- 11 21. Selton-Suty C, Célard M, Le Moing V, Doco-Lecompte T, Chirouze C, Iung B, Strady C,
12 Revest M, Vandenesch F, Bouvet A, Delahaye F, Alla F, Duval X, Hoen B. Preeminence
13 of *Staphylococcus aureus* in Infective Endocarditis: A 1-Year Population-Based Survey.
14 *Clin Infect Dis* 2012;54:1230–1239.
- 15 22. Fowler VG, Miro JM, Hoen B, Cabell CH, Abrutyn E, Rubinstein E, Corey GR, Spelman
16 D, Bradley SF, Barsic B, Pappas PA, Anstrom KJ, Wray D, Fortes CQ, Anguera I, Athan
17 E, Jones P, Meer JTM van der, Elliott TSJ, Levine DP, Bayer AS, ICE Investigators for
18 the. *Staphylococcus aureus* Endocarditis: A Consequence of Medical Progress. *JAMA*
19 2005;293:3012.
- 20 23. Di Salvo G, Habib G, Pergola V, Avierinos JF, Philip E, Casalta JP, Vailloud JM,
21 Derumeaux G, Gouvernet J, Ambrosi P, Lambert M, Ferracci A, Raoult D, Luccioni R.
22 Echocardiography predicts embolic events in infective endocarditis. *J Am Coll Cardiol*
23 2001;37:1069–1076.

- 1 24. Steckelberg JM. Emboli in Infective Endocarditis: The Prognostic Value of
2 Echocardiography. *Ann Intern Med* 1991;114:635.
- 3 25. Fernández- Hidalgo N, Almirante B, Tornos P, González- Alujas MT, Planes AM,
4 Larrosa MN, Sambola A, Igual A, Pahissa A. Prognosis of left- sided infective
5 endocarditis in patients transferred to a tertiary- care hospital—prospective analysis of
6 referral bias and influence of inadequate antimicrobial treatment. *Clin Microbiol Infect*
7 2011;17:769–775.
- 8 26. Habib G, (France), Badano L, (Italy), Tribouilloy C, (France), Vilacosta I, (Spain),
9 Zamorano JL, (Spain), Scientific Committee:, Galderisi M, (Italy), Voigt J-U, (Belgium),
10 Sicari R, (Italy), Document Reviewers:, Cosyns B, (Belgium), Fox K, (UK), Aakhus S,
11 (Norway), On behalf of the European Association of Echocardiography.
12 Recommendations for the practice of echocardiography in infective endocarditis. *Eur J*
13 *Echocardiogr* 2010;11:202–219.
- 14 27. Chu VH, Park LP, Athan E, Delahaye F, Freiburger T, Lamas C, Miro JM, Mudrick DW,
15 Strahilevitz J, Tribouilloy C, Durante-Mangoni E, Pericas JM, Fernández-Hidalgo N,
16 Nacinovich F, Rizk H, Krajcinovic V, Giannitsioti E, Hurley JP, Hannan MM, Wang A.
17 Association Between Surgical Indications, Operative Risk, and Clinical Outcome in
18 Infective Endocarditis: A Prospective Study From the International Collaboration on
19 Endocarditis. *Circulation* 2015;131:131–140.
- 20 28. López-Dupla M, Hernández S, Olona M, Mercé J, Lorenzo A, Tapiol J, Gómez F,
21 Santamaría J, García R, Auguet T, Richart C, Castells E, Bardají A, Vidal F. [Clinical
22 characteristics and outcome of infective endocarditis in individuals of the general
23 population managed at a teaching hospital without cardiac surgery facilities. Study of 120
24 cases]. *Rev Esp Cardiol* 2006;59:1131–1139.

- 1 29. Oliver L, Lavoute C, Giorgi R, Salaun E, Hubert S, Casalta J-P, Gouriet F, Renard S, Saby
2 L, Avierinos J-F, Maysou L-A, Riberi A, Grisoli D, Casalta A-C, Collart F, Raoult D,
3 Habib G. Infective endocarditis in octogenarians. *Heart* 2017;103:1602–1609.
- 4 30. Ramírez-Duque N, García-Cabrera E, Ivanova-Georgieva R, Noureddine M, Lomas JM,
5 Hidalgo-Tenorio C, Plata A, Gálvez-Acebal J, Ruíz-Morales J, Torre-Lima J de la,
6 Reguera JM, Martínez-Marcos FJ, Alarcón A de. Surgical treatment for infective
7 endocarditis in elderly patients. *J Infect* 2011;63:131–138.

8

9

10

11 **Figure legend**

12

13 **Figure 1 : Probability of survival according to health-care pathway**

14 A: Comparison of survival between the three groups

15 B: Comparison of survival between Referral Centers (RC) and Transferred to Referral Centers
16 (TRC) groups

17 C: Comparison of survival between RC and Non-Referral Centers (NRC) groups

18 D: Comparison of survival between groups RC and TRC together and NRC group

19

20

21

22

23

24

25

1 **Investigators and sites**

2 **La Timone Hospital Marseille:** Florent Arregle, MD, Mary Philip, MD, SanRedrine Hubert,
3 MD, Frédérique Gouriet, MD PhD, Jean-Paul Casalta, MD, Frédéric Collart, MD, Alberto
4 Riberi, MD, Hélène Martel, MD, Sébastien Renard, MD, Laurence Camoin ,MD, PhD, Anne
5 Claire Casalta , MD, Hubert Lepidi ,MD, PhD, Michel Drancourt , MD, PhD, Didier Raoult
6 ,MD, PhD, Gilbert Habib MD, PhD. **CHU Nord Marseille:** Franck Thuny. MD, Yves
7 Frances. MD, Karim Aissi. MD, Philippe Brouqui. MD, Matthieu Million. MD, **CHU de**
8 **Montpellier** Catherine Sportouch. MD, Stéphane Cade. MD, Frédéric Cransac, MD. **CHU de**
9 **Nîmes:** Patrick Messner-Pellenc. MD, Camille Soulier, MD, Jean-etienne Ricci, MD,
10 **Hôpital d’Instruction des Armées Laveran,** Marseille: Laurent Fourcade, MD, Pierre-
11 Laurent Massoure, MD, Jean-Marie Gil. MD, **Hôpital Ambroise Paré-Desbief, Marseille:**
12 Emmanuel Philip. MD, **Fondation Saint Joseph, Marseille:** Patrick Khanoyan. MD, Roger
13 Rosario, MD, Nicolas Michel, MD, **CH de Martigues:** Serge Yvorra. MD, Virginie
14 Chellini. MD, **CH d’Avignon:** Stéphanie Branger. MD, Stéphane Andrieux, MD, Saida
15 Cheggour. MD, Salah Aboukoudir, MD. **Clinique Rhône-Durance, Avignon.** Laurent
16 Meille. MD, Catherine Meulemans. MD, **CH Toulon:** Jean-Michel Tartière. MD, Danielle
17 Raufast. MD, Isabelle LeCardonnel. MD, **Hôpital d’Instruction des Armées Saint-Anne,**
18 **Toulon:** Christophe Jego. MD, Gilles Cellarier. MD, Raphael Poyet. MD, **Polyclinique Les**
19 **Fleurs, Ollioules:** Philippe Pietri. MD, Philippe Villain. MD, Jean-Paul Giorgi. MD,
20 Chirine Parsai, MD, Olivier Brissy. MD, **CH Aix-en-Provence:** Bernard Jouve. MD, Tewfik
21 Benchaa. MD, **CH d’Arles:** Salah Si Ahmed MD, François Saint-Pierre. MD, **CH de**
22 **Draguignan:** André-François Chaix. Pierrick Boyer. MD, **Clinique La Casamance,**
23 **Aubagne** Véronique Haddad. MD, **CH des Alpes du Sud, Gap-Sisteron:** Christian Tafani.
24 MD, Amar Hidoud, MD, Audrey Boudes , MD, Hakim Berguigua, MD, Fabien Devemy,
25 MD, Bernard d’Hautefeuille, MD, **CH de Manosque** Alain Rapuzzi. MD, **Clinique des**

- 1 **Franciscaines, Nîmes:** Vlad Ciobataru. MD, **Hôpital de Fréjus:** Xavier Lamit. MD,
- 2 **Hôpital de Bastia:** Aurelia Tho, MD, Thierry James, MD, Abdelkader Bensalah. MD.
- 3
- 4

Table 1: Characteristics of patients according to their health-care pathway

	Total (n=342)	RC group (n=119)	TRC group (n=111)	NRC group (n=112)	p
Male	236 (69)	82 (69)	83 (75)	71 (63)	0.19
Age (mean \pm SD, years)	68 +/- 15	67 +/- 14	66 +/-15	73 +/- 14	<0.001
Medical history					
History of IE	42 (12)	17 (14)	16 (14)	9 (8)	0.25
Intravenous drug abuse	16 (5)	5 (4)	9 (8)	2 (2)	0.08
Cancer	59 (17)	12 (10)	21 (19)	26 (23)	0.03
Renal failure	50 (15)	18 (15)	16 (14)	16 (14)	0.98
Diabetes	85 (25)	28 (24)	31 (28)	26 (23)	0.66
Hypertension	147 (43)	47 (40)	53 (48)	47 (40)	0.38
Atrial fibrillation	116 (34)	48 (40)	33 (30)	35 (31)	0.19
History of valvular surgery	114 (33)	53 (45)	41 (37)	20 (18)	<0.001
Charlson index (mean \pm SD)	3.9 +/- 2	3.4 +/- 2	3.4 +/- 2	4.9 +/- 2	<0.001
Euroscore II (mean \pm SD, %)	9.3 +/- 11	7.2 +/- 6	7.4 +/- 8	13.6 +/- 15	0.004
Treatment before episode					
Oral anticoagulant	126/233 (54)	52/78 (67)	38/72 (53)	36/83 (43)	0.01
VKA	93/227 (40)	40/75 (53)	30/71 (42)	23/81 (28)	0.006
Aspirin	77/227 (34)	23/75 (31)	21/71 (30)	33/81 (41)	0.27
Statin	87/227 (38)	25/75 (33)	33/71 (46)	29/81 (36)	0.23
Source of infection					
Community	284 (83)	97 (82)	97 (87)	90 (80)	
HC related - nosocomial	54 (16)	22 (18)	12 (11)	20 (18)	0.21
HC related - non nosocomial	4 (1)	0	2 (2)	2 (2)	
Biology					
Hemoglobin (mean \pm SD, g/L)	109 +/- 22	111 +/- 20	108 +/- 20	109 +/- 18	0.39
Leukocytes (mean \pm SD, G/L)	11 +/- 5	10 +/- 4	11 +/- 4	11 +/- 6	0.28
Platelets (mean \pm SD, G/L)	240 +/- 132	243 +/- 123	226 +/- 124	254 +/- 148	0.22
CRP (mean \pm SD, mg/L)	106 +/- 99	91 +/- 95	103 +/- 84	127 +/- 117	0.02
Serum cretinine (mean \pm SD, μ mol/L)	127 +/- 102	131 +/- 91	128 +/- 113	123 +/- 104	0.22
Positive rheumatoid factor	84 (25)	39 (33)	33 (30)	12 (11)	<0.001
Microbiology					
<i>Staphylococcus Aureus</i>	88 (26)	24 (20)	34 (31)	30 (27)	0.18
Coagulase-negative Staphylococci	24 (7)	13 (11)	4 (4)	7 (6)	0.89

Oral Streptococci	53 (15)	14 (12)	21 (19)	18 (16)	0.32
<i>Enterococcus faecalis</i>	53 (15)	17 (14)	19 (17)	17 (15)	0.83
<i>Streptococcus gallolyticus</i>	29 (8)	7 (6)	9 (8)	13 (12)	0.29
Gram negative bacilli	22 (6)	8 (7)	4 (4)	10 (9)	0.27
Others	24 (7)	10 (8)	9 (8)	5 (4)	0.43
Negative blood cultures	58 (17)	28 (24)	16 (14)	14 (13)	0.06
Clinical data					
Heart failure	95 (28)	33 (28)	40 (36)	22 (20)	0.02
Cardiogenic shock	13 (4)	12 (10)	1 (1)	0	<0.001
Septic shock	32 (9)	12 (10)	6 (5)	14 (13)	0.18
Systemic embolism	131 (38)	44 (37)	50 (45)	37 (33)	0.17
Silent embolism	67 (20)	31 (26)	25 (23)	11 (10)	0.01
Intracranial hemorrhage	24 (7)	8 (7)	8 (7)	8 (7)	0.99
Mycotic aneurysm	14 (4)	5 (4)	3 (3)	6 (5)	0.62
Spondylodiscitis	45 (13)	9 (8)	15 (14)	21 (19)	0.04
Echocardiographic data					
IE localization					
Prosthetic valve	113 (33)	53 (45)	40 (36)	20 (18)	<0.001
Aortic	148 (43)	54 (45)	56 (50)	38 (34)	0.04
Mitral	144 (42)	51 (43)	49 (44)	44 (39)	0.75
Tricuspid	23 (7)	11 (9)	8 (7)	4 (4)	0.22
Pulmonary	0	0	0	0	
Pacemaker / ICD	24 (7)	9 (8)	11 (10)	4 (4)	0.17
Vegetation	241 (70)	83 (70)	88 (79)	70 (63)	0.02
Vegetation length (mean, mm)	13 +/- 8	13 +/- 9	16 +/- 8	10 +/- 7	<0.001
Peri-valvular lesion	55 (16)	18 (15)	31 (28)	6 (5)	<0.001
Severe valvular regurgitation	129 (38)	59 (50)	50 (45)	20 (18)	<0.001
LVEF (mean \pm SD, %)	56 +/- 12	56 +/- 12	56 +/- 12	56 +/- 11	0.42
Surgery and outcome					
Indication for surgery / device extraction	200 (58)	81 (68)	91 (82)	28 (25)	<0.001
Surgery / device extraction	118 (35)	52 (44)	60 (55)	6 (5)	<0.001
Hospital mortality	44 (13)	13 (11)	9 (8)	22 (20)	0.03
One-year mortality	88 (26)	24 (20)	23 (21)	41 (37)	0.003
Recurrence at one year	13 (4)	8 (7)	2 (2)	3 (3)	0.16

Values are n(%). RC: Referral Center; TRC : Transferred to Referral Center; NRC: Non-Referral Center; HC: health-care. CRP: C-Reactive Protein. LVEF: left ventricular ejection fraction. ICD implantable cardioverter defibrillator

Table 2: Predictive factors of mortality in multivariate analysis

Multivariate analysis		
	HR 95 %-CI	<i>p</i> value
RC Group*	1	-
TRC Group	0.95 (0.53-1.69)	0.857
NRC Group	2.56 (1.44-4.55)	0.001
Age (per one year)	1.03 (1.01-1.05)	0.002
Heart failure - No*	1	-
Yes	2.4 (1.52-3.78)	<0.001
Peri-valvular lesion - No*	1	-
Yes	3.25 (1.86-5.68)	<0.001
LVEF (per one %)	0.97 (0.95-0.99)	0.001

*Reference category

RC : Referral Center ; TRC: Transferred to Referral Center; NRC: Non-Referral Center;


LVEF : Left Ventricular Ejection Fraction

Table 3: Implementation of European guidelines according to health-care pathway


	Total	RC group	TRC group	NRC group	<i>p</i>	<i>p</i> RC vs TRC	<i>p</i> RC vs NRC	<i>p</i> RC+TRC vs NRC
ESC guidelines implementation	178 (52)	67 (61)	65 (60)	46 (46)	0.04	0.91	0.03	0.04
TEE	313 (92)	115 (97)	107 (96)	91 (81)	<0.001	1	<0.001	<0.001
Surgery / device explantation	267 (78)	92 (77)	81 (73)	94 (84)	0.14	0.45	0.20	0.14
Antibiotic therapy	246 (72)	92 (84)	90 (87)	64 (65)	<0.001	0.66	0.001	<0.001

RC : Referral Center, TRC : Transferred to Referral Center; NRC: Non-Referral Center; TEE : Trans-Esophageal Echocardiography


A.




B.



C.



D.



Supplemental Data File (.doc, .tif, pdf, etc.)



Click here to access/download
Supplemental Data File (.doc, .tif, pdf, etc.)
Supplementary1.docx

