

Do neuroplasticity and genetic factors contribute to cognitive training in children?

Iris Menu, Qin He, Julie Victor, Gabriela Rezende, Lorna Le Stanc, Julie Vidal, Catherine Oppenheim, Edouard Duchesnay, Boris Chaumette, Olivier Houdé, et al.

▶ To cite this version:

Iris Menu, Qin He, Julie Victor, Gabriela Rezende, Lorna Le Stanc, et al.. Do neuroplasticity and genetic factors contribute to cognitive training in children?. OHBM (Organization for Human Brain Mapping) 2022 Annual Meeting, Jun 2022, Glasgow, United Kingdom. hal-04045235

HAL Id: hal-04045235

https://hal.science/hal-04045235

Submitted on 24 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Do neuroplasticity and genetic factors contribute to cognitive training in children?



Iris MENU¹, Qin HE², Julie VICTOR³, Gabriela REZENDE¹, Lorna LE STANC¹, Julie VIDAL¹, Catherine OPPENHEIM², Edouard DUCHESNAY³, Boris CHAUMETTE², Olivier HOUDÉ^{1,4}, Grégoire BORST^{1,4,*}, & Arnaud CACHIA^{1,2,*}



¹Université de Paris, LaPsyDÉ, CNRS, Paris, F-75005, ²Université de Paris, IPNP, INSERM, Paris, F-75005, ³Université Paris-Saclay, CEA, CNRS, Neurospin, Baobab, Gif-sur-Yvette, F-91191, ⁴Institut Universitaire de France, Paris, France

*Equal contribution

Contact: iris.menu@hotmail.fr / arnaud.cachia@u-paris.fr



Introduction

Inhibitory control (IC)

- important for academic achievement, physical and mental health (Diamond, 2013)
- can be improved with cognitive training programs (e.g., Diamond & Lee, 2011; Jaeggi et al., 2011)

Factors contribute to cognitive training gains

- basal cognitive level (Karbach & Unger, 2014)
- brain activity/anatomy (Baniqued et al., 2019; Chaddock-Heyman et al., 2020)
- genetic factors (Leckie et al., 2014)

Aim of this study: investigate the genetic, cerebral and cognitive factors contributing to the receptivity of IC training

Material & methods

Brain anatomy (pre- and post-training)

 Grey matter volumes in regions of interest (ROI) of the IC network

Cognition (pre- and post-training)

- color-word Stroop task (Stroop interference)
- stop signal task (stop signal reaction time, SSRT)

Genes (saliva sample, pre-training)

 Polygenic risk score (PRS) calculated with the summary statistics of UK Biobank GWAS (N=84,259) related to IC (Trail Making Test, TMT).

Multilevel Structural Equation Model (SEM):

Test simultaneously and globally a set of *a priori* defined hypothetical relationships, combining:

- 2 latent change score models (LCS; Kievit et al., 2018; McArdle, 2009) to investigate training-related changes at both cognitive and brain levels (in yellow)
- 2 mediation models to investigate whether the genetic effect on cognition was mediated by the grey matter volumebaseline (in blue) and change (in green)

Model selection for the ROI based on:

- specific IC-training effect
- good SEM fit indices
- significant training-related cognitive changes

Results

Raw grey matter volume (without spatial normalization)


- IC training: IC changes depend on genetic (PRS), brain (basal and changes of left ACC volume) and cognitive (baseline SSRT) features
- Control training: IC changes only depend on the cognitive (baseline SSRT) features

Normalized grey matter volumes (with spatial normalization to MNI)


- *IC training*: IC changes depend on **brain** (basal and changes of left Putamen volume) and **cognitive** (basal SSRT) features
- *Control training*: IC changes depend only explained on the **cognitive** (baseline SSRT) feature

111 children (9-10 years) 75 children (9-10 years) **RED GREEN** GM volume in 8 regions Training of interest: left and righ color-word Stroop at home on tablet ACC, left and right IFG, task: stop signal task: Stroop **RED GREEN** left and right **caudate**, <u>aMRI:</u> 5 sessions per week left and right **putamen** GM volume in 8 regions (in seconds) (1/day, 5d/w, 15 min) **BLUE** of interest: left and righ color-word Stroop 5 weeks = 25 sessions ACC, left and right IFG, left and right **caudate**, interference score post-training left and right **putamen** (in seconds) pre-training **Inhibitory Control** Active control training (IC) training (AC) 35 children 40 children whole genome sample: color-word Stroop task - trivia-like questions polygenic risk score - stop signal task

Without spatial normalization (left ACC)



With spatial normalization (left Putamen)



References: Baniqued, P. L., et al. (2018). *Frontiers in Aging Neuroscience*. Chaddock-Heyman, L., et al. (2020). *Frontiers in Human Neuroscience*. Diamond, A. (2013). *Annual Review of Psychology*. Diamond, A., et al. (2011). *Science*. Jaeggi, S. M., et al. (2011). *Proceedings of the National Academy of Sciences*. Karbach, J., et al. (2014). *Frontiers in Psychology*. Kievit, R. A., et al. (2018). *Developmental Cognitive Neuroscience*. Leckie, R. L., et al. (2014). *Frontiers in Human Neuroscience*. McArdle, J. J. (2009). *Annual Review of Psychology*.

→ Training-relative cognitive changes depends on the interrelation of factors at different levels: molecular (genetics), neural (basal and change of reginal grey matter volumes) and cognition (basal level)

