



HAL
open science

Conjectures analogous to the Collatz conjecture

Fabio Briscese, Francesco Calogero

► **To cite this version:**

Fabio Briscese, Francesco Calogero. Conjectures analogous to the Collatz conjecture. 2023. hal-04046113v3

HAL Id: hal-04046113

<https://hal.science/hal-04046113v3>

Preprint submitted on 15 May 2023 (v3), last revised 22 Jan 2024 (v6)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conjectures analogous to the Collatz conjecture

Fabio Briscese^{a,b,*} and Francesco Calogero^{b,c,d,#}

^a Università Roma Tre, Dipartimento di Architettura, Rome, Italy

^b Istituto Nazionale di Alta Matematica Francesco Severi, Gruppo Nazionale di Fisica Matematica, Italy

^c Physics Department, University of Rome "La Sapienza", Rome, Italy

^d Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, Rome, Italy

* fabio.briscese@uniroma3.it, briscese.phys@gmail.com

francesco.calogero@uniroma1.it, francesco.calogero@roma1.infn.it

This paper is dedicated to the memory of Mario Bruschi

(08.08.1948-01.07.2021)

Abstract

In this paper we introduce some conjectures analogous to the well-known Collatz conjecture.

1 Introduction

The Collatz conjecture (also called " $3n + 1$ conjecture", or with various other names) [1] [2] is, so far, neither *proven*, nor shown to be *unprovable*. It is considered "one of the most famous unsolved problems in mathematics" [1]. In this paper we introduce some analogous conjectures; and mention some related ideas and conjectures.

Mario Bruschi (MB) was a pupil (almost 13 years younger), and then friend, colleague and, over time, co-author of several scientific papers with the second author of this paper (FC); he was a very remarkable man, with many interests in life besides science; he also authored many scientific papers (as single author or with various colleagues); 2 of these papers were related to the Collatz conjecture [3, 4], and it was him who first brought the attention of FC to this topic, and who made the numerical calculations reported in a joint paper never published (the last typed copy of this paper is dated February 4th, 2019, and we now put it on arXiv [5]).

2 The Collatz conjecture

Consider the sequence of *positive integers* n_j ($j = 0, 1, 2, \dots$) generated by the following 2 simple rules (hereafter we assume these rules to be applied *sequentially*, so that they generate a *uniquely* defined sequence of *positive integers*): $n_{j+1} = n_j/2$ if n_j is *even*, $n_{j+1} = 3n_j + 1$ if n_j is *odd*. The Collatz conjectures

states that, for *every* positive integer *initial* value n_0 , this sequence ends up into the cycle $\{1, 4, 2, 1\}$. This conjecture is attributed to Lothar Collatz (1910-1990), who perhaps first thought of it in 1937 [2]; it has been checked (in 2020) for *all* positive integers numbers n_0 up to $2^{68} \approx 2.95 \cdot 10^{20}$, and some computer experts continue to labor in order to increase this upper bound [1]; it is *obviously true* for certain *arbitrarily large* initial values (for instance for $n_0 = 2^N$ with N any positive integer); but it has not yet been *proven* nor, of course, *disproved*, nor shown to be *unprovable*; recently it has been shown that it is *extremely probable* that this conjecture is *true*, in an appropriate mathematical framework where this statement has a *precise* meaning [6]. The reason why it might *not* be true is, of course, because there might exist an *initial* positive integer n_0 such that the sequence generated by the simple rules detailed above either *diverges* or leads to a *different* cycle than the *single* one mentioned above.

Many analogous conjectures have been proposed over time (see for instance the book [7]). In this paper we mention some, possibly new, analogous *conjectures*.

3 The "5n + 1 Conjecture"

Consider the sequence of *positive integers* n_j ($j = 0, 1, 2, \dots$) generated by the following 3 simple rules: $n_{j+1} = n_j/2$ if n_j is *even*, $n_{j+1} = n_j/3$ if n_j is *odd and divisible by 3*, $n_{j+1} = 5n_j + 1$ if n_j is *odd and not divisible by 3*. The "5n + 1 Conjecture" states that, for *every* positive integer *initial* value n_0 , this sequence ends up into the cycle $\{1, 6, 3, 1\}$. This conjecture is not new; we thought so when we wrote the previous version (<https://hal.science/hal-04046113>) of this paper; it is in fact sometimes called the *Takutani conjecture*, see for instance the Chapter entitled *Empirical verification of the 3x + 1 and related conjectures* by Tomás Oliveira e Silva (kindly made available to us by its author) in the book [7].

3.1 A modified version of the "5n + 1 Conjecture"

Note that, if we eliminate the *second* rule namely we do *not* divide n_j by 3 if n_j is divisible by 3, we find several cycles: (e.g. $\{3, 16, 8, 4, 2, 1, 6, 3\}$; $\{5, 26, 13, 66, 33, 166, 83, 416, 208, 104, 52, 26\}$; $\{13, 66, 33, 166, 83, 416, 208, 104, 52, 26, 13\}$); while starting from other numbers (e.g. 7, 11,...) the sequence *seems* to diverge.

4 The "7n + 1 Conjecture"

Consider the sequence of *positive integers* n_j ($j = 0, 1, 2, \dots$) generated by the following 4 simple rules: $n_{j+1} = n_j/2$ if n_j is *even*, $n_{j+1} = n_j/3$ if n_j is *odd*

and divisible by 3, $n_{j+1} = n_j/5$ if n_j is odd and divisible by 5, $n_{j+1} = 7n_j + 1$ if n_j is odd and not divisible by 3 and 5. The "7n + 1 Conjecture" states that, for every positive integer initial value n_0 , this sequence ends up into the cycle $\{1, 8, 4, 2, 1\}$. This conjecture is not new (we thought so when we wrote the previous version—see: <https://hal.science/hal-04046113>—of this paper); it is in fact sometimes also called the *Takutani conjecture*, see for instance again the Chapter by Tomás Oliveira e Silva in the book [7].

4.1 A modified version of the "7n + 1 Conjecture"

Consider the sequence of positive integers n_j ($j = 0, 1, 2, \dots$) generated by the following 3 simple rules: $n_{j+1} = n_j/2$ if n_j is even, $n_{j+1} = n_j/3$ if n_j is odd and divisible by 3, $n_{j+1} = 7n_j + 1$ if n_j is odd and not divisible by 3 and 5. Then, for every positive integer initial value n_0 , this sequence ends up into one of the two cycles $\{1, 8, 4, 2, 1\}$ and $\{19, 134, 67, 470, 235, 1646, 823, 5762, 2881, 20168, 10084, 5042, 2521, 17648, 8824, 4412, 2206, 1103, 7722, 3861, 1287, 429, 143, 1002, 501, 167, 1170, 585, 195, 65, 456, 228, 114, 57, 19\}$. We checked this statement (only) for all positive integers n_0 up to 10^9 . This is the first case we found, of a conjecture, somewhat analogous to the Collatz conjectures, which figures (only) 2 cycles.

5 The "11n + 1 Conjecture"

Following the same intuition that had led us to write the "5n + 1 Conjecture" and the "7n + 1 Conjecture", one might be tempted to propose many quite analogous conjectures (based on *primes*); even stating that, starting from any positive integer, one always reaches a unique cycle containing 1. However, the following observation disproves this naive idea (although some variations of this idea might merit further exploration; see below).

Indeed, consider the sequence of positive integer numbers n_j ($j = 0, 1, 2, \dots$) generated by the following 5 simple rules: $n_{j+1} = n_j/2$ if n_j is even, $n_{j+1} = n_j/3$ if n_j is odd and divisible by 3, $n_{j+1} = n_j/5$ if n_j is odd and divisible by 5, $n_{j+1} = n_j/7$ if n_j is odd and divisible by 7, $n_{j+1} = 11n_j + 1$ if n_j is odd and not divisible by 3, 5 and 7. The "11n + 1 Conjecture" states that, for every positive integer initial value n_0 , this sequence ends up into one of the two cycles $\{1, 12, 6, 3, 1\}$ and $\{17, 188, 94, 47, 518, 259, 37, 408, 204, 102, 51, 17\}$. We checked its validity (only) for all positive integers n_0 up to 10^9 .

5.1 The "Modified 11n + 1 Conjecture"

Surprisingly, if we modify the "11n + 1 Conjecture" by eliminating the *fourth* rule, we seem to only get 1 cycle. Indeed, consider the sequence of positive integer numbers n_j ($j = 0, 1, 2, \dots$) generated by the following 4 simple rules:

$n_{j+1} = n_j/2$ if n_j is even, $n_{j+1} = n_j/3$ if n_j is odd and divisible by 3, $n_{j+1} = n_j/5$ if n_j is odd and divisible by 5, $n_{j+1} = 11n_j + 1$ if n_j is odd and not divisible by 3 and 5. The "Modified $11n + 1$ Conjecture" states that, for every positive integer initial value n_0 , this sequence ends up into the cycle $\{1,12,6,3,1\}$. We checked this statement (only) for all positive integers n_0 up to 10^9 .

6 Final remarks

Note that, for the "Modified $7n + 1$ conjecture" and the " $11n + 1$ conjecture", one has 2 cycles. So—assuming these conjectures are indeed *true*—this entails the possibility to divide, in 2 different ways, all positive integers in 2 classes, identified by the property to generate—when used as initial value—a sequence that eventually ends up into one of these 2 cycles.

The results mentioned above clearly suggest the possibility to introduce many—indeed, presumably an *infinity*—of somewhat analogous conjectures to those described above. A very interesting open issue is whether the hypothesis that one of these conjecture be *true* (or, alternatively, *false*) might be shown to imply that another one of these conjectures is *as well true* (or, alternatively, *false*); and whether progress in this direction might eventually provide an avenue to finally *prove* or *disprove* the validity of some, or even of all, these conjectures.

Let us end this paper by recalling the simple idea [5], that if the Collatz conjecture is *true*, then the 3-times iterated Collatz procedure (as described in **Section 2**) implies that all positive integers can be divided into 3 distinct categories: those such that, when taken as initial values of the sequence generated by that procedure, entail that it converge eventually to 1, or to 2 or to 4; hence the related conjecture —relevant of course only if the Collatz conjecture is true— that these 3 categories of integers are equally populated, a statement meant to be true in the same sense as the assertion that there are as many even positive integers as odd positive integers [5]. Clearly analogous ideas and conjectures can be associated to all the conjectures formulated above.

References

- [1] See the item "Collatz conjecture" in Wikipedia, the References quoted there, and, if need be, the References quoted in those References (the full list would add up to several hundred items...).
- [2] J. C. Lagarias, "The $3n+1$ problem: an overview", arXiv:2111.02635v1 [math.NT] 4 Nov 2021.
- [3] M. Bruschi, "Two Cellular Automata for the $3x+1$ Map", arXiv:nlin/0502061v1 [nlin.CG] 26 Feb 2005.

- [4] M. Bruschi, "A generalization of the Collatz problem and conjecture", arXiv:0810.5169v1 [math.NT] 29 Oct 2008.
- [5] M. Bruschi and F. Calogero, "Variations on a Theme of Collatz", arXiv:2303.08141v1 [math.GM] 14 Mar 2023.
- [6] Terence Tao, "Almost all orbits of the Collatz map attain almost bounded values", arXiv:1909.03562v5 [math.PR] 15 Feb 2022.
- [7] J. C. Lagarias, *The Ultimate Challenge: The $3x + 1$ Problem*, American Mathematical Society 2010