



## Models based on best-available information support a low inbreeding load and potential for recovery in the vaquita

Christopher Kyriazis, Jacqueline Robinson, Sergio Nigenda-Morales, Annabel Beichman, Lorenzo Rojas-Bracho, Kelly Robertson, Michael C. Fontaine, Robert Wayne, Barbara Taylor, Kirk Lohmueller, et al.

### ► To cite this version:

Christopher Kyriazis, Jacqueline Robinson, Sergio Nigenda-Morales, Annabel Beichman, Lorenzo Rojas-Bracho, et al.. Models based on best-available information support a low inbreeding load and potential for recovery in the vaquita. *Heredity*, 2023, 130, pp.183-187. 10.1038/s41437-023-00608-7 . hal-04047612

**HAL Id: hal-04047612**

**<https://hal.science/hal-04047612>**

Submitted on 27 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Models based on best-available information support a low inbreeding load and potential for recovery in the vaquita**

**Authors:** Christopher C. Kyriazis<sup>1\*</sup>, Jacqueline A. Robinson<sup>2\*</sup>, Sergio F. Nigenda-Morales<sup>3</sup>, Annabel C. Beichman<sup>4</sup>, Lorenzo Rojas-Bracho<sup>5</sup>, Kelly M. Robertson<sup>6</sup>, Michael C. Fontaine<sup>7,8,9</sup>, Robert K. Wayne<sup>1#</sup>, Barbara L. Taylor<sup>6</sup>, Kirk E. Lohmueller<sup>1,10\*</sup>, and Phillip A. Morin<sup>6\*</sup>

**Affiliations:**

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of California, Los Angeles; Los Angeles, CA, USA.

<sup>2</sup>Institute for Human Genetics, University of California, San Francisco; San Francisco, CA, USA.

<sup>3</sup>Advanced Genomics Unit, National Laboratory of Genomics for Biodiversity (Langebio), Center for Research and Advanced Studies (Cinvestav); Irapuato, Guanajuato, Mexico.

<sup>4</sup>Department of Genome Sciences, University of Washington; Seattle, WA, USA.

<sup>5</sup>Ocean Wise, Vancouver, BC, Canada.

<sup>6</sup>Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA ; La Jolla, CA, USA.

<sup>7</sup>MIVEGEC, Université de Montpellier, CNRS, IRD; Montpellier, France.

<sup>8</sup>Centre de Recherche en Écologie et Évolution de la Santé (CREES); Montpellier, France.

<sup>9</sup>Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen; Groningen, The Netherlands.

<sup>10</sup>Department of Human Genetics, David Geffen School of Medicine, University of California, Los Angeles; Los Angeles, CA, USA.

<sup>#</sup>deceased 26 December 2022

\*Correspondence: ckyriazis@g.ucla.edu, jacqueline.robinson@ucsf.edu, klohmueLLer@g.ucla.edu, phillip.morin@noaa.gov

## Main text

In Robinson et al. (2022), we employed genomic and simulation analyses to demonstrate that the critically endangered vaquita porpoise has a reduced potential for future inbreeding depression (inbreeding load), and is therefore not doomed to extinction by deleterious genetic factors. Garcia-Dorado and Hedrick (2022) (hereafter, GD&H) critique our analysis for not sufficiently demonstrating that the vaquita has a low inbreeding load and a good chance of recovery in the absence of continued incidental mortality in fishing gillnets.

Our conclusion that the vaquita likely has a very low inbreeding load is supported, first and foremost, by the finding that there are very few deleterious mutations that can contribute to inbreeding depression segregating in the vaquita due to its small historical population size. There is widespread agreement that inbreeding depression is overwhelmingly due to the exposure of recessive deleterious mutations that are concealed as heterozygotes (Charlesworth and Willis 2009; Hedrick and Garcia-Dorado 2016). In the relative absence of such mutations, there is simply very little fuel for inbreeding depression. For example, our genomic analysis shows that vaquitas have ~17 heterozygous loss of function (LOF) mutations per individual, whereas the blue whale genome has ~248 of such mutations. Loss of function mutations can, in many cases, have severe effects on fitness because they disrupt the function of protein-coding genes (though see MacArthur and Tyler-Smith 2010). Under the assumption that such LOF mutations are (partially) recessive, a low count of heterozygous LOF mutations as observed in the vaquita implies a low inbreeding load, whereas a high count as observed in the blue whale implies a high inbreeding load.

GD&H are correct that we do not precisely know the selection ( $s$ ) or dominance ( $h$ ) coefficients for putatively deleterious mutations, such that our identification of few segregating deleterious mutations in vaquitas offers only qualitative insight into inbreeding load. This limitation motivated our complementary simulation analysis, where we employ a distribution of  $s$  for new mutations that was inferred from our genomic dataset. Although we agree that our model, like any model, makes assumptions that should be critically evaluated, our model is informed by the best-available information on selection and demographic parameters in the

vaquita. Below, we review some of the key components of the vaquita and our model that inform our conclusions.

### **What should be considered a “small” historical population size?**

GD&H assert that inbreeding load in the vaquita cannot be low since the species has a large historical population size. Although the historical population size of vaquita may seem large relative to the very small recent effective population sizes ( $N_e$ ) that are observed in endangered species, the historical size of the vaquita population is not large when considering the broader context of long-term  $N_e$  in mammals or other taxa. Long term  $N_e$  in mammals (defined here as  $N_e = \pi / (4\mu)$  where  $\pi$  is heterozygosity and  $\mu$  is the mutation rate) is typically on the order of tens to hundreds of thousands, and rarely below 5000. For example, using published estimates of genome-wide heterozygosity in 42 species of mammals, we recently estimated a median long-term  $N_e$  of 21,875 (Kyriazis et al. 2022). Based on our estimate of the mutation rate in vaquitas of  $\mu = 5.8e-9$  and estimate of  $\pi = 9.04e-5$  (Robinson et al. 2022), vaquitas are estimated to have a long-term  $N_e$  of 3,896, the second lowest of the 42 species in this dataset (Kyriazis et al. 2022). This finding is also supported by fitting more complex non-equilibrium demographic models to genomic datasets for the vaquita, which similarly estimate  $N_e < 5,000$  going back tens or hundreds of thousands of years (Morin et al. 2021; Robinson et al. 2022). The species with the smallest long-term  $N_e$  in this dataset, the Channel Island fox, has been previously shown to exhibit no phenotypic signs of inbreeding depression despite having experienced severe recent bottlenecks (Robinson et al. 2018). By contrast, North American gray wolves have a large long-term  $N_e$  of ~92,000 (assuming  $\mu = 4.5e-9$  (Koch et al. 2019) and  $\pi = 1.65e-3$  (Robinson et al. 2019)), which may help explain numerous observed instances of severe inbreeding depression in the species (Fredrickson et al. 2007; Räikkönen et al. 2009; Robinson et al. 2019). Thus, although the historical vaquita population size may seem large relative the current size of the population, it is still vastly smaller than the population sizes observed in most other species of mammals.

Although these long-term effective population sizes may not be representative of the current size of many threatened or endangered populations, they are essential for modelling

the inbreeding load in a species. This is because long-term demographic processes have a major impact on patterns of segregating (recessive) deleterious variation, the key determinant of the inbreeding load. Though recent declines can influence patterns of segregating variation and inbreeding load, these effects often take tens or hundreds of generations to manifest. For example, as we show in our analysis, the recent and dramatic decline in the vaquita over the past ~35 years or ~3 generations does not appear to have impacted genetic diversity or inbreeding load in the species (Robinson et al. 2022). This is likely the case for many other large mammals that have experienced declines over the past century, given the long generation times typical of these species. Thus, low inbreeding loads in species with long generation times are likely a product of small long-term historical population sizes, rather than recent human-mediated declines. However, the extent to which this is true for a given species will depend on the duration and severity of decline, something that can readily be assessed using simulations.

#### **What should be considered a “typical” inbreeding load?**

A useful approach for determining whether computational models of inbreeding depression are reasonable is to compare the predicted inbreeding load ( $B$ ) from such models to those that have been estimated from natural populations (note that we report values in terms of the diploid inbreeding load  $[2B]$  in Robinson et al. (2022) but report the haploid inbreeding load  $[B]$  here to be consistent with GD&H). GD&H cite an average estimate from O’Grady et al. (2006) of  $B = 6$ , derived from an analysis of 16 existing inbreeding load estimates, as being a typical inbreeding load for wild populations. Based on this result, they then claim that our model-based prediction of  $B = 0.48$  for the vaquita is unreasonably low. However, it has previously been shown that the inbreeding load estimate from O’Grady et al. (2006) is unreliable and upwardly biased, in part due to methodological issues associated with using generalized linear models with a logit link to estimate the inbreeding load (Nietlisbach et al. 2019). Specifically, Nietlisbach et al. (2019) found that the three highest inbreeding load values reported by O’Grady et al. (2006) are overestimates due to biased statistical models or issues with the original datasets (note that many of these same concerns apply to estimates reported in Hedrick and Garcia-Dorado (2016)). Moreover, the average estimate from O’Grady

et al. (2006) is also likely to be upwardly biased due to their approach of summing together inbreeding load estimates from different life stages, an approach that ignores widespread pleiotropy for mutations underlying fitness (Pickrell et al. 2016; Boyle et al. 2017). This approach can contribute to upward bias by potentially double or triple counting the effects of recessive deleterious mutations that contribute to inbreeding depression at different life stages (see Kyriazis et al. (2022) for further discussion).

Based on 22 estimates that are deemed to be reliable and unbiased, Nietlisbach et al. (2019) instead report a median inbreeding load for survival to sexual maturity in vertebrates of  $B = 2.25$ . Although our predicted inbreeding load of  $B = 0.48$  is somewhat lower than this median, this is expected given the small historical population size and low levels of segregating (recessive) deleterious variation in the vaquita. Importantly, we note that this estimate of  $B = 2.25$  is based on only 13 species, nearly all of which are birds, and should therefore be interpreted cautiously. Thus, obtaining additional high-quality estimates of the inbreeding load from wild populations represents a key area of future research that will enable better assessment of simulation models (see Kyriazis et al. 2022, for further discussion). Nevertheless, based on available evidence, our model predicts an inbreeding load that is consistent with reliable estimates from natural populations.

### **How should we estimate selection and mutation parameters?**

GD&H claim that our selection and dominance parameters are incorrect because they do not agree with estimates of selection and dominance parameters derived from experiments in *Drosophila* (e.g., Simmons and Crow 1977; Pérez-Pereira et al. 2021, 2022). However, such experimental estimates are well known to be biased towards strongly deleterious variation, as mutations with more mild effects cannot be observed in an experimental setting (Davies et al. 1999; Eyre-Walker and Keightley 2007). These experimental approaches are also limited due to issues of identifiability for selection and mutation parameters (Lynch et al. 1999; Eyre-Walker and Keightley 2007; Halligan and Keightley 2009). In other words, the distributions of fitness in the experimental populations can be explained by a high mutation rate and a low strength of selection ( $s$ ) or vice versa, making interpretation of these experimental results challenging.

143 Finally, experimental approaches are also only possible for laboratory organisms such as  
144 *Drosophila* or yeast, thus their relevance for understanding deleterious mutation parameters in  
145 natural populations of mammals such as the vaquita is unclear.

146 These limitations have motivated the increasing use of sequence-based estimates of the  
147 distribution of  $s$ , which leverage genetic variation datasets to estimate selection parameters  
148 while controlling for demography (Eyre-Walker and Keightley 2007). Such approaches are  
149 widely employed in population genetics (Eyre-Walker et al. 2006; Boyko et al. 2008; Ma et al.  
150 2013; Chen et al. 2017; Huber et al. 2017, 2018; Kim et al. 2017; Tataru et al. 2017; Huang et al.  
151 2021) and give a much more complete picture of the full spectrum of deleterious mutations,  
152 from strongly to mildly deleterious. Nevertheless, these approaches do have limitations in that  
153 they are not well suited for estimating the proportion of lethal mutations (Wade et al. 2022)  
154 and often assume additivity during inference (though see Huber et al. (2018)). Addressing these  
155 limitations is an area of ongoing research.

156 Given these limitations of sequence-based approaches, we explored a variety of  
157 dominance models in our analysis, as well as models with an additional proportion of lethal  
158 mutations (see Fig. S21 and Table S6 in Robinson et al. 2022). We also explored results when  
159 assuming a selection and dominance model proposed by Kardos et al. (2021) that is similar to  
160 that of Pérez-Pereira et al. (2022) (see Kyriazis et al. 2022, for a detailed comparison of these  
161 models). In all cases, we found that recovery was still the likely outcome when assuming a 90%  
162 reduction in bycatch mortality rates (Robinson et al. 2022). In fact, we observed much lower  
163 predicted extinction rates under the model proposed by Kardos et al. (2021), perhaps due to  
164 more efficient purging in models with a high fraction of lethal mutations (Robinson et al. 2022).  
165 In sum, models with a higher lethal mutation rate, like those favored by GD&H, also support our  
166 main conclusion that recovery is possible.

167 Finally, we note we did not consider the impact of non-coding deleterious mutations, as  
168 these mutations are generally inferred to be weakly deleterious ( $s$  on the order of  $1e-3$ ;  
169 (Torgerson et al. 2009; Murphy et al. 2021; Dukler et al. 2022)) and therefore not highly  
170 relevant to modelling inbreeding depression. Indeed, the analysis of Pérez-Pereira et al. (2021)  
171 supports this assumption by suggesting that such deleterious mutations, though relevant over

evolutionary timescales, are not particularly relevant in a conservation context. Future work should aim to refine estimates of the strength of selection against non-coding deleterious mutations and explore their potential impact on genetic load and extinction risk.

## **How can we validate simulation models?**

Although the above verbal arguments serve as justification for the simulation analysis we present in Robinson et al. (2022), several approaches can be employed to more quantitatively validate and compare selection and dominance models informed by experimental versus sequence-based studies. These include comparing proposed models in terms of (1) how well they agree with patterns of genetic variation in sequencing datasets and (2) how well they agree with reliable empirical estimates of the inbreeding load.

In Kyriazis et al. (2022), we undertook this task using humans as a focal species. Humans are useful for this exercise because there are extensive genetic variation datasets (Auton *et al.*, 2015), published demographic models (e.g., Gutenkunst et al. 2009; Gravel et al. 2011; Li and Durbin 2011; Tennessen et al. 2012), estimates of mutation rates and coding sequence length (Keightley 2012), estimates of the distribution of  $s$  (Eyre-Walker et al. 2006; Boyko et al. 2008; Kim et al. 2017), and estimates of the inbreeding load (Bittles and Neel 1994) and number of segregating recessive lethals per individual (Gao et al. 2015). Moreover, humans are much more closely related to the vaquita than *Drosophila* and have a long-term  $N_e$  that is typical for mammals ( $N_e \sim 17,000$ ). In Kyriazis et al. (2022), we leveraged these existing estimates of human demography, mutation rates, and coding sequence length to compare proposed selection and dominance models from Pérez-Pereira et al. (2022) and Kardos et al. (2021) to a model we previously presented in Kyriazis et al. (2021), which is similar to that of Robinson et al. (2022). In Kyriazis et al. (2022), we also propose a new model that better incorporates the impacts of recessive lethal mutations (Wade et al. 2022).

When comparing these various selection and dominance models, we found that sequence-based models, such as the model used in Robinson et al. (2022), make predictions that are consistent with empirical estimates of the inbreeding load in humans, whereas models based on experimental approaches do not (Kyriazis et al. 2022). For example, our model



proposed in Kyriazis et al. (2022) predicts an inbreeding load of  $B = 3.2$  and  $\sim 0.9$  recessive lethal mutations per diploid. These predictions are compatible with existing evidence in humans (note that Bittles and Neel (1994) estimate  $B = 0.7$  for humans, though this is likely to be an underestimate as it is based only on juvenile mortality). By contrast, the model proposed by Pérez-Pereira et al. (2022) predicts an inbreeding load of  $B = 14$ , vastly exceeding available estimates in humans. The Pérez-Pereira et al. (2022) model also predicts  $\sim 12$  recessive lethal mutations per individual, whereas available estimates are on the order of 0.6 mutations per diploid (Gao et al. 2015).

Comparing predicted patterns of genetic variation from these models to those observed in humans also provides support for sequence-based models. Specifically, the Pérez-Pereira et al. (2022) model predicts a large overabundance of rare mutations, with 72.8% of nonsynonymous mutations predicted to be singletons (variants with frequency  $1/2n$  in a sample). However, only 56.8% of such mutations are observed to be singletons in the European sample from the 1000 Genomes dataset (Auton et al. 2015; Kyriazis et al. 2022). This large excess of rare mutations predicted by Pérez-Pereira et al. (2022) is a consequence of the extreme strength of negative selection in the model, which results in deleterious mutations being kept at low frequency. Importantly, the model we propose in Kyriazis et al. (2022) predicts 57.3% of mutations to be singletons, in good agreement with observed patterns of genetic variation in humans (Auton et al. 2015).

Overall, these findings are consistent with previous findings that selection parameters derived from experimental studies in *Drosophila* and other taxa are biased towards strongly deleterious mutations (Davies et al. 1999; Eyre-Walker and Keightley 2007; Kyriazis et al. 2022). Thus, this analysis helps validate the use of sequence-based estimates of selection parameters, such as those employed in Robinson et al. (2022), given that they are consistent both with patterns of genetic variation and empirical estimates of the inbreeding load.

## **Other concerns**

GD&H critique our simulations for ignoring stochastic environmental and demographic factors and not modelling a loss of adaptive potential in the vaquita. However, our analysis

does incorporate demographic stochasticity, as we model survival and reproduction probabilistically. Although we do not model environmental stochasticity or loss of adaptive potential, the threat that these factors pose to the vaquita, if any, is entirely unknown. Moreover, we emphasize that the aim of our analysis was to test the assumption that the vaquita is doomed to extinction by inbreeding depression. We do not interpret our model projections beyond demonstrating the qualitative result that inbreeding depression alone does not impede recovery in the species, as we agree with the general view that population viability models should be interpreted cautiously in terms of their ability to accurately predict future population sizes (Beissinger and Westphal 1998). Though incorporating factors such as environmental stochasticity may influence our model predictions, they would not change our central conclusion that recovery remains possible.

## **Conclusions**

In conclusion, we agree with GD&H that predictive models should be critically evaluated. Indeed, the critical evaluation of proposed mutation and selection models that we present in Kyriazis et al. (2022) serves as justification for the type of sequence-based models we employed in Robinson et al. (2022). Moreover, this analysis demonstrates that models based on experimental results, such as those proposed by Pérez-Pereira et al. (2022) and Kardos et al. (2021), are biased towards strongly deleterious variation and are not consistent with patterns of genetic variation or empirical estimates of the inbreeding load in humans. Nevertheless, our analysis in Robinson et al. (2022) found that recovery was still the likely outcome when assuming models with a much higher fraction of strongly deleterious variation. However, we emphasize that all predictive models should be interpreted cautiously, given that there is often a fair amount of uncertainty in parameter estimates that can be challenging to validate with orthogonal sources of information (Beissinger and Westphal 1998). For instance, having field-based estimates of the inbreeding load for the vaquita would represent a valuable source of additional information for our study, however, such data do not currently exist, and may never exist given the perilous circumstances of the species. Future work should aim to validate the sorts of models we employ in Robinson et al. (2022) for wild species where field-based

estimates of the inbreeding load exist. Such work could greatly strengthen conclusions drawn from such predictive models.

All modelling considerations aside, our conclusion that recovery is possible is also supported by all field surveys since late 1990s, including those in 2019 and 2021, where active healthy vaquitas with calves have been sighted (Rojas-Bracho et al. 2022). Despite an almost certain increase in gillnetting within the small range where vaquitas remain, vaquitas continue to survive at higher numbers than expected (Rojas-Bracho et al. 2022). The possibility of recovery is also supported by numerous examples of recovery for species that have dwindled below 20 individuals (Wiedenfeld et al. 2021), many of which were once thought to be doomed to extinction. Although inbreeding depression may in many contexts contribute to population decline, the naïve assumption that it will inevitably doom small populations is a dangerous view that is harmful to species conservation.

## **Acknowledgements**

We are grateful to three anonymous reviewers for constructive comments on this manuscript. C.C.K. and K.E.L. were supported by National Institutes of Health (NIH) grant R35GM119856 (to K.E.L.). A.C.B. was supported by the Biological Mechanisms of Healthy Aging Training Program, NIH T32AG066574. S.F.N.-M. was supported by the Mexican National Council for Science and Technology (CONACYT) postdoctoral fellowship 724094 and the Mexican Secretariat of Agriculture and Rural Development postdoctoral fellowship. We dedicate this work to our mentor, colleague, and friend Bob Wayne, who passed away in December 2022.

## **Author contributions**

C.C.K. wrote the manuscript with input from all coauthors.

## **Conflict of interest**

The authors declare no conflict.

## References

- Auton A, Abecasis GR, Altshuler DM, Durbin RM, Bentley DR, Chakravarti A et al. (2015). A global reference for human genetic variation. *Nature* **526**: 68–74.
- Beissinger SR, Westphal MI (1998). On the Use of Demographic Models of Population Viability in Endangered Species Management. *J Wildl Manage* **62**: 821–841.
- Bittles AH, Neel JV (1994). The costs of human inbreeding and their implications for variations at the DNA level. *Nat Genet* **8**: 117–121.
- Boyko AR, Williamson SH, Indap AR, Degenhardt JD, Hernandez RD, Lohmueller KE et al. (2008). Assessing the evolutionary impact of amino acid mutations in the human genome. *PLoS Genet* **4**.
- Boyle EA, Li YI, Pritchard JK (2017). An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169**: 1177–1186.
- Charlesworth D, Willis JH (2009). The genetics of inbreeding depression. *Nat Rev Genet* **10**: 783–796.
- Chen J, Glémin S, Lascoux M (2017). Genetic diversity and the efficacy of purifying selection across plant and animal species. *Mol Biol Evol* **34**: 1417–1428.
- Davies EK, Peters AD, Keightley PD (1999). High frequency of cryptic deleterious mutations in *Caenorhabditis elegans*. *Science (80- )* **285**: 1748–1751.
- Dukler N, Mughal MR, Ramani R, Huang Y-F, Siepel A (2022). Extreme purifying selection against point mutations in the human genome. *Nat Commun* **13**: 1–12.
- Eyre-Walker A, Keightley PD (2007). The distribution of fitness effects of new mutations. *Nat Rev Genet* **8**: 610–618.
- Eyre-Walker A, Woolfit M, Phelps T (2006). The Distribution of Fitness Effects of New Deleterious Amino Acid Mutations in Humans. *Genetics* **173**: 891–900.
- Fredrickson RJ, Siminski P, Woolf M, Hedrick PW (2007). Genetic rescue and inbreeding depression in Mexican wolves. *Proc R Soc B Biol Sci* **274**: 2365–2371.
- Gao Z, Waggoner D, Stephens M, Ober C, Przeworski M (2015). An estimate of the average number of recessive lethal mutations carried by humans. *Genetics* **199**: 1243–1254.
- Garcia-Dorado A, Hedrick P (2022). Some hope and many concerns on the future of the vaquita. *Heredity (Edinb)*.
- Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG et al. (2011). Demographic history and rare allele sharing among human populations. *Proc Natl Acad Sci* **108**: 11983–11988.
- Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009). Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLoS Genet* **5**: 1–11.
- Halligan DL, Keightley PD (2009). Spontaneous mutation accumulation studies in evolutionary genetics. *Annu Rev Ecol Evol Syst* **40**: 151–172.
- Hedrick PW, Garcia-Dorado A (2016). Understanding Inbreeding Depression, Purging, and Genetic Rescue. *Trends Ecol Evol* **31**: 940–952.
- Huang X, Fortier AL, Coffman AJ, Struck TJ, Irby MN, James JE et al. (2021). Inferring Genome-Wide Correlations of Mutation Fitness Effects between Populations. *Mol Biol Evol* **38**: 4588–4602.
- Huber CD, Durvasula A, Hancock AM (2018). Gene expression drives the evolution of

dominance. *Nat Commun* **9**: 1–11.

Huber CD, Kim BY, Marsden CD, Lohmueller KE (2017). Determining the factors driving selective effects of new nonsynonymous mutations. *Proc Natl Acad Sci* **114**: 4465–4470.

Kardos M, Armstrong E, Fitzpatrick S, Hauser S, Hedrick P, Miller J et al. (2021). The crucial role of genome-wide genetic variation in conservation. *Proc Natl Acad Sci* **118**: 2021.07.05.451163.

Keightley PD (2012). Rates and fitness consequences of new mutations in humans. *Genetics* **190**: 295–304.

Kim BY, Huber CD, Lohmueller KE (2017). Inference of the Distribution of Selection Coefficients for New Nonsynonymous Mutations Using Large Samples. *Genetics* **206**: 345–361.

Koch EM, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, Wayne RK et al. (2019). De Novo Mutation Rate Estimation in Wolves of Known Pedigree. *Mol Biol Evol* **36**: 2536–2547.

Kyriazis CC, Robinson JA, Lohmueller KE (2022). Using computational simulations to quantify genetic load and predict extinction risk. *bioRxiv*: 1–29.

Kyriazis CC, Wayne RK, Lohmueller KE (2021). Strongly deleterious mutations are a primary determinant of extinction risk due to inbreeding depression. *Evol Lett* **5**: 33–47.

Li H, Durbin R (2011). Inference of human population history from individual whole-genome sequences. *Nature* **475**: 493–496.

Lynch M, Blanchard J, Houle D, Kibota T, Schultz S, Vassilieva L et al. (1999). Perspective: Spontaneous deleterious mutation. *Evolution (N Y)* **53**: 645–663.

Ma X, Kelley JL, Eilertson K, Musharoff S, Degenhardt JD, Martins AL et al. (2013). Population Genomic Analysis Reveals a Rich Speciation and Demographic History of Orang-utans (*Pongo pygmaeus* and *Pongo abelii*). *PLoS One* **8**.

MacArthur DG, Tyler-Smith C (2010). Loss-of-function variants in the genomes of healthy humans. *Hum Mol Genet* **19**: 125–130.

Morin PA, Archer FI, Avila CD, Balacco JR, Bukhman YV, Chow W et al. (2021). Reference genome and demographic history of the most endangered marine mammal, the vaquita. *Mol Ecol Resour* **21**: 1008–1020.

Murphy D, Elyashiv E, Amster G, Sella G (2021). Broad-scale variation in human genetic diversity levels is predicted by purifying selection on coding and non-coding elements. *bioRxiv*: 1–18.

Nietlisbach P, Muff S, Reid JM, Whitlock MC, Keller LF (2019). Nonequivalent lethal equivalents: Models and inbreeding metrics for unbiased estimation of inbreeding load. *Evol Appl* **12**: 266–279.

O’Grady JJ, Brook BW, Reed DH, Ballou JD, Tonkyn DW, Frankham R (2006). Realistic levels of inbreeding depression strongly affect extinction risk in wild populations. *Biol Conserv* **133**: 42–51.

Pérez-Pereira N, Caballero A, García-Dorado A (2022). Reviewing the consequences of genetic purging on the success of rescue programs. *Conserv Genet* **23**: 1–17.

Pérez-Pereira N, Pouso R, Rus A, Vilas A, López-Cortegano E, García-Dorado A et al. (2021). Long-term exhaustion of the inbreeding load in *Drosophila melanogaster*. *Heredity (Edinb)*: 1–11.

Pickrell JK, Berisa T, Liu JZ, Séguérel L, Tung JY, Hinds DA (2016). Detection and interpretation of shared genetic influences on 42 human traits. *Nat Genet* **48**: 709–717.

- Räikkönen J, Vucetich JA, Peterson RO, Nelson MP (2009). Congenital bone deformities and the inbred wolves ( *Canis lupus* ) of Isle Royale. *Biol Conserv* **142**: 1025–1031.
- Robinson JA, Brown C, Kim BY, Lohmueller KE, Wayne RK (2018). Purging of Strongly Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding Depression in Island Foxes. *Curr Biol* **28**: 3487-3494.e4.
- Robinson JA, Kyriazis CC, Nigenda-Morales SF, Beichman AC, Rojas-Bracho L, Robertson KM et al. (2022). The critically endangered vaquita is not doomed to extinction by inbreeding depression. *Science (80- )* **639**: 635–639.
- Robinson JA, Räikkönen J, Vucetich LM, Vucetich JA, Peterson RO, Lohmueller KE et al. (2019). Genomic signatures of extensive inbreeding in Isle Royale wolves, a population on the threshold of extinction. *Sci Adv* **5**: 1–13.
- Rojas-Bracho L, Taylor BL, Booth C, Thomas L, Jaramillo-Legorreta AM, Nieto-Garcia E et al. (2022). More vaquita porpoises survive than expected. *Endanger Species Res* **48**: 225–234.
- Simmons MJ, Crow JF (1977). Mutations affecting fitness in *Drosophila* populations. *Ann Rev Genet* **11**: 49–78.
- Tataru P, Mollion M, Glémin S, Bataillon T (2017). Inference of Distribution of Fitness Effects and Proportion of Adaptive Substitutions from Polymorphism Data. *Genetics* **207**: 1103–1119.
- Tennessen JA, Bigham AW, O'Connor TD, Fu W, Kenny EE, Gravel S et al. (2012). Evolution and Functional Impact of Rare Coding Variation from Deep Sequencing of Human Exomes. *Sci (New York, NY)* **337**: 64–69.
- Torgerson DG, Boyko AR, Hernandez RD, Indap A, Hu X, White TJ et al. (2009). Evolutionary processes acting on candidate cis-regulatory regions in humans inferred from patterns of polymorphism and divergence. *PLoS Genet* **5**.
- Wade EE, Kyriazis CC, Cavassim MIA, Lohmueller KE (2022). Quantifying the fraction of new mutations that are recessive lethal. *bioRxiv*: 1–24.
- Wiedenfeld DA, Alberts AC, Angulo A, Bennett EL, Byers O, Contreras-MacBeath T et al. (2021). Conservation resource allocation, small population resiliency, and the fallacy of conservation triage. *Conserv Biol* **35**: 1388–1395.