



HAL
open science

Decoding Trans-Saccadic Prediction Error

Louise Catheryne Barne, Jonathan Giordano, Thérèse Collins, Andrea Desantis

► **To cite this version:**

Louise Catheryne Barne, Jonathan Giordano, Thérèse Collins, Andrea Desantis. Decoding Trans-Saccadic Prediction Error. *Journal of Neuroscience*, 2023, 43 (11), pp.1933 - 1939. 10.1523/jneurosci.0563-22.2022 . hal-04049674v2

HAL Id: hal-04049674

<https://hal.science/hal-04049674v2>

Submitted on 28 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Decoding Trans-Saccadic Prediction Error

✉ Louise Catheryne Barne,^{1,3} Jonathan Giordano,² ✉ Thérèse Collins,^{2,*} and Andrea Desantis^{1,2,3*}

¹Département Traitement de l'Information et Systèmes, Office National d'Études et de Recherches Aérospatiales, Salon-de-Provence 13661, France,

²Integrative Neuroscience and Cognition Center (Unité Mixte de Recherche 8002), Centre National de la Recherche Scientifique, Université Paris Cité, Paris 75006, France, and ³Institut de Neurosciences de la Timone (Unité Mixte de Recherche 7289), Centre National de la Recherche Scientifique, Aix-Marseille Université, Marseille 13005, France

We are constantly sampling our environment by moving our eyes, but our subjective experience of the world is stable and constant. Stimulus displacement during or shortly after a saccade often goes unnoticed, a phenomenon called the saccadic suppression of displacement. Although we fail to notice such displacements, our oculomotor system computes the prediction errors and adequately adjusts the gaze and future saccadic execution, a phenomenon known as saccadic adaptation. In the present study, we aimed to find a brain signature of the trans-saccadic prediction error that informs the motor system but not explicit perception. We asked participants (either sex) to report whether a visual target was displaced during a saccade while recording electroencephalography (EEG). Using multivariate pattern analysis, we were able to differentiate displacements from no displacements, even when participants failed to report the displacement. In other words, we found that trans-saccadic prediction error is represented in the EEG signal 100 ms after the displacement presentation, mainly in occipital and parieto-occipital channels, even in the absence of explicit perception of the displacement.

Key words: decoding; EEG; saccadic suppression of displacement; trans-saccadic error

Significance Statement

Stability in vision occurs even while performing saccades. One suggested mechanism for this counterintuitive visual phenomenon is that external displacement is suppressed during the retinal remapping caused by a saccade. Here, we shed light on the mechanisms of trans-saccadic stability by showing that displacement information is not entirely suppressed and specifically present in the early stages of visual processing. Such a signal is relevant and computed for oculomotor adjustment despite being neglected for perception.

Introduction

Seeing the world requires moving the eyes such that the high-acuity fovea can sample regions of interest that would otherwise remain in the periphery. The benefits of eye movements for vision are undeniable; however, they also pose some problems. Indeed, each eye movement displaces the retinal position of objects, raising the issue of how trans-saccadic perceptual stability is achieved.

Trans-saccadic perceptual stability is studied in the laboratory using gaze-contingent techniques. Participants are asked to report whether a visual target occupies the same spatial location before and after a saccade. They are surprisingly bad at this task,

systematically missing displacements up to one-third the amplitude of the saccade, a phenomenon called saccadic suppression of displacement (Bridgeman et al., 1975; Deubel et al., 1996). Such observations led scientists to propose that the visual system has a default hypothesis or prior belief about stability (Niemeier et al., 2007). It has been argued that this is efficient given that objects tend to not move during a saccade. Moreover, it prevents the system from mistaking changes in the retinal position of objects because of eye movements for movements in the outside world. Saccade landing positions are generally normally distributed around the visual target, and although some of this variability is centrally planned (Collins et al., 2009), noise arises at lower levels and cannot be predicted (van Beers, 2007). Such a distinction between predicted and unpredicted saccade landing errors is crucial. Indeed, when the saccade does not land on the target but this was predicted (i.e., represented in the efference copy of the saccade), no prediction error occurs. However, in the case of a target displacement, targeting error occurs, but this was not predicted.

Thus, after a saccade, there can be a prediction error between the predicted and actual retinal position of the target, which is not necessarily the same as the target eccentricity. Although it

Received Mar. 21, 2022; revised Nov. 14, 2022; accepted Nov. 17, 2022.

Author contributions: T.C. and A.D. designed research; A.D. performed research; L.C.B. and J.G. analyzed data; and L.C.B., T.C., and A.D. wrote the paper.

This work was supported by Agence Nationale de la Recherche Grant ANR-18-CE10-0001 to A.D. We thank Dr. Qing Yang for help with data collection.

*T.C. and A.D. share senior authorship.

The authors declare no competing financial interests.

Correspondence should be addressed to Louise Catheryne Barne at louisec.barne@gmail.com.

<https://doi.org/10.1523/JNEUROSCI.0563-22.2022>

Copyright © 2023 the authors

does not lead to the perception of object displacement, it does lead to saccadic adaptation (McLaughlin, 1967; Collins and Wallman, 2012). Saccadic adaptation is the adjustment of saccade amplitude as a function of previous prediction errors. Initially observed in patients recovering from unilateral extraocular muscle paresis, adaptation can be observed in healthy participants by displacing the target during the saccade and observing the evolution of saccade amplitude; backward displacements lead to amplitude-decreasing adaptation, whereas forward displacements lead to amplitude-increasing adaptation. Adaptation also occurs on a trial-by-trial basis; an undershoot on one trial tends to be followed by a saccade of greater amplitude on the next. However, when the targeting errors are too large, there is little or no consequence for the next saccade target (Robinson et al., 2003). Moreover, there tends to be a better correlation between targeting error on one trial and amplitude adjustment on the subsequent trial when observers report not seeing the displacement (Collins, 2014).

The targeting errors that do not inform perception do inform the motor system. The present study aimed to identify a brain signature of the prediction errors in the absence of explicit perception of a physical object displacement. Healthy human participants were asked to perform a saccade toward a visual target and to report whether it moved during the saccade. We recorded their brain activity via electroencephalography (EEG). We used multivariate pattern analysis to distinguish the brain activity linked to target displacement, a proxy for the prediction error, grounded on the assumption that displacements generate unexpected targeting errors. We performed these analyses separately when observers reported a target displacement and when they reported no displacement. Our primary interest was the no-displacement response trials as decoding brain activity linked to these trials permits identifying whether and when the brain represents the prediction error when explicit perception is lacking.

Materials and Methods

Participants

Nineteen volunteers (age 25.42 ± 5.05 years, 10 female) took part in the experiment and received 15 euros per hour. All participants had a normal or corrected-to-normal vision and were not told of the hypotheses under investigation. They all gave written informed consent before participating in the experiment. This study was conducted in agreement with the requirements of the Declaration of Helsinki (2004) and approved by the local ethics committee (Comité d'éthique de la recherche) of Université Paris Cité.

Apparatus

Stimuli were presented on a CRT Sony GDM-C520 (100 Hz refresh rate) with a resolution of 1024×768 pixels and 40 cm width. Stimulus presentation and data collection were performed using MATLAB (MathWorks) with the Psychophysics Toolbox and EyeLink Toolbox extensions (Brainard, 1997; Cornelissen et al., 2002). EEG data were recorded with 64 Ag/AgCl electrodes mounted on an elastic cap and amplified by an actiCHamp amplifier (Brain Products). The sampling rate of the EEG recording was set to 500 Hz. Electrodes were arranged according to the international 10–20 system. Two of the electrodes (TP9 and TP10) were used to record horizontal eye movements [left and right horizontal electro-oculogram (HEOG), respectively], and two other electrodes (FT9 and FT10) were placed on the mastoids (M1 and M2, respectively). The right mastoid (M2) was used as the online reference. Viewing was binocular, and movements of the right eye were monitored with an EyeLink 1000 (SR Research) at a 1000 Hz sampling rate. Head movements were restrained with a chin rest located 51 cm from the screen.

Main experiment procedure

To start the trial, we instructed the participants to fixate an open black point, 0.2° of visual angle (dva) in diameter, displayed at the center of a

gray screen (Fig. 1A). After 1000 ms of fixation, plus a random jitter between 10 and 40 ms, a target was presented. The target was a vertically oriented Gabor patch (1.7 dva in diameter, three cycles) displayed at an eccentricity of 8 dva either to the left or to the right of fixation and at the same vertical level as the fixation dot. When the fixation point disappeared (i.e., after a random interval selected between 360 and 390 ms from target presentation), participants had up to 1000 ms to perform a saccade toward the target. If they performed the eye movement before or 1000 ms after the go signal (fixation dot disappearance), the trial was canceled, an error message (too early or too late) appeared on the screen, and a new trial was added to the end of the trial list.

A saccade was considered to have been initiated when gaze direction moved 2° away from the center of fixation. Once a saccade was detected, the target disappeared. It reappeared 50 ms later, either at the same location (on 50% of the trials), or at some distance (d) to the left or right of the initial target location. The 50 ms blank period between the action and the post-saccadic target presentation was introduced to decrease the overlapping of saccade and target-displacement EEG responses. The target was displayed for an additional 600 ms and then disappeared. After it disappeared, participants had to respond whether they believed the target had moved or not from its initial position. They used the up arrow key to denote “Yes the target moved” and the down key for “No the target did not move.” Responses were reported with the index finger of the right hand. After their response, participants had 1 s to blink before a new trial began. A break was proposed every 32 trials during which the experimenter was able to perform an eye-tracker calibration if necessary.

The distance d was set individually using a staircase procedure (described below), which each participant completed before the main experiment at the beginning of the session. More specifically, the distance d was adjusted to lead to an average performance accuracy ranging between 50 and 60%.

A series of operational errors could lead to the trial being canceled and added to the end of the trial list. This was the case if participants' saccade failed to land within 4 dva from the initial position of the target (a radius of half of target horizontal eccentricity), if participants responded before the target disappeared, or if they blinked at any moment between the beginning of the trial and the disappearance of the post-saccadic target.

The experiment ended when participants had performed a total of 640 valid trials (i.e., without any of the operational errors described above) resulting from 160 repetitions of the combinations of two displacements (present, absent) and two saccade directions (left, right). The task was slightly shorter for the first two participants, who completed 512 trials (2 displacements (present, absent) * 2 saccade directions (left, right) * 128 trials). Only valid trials were analyzed.

Staircase procedure

Before the main experiment, participants ran a staircase procedure. The task was identical to the one described above, except that during the staircase, no EEG data were recorded, and participants were required to report whether the target moved to the left or the right of its initial location using the left or right arrow key, respectively. We did not include trials with no displacement, and the direction (left or right) was selected randomly. Three interleaved staircases controlled the amplitude of the left/right target displacement. Each staircase started with a different displacement amplitude, that is, 0.5, 1, and 2 dva. If the observer's response was correct (incorrect), the amplitude of the displacement (left or right) decreased (increased) in the subsequent trial. The magnitude of the displacement was controlled by an accelerated stochastic approximation algorithm (Kesten, 1958) set to converge at the displacement amplitude leading to an average performance accuracy ranging between 50 and 60% correct responses. Each staircase stopped once the convergence level was reached. The convergence values obtained with the three staircases were averaged and used in the main experiment trials when a target movement was displayed. The staircase procedure lasted ~20 min.

Behavioral Analysis

Trials were divided based on the combination of participants' judgment (yes, no) and target displacement (displacement, no displacement),




Figure 1. *A*, Time course of a single trial. The red dot represents the current eye position. After a go signal, participants performed a saccade toward a target placed either on the left or on the right of a central fixation point. At saccade onset the target disappeared and reappeared 50 ms after at the same place (no displacement, 50%) or at a distance d away from the initial position (displacement condition, 25% backward, 25% forward). Participants reported whether the target moved. *B*, Average distance thresholds d in degrees of visual angle measured in the staircase task. *C*, Average sensitivity d' to target displacements. *D*, Decision criterion. All averages were calculated across participants, and each point represents an individual observer (*B–D*). *E*, Individual distributions of landing errors and the box plot of median error across participants. Negative values represent undershoots and positive value overshoots. *F*, Relationship between post-saccadic landing error and adjustment in subsequent saccadic amplitude. Lines represent their linear relationship for each participant, displacement condition (Backward/No displacement/Forward) and response type (No/Yes). *G*, Median amplitude residuals of the linear relationship of saccadic adjustment and post-saccadic landing errors separated by displacement condition (Backward, No-displacement, Forward) across participants. Post-saccadic errors were not enough for explaining the saccadic adjustment as the actual adjustment is below/above the expected in the forward/backward conditions, respectively. This intuition can be extracted from *F* as well, where at no adjustment ($y = 0$), participants systematically had more positive errors (overshoots) in backward trials than in no-displacement trials, and more negative errors (undershoots) in forward trials. *H*, Distribution of the median absolute post-saccadic error (PSE) for hits, false alarms (FA), misses, and correct rejections (CR) across participants.

leading to four experimental conditions corresponding to the following four possible outcomes of signal detection theory: hits (observers reported that the target moved and the target had indeed moved), false alarms (observers reported that the target moved but the target had not moved), correct rejections (observers reported that the target did not move, and the target indeed had not moved), and misses (observers reported that the target did not move, but the target had moved). Participants' performance was quantified by calculating detection sensitivity d' and decision criterion c .

Saccadic landing errors were calculated as the difference between the eye position when the target reappeared (i.e., 50 ms after the onset of the saccade) and the initial position of the target. In addition, the sign of the error was flipped for the left saccade trials such that positive errors expressed overshooting saccades and negative errors undershooting saccades.

The post-saccadic error was calculated as the difference between the eye position when the target reappeared and the final target position. To assess saccadic adaptation, we computed the change in saccadic amplitude between successive trials. More specifically, the saccadic amplitude at trial t_1 was subtracted from the amplitude observed at trial t_0 . Then we fitted a generalized linear model (GLM), implemented in statsmodels library (version 0.10.1) in Python for each participant to predict the saccadic adjustment. The function took into account the following factors: the post-saccadic error, displacement (-1 for backward, 0 for no displacement, 1 for forward), response (0 for no displacement, 1 for displacement), and interactions. Saccade amplitude and post-saccadic error were expressed in pixels. The statistical significance of the factors was evaluated at the group level via a one-sample t test ($\mu = 0$, $\alpha = 0.05$).

EEG Analyses

Preprocessing. Preprocessing of continuous EEG activity was conducted in Python (version 3.6.9) by using the Python-MNE (version 0.18.2) toolbox (Gramfort et al., 2014). Data were rereferenced to the average of left and right mastoids and then filtered with a fourth-order Butterworth noncausal filter between 0.5 and 30 Hz. Data were segmented from -200 to 600 ms relative to saccade onset. EEG segments

were then corrected with a 200 ms baseline time locked to the first target presentation in the trial (i.e., from -200 to 0 ms relative to target onset).

Multivariate pattern analysis. EEG data were analyzed using Multivariate Pattern Analyses (MVPA) performed in Python using the scikit-learn library (version 0.22.1). Different models were applied depending on the analyzed signal (EEG or eye-tracking data).

We were particularly interested in evaluating whether we could classify displacement and no-displacement target trials from brain activity. One model aimed at classifying displacement and no-displacement trials when participants reported a target displacement (i.e., hit vs false alarm trials), and another model aimed at dissociating the same trials but when participants did not report experiencing a displacement (i.e., correct rejection vs miss trials). Importantly, to ensure that classification accuracy was not driven by eye movements occurring after saccade landing, MVPA was performed on all trials, excluding those that contained microsaccades occurring within a time window from 50 to 200 ms after the reappearance of the target (see below, Control analyses, Eye position for more details).

Trial-by-trial EEG classification was performed with an L2-regularized logistic classifier with a liblinear solver. Features consisted of the EEG activity recorded from all electrodes, except the references and HEOG electrodes. Feature scaling was performed as a preprocessing step, namely, data were normalized (z -score transformation) for each time point across trials, using the parameters (mean and SD) of the training set. Our classification procedure implemented a stratified fivefold cross-validation approach for each time point of the segmented data. Scores were the area under the curve (AUC), which we calculated for each fold and reported their average.

For each time point, we computed the weight vectors from each linear model to assess the electrodes that contributed the most to classification accuracy. Later, they were transformed into activation patterns as implemented in MNE to provide a neurophysiological interpretation to the coefficients, implying that electrodes with nonzero values carry class-specific information (Haufe et al., 2014).

Finally, to assess significant classification from chance (AUC = 0.5) at the participant level, we used cluster-based permutation tests based on paired t scores (Maris and Oostenveld, 2007), with 10,000 random

permutations, alpha value as 0.05, implemented in MNE by the function `permutation_cluster_1samp_test`. In addition, for the evaluation of activation maps, we performed permutation *t* tests on electrode data as implemented in MNE by the function `permutation_t_test`, with 10,000 random permutations and alpha as 0.0125 (alpha of 0.05 divided by four time-window tests).

Control Analyses

Eye position. Even in the absence of conscious perception of target displacement, volunteers could move their gaze toward the displaced target, thus potentially contaminating the EEG signal. Therefore, we removed (off-line) the trials containing (micro)saccades within a time window ranging from 50 to 200 ms after the main saccade detection (proportion of excluded trials = 0.044 ± 0.062). We selected this time window because it exhibited the largest EEG decoding accuracy (see below, Results for more details). Microsaccades were identified using the Microsaccade Toolbox 0.9 implemented in R software, considering the eye-tracking sampling rate (1000 Hz) and 6 ms as the minimum duration criteria and velocity threshold scaling factor as 5. The MVPA on EEG data was performed on this set of trials.

Furthermore, to ensure that EEG decoding accuracy was not merely driven by eye movements, we conducted classification analyses to dissociate displacement from no-displacement trials using gaze position data. Gaze position data corresponded to the online horizontal and vertical eye position, which was recorded throughout the experiment. Specifically, as for the EEG data, we performed the classification of the displacement separately for the responses yes the target moved (i.e., hit vs false alarms trials), and no the target did not move (i.e., correct rejection vs miss trials). We used a nonlinear classifier (C-Support Vector Classification with RBF (radial basis function) kernel and gamma as auto) because better classification boundaries are generated as the target could move to two different directions (Fig. 3B). Features consisted of the gaze position in the vertical axis and the absolute distance of the gaze from the center of the screen in the horizontal axis. This transformation in the *x*-axis enabled us to mirror the left hemifield and aggregate gaze position of the left and right trials. Data were normalized using *z*-score transformation, and classification was performed with a stratified fivefold cross-validation approach for each time point. Classification accuracy was summarized by the area under the receiver operating characteristic curve.

Target eccentricity. To evaluate a possible role of saccadic amplitudes and post-saccadic error (i.e., the distance between the new target position and gaze location after the saccade) on the EEG decoding, the same EEG-MVPA protocol described above was implemented with the exception that classifiers were trained and tested on the activity of parieto-occipital electrodes (O1, Oz, O₂, PO7, PO3, POz, PO4, PO8), and within a time window from 100 to 200 ms after the saccade. Instead of averaging the activity in time for each electrode, we used the signal at each time point as an attribute, summing up to 408 independent variables (8 electrodes \times 51 time points). We focused on parieto-occipital channels as these electrodes were more strongly responsible for EEG decoding accuracy (see below, Results). Then, for each participant and each response model, we fitted the predicted EEG classifier labels (one for displacement and zero for no displacement) with a binomial GLM that included saccadic amplitude, target eccentricity (aka, absolute post-saccadic error), and their interaction as predictors. The statistical significance of beta values was assessed at the group level, with a one-sample *t* test ($\mu = 0$, $\alpha = 0.05$) for each response model.

Behavioral relevance of EEG decoding. In a second step, we aimed to verify whether the prediction error signal present in parieto-occipital electrodes was a variable explaining the saccadic amplitude adjustment. Thus, we fitted mixed models and calculated Akaike's information criterion (AIC) and Bayesian information criterion (BIC) to compare their goodness of fit. The mixed-models and goodness-of-fit calculations were performed with the `statsmodels` library (version 0.10.1) in Python.

The first mixed-model (null model) considered the post-saccadic error and response in trial *t*₀ as fixed factors (with interactions) and participants as the random factor to model the amplitude adjustment in trial *t*₁. A second mixed model (full model) considered the post-saccadic error, the

actual target displacement (backward, forward, and no displacement), and response as fixed factors (with interactions). The factor participants was included as a random intercept. The third and fourth models were identical to the second, except that rather than using the actual target displacement to predict saccadic adaptation, we used the target displacement labels predicted by the EEG decoders described below.

The same EEG-MVPA method reported below in Multivariate pattern analysis was performed with two exceptions. First, classifiers were trained and tested on the activity of parieto-occipital electrodes (O1, Oz, O₂, PO7, PO3, POz, PO4, PO8) observed within a time window from 100 to 200 ms after the saccade, corresponding to 408 independent variables (8 electrodes \times 51 time points). Second, the one-versus-rest approach was selected to decode backward displacement, forward displacement, and no-displacement trials. For the third mixed model (true decoding model), the displacement labels used to predict saccadic adaptation were the good guesses of the true classifier, that is, the classifier exposed to the proper labels during training. Instead, the fourth mixed model (shuffled decoding model) used the displacement labels predicted by a classifier trained on shuffled labels. In other words, for the shuffled classifier, we removed the actual correspondence between brain activity and actual target displacement. The model predicting saccadic adaptation from the output of the shuffled decoder was compared with the true decoding model by calculating the AIC and BIC estimators to evaluate whether the good guesses of the true decoder could predict saccadic adaptation. The AIC and BIC criteria favor models with the closest probability distribution to the fitted data while controlling for overfitting issues. In fact, simply adding a random parameter to a model may increase its likelihood. However, AIC and BIC attribute penalties for additional parameters (the penalty term is larger in BIC than in AIC). Therefore, we expected the shuffled decoder to be the worst model.

Data availability

All data and analysis scripts are freely available online at <https://osf.io/evnhx/>. The procedures and analysis were predetermined but not pre-registered on a public repository.

Results

Behavioral results

The individual displacement distances ranged from 0.38 to 1.39 dva ($Q_{1/4} = 0.59$, $Q_{2/4} = 0.73$, $Q_{3/4} = 0.85$; Fig. 1B), and as expected by the experimental design, participants exhibited low sensitivity *d'* to target displacement (Fig. 1C). They also showed positive decision criterion values indicating a tendency toward a conservative response bias ($Q_{1/4} = -0.08$, $Q_{2/4} = 0.17$, $Q_{3/4} = 0.32$; Wilcoxon signed-rank test; $Z = 2.374$, $p = 0.018$; Fig. 1D). Additionally, the hit rates for forward (outward) and backward (inward) displacements were not significantly different (hit rate, mean \pm SD, for backward = 0.512 ± 0.237 and forward = 0.456 ± 0.185 ; $t_{(18)} = -0.048$, $p = 0.962$). Overall, participants performed the task near their perceptual threshold (proportion of correct responses, $Q_{1/4} = 0.54$, $Q_{2/4} = 0.56$, $Q_{3/4} = 0.58$).

Saccades ended 15 ms (median time) after the target disappearance, so in most of the trials there was no target when the participants' eyes landed. The analysis of saccade landing errors showed that saccades undershot with respect to the initial position of the target (Fig. 1E). Notwithstanding, participants adapted their saccadic performance on a trial-by-trial basis. Post-saccadic errors (pse) resulting from saccade undershoots led to a saccade with larger amplitude in the following trial, whereas saccade overshoots resulted in subsequently smaller saccadic amplitudes (β mean \pm SD = -0.865 ± 0.094 ; $t_{(18)} = -40.189$, $p < 0.001$; Fig. 1F). This replicates previous work, except that we did not find a modulation in saccadic adaptation by participants' response (response, $\beta = -0.168 \pm 3.181$, $t_{(18)} = -0.231$, $p = 0.82$; pse-response interaction, $\beta = 0.017 \pm 0.099$, $t_{(18)} = 0.734$, $p =$




Figure 2. Decoding displacements from the EEG signal. Left, Dotted vertical lines represent saccade detection (at 0 s) and target reappearance (at 0.05 s). Thick lines represent time points at which classification performance differs from chance ($AUC = 0.5$). Right, Topographic activation maps at four time windows, with the significant channels highlighted. The color map ranges from -10 (cool colors) to 10 (warm colors) arbitrary units.

0.472; response–displacement interaction, $\beta = 0.002 \pm 3.386$, $t_{(18)} = 0.003$, $p = 0.998$; pse–response–displacement interaction, $\beta = -0.014 \pm 0.136$, $t_{(18)} = -0.448$, $p = 0.66$). Saccadic adaptation also depended on the target displacement ($\beta = -14.577 \pm 6.767$, $t_{(18)} = -9.389$, $p < 0.001$; Fig. 1G). Notably, when participants were presented with a backward (forward) displacement, they tended to execute larger (smaller) saccades in the subsequent trial. However, the actual adjustment was often smaller than the adjustment one would expect from the forward/backward conditions (e.g., if in a given trial the target was displaced by 1 dva backward or forward, in the subsequent trial the adjustment was smaller than 1 dva).

Finally, we observed no interaction between post-saccadic errors and displacement (pse–displacement, $\beta = 0.031 \pm 0.087$, $t_{(18)} = 1.557$, $p = 0.137$); that is, the strength of saccadic adaptation was comparable across the different displacement conditions (backward, pse $\beta = -0.878 \pm 0.123$; no displacement, pse $\beta = -0.832 \pm 0.076$; forward, pse $\beta = -0.846 \pm 0.127$; paired t test, backward–forward, $t_{(18)} = -0.994$, $p = 0.334$; paired t test, backward–no, $t_{(18)} = -1.543$, $p = 0.14$; paired t test, forward–no, $t_{(18)} = -0.534$, $p = 0.6$).

Unseen post-saccadic displacement is decodable at early stages of visual processing

Despite low sensitivity to target displacements, participants nevertheless adapted their saccades based on the post-saccadic displacement and errors. Therefore, we investigated whether displacement information was present in the EEG signal. We were particularly interested in whether we could decode displacement information (i.e., the target moved vs the target did not move) in the cases in which participants replied no the target did not move as these unseen displacements still led to trial-by-trial adaptation. Although classification score was not very high ($AUC < 0.6$), we could decode target displacement above chance (Fig. 2) both when participants reported that the target moved (cluster-based permutation tests, from 166 to 212 ms, $p = 0.026$; from 298 to 348 ms, $p = 0.02$; from 362 to 416 ms, $p = 0.021$; from 498 to 552 ms, $p = 0.028$) and when they reported that it did not move (cluster-based permutation tests, no response, from 128 to 338 ms, $p < 0.001$). Classification scores along time were comparable for the

response models (cluster-based permutation test, $p = 0.938$ and $p = 0.482$).

To understand which electrodes were contributing the most to the significant classification from 100 to 200 ms, we analyzed the activation patterns generated by the linear classifiers at four time windows within this period (115–135, 140–160, 165–185, 190–210 ms). The electrodes that contributed the most to displacement classification were occipital and parieto-occipital electrodes (permutation tests, response yes, 115–135 ms = Oz, O₂, PO3, POz; 140–160 ms = Oz, POz; 190–210 ms = CP1, Pz, P3, P1, PO3, POz, PO4, P2, CPz; response no, 115–135 ms = O1, Oz, O₂, PO7; 140–160 ms = O1, Oz, PO7, PO3; 190–210 ms = POz). This suggests that information about post-saccadic displacement is represented in cortical visual areas during the early visual processing stages, even when this information will not subsequently lead to a perception of displacement.

Post-saccadic displacement classification during early processing stages (100–200 ms) cannot be explained by eye movements as we excluded trials with microsaccades within this window. Additionally, when we performed MVPA, but using the gaze position data, scores were only different from chance from 200 ms onward (Fig. 3A; cluster-based permutation tests, response no, $p < 0.001$; response yes, $p = 0.363$), suggesting that saccade landing positions were similar across conditions, but after 150 ms the target reappearance in displaced trials participants start to correct their gaze toward the target (Fig. 3B). All considered, this confirms that the early decoding of displacement is not because of eye movements.

Another alternative explanation is that the significant classification was not related to an error-related signal but because of differences in post-saccadic errors (i.e., the distance between the new target position and gaze location after the saccade). Indeed, by design, displacement trials generated bigger post-saccadic errors than no-displacement trials (Fig. 1H). One way of dealing with this confound would be to consider comparable parts of the distribution of errors between conditions, but of course this would create a new confound because those trials would no longer be comparable with regard to saccadic amplitude. For this reason, a series of binomial GLMs investigated whether error size and saccadic amplitude could predict the displacement labels resulting from EEG classifiers (see above, Control analyses). The rationale was to verify whether the size of the post-saccadic error and/or the saccadic amplitude were the variables explaining the label assignment. If this was the case, then we would expect, for example, that small post-saccadic errors would be consistently associated with no-displacement labels and bigger errors with displacement labels.

Additionally, linear mixed models assessed whether the target displacement labels predicted by the EEG classifiers could predict saccadic adaptation over and above post-saccadic errors. Together, these analyses would highlight whether such brain activity contained specific information about trans-saccadic prediction error that was not merely reflected by the size of post-saccadic errors. As indicated above, for these analyses, classifiers were trained and tested on the activity of parieto-occipital electrodes (i.e., the electrodes most responsible for decoding accuracy) during the time window from 100 to 200 ms after the saccade (i.e., a




Figure 3. *A*, Decoding displacements from the eye position data. Dashed lines and shaded areas as in Figure 2. *B*, Eye position data of participant 1 at 50 ms (target reappearance) and at 600 ms (right before the response go signal). The y-axis is the vertical position on the screen, whereas the x-axis is the distance from the center of the screen, the initial fixation. The plus signs represent the three possible positions of the target across trials (backward displacement, no-displacement, forward displacement). The gaze of participant 1 in all trials is represented by the dots. The dot color represents if there was a displacement (red) or not (blue) in each trial. As in *A*, for participant 1, there is a clear boundary for classifying the displacement from the eye position at the end of the trial (600 ms) but not at the beginning when the target just reappeared (50 ms), meaning that participant 1 generally performed eye movements toward the target at some point of the trial.

time window in which EEG signals were not contaminated by eye movements).

Displacement classification remained significant using the signal from parieto-occipital channels during early time points (response yes, $AUC = 0.571 \pm 0.064$, $t_{(18)} = 4.802$, $p < 0.001$; response no, $AUC = 0.574 \pm 0.093$, $t_{(18)} = 3.505$, $p = 0.003$). Critically, the size of post-saccadic errors and saccadic amplitude did not predict the displacement labels provided by the EEG classifier. This was the case both in trials in which participants responded the target moved (error size, $\beta = -0.024 \pm 0.08$, $t_{(18)} = -1.31$, $p = 0.207$; amplitude, $\beta = 0 \pm 0.018$, $t_{(18)} = -0.083$, $p = 0.935$; interaction, $\beta = 2.03e-04 \pm 4.45e-04$, $t_{(18)} = 1.957$, $p = 0.066$), and the target did not move (error size, $\beta = 0.024 \pm 0.14$, $t_{(18)} = 0.761$, $p = 0.457$; amplitude, $\beta = 0.002 \pm 0.021$, $t_{(18)} = 0.456$, $p = 0.654$; interaction, $\beta = -6.14e-05 \pm 7.75e-04$, $t_{(18)} = -0.346$, $p = 0.734$). Similar results are seen when performing this analysis but on the subset of trials that were correctly classified, with the exception that in this case the interaction of saccadic amplitude and error size was significant in response yes trials ($\beta = -0.001 \pm 0.003$, $t_{(18)} = 2.137$, $p = 0.047$). We can thus exclude that post-saccadic target eccentricity and saccadic amplitude were the information source for displacement EEG classification. We do not claim that occipital and parietal activity does not reflect visual post-saccadic target representation, but our results suggest that it also reflects a representation of prediction error.

The following analyses investigated whether the displacement labels resulting from the EEG classification predicted saccadic adaptation, consequently, representing prediction errors. We fitted four mixed models for the amplitude adjustment and compared their goodness of fit. The null model included post-saccadic errors and response type as predictors. The other three models nested the null behavioral model plus a displacement

Table 1. Goodness of fit measured by AIC and BIC for the estimated mixed models

Model	AIC	BIC
Full behavioral	99012.465	99085.850
True decoding	101140.834	101214.219
Null behavioral	101210.495	101254.525
Shuffled decoding	101199.043	101272.427

Smaller values correspond to better fits.

variable. In the full behavioral model the displacement variable consisted of the actual target displacement observed in the trials. In the true decoding model, this variable contained the displacement labels predicted by the classifiers trained with the actual target displacement. Finally, in the shuffled decoding model, the displacement variable consisted of the displacement labels predicted by the shuffled decoders (i.e., these classifiers were trained on a set of trials in which we shuffled the actual target displacement labels). Table 1 summarizes AIC and BIC values of the models and shows again by another method that the full behavioral model had the best fit. The second-best model was the true decoding model, with lower AIC and BIC values than the null and shuffled decoding model. Additionally, we ran likelihood ratio tests comparing each full model against the null behavioral model. Once more, the analyses showed that displacement variables significantly improved the fit [full behavioral model, $\chi^2(4) = 2206.03$, $p < 0.001$; true decoding model, $\chi^2(4) = 77.66$, $p < 0.001$; shuffled decoding, $\chi^2(4) = 19.452$, $p = 0.001$]. In sum, these analyses showed that the signal observed from 100 to 200 ms in parieto-occipital sites contains information that is not merely reflected by post-saccadic errors and that is relevant for predicting the future saccadic behavior.

Discussion

This study investigated the brain signatures of trans-saccadic prediction errors in the absence of the conscious perception of target displacement. Participants performed a saccade toward a visual target that disappeared when the saccade was initiated and reappeared 50 ms later either in the same or in a different location. We decoded the presence versus absence of displacement both when participants were aware of a displacement and when they were not. We argue that the signal driving EEG decoding accuracy is the prediction error between the expected and observed saccade landing position and that this signal is present at early stages of processing, even when it drives motor learning but does not inform perception.

Saccadic suppression of displacement (SSD) refers to the absence of awareness of trans-saccadic displacements (Bridgeman et al., 1975). Such blindness was taken as evidence that the visual system discards or ignores these displacements. However, two findings show that this blindness is not because of low sensitivity. First, performance is near perfect when the post-saccadic target reappears at its displaced location several hundred milliseconds after the saccade has landed (Deubel et al., 1996, 1998). Second, saccade amplitude changes trial by trial as a function of these unseen displacements, suggesting that although they are ignored by explicit perception, they are not ignored by the motor system, which continually adjusts to make up for what it considers to be its own errors (Collins, 2014). We replicated this trial-by-trial saccadic adaptation in the current dataset. We further showed that the motor system has access to information about target displacement at early stages during visual processing. Indeed, EEG classifiers could dissociate displacement and no-displacement

trials from ~100 ms after saccade onset. Moreover, occipital and parieto-occipital channels were the electrodes mostly responsible for classification accuracy.

These results are compatible with previous studies reporting an involvement of parietal areas in error signals related to saccades (Collins and Jacquet, 2018). Such error signals are thought to represent a discrepancy between the predicted landing position and the observed landing position. We believe that the post-saccadic displacement and error that we measured here reflect such an error signal. Neuropsychological studies reported deficits in saccade tasks compatible with the involvement of posterior parietal cortex (PPC) in saccade-related error signals (Duhamel et al., 1992; Gaymard et al., 1994; Heide and Kömpf, 1998; Pisella et al., 2011), but the often large extent of the lesions and the relatively small sample sizes make it difficult to pinpoint a more specific locus. Imaging studies also point toward the involvement of the PPC in saccade-related error signals: PPC has an eye-centered organization that is updated when the eyes move (Medendorp et al., 2003; Bellebaum et al., 2005) according to saccadic goals and not to the coordinates of the visual stimulus (Medendorp et al., 2005).

Furthermore, some PPC neurons show predictive remapping of their receptive fields; just before a saccade, receptive fields become responsive to stimuli in the future retinotopic location (Duhamel et al., 1992; Hall and Colby, 2011). This means that the PPC receives information about saccade error and updates retinotopic visual maps based on this signal. The expected oculomotor error is transmitted via efference copy and is available ~30 ms before the onset of a saccade. It has been shown that this prediction is usually correct (Collins et al., 2009) and that saccadic adaptation depends mostly on its prediction error (Collins and Wallman, 2012).

Our interpretation of the EEG decoding relies on the assumptions that saccadic adaptation is driven by prediction error and that internal prediction of saccadic landing positions is usually correct so that prediction errors only occur when the target is displaced. Our interpretation would need to be revised if, for example, saccadic adaptation resulted from postdictive updates of space rather than prediction errors (Masselink and Lappe, 2021). The alternative explanation that adaptation would be because of post-saccadic target information (Bellebaum et al., 2005) cannot explain our results because we explicitly showed that decoding labels were not predicted by post-saccadic landing positions. Furthermore, EEG decoding labels provided relevant information that improved the prediction of saccadic adaptation. In sum, the prediction error account of the SSD phenomenon seems to best fit our results.

All things considered, our results are based on current assumptions about visuomotor learning and are coherent with these studies in that we were able to decode unseen displacements successfully. However, our results also go further than previous work showing that saccade-related error signals are also present in the early stages of visual processing.

References

- Bellebaum C, Hoffmann K-P, Daum I (2005) Post-saccadic updating of visual space in the posterior parietal cortex in humans. *Behav Brain Res* 163:194–203.
- Brainard DH (1997) The Psychophysics Toolbox. *Spat Vis* 10:433–436.
- Bridgeman B, Hendry D, Stark L (1975) Failure to detect displacement of the visual world during saccadic eye movements. *Vision Res* 15:719–722.
- Collins T (2014) Trade-off between spatiotopy and saccadic plasticity. *J Vis* 14(12):28.
- Collins T, Wallman J (2012) The relative importance of retinal error and prediction in saccadic adaptation. *J Neurophysiol* 107:3342–3348.
- Collins T, Jacquet PO (2018) TMS over posterior parietal cortex disrupts trans-saccadic visual stability. *Brain Stimul* 11:390–399.
- Collins T, Rolfs M, Deubel H, Cavanagh P (2009) Post-saccadic location judgments reveal remapping of saccade targets to non-foveal locations. *J Vis* 9:29 1–9.
- Cornelissen FW, Peters EM, Palmer J (2002) The EyeLink Toolbox: eye tracking with MATLAB and the Psychophysics Toolbox. *Behav Res Methods Instrum Comput* 34:613–617.
- Deubel H, Schneider WX, Bridgeman B (1996) Postsaccadic target blanking prevents saccadic suppression of image displacement. *Vision Res* 36:985–996.
- Deubel H, Bridgeman B, Schneider WX (1998) Immediate post-saccadic information mediates space constancy. *Vision Res* 38:3147–3159.
- Duhamel JR, Colby CL, Goldberg ME (1992) The updating of the representation of visual space in parietal cortex by intended eye movements. *Science* 255:90–92.
- Gaymard B, Rivaud S, Pierrot-Deseilligny C (1994) Impairment of extraretinal eye position signals after central thalamic lesions in humans. *Exp Brain Res* 102:1–9.
- Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Parkkonen L, Hämäläinen MS (2014) MNE software for processing MEG and EEG data. *Neuroimage* 86:446–460.
- Hall NJ, Colby CL (2011) Remapping for visual stability. *Philos Trans R Soc Lond B Biol Sci* 366:528–539.
- Haufe S, Meinecke F, Görgen K, Dähne S, Haynes J-D, Blankertz B, Bießmann F (2014) On the interpretation of weight vectors of linear models in multivariate neuroimaging. *Neuroimage* 87:96–110.
- Heide W, Kömpf D (1998) Combined deficits of saccades and visuo-spatial orientation after cortical lesions. *Exp Brain Res* 123:164–171.
- Kesten H (1958) Accelerated stochastic approximation. *Ann Inst Math Stat* 29:41–59.
- Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* 164:177–190.
- Masselink J, Lappe M (2021) Visuomotor learning from postdictive motor error. *Elife* 10:e64278.
- McLaughlin SC (1967) Parametric adjustment in saccadic eye movements. *Percept Psychophys* 2:359–362.
- Medendorp WP, Goltz HC, Vilis T, Crawford JD (2003) Gaze-centered updating of visual space in human parietal cortex. *J Neurosci* 23:6209–6214.
- Medendorp WP, Goltz HC, Crawford JD, Vilis T (2005) Integration of target and effector information in human posterior parietal cortex for the planning of action. *J Neurophysiol* 93:954–962.
- Niemeier M, Douglas Crawford J, Tweed DB (2007) Optimal inference explains dimension-specific contractions of spatial perception. *Exp Brain Res* 179:313–323.
- Pisella L, Alahyane N, Blangero A, Thery F, Blanc S, Pelisson D (2011) Right-hemispheric dominance for visual remapping in humans. *Philos Trans R Soc Lond B Biol Sci* 366:572–585.
- Robinson FR, Noto CT, Bevans SE (2003) Effect of visual error size on saccade adaptation in monkey. *J Neurophysiol* 90:1235–1244.
- van Beers RJ (2007) The sources of variability in saccadic eye movements. *J Neurosci* 27:8757–8770.