



HAL
open science

Interior transmission problems with coefficients of low regularity

Georgi Vodev

► **To cite this version:**

Georgi Vodev. Interior transmission problems with coefficients of low regularity. *Inverse Problems and Imaging*, 2025, 19 (1), pp.1-33. hal-04050353

HAL Id: hal-04050353

<https://hal.science/hal-04050353v1>

Submitted on 29 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTERIOR TRANSMISSION PROBLEMS WITH COEFFICIENTS OF LOW REGULARITY

GEORGI VODEV

ABSTRACT. We obtain parabolic transmission eigenvalue-free regions for both isotropic and anisotropic interior transmission problems with L^∞ coefficients which are Lipschitz near the boundary. We also suppose that the restrictions of the coefficients on the boundary are C^μ smooth with an integer $\mu \geq 2$ and we investigate the way in which the transmission eigenvalue-free regions depend on μ .

Key words: interior transmission problems, transmission eigenvalues.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be a bounded, connected domain with a C^∞ smooth boundary $\Gamma = \partial\Omega$, and consider the interior transmission problem

$$(1.1) \quad \begin{cases} (\nabla c_1(x)\nabla + \lambda^2 n_1(x))u_1 = 0 & \text{in } \Omega, \\ (\nabla c_2(x)\nabla + \lambda^2 n_2(x))u_2 = 0 & \text{in } \Omega, \\ u_1 = u_2, c_1 \partial_\nu u_1 = c_2 \partial_\nu u_2 & \text{on } \Gamma, \end{cases}$$

where $\lambda \in \mathbb{C}$, $\operatorname{Re} \lambda \geq 0$, ν denotes the Euclidean unit inner normal to Γ and $c_j, n_j \in L^\infty(\Omega)$, $j = 1, 2$, are real-valued functions satisfying $c_j(x) \geq b_0$, $n_j(x) \geq b_0$ for some constant $b_0 > 0$. We also suppose that the coefficients are Lipschitz near the boundary. Given a parameter $0 < \delta \ll 1$, set $\Omega_\delta = \{x \in \Omega : \operatorname{dist}(x, \Gamma) < \delta\}$. More precisely, we suppose that

$$(1.2) \quad c_j, n_j \in C^1(\overline{\Omega}_\delta), \quad j = 1, 2,$$

for some δ . Throughout this paper, given an integer $k \geq 1$, C^k will denote the space of the functions a such that $\partial_x^\alpha a \in L^\infty$ for all multi-indices α with $|\alpha| = k$.

In this paper we will consider two types of interior transmission problems. The isotropic one when we have the condition

$$(1.3) \quad c_1(x) \equiv c_2(x) \equiv 1 \quad \text{in } \Omega, \quad n_1(x) \neq n_2(x) \quad \text{on } \Gamma,$$

and the anisotropic one when we have

$$(1.4) \quad (c_1(x) - c_2(x))(c_1(x)n_1(x) - c_2(x)n_2(x)) \neq 0 \quad \text{on } \Gamma.$$

If the equation (1.1) has a non-trivial solution (u_1, u_2) the complex number λ is said to be an interior transmission eigenvalue. An important question in the theory of the transmission eigenvalues is that one of finding conditions on the coefficients that guarantee that the transmission eigenvalues form a discrete set. Various such conditions were found in the papers [1], [5], [6], [8], [12], the most general ones being those in [8]. It follows from the results in [8] that under the above conditions the transmission eigenvalues are discrete in both the isotropic and anisotropic cases, so it is natural to ask if the counting function of these transmission eigenvalues admits Weyl asymptotics. Indeed, such asymptotics were obtained in [10], [11] in the case of C^∞ smooth coefficients, and more recently in [3], [9] for coefficients of very low regularity

but with much worse remainder terms. The proof in [10] relies heavily on the location of the transmission eigenvalues on the complex plane and the C^∞ regularity of the coefficients is not essential. Roughly speaking, the result in [10] says that parabolic eigenvalue-free regions imply Weyl asymptotics with a remainder term depending on the shape of the eigenvalue-free region, namely, the bigger the eigenvalue-free region is, the smaller the remainder term is. In the C^∞ setting, parabolic eigenvalue-free regions were obtained in the papers [14], [15], [16] in both the isotropic and anisotropic cases. It follows from the results in [14] and [16] that under the condition (1.3) there are no transmission eigenvalues in the region

$$(1.5) \quad |\operatorname{Im} \lambda| \geq C$$

for some constant $C > 0$. Note that the eigenvalue-free region (1.5) is optimal as shown in Section 4 of [7]. In the one dimensional case (1.5) was obtained in [13]. In the anisotropic case it is proved in [16] that under the condition

$$(1.6) \quad (c_1(x) - c_2(x))(c_1(x)n_1(x) - c_2(x)n_2(x)) < 0 \quad \text{on } \Gamma,$$

there are no transmission eigenvalues in the region

$$(1.7) \quad \operatorname{Re} \lambda \geq 1, \quad |\operatorname{Im} \lambda| \geq C.$$

On the other hand, it is proved in [14] that under the condition

$$(1.8) \quad (c_1(x) - c_2(x))(c_1(x)n_1(x) - c_2(x)n_2(x)) > 0 \quad \text{on } \Gamma,$$

there are no transmission eigenvalues in the region

$$(1.9) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{3/5}.$$

Moreover, if in addition to (1.8) the condition

$$(1.10) \quad \frac{c_1(x)}{n_1(x)} \neq \frac{c_2(x)}{n_2(x)} \quad \text{on } \Gamma,$$

is assumed, it is proved in [16] that there are no transmission eigenvalues in the region

$$(1.11) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C \log |\lambda|.$$

Our goal in the present paper is to extend the above results to coefficients having as low regularity as possible. The eigenvalue-free regions we get are not as big as those in the C^∞ case, but they are still parabolic. This fact is important in view of possible applications to obtaining Weyl asymptotics as in [10]. Our main result is the following

Theorem 1.1. *Suppose that the coefficients satisfy (1.2) and that $c_j|_\Gamma, n_j|_\Gamma \in C^\mu(\Gamma)$ with some integer $\mu \geq 2$. Then, under the condition (1.3), there exists a constant $C > 2$ such that there are no transmission eigenvalues in the region*

$$(1.12) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{1-p_1} (\log |\lambda|)^{p_2},$$

where

$$p_1 = \frac{\mu}{2\mu + 2d - 1}, \quad p_2 = \frac{2}{2\mu + 2d - 1}, \quad \text{if } \mu \leq 2d - 1,$$

$$p_1 = \frac{\mu + 2}{2\mu + 2d + 5}, \quad p_2 = \frac{2}{2\mu + 2d + 5}, \quad \text{if } 2d - 1 < \mu \leq 4d,$$

and $p_1 = \frac{2}{5}, p_2 = 0$ if $\mu > 4d$. Under the condition (1.6), there are no transmission eigenvalues in the region

$$(1.13) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{1-p_1} (\log |\lambda|)^{p_2}, \quad \operatorname{Re} \lambda \geq C|\lambda|^{1-p_3} (\log |\lambda|)^{p_4},$$

where p_1 and p_2 are the same as above, and

$$p_3 = \frac{\mu}{2\mu + 2d + 2}, \quad p_4 = \frac{1}{\mu + d + 1}.$$

Under the condition (1.8), there are no transmission eigenvalues in the region (1.12) with

$$p_1 = \frac{\mu}{2\mu + 2d + 2}, \quad p_2 = \frac{1}{\mu + d + 1}, \quad \text{if } \mu \leq \frac{4}{3}(d + 1),$$

and $p_1 = \frac{2}{7}$, $p_2 = 0$ if $\mu > \frac{4}{3}(d + 1)$.

Remark. In fact, the theorem remains valid for all $\mu \geq 2$ not necessarily integers. However, we prefer assuming that μ is integer because this simplifies the exposition significantly.

This result seems to be new even in the radial case when $\Omega = \{x \in \mathbb{R}^d : |x| \leq 1\}$ and $c_j = c_j(r)$, $n_j = n_j(r)$ depend only on the radial variable $r = |x|$. Since in this case the restrictions of c_j and n_j on the boundary are constants, the above theorem implies the following

Corollary 1.2. *Suppose that the coefficients $c_j(r)$, $n_j(r)$ are Lipschitz on the interval $[1 - \delta, 1]$ for some $0 < \delta \ll 1$. Then, under the condition (1.3), there exists a constant $C > 2$ such that there are no transmission eigenvalues in the region*

$$(1.14) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{3/5}.$$

Under the condition (1.6), there are no transmission eigenvalues in the region

$$(1.15) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{3/5}, \quad \operatorname{Re} \lambda \geq C_\epsilon |\lambda|^{1/2+\epsilon},$$

for every $0 < \epsilon \ll 1$. Under the condition (1.8), there are no transmission eigenvalues in the region

$$(1.16) \quad |\lambda| \geq C, \quad |\operatorname{Im} \lambda| \geq C|\lambda|^{5/7}.$$

To prove Theorem 1.1 we adapt the methods developed in [14], [15], [16] for C^∞ smooth coefficients. In these papers the transmission eigenvalue-free regions were derived from suitable approximations of the interior Dirichlet-to-Neumann map outside a parabolic neighbourhood of the real axis. In other words, the problem of finding transmission eigenvalue-free regions was reduced to that one of finding as good as possible approximation of the interior Dirichlet-to-Neumann map by a semiclassical pseudodifferential operator ($h - \Psi$ DO) on the boundary (with a semiclassical parameter $h \sim |\lambda|^{-1}$) outside as small as possible parabolic neighbourhood of the real axis. With such an approximation in hands, in order to get the eigenvalue-free regions, one has to invert a semiclassical pseudodifferential operator whose symbol, say a , can be computed explicitly in terms of the restrictions of the coefficients on the boundary and the principal symbol of the Laplace-Beltrami operator on the boundary. Recall that the boundary Γ can be considered as a compact Riemannian manifold of dimension $d - 1$ without boundary with a Riemannian metric induced by the Euclidean one. Then the conditions (1.3) and (1.4) guarantee that the function a is invertible outside some parabolic neighbourhood of the real axis. Then to show that the semiclassical pseudodifferential operator $\operatorname{Op}_h(a)$ with symbol a is invertible, one has to make use of some pseudodifferential calculus which are well-known in the C^∞ setting. Thus one arrives at the conclusion that the eigenvalue-free regions correspond to the regions where the Dirichlet-to-Neumann map has a good approximation and where the operator $\operatorname{Op}_h(a)$ is invertible. Therefore, to get as big as possible eigenvalue-free regions one has to find as good as possible approximation of the interior Dirichlet-to-Neumann map. In the C^∞ case this is done in [14], [15], [16] by constructing semiclassical parametrices for the solutions of the equation (1.1) near the boundary mod $\mathcal{O}(h^\infty)$. Note that the C^∞ regularity of the coefficients

near the boundary is essential in the analysis in these papers. The main difficulty in constructing such parametrices is that one has to solve the eikonal equation mod $\mathcal{O}(x_1^\infty)$ and the transport equations mod $\mathcal{O}(x_1^\infty + h^\infty)$, where $0 < x_1 \ll 1$ is the normal coordinate near the boundary, that is the distance to Γ . This is especially difficult to do when the boundary data is microlocally supported near the glancing region. Of course, when the coefficients are of low regularity this analysis does not work any more. In this case we built (see Section 4) a less accurate parametrix by solving the eikonal equation mod $\mathcal{O}(x_1)$ and with no need to solve the transport equations. In other words, our parametrix is the simplest possible in this context. Nevertheless, this leads to some approximation of the Dirichlet-to-Neumann map, which of course is not as good as in the C^∞ case. Consequently, the eigenvalue-free regions we get by using this approximation are much smaller than those in the C^∞ case. In order to make this approach work, however, we need to use $h - \Psi$ DOs with symbols which are C^μ smooth with respect to the space variable. To this end, we develop in Section 2 some pseudodifferential calculus for such operators. In particular, we find some useful criteria for L^2 boundedness (see Proposition 2.4) and we also show how to invert such operators (see Proposition 2.6). Note finally that $\mu = 2$ is the lowest regularity that makes possible the construction of some parametrix.

The paper is organized as follows. In Section 2 we recall some basic properties of the $h - \Psi$ DOs with C^∞ symbols concerning the L^2 boundedness and the composition of two operators. We then extend these properties to $h - \Psi$ DOs with symbols which are C^μ smooth with respect to the space variable. This is done by using a suitable interpolation between symbols which are L^∞ and C^∞ smooth with respect to the space variable. To this end, we establish a criteria for L^2 boundedness for $h - \Psi$ DOs with L^∞ symbols (see Proposition 2.3). In Section 3 we prove a priori estimates which allow us to bound the norm of the difference between the Dirichlet-to-Neumann map and the parametrix. In Section 4 we build a parametrix for the solutions of the equation (1.1) near the boundary, which is a finite sum of h -FIOs. We then use it to get the parametrix for the Dirichlet-to-Neumann map. In Section 5 we improve our parametrix in the elliptic region in the isotropic case. This allows us to get a better approximation of the Dirichlet-to-Neumann map in this case. This improvement is crucial in order to get the eigenvalue-free regions in the isotropic case. In Section 6 we invert the operator $\text{Op}_h(a)$ with a symbol a as above outside some parabolic neighbourhoods of the real axis as well as outside some parabolic neighbourhoods of the imaginary axis when the condition (1.6) is assumed. This gives us the eigenvalue-free regions. Note that in our case a is C^μ smooth with respect to the space variable, so we need to use the pseudodifferential calculus developed in Section 2.

2. $h - \Psi$ DOs WITH SYMBOLS OF LOW REGULARITY

We begin this section by recalling some basic properties of the $h - \Psi$ DOs with C^∞ symbols. In what follows we will identify the cotangent space $T^*\mathbb{R}^{d-1}$ with $\mathbb{R}^{d-1} \times \mathbb{R}^{d-1}$. Introduce the space S^k , $k \in \mathbb{R}$, of all functions $a(x, \xi) \in C^\infty(\mathbb{R}^{d-1} \times \mathbb{R}^{d-1})$ satisfying

$$(2.1) \quad \left| \partial_\xi^\alpha \partial_x^\beta a(x, \xi) \right| \leq C_{\alpha, \beta} \langle \xi \rangle^{k - |\alpha|}$$

for all multi-indices α and β , where $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$. We define the $h - \Psi$ DO with a symbol $a \in S^k$ by

$$(\text{Op}_h(a)f)(x) = (2\pi h)^{-d+1} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^{d-1}} e^{-\frac{i}{h}\langle x-y, \xi \rangle} a(x, \xi) f(y) d\xi dy,$$

where $0 < h \leq 1$ is a semiclassical parameter. Denote by $H_h^k(\mathbb{R}^{d-1})$ the Sobolev space equipped with the h -semiclassical norm

$$\|u\|_{H_h^k} := \left\| \text{Op}_h(\langle \xi \rangle^k) u \right\|_{L^2},$$

where $\|\cdot\|_{L^2}$ denotes the norm in $L^2(\mathbb{R}^{d-1})$. It is well-known that the operator $\text{Op}_h(a) : H_h^k(\mathbb{R}^{d-1}) \rightarrow L^2(\mathbb{R}^{d-1})$ is bounded for symbols $a \in S^k$. In fact, we have a stronger result.

Proposition 2.1. *Let $a \in C^\infty(\mathbb{R}^{d-1} \times \mathbb{R}^{d-1})$ satisfy*

$$(2.2) \quad \left| \partial_\xi^\alpha \partial_x^\beta a(x, \xi) \right| \leq C_{\alpha, \beta} \langle \xi \rangle^k$$

for all multi-indices α and β . Then there exists an integer s_d depending only on the dimension such that we have the bound

$$(2.3) \quad \|\text{Op}_h(a)\|_{H_h^k \rightarrow L^2} \lesssim \sum_{|\alpha|+|\beta| \leq s_d} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha, \beta}.$$

Proof. Since $\text{Op}_h(a) = \text{Op}_h(a \langle \xi \rangle^{-k}) \text{Op}_h(\langle \xi \rangle^k)$ and the function $a \langle \xi \rangle^{-k}$ satisfies (2.2) with $k = 0$, it suffices to prove (2.3) for $k = 0$. In this case, it is easy to see that the norm in the left-hand side of (2.3) is equal to the norm $\|\text{Op}_1(a_h)\|_{L^2 \rightarrow L^2}$, where $a_h(x, \xi) = a(h^{1/2}x, h^{1/2}\xi)$. On the other hand, by Theorem 18.6.3 of [4] (see also Theorem 7.11 of [2]) the operator $\text{Op}_1(a_h) : L^2 \rightarrow L^2$ is bounded and

$$\|\text{Op}_1(a_h)\|_{L^2 \rightarrow L^2} \lesssim \sum_{|\alpha|+|\beta| \leq s_d} \sup \left| \partial_\xi^\alpha \partial_x^\beta a_h(x, \xi) \right|,$$

which clearly implies (2.3) with $k = 0$. \square

We will now use this proposition to prove the following

Proposition 2.2. *Let a_1 satisfy (2.2) with $k = 0$ and constants $C_{\alpha, \beta}^{(1)}$, and let a_2 satisfy (2.2) with constants $C_{\alpha, \beta}^{(2)}$. Then we have the bound*

$$(2.4) \quad \begin{aligned} & \|\text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2)\|_{H_h^k \rightarrow L^2} \\ & \lesssim \sum_{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2| \leq s'_d, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}, \end{aligned}$$

where $s'_d = 2d + 5 + s_d$.

Proof. We will use the fact that

$$\text{Op}_h(a_1)\text{Op}_h(a_2) = \text{Op}_h(b),$$

where

$$\begin{aligned} b(x, \xi) &= e^{ih \langle D_\eta, D_y \rangle} a_1(x, \eta) a_2(y, \xi) \Big|_{y=x, \eta=\xi} \\ &= (2\pi)^{-d+1} \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}^{d-1}} e^{i \langle \zeta_1, \zeta_2 \rangle} a_1(x, \xi - h^{1/2} \zeta_1) a_2(x - h^{1/2} \zeta_2, \xi) d\zeta_1 d\zeta_2, \end{aligned}$$

where we have put $D_\eta = -i\nabla_\eta$, $D_y = -i\nabla_y$. Set $\zeta = (\zeta_1, \zeta_2) \in \mathbb{R}^{2d-2}$ and $\varphi(\zeta) = \langle \zeta_1, \zeta_2 \rangle$. Clearly,

$$\nabla_\zeta \varphi = (\nabla_{\zeta_1} \varphi, \nabla_{\zeta_2} \varphi) = (\zeta_2, \zeta_1),$$

so we have $|\nabla_\zeta \varphi| = |\zeta|$. Consider now the function

$$g(t) = a_1(x, \xi - th^{1/2} \zeta_1) a_2(x - th^{1/2} \zeta_2, \xi), \quad 0 \leq t \leq 1.$$

An easy computation leads to the formulas

$$\begin{aligned} g'(t) &= -h^{1/2} \langle \zeta_1, \nabla_\xi a_1(x, \xi - th^{1/2} \zeta_1) \rangle a_2(x - th^{1/2} \zeta_2, \xi) \\ &\quad - h^{1/2} \langle \zeta_2, \nabla_x a_2(x - th^{1/2} \zeta_2, \xi) \rangle a_1(x, \xi - th^{1/2} \zeta_1), \end{aligned}$$

$$g''(t) = \mathcal{G}_1(t) + \mathcal{G}_2(t),$$

where

$$\begin{aligned} \mathcal{G}_1 &= h\langle \zeta_1, \nabla_\xi \langle \zeta_1, \nabla_\xi a_1(x, \xi - th^{1/2}\zeta_1) \rangle \rangle a_2(x, \xi) \\ &\quad + h\langle \zeta_2, \nabla_x \langle \zeta_2, \nabla_x a_2(x - th^{1/2}\zeta_2, \xi) \rangle \rangle a_1(x, \xi), \\ \mathcal{G}_2 &= h\langle \zeta_1, \nabla_\xi \langle \zeta_1, \nabla_\xi a_1(x, \xi - th^{1/2}\zeta_1) \rangle \rangle \left(a_2(x - th^{1/2}\zeta_2, \xi) - a_2(x, \xi) \right) \\ &\quad + h\langle \zeta_2, \nabla_x \langle \zeta_2, \nabla_x a_2(x - th^{1/2}\zeta_2, \xi) \rangle \rangle \left(a_1(x, \xi - th^{1/2}\zeta_1) - a_1(x, \xi) \right) \\ &\quad + 2h\langle \zeta_1, \nabla_\xi a_1(x, \xi - th^{1/2}\zeta_1) \rangle \langle \zeta_2, \nabla_x a_2(x - th^{1/2}\zeta_2, \xi) \rangle. \end{aligned}$$

Clearly, the function \mathcal{G}_2 satisfies the bounds

$$(2.5) \quad \left| \partial_\zeta^\gamma \mathcal{G}_2 \right| \lesssim \langle \xi \rangle^k \sum_{\ell=0}^3 |\zeta|^\ell \sum_{|\alpha|+|\beta|=|\gamma|+\ell, |\alpha|\geq 1, |\beta|\geq 1} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha,0}^{(1)} C_{0,\beta}^{(2)}$$

for all multi-indices γ , uniformly in t . Observe now that we can write

$$g(1) = g(0) + g'(0) + \int_0^1 (1-t)g''(t)dt,$$

where

$$\begin{aligned} g(0) &= a_1(x, \xi)a_2(x, \xi), \\ g'(0) &= -h^{1/2}\langle \zeta_1, \nabla_\xi a_1(x, \xi) \rangle a_2(x, \xi) - h^{1/2}\langle \zeta_2, \nabla_x a_2(x, \xi) \rangle a_1(x, \xi). \end{aligned}$$

On the other hand, given any function $\psi \in C^\infty(\mathbb{R}^{d-1})$, we have the formula

$$(2\pi)^{-d+1} \int_{\mathbb{R}^{2d-2}} e^{i\varphi(\zeta)} \psi(\zeta_j) d\zeta = \psi(0)$$

where $j = 1, 2$. Therefore we have

$$\int_{\mathbb{R}^{2d-2}} e^{i\varphi} g'(0) d\zeta = \int_{\mathbb{R}^{2d-2}} e^{i\varphi} \mathcal{G}_1(t) d\zeta = 0.$$

We get from the above formulas

$$(2.6) \quad b - a_1 a_2 = (2\pi)^{-d+1} \int_0^1 (1-t) \int_{\mathbb{R}^{2d-2}} e^{i\varphi} \mathcal{G}_2(t) d\zeta dt.$$

Let $\phi \in C_0^\infty(\mathbb{R})$, $\phi(\sigma) = 1$ for $|\sigma| \leq 1$, $\phi(\sigma) = 0$ for $|\sigma| \geq 2$. We will now bound the integrals

$$\begin{aligned} I_1 &= (2\pi)^{-d+1} \int_{\mathbb{R}^{2d-2}} e^{i\varphi} \phi(|\zeta|) \mathcal{G}_2(t) d\zeta, \\ I_2 &= (2\pi)^{-d+1} \int_{\mathbb{R}^{2d-2}} e^{i\varphi} (1-\phi)(|\zeta|) \mathcal{G}_2(t) d\zeta. \end{aligned}$$

By (2.5) with $\gamma = 0$, we have

$$(2.7) \quad |I_1| \lesssim \langle \xi \rangle^k \sum_{|\alpha|+|\beta|\leq 3, |\alpha|\geq 1, |\beta|\geq 1} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha,0}^{(1)} C_{0,\beta}^{(2)}$$

uniformly in t . To bound the second integral we will integrate by parts. To this end observe that

$$L_\zeta e^{i\varphi} = e^{i\varphi},$$

where

$$L_\zeta = -i|\zeta|^{-2} \langle \nabla_\zeta \varphi, \nabla_\zeta \rangle.$$

It is easy to see that, given any integer $m \geq 1$, the adjoint operator to L_ζ^m satisfies

$$(2.8) \quad |(L_\zeta^m)^* f(\zeta)| \lesssim |\zeta|^{-m} \sum_{|\gamma| \leq m} \left| \partial_\zeta^\gamma f(\zeta) \right|.$$

We can now write the integral I_2 in the form

$$I_2 = (2\pi)^{-d+1} \int_{\mathbb{R}^{2d-2}} e^{i\varphi} (L_\zeta^m)^* ((1-\phi)\mathcal{G}_2(t)) d\zeta.$$

Using (2.5) and (2.8) we obtain

$$(2.9) \quad \begin{aligned} |I_2| &\lesssim \langle \xi \rangle^k \sum_{\ell=0}^3 \int_{\zeta \in \mathbb{R}^{2d-2}; |\zeta| \geq 1} |\zeta|^{\ell-m} d\zeta \sum_{|\alpha|+|\beta| \leq m+\ell, |\alpha| \geq 1, |\beta| \geq 1} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha,0}^{(1)} C_{0,\beta}^{(2)} \\ &\lesssim \langle \xi \rangle^k \sum_{|\alpha|+|\beta| \leq 2d+5, |\alpha| \geq 1, |\beta| \geq 1} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha,0}^{(1)} C_{0,\beta}^{(2)} \end{aligned}$$

if we take $m = 2d + 2$. By (2.6), (2.7) and (2.9), we conclude

$$(2.10) \quad |b - a_1 a_2| \lesssim \langle \xi \rangle^k \sum_{|\alpha_1|+|\beta_2| \leq 2d+5, |\alpha_1| \geq 1, |\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_2|}{2}} C_{\alpha_1,0}^{(1)} C_{0,\beta_2}^{(2)}.$$

Similarly, given any multi-indices α and β , we can obtain the bounds

$$(2.11) \quad \begin{aligned} &\langle \xi \rangle^{-k} h^{\frac{|\alpha|+|\beta|}{2}} |\partial_\xi^\alpha \partial_x^\beta (b - a_1 a_2)| \\ &\lesssim \sum_{|\alpha_1|+|\alpha_2|+|\beta_1|+|\beta_2| \leq |\alpha|+|\beta|+2d+5, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\alpha_2|+|\beta_1|+|\beta_2|}{2}} C_{\alpha_1,\beta_1}^{(1)} C_{\alpha_2,\beta_2}^{(2)}. \end{aligned}$$

Clearly, the bound (2.4) follows from (2.11) and Proposition 2.1. \square

The above definition of the h - Ψ DOs makes sense also for symbols $a(x, \xi)$ which are smooth only with respect to the variable ξ and satisfy

$$(2.12) \quad \left| \partial_\xi^\alpha a(x, \xi) \right| \leq C_\alpha \langle \xi \rangle^{k-|\alpha|}$$

for all multi-indices α , where $k \in \mathbb{R}$. For such operators, we will prove the following

Proposition 2.3. *Let a satisfy (2.12) and let $k' > k$. Then we have*

$$(2.13) \quad \|\text{Op}_h(a)\|_{H_h^{k'} \rightarrow L^2} \lesssim \log(1 + h^{-1}) \sum_{|\alpha| \leq d} C_\alpha.$$

Proof. As above, it suffices to prove (2.13) for $k' = 0$. Then we have $k < 0$. Clearly, the kernel K of the operator $\text{Op}_h(a)$ is

$$K(x, y) = (2\pi h)^{-d+1} \int_{\mathbb{R}^{d-1}} e^{-\frac{i}{h} \langle x-y, \xi \rangle} a(x, \xi) d\xi.$$

In view of Schur's lemma, it suffices to bound from above the integrals

$$\int_{\mathbb{R}^{d-1}} |K(x, y)| dx + \int_{\mathbb{R}^{d-1}} |K(x, y)| dy.$$

Integrating by parts in the above integral leads to the bounds

$$|K(x, y)| \lesssim h^{-d+1} \left(\frac{h}{|x-y|} \right)^d \sum_{|\alpha|=d} \int_{\mathbb{R}^{d-1}} |\partial_\xi^\alpha a(x, \xi)| d\xi$$

$$\begin{aligned} &\lesssim \frac{h}{|x-y|^d} \sum_{|\alpha|=d} C_\alpha \int_{\mathbb{R}^{d-1}} \langle \xi \rangle^{k-d} d\xi \\ &\lesssim \frac{h}{|x-y|^d} \sum_{|\alpha|=d} C_\alpha. \end{aligned}$$

Hence

$$\begin{aligned} &\int_{|x-y|\geq h} |K(x,y)| dx + \int_{|x-y|\geq h} |K(x,y)| dy \\ &\lesssim h \sum_{|\alpha|=d} C_\alpha \int_{w \in \mathbb{R}^{d-1}; |w|\geq h} |w|^{-d} dw \\ &\lesssim h \sum_{|\alpha|=d} C_\alpha \int_h^\infty \sigma^{-2} d\sigma \lesssim \sum_{|\alpha|=d} C_\alpha. \end{aligned}$$

We will now bound the kernel for $|x-y| < h$. Set $A = \frac{h}{|x-y|} > 1$. Let the function ϕ be as above and decompose the kernel as $K = K_1 + K_2$, where

$$K_1(x,y) = (2\pi h)^{-d+1} \int_{\mathbb{R}^{d-1}} e^{-\frac{i}{h}\langle x-y, \xi \rangle} a(x, \xi) \phi(|\xi|/A) d\xi.$$

An integration by parts leads to the bounds

$$\begin{aligned} |K_1(x,y)| &\lesssim h^{-d+1} \left(\frac{h}{|x-y|} \right)^{d-2} \sum_{|\alpha|=d-2} \int_{\mathbb{R}^{d-1}} |\partial_\xi^\alpha (a(x, \xi) \phi(|\xi|/A))| d\xi \\ &\lesssim \frac{h^{-1}}{|x-y|^{d-2}} \sum_{|\alpha|\leq d-2} A^{|\alpha|+2-d} \int_{|\xi|\leq A} |\partial_\xi^\alpha a(x, \xi)| d\xi \\ &\lesssim \frac{h^{-1}}{|x-y|^{d-2}} \sum_{|\alpha|\leq d-2} C_\alpha A^{|\alpha|+2-d} \int_{|\xi|\leq A} \langle \xi \rangle^{k-|\alpha|} d\xi \\ &\lesssim \frac{h^{-1}}{|x-y|^{d-2}} \sum_{|\alpha|\leq d-2} C_\alpha A^{|\alpha|+2-d} \int_0^{A+1} \sigma^{k-|\alpha|+d-2} d\sigma \\ &\lesssim \frac{h^{-1} A^{1+k}}{|x-y|^{d-2}} \sum_{|\alpha|\leq d-2} C_\alpha \\ &\lesssim \frac{A^k}{|x-y|^{d-1}} \sum_{|\alpha|\leq d-2} C_\alpha. \end{aligned}$$

Set $\epsilon = (\log(1+h^{-1}))^{-1}$. Clearly, there is $0 < h_0 \ll 1$ such that $\epsilon < -k$ for $0 < h < h_0$, and hence $A^k < A^{-\epsilon}$. Thus, taking into account that $h^{-\epsilon} = e$, we get

$$|K_1(x,y)| \lesssim \frac{1}{|x-y|^{d-1-\epsilon}} \sum_{|\alpha|\leq d-2} C_\alpha$$

for $0 < h < h_0$. Clearly, for $h_0 \leq h \leq 1$ the above bound still holds with $0 < \epsilon \ll 1$ independent of h . Hence, in all cases, we have

$$\int_{|x-y|\leq h} |K_1(x,y)| dx + \int_{|x-y|\leq h} |K_1(x,y)| dy$$

$$\begin{aligned}
 &\lesssim \sum_{|\alpha| \leq d-2} C_\alpha \int_{w \in \mathbb{R}^{d-1}; |w| \leq h} |w|^{-d+1+\epsilon} dw \\
 &\lesssim \sum_{|\alpha| \leq d-2} C_\alpha \int_0^h \sigma^{-1+\epsilon} d\sigma \lesssim \epsilon^{-1} \sum_{|\alpha| \leq d-2} C_\alpha.
 \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 |K_2(x, y)| &\lesssim h^{-d+1} \left(\frac{h}{|x-y|} \right)^{d-1} \sum_{|\alpha|=d-1} \int_{\mathbb{R}^{d-1}} |\partial_\xi^\alpha (a(x, \xi)(1-\phi)(|\xi|/A))| d\xi \\
 &\lesssim \frac{1}{|x-y|^{d-1}} \sum_{|\alpha| \leq d-1} A^{|\alpha|+1-d} \int_{|\xi| \geq A} |\partial_\xi^\alpha a(x, \xi)| d\xi \\
 &\lesssim \frac{1}{|x-y|^{d-1}} \sum_{|\alpha| \leq d-1} C_\alpha A^{|\alpha|+1-d} \int_{|\xi| \geq A} \langle \xi \rangle^{k-|\alpha|} d\xi \\
 &\lesssim \frac{A^{-\epsilon}}{|x-y|^{d-1}} \sum_{|\alpha| \leq d-1} C_\alpha \int_{\mathbb{R}^{d-1}} \langle \xi \rangle^{k-d+1+\epsilon} d\xi \\
 &\lesssim \frac{1}{|x-y|^{d-1-\epsilon}} \sum_{|\alpha| \leq d-1} C_\alpha \int_0^\infty (\sigma+1)^{k-1+\epsilon} d\sigma \\
 &\lesssim \frac{1}{|x-y|^{d-1-\epsilon}} \sum_{|\alpha| \leq d-1} C_\alpha.
 \end{aligned}$$

As above, this implies

$$\int_{|x-y| \leq h} |K_2(x, y)| dx + \int_{|x-y| \leq h} |K_2(x, y)| dy \lesssim \epsilon^{-1} \sum_{|\alpha| \leq d-1} C_\alpha,$$

which clearly completes the proof of (2.13) with $k' = 0$. \square

The logarithmic term in the right-hand side of (2.13) can be removed if some regularity of the symbol in x is assumed. Indeed, in this case we can decompose a as a sum of a large symbol which is C^∞ in x and a small symbol which is L^∞ in x . Then we can apply (2.3) to the C^∞ part and (2.13) to the L^∞ part. More precisely, we have the following

Proposition 2.4. *Suppose that a satisfy (2.1) for all multi-indices α and β such that $|\beta| \leq \mu$, $\mu \geq 1$ being integer, and let $k' > k$. Then we have*

$$\begin{aligned}
 \|\text{Op}_h(a)\|_{H_h^{k'} \rightarrow L^2} &\lesssim \sum_{|\alpha|+|\beta| \leq s_d, |\beta| \leq \mu} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha, \beta} \\
 (2.14) \quad &+ h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha| \leq d, |\beta| \leq \mu} C_{\alpha, \beta}.
 \end{aligned}$$

Moreover, if $\mu \geq s_d$ the second sum in the right-hand side of (2.14) can be removed.

Proof. Let $\phi_0 \in C_0^\infty(\mathbb{R}^{d-1})$, $\phi_0 \geq 0$, be such that $\int_{\mathbb{R}^{d-1}} \phi_0(x) dx = 1$. Given a parameter $0 < t \leq 1$, set

$$a_t(x, \xi) = t^{-d+1} \int_{\mathbb{R}^{d-1}} \phi_0((y-x)/t) a(y, \xi) dy = \int_{\mathbb{R}^{d-1}} \phi_0(w) a(x+tw, \xi) dw.$$

Clearly, the function a_t is C^∞ smooth in x and satisfies (2.1) uniformly in t for all multi-indices α and β such that $|\beta| \leq \mu$, while for $|\beta| \geq \mu + 1$ it satisfies the bounds

$$(2.15) \quad \left| \partial_\xi^\alpha \partial_x^\beta a_t(x, \xi) \right| \leq t^{-|\beta|+\mu} \langle \xi \rangle^{k-|\alpha|} \sum_{|\beta'|=\mu} C_{\alpha, \beta'}.$$

The Taylor expansion of the function $a(x+tw, \xi)$ at $t=0$ yields

$$\begin{aligned} a(x+tw, \xi) &= \sum_{s=0}^{\mu-1} \frac{t^s}{s!} \partial_t^s a(x+tw, \xi)|_{t=0} + \frac{t^\mu}{\mu!} \partial_t^\mu a(x+tw, \xi)|_{t=0} \\ &= \sum_{s=0}^{\mu-1} \frac{t^s}{s!} \sum_{|\beta|=s} w^\beta \partial_x^\beta a(x, \xi) + \frac{t^\mu}{\mu!} \sum_{|\beta|=\mu} w^\beta \partial_x^\beta a(x+t'w, \xi) \end{aligned}$$

with some $0 \leq t' \leq t$. Clearly, the functions $\partial_x^\beta a(x+t'w, \xi)$ with $|\beta| = \mu$ satisfy (2.12) uniformly in $t'w$. Therefore, by Proposition 2.3 we have

$$(2.16) \quad \sum_{|\beta|=\mu} \left\| \text{Op}_h(\partial_x^\beta a(x+t'w, \xi)) \right\|_{H_h^{k'} \rightarrow L^2} \lesssim \log(1+h^{-1}) \sum_{|\alpha| \leq d, |\beta|=\mu} C_{\alpha, \beta}.$$

We now apply Proposition 2.1 with a_t . In view of (2.15) we get

$$(2.17) \quad \left\| \text{Op}_h(a_t) \right\|_{H_h^k \rightarrow L^2} \lesssim \sum_{|\alpha|+|\beta| \leq s_d, |\beta| \leq \mu} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha, \beta}$$

if we take $t = h^{1/2}$. Thus, by (2.16) and (2.17), we obtain

$$(2.18) \quad \begin{aligned} \left\| \text{Op}_h(a) \right\|_{H_h^{k'} \rightarrow L^2} &\lesssim \sum_{s=1}^{\mu-1} h^{s/2} \sum_{|\beta|=s} \left\| \text{Op}_h(\partial_x^\beta a) \right\|_{H_h^{k'} \rightarrow L^2} \\ &+ h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha| \leq d, |\beta|=\mu} C_{\alpha, \beta} + \sum_{|\alpha|+|\beta| \leq s_d, |\beta| \leq \mu} h^{\frac{|\alpha|+|\beta|}{2}} C_{\alpha, \beta}, \end{aligned}$$

where the first sum in the right-hand side is zero if $\mu = 1$. It is easy to see now that (2.14) follows from (2.18) by induction in μ . \square

We will next extend Proposition 2.2 to $h - \Psi$ DOs with C^μ smooth symbols, where $\mu \geq 1$ is integer. We will first prove the following

Proposition 2.5. *Let a_1 satisfy (2.1) with $k < 0$ for all multi-indices α and β such that $|\beta| \leq \mu$ with constants $C_{\alpha, \beta}^{(1)}$, and let a_2 satisfy (2.1) for all multi-indices α and β with constants $C_{\alpha, \beta}^{(2)}$. Then we have the bound*

$$\begin{aligned} &\left\| \text{Op}_h(a_1) \text{Op}_h(a_2) - \text{Op}_h(a_1 a_2) \right\|_{H_h^k \rightarrow L^2} \\ &\lesssim \sum_{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2| \leq s'_d, |\beta_1| \leq \mu, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \end{aligned}$$

$$\begin{aligned}
 & +h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1| = \mu, |\alpha_2| + |\beta_2| \leq s_d} h^{\frac{|\alpha_2| + |\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\
 (2.19) \quad & +h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha_1| + |\alpha_2| \leq d, |\beta_1| = \mu, \beta_2 = 0} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}.
 \end{aligned}$$

Let a_1 satisfy (2.1) with $k = 0$ for all multi-indices α and β with constants $C_{\alpha, \beta}^{(1)}$, and let a_2 satisfy (2.1) with $k < k'$ for all multi-indices α and β such that $|\beta| \leq \mu$ with constants $C_{\alpha, \beta}^{(2)}$. Then we have the bound

$$\begin{aligned}
 & \|\text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2)\|_{H_h^{k'} \rightarrow L^2} \\
 \lesssim & \sum_{|\alpha_1| + |\beta_1| + |\alpha_2| + |\beta_2| \leq s'_d, |\beta_2| \leq \mu, |\alpha_1| + |\beta_1| \geq 1, |\alpha_2| + |\beta_2| \geq 1} h^{\frac{|\alpha_1| + |\beta_1| + |\alpha_2| + |\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\
 & +h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha_1| + |\beta_1| \leq s_d, |\alpha_2| \leq d, |\beta_2| = \mu} h^{\frac{|\alpha_1| + |\beta_1|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\
 (2.20) \quad & +h^{\mu/2} \log(1+h^{-1}) \sum_{|\alpha_1| + |\alpha_2| \leq d, \beta_1 = 0, |\beta_2| = \mu} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}.
 \end{aligned}$$

Let a_1 satisfy (2.1) with $k = 0$ for all multi-indices α and β with constants $C_{\alpha, \beta}^{(1)}$, and let a_2 be independent of the variable ξ and satisfies

$$(2.21) \quad \left| \partial_x^\beta a_2(x) \right| \leq C_\beta^{(2)}$$

for all multi-indices β such that $|\beta| \leq \mu$. Then we have the bound

$$\begin{aligned}
 & \|\text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2)\|_{L^2 \rightarrow L^2} \\
 \lesssim & \sum_{|\alpha_1| + |\beta_1| + |\beta_2| \leq s'_d, |\alpha_1| + |\beta_1| \geq 1, 1 \leq |\beta_2| \leq \mu} h^{\frac{|\alpha_1| + |\beta_1| + |\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\beta_2}^{(2)} \\
 (2.22) \quad & +h^{\mu/2} \sum_{|\alpha_1| + |\beta_1| \leq s_d, |\beta_2| = \mu} h^{\frac{|\alpha_1| + |\beta_1|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\beta_2}^{(2)}.
 \end{aligned}$$

Proof. We will only prove (2.19) since (2.20) and (2.22) can be obtained in the same way. We approximate a_1 by the smooth function $a_{1,t}$ as in the proof of Proposition 2.4. Then the Taylor expansion allows us to write the difference $a_1 - a_{1,t}$ in the form

$$(2.23) \quad a_1 - a_{1,t} = \sum_{s=1}^{\mu-1} t^s a_1^{(s)} + t^\mu a_{1,t}^{(\mu)},$$

where

$$\begin{aligned}
 a_1^{(s)} &= \sum_{|\beta|=s} c_\beta \partial_x^\beta a_1(x, \xi), \quad 1 \leq s \leq \mu - 1, \\
 a_{1,t}^{(\mu)} &= (\mu!)^{-1} \sum_{|\beta|=\mu} \int_{\mathbb{R}^{d-1}} \phi_0(w) w^\beta \partial_x^\beta a_1(x + t'w, \xi) dw.
 \end{aligned}$$

Clearly, when $\mu = 1$ the sum in the right-hand side of (2.23) is zero. We also have that $a_1^{(s)}$, $1 \leq s \leq \mu - 1$, satisfy (2.1) with $k < 0$ for all multi-indices α and β such that $|\beta| \leq \mu - s$, while

$a_{1,t}^{(\mu)}$ satisfies (2.12) with $k < 0$ uniformly in t for all multi-indices α . In view of (2.23), we can write

$$(2.24) \quad \text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2) = \mathcal{A}_0 + \sum_{s=1}^{\mu-1} t^s \mathcal{A}_s + t^\mu \mathcal{A}_\mu,$$

where

$$\begin{aligned} \mathcal{A}_0 &= \text{Op}_h(a_{1,t})\text{Op}_h(a_2) - \text{Op}_h(a_{1,t} a_2), \\ \mathcal{A}_s &= \text{Op}_h(a_1^{(s)})\text{Op}_h(a_2) - \text{Op}_h(a_1^{(s)} a_2), \\ \mathcal{A}_\mu &= \text{Op}_h(a_{1,t}^{(\mu)})\text{Op}_h(a_2) - \text{Op}_h(a_{1,t}^{(\mu)} a_2). \end{aligned}$$

We take $t = h^{1/2}$ and use Proposition 2.2 to bound the norm of the operator \mathcal{A}_0 . Taking into account the bounds (2.15), we get

$$\begin{aligned} & \|\mathcal{A}_0\|_{H_h^k \rightarrow L^2} \\ & \lesssim \sum_{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2| \leq s'_d, |\beta_1| \leq \mu, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}. \end{aligned}$$

To bound the norm of the operator \mathcal{A}_μ we will use Propositions 2.1 and 2.3. We get

$$\begin{aligned} \|\mathcal{A}_\mu\|_{H_h^k \rightarrow L^2} & \leq \left\| \text{Op}_h(a_{1,t}^{(\mu)}) \right\|_{L^2 \rightarrow L^2} \|\text{Op}_h(a_2)\|_{H_h^k \rightarrow L^2} + \left\| \text{Op}_h(a_{1,t}^{(\mu)} a_2) \right\|_{H_h^k \rightarrow L^2} \\ & \lesssim \log(1 + h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1| = \mu} C_{\alpha_1, \beta_1}^{(1)} \sum_{|\alpha_2|+|\beta_2| \leq s_d} h^{\frac{|\alpha_2|+|\beta_2|}{2}} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + \log(1 + h^{-1}) \sum_{|\alpha_1|+|\alpha_2| \leq d, |\beta_1| = \mu, \beta_2 = 0} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}. \end{aligned}$$

Now (2.19) follows from the above bounds and (2.24) by induction in μ . \square

We will now use the above proposition to prove the following

Proposition 2.6. *Let a_1 satisfy (2.1) with $k < 0$ for all multi-indices α and β such that $|\beta| \leq \mu_1$ with constants $C_{\alpha, \beta}^{(1)}$, and let a_2 satisfy (2.1) with $k < k'$ for all multi-indices α and β such that $|\beta| \leq \mu_2$ with constants $C_{\alpha, \beta}^{(2)}$. Then we have the bound*

$$\begin{aligned} & \left\| \text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2) \right\|_{H_h^{k'} \rightarrow L^2} \\ & \lesssim \sum_{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2| \leq s'_d, |\beta_1| \leq \mu_1, |\beta_2| \leq \mu_2, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_1/2} \log(1 + h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1| = \mu_1, |\alpha_2|+|\beta_2| \leq s_d, |\beta_2| \leq \mu_2} h^{\frac{|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_1/2} \log(1 + h^{-1}) \sum_{|\alpha_1|+|\alpha_2| \leq d, |\beta_1| = \mu_1, \beta_2 = 0} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_2/2} \log(1 + h^{-1}) \sum_{|\alpha_1|+|\beta_1| \leq s_d, |\beta_1| \leq \mu_1, |\alpha_2| \leq d, |\beta_2| = \mu_2} h^{\frac{|\alpha_1|+|\beta_1|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_2/2} \log(1 + h^{-1}) \sum_{|\alpha_1|+|\alpha_2| \leq d, \beta_1 = 0, |\beta_2| = \mu_2} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \end{aligned}$$

$$(2.25) \quad +h^{(\mu_1+\mu_2)/2} \log^2(1+h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1|=\mu_1, |\alpha_2| \leq d, |\beta_2|=\mu_2} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}.$$

Moreover, if a_2 is independent of the variable ξ and satisfies (2.21) for all multi-indices β such that $|\beta| \leq \mu_2$, then we have the bound

$$(2.26) \quad \begin{aligned} & \|\text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2)\|_{L^2 \rightarrow L^2} \\ & \lesssim \sum_{|\alpha_1|+|\beta_1|+|\beta_2| \leq s'_d, |\alpha_1|+|\beta_1| \geq 1, |\beta_1| \leq \mu_1, 1 \leq |\beta_2| \leq \mu_2} h^{\frac{|\alpha_1|+|\beta_1|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\beta_2}^{(2)} \\ & \quad + h^{\mu_1/2} \log(1+h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1|=\mu_1} C_{\alpha_1, \beta_1}^{(1)} C_0^{(2)} \\ & \quad + h^{\mu_2/2} \sum_{|\alpha_1|+|\beta_1| \leq s_d, |\beta_1| \leq \mu_1, |\beta_2|=\mu_2} h^{\frac{|\alpha_1|+|\beta_1|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\beta_2}^{(2)}. \end{aligned}$$

Proof. The proof is similar to the proof of (2.19) and we keep the same notations replacing μ by μ_1 . The only difference is that a_2 is no longer C^∞ in x , so we have to bound the norms of the operators \mathcal{A}_0 and \mathcal{A}_{μ_1} differently. To bound the norm of \mathcal{A}_0 we will use Proposition 2.5. By (2.20), we have

$$\begin{aligned} & \|\mathcal{A}_0\|_{H_h^{k'} \rightarrow L^2} \\ & \lesssim \sum_{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2| \leq s'_d, |\beta_1| \leq \mu_1, |\beta_2| \leq \mu_2, |\alpha_1|+|\beta_1| \geq 1, |\alpha_2|+|\beta_2| \geq 1} h^{\frac{|\alpha_1|+|\beta_1|+|\alpha_2|+|\beta_2|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_2/2} \log(1+h^{-1}) \sum_{|\alpha_1|+|\beta_1| \leq s_d, |\alpha_2| \leq d, |\beta_1| \leq \mu_1, |\beta_2|=\mu_2} h^{\frac{|\alpha_1|+|\beta_1|}{2}} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_2/2} \log(1+h^{-1}) \sum_{|\alpha_1|+|\alpha_2| \leq d, \beta_1=0, |\beta_2|=\mu_2} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}. \end{aligned}$$

To bound the norm of the operator \mathcal{A}_{μ_1} we will use Propositions 2.3 and 2.4. We get

$$\begin{aligned} \|\mathcal{A}_{\mu_1}\|_{H_h^{k'} \rightarrow L^2} & \leq \left\| \text{Op}_h(a_{1,t}^{(\mu_1)}) \right\|_{L^2 \rightarrow L^2} \left\| \text{Op}_h(a_2) \right\|_{H_h^{k'} \rightarrow L^2} + \left\| \text{Op}_h(a_{1,t}^{(\mu_1)} a_2) \right\|_{H_h^{k'} \rightarrow L^2} \\ & \lesssim \log(1+h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1|=\mu_1} C_{\alpha_1, \beta_1}^{(1)} \sum_{|\alpha_2|+|\beta_2| \leq s_d, |\beta_2| \leq \mu_2} h^{\frac{|\alpha_2|+|\beta_2|}{2}} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + h^{\mu_2/2} \log^2(1+h^{-1}) \sum_{|\alpha_1| \leq d, |\beta_1|=\mu_1} C_{\alpha_1, \beta_1}^{(1)} \sum_{|\alpha_2| \leq d, |\beta_2|=\mu_2} C_{\alpha_2, \beta_2}^{(2)} \\ & \quad + \log(1+h^{-1}) \sum_{|\alpha_1|+|\alpha_2| \leq d, |\beta_1|=\mu_1, \beta_2=0} C_{\alpha_1, \beta_1}^{(1)} C_{\alpha_2, \beta_2}^{(2)}. \end{aligned}$$

Now (2.25) follows from the above bounds and (2.24) by induction in μ_1 . The bound (2.26) can be derived from (2.22) in the same way. \square

It is well-known that one can define h - Ψ DOs with C^∞ smooth symbols on an arbitrary compact manifold without boundary using the definition on the Euclidean space (e.g. see Chapter 18 of [4]). Roughly speaking, an h - Ψ DO on a manifold of dimension $d-1$, say Γ , is a finite sum of h - Ψ DOs on \mathbb{R}^{d-1} with symbols compactly supported in x . In other words, studying h - Ψ DOs on compact manifolds is reduced to studying h - Ψ DOs on the Euclidean space with symbols compactly supported in x . In the same way, we can define h - Ψ DOs on Γ with symbols

of low regularity in x . To be more precise, we cover Γ by a finite number of open sets U_j , $j = 1, \dots, J$, such that

$$\kappa_j : U_j \rightarrow \tilde{U}_j \subset \mathbb{R}^{d-1}$$

are diffeomorphisms. Then we can associate to κ_j a diffeomorphism

$$\mathcal{K}_j : T^*U_j \rightarrow T^*\tilde{U}_j \subset T^*\mathbb{R}^{d-1}$$

such that

$$\mathcal{K}_j(x, \xi) = (\kappa_j(x), {}^t\kappa'_j(x)\xi).$$

Let $\psi_j \in C^\infty(U_j)$ be such that $\sum_{j=1}^J \psi_j = 1$. If a is a function on $T^*\Gamma$, we can write it in the form $a = \sum_{j=1}^J a_j$, where $a_j = a\psi_j$. Define the function $a_j \circ \mathcal{K}_j^{-1}$ on $T^*\tilde{U}_j$ by

$$(a_j \circ \mathcal{K}_j^{-1})(y, \eta) = a_j(\mathcal{K}_j^{-1}(y, \eta)).$$

Then we define the operators $\text{Op}_h(a_j \circ \mathcal{K}_j^{-1})$ as in the beginning of the section. We now define $\text{Op}_h(a)$ in terms of these operators as follows

$$\text{Op}_h(a)f = \sum_{j=1}^J \sum_{\ell=1}^J \text{Op}_h(\psi_j a) \psi_\ell f = \sum_{j=1}^J \sum_{\ell=1}^J \left(\text{Op}_h(a_j \circ \mathcal{K}_j^{-1})((\psi_\ell f) \circ \kappa_\ell^{-1}) \right) \circ \kappa_j.$$

We will say that $a \in S^k(\Gamma)$ if $a_j \circ \mathcal{K}_j^{-1} \in S^k$ for all j . Similarly, we can extend the other classes of symbols on $T^*\mathbb{R}^{d-1}$ above to symbols on $T^*\Gamma$. It is then clear that the above propositions extend to h - Ψ DOs on Γ with the spaces L^2 and H_h^k replaced by $L^2(\Gamma)$ and $H_h^k(\Gamma)$, respectively. To simplify the notations in the formulas that follow, we will omit the diffeomorphisms and will identify T^*U_j with $T^*\tilde{U}_j$.

3. A PRIORI ESTIMATES

In this section we will prove a priori estimates for the solution to the equation

$$(3.1) \quad \begin{cases} (h^2 \nabla c(x) \nabla + zn(x))u = hv & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma, \end{cases}$$

where $c, n \in L^\infty(\Omega)$ are real-valued functions satisfying $c(x) \geq b_0$, $n(x) \geq b_0$ for some constant $b_0 > 0$, $0 < h \ll 1$ is a semiclassical parameter and $z \in Z^+ \cup Z^-$, where

$$Z^\pm = \{z \in \mathbb{C} : |z| = 1, \pm \text{Re } z \geq 0\}.$$

Set $\theta = |\text{Im } z|$ if $z \in Z^+$ and $\theta = 1$ if $z \in Z^-$. We will prove the following

Theorem 3.1. *Suppose that $c, n \in C^1(\bar{\Omega}_\delta)$ for some $0 < \delta \ll 1$. Let $\theta \geq h$ and let $u \in H^2(\Omega)$ satisfy equation (3.1). Then the function $g = h\partial_\nu u|_\Gamma$ satisfies the estimate*

$$(3.2) \quad \|g\|_{L^2(\Gamma)} \lesssim h^{1/2} \theta^{-1/2} \|v\|_{L^2(\Omega)}.$$

Proof. We will first prove the following

Lemma 3.2. *We have the estimate*

$$(3.3) \quad \|u\|_{H_h^1(\Omega)} \lesssim h\theta^{-1} \|v\|_{L^2(\Omega)}.$$

Proof. By the Green formula we have

$$(3.4) \quad \langle znu - hv, u \rangle_{L^2(\Omega)} = \langle -h^2 \nabla c \nabla u, u \rangle_{L^2(\Omega)} = \int_{\Omega} c |h \nabla u|^2.$$

Taking the imaginary part we get the identity

$$\operatorname{Im} z \left\| n^{1/2} u \right\|_{L^2(\Omega)}^2 = \operatorname{Im} \langle hv, u \rangle_{L^2(\Omega)},$$

which implies

$$(3.5) \quad \|u\|_{L^2(\Omega)} \lesssim h |\operatorname{Im} z|^{-1} \|v\|_{L^2(\Omega)}.$$

Taking the real part of (3.4) we get

$$(3.6) \quad \begin{aligned} \int_{\Omega} c |h \nabla u|^2 &= \operatorname{Re} z \langle nu, u \rangle_{L^2(\Omega)} - \operatorname{Re} \langle hv, u \rangle_{L^2(\Omega)} \\ &\leq (\operatorname{Re} z + \varepsilon) \langle nu, u \rangle_{L^2(\Omega)} + \mathcal{O}_{\varepsilon}(h^2) \|v\|_{L^2(\Omega)}^2 \end{aligned}$$

for every $0 < \varepsilon \leq 1$. In particular, (3.6) implies

$$(3.7) \quad \int_{\Omega} |h \nabla u|^2 \lesssim \|u\|_{L^2(\Omega)}^2 + h^2 \|v\|_{L^2(\Omega)}^2.$$

By (3.5) and (3.7) we obtain

$$(3.8) \quad \|u\|_{H_h^1(\Omega)} \lesssim h |\operatorname{Im} z|^{-1} \|v\|_{L^2(\Omega)}.$$

When $|\operatorname{Re} z| \leq 1/2$ we have $|\operatorname{Im} z| \geq 1/2$, so in this case (3.3) follows from (3.8). When $\operatorname{Re} z \geq 1/2$ we have $|\operatorname{Im} z| = \theta$, so in this case (3.3) again follows from (3.8). When $\operatorname{Re} z \leq -1/2$ we have $\theta = 1$, so in this case (3.3) follows from (3.6) with $\varepsilon = 1/4$. \square

Let $\mathcal{V} \subset \mathbb{R}^d$ be a small open domain such that $\mathcal{V}^0 := \mathcal{V} \cap \Gamma \neq \emptyset$. Let $(x_1, x') \in \mathcal{V}^+ := \mathcal{V} \cap \Omega$, $0 < x_1 \ll 1$, $x' = (x_2, \dots, x_d) \in \mathcal{V}^0$, be the local normal geodesic coordinates near the boundary. In these coordinates the principal symbol of the operator $-\Delta$ is equal to $\xi_1^2 + r(x, \xi')$, where (ξ_1, ξ') are the dual variables to (x_1, x') , and r is a homogeneous polynomial of order two and satisfies $C_1 |\xi'|^2 \leq r \leq C_2 |\xi'|^2$ with some constants $C_1, C_2 > 0$. Therefore, the principal symbol of the positive Laplace-Beltrami operator on Γ is equal to $r_0(x', \xi') = r(0, x', \xi')$.

Let $\mathcal{V}_1 \subset \mathcal{V}$ be a small open domain such that $\mathcal{V}_1^0 := \mathcal{V}_1 \cap \Gamma \neq \emptyset$. Choose a function $\psi \in C_0^\infty(\mathcal{V})$, $0 \leq \psi \leq 1$, such that $\psi = 1$ on \mathcal{V}_1 . Then the function $u^b := \psi u$ satisfies the equation

$$(3.9) \quad \begin{cases} (h^2 \nabla c(x) \nabla + zn) u^b = h v^b & \text{in } \Omega, \\ u^b = 0 & \text{on } \Gamma, \end{cases}$$

where $v^b = \psi v + h[\nabla c \nabla, \psi]u$ satisfies

$$(3.10) \quad \|v^b\|_{L^2(\Omega)} \lesssim \|v\|_{L^2(\Omega)} + \|u\|_{H_h^1(\Omega)}.$$

We will now write the operator $-h^2 \nabla c(x) \nabla$ in the coordinates $x = (x_1, x')$. Denote $\mathcal{D}_{x_j} = -ih \partial_{x_j}$. We can write

$$(3.11) \quad -h^2 \nabla c(x) \nabla = c(x) \mathcal{D}_{x_1}^2 + c(x) r(x, \mathcal{D}_{x'}) + h \mathcal{R}(x, \mathcal{D}_x),$$

where \mathcal{R} is a first-order differential operator with L^∞ coefficients. Denote

$$\langle f, g \rangle_0 = \int f \bar{g} dx', \quad \|f\|_0^2 = \int |f|^2 dx',$$

and introduce the function

$$F(x_1) = \left\| \mathcal{D}_{x_1} u^b \right\|_0^2 - \left\langle r(x_1, \cdot, \mathcal{D}_{x'} u^b), u^b \right\rangle_0 + \operatorname{Re} z \left\langle \tilde{n}(x_1, \cdot) u^b, u^b \right\rangle_0,$$

where $\tilde{n} = c^{-1}n \in C^1(\overline{\Omega}_\delta)$. Since $u^b|_{x_1=0} = 0$, we have

$$(3.12) \quad F(0) = \left\| \mathcal{D}_{x_1} u^b|_{x_1=0} \right\|_0^2.$$

On the other hand,

$$(3.13) \quad F(0) = - \int_0^\delta F'(x_1) dx_1$$

for some constant $\delta > 0$, where F' denotes the first derivative with respect to x_1 . We will now bound $F(0)$ from above. To this end we will compute $F'(x_1)$ using that u^b satisfies (3.9) together with (3.11). We have

$$\begin{aligned} F'(x_1) &= -2\operatorname{Re} \left\langle (\mathcal{D}_{x_1}^2 + r - \operatorname{Re} z \tilde{n}) u^b, \partial_{x_1} u^b \right\rangle_0 - \left\langle (r' - \operatorname{Re} z \tilde{n}') u^b, u^b \right\rangle_0 \\ &= 2h^{-1} \operatorname{Im} \left\langle (-h^2 \nabla c(x) \nabla + \operatorname{Re} z n - h\mathcal{R}) u^b, c^{-1} \mathcal{D}_{x_1} u^b \right\rangle_0 \\ &\quad - \left\langle (r' - \operatorname{Re} z \tilde{n}') u^b, u^b \right\rangle_0 \\ &= 2\operatorname{Im} \left\langle (v^b - ih^{-1} \operatorname{Im} z n u^b - \mathcal{R} u^b), c^{-1} \mathcal{D}_{x_1} u^b \right\rangle_0 \\ &\quad - \left\langle (r' - \operatorname{Re} z \tilde{n}') u^b, u^b \right\rangle_0. \end{aligned}$$

Hence

$$|F'(x_1)| \lesssim h\theta^{-1} \|v^b\|_0^2 + \theta h^{-1} \sum_{\ell=0}^1 \|\mathcal{D}_{x_1}^\ell u^b\|_0^2 + \sum_{|\alpha| \leq 1} \|\mathcal{D}_x^\alpha u^b\|_0^2.$$

Using this estimate together with (3.10), (3.13) and Lemma 3.2 we obtain

$$(3.14) \quad \begin{aligned} F(0) &\leq \int_0^\delta |F'(x_1)| dx_1 \lesssim h\theta^{-1} \|v\|_{L^2(\Omega)}^2 + (1 + \theta h^{-1}) \|u\|_{H_h^1(\Omega)}^2 \\ &\lesssim (h\theta^{-1} + h^2\theta^{-2}) \|v\|_{L^2(\Omega)}^2 \lesssim h\theta^{-1} \|v\|_{L^2(\Omega)}^2. \end{aligned}$$

Observe now that

$$\mathcal{D}_{x_1} u^b|_{x_1=0} = \psi_0 \mathcal{D}_{x_1} u|_{x_1=0}, \quad \mathcal{D}_{x'} u|_{x_1=0} = 0,$$

where $\psi_0 = \psi|_{x_1=0}$ is supported in \mathcal{V}^0 and such that $\psi_0 = 1$ on \mathcal{V}_1^0 . Therefore, by (3.12) and (3.14),

$$\|\psi_0 \mathcal{D}_{x_1} u|_{x_1=0}\|_0 \lesssim h^{1/2} \theta^{-1/2} \|v\|_{L^2(\Omega)},$$

which clearly implies

$$(3.15) \quad \|\psi_0 g\|_0 \lesssim h^{1/2} \theta^{-1/2} \|v\|_{L^2(\Omega)}.$$

Since Γ is compact, there exist a finite number of smooth functions ψ_i , $0 \leq \psi_i \leq 1$, $i = 1, \dots, I$, such that $1 = \sum_{i=1}^I \psi_i$ and (3.15) holds with ψ_0 replaced by each ψ_i . Therefore, the estimate (3.2) is obtained by summing up all such estimates (3.15). \square

4. APPROXIMATION OF THE DIRICHLET-TO-NEUMANN MAP

Given $f \in H^1(\Gamma)$ let u solve the equation

$$(4.1) \quad \begin{cases} (h^2 \nabla c(x) \nabla + zn(x))u = 0 & \text{in } \Omega, \\ u = f & \text{on } \Gamma, \end{cases}$$

where c, n, h, z are as in previous section. We define the semiclassical Dirichlet-to-Neumann map

$$\mathcal{N}(h, z) : H^1(\Gamma) \rightarrow L^2(\Gamma)$$

by

$$\mathcal{N}(h, z)f := -ih\partial_\nu u|_\Gamma.$$

We would like to approximate $\mathcal{N}(h, z)$ by an h - Ψ DO on Γ similarly to the C^∞ case. To this end, introduce the function

$$\rho(x', \xi', z) = \sqrt{-r_0(x', \xi') + z\tilde{n}_0(x')}, \quad \text{Im } \rho > 0,$$

where $\tilde{n}_0 = \tilde{n}|_\Gamma$, $\tilde{n} := c^{-1}n$, and r_0 is the principal symbol of the Laplace-Beltrami operator on Γ written in the coordinates $(x', \xi') \in T^*\Gamma$. Let $\eta \in C^\infty(T^*\Gamma)$ be such that $\eta = 1$ for $r_0 \leq C_0$, $\eta = 0$ for $r_0 \geq 2C_0$, where $C_0 > 0$ does not depend on h . It is easy to see (e.g. see Lemma 3.1 of [14]) that taking C_0 big enough we can arrange

$$(4.2) \quad C_1\theta^{1/2} \leq |\rho| \leq C_2, \quad \text{Im } \rho \geq C_3|\theta||\rho|^{-1} \geq C_4|\theta|$$

for $(x', \xi') \in \text{supp } \eta$, and

$$(4.3) \quad |\rho| \geq \text{Im } \rho \geq C_5|\xi'|$$

for $(x', \xi') \in \text{supp } (1 - \eta)$ with some constants $C_j > 0$. In other words, $\text{supp } (1 - \eta)$ is contained in the elliptic region of the boundary value problem (4.1). Our goal is to prove the following

Theorem 4.1. *Let $\theta \geq h^{2/5}$. Suppose that $c, n \in C^1(\Omega_\delta)$ and $\tilde{n}_0 \in C^\mu(\Gamma)$ with an integer $\mu \geq 2$. Then for every $f \in H^1(\Gamma)$ we have the estimate*

$$(4.4) \quad \begin{aligned} \|\mathcal{N}(z, h)f - \text{Op}_h(\rho)f\|_{L^2(\Gamma)} &\lesssim h^{3/5}\|f\|_{H_h^1(\Gamma)} \\ &+ h\theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right) \|f\|_{H_h^1(\Gamma)}. \end{aligned}$$

Suppose in addition that $c \equiv 1$. Then we have the better estimate

$$(4.5) \quad \|\mathcal{N}(z, h)f - \text{Op}_h(\rho + hq)f\|_{L^2(\Gamma)} \lesssim h\theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right) \|f\|_{H_h^{-1}(\Gamma)},$$

where $q \in C^\infty(T^*\Gamma)$ is independent of z, h and the function n .

Proof. We will build a parametrix for the solutions of the equation (4.1). This parametrix will not be as good as that one built in the C^∞ case, but will suffice for the proof of the above estimates. Let (x_1, x') be the local normal geodesic coordinates near the boundary. Take a function $\chi \in C^\infty(T^*\Gamma)$, $0 \leq \chi \leq 1$, such that $\pi_{x'}(\text{supp } \chi) \subset \mathcal{V}^0$, where $\pi_{x'} : T^*\Gamma \rightarrow \Gamma$ denotes the projection $(x', \xi') \rightarrow x'$. Moreover, we require that either χ is of compact support or $\chi \in S^0(\Gamma)$ with $\text{supp } \chi \subset \text{supp } (1 - \eta)$. We will be looking for a parametrix of the solution to equation (4.1) in the form

$$\tilde{u} = \phi(x_1/\delta)(2\pi h)^{-d+1} \int \int e^{\frac{i}{h}(\langle y', \xi' \rangle + \varphi(x, \xi', z))} a(x, \xi', z) f(y') d\xi' dy'$$

where $\phi \in C_0^\infty(\mathbb{R})$, $\phi(t) = 1$ for $|t| \leq 1/2$, $\phi(t) = 0$ for $|t| \geq 1$, $0 < \delta \ll 1$ being a parameter independent of h and z . We require that \tilde{u} satisfies the boundary condition $\tilde{u} = \text{Op}_h(\chi)f$ on $x_1 = 0$. We take $a = \chi(x', \xi')$. Furthermore, we choose the phase function in the form

$$\varphi = -\langle x', \xi' \rangle + x_1 \rho.$$

Then it is easy to see that φ satisfies the following eikonal equation

$$(4.6) \quad (\partial_{x_1} \varphi)^2 + r(x, \nabla_{x'} \varphi) - z \tilde{n}(x) = x_1 \Phi,$$

where $|\Phi|$ is bounded as $x_1 \rightarrow 0$. By assumption, we have

$$\tilde{n}(x) = \tilde{n}_0(x') + x_1 \tilde{n}^\sharp(x), \quad \tilde{n}_0 \in C^\mu, \tilde{n}^\sharp \in L^\infty.$$

Furthermore, it is clear that the function r can be written in the form

$$r(x, \xi') = \langle R(x) \xi', \xi' \rangle$$

where R is a C^∞ smooth $(d-1) \times (d-1)$ matrix-valued function. Hence we can write

$$R(x) = R_0(x') + x_1 R^\sharp(x)$$

where $R_0 = R|_{x_1=0}$ and R^\sharp are C^∞ smooth functions. Thus we obtain

$$\Phi = -z \tilde{n}^\sharp + \langle R^\sharp(x) \xi', \xi' \rangle - 2 \langle (R_0(x') + x_1 R^\sharp(x)) \xi', \nabla_{x'} \rho \rangle + x_1 \langle R(x) \nabla_{x'} \rho, \nabla_{x'} \rho \rangle.$$

Observe now that

$$(4.7) \quad \nabla_{x'} \rho = (2\rho)^{-1} (-\nabla_{x'} r_0 + z \nabla_{x'} \tilde{n}_0).$$

Hence we can write the function Φ in the form

$$\begin{aligned} \Phi &= -z \tilde{n}^\sharp + \langle R^\sharp(x) \xi', \xi' \rangle + \rho^{-1} \langle (R_0(x') + x_1 R^\sharp(x)) \xi', \nabla_{x'} r_0 - z \nabla_{x'} \tilde{n}_0 \rangle \\ &\quad + x_1 (2\rho)^{-2} \langle R(x) (\nabla_{x'} r_0 - z \nabla_{x'} \tilde{n}_0), \nabla_{x'} r_0 - z \nabla_{x'} \tilde{n}_0 \rangle. \end{aligned}$$

Observe now that \tilde{u} satisfies the equation

$$(h^2 \nabla c(x) \nabla + z n(x)) \tilde{u} = h \tilde{v},$$

where the function \tilde{v} is of the form

$$\begin{aligned} \tilde{v} &= (2\pi h)^{-d+1} \int \int e^{\frac{i}{h} \langle y' - x', \xi' \rangle} e^{ix_1 \rho/h} A(x, \xi', h, z) f(y') d\xi' dy' \\ &= \text{Op}_h \left(e^{ix_1 \rho/h} A \right) f \end{aligned}$$

where

$$A = h^{-1} e^{-i\varphi/h} (h^2 \nabla c(x) \nabla + z n(x)) \left(\phi(x_1/\delta) e^{i\varphi/h} a \right).$$

To compute the function A observe that the operator $-\nabla c \nabla$ can be written in the coordinates (x_1, x') in the form

$$-\nabla c \nabla = c \mathcal{D}_{x_1}^2 + c \langle R(x) \mathcal{D}_{x'}, \mathcal{D}_{x'} \rangle + \langle Q(x), \mathcal{D}_x \rangle,$$

where $\mathcal{D}_{x_1} = -i\partial_{x_1}$, $\mathcal{D}_{x'} = -i\nabla_{x'}$, $\mathcal{D}_x = -i\nabla_x = (\mathcal{D}_{x_1}, \mathcal{D}_{x'})$, R is the smooth matrix-valued function as above, and $Q = (Q_1, \dots, Q_d)$ with scalar-valued functions $Q_j \in L^\infty$. Denote $\tilde{Q} = (Q_2, \dots, Q_d)$. Hence we can write

$$-h^2 \nabla c \nabla = c \mathcal{D}_{x_1}^2 + h Q_1(x) \mathcal{D}_{x_1} + c \langle R(x) \mathcal{D}_{x'}, \mathcal{D}_{x'} \rangle + h \langle \tilde{Q}(x), \mathcal{D}_{x'} \rangle,$$

where $\mathcal{D}_{x_1} = h \mathcal{D}_{x_1}$, $\mathcal{D}_{x'} = h \mathcal{D}_{x'}$. Therefore, the function A can be decomposed as $A_1 + \phi(x_1/\delta) A_2$, where

$$A_1 = e^{-i\varphi/h} [ch^{-1} \mathcal{D}_{x_1}^2 + Q_1(x) \mathcal{D}_{x_1}, \phi(x_1/\delta)] \left(e^{i\varphi/h} a \right)$$

$$\begin{aligned}
 &= -2ic\delta^{-1}\phi'(x_1/\delta)e^{-i\varphi/h}\mathcal{D}_{x_1}\left(e^{i\varphi/h}a\right) \\
 &\quad -hc\delta^{-2}\phi''(x_1/\delta)a - ihQ_1\delta^{-1}\phi'(x_1/\delta)a \\
 &\quad = -2ic\delta^{-1}\phi'(x_1/\delta)(a\partial_{x_1}\varphi + \mathcal{D}_{x_1}a) \\
 &\quad \quad -hc\delta^{-2}\phi''(x_1/\delta)a - ihQ_1\delta^{-1}\phi'(x_1/\delta)a \\
 &= -2ic\delta^{-1}\phi'(x_1/\delta)\chi\rho - hc\delta^{-2}\phi''(x_1/\delta)\chi - ihQ_1\delta^{-1}\phi'(x_1/\delta)\chi,
 \end{aligned}$$

and

$$\begin{aligned}
 A_2 &= -h^{-1}cx_1\Phi - 2ic(\partial_{x_1}\varphi\partial_{x_1}a + \langle\nabla_{x'}\varphi, \nabla_{x'}a\rangle) \\
 &\quad + ica(D_{x_1}^2 + \langle R(x)D_{x'}, D_{x'}\rangle)\varphi + hc(D_{x_1}^2 + \langle R(x)D_{x'}, D_{x'}\rangle)a \\
 &\quad \quad + Q_1(a\partial_{x_1}\varphi + \mathcal{D}_{x_1}a) + \langle\tilde{Q}, \nabla_{x'}\varphi\rangle a + \langle\tilde{Q}, \mathcal{D}_{x'}a\rangle \\
 &= -h^{-1}cx_1\Phi - 2ic(-\xi' + x_1\nabla_{x'}\rho, \nabla_{x'}\chi) - icx_1\chi\langle R(x)\nabla_{x'}, \nabla_{x'}\rangle\rho \\
 &\quad - hc\langle R(x)\nabla_{x'}, \nabla_{x'}\rangle\chi + Q_1\chi\rho + \langle\tilde{Q}, -\xi' + x_1\nabla_{x'}\rho\rangle\chi - ih\langle\tilde{Q}, \nabla_{x'}\chi\rangle \\
 &\quad = -h^{-1}cx_1\Phi + A_3.
 \end{aligned}$$

In view of (4.7), the function A_3 can be written in the form

$$\begin{aligned}
 A_3 &= 2ic\langle\xi', \nabla_{x'}\chi\rangle + ix_1\rho^{-1}c\langle\nabla_{x'}r_0 - z\nabla_{x'}\tilde{n}_0, \nabla_{x'}\chi\rangle \\
 &\quad - hc\langle R(x)\nabla_{x'}, \nabla_{x'}\rangle\chi + Q_1\chi\rho - \langle\tilde{Q}, \xi'\rangle\chi - ih\langle\tilde{Q}, \nabla_{x'}\chi\rangle \\
 &\quad - x_1(2\rho)^{-1}\langle\tilde{Q}, \nabla_{x'}r_0 - z\nabla_{x'}\tilde{n}_0\rangle\chi - icx_1\chi\langle R(x)\nabla_{x'}, \nabla_{x'}\rangle\rho.
 \end{aligned}$$

The last term can also be expressed in terms of negative powers of ρ . Indeed, if $2 \leq \ell, j \leq d$, we have

$$\begin{aligned}
 \partial_{x_\ell}\partial_{x_j}\rho &= \partial_{x_\ell}((2\rho)^{-1}(-\partial_{x_j}r_0 + z\partial_{x_j}\tilde{n}_0)) \\
 &= (2\rho)^{-1}(-\partial_{x_\ell}\partial_{x_j}r_0 + z\partial_{x_\ell}\partial_{x_j}\tilde{n}_0) + 2^{-1}\rho^{-2}(\partial_{x_j}r_0 - z\partial_{x_j}\tilde{n}_0)\partial_{x_\ell}\rho \\
 &= (2\rho)^{-1}(-\partial_{x_\ell}\partial_{x_j}r_0 + z\partial_{x_\ell}\partial_{x_j}\tilde{n}_0) - 2^{-2}\rho^{-3}(\partial_{x_j}r_0 - z\partial_{x_j}\tilde{n}_0)(\partial_{x_\ell}r_0 - z\partial_{x_\ell}\tilde{n}_0).
 \end{aligned}$$

It is easy to see from the above expressions that we have the following

Lemma 4.2. *The functions Φ and A_3 are of the form*

$$(4.8) \quad \Phi = b_1 + b_2\rho^{-1} + x_1\rho^{-1}(b_3 + b_4\rho^{-1}),$$

$$(4.9) \quad A_3 = b_5 + hb_6 + b_7\rho + x_1\rho^{-1}(b_8 + b_9\rho^{-1} + b_{10}\rho^{-2}),$$

where b_j are functions on $T^*\Gamma$ independent of h, ρ and depending on x_1 and z . Moreover, each function b_j is of the form

$$(4.10) \quad b_j = \sum_{|\alpha| \leq \ell_j} \omega_{j,\alpha}(x, z)\chi_{j,\alpha}(x', \xi')\xi'^{\alpha},$$

where $\chi_{j,\alpha}$ are either χ or derivatives of χ , $\omega_{j,\alpha} \in L^\infty$ uniformly in z , and $0 \leq \ell_j \leq 4$. More precisely, we have $\ell_1 = 2$, $\ell_2 = \ell_3 = 3$, $\ell_4 = 4$, $\ell_5 = 1$, $\ell_6 = \ell_7 = 0$, $\ell_8 = 2$, $\ell_9 = 3$, $\ell_{10} = 4$.

Using this lemma we will prove the following

Proposition 4.3. *Let $0 < x_1 \leq \delta$. If χ is of compact support, there exists a constant $C > 0$ such that, given any $N \geq 0$ independent of h , we have the estimate*

$$(4.11) \quad \left\| \left(\text{Op}_h \left(e^{ix_1\rho/h} A_2 \right) f \right) (x_1, \cdot) \right\|_{L^2(\Gamma)} \\ \lesssim \left(1 + h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu} \right) \theta^{-3/2} e^{-Cx_1\theta/2h} \|f\|_{H_h^{-N}(\Gamma)}.$$

If $\text{supp } \chi \subset \text{supp}(1 - \eta)$, given any $0 < \epsilon < 1/2$ independent of h , we have the estimate

$$(4.12) \quad \left\| \left(\text{Op}_h \left(e^{ix_1\rho/h} A_2 \right) f \right) (x_1, \cdot) \right\|_{L^2(\Gamma)} \lesssim h^{(1-3\epsilon)/2} x_1^{-1/2+\epsilon} \|f\|_{H_h^1(\Gamma)}.$$

Proof. Clearly, $\tilde{\chi}_{j,\alpha} := \chi_{j,\alpha} \xi^{\prime\alpha} \in C^\infty(T^*\Gamma)$ is compactly supported if so is χ , and $\tilde{\chi}_{j,\alpha} \in S^{|\alpha|}(\Gamma)$, $\text{supp } \tilde{\chi}_{j,\alpha} \subset \text{supp}(1 - \eta)$ if $\chi \in S^0(\Gamma)$, $\text{supp } \chi \subset \text{supp}(1 - \eta)$. In view of Lemma 4.2 we can write

$$\begin{aligned} \text{Op}_h \left(e^{ix_1\rho/h} A_2 \right) &= -h^{-1} x_1 \sum_{|\alpha| \leq 2} c\omega_{1,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \tilde{\chi}_{1,\alpha} \right) \\ &\quad - h^{-1} x_1 \sum_{|\alpha| \leq 3} c\omega_{2,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-1} \tilde{\chi}_{2,\alpha} \right) \\ &\quad - h^{-1} x_1^2 \sum_{|\alpha| \leq 3} c\omega_{3,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-1} \tilde{\chi}_{3,\alpha} \right) \\ &\quad - h^{-1} x_1^2 \sum_{|\alpha| \leq 4} c\omega_{4,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-2} \tilde{\chi}_{4,\alpha} \right) \\ &+ \sum_{|\alpha| \leq 1} \omega_{5,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \tilde{\chi}_{5,\alpha} \right) + h\omega_{6,0} \text{Op}_h \left(e^{ix_1\rho/h} \tilde{\chi}_{6,0} \right) + \omega_{7,0} \text{Op}_h \left(e^{ix_1\rho/h} \rho \tilde{\chi}_{7,0} \right) \\ &\quad + x_1 \sum_{|\alpha| \leq 2} \omega_{8,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-1} \tilde{\chi}_{8,\alpha} \right) \\ &\quad + x_1 \sum_{|\alpha| \leq 3} \omega_{9,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-2} \tilde{\chi}_{9,\alpha} \right) \\ &\quad + x_1 \sum_{|\alpha| \leq 4} \omega_{10,\alpha} \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-3} \tilde{\chi}_{10,\alpha} \right). \end{aligned}$$

Hence

$$\begin{aligned} \left\| \text{Op}_h \left(e^{ix_1\rho/h} A_2 \right) f \right\| &\lesssim \sum_{\ell=1}^2 \sum_{k=\ell-1}^{\ell} h^{-1} x_1^\ell \sum_{|\alpha| \leq k+2} \left\| \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-k} \tilde{\chi}_{j(k,\ell),\alpha} \right) f \right\| \\ &+ \sum_{|\alpha| \leq 1} \left\| \text{Op}_h \left(e^{ix_1\rho/h} \tilde{\chi}_{5,\alpha} \right) f \right\| + h \left\| \text{Op}_h \left(e^{ix_1\rho/h} \tilde{\chi}_{6,0} \right) f \right\| + \left\| \text{Op}_h \left(e^{ix_1\rho/h} \rho \tilde{\chi}_{7,0} \right) f \right\| \\ &\quad + x_1 \sum_{k=1}^3 \sum_{|\alpha| \leq k+1} \left\| \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-k} \tilde{\chi}_{7+k,\alpha} \right) f \right\|. \end{aligned}$$

It is easy to see now that the proposition follows from

Proposition 4.4. *Let $k \geq 0$ be an integer or $k = -1$ and let $\ell \geq 0$. If $\tilde{\chi} \in C^\infty(T^*\Gamma)$ is of compact support, there exists a constant $C > 0$ such that, given any $N \geq 0$ independent of h , we have the bound*

$$(4.13) \quad \begin{aligned} & \left\| \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-k} \tilde{\chi} \right) \right\|_{H_h^{-N}(\Gamma) \rightarrow L^2(\Gamma)} \\ & \lesssim x_1^{-\ell} h^\ell \theta^{-\ell - \tilde{k}/2} \left(1 + h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu} \right) e^{-Cx_1\theta/2h}, \end{aligned}$$

where $\tilde{k} = 0$ if $k = -1$ and $\tilde{k} = k$ if $k \geq 0$. If $\tilde{\chi} \in S^p(\Gamma)$, $\text{supp } \tilde{\chi} \subset \text{supp}(1 - \eta)$, we have the bound

$$(4.14) \quad \left\| \text{Op}_h \left(e^{ix_1\rho/h} \rho^{-k} \tilde{\chi} \right) \right\|_{H_h^1(\Gamma) \rightarrow L^2(\Gamma)} \lesssim h^\ell x_1^{-\ell},$$

provided $p < k + \ell + 1$.

Proof. We will first prove the following

Lemma 4.5. *Let k and \tilde{k} be as in Proposition 4.4. Then, there exists a constant $C > 0$ such that we have the bounds*

$$(4.15) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta \left(e^{ix_1\rho/h} \rho^{-k} \right) \right| \leq C_\alpha \theta^{-\tilde{k}/2 - |\alpha| - |\beta|} e^{-Cx_1\theta/h} \quad \text{on } \text{supp } \eta,$$

$$(4.16) \quad \left| \partial_{\xi'}^\alpha \left(e^{ix_1\rho/h} \rho^{-k} \right) \right| \leq C_\alpha |\xi'|^{-k - |\alpha|} e^{-Cx_1|\xi'|/h} \quad \text{on } \text{supp}(1 - \eta),$$

for all multi-indices α and β such that $|\beta| \leq \mu$.

Proof. It follows from Lemma 3.2 of [14] that, if $k \geq 1$ is an integer or $k = -1$, we have the bounds

$$(4.17) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta (\rho^{-k}) \right| \leq \begin{cases} C_\alpha |\rho|^{-k - 2|\alpha| - 2|\beta|} & \text{on } \text{supp } \eta, \\ C_\alpha |\rho|^{-k - |\alpha|} & \text{on } \text{supp}(1 - \eta), \end{cases}$$

for all multi-indices α and β such that $|\beta| \leq \mu$. By (4.17) together with (4.2) and (4.3) we get that the function ρ^{-k} , $k \geq 1$ being integer, satisfies the bounds

$$(4.18) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta (\rho^{-k}) \right| \leq \begin{cases} C_\alpha \theta^{-k/2 - |\alpha| - |\beta|} & \text{on } \text{supp } \eta, \\ C_\alpha |\xi'|^{-k - |\alpha|} & \text{on } \text{supp}(1 - \eta), \end{cases}$$

while the function ρ satisfies

$$(4.19) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta \rho \right| \leq \begin{cases} C_\alpha |\rho|^{1 - 2|\alpha| - 2|\beta|} & \text{on } \text{supp } \eta, \\ C_\alpha |\xi'|^{1 - |\alpha|} & \text{on } \text{supp}(1 - \eta), \end{cases}$$

for all multi-indices α and β such that $|\beta| \leq \mu$. Clearly, the bounds (4.18) hold with $k = -1$, provided $|\alpha| + |\beta| \geq 1$. Therefore, to prove the lemma it suffices to prove the bounds (4.15) and (4.16) with $k = 0$. This in turn follows from Lemma 4.2 of [17], but we will sketch the proof here for the sake of completeness.

Let us see that the functions

$$c_{\alpha,\beta} = e^{-ix_1\rho/h} \partial_{\xi'}^\alpha \partial_{x'}^\beta \left(e^{ix_1\rho/h} \right), \quad |\alpha| + |\beta| \geq 1, \quad |\beta| \leq \mu,$$

satisfy the bounds

$$(4.20) \quad \left| \partial_{\xi'}^{\alpha'} \partial_{x'}^{\beta'} c_{\alpha,\beta} \right| \lesssim \sum_{j=1}^{|\alpha| + |\beta| + |\alpha'| + |\beta'|} \left(\frac{x_1}{h|\rho|} \right)^j |\rho|^{-2(|\alpha| + |\beta| + |\alpha'| + |\beta'| - j)}$$

on $\text{supp } \eta$, and

$$(4.21) \quad \left| \partial_{\xi'}^{\alpha'} \partial_{x'}^{\beta'} c_{\alpha, \beta} \right| \lesssim \sum_{j=1}^{|\alpha|+|\beta|+|\alpha'|+|\beta'|} \left(\frac{x_1}{h} \right)^j |\xi'|^{-(|\alpha|+|\alpha'|-j)}$$

on $\text{supp}(1 - \eta)$, for all multi-indices α' and β' such that $|\beta| + |\beta'| \leq \mu$. We will proceed by induction in $|\alpha| + |\beta|$. Let α_1 and β_1 be multi-indices such that $|\alpha_1| + |\beta_1| = 1$ and observe that

$$c_{\alpha+\alpha_1, \beta+\beta_1} = \partial_{\xi'}^{\alpha_1} \partial_{x'}^{\beta_1} c_{\alpha, \beta} + ix_1 h^{-1} c_{\alpha, \beta} \partial_{\xi'}^{\alpha_1} \partial_{x'}^{\beta_1} \rho.$$

More generally, we have

$$(4.22) \quad \partial_{\xi'}^{\alpha'} \partial_{x'}^{\beta'} c_{\alpha+\alpha_1, \beta+\beta_1} = \partial_{\xi'}^{\alpha_1+\alpha'} \partial_{x'}^{\beta_1+\beta'} c_{\alpha, \beta} + ix_1 h^{-1} \partial_{\xi'}^{\alpha'} \partial_{x'}^{\beta'} \left(c_{\alpha, \beta} \partial_{\xi'}^{\alpha_1} \partial_{x'}^{\beta_1} \rho \right).$$

By (4.19) and (4.22), it is easy to see that if (4.20) and (4.21) hold for $c_{\alpha, \beta}$, they hold for $c_{\alpha+\alpha_1, \beta+\beta_1}$ as well.

Using (4.20) together with (4.2) we obtain

$$\begin{aligned} \left| e^{ix_1 \rho/h} c_{\alpha, \beta} \right| &\lesssim \sum_{j=1}^{|\alpha|+|\beta|} \left(\frac{x_1}{h|\rho|} \right)^j |\rho|^{-2(|\alpha|+|\beta|-j)} e^{-2C\theta x_1(h|\rho|)^{-1}} \\ &\lesssim \sum_{j=1}^{|\alpha|+|\beta|} \theta^{-j} |\rho|^{-2(|\alpha|+|\beta|-j)} e^{-C x_1 \theta/h} \lesssim \theta^{-|\alpha|-|\beta|} e^{-C x_1 \theta/h}. \end{aligned}$$

Similarly, by (4.21) we obtain

$$\left| e^{ix_1 \rho/h} c_{\alpha, \beta} \right| \lesssim \sum_{j=1}^{|\alpha|+|\beta|} \left(\frac{x_1}{h} \right)^j |\xi'|^{-|\alpha|+j} e^{-2C x_1 |\xi'|/h} \lesssim |\xi'|^{-|\alpha|} e^{-C x_1 |\xi'|/h}.$$

□

In view of the inequality

$$e^{-C x_1 \theta/2h} \lesssim x_1^{-\ell} h^\ell \theta^{-\ell},$$

the bound (4.13) follows from Proposition 2.4 and (4.15). Since

$$e^{-C x_1 |\xi'|/h} \lesssim h^\ell x_1^{-\ell} |\xi'|^{-\ell},$$

the bound (4.14) follows from Proposition 2.3 and (4.16). □

Since the function A_1 is supported in $\delta/2 \leq x_1 \leq \delta$, the next lemma is an immediate consequence of Lemma 4.5 with $k = -1$.

Lemma 4.6. *For every $m \geq 0$ we have the estimates*

$$(4.23) \quad \left| \partial_{\xi'}^\alpha \left(e^{ix_1 \rho/h} A_1 \right) \right| \leq \begin{cases} C_{\alpha, m} h^m \theta^{-m-|\alpha|} & \text{on } \text{supp } \eta, \\ C_{\alpha, m} h^m |\xi'|^{-m-|\alpha|} & \text{on } \text{supp}(1 - \eta), \end{cases}$$

for all multi-indices α with constants $C_{\alpha, m} > 0$ independent of x_1 , θ , z and h .

This lemma together with Proposition 2.3 imply the following

Proposition 4.7. *Given any $N \geq 0$ independent of h , we have the estimate*

$$(4.24) \quad \left\| \left(\text{Op}_h \left(e^{ix_1 \rho/h} A_1 \right) f \right) (x_1, \cdot) \right\|_{L^2(\Gamma)} \lesssim h \|f\|_{H_h^{-N}(\Gamma)}.$$

Let u satisfy equation (4.1) with $u|_{\Gamma} = \text{Op}_h(\chi)f$. Then $u - \tilde{u}$ satisfies equation (3.1) with v replaced by \tilde{v} . Therefore, taking into account that

$$-ih\partial_{\nu}\tilde{u}|_{x_1=0} = \text{Op}_h(\rho\chi)f,$$

by (3.2) we get the estimate

$$(4.25) \quad \|\mathcal{N}(z, h)\text{Op}_h(\chi)f - \text{Op}_h(\rho\chi)f\|_{L^2(\Gamma)} \lesssim h^{1/2}\theta^{-1/2}\|\tilde{v}\|_{L^2(\Omega)}.$$

On the other hand, Propositions 4.3 and 4.7 imply the following

Lemma 4.8. *If χ is of compact support, given any $N \geq 0$ independent of h , we have the estimate*

$$(4.26) \quad \|\tilde{v}\|_{L^2(\Omega)} \lesssim h^{1/2}\theta^{-2} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right) \|f\|_{H_h^{-N}(\Gamma)}.$$

If $\text{supp } \chi \subset \text{supp}(1 - \eta)$, given any $0 < \epsilon < 1/2$ independent of h , we have the estimate

$$(4.27) \quad \|\tilde{v}\|_{L^2(\Omega)} \lesssim h^{(1-3\epsilon)/2} \|f\|_{H_h^1(\Gamma)}.$$

Proof. Clearly,

$$\|\tilde{v}\|_{L^2(\Omega)}^2 \lesssim \int_{\delta/2}^{\delta} \left\| \text{Op}_h \left(e^{ix_1\rho/h} A_1 \right) f \right\|_{L^2(\Gamma)}^2 dx_1 + \int_0^{\delta} \left\| \text{Op}_h \left(e^{ix_1\rho/h} A_2 \right) f \right\|_{L^2(\Gamma)}^2 dx_1.$$

Therefore, if χ is of compact support, by (4.11) and (4.24), we get

$$\begin{aligned} \|\tilde{v}\|_{L^2(\Omega)}^2 &\lesssim h \|f\|_{H_h^{-N}(\Gamma)}^2 \int_{\delta/2}^{\delta} dx_1 \\ &+ \theta^{-3} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right)^2 \|f\|_{H_h^{-N}(\Gamma)}^2 \int_0^{\delta} e^{-Cx_1\theta/h} dx_1 \\ &\lesssim h\theta^{-4} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right)^2 \|f\|_{H_h^{-N}(\Gamma)}^2 \end{aligned}$$

for every $N \geq 0$, which clearly implies (4.26). Similarly, if $\text{supp } \chi \subset \text{supp}(1 - \eta)$, by (4.12) and (4.24), we get

$$\|\tilde{v}\|_{L^2(\Omega)}^2 \lesssim h \|f\|_{H_h^{-N}(\Gamma)}^2 \int_{\delta/2}^{\delta} dx_1 + h^{1-3\epsilon} \|f\|_{H_h^1(\Gamma)}^2 \int_0^{\delta} x_1^{-1+2\epsilon} dx_1 \lesssim h^{1-3\epsilon} \|f\|_{H_h^1(\Gamma)}^2$$

which implies (4.27). \square

Combining (4.25) with Lemma 4.8 (with ϵ small enough) and taking into account that $\theta^{-1/2} \leq h^{-1/5}$, we obtain the following

Lemma 4.9. *If χ is of compact support, given any $N \geq 0$ independent of h , we have the estimate*

$$(4.28) \quad \|\mathcal{N}(z, h)\text{Op}_h(\chi)f - \text{Op}_h(\rho\chi)f\|_{L^2(\Gamma)} \lesssim h\theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right) \|f\|_{H_h^{-N}(\Gamma)}.$$

If $\text{supp } \chi \subset \text{supp}(1 - \eta)$, we have the estimate

$$(4.29) \quad \|\mathcal{N}(z, h)\text{Op}_h(\chi)f - \text{Op}_h(\rho\chi)f\|_{L^2(\Gamma)} \lesssim h^{3/5} \|f\|_{H_h^1(\Gamma)}.$$

Since the function η can be written as a finite sum of functions χ for which (4.28) holds and the function $1 - \eta$ can be written as a finite sum of functions χ for which (4.29) holds, Lemma 4.9 implies the following

Proposition 4.10. *Given any $N \geq 0$ independent of h , we have the estimates*

$$(4.30) \quad \|\mathcal{N}(z, h)\text{Op}_h(\eta)f - \text{Op}_h(\rho\eta)f\|_{L^2(\Gamma)} \lesssim h\theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu}\right) \|f\|_{H_h^{-N}(\Gamma)},$$

and

$$(4.31) \quad \|\mathcal{N}(z, h)\text{Op}_h(1 - \eta)f - \text{Op}_h(\rho(1 - \eta))f\|_{L^2(\Gamma)} \lesssim h^{3/5} \|f\|_{H_h^1(\Gamma)}.$$

Clearly, the estimate (4.4) follows from (4.30) and (4.31). To prove the estimate (4.5) we need to improve (4.31), only, assuming that $c \equiv 1$. To this end, we have to build a better parametrix in the elliptic region in this case. This will be carried out in the next section.

5. IMPROVED PARAMETRIX IN THE ELLIPTIC REGION

Our goal in this section is to prove the following

Proposition 5.1. *Suppose that $c \equiv 1$ and $n \in C^1(\Omega_\delta)$, $n|_\Gamma \in C^2(\Gamma)$. Then we have the estimate*

$$(5.1) \quad \|\mathcal{N}(z, h)\text{Op}_h(1 - \eta)f - \text{Op}_h(\rho(1 - \eta) + hq)f\|_{L^2(\Gamma)} \lesssim h \|f\|_{H_h^{-1}(\Gamma)},$$

where $q \in S^0(\Gamma)$ is independent of z , h and the function n .

Then the estimate (4.5) would follow from (4.30) and (5.1). To prove (5.1) we will improve our parametrix when $\chi \in S^0(\Gamma)$ with $\text{supp } \chi \subset \text{supp}(1 - \eta)$. To this end, we choose the phase function in the form

$$\varphi = -\langle x', \xi' \rangle + x_1\rho + x_1^2\varphi_2 + x_1^3\varphi_3,$$

where the functions $\varphi_2, \varphi_3 \in S^1(\Gamma)$ do not depend on x_1 , z and h , and will be chosen in such a way that the eikonal equation (4.6) is satisfied with $\Phi = \mathcal{O}(1 + x_1^2|\xi'|^2)$. Furthermore, we choose the amplitude in the form

$$a = a_0 + x_1a_1 + x_1^2a_2,$$

where $a_0 = \chi$, $a_1 = a_{1,0} + ha_{1,1}$ with functions $a_{1,0}, a_2 \in S^0(\Gamma)$, $a_{1,1} \in S^{-1}(\Gamma)$ independent of x_1 and h , and will be chosen in such a way that $A_3 = \mathcal{O}(x_1^2|\xi'| + hx_1)$. To find φ_2 and φ_3 , observe that the function Φ with the new phase can be written in the form

$$\begin{aligned} \Phi &= 2\rho(2\varphi_2 + 3x_1\varphi_3) + x_1(2\varphi_2 + 3x_1\varphi_3)^2 - zn^\sharp \\ &\quad - 2\langle R_0(x')\xi', \nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3) \rangle \\ &\quad + x_1\langle R_0(x')\nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3), \nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3) \rangle \\ &\quad + \langle R^\sharp(x)\xi', \xi' \rangle - 2x_1\langle R^\sharp(x)\xi', \nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3) \rangle \\ &\quad + x_1^2\langle R^\sharp(x)\nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3), \nabla_{x'}(\rho + x_1\varphi_2 + x_1^2\varphi_3) \rangle. \end{aligned}$$

We need now the following

Lemma 5.2. *On $\text{supp}(1 - \eta)$ we have the bounds*

$$(5.2) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta \left(\rho - ir_0^{1/2} \right) \right| \leq C_\alpha |\xi'|^{-1-|\alpha|},$$

$$(5.3) \quad \left| \partial_{\xi'}^\alpha \left(\rho^{-1} + ir_0^{-1/2} \right) \right| \leq C_\alpha |\xi'|^{-3-|\alpha|},$$

$$(5.4) \quad \left| \partial_{\xi'}^\alpha \left(\nabla_{x'}\rho - i2^{-1}r_0^{-1/2}\nabla_{x'}r_0 \right) \right| \leq C_\alpha |\xi'|^{-1-|\alpha|},$$

for all multi-indices α and β such that $|\beta| \leq 2$.

Proof. The bound (5.2) with $\beta = 0$ follows from the identity

$$\rho - ir_0^{1/2} = \frac{zn_0}{\rho + ir_0^{1/2}}$$

together with (4.19) and the fact that $|\rho + ir_0^{1/2}| \geq Cr_0^{1/2}$, $C > 0$, on $\text{supp}(1 - \eta)$. Since $n_0 \in C^2$, we can differentiate the above identity with respect to x' to get (5.2) with $1 \leq |\beta| \leq 2$ in the same way. Furthermore, since

$$\rho^{-1} + ir_0^{-1/2} = i\rho^{-1}r_0^{-1/2} \left(\rho - ir_0^{1/2} \right),$$

the bound (5.3) follows from (5.2) and (4.18). Finally, the bound (5.4) follows from the identity (4.7) together with the bounds (5.3) and (4.18). \square

Write the function R^\sharp as

$$R^\sharp(x) = R_0^\sharp(x') + x_1 R_1^\sharp(x') + x_1^2 R^b(x),$$

where R_0^\sharp , R_1^\sharp , R^b are smooth functions. Then the function Φ can be written in the form

$$\Phi = \Phi_0 + x_1 \Phi_1 + \mathcal{O}(1 + x_1^2 |\xi'|^2),$$

where

$$\begin{aligned} \Phi_0 &= 4ir_0^{1/2} \varphi_2 - ir_0^{-1/2} \langle R_0(x') \xi', \nabla_{x'} r_0 \rangle + \langle R_0^\sharp(x') \xi', \xi' \rangle, \\ \Phi_1 &= 6ir_0^{1/2} \varphi_3 + 4\varphi_2^2 - 2 \langle R_0(x') \xi', \nabla_{x'} \varphi_2 \rangle + \langle R_1^\sharp(x') \xi', \xi' \rangle \\ &\quad - (4r_0)^{-1} \langle R_0(x') \nabla_{x'} r_0, \nabla_{x'} r_0 \rangle - ir_0^{-1/2} \langle R_0^\sharp(x') \xi', \nabla_{x'} r_0 \rangle. \end{aligned}$$

We now choose the function φ_2 so that $\Phi_0 = 0$ and the function φ_3 so that $\Phi_1 = 0$. Clearly, the functions φ_2 and φ_3 are smooth and satisfy the estimates

$$(5.5) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta \varphi_j \right| \leq C_{\alpha, \beta} |\xi'|^{1 - |\alpha|}, \quad j = 2, 3,$$

on $\text{supp} \chi$, for all multi-indices α and β . It is easy to see that (5.5) together with Lemma 5.2 imply the following

Lemma 5.3. *For $0 < x_1 \leq \delta$, we have the bounds*

$$(5.6) \quad \left| \partial_{\xi'}^\alpha \Phi \right| \leq C_\alpha (1 + x_1^2 |\xi'|^2) |\xi'|^{-|\alpha|}$$

for all multi-indices α with constants $C_\alpha > 0$ independent of x_1 , z and h .

To find the functions a_1 and a_2 , observe that in this case the function A_3 is of the form

$$\begin{aligned} A_3 &= -2i(\rho + 2x_1 \varphi_2 + 3x_1^2 \varphi_3)(a_1 + 2x_1 a_2) \\ &\quad - 2i \langle -\xi' + x_1 \nabla_{x'}(\rho + x_1 \varphi_2 + x_1^2 \varphi_3), \nabla_{x'}(\chi + x_1 a_1 + x_1^2 a_2) \rangle \\ &\quad \quad - 2i(\chi + x_1 a_1 + x_1^2 a_2)(\varphi_2 + x_1 \varphi_3) \\ &\quad - ix_1(\chi + x_1 a_1 + x_1^2 a_2) \langle R(x) \nabla_{x'}, \nabla_{x'} \rangle (\rho + x_1 \varphi_2 + x_1^2 \varphi_3) \\ &\quad \quad - h \langle R(x) \nabla_{x'}, \nabla_{x'} \rangle (\chi + x_1 a_1 + x_1^2 a_2) \\ &\quad \quad + Q_1(\chi + x_1 a_1 + x_1^2 a_2)(\rho + 2x_1 \varphi_2 + 3x_1^2 \varphi_3) \\ &\quad + \left\langle \tilde{Q}, -\xi' + x_1 \nabla_{x'}(\rho + x_1 \varphi_2) \right\rangle (\chi + x_1 a_1 + x_1^2 a_2) \\ &\quad \quad - ih \left\langle \tilde{Q}, \nabla_{x'}(\chi + x_1 a_1 + x_1^2 a_2) \right\rangle. \end{aligned}$$

Since $c \equiv 1$, the functions Q_1 and \tilde{Q} are smooth, so we can write them in the form

$$Q_1(x) = Q_{1,0}(x') + x_1 Q_{1,1}(x') + x_1^2 Q_1^b(x),$$

$$\tilde{Q}(x) = \tilde{Q}_0(x') + x_1 \tilde{Q}_1(x') + x_1^2 \tilde{Q}^b(x),$$

where all functions are smooth. Hence we can write the function A_3 in the form

$$A_3 = A_{3,0} + x_1 A_{3,1} + \mathcal{O}(x_1^2 |\xi'| + hx_1),$$

where

$$\begin{aligned} A_{3,0} &= 2r_0^{1/2} a_1 + 2i \langle \xi', \nabla_{x'} \chi \rangle - 2i \chi \varphi_2 + i Q_1 \chi r_0^{1/2} - \langle \tilde{Q}, \xi' \rangle \chi \\ &\quad - h \langle R_0(x') \nabla_{x'}, \nabla_{x'} \rangle \chi - ih \langle \tilde{Q}, \nabla_{x'} \chi \rangle, \\ A_{3,1} &= 4r_0^{1/2} a_2 - 4ia_1 \varphi_2 + 2i \langle \xi', \nabla_{x'} a_{1,0} \rangle + r_0^{-1/2} \langle \nabla_{x'} r_0, \nabla_{x'} \chi \rangle \\ &\quad - 2i \chi \varphi_3 - 2ia_1 \varphi_2 + \chi \langle R_0(x') \nabla_{x'}, \nabla_{x'} \rangle r_0^{1/2} \\ &\quad + 2\chi Q_{1,0}(x') \varphi_2 + i Q_{1,0}(x') r_0^{1/2} a_{1,0} + i \chi Q_{1,1}(x') r_0^{1/2} \\ &\quad - \chi \langle \tilde{Q}_1(x'), \xi' \rangle - \langle \tilde{Q}_0(x'), \xi' \rangle a_{1,0} + i 2^{-1} r_0^{-1/2} \langle \tilde{Q}_0(x'), \nabla_{x'} r_0 \rangle \chi. \end{aligned}$$

We now choose the function a_1 so that $A_{3,0} = 0$ and the function a_2 so that $A_{3,1} = 0$. We get

$$a_{1,0} = -ir_0^{-1/2} \langle \xi', \nabla_{x'} \chi \rangle + ir_0^{-1/2} \chi \varphi_2 - i 2^{-1} Q_1 \chi + 2^{-1} r_0^{-1/2} \langle \tilde{Q}, \xi' \rangle \chi \in S^0(\Gamma),$$

$$a_{1,1} = 2^{-1} r_0^{-1/2} \langle R_0(x') \nabla_{x'}, \nabla_{x'} \rangle \chi + i 2^{-1} r_0^{-1/2} \langle \tilde{Q}, \nabla_{x'} \chi \rangle \in S^{-1}(\Gamma).$$

The next lemma follows easily from (5.5) and Lemma 5.2.

Lemma 5.4. *For $0 < x_1 \leq \delta$, we have the bounds*

$$(5.7) \quad |\partial_{\xi'}^\alpha A_3| \leq C_\alpha (hx_1 + x_1^2 |\xi'|) |\xi'|^{-|\alpha|}$$

for all multi-indices α with constants $C_\alpha > 0$ independent of x_1, z and h .

Set

$$\tilde{\rho} = \rho + x_1 \varphi_2 + x_1^2 \varphi_3 = \rho + \mathcal{O}(x_1 |\xi'|).$$

Taking δ small enough we can arrange that the inequalities in (4.3) still hold with ρ replaced by $\tilde{\rho}$ for all $0 < x_1 \leq \delta$. Therefore, the bound (4.16) (with $k = 0$) holds with $e^{ix_1 \rho/h}$ replaced by $e^{ix_1 \tilde{\rho}/h}$. This together with Lemmas 5.3 and 5.4 imply the following

Lemma 5.5. *For $0 < x_1 \leq \delta$ and for all $0 < \epsilon < 1/2$, we have the estimates*

$$(5.8) \quad \left| \partial_{\xi'}^\alpha \left(e^{ix_1 \tilde{\rho}/h} A_2 \right) \right| \leq C_\alpha h^{5/2-\epsilon} x_1^{-1/2+\epsilon} |\xi'|^{-3/2+\epsilon-|\alpha|}$$

for all multi-indices α with constants $C_\alpha > 0$ independent of x_1, z and h .

It is easy also to see that the new function A_1 satisfies the bound (4.23) (on $\text{supp}(1 - \eta)$) with $e^{ix_1 \rho/h}$ replaced by $e^{ix_1 \tilde{\rho}/h}$. Hence Proposition 4.7 still holds with $e^{ix_1 \rho/h}$ replaced by $e^{ix_1 \tilde{\rho}/h}$. Therefore, in the same way as in the previous section we can deduce from Lemma 5.5 and Proposition 2.3 that the function \tilde{v} satisfies the estimate

$$(5.9) \quad \|\tilde{v}\|_{L^2(\Omega)} \lesssim h^{(5-3\epsilon)/2} \|f\|_{H_h^{-1}(\Gamma)}.$$

We also have

$$-ih \partial_\nu \tilde{u}|_{x_1=0} = \text{Op}_h(\rho \chi - iha_1) f = \text{Op}_h(\rho \chi - iha_{1,0}) f - ih^2 \text{Op}_h(a_{1,1}) f$$

and the operator $\text{Op}_h(a_{1,1}) : H_h^{-1}(\Gamma) \rightarrow L^2(\Gamma)$ is uniformly bounded. Therefore, Proposition 5.1 follows from (3.2) and (5.9).

6. PROOF OF THEOREM 1.1

We have to show that if $\lambda \in \mathbb{C}$ belongs to the eigenvalue-free regions in Theorem 1.1, then the solution (u_1, u_2) of the equation (1.1) is identically zero. Clearly, it suffices to show that the function $f = u_1|_\Gamma = u_2|_\Gamma$ is identically zero. We may suppose that $|\lambda| \gg 1$. Set $h = |\lambda|^{-1} \ll 1$ and $z = (h\lambda)^2$. Clearly, $z \in Z^+$ if $\operatorname{Re} \lambda \geq |\operatorname{Im} \lambda|$ and $z \in Z^-$ if $|\operatorname{Im} \lambda| \geq \operatorname{Re} \lambda$, where Z^+ and Z^- are as in Section 3. Set $\theta = |\operatorname{Im} z|$ if $z \in Z^+$ and $\theta = 1$ if $z \in Z^-$. In order to simplify the notations, in what follows the restrictions of the functions c_j, n_j, \tilde{n}_j , $j = 1, 2$, on Γ will be again denoted by c_j, n_j, \tilde{n}_j , respectively. By assumption, we have that they belong to $C^\mu(\Gamma)$. Define the functions ρ_j , $j = 1, 2$, by replacing in the definition of ρ the function \tilde{n} by \tilde{n}_j . Denote by \mathcal{N}_j , $j = 1, 2$, the Dirichlet-to-Neumann map introduced in Section 4 associated to (c_j, n_j) , and set

$$T(z, h) = c_1 \mathcal{N}_1(z, h) - c_2 \mathcal{N}_2(z, h).$$

If λ is a transmission eigenvalue, then $Tf = 0$ on Γ . By Theorem 4.1 we get

$$(6.1) \quad \begin{aligned} & \|\operatorname{Op}_h(c_1 \rho_1 - c_2 \rho_2) f\|_{L^2(\Gamma)} \lesssim h^{3/5} \|f\|_{H_h^k(\Gamma)} \\ & + h \theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu}\right) \|f\|_{H_h^k(\Gamma)} \end{aligned}$$

provided $\theta \geq h^{2/5}$, where $k = 1$ in the anisotropic case and $k = -1$ in the isotropic case. Set

$$a_1 = (r_0 + 1)^{-1/2} (c_1 \rho_1 + c_2 \rho_2), \quad a_2 = c_1 \rho_1 - c_2 \rho_2.$$

We will prove now the following

Proposition 6.1. *Suppose that θ satisfies the condition*

$$(6.2) \quad h^{\mu/2} \log(h^{-1}) \theta^{1/2-d-\mu} \leq 1.$$

Then we have the bounds

$$(6.3) \quad \|\operatorname{Op}_h(a_1)\|_{L^2(\Gamma) \rightarrow L^2(\Gamma)} \lesssim 1,$$

$$(6.4) \quad \|\operatorname{Op}_h(a_1) \operatorname{Op}_h(a_2) - \operatorname{Op}_h(a_1 a_2)\|_{H_h^k(\Gamma) \rightarrow L^2(\Gamma)} \lesssim h \theta^{-1} + h^{\mu/2} \log(h^{-1}) \theta^{1/2-d-\mu}.$$

Proof. Clearly, the functions a_1 and a_2 are bounded on $\operatorname{supp} \eta$. Moreover, it follows from (4.19) and (4.2) that they satisfy the bounds

$$(6.5) \quad \left| \partial_{\xi'}^\alpha \partial_x^\beta a_j \right| \lesssim \theta^{1/2-|\alpha|-|\beta|} \quad \text{on } \operatorname{supp} \eta,$$

for all multi-indices α and β such that $|\alpha| + |\beta| \geq 1$ and $|\beta| \leq \mu$. On the other hand, since $\rho_j = i r_0^{1/2} + \mathcal{O}(r_0^{-1/2})$ for $r_0 \gg 1$, we have

$$\begin{aligned} a_1 &= i(c_1 + c_2) + \mathcal{O}(r_0^{-1}), \\ a_2 &= i(c_1 - c_2) r_0^{1/2} + \mathcal{O}(r_0^{-1/2}), \quad \text{if } k = 1, \\ a_2 &= \frac{z(n_1 - n_2)}{\rho_1 + \rho_2} = (2i)^{-1} z(n_1 - n_2) r_0^{-1/2} + \mathcal{O}(r_0^{-3/2}), \quad \text{if } k = -1. \end{aligned}$$

Set $b_1 = i(c_1 + c_2)$ and

$$b_2 = \begin{cases} i(c_1 - c_2) & \text{if } k = 1, \\ (2i)^{-1} z(n_1 - n_2) & \text{if } k = -1. \end{cases}$$

We now decompose the functions a_j as follows

$$a_j = \sum_{\ell=1}^3 a_j^{(\ell)},$$

where $a_j^{(1)} = \eta a_j$, $a_1^{(3)} = b_1(1-\eta)$ and $a_2^{(3)} = b_2(1-\eta)r_0^{k/2}$. Then the functions $a_j^{(2)}$ are supported on $\text{supp}(1-\eta)$ and $a_1^{(2)} = \mathcal{O}(r_0^{-1})$, $a_2^{(2)} = \mathcal{O}(r_0^{k/2-1})$. Moreover, they satisfy the bounds

$$(6.6) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta a_1^{(2)} \right| \lesssim \langle \xi' \rangle^{-2-|\alpha|},$$

$$(6.7) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta a_2^{(2)} \right| \lesssim \langle \xi' \rangle^{k-2-|\alpha|},$$

for all multi-indices α and β such that $|\beta| \leq \mu$. These bounds follow from the fact that (5.2) holds for all β such that $|\beta| \leq \mu$, which in turn can be easily proved by induction in μ .

We can write

$$\begin{aligned} \text{Op}_h(a_1)\text{Op}_h(a_2) - \text{Op}_h(a_1 a_2) &= \sum_{\ell_1=1}^3 \sum_{\ell_2=1}^3 \left(\text{Op}_h(a_1^{(\ell_1)})\text{Op}_h(a_2^{(\ell_2)}) - \text{Op}_h(a_1^{(\ell_1)} a_2^{(\ell_2)}) \right) \\ &= \sum_{\ell_1=1}^2 \sum_{\ell_2=1}^2 \left(\text{Op}_h(a_1^{(\ell_1)})\text{Op}_h(a_2^{(\ell_2)}) - \text{Op}_h(a_1^{(\ell_1)} a_2^{(\ell_2)}) \right) \\ &\quad + b_1 \sum_{\ell_2=1}^2 \left(\text{Op}_h((1-\eta))\text{Op}_h(a_2^{(\ell_2)}) - \text{Op}_h((1-\eta)a_2^{(\ell_2)}) \right) \\ &\quad + b_2 \sum_{\ell_1=1}^2 \left(\text{Op}_h(a_1^{(\ell_1)})\text{Op}_h((1-\eta)r_0^{k/2}) - \text{Op}_h(a_1^{(\ell_1)}(1-\eta)r_0^{k/2}) \right) \\ &\quad + \sum_{\ell_1=1}^2 \left[\text{Op}_h(a_1^{(\ell_1)}), b_2 \right] \text{Op}_h((1-\eta)r_0^{k/2}) \\ &\quad + b_1 [\text{Op}_h(1-\eta), b_2] \text{Op}_h((1-\eta)r_0^{k/2}) \\ &\quad + b_1 b_2 \left(\text{Op}_h(1-\eta)\text{Op}_h((1-\eta)r_0^{k/2}) - \text{Op}_h((1-\eta)^2 r_0^{k/2}) \right). \end{aligned}$$

Now we apply Proposition 2.6 to the operators in the double sum in the right-hand side, Proposition 2.5 to the operators in the next two sums and Proposition 2.2 to the operator in the last term. Furthermore, the operator

$$\text{Op}_h((1-\eta)r_0^{k/2}) : H_h^k(\Gamma) \rightarrow L^2(\Gamma)$$

is bounded, while the norm of the commutators

$$\left[\text{Op}_h(a_1^{(\ell_1)}), b_2 \right] : L^2(\Gamma) \rightarrow L^2(\Gamma)$$

can be bounded by using (2.26), and the norm of the commutator

$$[\text{Op}_h(1-\eta), b_2] = -[\text{Op}_h(\eta), b_2] : L^2(\Gamma) \rightarrow L^2(\Gamma)$$

can be bounded by using (2.22). Thus, taking into account the bounds (6.5), (6.6) and (6.7), we arrive at (6.4). The bound (6.3) follows similarly from Propositions 2.1 and 2.4. \square

Combining Proposition 6.1 with the estimate (6.1) we get

$$(6.8) \quad \|\mathrm{Op}_h(a_1 a_2) f\|_{L^2(\Gamma)} \lesssim h^{3/5} \|f\|_{H_h^k(\Gamma)} + \mathcal{E}_1(h, \theta) \|f\|_{H_h^k(\Gamma)}$$

provided $\theta \geq h^{2/5}$ and θ satisfies (6.2), where

$$\mathcal{E}_1 = h\theta^{-5/2} \left(1 + h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu}\right) + h^{\mu/2} \log(h^{-1}) \theta^{1/2-d-\mu}.$$

Set

$$\theta_1(h) = \begin{cases} (h^{\mu/2} \log(h^{-1}))^{1/(d+\mu-1/2)} & \text{if } \mu \leq 2d-1, \\ (h^{\mu/2+1} \log(h^{-1}))^{1/(d+\mu+5/2)} & \text{if } 2d-1 < \mu \leq 4d, \\ h^{2/5} & \text{if } \mu > 4d. \end{cases}$$

It is easy to check that $\theta \gg \theta_1$ implies $\mathcal{E}_1 \ll 1$ and $\theta \geq h^{2/5}$ satisfies (6.2). We would like to show that for such θ and h small enough the estimate (6.8) implies $f \equiv 0$ in the isotropic case. To do so, we will use the identity

$$(6.9) \quad (c_1 \rho_1 + c_2 \rho_2)(c_1 \rho_1 - c_2 \rho_2) = c_1^2 \rho_1^2 - c_2^2 \rho_2^2 = -(c_1^2 - c_2^2) r_0 + z(c_1 n_1 - c_2 n_2).$$

So, in the isotropic case (6.9) gives

$$\mathrm{Op}_h(a_1 a_2) = z(n_1 - n_2) \mathrm{Op}_h \left((r_0 + 1)^{-1/2} \right).$$

Recall that in this case we have $|n_1 - n_2| \geq c$ with some constant $c > 0$ by assumption. Since

$$\left\| \mathrm{Op}_h \left((r_0 + 1)^{-1/2} \right) f \right\|_{L^2(\Gamma)} \sim \|f\|_{H_h^{-1}(\Gamma)},$$

the estimate (6.8) implies

$$(6.10) \quad \|f\|_{H_h^{-1}(\Gamma)} \lesssim h^{3/5} \|f\|_{H_h^{-1}(\Gamma)} + \mathcal{E}_1 \|f\|_{H_h^{-1}(\Gamma)}.$$

Taking h and \mathcal{E}_1 small enough we deduce from (6.10) that $\|f\|_{H_h^{-1}(\Gamma)} = 0$, and hence $f \equiv 0$, as desired. Hence the region $\theta \gg \theta_1$ is an eigenvalue-free region. It is easy to see that this region corresponds to (1.12) on the λ plane.

Consider now the anisotropic case. Then the function

$$m = \frac{c_1 n_1 - c_2 n_2}{c_1^2 - c_2^2}$$

is strictly negative under the condition (1.6) and strictly positive under the condition (1.8). Moreover, we have $m \in C^\mu(\Gamma)$. Set

$$A_1 = (r_0 + 1)(r_0 - zm)^{-1}, \quad A_2 = (r_0 + 1)^{-1/2}(r_0 - zm).$$

Let $\tau = |\mathrm{Im} z|$ if $z \in Z^-$ and $\tau = 1$ if $z \in Z^+$. We need now the following

Proposition 6.2. *Suppose the condition (1.8) fulfilled. Suppose also that θ satisfies the condition*

$$(6.11) \quad h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu} \leq 1.$$

Then we have the bounds

$$(6.12) \quad \|\mathrm{Op}_h(A_1)\|_{L^2(\Gamma) \rightarrow L^2(\Gamma)} \lesssim \theta^{-1},$$

$$(6.13) \quad \|\mathrm{Op}_h(A_1) \mathrm{Op}_h(A_2) - \mathrm{Op}_h(A_1 A_2)\|_{H_h^1(\Gamma) \rightarrow L^2(\Gamma)} \lesssim h\theta^{-2} + h^{\mu/2} \log(h^{-1}) \theta^{-d-\mu-1}.$$

Suppose the condition (1.6) fulfilled. Suppose also that $\tau \geq h^{1/2}$ satisfies the condition

$$(6.14) \quad h^{\mu/2} \log(h^{-1}) \tau^{-d-\mu} \leq 1.$$

Then we have the bounds

$$(6.15) \quad \|\mathrm{Op}_h(A_1)\|_{L^2(\Gamma) \rightarrow L^2(\Gamma)} \lesssim \tau^{-1},$$

$$(6.16) \quad \|\mathrm{Op}_h(A_1)\mathrm{Op}_h(A_2) - \mathrm{Op}_h(A_1A_2)\|_{H_h^1(\Gamma) \rightarrow L^2(\Gamma)} \lesssim h\tau^{-2} + h^{\mu/2} \log(h^{-1})\tau^{-d-\mu-1}.$$

Proof. Clearly, the function A_2 satisfies the bounds

$$(6.17) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta A_2 \right| \lesssim \langle \xi' \rangle^{1-|\alpha|},$$

for all multi-indices α and β such that $|\beta| \leq \mu$. We would like to find similar bounds for the derivatives of A_1 . To this end, observe that we can arrange $|r_0 - zm| \geq Cr_0$, $C > 0$, on $\mathrm{supp}(1 - \eta)$, provided the constant C_0 in the definition of η is taken large enough (what we can do without loss of generality). The function A_1 satisfies the following

Lemma 6.3. *If $m\mathrm{Re} z \geq 0$ we have the bounds*

$$(6.18) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta A_1 \right| \lesssim \begin{cases} |\mathrm{Im} z|^{-1-|\alpha|-|\beta|} & \text{on } \mathrm{supp} \eta, \\ |\xi'|^{-|\alpha|} & \text{on } \mathrm{supp}(1 - \eta), \end{cases}$$

for all multi-indices α and β such that $|\beta| \leq \mu$. If $m\mathrm{Re} z \leq 0$ the bounds (6.18) hold with $|\mathrm{Im} z|$ replaced by 1.

Proof. We have

$$|r_0 - zm| \geq C(r_0 + 1),$$

if $m\mathrm{Re} z \leq 0$, and

$$\begin{aligned} |r_0 - zm| &\geq C|\mathrm{Im} z| && \text{on } \mathrm{supp} \eta, \\ |r_0 - zm| &\geq C|\xi'|^2 && \text{on } \mathrm{supp}(1 - \eta), \end{aligned}$$

if $m\mathrm{Re} z \geq 0$, where $C > 0$ is a constant. The bounds (6.18) can be easily derived from the above inequalities by induction in $|\alpha| + |\beta|$. \square

Decompose the functions A_j as follows

$$A_j = \sum_{\ell=1}^3 A_j^{(\ell)},$$

where $A_j^{(1)} = \eta A_j$, $A_1^{(3)} = (1 - \eta)$ and $A_2^{(3)} = (1 - \eta)r_0^{1/2}$. Then the functions $A_j^{(2)}$ are supported on $\mathrm{supp}(1 - \eta)$ and $A_1^{(2)} = \mathcal{O}(r_0^{-1})$, $A_2^{(2)} = \mathcal{O}(r_0^{-1/2})$. Moreover, it follows from Lemma 6.3 that they satisfy the bounds

$$(6.19) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta A_1^{(2)} \right| \lesssim \langle \xi' \rangle^{-2-|\alpha|},$$

$$(6.20) \quad \left| \partial_{\xi'}^\alpha \partial_{x'}^\beta A_2^{(2)} \right| \lesssim \langle \xi' \rangle^{-1-|\alpha|},$$

for all multi-indices α and β such that $|\beta| \leq \mu$. We can write

$$\begin{aligned} \mathrm{Op}_h(A_1)\mathrm{Op}_h(A_2) - \mathrm{Op}_h(A_1A_2) &= \sum_{\ell_1=1}^3 \sum_{\ell_2=1}^3 \left(\mathrm{Op}_h(A_1^{(\ell_1)})\mathrm{Op}_h(A_2^{(\ell_2)}) - \mathrm{Op}_h(A_1^{(\ell_1)}A_2^{(\ell_2)}) \right) \\ &= \sum_{\ell_1=1}^2 \sum_{\ell_2=1}^2 \left(\mathrm{Op}_h(A_1^{(\ell_1)})\mathrm{Op}_h(A_2^{(\ell_2)}) - \mathrm{Op}_h(A_1^{(\ell_1)}A_2^{(\ell_2)}) \right) \end{aligned}$$

$$\begin{aligned}
 & + \sum_{\ell_2=1}^2 \left(\text{Op}_h((1-\eta)) \text{Op}_h(A_2^{(\ell_2)}) - \text{Op}_h((1-\eta)A_2^{(\ell_2)}) \right) \\
 & + \sum_{\ell_1=1}^2 \left(\text{Op}_h(A_1^{(\ell_1)}) \text{Op}_h((1-\eta)r_0^{1/2}) - \text{Op}_h(A_1^{(\ell_1)}(1-\eta)r_0^{1/2}) \right) \\
 & + \left(\text{Op}_h(1-\eta) \text{Op}_h((1-\eta)r_0^{1/2}) - \text{Op}_h((1-\eta)^2 r_0^{1/2}) \right).
 \end{aligned}$$

Now we apply Proposition 2.6 to the operators in the double sum in the right-hand side, Proposition 2.5 to the operators in the next two sums and Proposition 2.2 to the operator in the last term. Thus, taking into account the bounds (6.17) and (6.18), we arrive at (6.13) and (6.16). The bounds (6.12) and (6.15) follow from Propositions 2.1 and 2.4. \square

In view of (6.9) we have $A_2 = -(c_1^2 - c_2^2)a_1a_2$. By assumption, $|c_1^2 - c_2^2| \geq c > 0$, so Proposition 6.2 together with the estimate (6.8) imply

$$(6.21) \quad \|\text{Op}_h(A_1A_2)f\|_{L^2(\Gamma)} \lesssim h^{1/5} \|f\|_{H_h^1(\Gamma)} + \mathcal{E}(h, \theta, \tau) \|f\|_{H_h^1(\Gamma)}$$

provided $\theta \geq h^{2/5}$, θ satisfies (6.2) and (6.11), and $\tau \geq h^{1/2}$, τ satisfies (6.14), where $\mathcal{E} = \mathcal{E}_2$ if (1.8) holds, $\mathcal{E} = \mathcal{E}_1$ if (1.6) holds and $z \in Z^+$, and $\mathcal{E} = \mathcal{E}_3$ if (1.6) holds and $z \in Z^-$, where

$$\mathcal{E}_2 = h\theta^{-7/2} + h^{\mu/2} \log(h^{-1})\theta^{-d-\mu-1},$$

$$\mathcal{E}_3 = h\tau^{-2} + h^{\mu/2} \log(h^{-1})\tau^{-d-\mu-1}.$$

Set

$$\theta_2(h) = \begin{cases} (h^{\mu/2} \log(h^{-1}))^{1/(d+\mu+1)} & \text{if } \mu \leq \frac{4}{3}(d+1), \\ h^{2/7} & \text{if } \mu > \frac{4}{3}(d+1), \end{cases}$$

and

$$\tau_3(h) = \left(h^{\mu/2} \log(h^{-1}) \right)^{1/(d+\mu+1)}.$$

It is easy to check that $\theta \gg \theta_2$ implies $\mathcal{E}_2 \ll 1$ and $\theta \geq h^{2/7}$ satisfies (6.2) and (6.11). Similarly, $\tau \gg \tau_3$ implies $\mathcal{E}_3 \ll 1$ and $\tau \geq h^{1/2}$ satisfies (6.14). Since $A_1A_2 = (r_0 + 1)^{1/2}$ and

$$\left\| \text{Op}_h \left((r_0 + 1)^{1/2} \right) f \right\|_{L^2(\Gamma)} \sim \|f\|_{H_h^1(\Gamma)},$$

the estimate (6.21) implies

$$(6.22) \quad \|f\|_{H_h^1(\Gamma)} \lesssim h^{1/5} \|f\|_{H_h^1(\Gamma)} + \mathcal{E} \|f\|_{H_h^1(\Gamma)}.$$

Taking h and \mathcal{E} small enough we deduce from (6.22) that $\|f\|_{H_h^1(\Gamma)} = 0$, and hence $f \equiv 0$, as desired. In other words, the regions $\theta \gg \theta_2$ when (1.8) holds, and $\theta \gg \theta_1$ and $\tau \gg \tau_3$ when (1.6) holds are eigenvalue-free regions. It is easy to see that these regions correspond to eigenvalue-free regions in Theorem 1.1 on the λ plane.

REFERENCES

- [1] A.S. BONNET-BEN DHIA, L. CHESNEL AND H. HADDAR, *On the use of T -coercivity to study interior transmission eigenvalue problem*, C. R. Math. Acad. Sci. Paris **349** (2011), 647-651.
- [2] M. DIMASSI AND J. SJÖSTRAND, *Spectral Asymptotics in Semi-classical Limit*, London Mathematical Society, Lecture Notes Series, Vol. 268, Cambridge University Press, 1999.
- [3] J. FORNEROD AND H-M. NGUYEN, *The Weyl law of transmission eigenvalues and the completeness of generalized transmission eigenvalues without complementing conditions*, preprint 2022.
- [4] L. HÖRMANDER, *The Analysis of Linear Partial Differential Operators, Vol.3*, Springer, Berlin, 1985.
- [5] E. LAKSHTANOV AND B. VAINBERG, *Ellipticity in the interior transmission problem in anisotropic media*, SIAM J. Math. Anal. **44** (2012), no. 2, 1165-1174.
- [6] E. LAKSHTANOV AND B. VAINBERG, *Applications of elliptic operator theory to the isotropic interior transmission eigenvalue problem*, Inverse Problems **29** (2013), 104003.
- [7] Y.-J. LEUNG AND D. COLTON, *Complex transmission eigenvalues for spherically stratified media*, Inverse Problems **28** (2012), 075005.
- [8] H-M. NGUYEN AND Q-H. NGUYEN, *Discreteness of interior transmission eigenvalues revisited*, Calc. Var. Partial Differential Equations **56** (2017), no. 2, Paper No. 51.
- [9] H-M. NGUYEN AND Q-H. NGUYEN, *The Weyl law of transmission eigenvalues and the completeness of generalized transmission eigenvalues*, J. Funct. Anal. **281** (2021), 109146.
- [10] V. PETKOV AND G. VODEV, *Asymptotics of the number of the interior transmission eigenvalues*, J. Spectral Theory **7** (2017), 1-31.
- [11] L. ROBBIANO, *Counting function for interior transmission eigenvalues*, Math. Control Relat. Fields **6** (2016), no. 1, 167-183.
- [12] J. SYLVESTER, *Discreteness of transmission eigenvalues via upper triangular compact operators*, SIAM J. Math. Anal. **44** (2012), no. 1, 341-354.
- [13] J. SYLVESTER, *Transmission eigenvalues in one dimension*, Inverse Problems **29** (2013), 104009.
- [14] G. VODEV, *Transmission eigenvalue-free regions*, Comm. Math. Phys. **336** (2015), 1141-1166.
- [15] G. VODEV, *Parabolic transmission eigenvalue-free regions in the degenerate isotropic case*, Asymptot. Analysis **106** (2018), 147-168.
- [16] G. VODEV, *High-frequency approximation of the interior Dirichlet-to-Neumann map and applications to the transmission eigenvalues*, Anal.PDE **11** (2018), 213-236.
- [17] G. VODEV, *Semiclassical parametrix for the Maxwell equation and applications to the electromagnetic transmission eigenvalues*, Res. Math. Sci. **8** (2021), Paper No. 35.

UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, 2 RUE DE LA HOUSSINIÈRE, BP 92208, 44322 NANTES CEDEX 03, FRANCE
 Email address: Georgi.Vodev@univ-nantes.fr