

Hopf algebras and alternating multiple zeta values in positive characteristic

Bo-Hae Im, Hojin Kim, Khac Nhuan Le, Tuan Ngo Dac, Lan Huong Pham

▶ To cite this version:

Bo-Hae Im, Hojin Kim, Khac Nhuan Le, Tuan Ngo Dac, Lan Huong Pham. Hopf algebras and alternating multiple zeta values in positive characteristic. 2023. hal-04052875

HAL Id: hal-04052875 https://hal.science/hal-04052875

Preprint submitted on 30 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOPF ALGEBRAS AND ALTERNATING MULTIPLE ZETA VALUES IN POSITIVE CHARACTERISTIC

BO-HAE IM, HOJIN KIM, KHAC NHUAN LE, TUAN NGO DAC, AND LAN HUONG PHAM

ABSTRACT. In [32] we presented a systematic study of algebra structures of multiple zeta values in positive characteristic introduced by Thakur as analogues of classical multiple zeta values of Euler. In this paper we construct algebra and Hopf algebra structures of alternating multiple zeta values introduced by Harada, extending our previous work. Our results could be considered as an analogue of those of Hoffman [28] and Racinet [42] in the classical setting. The proof is based on two new ingredients: the first one is a direct and explicit construction of the shuffle Hopf algebra structure, and the second one is the notion of horizontal maps.

CONTENTS

1.	Introduction	1
2.	Review on Hopf algebra	6
3.	Review of shuffle Hopf algebra of MZV's	9
4.	Alternating multiple zeta values	11
5.	Shuffle algebra of AMZV's	13
6.	Shuffle Hopf algebra of AMZV's	20
Re	eferences	35

1. INTRODUCTION

1.1. Classical setting.

Let $\mathbb{N} = \{1, 2, ...\}$ be the set of positive integers. The multiple zeta values (MZV's for short) are infinite sum defined by

$$\zeta(n_1, \dots, n_r) = \sum_{0 < k_1 < \dots < k_r} \frac{1}{k_1^{n_1} k_2^{n_2} \dots k_r^{n_r}},$$

for $n_i \in \mathbb{N}$ with $n_r \geq 2$ for convergence. Here r and $w := n_1 + \cdots + n_r$ are said to be the depth and weight of the presentation $\zeta(n_1, \ldots, n_r)$, respectively. In the 18th century, Euler [20] studied MZV's for depth 1 and 2 cases. After two centuries, Zagier [54] studied the general MZV's, initiating extensive works connected to various

Date: March 30, 2023.

²⁰¹⁰ Mathematics Subject Classification. Primary 11M32; Secondary 11M38, 16S10, 16T30, 11R58.

Key words and phrases. multiple zeta values, zeta and L-functions in characteristic p, Hopf algebras, shuffle algebras, horizontal maps, arithmetic of function fields.

fields in mathematics and physics, including number theory, K-theory, knot theory, and higher energy physics (see for example [9, 17, 18, 27, 35, 30, 38, 44]).

There exist more general objects called colored multiple zeta values. Letting $N \in \mathbb{N}$ be a positive integer and Γ_N be the group of N-th roots of unity, we define colored multiple zeta values by

$$\zeta \begin{pmatrix} \epsilon_1 & \dots & \epsilon_r \\ n_1 & \dots & n_r \end{pmatrix} = \sum_{0 < k_1 < \dots < k_r} \frac{\epsilon_1^{k_1} \dots \epsilon_r^{k_r}}{k_1^{n_1} \dots k_r^{n_r}}$$

for $\epsilon_i \in \Gamma_N$ and $n_i \in \mathbb{N}$ with $(n_r, \epsilon_r) \neq (1, 1)$ for convergence. These objects were first systematically studied by Arakawa-Kaneko, Deligne, Goncharov and Racinet (see [5, 6, 15, 17, 22, 23, 42]). Since then they have been also studied by a lot of mathematicians and physicists, including Broadhurst [7, 8], Hoffman [29], Kaneko-Tsumura [34], and Zhao [55] in various fields of mathematics and physics. For N = 1we recover MZV's. For N = 2 the colored multiple zeta values are also known as alternating multiple zeta values (AMZV's for short) or Euler sums.

The main goal of the theory of colored multiple zeta values is to find all linear and algebraic relations among colored multiple zeta values over a suitable coefficient field. It led to several important conjectures formulated by Ihara-Kaneko-Zagier [30], Zagier [54] and Hoffman [27]. We will explain below Ihara-Kaneko-Zagier's conjecture and refer the reader to the excellent book [11] for more details about Zagier and Hoffman's conjectures.

For MZV's (the case N = 1) Ihara-Kaneko-Zagier [30] introduced the double shuffle relations and regularization. Roughly speaking, they used Hoffman's algebra \mathfrak{h} (see [28]) and subalgebras $\mathfrak{h}^0 \subset \mathfrak{h}^1 \subset \mathfrak{h}$ which are related to the MZV's, and presented two different algebra structures ($\mathfrak{h}^1, *$) and (\mathfrak{h}, \sqcup). These structures can be extended to Hopf algebra structures (see for example the work of Hoffman [28]). By regularization one gets the double shuffle relations by 'comparing' two products in the level of MZV's. Double shuffle relations and regularization produce linear relations among MZV's, and Ihara-Kaneko-Zagier conjectured that these relations are enough to understand all of the linear relations among MZV's (see [30, Conjecture 1]).

For colored multiple zeta values the previous works of Ihara-Kanako-Zagier and Hoffman were extended by Racinet in [42]. However, we warn the reader that as mentioned in *loc. cit.*, for general N the double shuffle relations and regularization fail to exhaust all the linear relations among colored multiple zeta values.

1.2. Positive characteristic setting.

1.2.1. MZV's and AMZV's in positive characteristic.

Carlitz [12] and Thakur [45, 49] defined the zeta and multiple zeta values in positive characteristic via the function fields analogy (see [33, 39, 52]). Let q be a power of prime p and \mathbb{F}_q be a finite field of order q. We use the following notations:

- $A = \mathbb{F}_q[\theta]$ the ring of polynomials in the variable θ over \mathbb{F}_q ,
 - A_+ the set of monic polynomials in A,
- $K = \mathbb{F}_q(\theta)$ the field of fractions of A (with the rational point ∞),
- $K_{\infty} = \mathbb{F}_q((1/\theta))$ the completion of K at ∞ ,

 v_{∞} the discrete valuation on K associated to the place ∞ ,

normalized as
$$v_{\infty}(\theta) = -1$$
,

 $|\cdot|_{\infty} = q^{-v_{\infty}(\cdot)}$ the corresponding absolute value on K.

Carlitz [12] defined the Carlitz zeta values $\zeta_A(n)$ for $n \in \mathbb{N}$ as

$$\zeta_A(n) := \sum_{a \in A_+} \frac{1}{a^n} \in K_{\infty}.$$

Note that this definition is an analogue of the classical zeta values $\zeta(n) = \sum_{k \in \mathbb{N}} \frac{1}{k^n}$. Extending this, Thakur [45] defined the multiple zeta value in positive characteristic for tuples $\mathfrak{s} = (s_1, \ldots, s_r) \in \mathbb{N}^r$ as

$$\zeta_A(\mathfrak{s}) = \sum \frac{1}{a_1^{s_1} \dots a_r^{s_r}}.$$

Here the sum is over all tuples of monic polynomials with decreasing degree, i.e., $(a_1, \ldots, a_r) \in A_+^r$ with $\deg(a_1) > \cdots > \deg(a_r)$. One can see the analogy between the classical MZV's and MZV's in positive characteristic. We define the depth and weight of $\zeta_A(\mathfrak{s})$ by r and $w(\mathfrak{s}) := s_1 + \cdots + s_r$ respectively. Note that Carlitz zeta values are MZV's of depth one. Thakur showed some of their properties including non-vanishing [46], sum-shuffle relations [48], etc. For further references on the MZV's in positive characteristic, see [3, 4, 21, 24, 36, 37, 41, 45, 47, 48, 49, 50, 53].

In [25] Harada introduced the notion of alternating multiple zeta values in positive characteristic. Let $\mathfrak{s} = (s_1, \ldots, s_r) \in \mathbb{N}^r$ and $\boldsymbol{\varepsilon} = (\varepsilon_1, \ldots, \varepsilon_r) \in (\mathbb{F}_q^*)^r$, we define

$$\zeta_A \begin{pmatrix} \boldsymbol{\varepsilon} \\ \boldsymbol{\mathfrak{s}} \end{pmatrix} = \sum \frac{\varepsilon_1^{\deg a_1} \dots \varepsilon_r^{\deg a_r}}{a_1^{s_1} \dots a_r^{s_r}} \in K_{\infty}.$$

Here, the sum is over all tuples of monic polynomials with decreasing degree as MZV's, i.e., $(a_1, \ldots, a_r) \in A^r_+$ with $\deg(a_1) > \cdots > \deg(a_r)$. We define the depth and weight of $\zeta_A \begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix}$ in a similar manner, that is, r is the depth and $w(\mathfrak{s}) = s_1 + \cdots + s_r$ is the weight.

In *loc. cit.*, Harada also showed some fundamental properties of AMZV's which hold for MZV's including non-vanishing, sum-shuffle relations, period interpretation and linear independence.

1.2.2. Zagier-Hoffman's conjectures for AMZV's in positive characteristic.

The primary objective of the theory of MZV's and AMZV's in positive characteristic is to comprehend all of their K-linear relations as in the classical setting. The positive characteristic version of Zagier-Hoffman's conjectures for MZV's is suggested by Todd [51] and Thakur [49]. The algebraic part of these conjectures was solved by the fourth author [40] using ingredients of Chen [14], Thakur [48, 49] and Todd [51]. The transcendental part was solved for the small weight cases by the fourth author in [40] by applying Anderson's theory of motives [1, 10, 26], as well as Anderson-Brownawell-Papanikolas criterion [2] as a tool for transcendence.

In [31], by discovering a deep link among AMZV's and alternating Carlitz multiple polylogarithms, we solved both the MZV's and AMZV's version of the Zagier-Hoffman's conjectures in positive characteristic. We mention that the latter is much harder than the former (see [31, §0.3] for a detailed discussion). More precisely, we show that

$$\dim_K \mathcal{AZ}_w = s(w)$$

where \mathcal{AZ}_w is the vector space spanned by the AMZV's in the positive characteristic of weight w over K, and s(w) is a Fibonacci-like sequence defined by

$$s(w) = \begin{cases} 1 & \text{if } w = 0, \\ (q-1)q^{w-1} & \text{if } 1 \le w \le q-1, \\ (q-1)(q^{w-1}-1) & \text{if } w = q, \\ (q-1)\sum_{i=1}^{q-1} s(w-i) + s(w-q) & \text{if } w > q. \end{cases}$$

Further, we give an explicit basis over K of \mathcal{AZ}_w . Unlike the MZV's case, this basis consists of linear combinations of AMZV's, not AMZV's. This crucial point has been observed in the classical setting by Charlton [13] and Deligne [15].

1.2.3. Algebraic structures of MZV's in positive characteristic.

In our recent paper [32] we addressed a question raised by a referee of a previous paper [40] by the fourth author investigating the algebraic structure of MZV's in positive characteristic. It turns out¹ that in 2016, Shuji Yamamoto [49] gave a construction of the word algebra for the shuffle product, and Shi [43] equipped it with a conjectural Hopf algebra structure, but left some axioms for the Hopf algebra open (see [43, Conjecture 3.2.2 and 3.2.11]). Following this, Thakur wrote to Deligne, and he [16] responded with a helpful letter in 2017. In it, Deligne stated that he had never seen such a construction before and advised on how to proceed with proving and investigating it.

Conjecture 1.1. The set of MZV's in positive characteristic can be equipped with a shuffle Hopf algebra structure.

In [32] we proved Conjecture 1.1:

Theorem 1.2 ([32, Theorem B]). Conjecture 1.1 is true.

Further, based on the previous link among AMZV's and alternating Carlitz multiple polylogarithms introduced in [31], we deduced a Hopf algebra of MZV's for the stuffle product (see [32, §9]).

1.3. Main results and key ingredients.

1.3.1. Main results.

In this paper we present a systematic study of algebraic structures of AMZV's in positive characteristic, and hence extend our previous work for MZV's. It could be considered as an analogue of Hoffman's and Racinet's results mentioned above in the classical setting.

Let us stress that the extension from the setting of MZV's to that of AMZV's is highly nontrivial. This fact is somewhat expected as in [31] we have encountered a similar problem while extending the proof of Zagier-Hoffman's conjectures for MZV's to that for AMZV's.

Roughly speaking, in the case of MZV's, if we consider the corresponding word algebra denoted by \mathfrak{C} (see §3.2), then there is only one letter of fixed weight n

¹We are thankful to Thakur for his assistance with event orders.

which is denoted by x_n . Consequently, it follows that there is also a unique and canonical way to define the coproducts $\Delta(x_n)$ so that the compatibility for x_1 and x_n is satisfied for all $n \in \mathbb{N}$ (see §3.3). Unfortunately, when we work with AMZV's, the situation is much more complicated. Although it is not difficult to define the word algebra \mathfrak{D} of AMZV's and to show that it is associative, the definition of a coproduct is far from evident. The main issue is that in \mathfrak{D} there are q - 1 letters of fixed weight, and there are no canonical ways to define coproducts for letters by using the compatibility for letters of weight 1 and other letters. To move forward we would need some new ideas.

1.3.2. Key ingredients.

The solution that we give here differs from that for the MZV's case at some crucial points and requires two new ingredients.

The first new ingredient is giving direct and explicit formulas for the coproduct Δ (see §6.1). We do not define it by the recursive induction on weight as it is the case for MZV's.

The second new ingredient is introducing so-called horizontal maps which allow us to deal with letters of the same weight (see §5.2). In particular, if some property is verified for a word beginning with a specific letter, says x, then by using horizontal maps we are able to deduce this property for all words beginning with a letter of the same weight as x.

With these tools we prove that the coproduct verifies all the desired properties so that the word algebra \mathfrak{D} of AMZV's is indeed a connected graded Hopf algebra.

1.3.3. Generalization for colored MZV's in positive characteristic.

We now give some comments about our results.

First, as mentioned above we have discovered a connection among AMZV's and the counterpart of Carlitz multiple polylogarithms in our previous work [31, §1]. Using this connection we could define another Hopf algebra structure on \mathfrak{D} called the stuffle Hopf algebra of AMZV's which extends [32, §9]. As the extension can be done without difficulty, we leave it to the reader as an exercise.

Second, our construction of the Hopf algebra structure for the word algebra \mathfrak{D} of AMZV's can be extended in a straightforward manner to the word algebra \mathfrak{D}_N of colored MZV's for all $N \in \mathbb{N}$. More precisely, the letters of \mathfrak{D}_N consist of $x_{n,\varepsilon}$ where $n \in \mathbb{N}$ and ε is an N-th root of unity in $\overline{\mathbb{F}}_q$. However, we warn the reader that in general, colored MZV's do not belong to K_∞ but only to a finite extension of K_∞ .

1.3.4. Plan of the manuscript.

The structure of this paper is as follows.

- In §2 we review some notions on the Hopf algebra. Our presentation is identical to that given in [32].
- In §3, we recall the word algebra \mathfrak{C} of MZV's (called the composition space in [32]) and the shuffle Hopf algebra structure of \mathfrak{C} . Then we review main results of *loc. cit* and state all necessary results that we will need in this paper so that the reader could read it easily without referring to [32].
- In §4 we review the definitions and basic properties of the AMZV's in positive characteristic introduced by Harada [25].

- In §5 we introduce the word algebra or composition space \mathfrak{D} of AMZV's and equip it with the shuffle algebra structure. We then define horizontal maps in §5.2. Finally, using these maps we prove that \mathfrak{D} is an algebra (i.e., the associativity) and that there exists a shuffle map for AMZV's (see Theorem 5.9 and Theorem 5.11).
- In §6 we define the coproduct on \mathfrak{D} . This definition differs from that on \mathfrak{C} as it is direct, explicit and does not use any induction on weight. Using horizontal maps we prove the compatibility and coassociativity results (see Theorem 6.5 and Theorem 6.8, respectively), establishing the main theorem of this paper (see Theorem 6.9 for a precise statement):

Theorem 1.3. The word algebra \mathfrak{D} of AMZV's equipped with the shuffle product and the above coproduct is a connected graded Hopf algebra of finite type over \mathbb{F}_q .

Acknowledgments. We would like to thank Dinesh Thakur for his interest and guidance with literature and for helpful discussions. Bo-Hae Im was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by grant funded by the Korea government (MSIT) (NRF-2023R1A2C1002385). KN. Le and T. Ngo Dac were partially supported by the Excellence Research Chair "*L*-functions in positive characteristic and applications" financed by the Normandy Region. T. Ngo Dac and LH. Pham were partially supported by the Vietnam Academy of Science and Technology (VAST) under grant no. CTTH00.02/23-24 "Arithmetic and Geometry of schemes over function fields and applications".

2. Review on Hopf Algebra

We give a brief review on basic notions of Hopf algebras, referring to our previous paper [31, §2] on the Hopf algebra structure of MZV's in positive characteristic.

2.1. Hopf algebra.

Let k be a field. From now on, tensor products are taken over k. Let H be a k-vector space, and $\iota: H \otimes H \to H \otimes H$ be the transposition map, defined as $\iota(x \otimes y) = y \otimes x$ and to be linear.

Definition 2.1. An algebra over k is a triple (H, m, u) consisting of a k-vector space H together with k-linear maps $m: H \otimes H \to H$ called the multiplication and $u: k \to H$ called the unit such that the following diagrams are commutative:

(1) associativity

$$\begin{array}{ccc} H \otimes H \otimes H & \xrightarrow{m \otimes \mathrm{id}} & H \otimes H \\ \mathrm{id} \otimes m & & & \downarrow m \\ H \otimes H & \xrightarrow{m} & H \end{array}$$

(2) unitary

$$H \otimes k \xrightarrow{\operatorname{id} \otimes u} H \otimes H \xleftarrow{u \otimes \operatorname{id}} k \otimes H$$

$$\downarrow^m$$

$$H$$

where the diagonal arrows are canonical isomorphisms.

The algebra is said to be commutative if the following diagram is commutative:



Coalgebra over k is dual to the algebra over k, obtained by reversing the arrows.

Definition 2.2. A coalgebra over k is a triple (H, Δ, ϵ) consisting of a k-vector space H together with k-linear maps $\Delta: H \to H \otimes H$ called the coproduct and $\epsilon: H \to k$ called the counit such that the following diagrams are commutative:

(1) coassociativity



(2) counitary



where the diagonal arrows are canonical isomorphisms.

The coalgebra is said to be cocommutative if the following diagram is commutative:



Definition 2.3. A bialgebra over k is a 5-tuple $(H, m, u, \Delta, \epsilon)$ consisting of an algebra (H, m, u) over k and a coalgebra (H, Δ, ϵ) over k which are compatible, i.e., the following diagrams are commutative:

(1) product and coproduct

$$\begin{array}{ccc} H \otimes H & & \stackrel{m}{\longrightarrow} H & \stackrel{\Delta}{\longrightarrow} H \otimes H \\ & & & & & \uparrow \\ & & & & \uparrow \\ H \otimes H \otimes H \otimes H \otimes H & \stackrel{\operatorname{id} \otimes \iota \otimes \operatorname{id}}{\longrightarrow} H \otimes H \otimes H \otimes H \end{array}$$

(2) unit and coproduct

$$\begin{array}{c} H & \stackrel{\Delta}{\longrightarrow} & H \otimes H \\ u \uparrow & & \uparrow u \otimes u \\ k & \stackrel{\text{canonical}}{\longrightarrow} & k \otimes k \end{array}$$

(3) counit and product

$$\begin{array}{c} H \xleftarrow{m} H \otimes H \\ \epsilon \downarrow & \downarrow \epsilon \otimes \epsilon \\ k \xleftarrow{} canonical} k \otimes k \end{array}$$

(4) unit and counit



Definition 2.4. A Hopf algebra over k is a bialgebra $(H, m, u, \Delta, \epsilon)$ over k together with a k-linear map $S: H \to H$ called antipode such that the following diagram is commutative:



We note that there exist a bialgebra which does not admit an antipode (see [11,Exercise 3.83] for an example).

2.2. Graded Hopf algebras.

We recall the notions of connected graded bialgebras.

Definition 2.5.

(1) A bialgebra $(H, m, u, \Delta, \epsilon)$ over k is said to be graded if one can write H as a direct sum of k-subspaces

$$H = \bigoplus_{n=0}^{\infty} H_n,$$

such that for all integers $r, s \ge 0$, we have

$$m(H_r \otimes H_s) \subseteq H_{r+s}$$
 and $\Delta(H_r) \subseteq \bigoplus_{i+j=r} H_i \otimes H_j$

A graded bialgebra is said to be connected if $H_0 = k$.

- (2) A graded Hopf algebra is a Hopf algebra H whose underlying bialgebra is graded and the antipode S satisfies $S(H_n) \subseteq H_n$.
- (3) A graded Hopf algebra is said to be connected if $H_0 = k$.
- (4) A graded Hopf algebra is said to be of finite type if H_n is a k-vector space of finite dimension.

The following proposition shows that a connected graded bialgebra automatically admits an antipode, thus it is always a Hopf algebra.

Proposition 2.6. Let $(H, m, u, \Delta, \epsilon)$ be a connected graded bialgebra over k.

(1) For each element $x \in H_n$ with $n \ge 1$, we have

$$\Delta(x) = 1 \otimes x + x \otimes 1 + \sum x_{(1)} \otimes x_{(2)},$$

where $\sum x_{(1)} \otimes x_{(2)} \in \bigoplus_{\substack{i,j>0\\i+j=n}} H_i \otimes H_j$. Moreover, the counit ϵ vanishes on H_n

for all n > 1.

(2) We continue the notation as in (1) and define recursively a k-linear map $S: H \to H$ given by

$$S(x) = \begin{cases} x & \text{if } x \in H_0, \\ -x - \sum m(S(x_{(1)}) \otimes x_{(2)}) & \text{if } x \in H_n \text{ with } n \ge 1. \end{cases}$$

Then H is a graded Hopf algebra whose antipode is S.

Proof. See [19, Lemma 2.1].

3. Review of shuffle Hopf algebra of MZV's

We briefly review some results of our previous work [32] on algebra structures of MZV's in positive characteristic.

3.1. Alphabet.

Let *I* be a countable set and $A = \{x_n\}_{n \in I}$ be a set of variables indexed by *I*, equipped with weights $w(x_n) \in \mathbb{N}$. The set *A* will be called an alphabet and its elements will be called letters. A word over the alphabet *A* is a finite string of letters. In particular, the empty word will be denoted by 1. The depth depth(\mathfrak{a}) of a word \mathfrak{a} is the number of letters in the string of \mathfrak{a} , so that depth(1) = 0. The weight of a word is the sum of the weights of its letter and we put w(1) = 0. Let $\langle A \rangle$ denote the set of all words over *A*. We endow $\langle A \rangle$ with the concatenation product defined by the following formula:

$$x_{i_1} \dots x_{i_n} \cdot x_{j_1} \dots x_{j_m} = x_{i_1} \dots x_{i_n} x_{j_1} \dots x_{j_m}.$$

Let $\mathbb{F}_q\langle A\rangle$ be the free \mathbb{F}_q -vector space with basis $\langle A\rangle$. The concatenation product extends to $\mathbb{F}_q\langle A\rangle$ by linearity. For a letter $x_a \in A$ and an element $\mathfrak{a} \in \mathbb{F}_q\langle A\rangle$, we write simply $x_a\mathfrak{a}$ instead of $x_a \cdot \mathfrak{a}$. For each nonempty word $\mathfrak{a} \in \langle A\rangle$, we can write $\mathfrak{a} = x_a\mathfrak{a}_-$ where x_a is the first letter of \mathfrak{a} and \mathfrak{a}_- is the word obtained from \mathfrak{a} by removing x_a .

3.2. Alphabet and shuffle algebra associated to MZV's.

We recall the alphabet associated to MZV's as given in [32, §4.2]. The set I will be \mathbb{N} and we denote by $\Sigma = \{x_n\}_{n \in \mathbb{N}}$ with weight $w(x_n) = n$ the alphabet attached to MZV's. We put $\mathfrak{C} = \mathbb{F}_q \langle \Sigma \rangle$, i.e., the free \mathbb{F}_q -vector space with basis $\langle \Sigma \rangle$.

For positive integers r, s and for all positive integers i, j such that i + j = r + s, we put

(3.1)
$$\Delta_{r,s}^{i} = \begin{cases} (-1)^{r-1} {\binom{i-1}{r-1}} + (-1)^{s-1} {\binom{i-1}{s-1}} & \text{if } (q-1) \mid i, \\ 0 & \text{otherwise.} \end{cases}$$

Then we define recursively two products on $\mathfrak C$ as $\mathbb F_q\text{-bilinear maps}$

 $\diamond \colon \mathfrak{C} \times \mathfrak{C} \longrightarrow \mathfrak{C} \quad \text{and} \quad \sqcup \colon \mathfrak{C} \times \mathfrak{C} \longrightarrow \mathfrak{C}$

by setting $1 \diamond \mathfrak{a} = \mathfrak{a} \diamond 1 = \mathfrak{a}, 1 \sqcup \mathfrak{a} = \mathfrak{a} \sqcup 1 = \mathfrak{a}$ and

$$\begin{split} \mathfrak{a} \diamond \mathfrak{b} &= x_{a+b}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})), \\ \mathfrak{a} \sqcup \mathfrak{b} &= x_{a}(\mathfrak{a}_{-} \sqcup \mathfrak{b}) + x_{b}(\mathfrak{a} \sqcup \mathfrak{b}_{-}) + \mathfrak{a} \diamond \mathfrak{b} \end{split}$$

for any nonempty words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$. We call \diamond the diamond product and \sqcup the shuffle product. The unit $u : \mathbb{F}_q \to \mathfrak{C}$ is given by sending 1 to the empty word. One of the main results of [32] reads as follows (see [32, Theorem A]):

Theorem 3.1. The spaces (\mathfrak{C}, \diamond) and (\mathfrak{C}, \sqcup) are commutative \mathbb{F}_q -algebras.

3.3. Shuffle Hopf algebra associated to MZV's.

We briefly recall the coproduct as in [32, \$7.1]

$$\Delta: \mathfrak{C} \to \mathfrak{C} \otimes \mathfrak{C}.$$

We first define it on $\langle \Sigma \rangle$ by induction on weight and extend by \mathbb{F}_q -linearity to \mathfrak{C} . We put

$$\Delta(1) := 1 \otimes 1, \qquad \Delta(x_1) := 1 \otimes x_1 + x_1 \otimes 1.$$

Let $w \in \mathbb{N}$ and we suppose that we have defined $\Delta(\mathfrak{u})$ for all words \mathfrak{u} of weight $w(\mathfrak{u}) < w$. We now give a formula for $\Delta(\mathfrak{u})$ for all words \mathfrak{u} with $w(\mathfrak{u}) = w$. For such a word \mathfrak{u} with depth(\mathfrak{u}) > 1, we put $\mathfrak{u} = x_u \mathfrak{v}$ with $w(\mathfrak{v}) < w$. Since x_u and \mathfrak{v} are both of weight less than w, we have already defined

$$\Delta(x_u) = 1 \otimes x_u + \sum a_u \otimes b_u, \qquad \Delta(\mathfrak{v}) = \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}}.$$

Then we set

$$\Delta(\mathfrak{u}) := 1 \otimes \mathfrak{u} + \sum (a_u \triangleright a_v) \otimes (b_u \sqcup b_v).$$

Our last task is to define $\Delta(x_w)$. We know that

$$x_1 \sqcup x_{w-1} = x_w + x_1 x_{w-1} + x_{w-1} x_1 + \sum_{0 < j < w} \Delta^j_{1,w-1} x_{w-j} x_j$$

where all the words $x_{w-j}x_j$ have weight w and depth 2 and all $\Delta_{1,w-1}^j$ defined as in (3.1) belong to \mathbb{F}_q . Therefore, we set

$$\Delta(x_w) := \Delta(x_1) \sqcup \Delta(x_{w-1}) - \Delta(x_1 x_{w-1}) - \Delta(x_{w-1} x_1) - \sum_{0 < j < w} \Delta_{1,w-1}^j \Delta(x_{w-j} x_j)$$

By induction on weight, we can check that Δ preserves the grading.

Next, the counit $\epsilon : \mathfrak{C} \to \mathbb{F}_q$ is defined by putting $\epsilon(1) = 1$ and $\epsilon(\mathfrak{u}) = 0$ otherwise. We proved (see [32, Theorem B]):

Theorem 3.2. The connected graded bialgebra $(\mathfrak{C}, \sqcup, u, \Delta, \epsilon)$ is a connected graded Hopf algebra over \mathbb{F}_q .

3.4. Some properties of the coproduct.

In this section we collect some properties of the coproduct that will be useful in the sequel.

Lemma 3.3. For all words $\mathfrak{u} \in \mathfrak{C}$, we have

$$\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum a_{\mathfrak{u}} \otimes b_{\mathfrak{u}}, \ \text{where} \ a_{\mathfrak{u}} \neq 1.$$

Proof. See [32, Lemma 7.1].

Lemma 3.4. Let \mathfrak{u} and \mathfrak{v} be two nonempty words in $\langle \Sigma \rangle$. Suppose that

$$\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum \mathfrak{u}_{(1)} \otimes \mathfrak{u}_{(2)}, and \quad \Delta(\mathfrak{v}) = 1 \otimes \mathfrak{v} + \sum \mathfrak{v}_{(1)} \otimes \mathfrak{v}_{(2)}.$$

Then

$$\Delta(\mathfrak{u} \diamond \mathfrak{v}) = 1 \otimes (\mathfrak{u} \diamond \mathfrak{v}) + \sum (\mathfrak{u}_{(1)} \diamond \mathfrak{v}_{(1)}) \otimes (\mathfrak{u}_{(2)} \sqcup \mathfrak{v}_{(2)}).$$

Proof. See [32, Lemma 8.5].

Finally, there is a useful formula for $\Delta(x_n)$ for $x_n \in \Sigma$ that we briefly explain below.

Lemma 3.5. Let $n \in \mathbb{N}$. Then we have

(1) for all j < n,

$$\Delta_{1,n}^{j} = \begin{cases} 1 & \text{if } (q-1) \mid j \\ 0 & \text{otherwise.} \end{cases}$$

(2) $\Delta_{1,n}^n = 0.$

Proof. The result is straightforward from the definition of $\Delta_{1,n}^j$. See [32, Lemma 8.1].

Next, we introduce the bracket operator by the following formula: for any $\mathfrak{a} = x_{i_1} \dots x_{i_m} \in \langle \Sigma \rangle$,

(3.2)
$$[\mathfrak{a}] := (-1)^m \Delta_{1,w(\mathfrak{a})+1}^{i_1} \cdots \Delta_{1,w(\mathfrak{a})+1}^{i_m} x_{i_1} \sqcup \cdots \sqcup x_{i_m}.$$

As a matter of convention, we also agree that [1] = 1. Then we showed (see [32, Theorem C]):

Proposition 3.6. We keep the above notation. Then for all $n \in \mathbb{N}$, we have

$$\Delta(x_n) = 1 \otimes x_n + \sum_{\substack{r \in \mathbb{N}, \mathfrak{a} \in \langle \Sigma \rangle \\ r+w(\mathfrak{a})=n}} \binom{r + \operatorname{depth}(\mathfrak{a}) - 2}{\operatorname{depth}(\mathfrak{a})} x_r \otimes [\mathfrak{a}].$$

4. Alternating multiple zeta values

We recall some notions on MZV's and AMZV's in positive characteristic, referring to [25, 31, 32].

Letting $\mathfrak{s} = (s_1, \ldots, s_n) \in \mathbb{N}^n$ and $\mathfrak{e} = (\varepsilon_1, \ldots, \varepsilon_n) \in (\mathbb{F}_q^{\times})^n$, we set $\mathfrak{s}_- := (s_2, \ldots, s_n)$ and $\mathfrak{e}_- := (\varepsilon_2, \ldots, \varepsilon_n)$. A positive array $\begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix}$ is an array of the form $\begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix} = \begin{pmatrix} \varepsilon_1 & \cdots & \varepsilon_n \\ s_1 & \cdots & s_n \end{pmatrix}.$

We recall the power sums studied by Thakur [48]. For $d \in \mathbb{Z}$ and for $\mathfrak{s} = (s_1, \ldots, s_n) \in \mathbb{N}^n$ we introduce

$$S_d(\mathfrak{s}) = \sum_{\substack{a_1, \dots, a_n \in A_+ \\ d = \deg a_1 > \dots > \deg a_n \ge 0}} \frac{1}{a_1^{s_1} \dots a_n^{s_n}} \in K$$

and

$$S_{ \deg a_1 > \dots > \deg a_n \ge 0}} \frac{1}{a_1^{s_1} \dots a_n^{s_n}} \in K.$$

We also recall the following analogues of the power sums after Harada [25]. For a positive array $\begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix} = \begin{pmatrix} \varepsilon_1 & \cdots & \varepsilon_n \\ s_1 & \cdots & s_n \end{pmatrix}$, we introduce

$$S_d\begin{pmatrix}\varepsilon\\\mathfrak{s}\end{pmatrix} = \sum_{\substack{a_1,\dots,a_n \in A_+\\d = \deg a_1 > \dots > \deg a_n \ge 0}} \frac{\varepsilon_1^{\deg a_1} \dots \varepsilon_n^{\deg a_n}}{a_1^{s_1} \dots a_n^{s_n}} \in K$$

and

$$S_{ \deg a_1 > \dots > \deg a_n \ge 0}} \frac{\varepsilon_1^{\deg a_1} \dots \varepsilon_n^{\deg a_n}}{a_1^{s_1} \dots a_n^{s_n}} \in K$$

One verifies easily the following formulas:

$$S_d \begin{pmatrix} \varepsilon \\ s \end{pmatrix} = \varepsilon^d S_d(s),$$

$$S_d \begin{pmatrix} 1 & \cdots & 1 \\ s_1 & \cdots & s_n \end{pmatrix} = S_d(s_1, \dots, s_n),$$

$$S_{

$$S_d \begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix} = S_d \begin{pmatrix} \varepsilon_1 \\ s_1 \end{pmatrix} S_{$$$$

In [25] Harada introduced the alternating multiple zeta value (AMZV) in positive characteristic by setting

$$\zeta_A \begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix} := \sum_{d \ge 0} S_d \begin{pmatrix} \varepsilon \\ \mathfrak{s} \end{pmatrix} = \sum_{\substack{a_1, \dots, a_n \in A_+ \\ \deg a_1 > \dots > \deg a_n \ge 0}} \frac{\varepsilon_1^{\deg a_1} \dots \varepsilon_n^{\deg a_n}}{a_1^{s_1} \dots a_n^{s_n}} \in K_{\infty}.$$

Then he proved several fundamental properties of these values. In particular, it is shown that the product of two AMZV's can be expressed as a linear combination with coefficients in \mathbb{F}_q of AMZV's. To do so, we first recall Chen's formula in [14]. For integers a, b with $b \geq 0$, we recall the binomial number defined by

$$\binom{a}{b} := \frac{a(a-1)\dots(a-b+1)}{b!}$$

It should be remarked that $\binom{a}{b} = 0$ if $b > a \ge 0$. Then refining the work of Thakur [48], Chen [14, Theorem 3.1] showed that for positive integers r, s and for all $d \in \mathbb{N}$,

$$S_d(r)S_d(s) = S_d(r+s) + \sum_{i+j=r+s} \Delta^j_{r,s}S_d(i,j),$$

where the coefficients $\Delta_{r,s}^{j}$ are defined as in (3.1).

As a direct consequence of Chen's formula, we obtain

Lemma 4.1. Let α, β be two elements in \mathbb{F}_q^* , and let r, s be two positive integers. Then for all $d \in \mathbb{N}$, we have

$$S_d \begin{pmatrix} \alpha \\ r \end{pmatrix} S_d \begin{pmatrix} \beta \\ s \end{pmatrix} = S_d \begin{pmatrix} \alpha \beta \\ r+s \end{pmatrix} + \sum_{i+j=r+s} \Delta^j_{r,s} S_d \begin{pmatrix} \alpha \beta & 1 \\ i & j \end{pmatrix},$$

where the indices i, j are positive integers.

Proof. See for example [25, Lemma 2.3].

5. Shuffle algebra of AMZV's

In the following sections §5 and §6 we construct the Hopf shuffle algebra of AMZV's and note that the extension from the setting of MZV's to that of AMZV's is not straightforward as we have explained in the introduction. The key ingredient of our construction is the notion of so-called horizontal maps that we introduce in this section. These maps will allow us to deal with letters of the same weight (see §5.2). In the following section, we will give direct and explicit formulas for the coproduct and then prove that such formulas verify all desired properties.

5.1. Alphabet and shuffle algebra associated to AMZV's.

We put $I = \{(n, \varepsilon) : n \in \mathbb{N}, \varepsilon \in \mathbb{F}_q^*\}$. Let $\Gamma = \{x_{n,\varepsilon}\}_{n \in \mathbb{N}, \varepsilon \in \mathbb{F}_q^*}$ be the alphabet associated to AMZV's indexed by I (see §3.1 for notation). We define the weights by $w(x_{n,\varepsilon}) = n$. We define $\mathfrak{D} := \mathbb{F}_q \langle \Gamma \rangle$ to be the free \mathbb{F}_q -vector space with basis $\langle \Gamma \rangle$.

Inspired by §3 we define recursively two products on \mathfrak{D} as \mathbb{F}_q -bilinear maps

$$\diamond : \mathfrak{D} \times \mathfrak{D} \longrightarrow \mathfrak{D} \quad \text{and} \quad \sqcup : \mathfrak{D} \times \mathfrak{D} \longrightarrow \mathfrak{D}$$

by setting $1 \diamond \mathfrak{a} = \mathfrak{a} \diamond 1 = \mathfrak{a}, 1 \sqcup \mathfrak{a} = \mathfrak{a} \sqcup 1 = \mathfrak{a}$ and

$$\begin{split} \mathfrak{a} \diamond \mathfrak{b} &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j,1} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})), \\ \mathfrak{a} \sqcup \mathfrak{b} &= x_{a,\alpha}(\mathfrak{a}_{-} \sqcup \mathfrak{b}) + x_{b,\beta}(\mathfrak{a} \sqcup \mathfrak{b}_{-}) + \mathfrak{a} \diamond \mathfrak{b} \end{split}$$

any nonempty words $a h \in \langle \Gamma \rangle$ with a - r, a h - r, ch

for any nonempty words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_{-}, \mathfrak{b} = x_{b,\beta}\mathfrak{b}_{-}$. We call \diamond the diamond product and \sqcup the shuffle product.

Proposition 5.1. The diamond product and the shuffle product on \mathfrak{D} are commutative.

Proof. Let $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ be two arbitrary words. It suffices to show that

$$\mathfrak{a} \diamond \mathfrak{b} = \mathfrak{b} \diamond \mathfrak{a} \quad \text{and} \quad \mathfrak{a} \sqcup \mathfrak{b} = \mathfrak{b} \sqcup \mathfrak{a}$$

We proceed the proof by induction on depth(\mathfrak{a}) + depth(\mathfrak{b}). If one of \mathfrak{a} or \mathfrak{b} is the empty word, then (5.1) holds trivially. We assume that (5.1) holds when depth(\mathfrak{a}) + depth(\mathfrak{b}) < n with $n \in \mathbb{N}$ and $n \geq 2$. We need to show that (5.1) holds when depth(\mathfrak{a}) + depth(\mathfrak{b}) = n.

Indeed, assume that $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_-, \mathfrak{b} = x_{b,\beta}\mathfrak{b}_-$, we have

$$\begin{split} \mathfrak{a} \diamond \mathfrak{b} &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta_{a,b}^{j} x_{i,\alpha\beta}(x_{j,1} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})), \\ \mathfrak{b} \diamond \mathfrak{a} &= x_{b+a,\beta\alpha}(\mathfrak{b}_{-} \sqcup \mathfrak{a}_{-}) + \sum_{i+j=b+a} \Delta_{b,a}^{j} x_{i,\beta\alpha}(x_{j,1} \sqcup (\mathfrak{b}_{-} \sqcup \mathfrak{a}_{-})). \end{split}$$

It follows from the induction hypothesis that $\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-} = \mathfrak{b}_{-} \sqcup \mathfrak{a}_{-}$, hence $\mathfrak{a} \diamond \mathfrak{b} = \mathfrak{b} \diamond \mathfrak{a}$. On the other hand, we have

$$\mathfrak{a} \sqcup \mathfrak{b} = x_{a,\alpha}(\mathfrak{a}_{-} \sqcup \mathfrak{b}) + x_{b,\beta}(\mathfrak{a} \sqcup \mathfrak{b}_{-}) + \mathfrak{a} \diamond \mathfrak{b},$$

$$\mathfrak{b} \sqcup \mathfrak{a} = x_{b,\beta}(\mathfrak{b}_{-} \sqcup \mathfrak{a}) + x_{a,\alpha}(\mathfrak{b} \sqcup \mathfrak{a}_{-}) + \mathfrak{b} \diamond \mathfrak{a}.$$

It follows from the induction hypothesis and the above arguments that $\mathfrak{a}_{-} \sqcup \mathfrak{b} = \mathfrak{b} \sqcup \mathfrak{a}_{-}, \mathfrak{a} \sqcup \mathfrak{b}_{-} = \mathfrak{b}_{-} \sqcup \mathfrak{a}$ and $\mathfrak{a} \diamond \mathfrak{b} = \mathfrak{b} \diamond \mathfrak{a}$, hence $\mathfrak{a} \sqcup \mathfrak{b} = \mathfrak{b} \sqcup \mathfrak{a}$. This proves the proposition.

5.2. Horizontal maps.

In this section, we introduce horizontal maps which are crucial in the sequel. For each element $\alpha \in \mathbb{F}_q^*$, we consider the \mathbb{F}_q -linear map

$$\varphi_{\alpha} \colon \mathfrak{C} \longrightarrow \mathfrak{D},$$

which maps the empty word $1 \in \langle \Sigma \rangle$ to the empty word $1 \in \langle \Gamma \rangle$ and maps a nonempty word $x_{i_1}x_{i_2}\ldots x_{i_n}$ to $x_{i_1,\alpha}x_{i_2,1}\ldots x_{i_n,1}$. For all nonempty words $\mathfrak{a} \in \langle \Sigma \rangle$ with $\mathfrak{a} = x_a \mathfrak{a}_-$, we get

$$\varphi_{\alpha}(\mathfrak{a}) = \varphi_{\alpha}(x_a)\varphi_1(\mathfrak{a}_-) = x_{a,\alpha}\varphi_1(\mathfrak{a}_-).$$

As $w(\varphi_{\alpha}(\mathfrak{a})) = w(\mathfrak{a})$ for all $\alpha \in \mathbb{F}_q^*$ and $\mathfrak{a} \in \mathfrak{C}$, we call these maps horizontal maps. We note that they are all injective.

Lemma 5.2. Let $\mathfrak{a}, \mathfrak{b}$ be two words in $\langle \Sigma \rangle$. Then we have

$$\varphi_1(\mathfrak{a} \diamond \mathfrak{b}) = \varphi_1(\mathfrak{a}) \diamond \varphi_1(\mathfrak{b}) \quad and \quad \varphi_1(\mathfrak{a} \sqcup \mathfrak{b}) = \varphi_1(\mathfrak{a}) \sqcup \varphi_1(\mathfrak{b}).$$

Proof. The proof can be done by induction on depth(\mathfrak{a}) + depth(\mathfrak{b}). We omit the details.

As a direct consequence, φ_1 is a homomorphism of algebras from (\mathfrak{C},\diamond) to (\mathfrak{D},\diamond) (resp. from (\mathfrak{C},\sqcup) to (\mathfrak{D},\sqcup)).

Notation 5.3. Since φ_1 is injective, from now on for $\mathfrak{a} \in \mathfrak{C}$ we write $\mathfrak{a} \in \mathfrak{D}$ instead of $\varphi_1(\mathfrak{a}) \in \mathfrak{D}$.

Further, we extend the horizontal map as follows:

$$\varphi_{\alpha} \colon \mathfrak{D} \to \mathfrak{D}; \quad 1 \mapsto 1, \quad \mathfrak{u} \mapsto x_{u,\alpha\varepsilon}\mathfrak{u}_{-}$$

for nonempty words $\mathfrak{u} \in \langle \Gamma \rangle$ with $\mathfrak{u} = x_{u,\varepsilon}\mathfrak{u}_-$. This is a natural extension under the identification of $\mathfrak{a} \in \mathfrak{C}$ and $\varphi_1(\mathfrak{a}) \in \mathfrak{D}$. We note that if $\mathfrak{u} \in \langle \Gamma \rangle$ is a nonempty word written as $\mathfrak{u} = x_{u,\varepsilon}\mathfrak{u}_-$, then for all $\alpha \in \mathbb{F}_q^*$,

$$\varphi_{\alpha}(\mathfrak{u}) = \varphi_{\alpha}(x_{u,\varepsilon})\mathfrak{u}_{-}.$$

Under this extension, the above lemma holds for $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$.

Lemma 5.4. Let α, β be elements of \mathbb{F}_q^* , and let $\mathfrak{a}, \mathfrak{b}$ be nonempty words in $\langle \Sigma \rangle$. We have

$$\varphi_{\alpha}(\mathfrak{a}) \diamond \varphi_{\beta}(\mathfrak{b}) = \varphi_{\alpha\beta}(\mathfrak{a} \diamond \mathfrak{b}).$$

Further, for nonempty words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$,

$$\varphi_{\alpha}(\mathfrak{a}) \diamond \varphi_{\beta}(\mathfrak{b}) = \varphi_{\alpha\beta}(\mathfrak{a} \diamond \mathfrak{b}).$$

Proof. Consider the horizontal maps defined on $\langle \Sigma \rangle$. We write $\mathfrak{a} = x_a \mathfrak{a}_-, \mathfrak{b} = x_b \mathfrak{b}_-$ with $x_a, x_b \in \Sigma$ and $\mathfrak{a}_-, \mathfrak{b}_- \in \langle \Sigma \rangle$. From Lemma 5.2, we have

$$\begin{split} \varphi_{\alpha\beta}(\mathfrak{a} \diamond \mathfrak{b}) &= \varphi_{\alpha\beta} \left(x_{a+b}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \right) \\ &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \\ &= x_{a,\alpha} \mathfrak{a}_{-} \diamond x_{b,\beta} \mathfrak{b}_{-} \\ &= \varphi_{\alpha}(\mathfrak{a}) \diamond \varphi_{\beta}(\mathfrak{b}). \end{split}$$

The proof for the horizontal maps on $\langle \Gamma \rangle$ is parallel. Let $\mathfrak{a} = x_{a,\varepsilon_1}\mathfrak{a}_-, \mathfrak{b} = x_{b,\varepsilon_2}\mathfrak{b}_$ with $x_{a,\varepsilon_1}, x_{b,\varepsilon_2} \in \Gamma$ and $\mathfrak{a}_-, \mathfrak{b}_- \in \langle \Gamma \rangle$. From Lemma 5.2 and its following remark, we have

$$\begin{split} \varphi_{\alpha\beta}(\mathfrak{a} \diamond \mathfrak{b}) &= \varphi_{\alpha\beta} \left(x_{a+b,\varepsilon_1\varepsilon_2}(\mathfrak{a}_- \sqcup \mathfrak{b}_-) + \sum_{i+j=a+b} \Delta^j_{a,b} x_{i,\varepsilon_1\varepsilon_2}(x_j \sqcup (\mathfrak{a}_- \sqcup \mathfrak{b}_-)) \right) \\ &= x_{a+b,\alpha\beta\varepsilon_1\varepsilon_2}(\mathfrak{a}_- \sqcup \mathfrak{b}_-) + \sum_{i+j=a+b} \Delta^j_{a,b} x_{i,\alpha\beta\varepsilon_1\varepsilon_2}(x_j \sqcup (\mathfrak{a}_- \sqcup \mathfrak{b}_-)) \\ &= x_{a,\alpha\varepsilon_1}\mathfrak{a}_- \diamond x_{b,\beta\varepsilon_2}\mathfrak{b}_- \\ &= \varphi_\alpha(\mathfrak{a}) \diamond \varphi_\beta(\mathfrak{b}). \end{split}$$

This proves the lemma.

We end this section by noting that horizontal maps are associative. More precisely, we prove

Lemma 5.5. Let $\alpha, \beta \in \mathbb{F}_q^*$. For any word $\mathfrak{a} \in \langle \Gamma \rangle$, we have

$$\varphi_{\alpha\beta}(\mathfrak{a}) = \varphi_{\alpha}(\varphi_{\beta}(\mathfrak{a})).$$

Proof. The proof is straightforward from the definition of horizontal maps. \Box

5.3. Associativity.

In this subsection, we first show that the diamond product is associative over Γ (see Proposition 5.6), and then prove the associativity of the diamond product and the shuffle product on \mathfrak{D} (see Proposition 5.8).

Proposition 5.6. Let $x_{a,\alpha}, x_{b,\beta}, x_{c,\gamma}$ be letters in Γ . Then we have

$$(x_{a,\alpha} \diamond x_{b,\beta}) \diamond x_{c,\gamma} = x_{a,\alpha} \diamond (x_{b,\beta} \diamond x_{c,\gamma}).$$

Proof. From Lemma 5.4, we have

$$\begin{aligned} (x_{a,\alpha} \diamond x_{b,\beta}) \diamond x_{c,\gamma} &= (\varphi_{\alpha}(x_a) \diamond \varphi_{\beta}(x_b)) \diamond \varphi_{\gamma}(x_c) \\ &= \varphi_{\alpha\beta}(x_a \diamond x_b) \diamond \varphi_{\gamma}(x_c) \\ &= \varphi_{\alpha\beta\gamma}((x_a \diamond x_b) \diamond x_c), \\ x_{a,\alpha} \diamond (x_{b,\beta} \diamond x_{c,\gamma}) &= \varphi_{\alpha}(x_a) \diamond (\varphi_{\beta}(x_b) \diamond \varphi_{\gamma}(x_c)) \\ &= \varphi_{\alpha}(x_a) \diamond \varphi_{\beta\gamma}(x_b \diamond x_c) \\ &= \varphi_{\alpha\beta\gamma}(x_a \diamond (x_b \diamond x_c)). \end{aligned}$$

From the associativity of the diamond product on \mathfrak{C} (see Theorem 3.1), we have $(x_a \diamond x_b) \diamond x_c = x_a \diamond (x_b \diamond x_c)$. This proves the proposition.

Next, we define recursively a product on \mathfrak{D} as a \mathbb{F}_q -bilinear map

 $\triangleright \colon \mathfrak{D} \times \mathfrak{D} \longrightarrow \mathfrak{D}$

by setting $1 \triangleright \mathfrak{a} = \mathfrak{a} \triangleright 1 = \mathfrak{a}$ and

 $\mathfrak{a} \triangleright \mathfrak{b} = x_{a,\alpha}(\mathfrak{a}_{-} \sqcup \mathfrak{b})$

for any nonempty words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_-$. We call \triangleright the triangle product. The triangle product is neither commutative nor associative, as one verifies at once.

Lemma 5.7. For all words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_-, \mathfrak{b} = x_{b,\beta}\mathfrak{b}_-$, we have

(1) $\mathfrak{a} \diamond \mathfrak{b} = (x_{a,\alpha} \diamond x_{b,\beta}) \triangleright (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}).$ (2) $\mathfrak{a} \sqcup \mathfrak{b} = \mathfrak{a} \triangleright \mathfrak{b} + \mathfrak{b} \triangleright \mathfrak{a} + \mathfrak{a} \diamond \mathfrak{b}.$

Proof. We have

$$\begin{split} \mathfrak{a} \diamond \mathfrak{b} &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \\ &= x_{a+b,\alpha\beta} \triangleright (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b}(x_{i,\alpha\beta}x_{j}) \triangleright (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \\ &= \left(x_{a+b,\alpha\beta} + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta} x_{j} \right) \triangleright (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \\ &= (x_{a,\alpha} \diamond x_{b,\beta}) \triangleright (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}). \end{split}$$

This proves part (1). Part (2) is straightforward from the commutativity of the shuffle product (see Proposition 5.1). This completes the proof. \Box

We now ready to prove the main result of this subsection.

Proposition 5.8. The diamond product and the shuffle product on \mathfrak{D} are associative.

Proof. Let $\mathfrak{a}, \mathfrak{b}, \mathfrak{c} \in \langle \Gamma \rangle$ be arbitrary words. It suffices to show that

$$(\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{b} \diamond \mathfrak{c}) \text{ and } (\mathfrak{a} \sqcup \mathfrak{b}) \sqcup \mathfrak{c} = \mathfrak{a} \sqcup (\mathfrak{b} \sqcup \mathfrak{c}).$$

We proceed the proof by induction on depth(\mathfrak{a}) + depth(\mathfrak{b}) + depth(\mathfrak{c}). If one of $\mathfrak{a}, \mathfrak{b}$ or \mathfrak{c} is the empty word, then (5.3) holds trivially. We assume that (5.3) holds when depth(\mathfrak{a}) + depth(\mathfrak{b}) + depth(\mathfrak{c}) < n with $n \in \mathbb{N}$ and $n \geq 3$. We need to show that (5.3) holds when depth(\mathfrak{a}) + depth(\mathfrak{b}) + depth(\mathfrak{b}) + depth(\mathfrak{c}) = n.

We first show that $(\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{b} \diamond \mathfrak{c})$. Assume that $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_{-}, \mathfrak{b} = x_{b,\beta}\mathfrak{b}_{-} = \mathfrak{c} = x_{c,\gamma}\mathfrak{c}_{-}$. From Lemma 5.7, we have

$$\begin{split} &(\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} \\ &= \left(x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \hspace{-0.1cm} \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \hspace{-0.1cm} \mathfrak{b}_{-})) \right) \diamond \mathfrak{c} \\ &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \diamond \mathfrak{c} + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \hspace{-0.1cm} \mathfrak{b}_{-})) \diamond \mathfrak{c} \end{split}$$

$$= (x_{a+b,\alpha\beta} \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ + \sum_{i+j=a+b} \Delta^{j}_{a,b}(x_{i,\alpha\beta} \diamond x_{c,\gamma}) \triangleright ((x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \sqcup \mathfrak{c}_{-}).$$

For all $i, j \in \mathbb{N}$ with i + j = a + b, it follows from the induction hypothesis that

$$\begin{split} & (x_{i,\alpha\beta} \diamond x_{c,\gamma}) \triangleright ((x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \sqcup \mathfrak{c}_{-}) \\ &= (x_{i,\alpha\beta} \diamond x_{c,\gamma}) \triangleright (x_{j} \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-})) \\ &= \left(x_{i+c,\alpha\beta\gamma} + \sum_{i_{1}+j_{1}=i+c} \Delta_{i,c}^{j_{1}} x_{i_{1},\alpha\beta\gamma} x_{j_{1}} \right) \triangleright (x_{j} \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-})) \\ &= x_{i+c,\alpha\beta\gamma} \triangleright (x_{j} \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-})) \\ &+ \sum_{i_{1}+j_{1}=i+c} \Delta_{i,c}^{j_{1}} x_{i_{1},\alpha\beta\gamma} x_{j_{1}} \triangleright (x_{j} \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-})) \\ &+ \sum_{i_{1}+j_{1}=i+c} \Delta_{i,c}^{j_{1}} x_{i_{1},\alpha\beta\gamma} ((x_{j_{1}} \sqcup x_{j}) \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-})) \\ &= (x_{i+c,\alpha\beta\gamma} x_{j}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &+ \sum_{i_{1}+j_{1}=i+c} \Delta_{i,c}^{j_{1}} x_{i_{1},\alpha\beta\gamma} (x_{j_{1}} \sqcup x_{j}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &= \left(x_{i+c,\alpha\beta\gamma} x_{j} + \sum_{i_{1}+j_{1}=i+c} \Delta_{i,c}^{j_{1}} x_{i_{1},\alpha\beta\gamma} (x_{j_{1}} \sqcup x_{j}) \right) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &= ((x_{i,\alpha\beta\gamma} x_{j}) \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}). \end{split}$$

Thus

$$\begin{aligned} (\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} \\ &= (x_{a+b,\alpha\beta} \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &+ \sum_{i+j=a+b} \Delta^{j}_{a,b} ((x_{i,\alpha\beta} x_{j}) \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &= \left((x_{a+b,\alpha\beta} \diamond x_{c,\gamma}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} (x_{i,\alpha\beta} x_{j} \diamond x_{c,\gamma}) \right) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}) \\ &= ((x_{a,\alpha} \diamond x_{b,\beta}) \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-}). \end{aligned}$$

On the other hand, from Proposition 5.1 and the above arguments, we deduce that

$$\begin{split} \mathfrak{a} \diamond (\mathfrak{b} \diamond \mathfrak{c}) &= (\mathfrak{b} \diamond \mathfrak{c}) \diamond \mathfrak{a} \\ &= ((x_{b,\beta} \diamond x_{c,\gamma}) \diamond x_{a,\alpha}) \triangleright ((\mathfrak{b}_{-} \sqcup \hspace{-0.5mm} \mathfrak{c}_{-}) \sqcup \hspace{-0.5mm} \mathfrak{a}_{-}) \\ &= (x_{a,\alpha} \diamond (x_{b,\beta} \diamond x_{c,\gamma})) \triangleright (\mathfrak{a}_{-} \sqcup \hspace{-0.5mm} \mathfrak{(b}_{-} \sqcup \hspace{-0.5mm} \mathfrak{c}_{-})). \end{split}$$

It follows from Proposition 5.6 that $(x_{a,\alpha} \diamond x_{b,\beta}) \diamond x_{c,\gamma} = x_{a,\alpha} \diamond (x_{b,\beta} \diamond x_{c,\gamma})$. Moreover, it follows from the induction hypothesis that $(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}_{-} = \mathfrak{a}_{-} \sqcup (\mathfrak{b}_{-} \sqcup \mathfrak{c}_{-})$. We thus conclude that $(\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{b} \diamond \mathfrak{c})$.

Next, we show that $(\mathfrak{a} \sqcup \mathfrak{b}) \sqcup \mathfrak{c} = \mathfrak{a} \sqcup (\mathfrak{b} \sqcup \mathfrak{c})$. From Lemma 5.7, we have

 $(\mathfrak{a} \sqcup \mathfrak{b}) \sqcup \mathfrak{c} = (\mathfrak{a} \triangleright \mathfrak{b} + \mathfrak{b} \triangleright \mathfrak{a} + \mathfrak{a} \diamond \mathfrak{b}) \sqcup \mathfrak{c}$

$$= (\mathfrak{a} \triangleright \mathfrak{b}) \sqcup \mathfrak{c} + (\mathfrak{b} \triangleright \mathfrak{a}) \sqcup \mathfrak{c} + (\mathfrak{a} \diamond \mathfrak{b}) \sqcup \mathfrak{c}$$

$$= ((\mathfrak{a} \triangleright \mathfrak{b}) \triangleright \mathfrak{c} + \mathfrak{c} \triangleright (\mathfrak{a} \triangleright \mathfrak{b}) + (\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c})$$

$$+ ((\mathfrak{b} \triangleright \mathfrak{a}) \triangleright \mathfrak{c} + \mathfrak{c} \triangleright (\mathfrak{b} \triangleright \mathfrak{a}) + (\mathfrak{b} \triangleright \mathfrak{a}) \diamond \mathfrak{c})$$

$$+ ((\mathfrak{a} \diamond \mathfrak{b}) \triangleright \mathfrak{c} + \mathfrak{c} \triangleright (\mathfrak{a} \diamond \mathfrak{b}) + (\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c})$$

$$= (\mathfrak{a} \triangleright \mathfrak{b}) \triangleright \mathfrak{c} + (\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c} + (\mathfrak{b} \triangleright \mathfrak{a}) \triangleright \mathfrak{c} + (\mathfrak{b} \triangleright \mathfrak{a}) \diamond \mathfrak{c} + (\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c}$$

$$+ (\mathfrak{c} \triangleright (\mathfrak{a} \triangleright \mathfrak{b}) + \mathfrak{c} \triangleright (\mathfrak{b} \triangleright \mathfrak{a}) + \mathfrak{c} \triangleright (\mathfrak{a} \diamond \mathfrak{b})) + (\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c}$$

$$= (\mathfrak{a} \triangleright \mathfrak{b}) \triangleright \mathfrak{c} + (\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c} + (\mathfrak{b} \triangleright \mathfrak{a}) \triangleright \mathfrak{c} + (\mathfrak{b} \triangleright \mathfrak{a}) \diamond \mathfrak{c} + (\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c}$$

and

$$a \sqcup (b \sqcup c) = a \sqcup (b \triangleright c + c \triangleright b + b \diamond c)$$

$$= a \sqcup (b \triangleright c) + a \sqcup (c \triangleright b) + a \sqcup (b \diamond c)$$

$$= (a \triangleright (b \triangleright c) + (b \triangleright c) \triangleright a + a \diamond (b \triangleright c))$$

$$+ (a \triangleright (c \triangleright b) + (c \triangleright b) \triangleright a + a \diamond (c \triangleright b))$$

$$+ (a \triangleright (b \diamond c) + (b \diamond c) \triangleright a + a \diamond (b \diamond c))$$

$$= (b \triangleright c) \triangleright a + a \diamond (b \triangleright c) + (c \triangleright b) \triangleright a + a \diamond (c \triangleright b) + (b \diamond c) \triangleright a$$

$$+ (a \triangleright (b \triangleright c) + a \triangleright (c \triangleright b) + a \triangleright (b \diamond c)) + a \diamond (b \diamond c)$$

$$= (b \triangleright c) \triangleright a + a \diamond (b \triangleright c) + (c \triangleright b) \triangleright a + a \diamond (c \triangleright b) + (b \diamond c) \triangleright a$$

$$+ a \diamond (b \sqcup c) + a \diamond (b \diamond c).$$

We now compare the above expansions. We have showed that $(\mathfrak{a} \diamond \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{b} \diamond \mathfrak{c})$. On the other hand, we have

 $\mathfrak{c} \triangleright (\mathfrak{a} \sqcup \mathfrak{b}) = x_{c,\gamma}(\mathfrak{c}_{-} \sqcup (\mathfrak{a} \sqcup \mathfrak{b}))$ and $(\mathfrak{c} \triangleright \mathfrak{b}) \triangleright \mathfrak{a} = x_{c,\gamma}(\mathfrak{c}_{-} \sqcup \mathfrak{b}) \triangleright \mathfrak{a} = x_{c,\gamma}((\mathfrak{c}_{-} \sqcup \mathfrak{b}) \sqcup \mathfrak{a}).$ From the induction hypothesis and commutativity of shuffle product, one deduces that $\mathfrak{c} \triangleright (\mathfrak{a} \sqcup \mathfrak{b}) = (\mathfrak{c} \triangleright \mathfrak{b}) \triangleright \mathfrak{a}$. Similarly, one deduces that $(\mathfrak{a} \triangleright \mathfrak{b}) \triangleright \mathfrak{c} = \mathfrak{a} \triangleright (\mathfrak{b} \sqcup \mathfrak{c}).$

We have

$$\begin{split} (\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c} &= x_{a,\alpha} (\mathfrak{a}_{-} \sqcup \mathfrak{b}) \diamond \mathfrak{c} = (x_{a,\alpha} \diamond x_{c,\gamma}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}) \sqcup \mathfrak{c}_{-}), \\ \mathfrak{a} \diamond (\mathfrak{c} \triangleright \mathfrak{b}) &= \mathfrak{a} \diamond x_{c,\gamma} (\mathfrak{c}_{-} \sqcup \mathfrak{b}) = (x_{a,\alpha} \diamond x_{c,\gamma}) \triangleright (\mathfrak{a}_{-} \sqcup (\mathfrak{c}_{-} \sqcup \mathfrak{b})). \end{split}$$

From the induction hypothesis and commutativity of shuffle product, one deduces that $(\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{c} \triangleright \mathfrak{b}).$

We have

$$\begin{aligned} (\mathfrak{b} \triangleright \mathfrak{a}) \triangleright \mathfrak{c} &= x_{b,\beta}(\mathfrak{b}_{-} \sqcup \mathfrak{a}) \triangleright \mathfrak{c} \\ &= x_{b,\beta}((\mathfrak{b}_{-} \sqcup \mathfrak{a}) \sqcup \mathfrak{c}) \end{aligned} \quad \text{and} \quad \begin{aligned} (\mathfrak{b} \triangleright \mathfrak{c}) \triangleright \mathfrak{a} &= x_{b,\beta}((\mathfrak{b}_{-} \sqcup \mathfrak{c}) \triangleright \mathfrak{a}) \\ &= x_{b,\beta}((\mathfrak{b}_{-} \sqcup \mathfrak{c}) \sqcup \mathfrak{a}). \end{aligned}$$

From the induction hypothesis and commutativity of shuffle product, one deduces that $(\mathfrak{b} \triangleright \mathfrak{a}) \triangleright \mathfrak{c} = (\mathfrak{b} \triangleright \mathfrak{c}) \triangleright \mathfrak{a}$.

It follows from the induction hypothesis and that

$$\begin{split} (\mathfrak{a} \diamond \mathfrak{b}) \triangleright \mathfrak{c} &= \left(x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \right) \triangleright \mathfrak{c} \\ &= x_{a+b,\alpha\beta}(\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \triangleright \mathfrak{c} + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \triangleright \mathfrak{c} \end{split}$$

$$= x_{a+b,\alpha\beta}((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}((x_{j} \sqcup (\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-})) \sqcup \mathfrak{c})$$

$$= x_{a+b,\alpha\beta}((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta}(x_{j} \sqcup ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}))$$

$$= x_{a+b,\alpha\beta} \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c}) + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta} x_{j} \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c})$$

$$= \left(x_{a+b,\alpha\beta} + \sum_{i+j=a+b} \Delta^{j}_{a,b} x_{i,\alpha\beta} x_{j} \right) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c})$$

$$= (x_{a,\alpha} \diamond x_{b,\beta}) \triangleright ((\mathfrak{a}_{-} \sqcup \mathfrak{b}_{-}) \sqcup \mathfrak{c})$$

and

$$\mathfrak{a} \diamond (\mathfrak{b} \triangleright \mathfrak{c}) = \mathfrak{a} \diamond x_{b,\beta}(\mathfrak{b}_{-} \sqcup \mathfrak{c}) = (x_{a,\alpha} \diamond x_{b,\beta}) \triangleright (\mathfrak{a}_{-} \sqcup (\mathfrak{b}_{-} \sqcup \mathfrak{c})).$$

From the induction hypothesis, one deduces that $(\mathfrak{a} \diamond \mathfrak{b}) \triangleright \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{b} \triangleright \mathfrak{c})$. Similarly, one deduces that $(\mathfrak{b} \triangleright \mathfrak{a}) \diamond \mathfrak{c} = (\mathfrak{b} \diamond \mathfrak{c}) \triangleright \mathfrak{a}$.

From the above arguments, we conclude that $(\mathfrak{a} \sqcup \mathfrak{b}) \sqcup \mathfrak{c} = \mathfrak{a} \sqcup (\mathfrak{b} \sqcup \mathfrak{c})$. This completes the proof.

As for \mathfrak{C} , we define the unit $u \colon \mathbb{F}_q \to \mathfrak{D}$, sending $1 \in \mathbb{F}_q$ to an empty word $1 \in \mathfrak{D}$. Then, as a direct consequence of Proposition 5.1 and Proposition 5.8, we obtain the following result.

Theorem 5.9. The spaces (\mathfrak{D},\diamond) and (\mathfrak{D},\sqcup) are commutative \mathbb{F}_q -algebras.

The following proposition summarizes several properties of different products \triangleright , \diamond and \sqcup that are shown above and will be useful in the sequel.

Proposition 5.10. For all nonempty words $\mathfrak{a}, \mathfrak{b}, \mathfrak{c} \in \langle \Gamma \rangle$, we have

(1)
$$(\mathfrak{a} \triangleright \mathfrak{b}) \triangleright \mathfrak{c} = \mathfrak{a} \triangleright (\mathfrak{b} \sqcup \mathfrak{c}),$$

(2) $(\mathfrak{a} \triangleright \mathfrak{b}) \diamond \mathfrak{c} = \mathfrak{a} \diamond (\mathfrak{c} \triangleright \mathfrak{b}) = (\mathfrak{a} \diamond \mathfrak{c}) \triangleright \mathfrak{b}.$

5.4. Shuffle map for AMZV's in positive characteristic.

For each $d \in \mathbb{Z}$, we define the following two \mathbb{F}_q -linear maps

$$S_{\leq d} \colon \mathfrak{D} \to K_{\infty} \quad \text{and} \quad \zeta_A \colon \mathfrak{D} \to K_{\infty},$$

both of which map the empty word $1 \in \mathfrak{D}$ to $1 \in K_{\infty}$, and the word $x_{s_1,\varepsilon_1} \dots x_{s_r,\varepsilon_r}$ to $S_{\leq d} \begin{pmatrix} \varepsilon_1 & \cdots & \varepsilon_n \\ s_1 & \cdots & s_n \end{pmatrix}$ and $\zeta_A \begin{pmatrix} \varepsilon_1 & \cdots & \varepsilon_n \\ s_1 & \cdots & s_n \end{pmatrix}$ respectively, identifying presentations

$$S_{
$$\zeta_A(x_{s_1,\varepsilon_1}\dots x_{s_r,\varepsilon_r}) = \zeta_A\begin{pmatrix}\varepsilon_1 & \cdots & \varepsilon_n\\s_1 & \cdots & s_n\end{pmatrix}.$$$$

Theorem 5.11. For $\mathfrak{a}, \mathfrak{b} \in \mathfrak{D}$ and $d \in \mathbb{Z}$ we have

 $S_{< d}(\mathfrak{a} \sqcup \mathfrak{b}) = S_{< d}(\mathfrak{a})S_{< d}(\mathfrak{b}).$ $\zeta_A(\mathfrak{a} \sqcup \mathfrak{b}) = \zeta_A(\mathfrak{a})\zeta_A(\mathfrak{b}).$

Proof. The \sqcup product is defined to satisfy this equation. We can check that the \sqcup follows the inductive steps for product of power sums and ζ_A as follows.

When any of \mathfrak{a} and \mathfrak{b} is the empty word, the theorem holds trivially. Let $\mathfrak{a} = x_{a,\alpha}\mathfrak{a}_{-}$ and $\mathfrak{b} = x_{b,\beta}\mathfrak{b}_{-}$ be nonempty words. We proceed the proof by induction on depth(\mathfrak{a}) + depth(\mathfrak{b}). Following the proof of [25, Lemma 2.5] and applying Lemma 4.1,

$$\begin{split} S_{$$

This is parallel to the recursive definition of \sqcup . Proof is the same for ζ_A .

6. Shuffle Hopf Algebra of AMZV's

In §5 we have defined the shuffle algebra structure of AMZV's. This section is devoted to the construction of the Hopf algebra structure of AMZV's. In §6.1 we present the coproduct for the word algebra \mathfrak{D} . In §6.2 we prove the compatibility of the coproduct in Theorem 6.5. Unlike the MZV's case, the proof of this theorem uses the induction on depth instead of that on weight. Then in §6.3 we prove the coassociativity of the coproduct in Theorem 6.8. Putting all together we get the main result of this paper (see Theorem 6.9).

6.1. Coproduct.

This section is inspired by our previous work [32, §8]. We will introduce the coproduct

$$\Delta:\mathfrak{D}\to\mathfrak{D}\otimes\mathfrak{D}.$$

First, we define the coproduct of depth one. We put

$$\Delta(1) := 1 \otimes 1, \qquad \Delta(x_{1,\varepsilon}) := 1 \otimes x_{1,\varepsilon} + x_{1,\varepsilon} \otimes 1$$

for all $\varepsilon \in \mathbb{F}_q^*$.

Further, we define the coproduct of depth one as follows.

Definition 6.1. For all $n \in \mathbb{N}$ and $\varepsilon \in \mathbb{F}_q^*$, we set

$$\Delta(x_{n,\varepsilon}) = 1 \otimes x_{n,\varepsilon} + \sum_{\substack{r \in \mathbb{N}, \mathfrak{a} \in \langle \Sigma \rangle \\ r+w(\mathfrak{a})=n}} \binom{r + \operatorname{depth}(\mathfrak{a}) - 2}{\operatorname{depth}(\mathfrak{a})} x_{r,\varepsilon} \otimes [\mathfrak{a}].$$

Here we recall that $[\mathfrak{a}]$ is given as in (3.2) and we identify $\mathfrak{a} \in \Sigma$ with $\varphi_1(\mathfrak{a}) \in \Gamma$.

Next, we define the coproduct for words of depth greater than one. Let $n \in \mathbb{N}$ with n > 1 and we suppose that we have defined $\Delta(\mathfrak{u})$ for all words \mathfrak{u} of depth(\mathfrak{u}) < n. We now give a formula for $\Delta(\mathfrak{u})$ for all words \mathfrak{u} with depth(\mathfrak{u}) = n. For such a word \mathfrak{u} , we put $\mathfrak{u} = x_{u,\varepsilon}\mathfrak{v}$ with depth(\mathfrak{v}) = n - 1 < n. Since $x_{u,\varepsilon}$ and \mathfrak{v} are both of depth strictly less than n, we have already defined

$$\Delta(x_{u,\varepsilon}) = 1 \otimes x_{u,\varepsilon} + \sum a_{u,\varepsilon} \otimes b_{u,\varepsilon}, \text{ and } \quad \Delta(\mathfrak{v}) = \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}},$$

with $a_{u,\varepsilon} \in \Gamma$ and $b_{u,\varepsilon}, a_{\mathfrak{v}}, b_{\mathfrak{v}} \in \langle \Gamma \rangle$. Then we set

$$\Delta(\mathfrak{u}) := 1 \otimes \mathfrak{u} + \sum (a_{u,\varepsilon} \triangleright a_{\mathfrak{v}}) \otimes (b_{u,\varepsilon} \sqcup b_{\mathfrak{v}})$$
$$= 1 \otimes \mathfrak{u} + \sum (a_{u,\varepsilon}a_{\mathfrak{v}}) \otimes (b_{u,\varepsilon} \sqcup b_{\mathfrak{v}}).$$

Here the last equality holds as $a_{u,\varepsilon} \in \Gamma$.

We end this section by proving the following result:

Proposition 6.2. For any nonempty word $u \in \langle \Gamma \rangle$, we have

$$\Delta\left(\varphi_{\varepsilon}(\mathfrak{u})\right) = (\varphi_{\varepsilon} \otimes \mathrm{Id})\left(\Delta(\mathfrak{u})\right) + (\mathrm{Id} \otimes \varphi_{\varepsilon} - \varphi_{\varepsilon} \otimes \mathrm{Id})(1 \otimes \mathfrak{u}).$$

Proof. Note that for $\mathfrak{a} = x_a \mathfrak{a}_- \in \langle \Gamma \rangle$, $\mathfrak{b} \in \langle \Gamma \rangle$ and $\varepsilon \in \mathbb{F}_q^*$,

$$\varphi_{\varepsilon}(\mathfrak{a}) \triangleright \mathfrak{b} = x_{a,\varepsilon}(\mathfrak{a}_{-} \sqcup \mathfrak{b}) = \varphi_{\varepsilon}(\mathfrak{a} \triangleright \mathfrak{b}).$$

We mention that this does not hold when a = 1.

We first consider the case where depth(\mathfrak{u}) = 1, i.e., \mathfrak{u} belongs to Γ . We put $\mathfrak{u} = x_{n,\alpha}$ with $n \in \mathbb{N}$ and $\alpha \in \mathbb{F}_q^*$. Thus $\varphi_{\varepsilon}(\mathfrak{u}) = x_{n,\varepsilon\alpha} = \varphi_{\varepsilon\alpha}(x_n)$.

Recall that (see Proposition 3.6)

$$\Delta(x_n) = 1 \otimes x_n + \sum_{\substack{r \in \mathbb{N}, \mathfrak{a} \in \langle \Sigma \rangle \\ r+w(\mathfrak{a})=n}} \binom{r + \operatorname{depth}(\mathfrak{a}) - 2}{\operatorname{depth}(\mathfrak{a})} x_r \otimes [\mathfrak{a}].$$

Therefore, Definition 6.1 can be written as

$$\Delta (\mathfrak{u}) = \Delta (\varphi_{\alpha}(x_n))$$

= $(\varphi_{\alpha} \otimes \operatorname{Id}) (\Delta(x_n) - 1 \otimes x_n) + (\operatorname{Id} \otimes \varphi_{\alpha}) (1 \otimes x_n)$
= $(\varphi_{\alpha} \otimes \operatorname{Id}) (\Delta(x_n)) + (\operatorname{Id} \otimes \varphi_{\alpha} - \varphi_{\alpha} \otimes \operatorname{Id}) (1 \otimes x_n).$

Similarly,

$$\Delta(\varphi_{\varepsilon}(\mathfrak{u})) = \Delta(\varphi_{\varepsilon\alpha}(x_n))$$

= $(\varphi_{\varepsilon\alpha} \otimes \mathrm{Id}) (\Delta(x_n) - 1 \otimes x_n) + (\mathrm{Id} \otimes \varphi_{\varepsilon\alpha}) (1 \otimes x_n)$
= $(\varphi_{\varepsilon\alpha} \otimes \mathrm{Id}) (\Delta(x_n)) + (\mathrm{Id} \otimes \varphi_{\varepsilon\alpha} - \varphi_{\varepsilon\alpha} \otimes \mathrm{Id}) (1 \otimes x_n)$

By Lemma 5.5, we conclude that for $\mathfrak{u} = x_{n,\alpha}$,

$$\Delta(\varphi_{\varepsilon}(\mathfrak{u})) = (\varphi_{\varepsilon} \otimes \mathrm{Id})(\Delta(\mathfrak{u})) + (\mathrm{Id} \otimes \varphi_{\varepsilon} - \varphi_{\varepsilon} \otimes \mathrm{Id})(1 \otimes \mathfrak{u}).$$

Next, suppose that $\mathfrak{u} = x_{u,\alpha}\mathfrak{v} \in \langle \Gamma \rangle$ with depth $(\mathfrak{u}) \geq 2$. Let $\Delta(x_u) = 1 \otimes x_u + \sum a_u \otimes b_u$, and $\Delta(\mathfrak{v}) = \sum a_\mathfrak{v} \otimes b_\mathfrak{v}$. We have $a_u \in \Sigma$, $b_u \in \langle \Sigma \rangle$ with $a_u \neq 1$ by Lemma 3.3, and $a_\mathfrak{v}, b_\mathfrak{v} \in \langle \Gamma \rangle$. We have seen that

$$\Delta(x_{u,\alpha}) = 1 \otimes x_{u,\alpha} + \sum a_{u,\alpha} \otimes b_u$$

By definition,

$$\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum (a_{u,\alpha} \triangleright a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}})$$
$$= 1 \otimes \mathfrak{u} + \sum (a_{u,\alpha} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}}).$$

The last equality follows from the fact that $depth(a_{u,\alpha}) = 1$.

Since
$$\varphi_{\varepsilon}(\mathfrak{u}) = x_{u,\varepsilon\alpha}\mathfrak{v}$$
 and $\Delta(\varphi_{\varepsilon}(x_{u,\alpha})) = 1 \otimes \varphi_{\varepsilon}(x_{u,\alpha}) + \sum \varphi_{\varepsilon}(a_{u,\alpha}) \otimes b_{u,\varepsilon\alpha}$
 $\Delta(\varphi_{\varepsilon}(\mathfrak{u})) = \Delta(\varphi_{\varepsilon}(x_{u,\alpha})\mathfrak{v})$

$$\begin{split} \Delta \left(\varphi_{\varepsilon}(\mathfrak{u}) \right) &= \Delta \left(\varphi_{\varepsilon}(x_{u,\alpha}) \mathfrak{v} \right) \\ &= 1 \otimes \varphi_{\varepsilon}(\mathfrak{u}) + \sum \left(\varphi_{\varepsilon}(a_{u,\alpha}) a_{\mathfrak{v}} \right) \otimes \left(b_u \sqcup b_{\mathfrak{v}} \right) \\ &= 1 \otimes \varphi_{\varepsilon}(\mathfrak{u}) + \sum \varphi_{\varepsilon} \left(a_{u,\alpha} a_{\mathfrak{v}} \right) \otimes \left(b_u \sqcup b_{\mathfrak{v}} \right). \end{split}$$

Thus we get

$$\Delta\left(\varphi_{\varepsilon}(\mathfrak{u})\right) = (\varphi_{\varepsilon} \otimes \mathrm{Id})\left(\Delta(\mathfrak{u})\right) + (\mathrm{Id} \otimes \varphi_{\varepsilon} - \varphi_{\varepsilon} \otimes \mathrm{Id})(1 \otimes \mathfrak{u}).$$

6.2. Compatibility.

6.2.1. Compatibility for letters.

We want to prove the following result (for MZV's, see [32, Proposition 7.5]):

Proposition 6.3. Let $u, v \in \mathbb{N}$ and $\alpha, \beta \in \mathbb{F}_q^*$. Then

$$\Delta(x_{u,\alpha} \sqcup \sqcup x_{v,\beta}) = \Delta(x_{u,\alpha}) \sqcup \Delta(x_{v,\beta}).$$

Proof. Let

$$\Delta(x_u) = 1 \otimes x_u + \sum a_u \otimes b_u, \quad \Delta(x_v) = 1 \otimes x_v + \sum a_v \otimes b_v.$$

By Proposition 6.2, we have

$$\Delta(x_{u,\alpha}) = 1 \otimes x_{u,\alpha} + \sum a_{u,\alpha} \otimes b_u, \quad \Delta(x_{v,\beta}) = 1 \otimes x_{v,\beta} + \sum a_{v,\beta} \otimes b_v,$$

where $a_{u,\alpha} = \varphi_{\alpha}(a_u)$ and $a_{v,\beta} = \varphi_{\beta}(a_v)$. Here we recall that a_u and a_v are of depth 1.

Since $x_{u,\alpha} \sqcup \sqcup x_{v,\beta} = x_{u,\alpha} x_{v,\beta} + x_{v,\beta} x_{u,\alpha} + \varphi_{\alpha\beta} (x_u \diamond x_v)$ (by Lemma 5.4),

$$\Delta (x_{u,\alpha} \sqcup x_{v,\beta}) = 1 \otimes x_{u,\alpha} x_{v,\beta} + \sum a_{u,\alpha} \otimes (b_u \sqcup x_{v,\beta}) + \sum (a_{u,\alpha} a_{v,\beta}) \otimes (b_u \sqcup b_v) + 1 \otimes x_{v,\beta} x_{u,\alpha} + \sum a_{v,\beta} \otimes (b_v \sqcup x_{u,\alpha}) + \sum (a_{v,\beta} a_{u,\alpha}) \otimes (b_v \sqcup b_u) + \Delta (\varphi_{\alpha\beta} (x_u \diamond x_v)).$$

On the other hand,

$$\Delta(x_{u,\alpha}) \sqcup \Delta(x_{v,\beta}) = 1 \otimes (x_{u,\alpha} \sqcup x_{v,\beta}) + \sum a_{u,\alpha} \otimes (b_u \sqcup x_{v,\beta}) + \sum a_{v,\beta} \otimes (b_v \sqcup x_{u,\alpha}) + \sum (a_{u,\alpha} \sqcup a_{v,\beta}) \otimes (b_u \sqcup b_v).$$

By expanding \sqcup in each tensor and and canceling, the proposition holds when

$$\Delta(\varphi_{\alpha\beta}(x_u \diamond x_v)) = 1 \otimes \varphi_{\alpha\beta}(x_u \diamond x_v) + \sum (a_{u,\alpha} \diamond a_{v,\beta}) \otimes (b_u \sqcup b_v)$$

= 1 \otimes \varphi_{\alpha\beta}(x_u \otimes x_v) + (\varphi_{\alpha\beta} \otimes \operatorname{Id}) \left(\sum (a_u \otimes a_v) \otimes (b_u \leftur b_v) \right).

By Proposition 6.2, the proposition is reduced into

$$\Delta(x_u \diamond x_v) = 1 \otimes (x_u \diamond x_v) + \sum (a_u \diamond a_v) \otimes (b_u \sqcup b_v),$$

which is a special case of Lemma 3.4.

6.2.2. Compatibility for words of arbitrary depth.

Lemma 6.4. Let $\mathfrak{u}, \mathfrak{v} \in \langle \Gamma \rangle$. Then $\mathfrak{u} \sqcup \mathfrak{v}$ can be presented as the sum of words whose depths are all less than or equal to depth(\mathfrak{u}) + depth(\mathfrak{v}).

Proof. We proceed with the induction on the total depth depth(\mathfrak{u}) + depth(\mathfrak{v}).

The lemma is trivial when \mathfrak{u} or \mathfrak{v} is trivial. If $\mathfrak{u} = x_{u,\alpha}$ and $\mathfrak{v} = x_{v,\beta}$, then the lemma is also immediate from the expression

$$\mathfrak{u} \sqcup \mathfrak{v} = x_{u+v,\alpha\beta} + x_{u,\alpha} x_{v,\beta} + x_{v,\beta} x_{u,\alpha} + \sum_{i+j=u+v} \Delta^j_{u,v} x_{i,\alpha\beta} x_j.$$

So we assume that $\operatorname{depth}(\mathfrak{u}), \operatorname{depth}(\mathfrak{v}) \geq 1$ with at least one of \mathfrak{u} and \mathfrak{v} has $\operatorname{depth} > 1$.

Now assume the lemma holds for all $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ such that $\operatorname{depth}(\mathfrak{a}) + \operatorname{depth}(\mathfrak{b}) < \operatorname{depth}(\mathfrak{u}) + \operatorname{depth}(\mathfrak{b})$. Let $\mathfrak{u} = x_{u,\alpha}\mathfrak{u}_{-}, \mathfrak{v} = x_{v,\beta}\mathfrak{v}_{-}$. Then

$$\begin{aligned} x_{u,\alpha}\mathfrak{u}_{-} \amalg x_{v,\beta}\mathfrak{v}_{-} &= x_{u+v,\alpha\beta}(\mathfrak{u}_{-} \amalg \mathfrak{v}_{-}) + x_{u,\alpha}(\mathfrak{u}_{-} \amalg \mathfrak{v}) + x_{v,\beta}(\mathfrak{u} \amalg \mathfrak{v}_{-}) \\ &+ \sum_{i+j=u+v} \Delta^{j}_{u,v} x_{i,\alpha\beta}(x_{j} \amalg \mathfrak{u}_{-} \amalg \mathfrak{v}_{-}) \end{aligned}$$

From the induction hypothesis, the first three terms can be presented as a sum of words with depths less than or equal to depth(\mathfrak{u}) + depth(\mathfrak{v}). For each j, $x_j \sqcup \mathfrak{u}_- = \mathfrak{u}_- \sqcup x_j$ can be presented as a sum of words whose depths are less than or equal to depth(\mathfrak{u}) from the induction hypothesis. Therefore, $x_i(x_j \sqcup \mathfrak{u}_- \sqcup \mathfrak{v}_-)$ can be presented as sum of words with depth depth(\mathfrak{u}) + depth(\mathfrak{v}). This proves the lemma.

The proof of the compatibility of Δ and \sqcup for $\langle \Gamma \rangle$ is parallel to that for $\langle \Sigma \rangle$.

Theorem 6.5. Let $\mathfrak{u}, \mathfrak{v} \in \langle \Gamma \rangle$. Then Δ and \sqcup are compatible for $\mathfrak{u}, \mathfrak{v}$, that means

$$\Delta(\mathfrak{u} \sqcup \mathfrak{v}) = \Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}).$$

Note that the theorem is immediate when \mathfrak{u} or \mathfrak{v} is trivial. Also, depth(\mathfrak{u}) = depth(\mathfrak{v}) = 1 case is shown in Proposition 6.3. Before proving this theorem, we need some preparatory results.

Lemma 6.6. Let $\mathfrak{u} = x_{u,\alpha}\mathfrak{u}_-$, $\mathfrak{v} = x_{v,\beta}\mathfrak{v}_-$ be two words in $\langle \Gamma \rangle$, whose depths are ≥ 1 . Assume that Δ and \sqcup are compatible for all words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with depth(\mathfrak{a}) + depth(\mathfrak{b}) < depth(\mathfrak{u}) + depth(\mathfrak{v}). Further, we suppose that $\Delta(x_u\mathfrak{u}_- \sqcup x_v\mathfrak{v}_-) = \Delta(x_u\mathfrak{u}_-) \sqcup \Delta(x_v\mathfrak{v}_-)$.

Then we have

$$\Delta(\mathfrak{u} \sqcup \mathfrak{v}) = \Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}).$$

Proof. Let

$$\begin{split} \Delta(x_u) &= 1 \otimes x_u + \sum a_u \otimes b_u, \qquad \quad \Delta(x_v) = 1 \otimes x_v + \sum a_v \otimes b_v, \\ \Delta(\mathfrak{u}_-) &= \sum a_{\mathfrak{u}_-} \otimes b_{\mathfrak{u}_-}, \qquad \qquad \Delta(\mathfrak{v}_-) = \sum a_{\mathfrak{v}_-} \otimes b_{\mathfrak{v}_-}, \end{split}$$

with $x_u, x_v, a_u, a_v \in \Sigma$, $b_u, b_v \in \langle \Sigma \rangle$ and $a_{\mathfrak{u}_-}, b_{\mathfrak{u}_-}, a_{\mathfrak{v}_-}, b_{\mathfrak{v}_-} \in \langle \Gamma \rangle$. We have

$$\Delta(x_{u,\alpha}) = 1 \otimes x_{u,\alpha} + \sum a_{u,\alpha} \otimes b_u,$$

$$\Delta(x_{v,\beta}) = 1 \otimes x_{v,\beta} + \sum a_{v,\beta} \otimes b_v,$$

where $a_{u,\alpha} = \varphi_{\alpha}(a_u), a_{v,\beta} = \varphi_{\beta}(a_v)$. Note that

$$\Delta(x_{u,\alpha}\mathfrak{u}_{-}) = 1 \otimes x_{u,\alpha}\mathfrak{u}_{-} + \sum a_{u,\alpha}a_{\mathfrak{u}_{-}} \otimes (b_{u} \sqcup b_{\mathfrak{u}_{-}}),$$

$$\Delta(x_{v,\beta}\mathfrak{v}_{-}) = 1 \otimes x_{v,\beta}\mathfrak{v}_{-} + \sum a_{v,\beta}a_{\mathfrak{v}_{-}} \otimes (b_{v} \sqcup b_{\mathfrak{v}_{-}}).$$

Therefore, we have

(6.1)

$$\Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}) = \Delta(x_{u,\alpha}\mathfrak{u}_{-}) \sqcup \Delta(x_{v,\beta}\mathfrak{v}_{-})$$

$$= \left(1 \otimes x_{u,\alpha}\mathfrak{u}_{-} + \sum a_{u,\alpha}a_{\mathfrak{u}_{-}} \otimes (b_{u} \sqcup b_{\mathfrak{u}_{-}})\right) \sqcup \left(1 \otimes x_{v,\beta}\mathfrak{v}_{-} + \sum a_{v,\beta}a_{\mathfrak{v}_{-}} \otimes (b_{v} \sqcup b_{\mathfrak{v}_{-}})\right)$$

$$= 1 \otimes (x_{u,\alpha}\mathfrak{u}_{-} \sqcup x_{v,\beta}\mathfrak{v}_{-}) + \sum a_{v,\beta}a_{\mathfrak{v}_{-}} \otimes (x_{u,\alpha}\mathfrak{u}_{-} \sqcup b_{v} \sqcup b_{\mathfrak{v}_{-}})$$

$$+ \sum a_{u,\alpha}a_{\mathfrak{u}_{-}} \otimes (x_{v,\beta}\mathfrak{v}_{-} \sqcup b_{u} \sqcup b_{\mathfrak{u}_{-}}) + \sum (a_{u,\alpha}a_{\mathfrak{u}_{-}} \sqcup a_{v,\beta}a_{\mathfrak{v}_{-}}) \otimes (b_{u} \sqcup b_{v} \sqcup b_{\mathfrak{u}_{-}} \sqcup b_{\mathfrak{v}_{-}}).$$
On the other hand, from Lemma 5.4 and Lemma 5.7

 $\mathfrak{u} \sqcup \mathfrak{v} = \mathfrak{u} \triangleright \mathfrak{v} + \mathfrak{v} \triangleright \mathfrak{u} + \varphi_{\alpha\beta}(x_u \mathfrak{u}_- \diamond x_v \mathfrak{v}_-),$

thus we have

(6.2)
$$\Delta(\mathfrak{u}\sqcup\mathfrak{v}) = S_1 + S_2 + S_3$$

where

$$\begin{split} S_1 &= \Delta(\mathfrak{u} \triangleright \mathfrak{v}) = \Delta(x_{u,\alpha}(\mathfrak{u}_- \sqcup \mathfrak{v})), \\ S_2 &= \Delta(\mathfrak{v} \triangleright \mathfrak{u}) = \Delta(x_{v,\beta}(\mathfrak{v}_- \sqcup \iota \mathfrak{u})), \\ S_3 &= \Delta(\varphi_{\alpha\beta}(x_u\mathfrak{u}_- \diamond x_v\mathfrak{v}_-)) \\ &= (\varphi_{\alpha\beta} \otimes \mathrm{Id})\Delta(x_u\mathfrak{u}_- \diamond x_v\mathfrak{v}_-) + (\mathrm{Id} \otimes \varphi_{\alpha\beta} - \varphi_{\alpha\beta} \otimes \mathrm{Id})(1 \otimes (x_u\mathfrak{u}_- \diamond x_v\mathfrak{v}_-)). \end{split}$$

We calculate S_1 and S_2 . Since $depth(\mathfrak{u}_-) + depth(\mathfrak{v}) < depth(\mathfrak{u}) + depth(\mathfrak{v})$, the induction hypothesis yields

$$\begin{aligned} \Delta(\mathfrak{u}_{-} \sqcup \mathfrak{v}) &= \Delta(\mathfrak{u}_{-}) \sqcup \Delta(x_{v,\beta}\mathfrak{v}_{-}) \\ &= \left(\sum a_{\mathfrak{u}_{-}} \otimes b_{\mathfrak{u}_{-}}\right) \sqcup \left(1 \otimes x_{v,\beta}\mathfrak{v}_{-} + \sum a_{v,\beta}a_{\mathfrak{v}_{-}} \otimes (b_{v} \sqcup b_{\mathfrak{v}_{-}})\right) \\ &= \sum a_{\mathfrak{u}_{-}} \otimes (b_{\mathfrak{u}_{-}} \sqcup x_{v,\beta}\mathfrak{v}_{-}) + \sum (a_{\mathfrak{u}_{-}} \sqcup a_{v,\beta}a_{\mathfrak{v}_{-}}) \otimes (b_{v} \sqcup b_{\mathfrak{u}_{-}} \sqcup b_{\mathfrak{v}_{-}}), \end{aligned}$$

 \mathbf{SO}

$$S_{1} = 1 \otimes (x_{u,\alpha}(\mathfrak{u}_{-} \sqcup \mathfrak{v})) + \sum a_{u,\alpha}a_{\mathfrak{u}_{-}} \otimes (b_{u} \sqcup b_{\mathfrak{u}_{-}} \sqcup x_{v,\beta}\mathfrak{v}_{-}) + \sum a_{u,\alpha}(a_{\mathfrak{u}_{-}} \sqcup a_{v,\beta}a_{\mathfrak{v}_{-}}) \otimes (b_{u} \sqcup b_{v} \sqcup b_{\mathfrak{u}_{-}} \sqcup b_{\mathfrak{v}_{-}}).$$

A similar calculation yields

$$S_{2} = 1 \otimes (x_{v,\beta}(\mathfrak{u} \sqcup \mathfrak{v}_{-})) + \sum a_{v,\beta}a_{\mathfrak{v}_{-}} \otimes (b_{v} \sqcup b_{\mathfrak{v}_{-}} \sqcup x_{u,\alpha}\mathfrak{u}_{-})$$
$$+ \sum a_{v,\beta}(a_{\mathfrak{v}_{-}} \sqcup u_{u,\alpha}a_{\mathfrak{u}_{-}}) \otimes (b_{u} \sqcup b_{v} \sqcup b_{\mathfrak{u}_{-}} \sqcup b_{\mathfrak{v}_{-}}).$$

What we want to show is that two equations (6.1) and (6.2) coincide, that is, $X := \Delta(\mathfrak{u} \sqcup \mathfrak{v}) - \Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}) = 0.$

First, we collect the terms of the form $1 \otimes \Box$. The equation for the right tensor ands of such terms in X is

$$\begin{aligned} x_{u,\alpha}(\mathfrak{u}_{-}\sqcup \mathfrak{v}) + x_{v,\beta}(\mathfrak{u}\sqcup \mathfrak{v}_{-}) + \varphi_{\alpha\beta}(x_{u}\mathfrak{u}_{-}\diamond x_{v}\mathfrak{v}_{-}) - (x_{u}\mathfrak{u}_{-}\diamond x_{v}\mathfrak{v}_{-}) - \mathfrak{u}\sqcup \mathfrak{v} \\ &= x_{u,\alpha}(\mathfrak{u}_{-}\sqcup \mathfrak{v}) + x_{v,\beta}(\mathfrak{u}\sqcup \mathfrak{v}_{-}) + \varphi_{\alpha\beta}(x_{u}\mathfrak{u}_{-}\diamond x_{v}\mathfrak{v}_{-}) - (x_{u}\mathfrak{u}_{-}\diamond x_{v}\mathfrak{v}_{-}) \\ &- \mathfrak{u} \triangleright \mathfrak{v} - \mathfrak{v} \triangleright \mathfrak{u} - \mathfrak{u} \diamond \mathfrak{v} \\ &= -(x_{u}\mathfrak{u}_{-}\diamond x_{v}\mathfrak{v}_{-}) \end{aligned}$$

by Lemma 5.4, that is, the terms of the form $1 \otimes \Box$ in X are collected as

$$-1\otimes(x_u\mathfrak{u}_-\diamond x_v\mathfrak{v}_-).$$

Next we collect the terms of the form $\sum \Box \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}_-} \sqcup b_{\mathfrak{v}_-})$. For fixed indices, the equation for the left tensorands of such terms in X is

$$\begin{aligned} a_{u,\alpha}(a_{\mathfrak{u}_{-}} \sqcup u_{v,\beta}a_{\mathfrak{v}_{-}}) + a_{v,\beta}(a_{\mathfrak{v}_{-}} \sqcup u_{u,\alpha}a_{\mathfrak{u}_{-}}) - (a_{u,\alpha}a_{\mathfrak{u}_{-}} \sqcup u_{v,\beta}a_{\mathfrak{v}_{-}}) \\ &= -\varphi_{\alpha\beta}(a_{u}a_{\mathfrak{u}_{-}} \diamond a_{v}a_{\mathfrak{v}_{-}}), \end{aligned}$$

by expanding the shuffle product of the third term and reducing it. That is, the terms of such a form in X are collected as

$$-(\varphi_{\alpha\beta}\otimes \mathrm{Id})\sum(a_ua_{\mathfrak{u}_-}\diamond a_va_{\mathfrak{v}_-})\otimes(b_u\sqcup b_v\sqcup b_{\mathfrak{u}_-}\sqcup b_{\mathfrak{v}_-}).$$

Some of the remaining terms cancel each other. One can check that

$$\begin{aligned} X = & (\varphi_{\alpha\beta} \otimes \mathrm{Id}) \Delta(x_u \mathfrak{u}_- \diamond x_v \mathfrak{v}_-) - 1 \otimes (x_u \mathfrak{u}_- \diamond x_v \mathfrak{v}_-) \\ & - (\varphi_{\alpha\beta} \otimes \mathrm{Id}) \sum (a_u a_{\mathfrak{u}_-} \diamond a_v a_{\mathfrak{v}_-}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}_-} \sqcup b_{\mathfrak{v}_-}), \end{aligned}$$

which vanishes when

(6.3)

$$\Delta(x_u\mathfrak{u}_-\diamond x_v\mathfrak{v}_-)=1\otimes(x_u\mathfrak{u}_-\diamond x_v\mathfrak{v}_-)+\sum(a_ua_{\mathfrak{u}_-}\diamond a_va_{\mathfrak{v}_-})\otimes(b_u\sqcup b_v\sqcup b_{\mathfrak{u}_-}\sqcup b_{\mathfrak{v}_-})$$

Recall the assumption of the lemma, $\Delta(x_u\mathfrak{u}_-\sqcup x_v\mathfrak{v}_-) = \Delta(x_u\mathfrak{u}_-)\sqcup\Delta(x_v\mathfrak{v}_-)$. Expanding this by using definitions of \sqcup , \diamond and calculating, we can conclude the desired Equation (6.3). Thus the lemma holds.

Proposition 6.7. Let $x_u \mathfrak{u}$, $x_v \mathfrak{v}$ be two words in $\langle \Gamma \rangle$, whose first letters are with trivial character $1 \in \mathbb{F}_q^*$. If Δ and \sqcup are compatible for all words $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with depth (\mathfrak{a}) + depth (\mathfrak{b}) < depth $(x_u \mathfrak{u})$ + depth $(x_v \mathfrak{v})$, then

$$\Delta(x_u\mathfrak{u}\sqcup u \sqcup x_v\mathfrak{v}) = \Delta(x_u\mathfrak{u}) \sqcup \Delta(x_v\mathfrak{v}).$$

The proof closely follows that of [32, Proposition 7.4], with the only difference being that the induction step is now based on depth rather than weight induction. For the sake of clarity, we include a complete proof.

Note that the proof is valid for $\mathfrak{u} = 1$ or $\mathfrak{v} = 1$ cases. In those cases, some sum presentation is vacuous, i.e., if $\mathfrak{u} = 1$, the indexed set $\{(a_{\mathfrak{u}}, b_{\mathfrak{u}})\}$ is the empty set in the sum presentation of $\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum a_{\mathfrak{u}} \otimes b_{\mathfrak{u}}$.

Proof of Proposition 6.7. By the definition of the \square and Lemma 5.7, we have (6.4)

 $x_u \mathfrak{u} \sqcup \sqcup x_v \mathfrak{v}$

$$\begin{split} &= (x_u \mathfrak{u}) \triangleright (x_v \mathfrak{v}) + (x_v \mathfrak{v}) \triangleright (x_u \mathfrak{u}) + (x_u \mathfrak{u}) \diamond (x_v \mathfrak{v}) \\ &= x_v (x_u \mathfrak{u} \sqcup \mathfrak{v}) + x_u (\mathfrak{u} \sqcup (x_v \mathfrak{v})) + (x_u \diamond x_v) \triangleright (\mathfrak{u} \sqcup \mathfrak{v}) \\ &= x_v (x_u \mathfrak{u} \sqcup \mathfrak{v}) + x_u (\mathfrak{u} \sqcup (x_v \mathfrak{v})) + x_{u+v} (\mathfrak{u} \sqcup \mathfrak{v}) + \sum_{i+j=u+v} \Delta^j_{u,v} x_i (x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}). \end{split}$$

Therefore, we get

(6.5)
$$\Delta(x_{u}\mathfrak{u} \sqcup x_{v}\mathfrak{v}) - \Delta(x_{u}\mathfrak{u}) \sqcup \Delta(x_{v}\mathfrak{v})$$
$$= \Delta(x_{v}(x_{u}\mathfrak{u} \sqcup \mathfrak{v})) + \Delta(x_{u}(\mathfrak{u} \sqcup x_{v}\mathfrak{v})) + \Delta(x_{u+v}(\mathfrak{u} \sqcup \mathfrak{v}))$$
$$+ \sum_{i+j=u+v} \Delta_{u,v}^{j} \Delta(x_{i}(x_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v})) - \Delta(x_{u}\mathfrak{u}) \sqcup \Delta(x_{v}\mathfrak{v}).$$

We calculate each term on the right hand side to conclude that this vanishes. We put

$$\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum a_{\mathfrak{u}} \otimes b_{\mathfrak{u}}, \text{ and } \Delta(\mathfrak{v}) = 1 \otimes \mathfrak{v} + \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}}.$$

Also for all $j \in \mathbb{N}$, we put

$$\Delta(x_j) = 1 \otimes x_j + \sum a_j \otimes b_j.$$

In particular,

$$\Delta(x_u) = 1 \otimes x_u + \sum a_u \otimes b_u$$
, and $\Delta(x_v) = 1 \otimes x_v + \sum a_v \otimes b_v$.

The first term $\Delta(x_v(x_u\mathfrak{u}\sqcup\mathfrak{v}))$.

From the definition of Δ ,

(6.6)
$$\Delta(x_u\mathfrak{u}) = 1 \otimes x_u\mathfrak{u} + \sum a_u \otimes (b_u \sqcup \mathfrak{u}) + \sum (a_u \triangleright a_\mathfrak{u}) \otimes (b_u \sqcup b_\mathfrak{u}).$$

As $depth(x_u \mathfrak{u}) + depth(\mathfrak{v}) < depth(x_u \mathfrak{u}) + depth(x_v \mathfrak{v})$, by the induction hypothesis we obtain

$$\begin{split} \Delta(x_{u}\mathfrak{u} \sqcup \mathfrak{v}) = &\Delta(x_{u}\mathfrak{u}) \sqcup \Delta(\mathfrak{v}) \\ = &1 \otimes ((x_{u}\mathfrak{u}) \sqcup \mathfrak{v}) + \sum a_{u} \otimes (b_{u} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum a_{\mathfrak{v}} \otimes ((x_{u}\mathfrak{u}) \sqcup b_{\mathfrak{v}}) \\ &+ \sum (a_{u} \triangleright a_{\mathfrak{u}}) \otimes (b_{u} \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) + \sum (a_{u} \sqcup a_{\mathfrak{v}}) \otimes (b_{u} \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) \\ &+ \sum ((a_{u} \triangleright a_{\mathfrak{u}}) \sqcup a_{\mathfrak{v}}) \otimes (b_{u} \sqcup b_{\mathfrak{u}} \sqcup b_{\mathfrak{v}}). \end{split}$$

Thus

$$(6.7) \qquad \Delta(x_v(x_u\mathfrak{u} \sqcup \mathfrak{v})) = 1 \otimes x_v(x_u\mathfrak{u} \sqcup x_v) + \sum a_v \otimes (b_v \sqcup (x_u\mathfrak{u}) \sqcup \mathfrak{v}) + \sum (a_v \triangleright a_u) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum (a_v \triangleright (a_u \triangleright a_u)) \otimes (b_u \sqcup b_v \sqcup b_u \sqcup \mathfrak{v}) + \sum (a_v \triangleright a_v) \otimes (b_v \sqcup (x_u\mathfrak{u}) \sqcup b_v) + \sum (a_v \triangleright (a_u \sqcup a_v)) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_v) + \sum (a_v \triangleright ((a_u \triangleright a_u)) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_v).$$

The second term $\Delta(x_u(\mathfrak{u} \sqcup (x_v \mathfrak{v})))$.

Similarly, we get

$$(6.8) \qquad \Delta(x_u(\mathfrak{u} \sqcup (x_v \mathfrak{v}))) = 1 \otimes x_u(\mathfrak{u} \sqcup x_v \mathfrak{v}) + \sum a_u \otimes (b_u \sqcup \mathfrak{u} \sqcup (x_v \mathfrak{v})) + \sum ((a_u \triangleright a_v) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum (a_u \triangleright (a_v \triangleright a_v)) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_v) + \sum (a_u \triangleright a_u) \otimes (b_u \sqcup b_u \sqcup (x_v \mathfrak{v})) + \sum (a_u \triangleright (a_u \sqcup a_v)) \otimes (b_u \sqcup b_u \sqcup b_v \sqcup \mathfrak{v}) + \sum (a_u \triangleright (a_u \sqcup (a_v \triangleright a_v)) \otimes (b_u \sqcup b_v \sqcup b_u \sqcup b_v).$$

The third term $\Delta(x_{u+v}(\mathfrak{u} \sqcup \mathfrak{v}))$.

We put

$$\Delta(\mathfrak{u} \sqcup \mathfrak{v}) = 1 \otimes (\mathfrak{u} \sqcup \mathfrak{v}) + \sum a_{\mathfrak{u} \sqcup \mathfrak{v}} \otimes b_{\mathfrak{u} \sqcup \mathfrak{v}}$$

As $depth(\mathfrak{u}) + depth(\mathfrak{v}) < depth(x_u\mathfrak{u}) + depth(x_v\mathfrak{v})$, the induction hypothesis implies that

$$\Delta(\mathfrak{u} \sqcup \mathfrak{v}) = \Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}) = \left(1 \otimes \mathfrak{u} + \sum a_{\mathfrak{u}} \otimes b_{\mathfrak{u}}\right) \sqcup \left(1 \otimes \mathfrak{v} + \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}}\right).$$

Thus

$$1 \otimes (\mathfrak{u} \sqcup \mathfrak{v}) + \sum a_{\mathfrak{u} \sqcup \mathfrak{v}} \otimes b_{\mathfrak{u} \sqcup \mathfrak{v}} = \left(1 \otimes \mathfrak{u} + \sum a_{\mathfrak{u}} \otimes b_{\mathfrak{u}} \right) \sqcup \left(1 \otimes \mathfrak{v} + \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}} \right)$$
$$= 1 \otimes (\mathfrak{u} \sqcup \mathfrak{v}) + \sum a_{\mathfrak{u}} \otimes (b_{\mathfrak{u}} \sqcup \mathfrak{v})$$
$$+ \sum a_{\mathfrak{v}} \otimes (\mathfrak{u} \sqcup b_{\mathfrak{v}}) + \sum (a_{\mathfrak{u}} \sqcup a_{\mathfrak{v}}) \otimes (b_{\mathfrak{u}} \sqcup b_{\mathfrak{v}}),$$

which implies

(6.9)
$$\sum a_{\mathfrak{u}\sqcup\mathfrak{v}}\otimes b_{\mathfrak{u}\sqcup\mathfrak{v}} = \sum a_{\mathfrak{u}}\otimes (b_{\mathfrak{u}}\sqcup\mathfrak{v}) + \sum a_{\mathfrak{v}}\otimes (\mathfrak{u}\sqcup b_{\mathfrak{v}}) + \sum (a_{\mathfrak{u}}\sqcup a_{\mathfrak{v}})\otimes (b_{\mathfrak{u}}\sqcup b_{\mathfrak{v}}).$$

Thus we have

(6.10)
$$\Delta(x_{u+v}(\mathfrak{u}\sqcup\mathfrak{v})) = 1 \otimes (x_{u+v}(\mathfrak{u}\sqcup\mathfrak{v})) + \sum a_{u+v} \otimes (b_{u+v}\sqcup\mathfrak{u}\sqcup\mathfrak{v}) + \sum (a_{u+v} \triangleright a_{u\sqcup\mathfrak{v}}) \otimes (b_{u+v}\sqcup b_{u\sqcup\mathfrak{v}}).$$

The fourth terms $\Delta(x_i(x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}))$ for all $0 \le i, j \le u + v$ with i + j = u + v.

As depth(\mathfrak{u}) + depth(x_j) < depth($x_u\mathfrak{u}$) + depth($x_v\mathfrak{v}$), we have $\Delta(x_j \sqcup \mathfrak{u}) = \Delta(\mathfrak{u} \sqcup x_j) = \Delta(\mathfrak{u}) \sqcup \Delta(x_j)$ by induction hypothesis. Further by Lemma 6.4, $x_j \sqcup \mathfrak{u}$ can be represented as the sum of words with depth less than or equal to depth($x_j\mathfrak{u}$). Thus $\Delta(x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) = \Delta(x_j \sqcup \mathfrak{u}) \sqcup \Delta(\mathfrak{v}) = \Delta(x_j) \sqcup \Delta(\mathfrak{u}) \sqcup \Delta(\mathfrak{v}) = \Delta(x_j) \sqcup \Delta(\mathfrak{u} \sqcup \mathfrak{v})$ by the induction hypothesis again.

Therefore,

$$\begin{split} \Delta(x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) &= \Delta(x_j) \sqcup \Delta(\mathfrak{u} \sqcup \mathfrak{v}) \\ &= \left(1 \otimes x_j + \sum a_j \otimes b_j \right) \sqcup \left(1 \otimes (\mathfrak{u} \sqcup \mathfrak{v}) + \sum a_{\mathfrak{u} \sqcup \mathfrak{v}} \otimes b_{\mathfrak{u} \sqcup \mathfrak{v}} \right) \\ &= 1 \otimes (x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum a_j \otimes (b_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) \\ &+ \sum a_{\mathfrak{u} \sqcup \mathfrak{v}} \otimes (x_j \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}) + \sum (a_j \sqcup a_{\mathfrak{u} \sqcup \mathfrak{v}}) \otimes (b_j \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}). \end{split}$$

This leads to

$$(6.11) \quad \Delta(x_i(x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v})) = 1 \otimes (x_i(x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v})) + \sum a_i \otimes (b_i \sqcup x_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum (a_i \triangleright a_j) \otimes (b_i \sqcup b_j \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum (a_i \triangleright a_{\mathfrak{u} \sqcup \mathfrak{v}}) \otimes (b_i \sqcup x_j \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}) + \sum (a_i \triangleright (a_j \sqcup a_{\mathfrak{u} \sqcup \mathfrak{v}})) \otimes (b_i \sqcup b_j \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}).$$

The last term $\Delta(x_u \mathfrak{u}) \sqcup \Delta(x_v \mathfrak{v})$.

Note that $\Delta(x_u \mathfrak{u})$ is given by (6.6). A parallel calculation gives

$$\Delta(x_{v}\mathfrak{v}) = 1 \otimes x_{v}\mathfrak{v} + \sum a_{v} \otimes (b_{v} \sqcup \mathfrak{v}) + \sum (a_{v} \triangleright a_{\mathfrak{v}}) \otimes (b_{v} \sqcup b_{\mathfrak{v}}).$$

Thus

$$(6.12)$$

$$\Delta(x_{u}\mathfrak{u}) \sqcup \Delta(x_{v}\mathfrak{v})$$

$$= \left(1 \otimes x_{u}\mathfrak{u} + \sum a_{u} \otimes (b_{u} \sqcup \mathfrak{u}) + \sum (a_{u} \triangleright a_{\mathfrak{u}}) \otimes (b_{u} \sqcup b_{\mathfrak{u}})\right)$$

$$\amalg \left(1 \otimes x_{v}\mathfrak{v} + \sum a_{v} \otimes (b_{v} \sqcup \mathfrak{v}) + \sum (a_{v} \triangleright a_{\mathfrak{v}}) \otimes (b_{v} \sqcup b_{\mathfrak{v}})\right)$$

$$= 1 \otimes (x_{u}\mathfrak{u} \sqcup x_{v}\mathfrak{v})$$

$$+ \sum a_{v} \otimes ((x_{u}\mathfrak{u}) \sqcup b_{v} \sqcup \mathfrak{v}) + \sum (a_{v} \triangleright a_{\mathfrak{v}}) \otimes ((x_{u}\mathfrak{u}) \sqcup b_{v} \sqcup b_{\mathfrak{v}})$$

$$+ \sum a_{u} \otimes (b_{u} \sqcup \mathfrak{u} \sqcup (x_{v}\mathfrak{v})) + \sum (a_{u} \sqcup a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u})$$

$$+ \sum (a_{u} \sqcup (a_{v} \triangleright a_{\mathfrak{v}})) \otimes (b_{u} \sqcup \mathfrak{u} \sqcup b_{v} \sqcup b_{\mathfrak{v}})$$

$$+ \sum (a_{u} \sqcup a_{u}) \otimes (b_{u} \sqcup b_{u} \sqcup (x_{v}\mathfrak{v})) + \sum ((a_{u} \triangleright a_{u}) \sqcup a_{v}) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup \mathfrak{v})$$

$$+ \sum ((a_{u} \triangleright a_{u}) \sqcup (a_{v} \triangleright a_{\mathfrak{v}})) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{\mathfrak{v}}).$$

Plugging the equations (6.7), (6.8), (6.10), (6.11), (6.12) into (6.5) yields

$$\Delta(x_u\mathfrak{u}\sqcup x_v\mathfrak{v}) - \Delta(x_u\mathfrak{u})\sqcup \Delta(x_v\mathfrak{v}) = S_0 - S_1 - S_2 - S_\mathfrak{u} - S_\mathfrak{v} + S_3 + S_4.$$

Here the sums S_0 , S_1 , S_2 , S_3 , S_4 , $S_{\mathfrak{u}}$ and $S_{\mathfrak{v}}$ are given as follows: $S_0 = 1 \otimes r_n(r_n\mathfrak{u} \sqcup \mathfrak{u}) + 1 \otimes r_n(\mathfrak{u} \sqcup \mathfrak{u}(r_n\mathfrak{v})) + 1 \otimes r_{n+n}(\mathfrak{u} \sqcup \mathfrak{u})$

$$S_{0} = 1 \otimes x_{v}(x_{u}\mathfrak{u} \sqcup \mathfrak{v}) + 1 \otimes x_{u}(\mathfrak{u} \sqcup (x_{v}\mathfrak{v})) + 1 \otimes x_{u+v}(\mathfrak{u} \sqcup \mathfrak{v})$$

$$+ \sum_{i+j=u+v} \Delta_{u,v}^{j} 1 \otimes x_{i}(x_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) - 1 \otimes ((x_{u}\mathfrak{u}) \sqcup (x_{v}\mathfrak{v})).$$

$$S_{1} = \sum (a_{u} \sqcup a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) - \sum (a_{u} \triangleright a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v})$$

$$- \sum (a_{v} \triangleright a_{u}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}).$$

$$S_{2} = \sum ((a_{u} \triangleright a_{u}) \sqcup (a_{v} \triangleright a_{v})) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$- \sum (a_{u} \triangleright (a_{u} \sqcup (a_{v} \triangleright a_{v}))) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$- \sum (a_{v} \triangleright ((a_{u} \triangleright a_{u}) \sqcup a_{v}) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v}).$$

$$S_{3} = \sum a_{u+v} \otimes (b_{u+v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum a_{i} \otimes (b_{i} \sqcup x_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v})$$

$$+ \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum (a_{i} \triangleright a_{j}) \otimes (b_{i} \sqcup b_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v}).$$

$$S_{4} = \sum (a_{u+v} \triangleright a_{\mathfrak{u}\sqcup\mathfrak{v}}) \otimes (b_{u+v} \sqcup b_{\mathfrak{u}\sqcup\mathfrak{v}}) + \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum (a_{i} \triangleright a_{\mathfrak{u}\sqcup\mathfrak{v}}) \otimes (b_{i} \sqcup x_{j} \sqcup b_{\mathfrak{u}\sqcup\mathfrak{v}})$$

$$+ \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum (a_{i} \triangleright (a_{j} \sqcup a_{\mathfrak{u}\sqcup\mathfrak{v}})) \otimes (b_{i} \sqcup b_{j} \sqcup b_{\mathfrak{u}\sqcup\mathfrak{v}}),$$
and

and

$$\begin{split} S_{\mathfrak{u}} &= \sum (a_u \sqcup (a_v \triangleright a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) - \sum (a_u \triangleright (a_v \triangleright a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) \\ &- \sum (a_v \triangleright (a_u \sqcup a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}), \\ S_{\mathfrak{v}} &= \sum ((a_u \triangleright a_{\mathfrak{u}}) \sqcup a_v) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) - \sum (a_u \triangleright (a_{\mathfrak{u}} \sqcup a_v)) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) \\ &- \sum (a_v \triangleright (a_u \triangleright a_{\mathfrak{u}})) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}). \end{split}$$

We claim that

(1)
$$S_0 = 0.$$

(2) $S_1 - S_3 = 0.$
(3) $S_2 + S_{\mathfrak{u}} + S_{\mathfrak{v}} - S_4 = 0.$

If these claims are true, the proposition is also true.

We prove three claims below. The claim (1), $S_0 = 0$, is immediate from (6.4). We show the claim (2). We will show that

$$S_1 = S_3 = \sum (a_u \diamond a_v) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup \mathfrak{v}).$$

With Lemma 5.7, we have

$$S_{1} = \sum (a_{u} \sqcup a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) - \sum (a_{u} \triangleright a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) - \sum (a_{v} \triangleright a_{u}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) = \sum (a_{u} \sqcup a_{v} - a_{u} \triangleright a_{v} - a_{v} \triangleright a_{u}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) = \sum (a_{u} \diamond a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}).$$

Now we calculate S_3 . Recall $x_u \diamond x_v = x_{u+v} + \sum_{i+j=u+v} \Delta^j_{u,v} x_i x_j$. Lemma 3.4 gives that

$$\Delta(x_u \diamond x_v) = 1 \otimes (x_u \diamond x_v) + \sum (a_u \diamond a_v) \otimes (b_u \sqcup b_v),$$

 \mathbf{SO}

$$1 \otimes x_{u+v} + \sum_{u+v} a_{u+v} \otimes b_{u+v} + \sum_{i+j=u+v} \Delta^j_{u,v} \left(1 \otimes x_i x_j + \sum_{u+v} a_i \otimes (b_i \sqcup u_j) + \sum_{u+v} (a_i \triangleright a_j) \otimes (b_i \sqcup u_j) \right)$$
$$= 1 \otimes (x_u \diamond x_v) + \sum_{u+v} (a_u \diamond a_v) \otimes (b_u \sqcup u_v),$$

therefore

(6.13)

$$\sum_{u+v} a_{u+v} \otimes b_{u+v} + \sum_{i+j=u+v} \Delta_{u,v}^j \sum_{u+v} a_i \otimes (b_i \sqcup x_j) + \sum_{i+j=u+v} \Delta_{u,v}^j \sum_{u+v} (a_i \triangleright a_j) \otimes (b_i \sqcup b_j)$$

$$= \sum (a_u \diamond a_v) \otimes (b_u \sqcup b_v).$$

We conclude that

$$S_{3} = \sum a_{u+v} \otimes (b_{u+v} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum a_{i} \otimes (b_{i} \sqcup x_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) + \sum_{i+j=u+v} \Delta_{u,v}^{j} \sum (a_{i} \triangleright a_{j}) \otimes (b_{i} \sqcup b_{j} \sqcup \mathfrak{u} \sqcup \mathfrak{v}) = \sum (a_{u} \diamond a_{v}) \otimes (b_{u} \sqcup b_{v} \sqcup \mathfrak{u} \sqcup \mathfrak{v})$$

and thus $S_1 = S_3$ as desired.

We now show the claim (3). Precisely, we show that

$$S_2 + S_{\mathfrak{u}} + S_{\mathfrak{v}} = S_4 = \sum ((a_u \diamond a_v) \triangleright a_{\mathfrak{u} \sqcup \mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}).$$

We have

$$S_{2} = \sum ((a_{u} \triangleright a_{u}) \sqcup (a_{v} \triangleright a_{v})) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$- \sum (a_{u} \triangleright (a_{u} \sqcup (a_{v} \triangleright a_{v}))) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$- \sum (a_{v} \triangleright ((a_{u} \triangleright a_{u}) \sqcup a_{v}) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$= \sum ((a_{u} \triangleright a_{u}) \sqcup (a_{v} \triangleright a_{v}) - a_{u} \triangleright (a_{u} \sqcup (a_{v} \triangleright a_{v})) - a_{v} \triangleright ((a_{u} \triangleright a_{u}) \sqcup a_{v}))$$

$$\otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v})$$

$$= \sum ((a_{u} \triangleright a_{u}) \diamond (a_{v} \triangleright a_{v})) \otimes (b_{u} \sqcup b_{u} \sqcup b_{v} \sqcup b_{v}).$$

Here the last equality is from Lemma 5.7 and Proposition 5.10 as

$$\begin{aligned} (a_{u} \triangleright a_{\mathfrak{u}}) & \sqcup (a_{v} \triangleright a_{\mathfrak{v}}) \\ &= (a_{u} \triangleright a_{\mathfrak{u}}) \triangleright (a_{v} \triangleright a_{\mathfrak{v}}) + (a_{v} \triangleright a_{\mathfrak{v}}) \triangleright (a_{u} \triangleright a_{\mathfrak{u}}) + (a_{u} \triangleright a_{\mathfrak{u}}) \diamond (a_{v} \triangleright a_{\mathfrak{v}}) \\ &= a_{u} \triangleright (a_{\mathfrak{u}} \sqcup (a_{v} \triangleright a_{\mathfrak{v}})) + a_{v} \triangleright ((a_{u} \triangleright a_{\mathfrak{u}}) \sqcup a_{\mathfrak{v}})) + (a_{u} \triangleright a_{\mathfrak{u}}) \diamond (a_{v} \triangleright a_{\mathfrak{v}}). \end{aligned}$$

By Lemma 5.7 and Proposition 5.10 again,

$$S_{\mathfrak{u}} = \sum (a_u \sqcup (a_v \triangleright a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) - \sum (a_u \triangleright (a_v \triangleright a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) - \sum (a_v \triangleright (a_u \sqcup a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) = \sum (a_u \sqcup (a_v \triangleright a_{\mathfrak{v}}) - a_u \triangleright (a_v \triangleright a_{\mathfrak{v}}) - a_v \triangleright (a_u \sqcup a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}) = \sum (a_u \diamond (a_v \triangleright a_{\mathfrak{v}})) \otimes (b_u \sqcup b_v \sqcup \mathfrak{u} \sqcup b_{\mathfrak{v}}),$$

and

$$S_{\mathfrak{v}} = \sum ((a_u \triangleright a_{\mathfrak{u}}) \sqcup a_v) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) - \sum (a_u \triangleright (a_u \sqcup a_v)) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) - \sum (a_v \triangleright (a_u \triangleright a_{\mathfrak{u}})) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}) = \sum ((a_u \triangleright a_{\mathfrak{u}}) \diamond a_v) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u}} \sqcup \mathfrak{v}).$$

With the equations for S_2 , $S_{\mathfrak{u}}$, $S_{\mathfrak{v}}$ and (6.9), again we can apply Lemma 5.7 and Proposition 5.10 to have

$$S_2 + S_{\mathfrak{u}} + S_{\mathfrak{v}} = \sum ((a_u \diamond a_v) \triangleright a_{\mathfrak{u} \sqcup \mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{u} \sqcup \mathfrak{v}}).$$

We now consider the term S_4 . We have

$$\begin{split} S_4 &= \sum (a_{u+v} \triangleright a_{u \sqcup v}) \otimes (b_{u+v} \sqcup b_{u \sqcup v}) + \sum_{i+j=u+v} \Delta^j_{u,v} \sum (a_i \triangleright a_{u \sqcup v}) \otimes (b_i \sqcup x_j \sqcup b_{u \sqcup v}) \\ &+ \sum_{i+j=u+v} \Delta^j_{u,v} \sum (a_i \triangleright (a_j \sqcup a_{u \sqcup v})) \otimes (b_i \sqcup b_j \sqcup b_u \sqcup v) \\ &= \sum (a_{u+v} \triangleright a_{u \sqcup v}) \otimes (b_{u+v} \sqcup b_{u \sqcup v}) + \sum_{i+j=u+v} \Delta^j_{u,v} \sum (a_i \triangleright a_{u \sqcup v}) \otimes (b_i \sqcup x_j \sqcup b_{u \sqcup v}) \\ &+ \sum_{i+j=u+v} \Delta^j_{u,v} \sum ((a_i \triangleright a_j) \triangleright a_{u \sqcup v}) \otimes (b_i \sqcup b_j \sqcup b_u \sqcup v) \\ &= \sum ((a_u \diamond a_v) \triangleright a_{u \sqcup v}) \otimes (b_u \sqcup b_v \sqcup b_u \sqcup v). \end{split}$$

Here the second equality follows from Proposition 5.10. The third one is a direct consequence of (6.13). This proves $S_2 + S_{\mathfrak{u}} + S_{\mathfrak{v}} = S_4$ as claimed.

Now Proposition 6.7 is proven with the claims (1), (2), and (3), as

$$\Delta(x_u \mathfrak{u} \sqcup x_v \mathfrak{v}) - \Delta(x_u \mathfrak{u}) \sqcup \Delta(x_v \mathfrak{v}) = S_0 - S_1 - S_2 - S_\mathfrak{u} - S_\mathfrak{v} + S_3 + S_4$$

= $S_0 - (S_1 - S_3) - (S_2 + S_\mathfrak{u} + S_\mathfrak{v} - S_4)$
= $0.$

Proof of Theorem 6.5. By combining Lemma 6.6 and Proposition 6.7 we conclude that, if Theorem 6.5 holds for all words $(\mathfrak{a}, \mathfrak{b}) \in \langle \Gamma \rangle \times \langle \Gamma \rangle$ with depth (\mathfrak{a}) +depth $(\mathfrak{b}) < n$ for some $n \in \mathbb{N}$ then the theorem holds for all words $(x_u\mathfrak{u}, x_v\mathfrak{v})$ such that depth $(x_u\mathfrak{u})$ +depth $(x_v\mathfrak{v}) = n$, and thus for $(x_{u,\alpha}\mathfrak{u}, x_{v,\beta}\mathfrak{v})$ such that depth $(x_{u,\alpha}\mathfrak{u})$ + depth $(x_{v,\beta}\mathfrak{v}) = n$, for all $u, v \in \mathbb{N}$ and $\alpha, \beta \in \mathbb{F}_q^*$. Since the theorem holds for the initial cases $(1,\mathfrak{b})$ and $(\mathfrak{a}, 1)$ for all $\mathfrak{a}, \mathfrak{b} \in \langle \Gamma \rangle$ with the arbitrary depths, the theorem holds for all $\mathfrak{u}, \mathfrak{v} \in \langle \Gamma \rangle$.

6.3. Coassociativity.

Next we want to prove:

Theorem 6.8. Let $\mathfrak{u} \in \langle \Gamma \rangle$. Then we have

$$(\mathrm{Id}\otimes\Delta)\Delta(\mathfrak{u}) = (\Delta\otimes\mathrm{Id})\Delta(\mathfrak{u}).$$

Proof. The proof is formed into two parts as follows:

- Part (1): if the coassociativity holds for $\mathfrak{u} = x_u \mathfrak{v}$, then coassociativity holds for $\mathfrak{u} = x_{u,\varepsilon}\mathfrak{v}$, for $\mathfrak{v} \in \langle \Gamma \rangle$ with depth(\mathfrak{v}) ≥ 0 , and
- Part (2): if the coassociativity holds for all \mathfrak{u} with depth(\mathfrak{u}) $\leq n$, then the coassociativity holds for $\mathfrak{u} = x_u \mathfrak{v}$ for $u \in \mathbb{N}$ and depth $\mathfrak{v} = n$.

Note that the initial case (i.e., when $\mathfrak{u} = x_{u,\varepsilon}$) is established by the coassociativity in \mathfrak{C} and Part (1).

For Part (1), we assume that $(\mathrm{Id} \otimes \Delta) \Delta(x_u \mathfrak{v}) = (\Delta \otimes \mathrm{Id}) \Delta(x_u \mathfrak{v})$. Let

$$\Delta(x_u) = 1 \otimes x_u + \sum a_u \otimes b_u, \qquad \Delta(\mathfrak{v}) = \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}}.$$

Here $a_u \in \Sigma$, $b_u \in \langle \Sigma \rangle$, and $a_v, b_v \in \langle \Gamma \rangle$. Thus

$$\Delta(x_u \mathfrak{v}) = 1 \otimes x_u \mathfrak{v} + \sum a_u a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}),$$

and

$$(\mathrm{Id} \otimes \Delta) \Delta(x_u \mathfrak{v}) = 1 \otimes 1 \otimes x_u \mathfrak{v} + \sum 1 \otimes a_u a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}) + \sum a_u a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_{\mathfrak{v}}),$$
$$(\Delta \otimes \mathrm{Id}) \Delta(x_u \mathfrak{v}) = 1 \otimes 1 \otimes x_u \mathfrak{v} + \sum \Delta(a_u a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}}).$$

Then the induction hypothesis that is the coassociativity for $x_u \mathfrak{v}$ is equivalent to (6.14)

$$\sum 1 \otimes a_u a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}) + \sum a_u a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_{\mathfrak{v}}) = \sum \Delta(a_u a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}}).$$

Now let $\mathfrak{u} = x_{u,\varepsilon}\mathfrak{v}$. We have

$$\Delta(x_{u,\varepsilon}) = 1 \otimes x_{u,\varepsilon} + \sum a_{u,\varepsilon} \otimes b_u$$

where $a_{u,\varepsilon} = \varphi_{\varepsilon}(a_u)$. By definition, we have

$$\Delta(\mathfrak{u}) = 1 \otimes \mathfrak{u} + \sum a_{u,\varepsilon} a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}).$$

One can calculate

$$(\mathrm{Id} \otimes \Delta)\Delta(\mathfrak{u}) = 1 \otimes \Delta(\mathfrak{u}) + \sum a_{u,\varepsilon}a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_{\mathfrak{v}})$$

$$= 1 \otimes 1 \otimes \mathfrak{u} + \sum 1 \otimes a_{u,\varepsilon}a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}) + \sum a_{u,\varepsilon}a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_{\mathfrak{v}}),$$

$$(\Delta \otimes \mathrm{Id})\Delta(\mathfrak{u}) = 1 \otimes 1 \otimes \mathfrak{u} + \sum \Delta(a_{u,\varepsilon}a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}})$$

$$= 1 \otimes 1 \otimes \mathfrak{u} + \sum (\varphi_{\varepsilon} \otimes \mathrm{Id})\Delta(a_u a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}})$$

$$+ \sum 1 \otimes a_{u,\varepsilon}a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}) - \sum 1 \otimes a_u a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}}).$$

Thus the coassociativity holds for $x_{u,\varepsilon}\mathfrak{v}$ if

$$\sum (\varphi_{\varepsilon} \otimes \mathrm{Id}) \Delta(a_u a_{\mathfrak{v}}) \otimes (b_u \sqcup b_{\mathfrak{v}}) = \sum a_{u,\varepsilon} a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_{\mathfrak{v}}) + \sum 1 \otimes a_u a_{\mathfrak{v}} \otimes (b_u \sqcup b_{\mathfrak{v}})$$

which is obtained by applying $\varphi_{\varepsilon} \otimes \operatorname{Id} \otimes \operatorname{Id}$ on Equation (6.14). This proves that the coassociativity for $x_u \mathfrak{v}$ implies the coassociativity for $x_{u,\varepsilon}\mathfrak{v}$.

We now prove Part (2). Let $n \ge 1$. We assume that the coassociativity holds for all words with depth $\le n$. Let $\mathfrak{u} = x_{t,\varepsilon}\mathfrak{t}$ with depth $\le n$. We write

$$\Delta(x_{t,\varepsilon}) = 1 \otimes x_{t,\varepsilon} + \sum_{t,\varepsilon} a_{t,\varepsilon} \otimes b_t, \text{ and } \Delta(\mathfrak{t}) = \sum_{t} a_{\mathfrak{t}} \otimes b_{\mathfrak{t}}.$$

where $a_{t,\varepsilon} \in \Gamma$ and $b_{t,\varepsilon}, a_t, b_t \in \langle \Gamma \rangle$. Then

$$\Delta(x_{t,\varepsilon}\mathfrak{t}) = 1 \otimes x_{t,\varepsilon} + \sum_{t,\varepsilon} a_{t,\varepsilon} a_{t} \otimes (b_t \sqcup b_t),$$

and the coassociativity for $x_{t,\varepsilon} \mathfrak{t}$ is reduced into

(6.15)
$$\sum \left(\Delta(a_{t,\varepsilon}a_{\mathfrak{t}}) - 1 \otimes a_{t,\varepsilon}a_{\mathfrak{t}} \right) \otimes (b_t \sqcup b_{\mathfrak{t}}) = \sum a_{t,\varepsilon}a_{\mathfrak{t}} \otimes (\Delta(b_t) \sqcup \Delta(b_{\mathfrak{t}})).$$
As a special case, let $\mathfrak{t} = 1$. Then $\Delta(\mathfrak{t}) = 1 \otimes 1$, and

$$\sum \left(\Delta(a_{t,\varepsilon}) - 1 \otimes a_{t,\varepsilon} \right) \otimes b_t = \sum a_{t,\varepsilon} \otimes \Delta(b_t).$$

Now assume $\mathfrak{u} = x_u x_{v,\beta} \mathfrak{v}$ with depth n + 1. Say

$$\begin{split} \Delta(x_u) &= 1 \otimes x_u + \sum a_u \otimes b_u, \\ \Delta(x_v) &= 1 \otimes x_v + \sum a_v \otimes b_v, \quad \text{so } \Delta(x_{v,\beta}) = 1 \otimes x_{v,\beta} + \sum a_{u,\beta} \otimes b_v, \\ \Delta(\mathfrak{v}) &= \sum a_{\mathfrak{v}} \otimes b_{\mathfrak{v}}, \end{split}$$

where $a_u, a_v \in \Sigma, b_u, b_v \in \langle \Sigma \rangle, a_{\mathfrak{v}}, b_{\mathfrak{v}} \in \langle \Gamma \rangle$, and $a_{u,\beta} = \varphi_{\beta}(a_u)$. Then,

$$\Delta(x_{v,\beta}\mathfrak{v}) = 1 \otimes x_{v,\beta}\mathfrak{v} + \sum a_{v,\beta}a_{\mathfrak{v}} \otimes (b_v \sqcup b_{\mathfrak{v}}).$$

It follows that

$$\Delta(x_u x_{v,\beta} \mathfrak{v}) = 1 \otimes x_u x_{v,\beta} \mathfrak{v} + \sum a_u \otimes (b_u \sqcup u x_{v,\beta} \mathfrak{v}) + \sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}),$$

thus

$$(\mathrm{Id} \otimes \Delta) \Delta(x_u x_{v,\beta} \mathfrak{v})$$

$$=1 \otimes 1 \otimes x_u x_{v,\beta} \mathfrak{v} + \sum 1 \otimes a_u \otimes (b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum 1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}})$$

$$+ \sum a_u \otimes \Delta(b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_v \sqcup b_{\mathfrak{v}}),$$

$$(\Delta \otimes \mathrm{Id}) \Delta(x_u x_{v,\beta} \mathfrak{v})$$

$$=1 \otimes 1 \otimes x_u x_{v,\beta} \mathfrak{v} + \sum \Delta(a_u) \otimes (b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum \Delta(a_u a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}),$$
so what we want to show is that

(6.16)
$$\sum 1 \otimes a_u \otimes (b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum 1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) + \sum a_u \otimes \Delta (b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes \Delta (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}), = \sum \Delta (a_u) \otimes (b_u \sqcup x_{v,\beta} \mathfrak{v}) + \sum \Delta (a_u a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}).$$

Note that, by Equation (6.15) we have

$$\begin{split} &\sum 1 \otimes a_u \otimes (b_u \amalg x_{v,\beta} \mathfrak{v}) + \sum a_u \otimes \Delta (b_u \amalg x_{v,\beta} \mathfrak{v}) - \sum \Delta (a_u) \otimes (b_u \amalg x_{v,\beta} \mathfrak{v}) \\ &= \sum a_u \otimes \Delta (b_u \amalg x_{v,\beta} \mathfrak{v}) - \left(\sum (\Delta (a_u) - 1 \otimes a_u) \otimes b_u \right) \amalg (1 \otimes 1 \otimes x_{v,\beta} \mathfrak{v}) \\ &= \sum a_u \otimes \Delta (b_u \amalg x_{v,\beta} \mathfrak{v}) - \left(\sum a_u \otimes \Delta (b_u) \right) \amalg (1 \otimes 1 \otimes x_{v,\beta} \mathfrak{v}) \\ &= \sum a_u \otimes (\Delta (b_u) \amalg \Delta (x_{v,\beta} \mathfrak{v})) - \sum a_u \otimes (\Delta (b_u) \amalg (1 \otimes x_{v,\beta} \mathfrak{v})) \\ &= \sum a_u \otimes (\Delta (b_u) \amalg (\Delta (x_{v,\beta} \mathfrak{v}) - 1 \otimes x_{v,\beta} \mathfrak{v})) \\ &= \sum a_u \otimes (\Delta (b_u) \amalg (a_{v,\beta} a_u \otimes (b_v \amalg b_v))) \end{split}$$

so Equation (6.16) is equivalent to

(6.17)
$$\sum \Delta(a_u a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) = \sum 1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) + \sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes \Delta(b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) + \sum a_u \otimes (\Delta(b_u) \sqcup (a_{v,\beta} a_{\mathfrak{v}} \otimes (b_v \sqcup b_{\mathfrak{v}}))).$$

From now on, tracing the indices is important. When needed, we use the notation \sum_u which stands for the sum over the pairs (a_u, b_u) , and similarly $\sum_{(v,\beta)}, \sum_{v}$, etc. We now fix a_u, b_u . The last two terms of (6.17) become

$$\begin{split} &\sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (\Delta(b_u) \sqcup \Delta(b_v) \sqcup \Delta(b_{\mathfrak{v}})) + \sum a_u \otimes (\Delta(b_u) \sqcup (a_{v,\beta} a_{\mathfrak{v}} \otimes (b_v \sqcup b_{\mathfrak{v}}))) \\ &= (1 \otimes \Delta(b_u)) \sqcup \left(\sum a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (\Delta(b_v) \sqcup \Delta(b_{\mathfrak{v}})) + \sum a_u \otimes a_{v,\beta} a_{\mathfrak{v}} \otimes (b_v \sqcup b_{\mathfrak{v}}) \right) \\ &= (a_u \otimes 1 \otimes 1) \cdot \left((1 \otimes \Delta(b_u)) \sqcup \left(\sum a_{v,\beta} a_{\mathfrak{v}} \otimes (\Delta(b_v) \sqcup \Delta(b_{\mathfrak{v}})) + \sum 1 \otimes a_{v,\beta} a_{\mathfrak{v}} \otimes (b_v \sqcup b_{\mathfrak{v}}) \right) \right) \end{split}$$

$$= (a_u \otimes 1 \otimes 1) \cdot \left((1 \otimes \Delta(b_u)) \sqcup \left(\sum \Delta(a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_v \sqcup b_{\mathfrak{v}}) \right) \right)$$

Recall that \cdot is the concatenation. The last equality is from the induction hypothesis, Equation (6.15). Equation (6.17) becomes

$$(6.18)$$

$$\sum \Delta(a_u a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}})$$

$$= \sum 1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) + \sum (a_u \otimes 1 \otimes 1) \cdot \left((1 \otimes \Delta(b_u)) \sqcup \left(\sum \Delta(a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_v \sqcup b_{\mathfrak{v}}) \right) \right).$$

Next we fix $a_u, a_{v,\beta}, a_v$. So the corresponding b's are also fixed. Let

$$\Delta(a_u) = 1 \otimes a_u + \sum_{(1)} c_{(1)} \otimes d_{(1)}, \quad \Delta(a_{v,\beta}a_{v}) = \sum_{(2)} c_{(2)} \otimes d_{(2)}.$$

Then, we have

$$\Delta(a_u a_{v,\beta} a_{\mathfrak{v}}) = 1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} + \sum_{(1),(2)} c_{(1)} c_{(2)} \otimes (d_{(1)} \sqcup d_{(2)}).$$

The summand of $\sum \Delta(a_u a_{v,\beta} a_{\mathfrak{v}}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}})$ for fixed $a_u, a_{v,\beta}, a_{\mathfrak{v}}$ is

$$\left(1 \otimes a_u a_{v,\beta} a_{\mathfrak{v}} + \left(\sum_{(1),(2)} c_{(1)} c_{(2)} \otimes (d_{(1)} \sqcup d_{(2)})\right)\right) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}).$$

Plugging this and the sum representation of $\Delta(a_{v,\beta}a_{\mathfrak{v}})$ into the equation (6.18) yields

$$\sum_{u,(v,\beta),\mathfrak{v}} \left(\sum_{(1),(2)} c_{(1)}c_{(2)} \otimes (d_{(1)} \sqcup d_{(2)}) \right) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) \\ - \sum a_u c_{(2)} \otimes (\Delta(b_u) \sqcup (d_{(2)} \otimes b_v \sqcup b_{\mathfrak{v}})).$$

Finally, we fix a_u , $a_{v,\beta}$, $a_{\mathfrak{v}}$, $c_{(2)}$. The corresponding summands are

$$\sum_{(1)} c_{(1)}c_{(2)} \otimes (d_{(1)} \sqcup d_{(2)}) \otimes (b_u \sqcup b_v \sqcup b_{\mathfrak{v}}) - a_u c_{(2)} \otimes (\Delta(b_u) \sqcup (d_{(2)} \otimes b_v \sqcup b_{\mathfrak{v}})).$$

By direct calculations, we get

$$\sum_{(1)} c_{(1)}c_{(2)} \otimes (d_{(1)} \sqcup d_{(2)}) \otimes (b_u \sqcup b_v \sqcup b_v) - a_u c_{(2)} \otimes (\Delta(b_u) \sqcup (d_{(2)} \otimes b_v \sqcup b_v))$$
$$= \left(\left(\sum_{(1)} c_{(1)} \otimes d_{(1)} \otimes b_u - a_u \otimes \Delta(b_u) \right) \sqcup (1 \otimes d_{(2)} \otimes b_v \sqcup b_v) \right) \cdot (c_{(2)} \otimes 1 \otimes 1),$$

whose inner sum is reduced to

$$\sum_{(1)} c_{(1)} \otimes d_{(1)} \otimes b_u - a_u \otimes 1 \otimes \Delta(b_u) = (\Delta(a_u) - 1 \otimes a_u) \otimes b_u - a_u \otimes 1 \otimes \Delta(b_u).$$

Now we can apply Equation (6.15) for the sum of the summand over a_u to have

$$\sum_{u} \left(\Delta(a_u) \otimes b_u - 1 \otimes a_u \otimes b_u - a_u \otimes 1 \otimes \Delta(b_u) \right) = 0,$$

that is, Equation (6.18) is true. Thus the induction step is proved.

From the coassociativity of \mathfrak{C} , we have the coassociativity for $\mathfrak{u} \in \Sigma$, and then for $\mathfrak{u} \in \Gamma$ by Part (1). By applying Parts (2) and (1) repeatedly, we can cover arbitrary depth of $\mathfrak{u} \in \langle \Gamma \rangle$. Hence this completes the proof.

6.4. Hopf algebra structure on \mathfrak{D} .

Let the counit $\epsilon \colon \mathfrak{D} \to \mathbb{F}_q$ defined as $\epsilon(1) = 1$; $\epsilon(\mathfrak{u}) = 0$ otherwise.

We already checked that $(\mathfrak{D}, \sqcup, u, \Delta, \epsilon)$ is a bialgebra by Theorem 5.9, Theorem 6.5, and Theorem 6.8. By induction it is easily verified that \sqcup and Δ preserve the weight, and the weight zero subset is isomorphic to \mathbb{F}_q . Therefore, $(\mathfrak{D}, \sqcup, u, \Delta, \epsilon)$ is a connected graded bialgebra. By Theorem 2.6, we have the following theorem.

Theorem 6.9. $(\mathfrak{D}, \sqcup, u, \Delta, \epsilon)$ is a connected graded Hopf algebra of finite type over \mathbb{F}_q .

References

- G. Anderson. t-motives. Duke Math. J., 53(2):457–502, 1986.
- [2] G. Anderson, W. D. Brownawell, and M. Papanikolas. Determination of the algebraic relations among special Γ-values in positive characteristic. Ann. of Math. (2), 160(1):237–313, 2004.
- [3] G. Anderson and D. Thakur. Tensor powers of the Carlitz module and zeta values. Ann. of Math. (2), 132(1):159–191, 1990.
- [4] G. Anderson and D. Thakur. Multizeta values for $\mathbb{F}_q[t]$, their period interpretation, and relations between them. Int. Math. Res. Not. IMRN, (11):2038–2055, 2009.
- [5] T. Arakawa and M. Kaneko. Multiple zeta values, poly-Bernoulli numbers, and related zeta functions. Nagoya Math. J., 153:189–209, 1999.
- [6] T. Arakawa and M. Kaneko. On multiple L-values. J. Math. Soc. Japan, 56(4):967–991, 2004.
- [7] D. J. Broadhurst. On the enumeration of irreducible k-fold Euler sums and their roles in knot theory and field theory. available at https://arxiv.org/abs/hep-th/9604128, 1996
- [8] D. J. Broadhurst. Conjectured Enumeration of irreducible Multiple Zeta Values, from Knots and Feynman Diagrams. available at https://arxiv.org/abs/hep-th/9612012, 1996
- [9] F. Brown. Mixed Tate motives over Z. Ann. of Math. (2), 175:949–976, 2012.
- [10] D. Brownawell and M. Papanikolas. A rapid introduction to Drinfeld modules, t-modules and t-motives. In G. Böckle, D. Goss, U. Hartl, and M. Papanikolas, editors, t-motives: Hodge structures, transcendence and other motivic aspects", EMS Series of Congress Reports, pages 3–30. European Mathematical Society, 2020.
- [11] J. Burgos Gil and J. Fresan. Multiple zeta values: from numbers to motives. to appear, Clay Mathematics Proceedings.
- [12] L. Carlitz. On certain functions connected with polynomials in Galois field. Duke Math. J., 1(2):137–168, 1935.
- [13] S. Charlton. On motivic multiple t values, Saha's basis conjecture, and generators of alternating MZV's. arXiv:2112.14613v1, 2021.
- [14] H.-J. Chen. On shuffle of double zeta values over $\mathbb{F}_q[t]$. J. Number Theory, 148:153–163, 2015.
- [15] P. Deligne. Le groupe fondamental unipotent motivique de $G_m \mu_N$, pour N = 2, 3, 4, 6 ou 8. Publ. Math. Inst. Hautes Études Sci., 112:101–141, 2010.
- [16] P. Deligne. Letter to Dinesh Thakur, 27 May 2017.
- [17] P. Deligne and A. Goncharov. Groupes fondamentaux motiviques de Tate mixte. Ann. Sci. École Norm. Sup. (4), 38(1):1–56, 2005.
- [18] V. Drinfeld. On quasitriangular quasi-Hopf algebras and on a group that is closely connected with Gal(Q/Q). Algebra i Analiz, 2(4):149–181, 1990.
- [19] R. Ehrenborg. On posets and Hopf algebras. Adv. Math., 119(1):1–25, 1996.
- [20] L. Euler. Meditationes circa singulare serierum genus. Novi commentarii academiae scientiarum Petropolitanae, 140–186, 1776.
- [21] O. Gezmis and F. Pellarin. Trivial multiple zeta values in Tate algebras. International Mathematics Research Notices, to appear, rnab104, 2021.
- [22] A. Goncharov. Multiple polylogarithms and mixed tate motives. arXiv:math/0103059v4, 2001.

- [23] A. Goncharov. Galois symmetries of fundamental groupoids and noncommutative geometry. Duke Math. J., 128(2):209–284, 2005.
- [24] D. Goss. Basic Structures of function field arithmetic, volume 35 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3). Springer-Verlag, Berlin, 1996.
- [25] R. Harada. Alternating multizeta values in positive characteristic. Math. Z., 298(3-4):1263– 1291, 2021.
- [26] U. Hartl and A. K. Juschka. Pink's theory of Hodge structures and the Hodge conjectures over function fields. In G. Böckle, D. Goss, U. Hartl, and M. Papanikolas, editors, t-motives: Hodge structures, transcendence and other motivic aspects", EMS Series of Congress Reports, pages 31–182. European Mathematical Society, 2020.
- [27] M. Hoffman. The algebra of multiple harmonic series. J. Algebra, 194:477–495, 1997.
- [28] M. Hoffman. Quasi-shuffle products. J. Algebraic Combin., 11(1):49–68, 2000.
- [29] M. Hoffman. An odd variant of multiple zeta values. Commun. Number Theory Phys., 13(3):529–567, 2019.
- [30] K. Ihara, M. Kaneko, and D. Zagier. Derivation and double shuffle relations for multiple zeta values. Compos. Math., 142(2):307–338, 2006.
- [31] B.-H. Im, H. Kim, K. N. Le, T. Ngo Dac, and L. H. Pham. Zagier-Hoffman's conjectures in positive characteristic. available at https://hal.archives-ouvertes.fr/hal-03667755, 2022.
- [32] B.-H. Im, H. Kim, K. N. Le, T. Ngo Dac, and L. H. Pham. Hopf algebras and multiple zeta values in positive characteristic. available at https://hal.archives-ouvertes.fr/hal-03937702, 2023.
- [33] K. Iwasawa. Analogies between number fields and function fields. In Some Recent Advances in the Basic Sciences, Vol. 2 (Proc. Annual Sci. Conf., Belfer Grad. School Sci., Yeshiva Univ., New York, 1965-1966), pages 203–208. Yeshiva Univ., Belfer Graduate School of Science, New York, 1969.
- [34] M. Kaneko, K. Tasaka. Double zeta values, double Eisenstein series, and modular forms of level. Math. Ann., 357(3): 1091--1118, 2013.
- [35] M. Kontsevich. Operads and motives in deformation quantization. volume 48, pages 35–72. 1999. Moshé Flato (1937–1998).
- [36] J. A. Lara Rodriguez and D. Thakur. Zeta-like multizeta values for $\mathbb{F}_q[t]$. Indian J. Pure Appl. Math., 45(5):787–801, 2014.
- [37] J. A. Lara Rodriguez and D. Thakur. Zeta-like multizeta values for higher genus curves. J. Théor. Nombres Bordeaux, 33(2):553–581, 2021.
- [38] T. T. Q. Le and J. Murakami. Kontsevich's integral for the Kauffman polynomial. Nagoya Math. J., 142:39–65, 1996.
- [39] B. Mazur and A. Wiles. Analogies between function fields and number fields. Amer. J. Math., 105(2):507–521, 1983.
- [40] T. Ngo Dac. On Zagier-Hoffman's conjectures in positive characteristic. Ann. of Math. (2), 194(1):361–392, 2021.
- [41] F. Pellarin. Values of certain L-series in positive characteristic. Ann. of Math. (2), 176(3):2055–2093, 2012.
- [42] G. Racinet. Doubles mélanges des polylogarithmes multiples aux racines de l'unité. Publ. Math. Inst. Hautes Études Sci., (95):185–231, 2002.
- [43] S. Shi. Multiple zeta values over F_q[t]. PhD thesis, University of Rochester (available at http://hdl.handle.net/1802/33859), November 2018.
- [44] T. Terasoma. Mixed Tate motives and multiple zeta values. Invent. Math., 149(2):339–369, 2002.
- [45] D. Thakur. Function field arithmetic. World Scientific Publishing Co., Inc., River Edge, NJ, 2004.
- [46] D. Thakur. Power sums with applications to multizeta and zeta zero distribution for $\mathbb{F}_q[t]$. Finite Fields Appl., 15(4):534–552, 2009.
- [47] D. Thakur. Relations between multizeta values for $\mathbb{F}_q[t]$. Int. Math. Res. Not., (12):2318–2346, 2009.
- [48] D. Thakur. Shuffle relations for function field multizeta values. Int. Math. Res. Not. IMRN, (11):1973–1980, 2010.
- [49] D. Thakur. Multizeta values for function fields: a survey. J. Théor. Nombres Bordeaux, 29(3):997–1023, 2017.

- [50] D. Thakur. Multizeta in function field arithmetic. In G. Böckle, D. Goss, U. Hartl, and M. Papanikolas, editors, t-motives: Hodge structures, transcendence and other motivic aspects", EMS Series of Congress Reports, pages 441–452. European Mathematical Society, 2020.
- [51] G. Todd. A conjectural characterization for $\mathbb{F}_q(t)$ -linear relations between multizeta values. J. Number Theory, 187:264–28, 2018.
- [52] A. Weil. Sur l'analogie entre les corps de nombres algébrique et les corps de fonctions algébrique. Revue Scient., 77:104–106, 1939.
- [53] J. Yu. Transcendence and special zeta values in characteristic p. Ann. of Math. (2), 134(1):1– 23, 1991.
- [54] D. Zagier. Values of zeta functions and their applications. In First European Congress of Mathematics, Vol. II Paris, 1992), volume 120 of Progr. Math., pages 497–512. Birkhäuser, Basel, 1994.
- [55] J. Zhao. Multiple zeta functions, multiple polylogarithms and their special values, volume 12 of Series on Number Theory and its Applications. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2016.

Dept. of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, South Korea

Email address: bhim@kaist.ac.kr

Dept. of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, South Korea

Email address: hojinkim@kaist.ac.kr

NORMANDIE UNIVERSITÉ, UNIVERSITÉ DE CAEN NORMANDIE - CNRS, LABORATOIRE DE MATHÉMATIQUES NICOLAS ORESME (LMNO), UMR 6139, 14000 CAEN, FRANCE.

Email address: khac-nhuan.le@unicaen.fr

NORMANDIE UNIVERSITÉ, UNIVERSITÉ DE CAEN NORMANDIE - CNRS, LABORATOIRE DE MATHÉMATIQUES NICOLAS ORESME (LMNO), UMR 6139, 14000 CAEN, FRANCE.

Email address: tuan.ngodac@unicaen.fr

Institute of Mathematics, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, 10307 Hanoi, Viet Nam

Email address: plhuong@math.ac.vn