



HAL
open science

LARGE TIME CONVERGENCE FOR A CHEMOTAXIS MODEL WITH DEGENERATE LOCAL SENSING AND CONSUMPTION

Philippe Laurençot

► **To cite this version:**

Philippe Laurençot. LARGE TIME CONVERGENCE FOR A CHEMOTAXIS MODEL WITH DEGENERATE LOCAL SENSING AND CONSUMPTION. Bulletin of the Korean Mathematical Society, 2024, 61 (2), pp.479–488. 10.4134/BKMS.b230173 . hal-04053626

HAL Id: hal-04053626

<https://hal.science/hal-04053626>

Submitted on 31 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LARGE TIME CONVERGENCE FOR A CHEMOTAXIS MODEL WITH DEGENERATE LOCAL SENSING AND CONSUMPTION

PHILIPPE LAURENÇOT

ABSTRACT. Convergence to a steady state in the long term limit is established for global weak solutions to a chemotaxis model with degenerate local sensing and consumption, when the motility function is C^1 -smooth on $[0, \infty)$, vanishes at zero, and is positive on $(0, \infty)$. A condition excluding that the large time limit is spatially homogeneous is also provided. These results extend previous ones derived for motility functions vanishing algebraically at zero and rely on a completely different approach.

1. INTRODUCTION

The chemotaxis system with local sensing and consumption

$$\partial_t u = \Delta(u\gamma(v)) \quad \text{in } (0, \infty) \times \Omega, \quad (1.1a)$$

$$\partial_t v = \Delta v - uv \quad \text{in } (0, \infty) \times \Omega, \quad (1.1b)$$

$$\nabla(u\gamma(v)) \cdot \mathbf{n} = \nabla v \cdot \mathbf{n} = 0 \quad \text{on } (0, \infty) \times \partial\Omega, \quad (1.1c)$$

$$(u, v)(0) = (u^{in}, v^{in}) \quad \text{in } \Omega, \quad (1.1d)$$

describes the dynamics of a population of cells with density $u \geq 0$ living on a nutrient with concentration $v \geq 0$ and moving in space under the combined effects of a nutrient-dependent diffusion and a nutrient-induced chemotactic bias [12]. Here, Ω is a bounded domain of \mathbb{R}^n , $n \geq 1$, and the motility γ is a smooth function which is positive on $(0, \infty)$. Unlike the classical Keller-Segel system with local sensing

$$\partial_t u = \Delta(u\gamma(v)) \quad \text{in } (0, \infty) \times \Omega, \quad (1.2a)$$

$$\partial_t v = \Delta v - v + u \quad \text{in } (0, \infty) \times \Omega, \quad (1.2b)$$

$$\nabla(u\gamma(v)) \cdot \mathbf{n} = \nabla v \cdot \mathbf{n} = 0 \quad \text{on } (0, \infty) \times \partial\Omega, \quad (1.2c)$$

$$(u, v)(0) = (u^{in}, v^{in}) \quad \text{in } \Omega, \quad (1.2d)$$

Date: March 31, 2023.

2020 Mathematics Subject Classification. 35B40 - 35K65 - 35K51 - 35Q92.

Key words and phrases. convergence - Liapunov functional - chemotaxis-consumption model - local sensing.

in which the variable v is rather the concentration of a signalling chemical produced by the cells as accounted for in (1.2b) [11], the nutrient is consumed by the cells in the model (1.1) according to the nonlinear absorption term $-uv$ in (1.1b). The dynamics of the two models is thus expected to differ significantly. On the one hand, the long term behaviour of solutions to (1.2) is far from being completely understood. Convergence of (u, v) to the spatially homogeneous steady state $(\|u^{in}\|_1/|\Omega|, \|u^{in}\|_1/|\Omega|)$ is established in [3, Theorem 1.5] when $\gamma'(s) \leq 0 \leq s\gamma'(s) + \gamma(s)$ for $s \geq 0$ and a similar result is likely to be true when $\gamma' \geq 0$. Such a simple dynamics is however unlikely to be the generic one, as there exist non-constant stationary solutions to (1.2) in some domains, see [18, 20]. Besides, though solutions to (1.2) are global and even bounded for a large class of motility functions γ [1–5, 7, 9, 17, 19, 25, 26], unbounded solutions do exist [6, 8, 10].

The situation is somewhat simpler for (1.1) as the set of stationary solutions can easily be identified and seen to depend heavily on the value of $\gamma(0)$ [16, 24]. Indeed, if $\gamma(0) > 0$, then the only steady states to (1.1) are the spatially homogeneous solutions $(M, 0)$, $M \geq 0$, and they attract the dynamics [13, 16]. In contrast, the set of stationary solutions to (1.1) is much larger when $\gamma(0) = 0$, as $(\bar{u}, 0)$ is a stationary solution to (1.1) for any (sufficiently smooth) function \bar{u} . Despite this wealth of stationary solutions, Winkler proves in [22, 24] that the dynamics of (1.1) selects one and only one steady state in the large time limit: more precisely, given

$$\gamma \in C^1([0, \infty)), \quad \gamma(0) = 0, \quad \gamma > 0 \quad \text{on} \quad (0, \infty), \quad (1.3)$$

satisfying

$$\gamma \in C^3((0, \infty)) \quad \text{and} \quad \liminf_{s \rightarrow 0} \left\{ \frac{\gamma(s)}{s^\alpha} \right\} > 0 \quad (1.4)$$

for some $\alpha \geq 1$ and a suitably constructed global weak solution (u, v) to (1.1) [21, 23], there is a non-negative measurable function u_∞ such that

$$(u(t), v(t)) \quad \text{converges to} \quad (u_\infty, 0) \quad \text{in a suitable topology}$$

if one of the following additional assumptions holds true:

- (I): $n \in \{1, 2\}$ and $\gamma'(0) > 0$, see [22, Theorem 1.4],
- (II): $n \geq 3$ and there is $\alpha' \in (1, 2]$ such that

$$\limsup_{s \rightarrow 0} \left\{ s^{2-\alpha'} |\gamma''(s)| \right\} < \infty, \quad (1.5)$$

see [24, Theorem 1.2].

Observe that, for $\alpha \geq 1$, the function $\gamma(s) = s^\alpha$, $s \geq 0$, satisfies (1.3), (1.4), and (1.5) with $\alpha' = \min\{\alpha, 2\}$.

Besides, conditions on γ , u^{in} , and v^{in} are provided to guarantee that u_∞ is not a constant, see [22, Theorem 1.5] and [24, Corollary 1.4].

The aim of this note is to prove that these results are actually valid under the sole assumption (1.3) on γ , without assuming an algebraic behaviour of γ near zero. However, the convergence established below takes place in a weaker topology for the u -component than the one obtained in [22, 24]. In

addition, the approach used herein is different and thus provides an alternative viewpoint on the stabilization issue for (1.1). We first state the convergence result.

Theorem 1.1. *Assume that γ satisfies (1.3) and consider $(u^{in}, v^{in}) \in L_+^\infty(\Omega, \mathbb{R}^2)$, where E_+ denotes the positive cone of the Banach lattice E . If (u, v) is a global weak solution to (1.1) in the sense of Definition 2.1 below, then*

$$A_\infty(x) := \int_0^\infty (u\gamma(v))(s, x) \, ds \in H_+^1(\Omega), \quad x \in \Omega,$$

and

$$u(t) \rightharpoonup u_\infty := \Delta A_\infty + u^{in} \quad \text{in } H^1(\Omega)' \quad \text{as } t \rightarrow \infty, \quad (1.6)$$

$$\lim_{t \rightarrow \infty} \|v(t)\|_p = 0, \quad p \in [1, \infty). \quad (1.7)$$

Moreover, $\langle u_\infty, \vartheta \rangle_{(H^1)', H^1} \geq 0$ for all $\vartheta \in H_+^1(\Omega)$ and $\langle u_\infty, 1 \rangle_{(H^1)', H^1} = \langle u^{in}, 1 \rangle_{(H^1)', H^1}$.

The cornerstone of our approach to the study of the long term behaviour of global weak solutions to (1.1) is the observation that the dynamics of the system (1.1) is somewhat encoded in that of the auxiliary function

$$A(t, x) := \int_0^t (u\gamma(v))(s, x) \, ds, \quad (t, x) \in (0, \infty) \times \Omega,$$

which bears several interesting properties collected in Lemma 2.3. Among others, $t \mapsto A(t, x)$ is non-decreasing for a.e. $x \in \Omega$ and the trajectory $\{A(t) : t \geq 0\}$ is bounded in $H^1(\Omega)$, two features which guarantee in particular that the function A_∞ introduced in Theorem 1.1 is well-defined and lies in $H^1(\Omega)$. Moreover, for all $t \geq 0$, the function $A(t)$ is a variational solution of the elliptic equation

$$-\Delta A(t) = u^{in} - u(t) \quad \text{in } \Omega, \quad \nabla A(t) \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega,$$

so that the large time behaviour of $u(t)$ is driven by that of $A(t)$.

The second contribution of this paper is in the spirit of [24, Theorem 1.3] and provides an estimate on the distance in $H^1(\Omega)'$ between the initial condition u^{in} and the final state u_∞ of the u -component of (1.1). Its statement requires additional notation, which we introduce now: for $z \in H^1(\Omega)'$, we set $\langle z \rangle := \langle z, 1 \rangle_{(H^1)', H^1} / |\Omega|$ and note that

$$\langle z \rangle = \frac{1}{|\Omega|} \int_\Omega z(x) \, dx \quad \text{for } z \in H^1(\Omega)' \cap L^1(\Omega).$$

Now, for $z \in H^1(\Omega)'$ with $\langle z \rangle = 0$, we define $\mathcal{K}[z] \in H^1(\Omega)$ as the unique (variational) solution to

$$-\Delta \mathcal{K}[z] = z \quad \text{in } \Omega, \quad \nabla \mathcal{K}[z] \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega, \quad (1.8a)$$

satisfying

$$\langle \mathcal{K}[z] \rangle = 0. \quad (1.8b)$$

We then choose the following norm $\|\cdot\|_{(H^1)'}$ on $H^1(\Omega)'$:

$$\|z\|_{(H^1)'} := \|\nabla \mathcal{K}[z - \langle z \rangle]\|_2 + |\langle z \rangle|, \quad z \in (H^1(\Omega))'.$$

Proposition 1.2. *Assume that γ satisfies (1.3) and consider $(u^{in}, v^{in}) \in L_+^\infty(\Omega, \mathbb{R}^2)$. If (u, v) is a global weak solution to (1.1) in the sense of Definition 2.1 below and u_∞ denotes the weak limit in $H^1(\Omega)'$ of $u(t)$ as $t \rightarrow \infty$ given by Theorem 1.1, then*

$$\|u_\infty - u^{in}\|_{(H^1)'}^2 \leq \|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, \|v^{in}\|_\infty)}. \quad (1.9)$$

In particular, if

$$\|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, \|v^{in}\|_\infty)} < \|u^{in} - \langle u^{in} \rangle\|_{(H^1)'}^2, \quad (1.10)$$

then u_∞ is not a constant.

An immediate consequence of Proposition 1.2 is that, given $u^{in} \in L_+^\infty(\Omega)$ with $u^{in} \not\equiv \langle u^{in} \rangle$ and a sufficient small $v^{in} \in L_+^\infty(\Omega)$, the first component of the corresponding global weak solution (u, v) to (1.1) has a non-constant limit. A quantitative estimate on the required smallness of v^{in} in $L^\infty(\Omega)$ is provided by (1.10) and reads

$$\|v^{in}\|_\infty \|\gamma'\|_{L^\infty(0, \|v^{in}\|_\infty)} < \frac{\|u^{in} - \langle u^{in} \rangle\|_{(H^1)'}^2}{|\Omega| \|u^{in}\|_\infty}.$$

Such a result is obviously connected with the fact that the solution (u, v) to (1.1) with initial condition $(u^{in}, 0)$ is the stationary solution $(u, v) = (u^{in}, 0)$, as already mentioned.

2. PROOFS

Let us first make precise the notion of global weak solution to (1.1) to be used in this paper. We emphasize here that, since $\gamma(0) = 0$, the equation (1.1a) is degenerate, so that we cannot expect much regularity on u .

Definition 2.1. *Assume that γ satisfies (1.3) and consider $(u^{in}, v^{in}) \in L_+^\infty(\Omega, \mathbb{R}^2)$. A global weak solution to (1.1) is a pair of non-negative functions (u, v) such that*

$$\begin{aligned} u &\in C_w([0, \infty), H^1(\Omega)') \cap L^\infty((0, \infty), L_+^1(\Omega)), \\ v &\in C([0, \infty), L_+^1(\Omega)) \cap L^\infty((0, \infty) \times \Omega) \cap L_{\text{loc}}^2([0, \infty), H^1(\Omega)), \\ u\sqrt{\gamma(v)} &\in L_{\text{loc}}^2([0, \infty), L^2(\Omega)), \end{aligned}$$

which satisfies

$$\begin{aligned} \langle u(t), \vartheta(t) \rangle_{(H^1)', H^1} - \int_\Omega u^{in} \vartheta(0) \, dx &= \int_0^t \int_\Omega u(s) \gamma(v(s)) \Delta \vartheta(s) \, dx ds \\ &\quad + \int_0^t \langle u(s), \partial_t \vartheta(s) \rangle_{(H^1)', H^1} \, ds \end{aligned}$$

for $\vartheta \in L^2((0, t), H_N^2(\Omega)) \cap W^{1,2}((0, t), H^1(\Omega))$ and $t \geq 0$, where

$$H_N^2(\Omega) := \{z \in H^2(\Omega) : \nabla z \cdot \mathbf{n} = 0 \text{ on } \partial\Omega\},$$

as well as

$$\begin{aligned} \int_{\Omega} (v(t)\vartheta(t) - v^{in}\vartheta(0)) \, dx + \int_0^t \int_{\Omega} \nabla v(s) \cdot \nabla \vartheta(s) \, dx ds + \int_0^t \int_{\Omega} (uv)(s)\vartheta(s) \, dx ds \\ = \int_0^t \int_{\Omega} v(s)\partial_t \vartheta(s) \, dx ds \end{aligned}$$

for $\vartheta \in L^2((0, t), H^1(\Omega)) \cap L^\infty((0, t) \times \Omega)$ and $t \geq 0$.

We shall not address the existence issue here and refer to [21–23] for results in that direction, the main assumption on γ being that it vanishes in an algebraic way at zero. In the non-degenerate case $\gamma > 0$ on $[0, \infty)$, implying in particular that $\gamma(0) > 0$, existence results are also available, see [14–16].

We now fix γ satisfying (1.3) and consider $(u^{in}, v^{in}) \in L_+^\infty(\Omega, \mathbb{R}^2)$, along with a global weak solution (u, v) to (1.1) in the sense of Definition 2.1. As a first step towards the identification of the large time limit of (u, v) , we collect obvious consequences of (1.1), the non-negativity of u and v , and the comparison principle.

Lemma 2.2. *For $t \geq 0$,*

$$\langle u(t) \rangle = M := \langle u^{in} \rangle \quad \text{and} \quad \|v(t)\|_\infty \leq V := \|v^{in}\|_\infty. \quad (2.1)$$

Moreover,

$$\int_0^\infty \|(uv)(t)\|_1 \, dt \leq \|v^{in}\|_1. \quad (2.2)$$

Proof. We integrate (1.1a) with respect to space and time and use the no-flux boundary conditions (1.1c) to obtain the first identity in (2.1). Similarly, we infer from (1.1b) and (1.1c) that

$$\frac{d}{dt} \|v(t)\|_1 + \int_0^t \|(uv)(s)\|_1 \, ds = \|v^{in}\|_1, \quad t \geq 0, \quad (2.3)$$

from which (2.2) readily follows. Finally, we use the comparison principle to deduce from (1.1b), (1.1c), and the non-negativity of uv that $v(t, x) \leq V$ for $(t, x) \in [0, \infty) \times \bar{\Omega}$, thereby completing the proof of (2.1). \square

We now define the auxiliary function

$$A(t, x) := \int_0^t (u\gamma(v))(s, x) \, ds, \quad (t, x) \in (0, \infty) \times \Omega,$$

and devote the next lemma to its properties.

Lemma 2.3. *The function A belongs to $L^\infty((0, \infty), H^1(\Omega))$ with*

$$\|A(t)\|_1 \leq \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, V)}, \quad t \geq 0, \quad (2.4a)$$

$$\|\nabla A(t)\|_2^2 \leq \|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, V)}, \quad t \geq 0. \quad (2.4b)$$

In addition, $t \mapsto A(t, x)$ is a non-decreasing function for a.e. $x \in \Omega$ and

$$A_\infty(x) := \sup_{t \geq 0} \{A(t, x)\} = \int_0^\infty (u\gamma(v))(s, x) \, ds, \quad x \in \Omega, \quad (2.5)$$

is well-defined and belongs to $H_+^1(\Omega)$. Also, for any $\vartheta \in H^1(\Omega)$,

$$\lim_{t \rightarrow \infty} \|A(t) - A_\infty\|_2 = \lim_{t \rightarrow \infty} \int_\Omega \nabla \vartheta \cdot \nabla (A(t) - A_\infty) \, dx = 0. \quad (2.6)$$

Proof. Owing to the non-negativity of u and γ ,

$$A(t_1, x) \leq A(t_2, x), \quad 0 \leq t_1 \leq t_2, \quad x \in \Omega,$$

while, for $t \geq 0$, it follows from (1.3), (2.1), and (2.2) that

$$\begin{aligned} \|A(t)\|_1 &= \int_0^t \int_\Omega (u\gamma(v))(s, x) \, dx ds \leq \|\gamma'\|_{L^\infty(0, V)} \int_0^t \int_\Omega (uv)(s, x) \, dx ds \\ &\leq \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, V)}, \end{aligned}$$

which proves (2.4a). Furthermore, the monotone convergence theorem implies that the function A_∞ defined by (2.5) belongs to $L_+^1(\Omega)$ and

$$\lim_{t \rightarrow \infty} \|A(t) - A_\infty\|_1 = 0. \quad (2.7)$$

We next infer from (1.1a), (1.1c), and the definition of A that, for $t \geq 0$,

$$u(t) - \Delta A(t) = u^{in} \quad \text{in } H^1(\Omega)'. \quad (2.8)$$

In particular, $A(t) - \langle A(t) \rangle = \mathcal{K}[u^{in} - u(t)]$ belongs to $H^1(\Omega)$ and we infer from (2.8) and the non-negativity of $u(t)$ and $A(t)$ that

$$\begin{aligned} \|\nabla A(t)\|_2^2 &= \langle -\Delta A(t), A(t) \rangle_{(H^1)', H^1} \\ &\leq \langle u(t) - \Delta A(t), A(t) \rangle_{(H^1)', H^1} = \int_\Omega u^{in} A(t) \, dx \\ &\leq \|u^{in}\|_\infty \|A(t)\|_1. \end{aligned}$$

Hence, by (2.4a),

$$\|\nabla A(t)\|_2^2 \leq \|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0, V)},$$

from which (2.4b) follows. Finally, we deduce the H^1 -regularity of A_∞ from (2.4b) by a weak compactness argument, whereas the convergence (2.6) is an immediate consequence of (2.4a), (2.4b), and (2.7). \square

We next turn to the convergence of v and begin with a classical energy estimate, which is available here thanks to the non-negativity of the right hand side of (1.1b).

Lemma 2.4. *For $t \geq 0$,*

$$\frac{d}{dt} \|v\|_2^2 + 2\|\nabla v\|_2^2 + 2\|v\sqrt{u}\|_2^2 = 0.$$

Lemma 2.5. *For each $p \in [1, \infty)$,*

$$\lim_{t \rightarrow \infty} \|v(t)\|_p = 0.$$

Proof. Introducing $P := \mathcal{K}[u - M]$ and $P^{in} := \mathcal{K}[u^{in} - M]$, we observe that $P = P^{in} - A$, so that

$$\|\nabla P\|_2 \leq \|\nabla P^{in}\|_2 + \|\nabla A\|_2 \leq c_1 := \|u^{in}\|_2 + \sqrt{\|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0,V)}}. \quad (2.9)$$

We next infer from (1.8a) and (2.3) that

$$\frac{d}{dt} \|v\|_1 = - \int_\Omega uv \, dx = - \int_\Omega (M - \Delta P)v \, dx = -M\|v\|_1 + \int_\Omega \nabla v \cdot \nabla P \, dx.$$

Hence, using (2.9) and Hölder's inequality,

$$\frac{d}{dt} \|v\|_1 + M\|v\|_1 \leq \|\nabla v\|_2 \|\nabla P\|_2 \leq c_1 \|\nabla v\|_2.$$

After integration with respect to time, we obtain

$$\|v(t)\|_1 \leq \|v^{in}\|_1 e^{-Mt} + c_1 \int_0^t e^{M(s-t)} \|\nabla v(s)\|_2 \, ds, \quad t \geq 0. \quad (2.10)$$

Now, by Hölder's inequality,

$$\begin{aligned} \int_0^t e^{M(s-t)} \|\nabla v(s)\|_2 \, ds &\leq \left(\int_0^t e^{M(s-t)} \, ds \right)^{1/2} \left(\int_0^t e^{M(s-t)} \|\nabla v(s)\|_2^2 \, ds \right)^{1/2} \\ &\leq \frac{1}{\sqrt{M}} \left(\int_0^t e^{M(s-t)} \|\nabla v(s)\|_2^2 \, ds \right)^{1/2}. \end{aligned}$$

Since

$$2 \int_0^\infty \|\nabla v(s)\|_2^2 \, ds \leq \|v^{in}\|_2^2$$

by Lemma 2.4, we deduce from the Lebesgue dominated convergence theorem that

$$\lim_{t \rightarrow \infty} \int_0^t e^{M(s-t)} \|\nabla v(s)\|_2^2 \, ds = 0.$$

Consequently,

$$\int_0^t e^{M(s-t)} \|\nabla v(s)\|_2 \, ds = 0 \quad (2.11)$$

and (2.10) and (2.11) entail that

$$\lim_{t \rightarrow \infty} \|v(t)\|_1 = 0,$$

thereby proving Lemma 2.5 for $p = 1$. To complete the proof, we use the above convergence, along with (2.1) and Hölder's inequality. \square

Thanks to the above analysis, we are now in a position to prove Theorem 1.1 and Proposition 1.2.

Proof of Theorem 1.1. According to Lemma 2.3, the function A_∞ introduced in Theorem 1.1 is well-defined and belongs to $H_+^1(\Omega)$. Setting $u_\infty = u^{in} + \Delta A_\infty \in H^1(\Omega)'$, we infer from (2.8) that, for $t \geq 0$ and $\vartheta \in H^1(\Omega)$,

$$\begin{aligned} \langle u(t) - u_\infty, \vartheta \rangle_{(H^1)', H^1} &= \langle u(t) - u^{in} + u^{in} - u_\infty, \vartheta \rangle_{(H^1)', H^1} \\ &= \langle \Delta(A(t) - A_\infty), \vartheta \rangle_{(H^1)', H^1} \\ &= - \int_{\Omega} \nabla(A(t) - A_\infty) \cdot \nabla \vartheta \, dx, \end{aligned}$$

and the right hand side of the above identity converges to zero as $t \rightarrow \infty$ due to (2.6). We have thus proved the convergence (1.6), whereas the convergence (1.7) is established in Lemma 2.5. As for the properties of u_∞ stated at the end of Theorem 1.1, they readily follow from (2.1), the non-negativity, and the convergence (1.6) \square

Proof of Proposition 1.2. The starting point is the estimate (2.4b) and the convergences (2.6) which imply that

$$\|\nabla A_\infty\|_2^2 \leq \|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0,V)}, \quad (2.12)$$

recalling that $V = \|v^{in}\|_\infty$. Since $\langle u_\infty \rangle = \langle u^{in} \rangle$ by Theorem 1.1, it follows from (2.12) and the definition of u_∞ that

$$\|u_\infty - u^{in}\|_{(H^1)'}^2 = \|\nabla \mathcal{K}[u_\infty - u^{in}]\|_2^2 = \|\nabla A_\infty\|_2^2 \leq \|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0,V)},$$

as stated in (1.9).

Assume now that (u^{in}, v^{in}) satisfies (1.10). It follows from (1.9) and (1.10) that

$$\begin{aligned} \|u_\infty - \langle u^{in} \rangle\|_{(H^1)'} &= \|u_\infty - u^{in} + u^{in} - \langle u^{in} \rangle\|_{(H^1)'} \\ &\geq \|u^{in} - \langle u^{in} \rangle\|_{(H^1)'} - \|u_\infty - u^{in}\|_{(H^1)'} \\ &> [\|u^{in}\|_\infty \|v^{in}\|_1 \|\gamma'\|_{L^\infty(0,V)}]^{1/2} - \|u_\infty - u^{in}\|_{(H^1)'} > 0. \end{aligned}$$

Consequently, $u_\infty \neq \langle u^{in} \rangle$, which completes the proof after noticing that the property $\langle u_\infty \rangle = \langle u^{in} \rangle$ established in Theorem 1.1 excludes that u_∞ coincides with any other constant. \square

ACKNOWLEDGMENTS

Enlightening (electronic) discussions with Michael Winkler on the topic studied in this paper are gratefully acknowledged. Part of this work was done while enjoying the kind hospitality of the Department of Mathematics, Indian Institute of Technology Roorkee.

REFERENCES

- [1] M. BURGER, PH. LAURENÇOT, AND A. TRESCASES, *Delayed blow-up for chemotaxis models with local sensing*, J. Lond. Math. Soc. (2), 103 (2021), pp. 1596–1617.
- [2] L. DESVILLETES, Y.-J. KIM, A. TRESCASES, AND C. YOON, *A logarithmic chemotaxis model featuring global existence and aggregation*, Nonlinear Anal. Real World Appl., 50 (2019), pp. 562–582.

- [3] L. DESVILLETES, PH. LAURENÇOT, A. TRESCASES, AND M. WINKLER, *Weak solutions to triangular cross diffusion systems modeling chemotaxis with local sensing*, *Nonlinear Anal.*, 226 (2023), pp. Paper No. 113153, 26.
- [4] K. FUJIE AND J. JIANG, *Boundedness of classical solutions to a degenerate Keller-Segel type model with signal-dependent motilities*, *Acta Appl. Math.*, 176 (2021), pp. Paper No. 3, 36.
- [5] K. FUJIE AND J. JIANG, *Comparison methods for a Keller-Segel-type model of pattern formations with density-suppressed motilities*, *Calc. Var. Partial Differential Equations*, 60 (2021), pp. Paper No. 92, 37.
- [6] K. FUJIE AND J. JIANG, *A note on construction of nonnegative initial data inducing unbounded solutions to some two-dimensional Keller-Segel systems*, *Math. Eng.*, 4 (2022), pp. Paper No. 045, 12.
- [7] K. FUJIE AND T. SENBA, *Global boundedness of solutions to a parabolic-parabolic chemotaxis system with local sensing in higher dimensions*, *Nonlinearity*, 35 (2022), pp. 3777–3811.
- [8] ———, *Global existence and infinite time blow-up of classical solutions to chemotaxis systems of local sensing in higher dimensions*, *Nonlinear Anal.*, 222 (2022), pp. Paper No. 112987, 7.
- [9] J. JIANG, PH. LAURENÇOT, AND Y. ZHANG, *Global existence, uniform boundedness, and stabilization in a chemotaxis system with density-suppressed motility and nutrient consumption*, *Comm. Partial Differential Equations*, 47 (2022), pp. 1024–1069.
- [10] H.-Y. JIN AND Z.-A. WANG, *Critical mass on the Keller-Segel system with signal-dependent motility*, *Proc. Amer. Math. Soc.*, 148 (2020), pp. 4855–4873.
- [11] E. F. KELLER AND L. A. SEGEL, *Initiation of slime mold aggregation viewed as an instability*, *J. Theoret. Biol.*, 26 (1970), pp. 399–415.
- [12] ———, *Traveling bands of chemotactic bacteria: a theoretical analysis*, *J. Theoret. Biol.*, 30 (1971), pp. 235–248.
- [13] PH. LAURENÇOT, *Long term spatial homogeneity for a chemotaxis model with local sensing and consumption*. preprint, 2023.
- [14] D. LI AND J. ZHAO, *Global boundedness and large time behavior of solutions to a chemotaxis-consumption system with signal-dependent motility*, *Z. Angew. Math. Phys.*, 72 (2021), p. 21. Id/No 57.
- [15] G. LI AND M. WINKLER, *Refined regularity analysis for a Keller-Segel-consumption system involving signal-dependent motilities*. arXiv: 2206.13327, to appear in *Applicable Anal.*, 2022.
- [16] ———, *Relaxation in a Keller-Segel-consumption system involving signal-dependent motilities*, *Commun. Math. Sci.*, 21 (2023), pp. 299–322.
- [17] H. LI AND J. JIANG, *Global existence of weak solutions to a signal-dependent Keller-Segel model for local sensing chemotaxis*, *Nonlinear Anal., Real World Appl.*, 61 (2021), p. 14. Id/No 103338.
- [18] C.-S. LIN, W.-M. NI, AND I. TAKAGI, *Large amplitude stationary solutions to a chemotaxis system*, *J. Differential Equations*, 72 (1988), pp. 1–27.
- [19] Y. TAO AND M. WINKLER, *Effects of signal-dependent motilities in a Keller-Segel-type reaction-diffusion system*, *Math. Models Methods Appl. Sci.*, 27 (2017), pp. 1645–1683.
- [20] Z.-A. WANG AND X. XU, *Steady states and pattern formation of the density-suppressed motility model*, *IMA J. Appl. Math.*, 86 (2021), pp. 577–603.
- [21] M. WINKLER, *Application of the Moser-Trudinger inequality in the construction of global solutions to a strongly degenerate migration model*, *Bull. Math. Sci.*, (2022), p. 16. Id/No 2250012.
- [22] ———, *A quantitative strong parabolic maximum principle and application to a taxis-type migration-consumption model involving signal-dependent degenerate diffusion*. arXiv: 2209.12724, to appear in *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 2022.
- [23] ———, *Global generalized solvability in a strongly degenerate taxis-type parabolic system modeling migration-consumption interaction*, *Z. Angew. Math. Phys.*, 74 (2023), p. 20. Id/No 32.
- [24] ———, *Stabilization despite pervasive strong cross-degeneracies in a nonlinear diffusion model for migration-consumption interaction*. preprint, 2023.
- [25] Y. XIAO AND J. JIANG, *Global existence and uniform boundedness in a fully parabolic Keller-Segel system with non-monotonic signal-dependent motility*, *J. Differ. Equations*, 354 (2023), pp. 403–429.

- [26] C. YOON AND Y.-J. KIM, *Global existence and aggregation in a Keller-Segel model with Fokker-Planck diffusion*, Acta Appl. Math., 149 (2017), pp. 101–123.

LABORATOIRE DE MATHÉMATIQUES (LAMA) UMR 5127, UNIVERSITÉ SAVOIE MONT BLANC, CNRS, F-73000
CHAMBÉRY, FRANCE

Email address: philippe.laurencot@univ-smb.fr