



**HAL**  
open science

## **Landiras virus, a novel hantavirus hosted by *Talpa aquitania* n.sp, a recently discovered South Eeuropean mole species**

Jean-Pierre André James Hugot, Bert Vanmechelen, Piet Maes

### ► **To cite this version:**

Jean-Pierre André James Hugot, Bert Vanmechelen, Piet Maes. Landiras virus, a novel hantavirus hosted by *Talpa aquitania* n.sp, a recently discovered South Eeuropean mole species. Bulletin de l'Académie Vétérinaire de France, 2023, 176, <10.3406/bavf.2023.71022>. <hal-04054158>

**HAL Id: hal-04054158**

**<https://hal.science/hal-04054158v1>**

Submitted on 31 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# LANDIRAS VIRUS, A NOVEL HANTAVIRUS HOSTED BY *TALPA AQUITANIA* N.SP, A RECENTLY DISCOVERED SOUTH EUROPEAN MOLE SPECIES

## LE VIRUS LANDIRAS, UN NOUVEL HANTAVIRUS HÉBERGÉ PAR *TALPA AQUITANIA* N. SP. RÉCEMMENT DÉCOUVERTE AU SUD-OUEST DE L'EUROPE

Par Jean-Pierre HUGOT, Bert VANMECHELEN, Piet MAES

(manuscrit soumis le 25 octobre 2022 et accepté le 24 janvier 2023)

### ABSTRACT

**Keywords:** *Hantavirus*, *Mammantavirinae*, *Orthohantavirus*, *Landiras virus*, *Talpa aquitania*

### INTRODUCTION

During the last 20 years, numerous new hantaviruses belonging to different species have been discovered in non-rodent micro-mammals. In the following, using the results of a Bayesian analysis, we describe *Landiras virus* (LDRV), a new *Orthohantavirus* discovered in a recently described new mole species: *Talpa aquitania* (Nicolas *et al.* 2015, 2017). The new virus species is compared with other hantaviruses, collected from moles, rodents or shrews, which are classified in the subfamily Mammantavirinae.

### MATERIALS AND METHODS

#### Sample collection

From 2011 to 2020, several hundred moles were captured in France and Northern Spain. The aim of this work was the search for hantaviruses likely to be hosted by moles. Moles are classifed pest species. This status authorizes different control methods, including trap killing, and no permits are required for fieldwork. Immediately after trapping, moles were stored at -20° C until further processing. Lung, kidney, heart, liver, and spleen tissues were aseptically removed and stored in RNAlater Stabilization Solution (Ambion). The genetic study of the

samples allowed the discovery of a new species of mole, present in France and in the north of Spain: *Talpa aquitania* (Hugot *et al.* 2014; Nicolas *et al.* 2017).

#### Hantavirus screening

One hundred and seventy (170) lung samples collected in mole specimens identified as *T. aquitania* were sent to the Virology laboratory of Leuven University (Belgium). Total RNA was extracted from mole tissue with the RNeasy Mini kit (Qiagen) according to the manufacturer's instructions. A nested degenerate RT-PCR was performed using the OneStep RT-PCR kit (Qiagen) with primers targeting a conserved region in the polymerase gene, as described previously (Klempa *et al.* 2006). PCR amplicons were purified using ExoSAP-IT PCR Product Cleanup Reagent (Affymetrix) and sent to Macrogen for Sanger sequencing. Sequences were manually inspected using Chromas 2.6 (Technelysium) and a consensus sequence was derived with Seqman 7.0 (DNASTar). Two samples were found positive for Hantaviruses. Both were collected on July 17<sup>th</sup>, 2016 in Landiras (Gironde department) in the south-west of France (N-44,5667; W-0,4167), identified as *T. aquitania* in the French National Museum of Natural History and referenced in the collections as MNHN-ZM-2017-2257 and MNHN-ZM-2017-2258.

1. Department of Systematics and Evolution, Biodiversity, Museum National d'Histoire Naturelle, Paris, France, E-mail: hugot@mnhn.fr

2. Laboratory of Clinical and Epidemiological Virology, Department of Microbiology and Immunology, Rega Institute for Medical Research, KU Leuven, Belgium



## S segment sequencing

To obtain the complete sequence of the S (nucleocapsid) gene, an additional nested RT-PCR was performed on the same total RNA extract, using the OneStep RT-PCR kit. The first round of this PCR used a custom primer that contains an 8-nt motif targeting the conserved panhandle sequence present on both ends of the S segment and a long 5' overhang (5'-GTTTCCCCTGGAGGATATAGTAGTA-3'). The inner PCR reaction was performed using a primer targeting the overhang of the first-round primer (5'-GTTTCCCCTGGAGGATA-3'). The resulting amplicon was used to prepare a sequencing library with the SQK-LSK109 and EXP-NBD104 kits (Oxford Nanopore Technologies) according to the manufacturer's instructions. The resulting library was run on a MinION R9.4.1 flow cell and the data were base called using the HAC model of Guppy v3.0.3. Fastq-files were assembled using Canu v1.8 (Koren *et al.* 2017) and the resulting contig was polished using Medaka v0.8. The resulting contig was checked for errors by mapping back the sequencing reads using CLC genomics work-bench 20.0.2 (Qiagen). Only MNHN-ZM-2017-2257 allowed to obtain a complete sequence (1966 nt) were submitted to GeneBank as ON944104 access number and used for the phylogenetic analysis.

## Phylogenetic analysis

Multiple sequence alignments of hantavirus sequences were generated in MAFFT and manually corrected using SeaView 4.6.3 (Katoh *et al.* 2002; Gouy *et al.* 2010). An alignment of the complete coding region of the S segment was used in subsequent Bayesian inference using MrBayes (Huelsenbeck and Ronquist 2001) using a General Time Reversible (GTR) substitution model and a discretized distribution with default priors. Two independent Markov Chain Monte Carlo analyses, each using 3 heated and 1 cold chain, were run until adequate effective sample sizes (ESS > 200) were obtained. A consensus tree was built employing a burn-in of 25% and rooted using the "midpoint root" option.

## RESULTS AND DISCUSSION

Analysis of the S (nucleocapsid) gene of *Landiras virus* shows that it is close to the *Orthohantavirus Bruges* (NC034394) previously described in several specimens of *T. europaea* captured in Belgium, Germany and the United Kingdom (Laenen *et al.* 2017) (Figure 1). Both are very different from Nova virus (FJ539168) also discovered in *T. europaea* (Kang *et al.* 2009; Hugot *et al.* 2014; Hun Gu *et al.* 2014). As *T. europaea* and *T. aquitania* have contiguous ranges, this raises the question of a possible virus host switching between the two mole species. Further investigations are needed to gain insights into hantaviruses ecology, transmission dynamics and virus-host evolution.




Figure 1 : Bayesian phylogenetic analysis of the nucleocapsid (S) gene. Numbers at the nodes indicate posterior clade probability values.

## REMERCIEMENTS

The results presented here could not have been obtained without the help of the French network *Taupe'Green* and its founder, Jerome Dormion. We particularly thank Jean-Luc Laclamette who collected the material from which the *Landiras virus* was isolated.

## BIBLIOGRAPHIE

- Gouy M, Guindon S, Gascuel O. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* 2010; 27:221–224
- Hugot JP, Hun Gu S, Feliu C, Ventura J, Ribas A, Dormion J *et al.* Genetic variability of *Talpa europaea* and Nova hantavirus (NVAV) in France. *Bull. Acad. Vét. France* 2014; 167(3): 277-284.
- Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 2001; 17:754–755.
- Hun Gu S, Dormion J, Hugot JP, Yanagihara R. High prevalence of Nova hantavirus infection in the European mole (*Talpa europaea*) in France. *Epidemiol Infect.* 2014; 142(6): 1167–1171.
- Kang HJ, Bennett SN, Sumibcay L, Satoru A, Hope AG, Mocz G, Song JW, Cook JA, Yanagihara R. Evolutionary insights from a genetically divergent hantavirus harbored by the European common mole (*Talpa europaea*). *PLOS One* 2009; 4(7): e6149.
- Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 2002; 30(14):3059–3066.
- Klempa B, *et al.* Hantavirus in African wood mouse, Guinea. *Emerg Infect Dis.* 2006; 12(5):838–840.
- Koren S, Walenz BP, Berlin K, Miller JR, Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Research* 2017; doi:10.1101/gr.215087.116
- Laenen L, Vergote V, Kafetzopoulou LE, Wawina Bokalinga T, Vassou D, Cook JA *et al.* Bruges virus, a novel European mole hantavirus, co-infects its host together with Nova virus without signs of genetic interaction. *GBE* 2017; 10(1): 45–55. doi:10.1093/gbe/evx268
- Nicolas V, Martinez-Vargas J, Hugot JP. *Talpa aquitania* nov. sp. (Talpidae, Soricomorpha) a new mole species from southwest France and north Spain. *Bull. Acad. Vét. France* 2015; 168(4): 1-6.
- Nicolas V, Martinez-Vargas J, Hugot JP. *Talpa aquitania* sp. nov. (Talpidae, Soricomorpha) a new mole species from SW France and North Spain. *Mammalia* 2017; 81(6): 641-642.