



**HAL**  
open science

## The role of fire disturbances, human activities and climate change for long-term forest dynamics in upper-montane forests of the central Dinaric Alps

Eleonora Cagliero, Laure Paradis, Niccolò Marchi, Zsuzsa Lisztes-Szabó, Mihály Braun, Katalin Hubay, Pierre Sabatier, Milić Čurović, Renzo Motta, Velibor Spalevic, et al.

### ► To cite this version:

Eleonora Cagliero, Laure Paradis, Niccolò Marchi, Zsuzsa Lisztes-Szabó, Mihály Braun, et al.. The role of fire disturbances, human activities and climate change for long-term forest dynamics in upper-montane forests of the central Dinaric Alps. *The Holocene*, 2023, 33 (7), pp.827-841. 10.1177/09596836231163515 . hal-04059282

**HAL Id: hal-04059282**

**<https://hal.science/hal-04059282>**

Submitted on 5 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Accepted Manuscript version of**

2 Cagliero E, Paradis L, Marchi N, Lisztes-Szabó Z, Braun M, Hubay K, et al. ( ( n.d.)) The role  
3 of fire disturbances, human activities and climate change for long-term forest dynamics in  
4 upper-montane forests of the central Dinaric Alps. *The Holocene*.

5 <https://doi.org/10.1177/095968362311635>

7 **The role of fire disturbances, human activities and climate change for long-term forest**  
8 **dynamics in upper-montane forests of the central Dinaric Alps**

9 Eleonora Cagliero<sup>a,b</sup>, Laure Paradis<sup>a</sup>, Niccolò Marchi<sup>b</sup>, Zsuzsa Lisztes-Szabó<sup>c</sup>, Mihály Braun<sup>c</sup>, Katalin  
10 Hubay<sup>c</sup>, Pierre Sabatier<sup>d</sup>, Milić Ćurović<sup>e</sup>, Velibor Spalevic<sup>e</sup>, Renzo Motta<sup>f</sup>, Emanuele Lingua<sup>b</sup>, Walter  
11 Finsinger<sup>a</sup>

13 <sup>a</sup> ISEM, University of Montpellier, CNRS, IRD, EPHE, Montpellier, France

14 <sup>b</sup> Department of Land, Environment, Agriculture and Forestry (TESAF), University of Padova, 35020 Legnaro (PD), Italy

15 <sup>c</sup> Isotope Climatology and Environmental Research Centre (ICER), Institute for Nuclear Research, 4026 Debrecen, Hungary

16 <sup>d</sup> EDYTEM, Université Savoie-Mont Blanc, CNRS, 73370 Le Bourget du Lac, France

17 <sup>e</sup> Biotechnical Faculty, University of Montenegro, 81000 Podgorica, Montenegro

18 <sup>f</sup> Department of Agricultural, Forest and Food Sciences (DISAFA), University of Torino, 10095 Grugliasco (TO), Italy

19  
20 **Abstract**

21 We present the first high-resolution Holocene pollen, plant-macrofossil, and charcoal records from the  
22 upper-montane zone in the central Dinaric Alps. Drawing on these new records from well-dated  
23 lacustrine sediments of Zminje Jezero (ca. 1500 m a.s.l.; Montenegro) and on independent chironomid-  
24 inferred summer temperatures, we explore long-term ecosystem responses to variations in climate, fire  
25 disturbances and land use, as well as legacy effects of past environmental changes.

26 A mixed spruce-fir forest established in the upper-montane zone around 9500 cal BP, and *Fagus*  
27 *sylvatica* became co-dominant with the two conifers after 5000 cal BP. Prehistoric land-use pressure  
28 was overall remarkably low, but increased since 2000 cal BP and was highest after the Middle Ages.

29 We found a significant positive relationship between biomass burning and summer temperature,  
30 indicating that fires were mostly climate driven. *Picea abies* was insensitive to summer temperature,  
31 biomass burning and human impact, which supports the view that spruce forests may not be significantly  
32 impacted by fire. In contrast, *Abies alba* and other disturbance-sensitive trees (*Tilia*, *Ulmus*, *Fraxinus*  
33 *excelsior*-type) show significant negative responses to land-use pressure and positive responses to  
34 summer temperature. This supports the notion that these species may be well-adapted to warmer-than  
35 present summer temperatures and that their populations declined in recent millennia due to land-use  
36 activities. Conversely, *F. sylvatica* was sensitive to summer temperatures but was promoted by low  
37 biomass burning, indicating that its expansion in the spruce-fir dominated forest was enhanced by the  
38 onset of cooler and possibly also moister climatic conditions as well as by fire disturbances.

39  
40 **Keywords**

41 Holocene, vegetation dynamics, mountain forests, fire history, vegetation-fire interactions, Montenegro

42  
43 **Corresponding author:**

44 Walter Finsinger

45 Institut des Sciences de l'Évolution de Montpellier (ISEM), Université de Montpellier

46 2, place Eugène Bataillon, 34095 Montpellier, France

47 E-mail address: [walter.finsinger@umontpellier.fr](mailto:walter.finsinger@umontpellier.fr)

## 48 **Introduction**

49 Assessing the responses of European mountain forests to changing environmental conditions is crucial  
50 to develop adaptation and management strategies to possible ecosystem shifts that may cause the loss  
51 of important ecosystem services, such as risk prevention (from avalanches, landslides or rockfall),  
52 recreation, and the maintenance and promotion of biodiversity (Klopčič et al., 2017). Such assessments  
53 are particularly important given that there is strong support for the hypothesis that climate change could  
54 markedly modify disturbance regimes, with a likely increase of some disturbances (e.g. fire) in a  
55 warming world (Seidl et al., 2017).

56 However, understanding natural ecological processes is challenging as processes often play out over  
57 long time scales (centuries, millennia), particularly when long-lived species such as trees are involved.  
58 The long time horizon of palaeoecological records permits to study ecosystem responses under  
59 substantially different environmental conditions than the present ones, including temperatures analogous  
60 to those predicted by future climate changes (approximately 1-2°C higher than present; Carter et al.,  
61 2018; Morales-Molino et al., 2021; Samartin et al., 2017). Moreover, current ecosystems and ecological  
62 processes often carry legacy effects of past environmental changes and anthropogenic impacts (Cagliero  
63 et al., 2022; Feurdean et al., 2009; Grindean et al., 2019; Morales-Molino et al., 2022), which have often  
64 altered species-environment relationships (Tinner et al., 2013). Thus, taking into account long-term  
65 records may be important to gain more accurate environmental-change response assessments, especially  
66 in Europe, where anthropogenic manipulation of nature started several millennia ago (Birks and Tinner,  
67 2016).

68 The long-term history of mountain ecosystems varies across regions and is arguably contingent on  
69 patterns of human settlement, land use, and socioeconomic development. In this context, the central  
70 Dinaric Alps are an interesting area as its land-use history may differ from those of other European  
71 mountain regions. There is little doubt that the Balkan Peninsula acted as a land bridge from the Near  
72 East, across Asia Minor and towards Central and Western Europe (Forenbaher et al., 2013). While it  
73 seems that the Neolithic spread mainly across the fertile plains along the Danube valley in the east  
74 (Starčevo culture) and along the Adriatic coastal strip in the west (Impresso culture) around the 6<sup>th</sup>  
75 millennium BCE (8000-7000 cal BP; Borić et al., 2019), there is evidence to suggest that the

76 mountainous hinterland of the central Dinaric Alps remained an agricultural frontier zone, eventually  
77 even until the Late Neolithic (Forenbaier and Miracle, 2005). The frontier-zone hypothesis is consistent  
78 with modelled anthropogenic deforestation (Kaplan et al., 2009), which suggests that during the past  
79 3000 years relatively low forest clearance occurred when compared to other regions of Europe, and that  
80 a large fraction of usable land in the region was not heavily exploited until the Middle Ages (600-1460  
81 CE; 1350-490 cal BP). By contrast, charcoal production, mining activities, and grazing likely occurred  
82 in the interior area of the Dinaric Alps since the Bronze Age (3500-600 BCE; 5450-2550 cal BP), and  
83 intensified during the Iron Age (600-200 BCE; 2550-2150 cal BP) and the Ottoman Empire (1460-1800  
84 CE; 490-150 cal BP) (Kranjc, 2009; Longman et al., 2018). However, it is not clear if these activities  
85 were widespread across the mountain region, or whether human activities rather concentrated in some  
86 areas. In the latter case, large forested regions in the interior range may have remained relatively intact  
87 until the present (Nagel et al., 2017) or managed with low intensity silvicultural systems during recent  
88 centuries (Boncina, 2011). Low historical forest clearance in the montane belt may have preserved fir  
89 (*Abies alba* Mill.), beech (*Fagus sylvatica* L.), and spruce (*Picea abies* (L.) Karst.) old-growth forests  
90 (Cagliero et al., 2022; Motta et al., 2011; Sabatini et al., 2018). As these tree species are widespread and  
91 often form mixed forests in European mountains (Hilmers et al., 2019), exploring their long-term  
92 responses under low land-use pressure may be useful to improve knowledge on the role of disturbances  
93 by fire and climate. However, the impact of Neolithic and of more recent cultures on the interior  
94 mountain region is still weakly constrained. In contrast to other European mountain regions, the area is  
95 both archeologically and palaeoecologically under-documented (Borić et al., 2019; Finsinger et al.,  
96 2017; Vander Linden et al., 2014).

97 Thus, to provide new insights into the Holocene vegetation dynamics and into vegetation responses to  
98 disturbances by fire and to changes in climate and land use we studied sediments from Zminje Jezero  
99 (Fig. 1), a lake at 1535 m a.s.l., thus in the montane belt of the central Dinaric Alps. Our main aims are  
100 to (1) assess the long-term Holocene vegetation dynamics of mixed montane forests, (2) characterize  
101 the Holocene land-use and fire histories, (3) investigate the influence of the main drivers of fire  
102 dynamics (climate and human impact), and (4) track responses of the dominant montane trees (*A. alba*,

103 *F. sylvatica*, *P. abies*, as well as of other disturbance-sensitive trees such as *Ulmus*, *Tilia*, and *Fraxinus*  
104 *excelsior*) to variations in climate, fire and human impact.

105

106

## 107 **Material and methods**

108

### 109 *Study area and study site*

110 In the Dinaric Alps, vegetation is influenced by the interaction of mountainous reliefs, pedological  
111 conditions, and proximity to the Adriatic Sea (Horvat et al., 1974; Nagel et al., 2017). Mediterranean  
112 evergreen sclerophyllous vegetation dominated by *Pinus halepensis* Mill., *Quercus ilex* L. and *Olea*  
113 *europaea* L. occurs near the coast. Mixed submediterranean deciduous woodlands with *Carpinus betulus*  
114 L., *Fraxinus ornus* L., deciduous *Quercus* L., *Ostrya carpinifolia* Scop., *Carpinus orientalis* Mill., and  
115 *Castanea sativa* Mill. occur up to c. 700 m asl. The valley floodplains are commonly occupied by *Alnus*  
116 *glutinosa* (L.) Gaertn., *Quercus robur* L., *Fraxinus* sp. and *Ulmus minor* Mill.. In the montane belt (from  
117 700 to 2000 m asl), forests are dominated by *Pinus nigra* J.F. Arnold, *F. sylvatica*, and mixed beech-fir-  
118 spruce forests. Other species occurring in the montane belt include *Pinus sylvestris* L., *O. carpinifolia*,  
119 *Acer* L., *Ulmus* L., *Tilia* L., and *Fraxinus excelsior* L.. The subalpine vegetation belt is generally above  
120 c. 2000 m asl and is dominated by *Pinus mugo* Turra, *Juniperus communis* L., and *Alnus viridis* (Chaix)  
121 DC.

122 Zminje Jezero (Snake lake; 43°09'21" N 19°04'14" E; 1535 m asl) is a small 1.2-ha large and 9.5-m  
123 deep glacial-origin lake (Fig. 1) whose water is drained into the Crno Jezero (1460 m asl) by a small  
124 stream. The lake is located on the northwest part of a high karst plateau in the Durmitor massif (Annys  
125 et al., 2014) in a valley adjacent to the one currently hosting the last surviving glacier in Montenegro,  
126 Debeli namet (Hughes et al., 2011). Given its elevation, Zminje Jezero is within the limits of terminal  
127 moraines that were deposited before the Lateglacial Interstadial (Hughes et al., 2011).

128 The vegetation in the surroundings of the lake is dominated by a dense spruce-fir forest (Fig. 1) with  
129 lesser amounts of beech, *Acer* sp., *Sorbus aucuparia* L., *Rhamnus fallax* Boiss., and *Vaccinium myrtillus*  
130 L.. However, *P. sylvestris*, *Pinus nigra* subsp. *nigra*, *Betula pendula* Roth, and *Pinus heldreichii* H.

131 Christ are also well represented in the Durmitor massif. To the east, there is a town (Žabljak; 1450 m  
132 asl) with pastures and agriculture lands. The lake shores host a rich community of wetland plants,  
133 including *Molinia caerulea* (L.) Moench, *Comarum palustre* L., *Carex* sp., *Sphagnum* sp., *Parnassia*  
134 *palustris* L., *Filipendula ulmaria* (L.) Maxim., and *Eriophorum vaginatum* L.. The current tree line is at  
135 c. 1950 m asl and is mainly formed by spruce with beech (Bui, 1975). Above the tree line are *P. mugo*  
136 and *J. communis* shrublands, alpine meadows, and rocks.

137 Climate is continental with a mean annual air temperature of 5.1°C and a mean annual precipitation of  
138 c. 1450 mm at Žabljak (Annys et al., 2014). In the northern and central parts of the mountain range,  
139 bedrock consists of Mesozoic sedimentary rocks and thick Middle and Upper Triassic and Upper  
140 Jurassic limestones, while in the southern part Upper Cretaceous flysch are predominant (Mirković,  
141 1985).

142 Zminje Jezero is located in a zone of special protection of the Durmitor National Park, where only  
143 cutting of naturally collapsed trees (so-called “sanitary cuttings”), clearing of forest roads, and tourism  
144 are allowed (Srdanović and Pavić, 2013; UNESCO and IUCN, 2018). Before the establishment of the  
145 National Park in 1952 CE, the area was informally protected since 1907 CE.

146

#### 147 *Field work and sediment-core correlation*

148 We studied the lake floor with an echo sounder connected to a Garmin GPS device (Fig. 1d) and  
149 collected two overlapping sediment cores (sections 1-m long and 6 cm in diameter) from the deepest  
150 part of the lake at a water depth of 9.3 m using a modified Livingstone piston corer and an UWITEC  
151 gravity corer. All core sections were split longitudinally and photographed at ISEM (University of  
152 Montpellier) to visually describe the sediments and align the sections with the aid of Corelyzer v2.1.1  
153 (CSDCO/LacCore-University of Minnesota). To assess sediment composition, we used an X-ray  
154 fluorescence (XRF) AVAATECH core scanner at 5 mm resolution (with the two following settings: 10  
155 kV/0.09 mA for 15 s and 30 kV/0.09 mA for 30 s), and then normalised the values using the centre-log-  
156 ratio method (CLR, Weltje et al., 2015). We correlated the sections based on marker layers and on  
157 selected XRF records. Thereafter, one core half was cut into 1-cm slices and samples were stored in zip-  
158 lock bags at 4°C.

159

160 *Chronology*

161 We modelled the depth-age relationship (Fig. 2) with RBacon v2.4.1 (Blaauw and Christen, 2011) using  
162 41 control points (Table 1 and Table S1), including 19 AMS  $^{14}\text{C}$  dates from terrestrial plant macrofossils  
163 calibrated using the IntCal20 dataset (Reimer et al., 2020), one pollen-inferred age (onset of the  
164 Holocene; Giesecke et al., 2014), and 21 control points derived from a chronology of short-lived  
165 radionuclides ( $^{210}\text{Pb}$  and  $^{137}\text{Cs}$ ; Fig. S1) that was built with the constant flux constant sedimentation  
166 (CFCS) using the serac package (Brueel and Sabatier, 2020). We excised visually identified event-deposit  
167 layers matching peaks of K, Ti, Rb, Zr, and Sr (Fig. 2), as they most probably represent sediment  
168 deposition that occurred over very short time spans (Finsinger et al., 2021; Heiri et al., 2003).

169

170 *Pollen, spores and plant-macrofossils analyses*

171 We processed 78  $1\text{-cm}^3$  samples for pollen analysis following standard physical and chemical  
172 treatments, including treatments with HCl, KOH, sieving with a  $500\text{-}\mu\text{m}$  mesh and decanting, HF,  
173 acetolysis, and mounted the fuchsin-stained residues on slides with glycerol. To calculate influxes ( $\#\text{cm}^{-2}\text{yr}^{-1}$ ), we added *Lycopodium* tablets (Stockmarr, 1971). We identified and counted pollen, stomata  
174 and the dung-fungi *Sporormiella* spore using determination keys and photographic atlases (Beug, 2004;  
175 Cagliero et al., 2022; Davis, 1987; Finsinger and Tinner, 2005; Moore et al., 1998; Reille, 1992) as well  
176 as the pollen and stomata reference collections at ISEM (University of Montpellier). We counted at least  
177 300 terrestrial pollen grains at  $400\times$  magnification and calculated percentages relative to the terrestrial  
178 pollen sum, which includes pollen of trees, shrubs, upland herbs, ferns, and vascular cryptogams and  
179 excludes pollen of obligate aquatic plants.

181 The pollen diagram was divided into pollen assemblage zones using optimal partitioning by sums-of-  
182 squares with square-root-transformed proportions of pollen types included in the terrestrial pollen sum,  
183 and exclusion of rare taxa (abundance  $<5\%$ ; Birks and Gordon 1985). The number of statistically  
184 significant zones was determined by comparison with the broken-stick model (Bennett, 1996) with  
185 Psimpoll v4.26 (Bennett, 2008).

186 For plant-macrofossils analyses, we processed 203 samples. We measured sample volumes ( $5.6 \pm 1.0$   
187  $\text{cm}^3$ ) by water displacement and then sieved them with a 100- $\mu\text{m}$  mesh sieve. We identified and counted  
188 macrofossils with a stereomicroscope at 7.5-60x magnification using atlases (Birks, 2017; Katz et al.,  
189 1965; Schoch et al., 1988) and the reference collection at ISEM (University of Montpellier) and stored  
190 them at  $-18^\circ\text{C}$  in Milli-Q water. Plant-macrofossil abundances are expressed as concentrations ( $\# \text{cm}^{-3}$ ).  
191 To assess land-use pressure, we used the abundances of *Secale cereale* pollen (a primary anthropogenic  
192 indicator), secondary indicators (as the sum of adventives and apophytes), and cultural indicators (as the  
193 sum of Cerealia-type and *Plantago lanceolata*-type pollen) (Behre, 1981; Tinner et al., 2003). These  
194 indexes include pollen of cultivated plants, ruderals, and plants of meadows (Deza-Araujo et al., 2020).  
195 *Veratrum album* was added to the apophyte category, as it is avoided by cattle due to its toxicity  
196 (Spiegelberger et al., 2006). We also assessed land-use pressure in the wider region using the OJC index  
197 (Mercuri et al., 2013), which is based on the abundance of woody crops (*Olea europaea* L., *Juglans*  
198 *regia* L., and *Castanea sativa* Mill.) that grow at lower elevations and along the coast of the Adriatic  
199 Sea. Despite their presence in uncultivated areas, an increase in their abundances is often due to broad-  
200 scale cultivation (Deza-Araujo et al., 2020).

201

#### 202 *Charcoal analyses*

203 We counted microscopic charcoal particles in 42 pollen slides spread along the sediment core (Finsinger  
204 and Tinner, 2005; Tinner and Hu, 2003) to obtain a low-resolution record of regional fire activity. For  
205 macroscopic charcoal analyses, we took contiguous 1- $\text{cm}^3$  samples and treated them without the use of  
206 an orbital shaker with a 5%  $\text{NaPO}_3$  solution overnight and then with a 15%  $\text{H}_2\text{O}_2$  solution for 24h  
207 (Schlachter and Horn, 2010). Subsequently, samples were sieved through a 150- $\mu\text{m}$  mesh (Higuera et  
208 al., 2005) and charcoal particles were identified and counted under a stereomicroscope at 7.5-60x  
209 magnification. The area of charcoal particles was not estimated since it is generally highly correlated to  
210 charcoal counts (Finsinger et al., 2014; Tinner et al., 1998). To account for variations in sedimentation  
211 rate (Whitlock and Larsen, 2001), both microscopic and macroscopic charcoal counts were converted  
212 to charcoal-accumulation rates (influx;  $\# \text{cm}^{-2} \text{yr}^{-1}$ ), hereafter denoted mCHAR and MCHAR,  
213 respectively.



214

215 *Fire episodes and fire-regime changes*

216 To reconstruct regional fire activity and landscape-scale biomass burning we used the long-term trends  
217 of two complementary proxies, mCHAR and MCHAR. While their potential source areas are similar (c.  
218 40-km distance from study sites; Adolf et al., 2018), the records may differ, with MCHAR potentially  
219 reflecting fires within a smaller distance (Finsinger et al., 2017). To determine significant changes in  
220 landscape-scale biomass burning, we identified change points based on the mean and the variance in  
221 MCHAR and tested their independence from variations in the modelled sediment-accumulation rates  
222 (Finsinger et al., 2018).

223 To detect fire episodes within a few km distance around the study site (Higuera et al., 2010), we analysed  
224 the MCHAR record with tapas v0.1.2 (Finsinger and Bonnici, 2022). This analysis involved resampling  
225 the record to a constant resolution of 33 years (i.e. the 3<sup>rd</sup> quartile of the sampling-interval distribution).  
226 Thereafter, we decomposed the record into a low-frequency background and a high-frequency peak  
227 component using a loess robust to outliers with a smoothing-window width of 2000 years. The suitability  
228 of the record for peak detection was evaluated through the signal-to-noise index (Kelly et al., 2011).  
229 Peaks were evaluated using the 95<sup>th</sup> percentile of the modelled noise distribution obtained with locally  
230 defined Gaussian mixture models for each 2000-year portion of the record (Higuera et al., 2009). The  
231 peaks were further screened (Higuera et al., 2010) before calculating the fire return intervals (FRI: years  
232 between adjacent fire episodes).

233

234 *Ecosystem responses*

235 We used generalized additive models (GAMs) to investigate ecosystem responses to variations in  
236 climate, fire and land use. GAMs are suited for this purpose as they can unfold nonlinear relationships  
237 between a response variable and a smoothed function of the predictor variable (Hastie and Tibshirani,  
238 1986). Specifically, we explored the responses of fires to variations in climate and land-use pressure,  
239 and the responses of dominant forest canopy taxa (*P. abies*, *A. alba*, *F. sylvatica*, *F. excelsior*-type, *Tilia*,  
240 and *Ulmus*) to variations in climate, fires and land-use pressure (Carter et al., 2020; Colombaroli et al.,  
241 2010; Morales-Molino et al., 2021).

242 As climate data, we used chironomid-inferred summer (July) surface-air temperature anomalies from  
243 Lago Verdarolo (1390 m asl) (Samartin et al., 2017) that is in a biogeographically comparable ecosystem  
244 in the Tuscan-Emilian Apennines where current vegetation is dominated by fir and beech. Comparisons  
245 between instrumental data and climate model runs (Samartin et al., 2017) show that at interannual to  
246 multidecadal timescales summer temperature variations in the Tuscan-Emilian Apennines are well  
247 correlated with grid cells in Montenegro (Pearson's correlation coefficients are 0.8-1 for multidecadal  
248 variability and 0.6-0.7 for interannual variability).

249 We assumed a quasi-Poisson distribution, log-transformed both the response and the predictor variables  
250 to stabilize their variances, and fitted GAMs using restricted maximum likelihood (REML) smoothness  
251 selection (Carter et al., 2020) with mgcv v1.8-38 (Wood, 2017). We chose a base period from 8000 cal  
252 BP to the present because secondary anthropogenic indicators were very abundant during the early  
253 Holocene, when human impact was negligible (Carter et al., 2020).

254

255

## 256 **Results and interpretation**

257

### 258 *Chronology and sediment composition*

259 The sediments cover part of the Younger Dryas and the entire Holocene, from c. 12,340 cal BP to present  
260 (Fig. 2). The reliability of the chronology in its most recent part is broadly confirmed (Fig. S2) by the  
261 match between the expected and modelled ages of Pb peaks during the Roman period and the Middle  
262 Ages (McConnell et al., 2018; Renberg et al., 2001). Sediment deposition times vary between 6-65 years  
263  $\text{cm}^{-1}$  (median = 21.6 years  $\text{cm}^{-1}$ ).

264 At the base, from 615 cm to 575 cm (12,340-11,640 cal BP), the sediments consist of light-brown sandy  
265 and silty clay (Fig. 2). From 575 cm to 479 cm (11,640-10,300 cal BP), there are 1-2-cm thick light grey  
266 silty-clay layers that alternate with darker and organic-richer layers (gradual boundaries), and a c. 28-  
267 cm thick sequence of five graded deposits (529-500 cm), each one characterized by a distinct fining-  
268 upward trend. Above 479 cm, sediments are mainly composed of dark-brown organic gyttja, which is  
269 interrupted by three minerogenic layers of fine grey detrital clay with sharp stratigraphic boundaries at

270 445-441 cm, 413-410 cm, and 283-281 cm depth. In keeping with the visual assessment of the sediments,  
271 minerogenic elements (K, Ti, Rb, Zr, and Sr) are most abundant below 479 cm (10,300 cal BP) as well  
272 as in conjunction with the minerogenic layers.

273

#### 274 *Long-term vegetation dynamics*

275 The pollen record (Figs. 3-4) was divided into four statistically significant assemblage zones at 566.5,  
276 455.5, and 359.5 cm (corresponding to 11,500, 9410, and 6400 cal BP) and three non-significant  
277 subzones with boundaries at 200.5 cm and 58.5 cm (2090 and 430 cal BP).

278 Pollen assemblages from 12,340 to 11,500 cal BP (ZMN-1; 615-566.5 cm) are dominated by *Pinus* and  
279 herb pollen (mainly *Artemisia*, Poaceae and Chenopodiaceae). Both stomata and plant macrofossils  
280 attest the local presence of *Betula* and *Pinus*, the latter presumably as low-density stands of dwarf  
281 mountain pine (*P. mugo*) or pine trees (*P. sylvestris*, *P. heldreichii*, or *P. nigra*). Cerealia-type pollen  
282 are likely associated with wild Poaceae, as Cerealia-type pollen were also found elsewhere long before  
283 the development of agriculture (Dörfler, 2013). Such assemblages are typical of the Central European  
284 Younger Dryas (12,850-11,650 cal BP; Rasmussen et al., 2014) and are indicative of an open and arid  
285 steppe-tundra with cold-tolerant pine woodlands. Rising arboreal pollen (from c. 45 to 65%) indicate a  
286 gradual closure of the woodland during the Younger Dryas cold stage, including mixed oak woodlands  
287 at lower altitude.

288 *Picea abies* stomata unequivocally indicate a rapid altitudinal shift of this tree species at the Younger  
289 Dryas-Holocene transition, when a rapid climatic change including a warming of c. 2-4°C occurred  
290 (Heiri et al., 2014; Tóth et al., 2015). The woodlands closed further from 11,500 to 9410 cal BP (ZMN-  
291 2; 566.5-455.5 cm), as attested by an increase of arboreal pollen (from c. 65% to 80%). *F. excelsior*-  
292 type, *Ulmus*, and *Tilia* pollen first appeared at 11,500 cal BP marking the expansion of meso-  
293 thermophilous trees at lower altitude, a characteristic feature of pollen records from the Balkan  
294 Peninsula and the adjacent European Alps (Caf et al., 2022; Vescovi et al., 2007; Willis, 1994). *Betula*  
295 populations were denser than during the Younger Dryas, as shown by both pollen and plant macrofossils.  
296 However, the abundance of light-demanding pioneer trees and shrubs (*Ephedra*, *Juniperus* and *Betula*)  
297 decreased in conjunction with the local expansion of spruce and fir populations starting between 10,400

298 and 9600 cal BP. Finds of *Sporormiella* dung-fungi spores indicate the presence of wild herbivores  
299 around the lake between 11,500 and 10,400 cal BP, as also observed on the Swiss Plateau (Rey et al.,  
300 2017) and in the Maritime Alps (Finsinger et al., 2021).

301 The shift to a *Picea-Abies* dominated forest was mostly concluded shortly after 9410 cal BP (onset of  
302 ZMN-3; 455.5-359.5 cm), as also observed at Prokoško Jezero (1670 m asl, Bosnia Herzegovina;  
303 Dörfler, 2013). Plant-macrofossils show that both *Pinus* and *Betula* were still intermixed in the spruce-  
304 fir dominated forest until around 8000 cal BP, when a low-density population of beech likely  
305 established, as suggested by the continuous pollen curve. At lower altitudes, vegetation was  
306 characterized by mixed oak forests with a higher share of *Corylus* between 9410 and 6000 cal BP.  
307 Around 7250 cal BP, *Ostrya*-type pollen increased indicating the expansion of *O. carpinifolia*, *C.*  
308 *orientalis*, or of both species.

309 This expansion was followed by an increase of *Fagus* and *C. betulus* pollen starting around 6400 cal BP  
310 (onset of ZMN-4a; 359.5-200.5 cm). Although the *Fagus* expansion at Zminje Jezero occurred  
311 approximately 2500-1000 years later than in pollen records from the Central Dinaric mountain area  
312 (Prokoško Jezero, Crveni Potok), it broadly coincided with the expansion of *Ostrya* and *Carpinus*  
313 *betulus*, as often seen in pollen records from the region (Prokoško Jezero, Lake Ohrid, Lake Prespa, and  
314 Crveni Potok; Dörfler, 2013; Finsinger et al., 2017; Panagiotopoulos et al., 2013; Wagner et al., 2009).

315 The admixture of *Fagus* in the *Picea-Abies* dominated forest involved a c. 2000-years long build-up  
316 phase (6400-4600 cal BP), as attested by rising pollen abundance, after which *Fagus* plant macrofossils  
317 and stomata regularly occur. As both *C. betulus* and *Fagus* are shade-tolerant and favoured by moist  
318 habitats on upland soils (Houston Durrant et al., 2016; Sikkema et al., 2016), an increase in humidity  
319 during the growing season may have occurred (Finsinger et al., 2017). In keeping with this, plant  
320 macrofossils of more light-demanding and better drought-resistant *Pinus* are absent in this zone, though  
321 stomata occasionally indicate their scattered presence. *Secale cereale* pollen first appear in this zone and  
322 occur mostly between 5800 and 3800 cal BP, suggesting its presence as a weed among other cereals  
323 during the Bronze Age (Behre, 1992). Some secondary anthropogenic-indicator pollen types (e.g. *P.*  
324 *lanceolata*-type, *Melampyrum*) increased as well. However, tree pollen abundance was highest in this  
325 time interval, suggesting that land-use pressure was minor. Similarly, a slightly lower tree-pollen

326 abundance between 3800 and 2100 cal BP suggests moderate presence of forest openings that are,  
327 however, not matched by rising abundance of anthropogenic-indicator pollen types.

328 By contrast, anthropogenic-indicator pollen document higher land-use pressures from Roman times  
329 onwards (*c.* after 2100 cal BP; ZMN-4b and ZMN-4c). *Secale cereale* pollen is mostly present around  
330 the Roman period and regularly occurs since the Ottoman Empire (ZMN-4c; 58.5-0 cm; 430 cal BP to  
331 present). Similarly, both cultural and secondary indicators are slightly higher during the Roman period  
332 and substantially rise since 430 cal BP in conjunction with a substantial tree-pollen decrease. Thus,  
333 pollen overall document two land-use phases characterized respectively by moderate land-use pressure  
334 (Roman period) and increasing and reaching highest land-use pressure since the Ottoman Empire. These  
335 two land-use phases were interrupted by a phase with minor land-use pressure during the Middle Ages  
336 (ZMN-4b; 200.5-58.5 cm; 2090-430 cal BP).

337 The frequent occurrence of *C. sativa* and *O. europaea* pollen since 1800 cal BP is consistent with a  
338 human-mediated spread of chestnut populations since Roman times (Conedera et al., 2004) and the  
339 cultivation of olive trees in Roman territories (Mercuri et al., 2013), including Montenegro, which was  
340 conquered by the Romans during the 2<sup>nd</sup> century BCE. Further, as in other records in the region (Dörfler,  
341 2013; Finsinger et al., 2017), pollen of *Juglans* first appear at *c.* 1500 cal BP and occur more regularly  
342 after 500 cal BP.

343 Although *Abies* pollen abundance was substantially lower after 2100 cal BP, a mixed *Abies-Picea-Fagus*  
344 was still present locally until 430 cal BP, as confirmed by stomata and plant macrofossils. However,  
345 lower pollen percentages and the absence of plant macrofossils of *Fagus* in conjunction with rare finds  
346 of *Abies* macrofossils since 430 cal BP suggest a major shift towards a *Picea*-dominated forest. Forest  
347 cover increased during the last century, perhaps due to the protection of the area since 1907 CE.

348

#### 349 *Charcoal-inferred fire history*

350 Regional fire activity, inferred based on microscopic charcoal (mCHAR), was highest over the past 2000  
351 years and was moderately high between 12,000 and 10,000 cal BP (Figs. 3-4). By contrast, landscape-  
352 scale biomass burning (MCHAR trend) was highest between 11,400 and 6000 cal BP, and overall  
353 decreasing from 6000 cal BP to the present (Figs. 3-4 and S4). The signal-to-noise index (Fig. S5) shows

354 that the macroscopic charcoal-accumulation rate (MCHAR) record is suitable for peak-detection  
355 analysis (Kelly et al., 2011). A total of 35 fire episodes were identified over the past 12,500 years, with  
356 a median fire-return interval (mFRI) of 260 years (FRI range: 79-790 years). Longest FRI values (>530  
357 years) occur before 8000 and after 2000 cal BP. Charred *A. alba* and *P. abies* needles document the  
358 occurrence of local fires in the conifer-dominated mountain ecosystem, mostly between 5600 and 500  
359 cal BP (Figs. 3-4).

360

#### 361 *Response of fire to variations in climate and land-use pressure*

362 Landscape-scale biomass burning (MCHAR trend) and regional fire activity (mCHAR) show different  
363 responses to climate and land-use pressure. Whereas biomass burning significantly increases in response  
364 to summer temperatures, fire activity significantly decreases (Fig. 5a; Table 2). Similarly, whereas  
365 biomass burning significantly decreases in response to land-use pressure, regional fire activity shows a  
366 hump-shaped response including an increase for the lower range of anthropogenic indicator pollen  
367 values (Fig. 5b).

368

#### 369 *Response of forest canopy species to variations in climate, fire and land use*

370 *Picea*'s responses to variations in climate, land use and biomass burning are not statistically significant  
371 (Fig. 5c-e). *Fagus* shows a significant negative response, and *Abies*, *F. excelsior*-type, *Ulmus*, and *Tilia*  
372 show significant positive responses to both summer temperature and biomass burning (Fig. 5c and e;  
373 Table 2). *Abies*, *F. excelsior*-type and *Ulmus* show significant negative responses to land-use pressure  
374 (Fig. 5d).

375

376

## 377 **Discussion**

378

#### 379 *Regional vs local land-use*

380 The investigation of long-term vegetation dynamics in relation to land-use pressure and disturbances  
381 requires a detailed understanding of land-use history. This is particularly important for southern Europe,

382 where forest clearances and agriculture often involved the use of fire. Moreover, land-use phases were  
383 often interspersed by periods of abandonment and secondary forest establishment as the result of the  
384 development and expansion of more permanent land-use practices, such as animal husbandry,  
385 ploughing, crop cultivation, and woodland management (Birks and Tinner, 2016). However, identifying  
386 the start and inferring the intensity of land-use pressure using palaeoecological records is challenging,  
387 particularly in regions where independent documentation by archaeological evidence is poor (Deza-  
388 Araujo et al., 2020), as is the case for the Durmitor.

389 Stomata and plant macrofossils indicate the local presence of taxa, mostly within some decametres  
390 around a site (Ammann et al., 2014; Birks, 2017). Thus, the continuous presence of plant macrofossils  
391 and stomata of trees (*Picea*, *Abies*, and *Fagus*) indicates that tree cover persisted in the surroundings of  
392 Zminje Jezero at least throughout the past 9000 years (Fig. 3). In keeping with this, high arboreal pollen  
393 (generally >80%) and AP/NAP ratio values (generally >4; Fig. 4) qualitatively indicate the presence of  
394 closed forest (Deza-Araujo et al., 2020). A persistently high forest cover was also found in the internal  
395 Dinaric Alps at Crveni potok (Finsinger et al., 2017), but contrasts with results from Prokoško Jezero  
396 (Dörfler, 2013) where arboreal pollen values decreased to 50% during the past 2000 years. This  
397 contrasting evidence would support the hypothesis that some of the forested regions in the interior range  
398 of the Dinaric Alps have been left relatively intact until the present (Nagel et al., 2017).

399 Conversely, if land-use pressure was regionally heterogeneous, reconstructions based on pollen may not  
400 necessarily covary with those based on plant macrofossils and stomata. Pollen from the wider region  
401 may reach small lakes such as Zminje Jezero even if their relevant pollen source area is relatively small  
402 (some kilometres at most; Sugita, 1994). Indeed, despite persistence of closed forest in the surroundings  
403 of Zminje Jezero, anthropogenic indicators unambiguously document increasing land-use pressure in  
404 the region since the Roman period. Cultural indicators, secondary indicators, as well as OJC-index  
405 values rise around 2000 cal BP, when the region currently belonging to Montenegro was conquered by  
406 the Romans (2<sup>nd</sup> century BCE). At that time, land-use pressure probably reached the Durmitor, as  
407 attested by the construction of roads, including the one connecting Nikšić and Komini. The further rise  
408 of anthropogenic indicators around 500 cal BP (15<sup>th</sup> century CE) is coherent with the first local  
409 settlements in the Durmitor. For instance, the so-called stećci sites (medieval tombstones and graveyards

410 from the 13th-17th century; Erdeljan, 2018) attest to the local presence of settlements on the Durmitor  
411 during the Middle Ages and the Ottoman Empire, and churches and monasteries from the 15<sup>th</sup>-19<sup>th</sup>  
412 century CE document the presence of local settlements during the Ottoman Empire. Thus, these results  
413 add a finer-scale scale documentation of past land-use pressure in comparison to modelled  
414 anthropogenic deforestation patterns (Kaplan et al., 2009), which suggest that relatively low forest  
415 clearance occurred during the past 3000 years and that the region was not heavily exploited until the  
416 Middle Ages.

417 Similarly, regional heterogeneity of land-use pressure may also have occurred in pre-Roman times.  
418 Anthropogenic indicators do not show distinct signs of Bronze and Iron Age land use, despite the  
419 probable connection between coastal and inland settlements (Bulatović et al., 2020) and Illyrian and  
420 Celtic Iron Age settlements (Cozzolino et al., 2020). However, *Secale cereale* pollen finds at Zminje  
421 Jezero suggest its presence as a weed among other cereals (Behre, 1992) since about 6000 cal BP, as  
422 also observed in archaeobotanical samples from Neolithic sites in the mountain hinterland (e.g. at  
423 Okolište, Bosnia-Herzegovina; de Vareilles et al., 2022). While this is about 1000 years later than the  
424 age of Neolithic settlements on the shores of the lowland Lake Ohrid (Hafner et al., 2021) and 2000  
425 years later than the earliest Starčevo and Butmir culture Neolithic settlements in the inland of Bosnia-  
426 Herzegovina (Vander Linden et al., 2014), the *Secale* pollen finds support the view that Neolithic  
427 novelties, albeit with a certain time lag, spread in the hinterland of Montenegro (Borić et al., 2019).  
428 Although Neolithic sites have not yet been found in the Durmitor area, there is evidence of Neolithic  
429 occupation sites at 20-35 km distance (Odmut rock shelter; Borić et al., 2019), suggesting that inland  
430 mountain areas may be archeologically under surveyed (Vander Linden et al., 2014). Thus, the  
431 palaeoecological records agree with the hypothesis that the mountain hinterland remained an agricultural  
432 frontier zone for much of the Neolithic (Forenbaher and Miracle, 2005) and possibly even until the  
433 Roman Age.

434

#### 435 *Regional vs local fires*

436 We found contrasting responses of fire to climate and land-use pressure over the past 8500 years (Fig.  
437 5a-b), pointing to different drivers of fire dynamics at landscape and regional scales. Landscape-scale



438 biomass burning, as inferred based on MCHAR, responded positively and significantly to summer  
439 temperature, as expected based on the strong influence (summer) temperature has on fire activity (Jain  
440 et al., 2022; Westerling et al., 2006) and as often observed with sedimentary charcoal records (Daniau  
441 et al., 2012; Power et al., 2008). As fuel was not a limiting factor in the moist and productive mountain  
442 forests, higher mean July temperatures perhaps coupled with prolonged dry seasons (Holocene Thermal  
443 Maximum; Samartin et al., 2017) likely improved fuel flammability (Pausas and Paula, 2012). By  
444 contrast, surprisingly regional fire activity, as inferred based on mCHAR, responded negatively to  
445 summer temperature. A disconnect between fire-history reconstructions can arise both when comparing  
446 paleofire records from sites located near each other (Finsinger et al., 2018; Gavin et al., 2006), as well  
447 as different paleofire proxies from the same site (Finsinger et al., 2017) if fire occurrence was not  
448 spatially homogeneous. While charcoal records from additional sites may be necessary to better  
449 constrain the spatial heterogeneity of fire occurrence in the region, it is likely that mCHAR captures fire  
450 activity at greater distance from the site than MCHAR (Finsinger et al., 2017). This interpretation is  
451 supported by the positive response of fire activity (mCHAR) to land-use pressure (Fig. 5b), which likely  
452 occurred in a wider region, including lower-elevation areas where submediterranean deciduous  
453 woodlands occur. The slightly earlier rise of regional fire activity, which started around the Iron Age  
454 (Fig. 4), may be connected to land use in the region that remained undetected by the anthropogenic  
455 indicators from Zminje Jezero. Biomass burning, instead, responded negatively to land-use pressure  
456 (Fig. 5c). It is unlikely that this relationship is the result of feedbacks arising from human-induced  
457 deforestation (e.g. lower fuel availability, fragmentation; Bowman et al., 2011) because biomass burning  
458 decreased predominantly at the Neolithic/Bronze Age transition, when land-use pressure was low. The  
459 overall decreasing biomass burning suggests that humans did not extensively use fire in the Durmitor.  
460 While this overall contrasts with paleofire evidence from other southern European mountains (e.g. the  
461 Apennines; Morales-Molino et al., 2021), the lower biomass burning rates in recent millennia fit with  
462 the notion that fires are currently rare in the Dinaric Alps (Nagel et al., 2017). However, charred *Picea*  
463 and *Abies* needles unambiguously document the occurrence of fires in the mixed spruce-fir-beech  
464 mountain forests of the Durmitor.

465

466 *Species-response curves and legacies of past environmental changes*

467 Species-response curves are useful to assess factors that have driven long-term vegetation dynamics,  
468 thereby unfolding legacies of past environmental changes (Carter et al., 2020; Colombaroli et al., 2010;  
469 Morales-Molino et al., 2021). We found similar species responses to summer temperature and biomass  
470 burning (Fig. 5c and 5e). Given the strong and positive relationship among these two predictors (Fig.  
471 5a), it is difficult to disentangle their roles separately. However, as (summer) temperature currently has  
472 a strong influence on fire activity (Jain et al., 2022; Westerling et al., 2006), the response curves may  
473 yield valuable insights into the long-term fire ecology of the analysed species (*P. abies*, *F. sylvatica*, *A.*  
474 *alba*, *Ulmus*, *Fraxinus excelsior*, and *Tilia*), which are widespread and often dominant in European  
475 mountain forests. The right-skewed responses of *Fagus* to biomass burning and temperature (Fig. 5c  
476 and 5e) suggest that it was favoured under low biomass-burning and cooler conditions. Overall, these  
477 response curves support the notion that *Fagus* became more competitive in mixed fir-spruce forests with  
478 the onset of cooler and moister summers and fire disturbance, as observed in the ecologically similar  
479 Northern Apennines (Morales-Molino et al., 2021). Indeed, *Fagus* requires moist summers for its  
480 germination (Giesecke et al., 2017) and is more sensitive than *Abies* to drought due to its shallow root  
481 system (Cheddadi et al., 2016; Tinner et al., 2013). *Fagus* is a shade-tolerant species and its regeneration  
482 can establish and grow with low light levels under the forest canopy for years before rapidly expanding  
483 and achieving dominance when light conditions improve after disturbances (Gardner and Willis, 1999).  
484 Thus, fire occurrence after 5000 cal BP may have promoted the recruitment of *Fagus* and its admixture  
485 into the mixed *Picea-Abies* forest. This matches with evidence suggesting that during the initial phase  
486 of stand establishment, *Fagus* is favoured by mixed-severity fire and intermediate human disturbances  
487 (Bradshaw and Lindbladh, 2005; Giesecke et al., 2007; Tinner and Lotter, 2006) thanks to its capacity  
488 to occupy newly exposed soils and canopy openings (Carter et al., 2018) more rapidly than its two main  
489 competitors (*Abies* and *Picea*). Despite lacking ecological adaptations to fire and being identified as a  
490 fire-sensitive species that is affected by higher biomass burning and temperatures (Carter et al., 2018;  
491 Morales-Molino et al., 2021; Tinner et al., 2000), *Fagus* can be advantaged by fire incidence as long as  
492 it is not excessive (Maringer et al., 2020). In keeping with this, the change-point analysis shows a  
493 decrease in biomass burning during the establishment of *Fagus* (Fig. 4), corroborating the view that

494 beech establishment occurs in conjunction with a shift to a lower fire activity (Bobek et al., 2019;  
495 Feurdean et al., 2017).

496 Further, similar to observations from the ecologically similar Northern Apennines (Morales-Molino et  
497 al., 2021), we found that *Abies* and disturbance-sensitive deciduous taxa such as *Ulmus*, *F. excelsior*,  
498 and *Tilia* were promoted by warmer-than-present summer temperatures (up to ca. 2.5°C; Fig. 5c).  
499 However, our findings also suggest that these taxa responded positively to increasing biomass burning  
500 (Fig. 5e). This contrasts with several records showing that both high-severity infrequent fires and low-  
501 severity frequent fires led to their rapid decline and to local extinctions in recent millennia (Morales-  
502 Molino et al., 2021; Tinner et al., 2000, 2013). On the one hand, the striking positive response of these  
503 disturbance-sensitive taxa to biomass burning could simply arise from the positive relationship between  
504 landscape-scale biomass burning and summer temperatures (Fig. 5a). On the other hand, the significant  
505 negative response of *Abies*, *Ulmus*, and *F. excelsior* to land-use pressure (Fig. 5d) as well as to regional  
506 fire activity (Fig. S6 and Table S2) indicates that their populations declined even in the wider region  
507 due to land use, probably with the use of fires. Indeed, the decline of *Abies* coincides with the rise in  
508 land-use pressure (pasturing and small-scale cultivation) and of regional fire activity since 2000 cal BP.  
509 Conversely, the local persistence of *Abies* in the surroundings of Zminje Jezero until present-day  
510 suggests that land-use pressure was not widespread across the montane forest belt, leaving relatively  
511 undisturbed patches of forests in a landscape mosaic (Fig. 1). Indeed, *Abies* can be resistant to fire  
512 disturbances when fires are rare (Henne et al., 2013) and its populations strongly reduce under excessive  
513 human-induced fire activity (Feurdean and Willis, 2008; Finsinger et al., 2021; Tinner et al., 2013),  
514 which was not the case around Zminje Jezero but may have been the case to the East of Žabljak and  
515 elsewhere in the upper-montane zone of the central Dinaric Alps.

516 *Picea*'s response curves suggest that neither temperature nor biomass burning and land use significantly  
517 influenced its populations (Fig. 5c-e). Indeed, *Picea* remained relatively stable even under higher-than-  
518 present summer temperatures (Fig. 4), as also observed with palaeoecological records from the  
519 Carpathians (Carter et al., 2018). Its insensitivity to summer temperatures may simply imply that past  
520 climatic conditions did not exceed the species' tolerance range. Moreover, the lack of a significant  
521 response to biomass burning agrees with evidence showing that *Picea* can persist thanks to vigorous

522 post-fire regeneration from wind-dispersed seeds (Bobek et al., 2018) even with median FRI as low as  
523 260 years as at Zminje Jezero (Brown and Giesecke, 2014; Feurdean et al., 2017; Finsinger et al., 2018).  
524 Although fires may have been fuelled by *Picea*'s traits that promote fire spread (high resin content,  
525 flammable litter, and a ladder-canopy structure), its well-established populations persisted at Zminje  
526 Jezero despite fire occurrence in the mixed spruce-fir-beech forests.

527 The species-response curves may also be useful to explain the relatively recent shift from a *Picea-Abies-*  
528 *Fagus* mixed forest to one dominated by *Picea* that occurred in the surroundings of Zminje Jezero  
529 around 430 cal BP (15<sup>th</sup> century CE; Fig. 4). Fir populations, which are more sensitive to land-use  
530 pressure than *Picea* and *Fagus* (Fig. 5d), could have been reduced due to the establishment of the local  
531 pastoral katun society (c. 1477 CE; Turkish Land Registry). Around 1400 CE, the Balkan Peninsula was  
532 under the influence of the Republic of Venice and from 1496 CE Montenegro fell under the domain of  
533 the Ottoman Empire (Rastoder, 2003). In the mid-16<sup>th</sup> century, local land-owners started seizing  
534 monastery estates and hunting grounds in the lower parts of Montenegro. This resulted in the retreat of  
535 considerable parts of the population in mountain areas (Rastoder, 2003), including the Durmitor, whose  
536 forests probably started to be more intensively exploited. An alternative, or additional explanation could  
537 involve the lower sensitivity of spruce to declining winter temperatures (Cheddadi et al., 2016). In this  
538 scenario, the development of a *Picea*-dominated forest could be linked to the onset of cooler conditions  
539 during the Little Ice Age (1450-1850 CE). Indeed, due to its shallow root system *Picea* is more  
540 vulnerable to drought than *Abies* (Henne et al., 2011), and spruce-dominated stands in the Dinaric Alps  
541 mostly occur in habitats with significantly colder and moister microclimates (Nagel et al., 2017).

542

543

## 544 **Conclusions**

545 Vegetation of forest ecosystems is often the result of complex interactions between vegetation dynamics  
546 and variations in climate, disturbances (e.g. fire), and human activities. However, understanding the  
547 ecological processes underlying these interactions is challenging as processes often play out over long  
548 time scales (centuries, millennia), particularly for forests where long-lived species such as trees are

549 involved. The long-term perspective of palaeoecological records permits to study ecosystem responses  
550 under substantially different environmental conditions than the present ones.

551 In this study, we sought to investigate the Holocene vegetation and fire history in the upper-montane  
552 zone of the central Dinaric Alps, where fires are currently rare and forests mainly include beech-  
553 dominated stands and mixed forests dominated by varying amounts of beech, fir, and occasionally  
554 spruce (Nagel et al., 2017). While records from different elevations may be necessary to better support  
555 our interpretation, our results support the view that at least some parts of the central Dinaric Alps  
556 remained an agricultural frontier zone during the Neolithic (Forenbaher and Miracle, 2005) and that  
557 land-use pressure was low until the Middle Ages (Kaplan et al., 2009).

558 Response curves indicate that fires in the upper-montane zone were mostly driven by variations in  
559 climatic conditions, as expected based on the strong influence (summer) temperature has on fire activity  
560 today (Jain et al., 2022; Westerling et al., 2006). *Picea* was insensitive to variations in summer  
561 temperature, biomass burning and human impact, which supports the view that spruce forests may not  
562 be significantly impacted by fire (Carter et al., 2018). In contrast, *Abies* and other disturbance-sensitive  
563 deciduous trees (*F. excelsior*-type, *Tilia*, and *Ulmus*) show a significant positive relationship with  
564 summer temperatures and a significant negative relationship with land-use pressure. This supports the  
565 notion that these tree species may be well-adapted to warmer-than-present summer temperatures and  
566 that their populations declined in recent millennia due to land-use activities (Morales-Molino et al.,  
567 2020; Tinner et al., 2000, 2013). Conversely, *Fagus* emerges as a species sensitive to summer  
568 temperatures whose expansion in the *Picea-Abies* dominated forest was enhanced by the onset of cooler  
569 and possibly moister climatic conditions as well as by fire disturbances.

570

## 571 **Acknowledgements**

572 We are grateful to the Environment protection Agency of Montenegro for granting permission to collect  
573 sediments from the lake, to S. Rouland, S. Canal, and A. Haliuc for help during the lab work, to I.  
574 Figueral for help with identification of plant-macrofossils and wood, and to the *Laboratoire Souterrain*  
575 *de Modane (LSM)* facilities for the gamma spectrometry measurements. We are further thankful to O.  
576 Heiri (University of Basel, Switzerland) for funding some of the radiocarbon dates. MB, KH and ZLS

577 were supported by the European Union and the State of Hungary, co-financed by the European Regional  
578 Development Fund (GINOP-2.3.2.-15-2016-00009 'ICER').

579

580


#### 581 **Funding**

582 This research was funded by ISEM (University of Montpellier, CNRS, IRD, EPHE) and by the  
583 University of Padova [grant number DOR1957359/19].


584

#### 585 **Data**

586 The palaeoecological data (pollen, spores, stomata, plant macrofossils, charcoal, and XRF records) will  
587 be uploaded to the Neotoma database (DOI: 10.21233/XYZZ-XYZZ) through the European Pollen  
588 Database.




590  
 591 **Fig. 1** a-c) Location of the study site at decreasing spatial scales. b-c) Map showing the distribution of  
 592 land cover types (CORINE Land Cover 2018) in the Durmitor National Park and surroundings. d)  
 593 bathymetric map of Zminje Jezero with contour lines at 1-m intervals and coring locations (red circles).  
 594



595  
 596 **Fig. 2** Depth-age model for the Zminje Jezero sediment core. The active control points used to constrain  
 597 the model (Table 1) are based on  $^{14}\text{C}$ ,  $^{210}\text{Pb}$ , and  $^{137}\text{Cs}$  dates as well as on a pollen-inferred age estimate  
 598 (onset of the Holocene). Passive control points based on local maxima in the XRF-Pb record (McConnell  
 599 et al., 2018; Renberg et al., 2001) are shown in green (see also Fig. S2). Grey horizontal bands indicate  
 600 turbidite layers that were excluded from the model. Lithological and XRF geochemical profiles of the  
 601 sediment core are shown on the right. The inset shows the distribution of sediment deposition time  
 602 within the sequence (median = 21 years  $\text{cm}^{-1}$ )  
 603








616

617 **Fig. 4** Comparison of the main proxies for vegetation dynamics, fire and land-use history from Zminje  
 618 Jezero, and chironomid-inferred July-air temperature ( $T_{July}$ ) anomalies from Lago Verdarolo (Samartin  
 619 et al., 2017). Microscopic charcoal (mCHAR, thick dark-grey line) documenting regional fire activity  
 620 and macroscopic charcoal (MCHAR, black filled area) documenting landscape-scale biomass burning  
 621 (red continuous line), local fire episodes (black crosses), periods of biomass burning as determined by  
 622 the change-point analysis (grey-shaded areas), and charred *Picea* and *Abies* needles (red diamonds).  
 623 Temporal changes of dominant forest canopy taxa and of anthropogenic indicator pollen types (filled  
 624 polygons: pollen percentages; black vertical bars: plant macrofossils concentrations; grey circles:  
 625 stomata). Grey vertical shaded areas represent pollen zone boundaries (see Fig. 3).

626



627  
 628 **Fig. 5** Generalized Additive Models showing (a-b) the responses of landscape-scale biomass burning  
 629 and regional fire activity to climate and land-use pressure, and (c-e) the responses of dominant forest  
 630 canopy taxa to climate, land-use pressure, and biomass burning.

**Table 1** Control points used to establish the depth-age model for the ZMN-19 sediment core from Zminje jezero (Durmitor Massif, Montenegro). For a detailed description of the terrestrial plant macrofossils used for radiocarbon dating see Table S1.

Depth (cm)	Lab-code	Material	Dry weight (mg)	Sample Thickness (cm)	Age ( $^{14}\text{C}$ yr BP) (cal yr BP)*
0	$^{210}\text{Pb}$	Bulk sediment	/	0	$-69 \pm 5^*$
0.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-65 \pm 1^*$
1.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-62 \pm 1^*$
2.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-58 \pm 1^*$
3.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-54 \pm 1^*$
4.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-51 \pm 1^*$
5.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-47 \pm 2^*$
6.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-44 \pm 2^*$
7.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-40 \pm 2^*$
8.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-36 \pm 3^*$
9.5	$^{137}\text{Cs}$ peak	Bulk sediment	/	0	$-36 \pm 1^*$
9.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-33 \pm 3^*$
10.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-29 \pm 3^*$
12.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-22 \pm 4^*$
13.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-19 \pm 4^*$
14.5	NWT $^{137}\text{Cs}$	Bulk sediment	/	0	$-15 \pm 5^*$
14.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-13 \pm 1^*$
15.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-11 \pm 5^*$
16.5	$^{137}\text{Cs}$	Bulk sediment	/	0	$-8 \pm 5^*$
16.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-5 \pm 5^*$
17.5	$^{210}\text{Pb}$	Bulk sediment	/	0	$-4 \pm 6^*$
71.5	DeA-33892	Terrestrial plant macrofossils	2.25	1	$565 \pm 19$
118.5	DeA-23961	Terrestrial plant macrofossils	20	1	$1054 \pm 24$
126.5	DeA-32618	Terrestrial plant macrofossils	16.8	1	$1427 \pm 20$
180.5	DeA-31441	Terrestrial plant macrofossils	9.8	1	$1844 \pm 23$
219.5	DeA-33893	Terrestrial plant macrofossils	3.5	1	$2335 \pm 55$
250.5	DeA-31442	Terrestrial plant macrofossils	8	1	$2755 \pm 22$
290.5	DeA-31091	Terrestrial plant macrofossils	8.5	1	$3762 \pm 22$
307.5	DeA-23962	Terrestrial plant macrofossils	194	1	$4108 \pm 30$
324.5	DeA-31093	Terrestrial plant macrofossils	17.8	1	$4532 \pm 25$
347.5	DeA-33890	Terrestrial plant macrofossils	1.18	1	$5212 \pm 51$
357.5	DeA-23963	Terrestrial plant macrofossils	266.5	1	$3980 \pm 30$
369.5	DeA-31440	Terrestrial plant macrofossils	17.6	1	$6060 \pm 28$
384.5	DeA-23964	Terrestrial plant macrofossils	170.2	1	$6552 \pm 37$
408.5	DeA-33891	Terrestrial plant macrofossils	1.6	1	$7449 \pm 60$
439.5	DeA-23967	Terrestrial plant macrofossils	148.5	1	$7908 \pm 39$
476	DeA-23965	Terrestrial plant macrofossils	532.3	0.5	$9139 \pm 44$
566.5	/	Pollen inferred (onset Holocene)	/	1	$11500 \pm 250^*$
581	Poz-10170	Terrestrial plant macrofossils	0.4	2	$10170 \pm 60$
594	Poz-143126	Terrestrial plant macrofossils	3	2	$10260 \pm 60$
601.5	Poz-143573	Terrestrial plant macrofossils	0.08	3	$9190 \pm 120$

**Table 2** Ecosystem responses to variations in climate, fire and land use at Zminje jezero (Durmitor Massif, Montenegro), as determined by Generalized Additive Models (GAMs) based on the pollen percentages and charcoal-accumulation rate records (this study) as well as chironomid-inferred July temperatures from Lago Verdarolo (Samartin et al., 2017). Effective degrees of freedom indicate the complexity of the smooth (a value = 1 indicates a straight line); significance of smooth terms: \* =  $P \leq 0.05$ , \*\* =  $P \leq 0.01$ , \*\*\* =  $P \leq 0.001$ .

Independent variable	Response variable	Effective degrees of freedom	Reference degrees of freedom	F-value	Deviance explained (%)
Climate (chironomid-inferred July temperature anomalies)	Biomass burning***	2.54	3.08	11.2	52.2
	Regional fire**	1	1	12.18	28.3
	<i>P. abies</i>	1	1	2.2	4.07
	<i>A. alba</i> ***	1	1	14.31	22.4
	<i>F. sylvatica</i> ***	3.07	3.88	11.47	57
	<i>F. excelsior-type</i> ***	1	1	21.72	28.9
	<i>Tilia</i> ***	1.88	2.35	10.02	31.1
	<i>Ulmus</i> ***	1	1	33.14	37.2
Biomass burning (long-term trend of MCHAR)	<i>P. abies</i>	1.0	1.0	3.88	7.1
	<i>A. alba</i> ***	2.32	2.78	10.65	39.1
	<i>F. sylvatica</i> ***	4.33	5.2	5.61	53
	<i>F. excelsior-type</i> ***	1.83	2.21	0.2	32.4
	<i>Tilia</i> ***	1.79	2.16	12.26	37
	<i>Ulmus</i> ***	1.65	1.99	23	46.5
Land-use pressure (anthropogenic indicator pollen)	Biomass burning**	1.89	2.37	6.55	26.3
	Regional fire***	3.32	4.07	7.13	38.4
	<i>P. abies</i>	1	1	0.11	0.21
	<i>A. alba</i> ***	2.26	2.83	11.14	41.7
	<i>F. sylvatica</i>	1.34	1.61	0.12	1.6
	<i>F. excelsior-type</i> **	1.08	1.16	10.37	25.2
	<i>Tilia</i>	1.48	1.82	2.59	13.7
	<i>Ulmus</i> ***	1.03	1.05	12.8	30.7

634 **References**

- 635 Adolf C, Wunderle S, Colombaroli D, Weber H, Gobet E, Heiri O, et al. (2018) The sedimentary and  
636 remote-sensing reflection of biomass burning in Europe. *Global Ecology and Biogeography* 27: 199–  
637 212: doi:10.1111/geb.12682.
- 638 Ammann B, Knaap WO van der, Lang G, Gaillard M-J, Kaltenrieder P, Rösch M, et al. (2014) The  
639 potential of stomata analysis in conifers to estimate presence of conifer trees: examples from the Alps.  
640 *Vegetation History and Archaeobotany* 23(3): 249–264: doi:10.1007/s00334-014-0431-9.
- 641 Annys K, Frankl A, Spalević V, Čurović M, Borota D and Nyssen J (2014) Geomorphology of the  
642 Durmitor Mountains and surrounding plateau Jezerska Površ (Montenegro). *Journal of Maps* 10(4):  
643 600–611: doi:10.1080/17445647.2014.909338.
- 644 Behre K-E (1981) The interpretation of anthropogenic indicators in pollen diagrams. *Pollen et Spores*  
645 23: 225–245.
- 646 Behre K-E (1992) The history of rye cultivation in Europe. *Vegetation History and Archaeobotany*  
647 1(3). Available at: <http://link.springer.com/10.1007/BF00191554>: doi:10.1007/BF00191554.
- 648 Bennett KD (1996) Determination of the number of zones in a biostratigraphical sequence. *New*  
649 *Phytologist* 132: 155–170.
- 650 Bennett KD (2008) Psimpoll v. 4.26. . Available at: <http://chrono.qub.ac.uk/psimpoll/psimpoll.html>.
- 651 Beug H-J (2004) *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*.  
652 München: Verlag Dr. Friedrich Pfeil.
- 653 Birks H (2017) *Plant Macrofossil Introduction*. : doi:10.1016/B978-0-12-409548-9.10499-3.
- 654 Birks HJB and Tinner W (2016) Past forests of Europe. In: San-Miguel-Ayanz J, de Rigo D, Caudullo  
655 G, Houston Durrant T and Mauri A (eds) *European Atlas of Forest Tree Species*. Luxembourg:  
656 Publications Office of the European Union, 36–39.
- 657 Blaauw M and Christen JA (2011) Flexible Paleoclimate Age-Depth Models Using an Autoregressive  
658 Gamma Process. *Bayesian Analysis* 6: 457–474: doi:10.1214/11-BA618.
- 659 Bobek P, Šamonil P and Jamrichová E (2018) Biotic controls on Holocene fire frequency in a  
660 temperate mountain forest, Czech Republic. *Journal of Quaternary Science* 33(8): 892–904:  
661 doi:10.1002/jqs.3067.
- 662 Bobek P, Svobodová-Svitavská H, Pokorný P, Šamonil P, Kuneš P, Kozáková R, et al. (2019)  
663 Divergent fire history trajectories in Central European temperate forests revealed a pronounced  
664 influence of broadleaved trees on fire dynamics. *Quaternary Science Reviews* 222: 105865:  
665 doi:10.1016/j.quascirev.2019.105865.
- 666 Boncina A (2011) History, current status and future prospects of uneven-aged forest management in  
667 the Dinaric region: an overview. *Forestry: An International Journal of Forest Research* 84(5): 467–  
668 478: doi:10.1093/forestry/cpr023.
- 669 Borić D, Borovinić N, Đuričić L, Bulatović J, Gerometta K, Filipović D, et al. (2019) Spearheading  
670 into the Neolithic: Last Foragers and First Farmers in the Dinaric Alps of Montenegro. *European*  
671 *Journal of Archaeology* 22(4): 470–498: doi:10.1017/eea.2019.14.

- 672 Bowman DMJS, Balch J, Artaxo P, Bond WJ, Cochrane MA, D'Antonio CM, et al. (2011) The human  
673 dimension of fire regimes on Earth. *Journal of Biogeography* 38(12): 2223–2236: doi:10.1111/j.1365-  
674 2699.2011.02595.x.
- 675 Bradshaw RHW and Lindbladh M (2005) Regional spread and stand-scale establishment of *Fagus*  
676 *sylvatica* and *Picea abies* in Scandinavia. *Ecology* 86: 1679–1686.
- 677 Brown KJ and Giesecke T (2014) Holocene fire disturbance in the boreal forest of central Sweden.  
678 *Boreas* 43: 639–651: doi:10.1111/bor.12056.
- 679 Bruel R and Sabatier P (2020) serac: an R package for ShortlivEd RADionuclide chronology of recent  
680 sediment cores. *Journal of Environmental Radioactivity* 225: 106449:  
681 doi:10.1016/j.jenvrad.2020.106449.
- 682 Bui B (1975) Aperçu sur la végétation des montagnes de la Yougoslavie centrale. *Méditerranée* 20(1):  
683 81–87: doi:10.3406/medit.1975.1589.
- 684 Bulatović A, Gori M and Vander Linden M (2020) Radiocarbon dating the 3rd millennium BC in the  
685 central Balkans: a re-examination of the early Bronze Age sequence. *Radiocarbon* 62(5): 1163–1191:  
686 doi:10.1017/RDC.2020.61.
- 687 Caf N, Sabatier P, Šmuc A, Ogrinc N, Dolenc M, Rapuc W, et al. (2022) Multi-proxy reconstruction  
688 of the Holocene vegetation and land use dynamics in the Julian Alps, north-west Slovenia. *Journal of*  
689 *Quaternary Science* jqs.3461: doi:10.1002/jqs.3461.
- 690 Cagliero E, Morresi D, Paradis L, Curović M, Spalević V, Marchi N, et al. (2022) Legacies of past  
691 human activities on one of the largest old-growth forests in south-east European mountains.  
692 *Vegetation History and Archaeobotany* 31: 415–430: doi:10.1007/s00334-021-00862-x.
- 693 Carter VA, Bobek P, Moravcová A, Šolcová A, Chiverrell RC, Clear JL, et al. (2020) The role of  
694 climate-fuel feedbacks on Holocene biomass burning in upper-montane Carpathian forests. *Global and*  
695 *Planetary Change* 103264: doi:10.1016/j.gloplacha.2020.103264.
- 696 Carter VA, Moravcová A, Chiverrell RC, Clear JL, Finsinger W, Dreslerová D, et al. (2018)  
697 Holocene-scale fire dynamics of central European temperate spruce-beech forests. *Quaternary Science*  
698 *Reviews* 191: 15–30: doi:10.1016/j.quascirev.2018.05.001.
- 699 Cheddadi R, Araújo MB, Maiorano L, Edwards M, Guisan A, Carré M, et al. (2016) Temperature  
700 Range Shifts for Three European Tree Species over the Last 10,000 Years. *Frontiers in Plant Science*  
701 7: 1581: doi:10.3389/fpls.2016.01581.
- 702 Colombaroli D, Henne PD, Kaltenrieder P, Gobet E and Tinner W (2010) Species responses to fire,  
703 climate and human impact at tree line in the Alps as evidenced by palaeo-environmental records and a  
704 dynamic simulation model. *Journal of Ecology* 98(6): 1346–1357: doi:10.1111/j.1365-  
705 2745.2010.01723.x.
- 706 Conedera M, Krebs P, Tinner W, Pradella M and Torriani D (2004) The cultivation of *Castanea sativa*  
707 (Mill.) in Europe, from its origin to its diffusion on a continental scale. *Vegetation History and*  
708 *Archaeobotany* 13: 161–179.
- 709 Cozzolino M, Baković M, Borovinić N, Galli G, Gentile V, Jabučanin M, et al. (2020) The  
710 Contribution of Geophysics to the Knowledge of the Hidden Archaeological Heritage of Montenegro.  
711 *Geosciences*. Multidisciplinary Digital Publishing Institute 10(5): 187:  
712 doi:10.3390/geosciences10050187.

- 713 Daniau A-L, Bartlein PJ, Harrison SP, Prentice IC, Brewer S, Friedlingstein P, et al. (2012)  
714 Predictability of biomass burning in response to climate changes. *Global Biogeochemical Cycles* 26:  
715 GB4007: doi:10.1029/2011GB004249.
- 716 Davis OK (1987) Spores of the dung fungus *Sporormiella*: increased abundance in historic sediments  
717 and before Pleistocene megafaunal extinction. *Quaternary Research* 28: 290–294.
- 718 de Vareilles A, Filipović D, Obradović D and Vander Linden M (2022) Along the Rivers and into the  
719 Plain: Early Crop Diversity in the Central and Western Balkans and Its Relationship with  
720 Environmental and Cultural Variables. *Quaternary* 5(1): 6: doi:10.3390/quat5010006.
- 721 Deza-Araujo M, Morales-Molino C, Tinner W, Henne PD, Heitz C, Pezzatti GB, et al. (2020) A  
722 critical assessment of human-impact indices based on anthropogenic pollen indicators. *Quaternary*  
723 *Science Reviews* 236: 106291: doi:10.1016/j.quascirev.2020.106291.
- 724 Dörfler W (2013) Prokoško Jezero: An environmental record from a subalpine lake in Bosnia-  
725 Herzegovina. In: Müller J, Rassmann K and Hofmann R (eds) *Okolište I – Untersuchungen einer*  
726 *spätneolithischen Siedlungskammer in Zentralbosnien*. Bonn: Habelt Verlag, 311–340.
- 727 Erdeljan J (2018) Cultural-historical heritage - a key to sustainable development of mountain areas. In:  
728 Jaćimović M (ed) *Sustainable development of mountain areas - experiences, challenges and*  
729 *perspectives*. Podgorica: Montenegrin Academy of Sciences and Arts, 185–189. Available at:  
730 <https://canupub.me/r1vr>.
- 731 Feurdean A, Florescu G, Vannièrè B, Tanțău I, O'Hara RB, Pfeiffer M, et al. (2017) Fire has been an  
732 important driver of forest dynamics in the Carpathian Mountains during the Holocene. *Forest Ecology*  
733 *and Management* 389: 15–26: doi:10.1016/j.foreco.2016.11.046.
- 734 Feurdean A and Willis KJ (2008) Long-term variability of *Abies alba* in NW Romania: implications  
735 for its conservation management. *Diversity and Distributions* 14: 1004–1017.
- 736 Feurdean AN, Willis KJ and Astaloş C (2009) Legacy of the past land-use changes and management  
737 on the 'natural' upland forest composition in the Apuseni Natural Park, Romania. *The Holocene* 19(6):  
738 967–981: doi:10.1177/0959683609337358.
- 739 Finsinger W and Bonnici I (2022) tapas: an R package to perform trend and peaks analysis. Zenodo.  
740 Available at: <https://zenodo.org/record/6344463>: doi:10.5281/zenodo.6344463.
- 741 Finsinger W, Fevre J, Orbán I, Pál I, Vincze I, Hubay K, et al. (2018) Holocene fire-regime changes  
742 near the treeline in the Retezat Mts. (Southern Carpathians, Romania). *Quaternary International* 477:  
743 94–105: doi:10.1016/j.quaint.2016.04.029.
- 744 Finsinger W, Kelly R, Fevre J and Magyari EK (2014) A guide to screening charcoal peaks in  
745 macrocharcoal-area records for fire-episodes reconstructions. *The Holocene* 24: 1002–1008:  
746 doi:10.1177/0959683614534737.
- 747 Finsinger W, Morales-Molino C, Gałka M, Valsecchi V, Bojovic S and Tinner W (2017) Holocene  
748 vegetation and fire dynamics at Crveni Potok, a small mire in the Dinaric Alps (Tara National Park,  
749 Serbia). *Quaternary Science Reviews* 167: 63–77: doi:10.1016/j.quascirev.2017.04.032.
- 750 Finsinger W and Tinner W (2005) Minimum count sums for charcoal-concentration estimates in  
751 pollen slides: reliability and potential errors. *The Holocene* 15: 293–297.
- 752 Finsinger W, Vanel Q, Ribolini A and Tinner W (2021) Early to late Holocene vegetation and fire  
753 dynamics at the treeline in the Maritime Alps. *Vegetation History and Archaeobotany* 30: 507–524:  
754 doi:10.1007/s00334-020-00795-x.



- 755 Forenbahe S, Kaiser T and Miracle PT (2013) Dating the East Adriatic Neolithic. *European Journal of Archaeology* 16(4): 589–609: doi:10.1179/1461957113Y.0000000038.
- 756
- 757 Forenbahe S and Miracle PT (2005) The spread of farming in the Eastern Adriatic. *Antiquity* 79: 514–528: doi:10.1017/S0003598X00114474.
- 758
- 759 Gardner AR and Willis KJ (1999) Prehistoric farming and the postglacial expansion of beech and hombeam: a comment on Küster. *The Holocene* 9(1): 119–121: doi:10.1191/095968399668254353.
- 760
- 761 Gavin DG, Hu FS, Lertzman K and Corbett P (2006) Weak climatic control of stand-scale fire history during the late Holocene. *Ecology* 87(7): 1722–1732.
- 762
- 763 Giesecke T, Brewer S, Finsinger W, Leydet M and Bradshaw RHW (2017) Patterns and dynamics of European vegetation change over the last 15,000 years. *Journal of Biogeography* 44: 1441–1456: doi:10.1111/jbi.12974.
- 764
- 765
- 766 Giesecke T, Davis B, Brewer S, Finsinger W, Wolters S, Blaauw M, et al. (2014) Towards mapping the late Quaternary vegetation change of Europe. *Vegetation History and Archaeobotany* 23: 75–86: doi:10.1007/s00334-012-0390-y.
- 767
- 768
- 769 Giesecke T, Hickler T, Kunkel T, Sykes MT and Bradshaw RHW (2007) Towards an understanding of the Holocene distribution of *Fagus sylvatica* L. *Journal of Biogeography* 34(1): 118–131: doi:10.1111/j.1365-2699.2006.01580.x.
- 770
- 771
- 772 Grindean R, Tanțău I and Feurdean A (2019) Linking vegetation dynamics and stability in the old-growth forests of Central Eastern Europe: Implications for forest conservation and management. *Biological Conservation* 229: 160–169: doi:10.1016/j.biocon.2018.11.019.
- 773
- 774
- 775 Hafner A, Reich J, Ballmer A, Bolliger M, Antolín F, Charles M, et al. (2021) First absolute chronologies of neolithic and bronze age settlements at Lake Ohrid based on dendrochronology and radiocarbon dating. *Journal of Archaeological Science: Reports* 38: 103107: doi:10.1016/j.jasrep.2021.103107.
- 776
- 777
- 778
- 779 Hastie T and Tibshirani R (1986) Generalized Additive Models. *Statistical Science*. Institute of Mathematical Statistics 1(3): 297–310: doi:10.1214/ss/1177013604.
- 780
- 781 Heiri O, Brooks SJ, Renssen H, Bedford A, Hazekamp M, Ilyashuk B, et al. (2014) Validation of climate model-inferred regional temperature change for late-glacial Europe. *Nature Communications* 5: 4914: doi:10.1038/ncomms5914.
- 782
- 783
- 784 Heiri O, Wick L, van Leeuwen JFN, van der Knaap WO and Lotter AF (2003) Holocene tree immigration and the chironomid fauna of a small Swiss subalpine lake (Hinterburgsee, 1515 m asl). *Palaeogeography Palaeoclimatology Palaeoecology* 189: 35–53.
- 785
- 786
- 787 Henne PD, Elkin C, Colombaroli D, Samartin S, Bugmann H, Heiri O, et al. (2013) Impacts of changing climate and land use on vegetation dynamics in a Mediterranean ecosystem: insights from paleoecology and dynamic modeling. *Landscape Ecology* 28: 819–833: doi:10.1007/s10980-012-9782-8.
- 788
- 789
- 790
- 791 Henne PD, Elkin CM, Reineking B, Bugmann H and Tinner W (2011) Did soil development limit spruce (*Picea abies*) expansion in the Central Alps during the Holocene? Testing a palaeobotanical hypothesis with a dynamic landscape model. *Journal of Biogeography* 38(5): 933–949: doi:10.1111/j.1365-2699.2010.02460.x.
- 792
- 793
- 794

- 795 Higuera PE, Brubaker LB, Anderson PM, Hu FS and Brown TA (2009) Vegetation mediated the  
796 impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska.  
797 *Ecological Monographs* 79: 201–219.
- 798 Higuera PE, Gavin DG, Bartlein PJ and Hallett DJ (2010) Peak detection in sediment–charcoal  
799 records: impacts of alternative data analysis methods on fire-history interpretations. *International*  
800 *Journal of Wildland Fire* 19(8): 996: doi:10.1071/WF09134.
- 801 Higuera PE, Sprugel DG and Brubaker LB (2005) Reconstructing fire regimes with charcoal from  
802 small-hollow sediments: a calibration with tree-ring records of fire. *The Holocene* 15: 238–251.
- 803 Hilmers T, Avdagić A, Bartkowicz L, Bielak K, Binder F, Bončina A, et al. (2019) The productivity of  
804 mixed mountain forests comprised of *Fagus sylvatica*, *Picea abies*, and *Abies alba* across Europe.  
805 *Forestry: An International Journal of Forest Research* 92(5): 512–522: doi:10.1093/forestry/cpz035.
- 806 Horvat I, Glavač V and Ellenberg H (1974) *Vegetation of Southeast Europe*. Stuttgart: Gustav Fischer  
807 Verlag.
- 808 Houston Durrant T, de Rigo D and Caudullo G (2016) *Fagus sylvatica* in Europe: distribution, habitat,  
809 usage and threats. In: San Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T and Mauri A  
810 (eds) *European Atlas of Forest Tree Species*. Luxembourg: Publ. Off. EU, e012b90+.
- 811 Hughes PD, Woodward JC, van Calsteren PC and Thomas LE (2011) The glacial history of the  
812 Dinaric Alps, Montenegro. *Quaternary Science Reviews* 30(23): 3393–3412:  
813 doi:10.1016/j.quascirev.2011.08.016.
- 814 Jain P, Castellanos-Acuna D, Coogan SCP, Abatzoglou JT and Flannigan MD (2022) Observed  
815 increases in extreme fire weather driven by atmospheric humidity and temperature. *Nature Climate*  
816 *Change* 12(1): 63–70: doi:10.1038/s41558-021-01224-1.
- 817 Kaplan JO, Krumhardt KM and Zimmermann N (2009) The prehistoric and preindustrial deforestation  
818 of Europe. *Quaternary Science Reviews* 28(27–28): 3016–3034: doi:10.1016/j.quascirev.2009.09.028.
- 819 Katz NJ, Katz SV and Kipiani MG (1965) *Atlas and keys of fruits and seeds occurring in the*  
820 *Quaternary deposits of the USSR*. Moscow: NAUKA.
- 821 Kelly R, Higuera PE, Barrett CM and Hu FS (2011) A signal-to-noise index to quantify the potential  
822 for peak detection in sediment–charcoal records. *Quaternary Research* 75(1): 11–17:  
823 doi:10.1016/j.yqres.2010.07.011.
- 824 Klopčič M, Mina M, Bugmann H and Bončina A (2017) The prospects of silver fir (*Abies alba* Mill.)  
825 and Norway spruce (*Picea abies* (L.) Karst) in mixed mountain forests under various management  
826 strategies, climate change and high browsing pressure. *European Journal of Forest Research* 136(5):  
827 1071–1090: doi:10.1007/s10342-017-1052-5.
- 828 Kranjc A (2009) History of Deforestation and Reforestation in the Dinaric Karst. *Geographical*  
829 *Research* 47(1): 15–23: doi:10.1111/j.1745-5871.2008.00552.x.
- 830 Longman J, Veres D, Finsinger W and Ersek V (2018) Exceptionally high levels of lead pollution in  
831 the Balkans from the Early Bronze Age to the Industrial Revolution. *Proceedings of the National*  
832 *Academy of Sciences* 1721546115: doi:10.1073/pnas.1721546115.
- 833 Maringer J, Wohlgemuth T, Hacket-Pain A, Ascoli D, Berretti R and Conedera M (2020) Drivers of  
834 persistent post-fire recruitment in European beech forests. *Science of The Total Environment* 699:  
835 134006: doi:10.1016/j.scitotenv.2019.134006.

- 836 McConnell JR, Wilson AI, Stohl A, Arienzo MM, Chellman NJ, Eckhardt S, et al. (2018) Lead  
837 pollution recorded in Greenland ice indicates European emissions tracked plagues, wars, and imperial  
838 expansion during antiquity. *Proceedings of the National Academy of Sciences*. National Academy of  
839 Sciences 115(22): 5726–5731: doi:10.1073/pnas.1721818115.
- 840 Mercuri AM, Bandini Mazzanti M, Florenzano A, Montecchi MC and Rattighieri E (2013) Olea,  
841 Juglans and Castanea: The OJC group as pollen evidence of the development of human-induced  
842 environments in the Italian peninsula. *Quaternary International* 303: 24–42:  
843 doi:10.1016/j.quaint.2013.01.005.
- 844 Mirković M (1985) Geological Map of southern Socialist Republic of Montenegro, 1:200,000.  
845 Titograd: The Republic self-Managing Community of Interest for Geological Exploration SR  
846 Montenegro.
- 847 Moore PD, Webb JA and Collinson ME (1998) *Pollen analysis*. Oxford: Blackwell Science Ltd.
- 848 Morales-Molino C, Leunda M, Morellón M, Gardoki J, Ezquerro FJ, Muñoz Sobrino C, et al. (2022)  
849 Millennial land use explains modern high-elevation vegetation in the submediterranean mountains of  
850 Southern Europe. *Journal of Biogeography* 49(10): 1779–1792: doi:10.1111/jbi.14472.
- 851 Morales-Molino C, Steffen M, Samartin S, van Leeuwen JFN, Hürlimann D, Vescovi E, et al. (2021)  
852 Long-term responses of mediterranean mountain forests to climate change, fire and human activities in  
853 the northern Apennines (Italy). *Ecosystems* 24: 1361–1377: doi:10.1007/s10021-020-00587-4.
- 854 Motta R, Berretti R, Castagneri D, Dukić V, Garbarino M, Govedar Z, et al. (2011) Toward a  
855 definition of the range of variability of central European mixed *Fagus – Abies – Picea* forests: the  
856 nearly steady-state forest of Lom (Bosnia and Herzegovina). *Canadian Journal of Forest Research*  
857 41(9): 1871–1884: doi:10.1139/x11-098.
- 858 Nagel TA, Mikac S, Dolinar M, Klopčič M, Keren S, Svoboda M, et al. (2017) The natural  
859 disturbance regime in forests of the Dinaric Mountains: A synthesis of evidence. *Forest Ecology and*  
860 *Management* 388: 29–42: doi:10.1016/j.foreco.2016.07.047.
- 861 Panagiotopoulos K, Aufgebauer A, Schäbitz F and Wagner B (2013) Vegetation and climate history of  
862 the Lake Prespa region since the Lateglacial. *Quaternary International* 293: 157–169:  
863 doi:10.1016/j.quaint.2012.05.048.
- 864 Pausas JG and Paula S (2012) Fuel shapes the fire-climate relationship: evidence from Mediterranean  
865 ecosystems: Fuel shapes the fire-climate relationship. *Global Ecology and Biogeography* 21(11):  
866 1074–1082: doi:10.1111/j.1466-8238.2012.00769.x.
- 867 Power MJ, Marlon J, Ortiz N, Bartlein PJ, Harrison SP, Mayle FE, et al. (2008) Changes in fire  
868 regimes since the Last Glacial Maximum: an assessment based on a global synthesis and analysis of  
869 charcoal data. *Climate Dynamics* 30: 887–907. doi:10.1007/s00382-007-0334-x.
- 870 Rasmussen SO, Bigler M, Blockley SP, Blunier T, Buchardt SL, Clausen HB, et al. (2014) A  
871 stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three  
872 synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy.  
873 *Quaternary Science Reviews* 106: 14–28: doi:10.1016/j.quascirev.2014.09.007.
- 874 Rastoder Š (2003) A short review of the history of Montenegro. In: Bieber F (ed) *Montenegro in*  
875 *transition: problems of identity and statehood*. Baden-Baden: Nomos Verlagsgesellschaft, 107–140.
- 876 Reille M (1992) *Pollen et spores d'Europe et d'Afrique du nord*. Marseille: Laboratoire de Botanique  
877 Historique et Palynologie.

- 878 Reimer PJ, Austin WEN, Bard E, Bayliss A, Blackwell PG, Ramsey CB, et al. (2020) The IntCal20  
879 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*. Cambridge  
880 University Press 62(4): 725–757: doi:10.1017/RDC.2020.41.
- 881 Renberg I, Bindler R and Brännvall M-L (2001) Using the historical atmospheric lead-deposition  
882 record as a chronological marker in sediment deposits in Europe. *The Holocene* 11(5): 511–516:  
883 doi:10.1191/095968301680223468.
- 884 Rey F, Gobet E, van Leeuwen JFN, Gilli A, van Raden UJ, Hafner A, et al. (2017) Vegetational and  
885 agricultural dynamics at Burgäschisee (Swiss Plateau) recorded for 18,700 years by multi-proxy  
886 evidence from partly varved sediments. *Vegetation History and Archaeobotany* 26(6): 571–586:  
887 doi:10.1007/s00334-017-0635-x.
- 888 Sabatini FM, Burrascano S, Keeton WS, Levers C, Lindner M, Pötzschner F, et al. (2018) Where are  
889 Europe’s last primary forests? *Diversity and Distributions* 24(10): 1426–1439: doi:10.1111/ddi.12778.
- 890 Samartin S, Heiri O, Joos F, Renssen H, Franke J, Brönnimann S, et al. (2017) Warm Mediterranean  
891 mid-Holocene summers inferred from fossil midge assemblages. *Nature Geoscience* 10: 207–212:  
892 doi:10.1038/ngeo2891.
- 893 Schlachter KJ and Horn SP (2010) Sample preparation methods and replicability in macroscopic  
894 charcoal analysis. *Journal of Paleolimnology* 44(2): 701–708: doi:10.1007/s10933-009-9305-z.
- 895 Schoch WH, Pawlik B and Schweingruber FH (1988) *Botanical macro-remains: An atlas for the*  
896 *determination of frequently encountered and ecologically important plant seeds*. Bern, Switzerland:  
897 Paul Haupt.
- 898 Seidl R, Thom D, Kautz M, Martin-Benito D, Peltoniemi M, Vacchiano G, et al. (2017) Forest  
899 disturbances under climate change. *Nature Climate Change* 7(6): 395–402: doi:10.1038/nclimate3303.
- 900 Sikkema R, Caudullo G and de Rigo D (2016) *Carpinus betulus* in Europe: distribution, habitat, usage  
901 and threats. In: San Miguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T and Mauri A (eds)  
902 *European Atlas of Forest Tree Species*. Luxembourg: Publ. Off. EU, e01d8cf+.
- 903 Spiegelberger T, Matthies D, Müller-Schärer H and Schaffner U (2006) Scale-dependent effects of  
904 land use on plant species richness of mountain grassland in the European Alps. *Ecography* 29(4): 541–  
905 548: doi:10.1111/j.0906-7590.2006.04631.x.
- 906 Srdanović M and Pavić D (2013) Environmental protection and major ecological problems in the  
907 National Park Durmitor. *Researches Reviews of the Department of Geography, Tourism and Hotel*  
908 *Management* 42: 9.
- 909 Stockmarr J (1971) Tablets with spores used in absolute Pollen Analysis. *Pollen et Spores* 13: 615–  
910 621.
- 911 Sugita S (1994) Pollen representation of vegetation in Quaternary sediments - theory and method in  
912 patchy vegetation. *Journal of Ecology* 82: 881–897.
- 913 Tinner W, Colombaroli D, Heiri O, Henne PD, Steinacher M, Untenecker J, et al. (2013) The past  
914 ecology of *Abies alba* provides new perspectives on future responses of silver fir forests to global  
915 warming. *Ecological Monographs* 83(4): 419–439: doi:10.1890/12-2231.1.
- 916 Tinner W, Conedera M, Ammann B, Gaggeler HW, Gedye S, Jones R, et al. (1998) Pollen and  
917 charcoal in lake sediments compared with historically documented forest fires in southern Switzerland  
918 since AD 1920. *The Holocene* 8(1): 31–42.

- 919 Tinner W, Conedera M, Gobet E, Hubschmid P, Wehrli M and Ammann B (2000) A palaeoecological  
920 attempt to classify fire sensitivity of trees in the southern Alps. *The Holocene* 10: 565–574.
- 921 Tinner W and Hu FS (2003) Size parameters, size-class distribution and area-number relationship of  
922 microscopic charcoal: relevance for fire reconstruction. *The Holocene* 13: 499–505.
- 923 Tinner W and Lotter AF (2006) Holocene expansions of *Fagus sylvatica* and *Abies alba* in Central  
924 Europe: where are we after eight decades of debate? *Quaternary Science Reviews* 25: 526–549.
- 925 Tinner W, Lotter AF, Ammann B, Conedera M, Hubschmid P, van Leeuwen JFN, et al. (2003)  
926 Climatic change and contemporaneous land-use phases north and south of the Alps 2300 BC to 800  
927 AD. *Quaternary Science Reviews* 22: 1447–1460.
- 928 Tóth M, Buzckó K, Magyari EK, Braun M, Panagiotopoulos K and Heiri O (2015) Chironomid-  
929 inferred Holocene temperature changes in the South Carpathians (Romania). *The Holocene* 25: 569–  
930 582: doi:10.1177/0959683614565953.
- 931 UNESCO and IUCN (2018) *Joint World Heritage Centre/IUCN Advisory mission to the World*  
932 *Heritage property “Durmitor National Park” (Montenegro)*. Paris, France and Gland, Switzerland:  
933 UNESCO World Heritage Centre and IUCN. Available at:  
934 <https://whc.unesco.org/en/documents/174707/>.
- 935 Vander Linden M, Pandžić I and Orton D (2014) New radiocarbon dates for the Neolithic period in  
936 Bosnia & Herzegovina. *Godišnjak Centra za balkanološka ispitivanja* 43: 7–34:  
937 doi:10.5644/Godisnjak.CBI.ANUBiH-43.35.
- 938 Vescovi E, Ravazzi C, Arpent E, Finsinger W, Pini R, Valsecchi V, et al. (2007) Interactions between  
939 climate and vegetation during the Lateglacial period as recorded by lake and mire sediment archives in  
940 Northern Italy and Southern Switzerland. *Quaternary Science Reviews* 26: 1650–1669: doi:Article.
- 941 Wagner B, Lotter AF, Nowaczyk N, Reed JM, Schwalb A, Sulpizio R, et al. (2009) A 40,000-year  
942 record of environmental change from ancient Lake Ohrid (Albania and Macedonia). *Journal of*  
943 *Paleolimnology* 41(3): 407–430: doi:10.1007/s10933-008-9234-2.
- 944 Weltje GJ, Bloemsma MR, Tjallingii R, Heslop D, Röhl U and Croudace IW (2015) Prediction of  
945 Geochemical Composition from XRF Core Scanner Data: A New Multivariate Approach Including  
946 Automatic Selection of Calibration Samples and Quantification of Uncertainties. In: Croudace IW and  
947 Rothwell RG (eds) *Micro-XRF Studies of Sediment Cores: Applications of a non-destructive tool for*  
948 *the environmental sciences*. Dordrecht: Springer Netherlands, 507–534. Available at:  
949 [https://doi.org/10.1007/978-94-017-9849-5\\_21](https://doi.org/10.1007/978-94-017-9849-5_21): doi:10.1007/978-94-017-9849-5\_21.
- 950 Westerling AL, Hidalgo HG, Cayan DR and Swetnam TW (2006) Warming and Earlier Spring  
951 Increase Western U.S. Forest Wildfire Activity. *Science*. American Association for the Advancement  
952 of Science 313(5789): 940–943: doi:10.1126/science.1128834.
- 953 Whitlock C and Larsen C (2001) Charcoal as a fire proxy. In: Smol JP, Birks HJB, Last WM and Last  
954 WM (eds) *Terrestrial, Algal, and Siliceous Indicators*. Dordrecht, The Netherlands: Kluwer Academic  
955 Publishers, 75–97.
- 956 Willis KJ (1994) The vegetational history of the Balkans. *Quaternary Science Reviews* (13): 769–788.
- 957 Wood SN (2017) *Generalized Additive Models: An Introduction with R, Second Edition*. New York:  
958 Chapman and Hall/CRC: doi:10.1201/9781315370279.

959 

<p><b>Table 2</b> Ecos Montenegro), charcoal-accu Verdarolo (Sa 1 indicates a s</p>
---