



HAL
open science

ABO blood types and SARS-CoV-2 infection assessed using seroprevalence data in a large population-based sample: the SAPRIS-SERO multi-cohort study

Mélanie Deschasaux-Tanguy, Fabien Szabo de Edelenyi, Nathalie Druesne-Pecollo, Younes Esseddik, Julien Allègre, Bernard Srour, Pilar Galan, Serge Hercberg, Gianluca Severi, Marie Zins, et al.

► To cite this version:

Mélanie Deschasaux-Tanguy, Fabien Szabo de Edelenyi, Nathalie Druesne-Pecollo, Younes Esseddik, Julien Allègre, et al.. ABO blood types and SARS-CoV-2 infection assessed using seroprevalence data in a large population-based sample: the SAPRIS-SERO multi-cohort study. *Scientific Reports*, 2023, 13 (1), pp.4775. 10.1038/s41598-023-30714-9 . hal-04059423

HAL Id: hal-04059423

<https://hal.science/hal-04059423>

Submitted on 30 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN

ABO blood types and SARS-CoV-2 infection assessed using seroprevalence data in a large population-based sample: the SAPRIS-SERO multi-cohort study

Mélanie Deschasaux-Tanguy^{1✉}, Fabien Szabo de Edelenyi¹, Nathalie Druesne-Pecollo¹, Younes Esseddik¹, Julien Allègre¹, Bernard Srour¹, Pilar Galan¹, Serge Hercberg¹, Gianluca Severi^{2,3}, Marie Zins⁴, Emmanuel Wiernik⁴, SAPRIS-SERO study group^{*}, Xavier de Lamballerie⁵, Fabrice Carrat^{6,7} & Mathilde Touvier¹

ABO blood type has been reported as a potential factor influencing SARS-CoV-2 infection, but so far mostly in studies that involved small samples, selected population and/or used PCR test results. In contrast our study aimed to assess the association between ABO blood types and SARS-CoV-2 infection using seroprevalence data (independent of whether or not individuals had symptoms or sought for testing) in a large population-based sample. Our study included 67,340 French participants to the SAPRIS-SERO multi-cohort project. Anti-SARS-CoV-2 antibodies were detected using ELISA (targeting the proteins spike (S) and nucleocapsid (NP)) and seroneutralisation (SN) tests on dried blood spots collected in May–November 2020. Non-O individuals (and especially types A and AB) were more likely to bear anti SARS-CoV-2 antibodies (ELISA-S, 2964 positive cases: $OR_{non-Ovs.O} = 1.09[1.01–1.17]$, $OR_{Avs.O} = 1.08[1.00–1.17]$; ELISA-S/ELISA-NP/SN, 678 triple positive cases: $OR_{non-Ovs.O} = 1.19[1.02–1.39]$, $OR_{Avs.O} = 1.19[1.01–1.41]$, $OR_{ABvs.O} = 1.43[1.01–2.03]$). Hence, our results provided additional insights into the dynamic of SARS-CoV-2 infection, highlighting a higher susceptibility of infection for individuals of blood types A and AB and a lesser risk for blood type O.

Since 2020, the world has been struggling with the COVID-19 (Coronavirus Disease 2019) pandemic caused by the SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2). The research community has mobilized to identify potential risk factors associated with SARS-CoV-2 infection and COVID-19 severity, in an effort to better understand the dynamic of infection, pinpoint individuals at higher risk and help prevent the disease and related adverse outcomes. Early reports comparing the distribution of ABO blood types between individuals infected or not by the SARS-CoV-2, suggested that ABO blood types could be related to the risk of SARS-CoV-2 infection^{1–4}, as previously observed for SARS-CoV⁵ but also for other infectious diseases^{6,7}. These studies and others published since then^{8–11} suggested that individuals with blood type O had a lower risk of SARS-CoV-2

¹Inserm U1153, Inrae U1125, Cnam, Nutritional Epidemiology Research Team (EREN), Epidemiology and Statistics Research Center – Université Paris Cité (CRESS), Sorbonne Paris Nord University, Bobigny, France. ²UVSQ, Inserm, Gustave Roussy, “Exposome and Heredity” team, CESP UMR1018, Paris-Saclay University, Villejuif, France. ³Department of Statistics, Computer Science and Applications “G. Parenti”, University of Florence, Florence, Italy. ⁴Population-based Cohorts Unit, INSERM, UMS 011, Paris Saclay University, Université de Versailles Saint-Quentin-en-Yvelines, Université Paris Cité, Paris, France. ⁵Unité des Virus Emergents (UVE), IRD 190, INSERM 1207, Aix Marseille Univ, Marseille, France. ⁶Inserm, Institut Pierre-Louis d’Epidémiologie et de Santé Publique, Sorbonne Université, Paris, France. ⁷Département de Santé Publique, APHP, Sorbonne Université, Paris, France. ^{*}A list of authors and their affiliations appears at the end of the paper. ✉email: m.deschasaux@eren.smbh.univ-paris13.fr

infection and that type A individuals would be more at risk, with yet some remaining inconsistencies, especially for B or AB types for which results are less robust. However, these results were obtained in selected populations (e.g., patients, blood donors) of various sample size (mostly under 15,000) and mostly using PCR test results (e.g., from questionnaires or registries) hence reflecting a population that sought for testing, with the exception of a recent report using seroprevalence data in $\approx 32,600$ blood donors¹¹. In contrast, this short report provides insights into the associations between ABO blood types and SARS-CoV-2 infection assessed from seroprevalence data obtained between May and November 2020 through the systematic screening for anti-SARS-CoV-2 antibodies in a large population-based sample of French adults.

Methods

The SAPRIS (“Santé, Perception, pratiques, Relations et Inégalités Sociales en population générale pendant la crise COVID-19”) study was set up in April 2020 to investigate various aspects of the COVID-19 crisis (COVID-19 infection/diagnosis and experience of lockdown), based on a consortium of existing French prospective cohort studies. In May 2020, participants from the Constances, E3N-E4N and NutriNet-Santé cohort studies answering SAPRIS questionnaires were also invited to take part in the SAPRIS-SERO project which aimed to estimate the seroprevalence of antibodies against SARS-CoV-2 at the population level, as previously described^{12,13}. Briefly, between May and November 2020 (during or at the end of the first wave, pre-vaccine period), participants received self-sampling kits by mail to collect dried blood spots from which were detected anti-SARS-CoV-2 antibodies (IgG) directed against the S1 domain of the spike protein (S) and the nucleocapsid protein (NP) using ELISA (Euroimmun®, Lübeck, Germany), as well as neutralizing anti-SARS-CoV-2 antibodies (SN) using an in-house micro-neutralization assay. ELISA-S and ELISA-NP tests were considered to be positive with an optical density ratio ≥ 1.1 , indeterminate between 0.8 and 1.1, and negative < 0.8 . The SN test was considered to be positive with a titre ≥ 40 . ELISA-NP and SN tests were performed whenever the optical density ratio for ELISA-S was ≥ 0.7 . Positivity (vs. negativity) to ELISA-S test and positivity (vs. negativity) to all three ELISA-S, ELISA-NP and SN tests (associated to more symptomatic COVID-19 in a previous study in SAPRIS-SERO¹³) were considered as outcomes. Characteristics of the participants including ABO and RhD blood types were collected from questionnaires as part of each cohort follow-up or from the shared SAPRIS questionnaire. Associations between the seroprevalence of antibodies against SARS-CoV-2 and blood types were studied using multi-adjusted logistic regression models stratified by cohort and including the following covariates: sex (men/women), age (≥ 18 – < 40 , 40–49, 50–59, 60–69, ≥ 70 years old), month of blood collection, residential area during the lockdown: city size (rural, $< 20,000$, 20–100,000, $\geq 100,000$ inhabitants) and French administrative region. Additional covariates linked to the risk of SARS-CoV-2 infection were included in sensitivity analyses: weight status, smoking status, educational level, professional activity during lockdown, socio-professional category, number of individuals at home during lockdown and alcohol intake. All tests were two-sided and $P < 0.05$ was considered statistically significant. Analyses were carried out using SAS 9.4 (SAS Institute Inc., USA).

Ethics approval and informed consent. The SAPRIS-SERO study (registered #NCT04392388) was conducted in accordance with the relevant guidelines and regulations and was approved by the ethics committee CPP Sud-Méditerranée III on April 27th 2020 and by the CNIL #920,193. All participants provided written or electronic informed consent for participation in each cohort and a specific electronic informed consent for participation in the SAPRIS-SERO study.

Results

Our study included 67,340 participants, among which 2964 with a positive seroprevalence of anti SARS-CoV-2 antibodies according to the ELISA-S test and 64,376 participants who tested negative (respectively 678 triple positive and 64,011 triple negative to ELISA-S, ELISA-NP and SN tests). Characteristics of participants are shown in Table 1.

Higher odds of presenting anti SARS-CoV-2 antibodies (Table 2) were observed in participants belonging to non-O blood types (ELISA-S: OR_{non-O vs. O} = 1.09 [1.01–1.17], ELISA-S/ELISA-NP/SN: OR_{non-O vs. O} = 1.19 [1.02–1.39]) and especially to blood type A (ELISA-S: OR_{A vs. O} = 1.08 [1.00–1.17], ELISA-S/ELISA-NP/SN: OR_{A vs. O} = 1.19 [1.01–1.41]) and AB (ELISA-S/ELISA-NP/SN: OR_{AB vs. O} = 1.43 [1.01–2.03]). In analyses combining ABO and RhD blood types, higher odds were observed in participants with blood types A- (ELISA-S: OR_{A- vs. O+} = 1.16 [1.00–1.35]), A+ (ELISA-S/ELISA-NP/SN: OR_{A+ vs. O+} = 1.24 [1.03–1.48]) and AB+ (which represented the majority of AB participants, ELISA-S: OR_{AB+ vs. O+} = 1.24 [1.02–1.51], ELISA-S/ELISA-NP/SN: OR_{AB+ vs. O+} = 1.60 [1.11–2.30]). No association was observed comparing RhD positive and negative blood types overall.

Additional adjustments in sensitivity analyses did not change the results of increased odds of positive seroprevalence in non-O participants (ELISA-S: OR_{non-O vs. O} = 1.09 [1.01–1.18], ELISA-S/ELISA-NP/SN: OR_{non-O vs. O} = 1.19 [1.02–1.39]).

Discussion

In this study, we used seroprevalence data obtained from screening a large population-based sample from 3 French prospective cohorts to assess the associations between ABO blood types and the risk of SARS-CoV-2 infection. Our results showed a decreased seroprevalence of anti-SARS-CoV-2 antibodies in type O individuals compared to all others. Among non-O blood types, types A and AB were more likely positive for SARS-CoV-2 infection. No association was observed with type B or RhD types.

These results confirmed prior observations obtained in other population settings, namely a decrease risk of infection for type O individuals and an increase risk for type A individuals, with some indication of increased

| | ELISA-S negative (N = 64,376) | | ELISA-S positive (N = 2,964) | | P |
|--|----------------------------------|--------|---------------------------------|--------|--------|
| | N | % | N | % | |
| Gender | | | | | <0.001 |
| Male | 21,174 | (32.9) | 832 | (28.1) | |
| Female | 43,202 | (67.1) | 2132 | (71.9) | |
| Age (years) | | | | | <0.001 |
| ≥ 18– < 40 | 6983 | (10.8) | 652 | (22.0) | |
| 40–49 | 10,340 | (16.1) | 988 | (33.3) | |
| 50–59 | 11,222 | (17.4) | 505 | (17.0) | |
| 60–69 | 13,224 | (20.5) | 334 | (11.3) | |
| ≥ 70 | 22,607 | (35.1) | 485 | (16.4) | |
| Weight status ^a | | | | | 0.006 |
| Underweight | 1992 | (3.1) | 110 | (3.7) | |
| Normal weight | 37,189 | (57.8) | 1749 | (59.0) | |
| Overweight | 17,763 | (27.6) | 745 | (25.1) | |
| Obesity | 5917 | (9.2) | 273 | (9.2) | |
| Smoking status ^a | | | | | <0.001 |
| Non smoker | 29,586 | (46.0) | 1605 | (54.1) | |
| Smoker | 5958 | (9.3) | 237 | (8.0) | |
| Ex-smoker | 26,351 | (40.9) | 997 | (33.6) | |
| Educational level ^a | | | | | <0.001 |
| < High school | 7262 | (11.3) | 171 | (5.8) | |
| High school—undergraduate | 28,063 | (43.6) | 1188 | (40.1) | |
| Graduate or doctorate | 24,074 | (37.4) | 1307 | (44.1) | |
| Month of blood collection | | | | | <.0001 |
| May | 8004 | (12.4) | 663 | (22.4) | |
| June | 427 | (0.7) | 25 | (0.7) | |
| July | 38,143 | (59.3) | 1556 | (52.5) | |
| August | 13,120 | (20.4) | 540 | (18.2) | |
| September | 4607 | (7.2) | 178 | (6.0) | |
| October | 72 | (0.1) | 1 | (0.0) | |
| November | 3 | (0.0) | 1 | (0.0) | |
| Residential area during the lockdown : city size ^a | | | | | <0.001 |
| Rural area | 14,876 | (23.1) | 555 | (18.7) | |
| < 20,000 inhabitants | 10,923 | (17.0) | 441 | (14.9) | |
| 20–100,000 inhabitants | 10,600 | (16.5) | 508 | (17.1) | |
| ≥ 100,000 inhabitants | 27,230 | (42.3) | 1441 | (48.6) | |
| Residential area during the lockdown : French administrative region ^a | | | | | <.0001 |
| Haut de France | 4116 | (6.4) | 179 | (6.0) | |
| Auvergne- Rhône-Alpes | 8316 | (12.9) | 351 | (11.8) | |
| Occitanie | 6979 | (10.9) | 188 | (6.3) | |
| PACA | 4306 | (6.7) | 133 | (4.5) | |
| Corse | 77 | (0.1) | 3 | (0.1) | |
| Normandie | 2463 | (3.8) | 87 | (2.9) | |
| Ile-de-France | 10,492 | (16.3) | 818 | (27.6) | |
| Grand-Est | 4679 | (7.3) | 389 | (13.1) | |
| Bretagne | 5066 | (7.9) | 143 | (4.8) | |
| Pays de la Loire | 3952 | (6.1) | 144 | (4.9) | |
| Centre-Val-de-Loire | 4224 | (6.6) | 174 | (5.9) | |
| Bourgogne | 2463 | (3.8) | 111 | (3.7) | |
| Nouvelle Aquitaine | 7228 | (11.2) | 243 | (8.2) | |
| DOM-TOM | 8 | (0.0) | 1 | (0.0) | |

Table 1. Characteristics of the population, SAPRIS-SERO multi-cohort study, 2020. ^aMissing values: N = 1602 (2.4%) for weight status, N = 2606 (3.9%) for smoking status, N = 5275 (7.8%) for educational level, N = 766 (1.1%) for city size and N = 7 (0.01%) for French administrative region.

| | N (%) | ELISA-S | | | | ELISA-S, ELISA-NP, SN | | |
|---------|---------------|-------------------------|-------------------------|------|---------------|--------------------------------------|------------------|------|
| | | N positive/ negative | OR [95%CI] ^a | P | | N positive/ negative ^b | OR [95%CI] | P |
| ABO | | | | | | | | 0.08 |
| O | 28,752 (42.7) | 1216/27,536 | Ref | | 27,648 (42.7) | 260/27,388 | Ref | |
| A | 28,742 (44.4) | 1342/28,574 | 1.08 [1.00–1.17] | | 28,742 (44.4) | 323/28,419 | 1.19 [1.01–1.41] | |
| B | 5,874 (8.7) | 275/5599 | 1.10 [0.96–1.27] | | 5,613 (8.7) | 58/5555 | 1.06 [0.80–1.42] | |
| AB | 2,798 (4.2) | 131/2667 | 1.14 [0.95–1.38] | | 2,686 (4.2) | 37/2649 | 1.43 [1.01–2.03] | |
| Non-O | 38,588 (57.3) | 1748/36,840 | 1.09 [1.01–1.17] | 0.03 | 37,041 (57.3) | 418/36,623 | 1.19 [1.02–1.39] | 0.03 |
| ABO/RhD | | | | | | | | 0.05 |
| O+ | 23,703 (35.2) | 995/22,708 | Ref | | 22,804 (35.3) | 217/22,587 | Ref | |
| O– | 5,049 (7.5) | 221/4828 | 1.03 [0.89–1.20] | | 4,844 (7.5) | 43/4801 | 0.96 [0.69–1.33] | |
| A+ | 25,315 (37.6) | 1120/24,195 | 1.07 [0.98–1.17] | | 24,344 (37.6) | 286/24,058 | 1.24 [1.03–1.48] | |
| A– | 4,601 (6.8) | 222/4379 | 1.16 [1.00–1.35] | | 4,398 (6.8) | 37/4361 | 0.91 [0.64–1.29] | |
| B+ | 4,886 (7.3) | 224/4662 | 1.09 [0.94–1.27] | | 4,675 (7.2) | 48/4627 | 1.04 [0.76–1.43] | |
| B– | 988 (1.5) | 51/937 | 1.21 [0.90–1.63] | | 938 (1.5) | 10/928 | 1.14 [0.60–2.17] | |
| AB+ | 2,363 (3.5) | 118/2245 | 1.24 [1.02–1.51] | | 2,266 (3.5) | 35/2231 | 1.60 [1.11–2.30] | |
| AB– | 435 (0.6) | 13/422 | 0.70 [0.40–1.22] | | 420 (0.6) | 2/418 | 0.48 [0.12–1.93] | |
| RhD | | | | 0.48 | | | | 0.08 |
| + | 56,267 (83.6) | 2457/53,810 | Ref | | 54,089 (83.6) | 586/53,503 | Ref | |
| – | 11,073 (16.4) | 507/10,566 | 0.97 [0.87–1.07] | | 10,600 (16.4) | 92/10,508 | 1.22 [0.98–1.52] | |

Table 2. Associations between ABO and RhD blood types and the seroprevalence of anti-SARS-CoV-2 antibodies, SAPRIS-SERO multi-cohort study, 2020. ^aOR and 95%CI obtained from a multi-adjusted logistic regression model stratified by cohort and including the following covariates: sex (men/women), age (<40ans, 40–49, 50–59, 60–69, ≥70 years old), month of blood collection, residential area during the lockdown: city size (rural, <20,000, 20–10,000, ≥100,000 inhabitants) and French administrative region. ^bA participant was considered positive (respectively negative) if positive (respectively negative) to all three ELISA-S, ELISA-NP, and SN tests. Among the N = 2964 individuals with a positive seroprevalence of anti SARS Cov-2 antibodies according to the ELISA-S test, N = 1185 were also positive according to the ELISA-NP test and 678 were positive according to all three ELISA-S, ELISA-NP, and SN tests.

risk for type AB individuals as well, thus strengthening the evidence towards a differential susceptibility of SARS-CoV-2 infection according to ABO blood types. Yet, most studies so far involved registry or hospital data of SARS-CoV-2 positivity from PCR test results in patients or blood donors in comparison to the general population^{8–10} and fewer studies used seroprevalence data, mostly in small samples, with the exception of a recent report in 32,600 blood donors¹¹. In turn, our study used seroprevalence data and a screening design whereby a large population-based sample of 67,340 participants involved in the SAPRIS-SERO project received a dried blood spot sampling kit. This allowed us to test for SARS-CoV-2 infection in a single general population, regardless whether the participants sought testing or had symptoms, allowing detecting a large panel of infected cases.

This differential susceptibility to SARS-CoV-2 infection according to blood types could be due to the incorporation of ABH antigens on the envelop of SARS-CoV-2 virions, leading to a likelihood of infection from one individual to another following ABO incompatibility^{9,10}. As type O individuals carry both anti-A and anti-B antibodies, they would be less susceptible to primary infection by SARS-CoV-2 viruses enveloped with type A or B-like patterns such as those excreted by non-O individuals. In contrast, type A individuals, carrying anti-B antibodies, would be susceptible to infection by viruses excreted by type O and type A individuals (i.e., 42% and 44% of the French population respectively), while type AB individuals carrying neither anti-A or anti-B antibodies would be susceptible to infection by individuals of all group types, which is why type A or AB are particularly at risk for SARS-CoV-2 infection. Type B individuals carrying anti-A antibodies are susceptible to infection by viruses excreted by type O and type B individuals but type B is less represented in the French population (10%) which may explain why no significant association was observed for these individuals. Another hypothesis, yet less supported by available data relates to the entry of SARS-CoV-2 in host's cells, that would be facilitated by antigen A^{9,14}. Finally, non-O individuals are at higher risk of thromboembolism, a common complication of COVID-19, hence beyond infection, non-O individuals could also be at higher risk of severe cases of COVID-19^{9,10}.

Our study strengths pertained to the systematic screening for anti-SARS-CoV-2 antibodies targeting a large well-characterized sample, hence providing data from positive and negative cases within the same population, as well as the use of seroprevalence allowing the detection of cases that may or may not have symptoms or sought testing. Yet, some limitations should also be acknowledged. First, although large, our sample size may have still been limited to have sufficient power in analyses combining ABO and RhD blood types, especially for rarer blood types. Next, misclassification of individuals as 'negative' may have arisen from imperfect sensitivity of the ELISA-S test (85–90%, the specificity being >95%)^{15,16} and from the decrease of anti-SARS-CoV-2 antibodies over time (with no detectable antibodies 4–5 months post-infection in half of participants in a recent report¹¹). Yet, the

sample collection was performed between May and November 2020, the first and second waves of the pandemic occurring in spring and fall 2020 in France, so that antibodies should still be detectable in most participants. Finally, the observational nature of the study makes it difficult to assess the extent to which participants have actually been exposed to the virus. Although lifestyle-related exposure to the virus is mostly unlikely to have been differential according to blood types, we adjusted our models on several covariates related to the spatial distribution of the epidemic since it could also relate to blood types, considering the distribution of ABO blood types in different regions or populations. However, we did not have access to racial and ethnic data as the collection of such data is generally not permitted in the framework of French cohort studies. Finally, our data only reflect infections occurring before November 2020, that is, before the emergence of variants but also before the large-scale vaccination campaigns.

To conclude, this study adds to the current body of evidence providing data on the association between ABO blood types and the risk of SARS-CoV-2 infection from screening a large sample from the general population using seroprevalence data. Our results confirmed a higher susceptibility of infection for individuals of blood types A and AB and a lesser risk for blood type O, thereby providing additional insights into the dynamic of SARS-CoV-2 infection. However, considering the magnitude of association and the current mechanistic hypotheses related to blood type incompatibility (hence a collective rather than individual type of protection), type O individuals are not without risk of infection and protective measures remain the best way of preventing SARS-CoV-2 transmission and infection.

Data availability

Data from the study are protected under the protection of health data regulation set by the French National Commission on Informatics and Liberty (Commission Nationale de l'Informatique et des Libertés, CNIL). The data can be made available upon reasonable request to the corresponding author (m.deschasaux@eren.smbh.univ-paris13.fr), after a consultation with the steering committee of the SAPRIS-SERO study. The French law forbids us to provide free access to SAPRIS-SERO data; access could however be given by the steering committee after legal verification of the use of the data. Please, feel free to come back to us should you have any additional question.

Received: 18 October 2022; Accepted: 28 February 2023

Published online: 23 March 2023

References

1. Zhao, J. *et al.* Relationship between the ABO blood group and the coronavirus disease 2019 (COVID-19) susceptibility. *Clin. Infect. Dis.* **73**, 328–331 (2021).
2. Göker, H. *et al.* The effects of blood group types on the risk of COVID-19 infection and its clinical outcome. *Turk J. Med. Sci.* **50**, 679–683 (2020).
3. Abdollahi, A., Mahmoudi-Aliabadi, M., Mehrdash, V., Jafarzadeh, B. & Salehi, M. The novel coronavirus SARS-CoV-2 vulnerability association with ABO/Rh blood types. *Iran J. Pathol.* **15**, 156–160 (2020).
4. Gallian, P. *et al.* Lower prevalence of antibodies neutralizing SARS-CoV-2 in group O French blood donors. *Antivir. Res.* **181**, 104880 (2020).
5. Cheng, Y. *et al.* ABO blood group and susceptibility to severe acute respiratory syndrome. *JAMA* **293**, 1450–1451 (2005).
6. Lindesmith, L. *et al.* Human susceptibility and resistance to Norwalk virus infection. *Nat. Med.* **9**, 548–553 (2003).
7. Batool, Z., Durrani, S. H. & Tariq, S. Association of ABO and Rh blood group types to hepatitis B, hepatitis C, HIV and Syphilis infection, a five year experience in healthy blood donors in a tertiary care hospital. *J. Ayub. Med. Coll. Abbottabad* **29**, 90–92 (2017).
8. Banchelli, F. *et al.* The role of ABO blood type in patients with SARS-CoV-2 infection: A systematic review. *J. Clin. Med.* **11**, 3029 (2022).
9. Pereira, E. *et al.* ABO blood group and link to COVID-19: A comprehensive review of the reported associations and their possible underlying mechanisms. *Microb. Pathog.* **169**, 105658 (2022).
10. Pendu, J. L., Breiman, A., Rocher, J., Dion, M. & Ruvoën-Clouet, N. ABO blood types and COVID-19: Spurious, anecdotal, or truly important relationships? A reasoned review of available data. *Viruses* **13**, 160 (2021).
11. Gallian, P. *et al.* SARS-CoV-2 IgG seroprevalence surveys in blood donors before the vaccination campaign, France 2020–2021. (2022). <https://doi.org/10.1101/2022.07.29.22278190>.
12. Carrat, F. *et al.* Age, COVID-19-like symptoms and SARS-CoV-2 seropositivity profiles after the first wave of the pandemic in France. *Infection* **50**, 257–262 (2022).
13. Carrat, F. *et al.* Antibody status and cumulative incidence of SARS-CoV-2 infection among adults in three regions of France following the first lockdown and associated risk factors: A multicohort study. *Int. J. Epidemiol.* **50**, 1458–1472 (2021).
14. Schetelig, J. *et al.* Blood group A epitopes do not facilitate entry of SARS-CoV-2. *J. Intern. Med.* **290**, 223–226 (2021).
15. Patel, E. U. *et al.* Comparative performance of five commercially available serologic assays to detect antibodies to SARS-CoV-2 and identify individuals with high neutralizing titers. *J. Clin. Microbiol.* **59**, 02257–02320 (2021).
16. Rikhtegaran Tehrani, Z. *et al.* Performance of nucleocapsid and spike-based SARS-CoV-2 serologic assays. *PLoS ONE* **15**, e0237828 (2020).

Acknowledgements

The authors warmly thank all the volunteers of the Constances, E3N-E4N and NutriNet-Santé cohorts for their continuous participation in the study and for participating in the SAPRIS-SERO project. We thank the CEPH-Biobank staff for their adaptability and the quality of their work. In the virology department, we thank Dr Nadège Brisbarre and the technical staff for impeccable management of samples and serological assays.

Author contributions

The authors' contributions were as follows – M.D.T., M.T. conceptualized the study and defined the analytical strategy; F.S.E.: performed statistical analyses; M.D.T. supervised statistical analyses and drafted the manuscript; M.T.: supervised statistical analyses and writing; F.S.E., N.D.P., Y.E., J.A., P.G., S.H., G.S., M.Z., E.W., X.dL, F.C., M.T., SAPRIS-SERO study group: played a key role in the acquisition of the data. All authors critically helped in

the interpretation of results, revised the manuscript and provided relevant intellectual input. They all read and approved the final manuscript. M.D.T. and M.T. had primary responsibility for the final content.

Funding

The SAPRIS/SAPRIS-SERO projects received funding from ANR (Agence Nationale de la Recherche, #ANR-20-COVI-000, #ANR-10-COHO-06), Fondation pour la Recherche Médicale (#20RR052-00), and Inserm (Institut National de la Santé et de la Recherche Médicale, #C20-26). The NutriNet-Santé cohort study was supported by the following public institutions: Ministère de la Santé, Santé Publique France, Institut National de la Santé et de la Recherche Médicale (INSERM), Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), Conservatoire National des Arts et Métiers (CNAM) and Université Sorbonne Paris Nord. The CONSTANCES cohort benefits from grant ANR-11-INBS-0002 from the French National Research Agency. CONSTANCES is supported by the Caisse Nationale d'Assurance Maladie, the French Ministry of Health, the Ministry of Research, and the Institut National de la Santé et de la Recherche Médicale (INSERM). CONSTANCES is also partly funded by AstraZeneca, Lundbeck, L'Oréal, and Merck Sharp & Dohme Corp. The E3N-E4N cohort is supported by the following institutions: Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation, INSERM, University Paris-Saclay, Gustave Roussy, the MGEN, and the French League Against Cancer. Study investigators are independent from the funders. Funders had no role in the study design, the collection, analysis, and interpretation of data, the writing of the manuscript, or the decision to submit the article for publication.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.D.-T.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

SAPRIS-SERO study group

SAPRIS-SERO coordination

Fabrice Carrat^{6,7}, Pierre-Yves Ancel⁸, Nathalie Bajos¹⁰, Marie-Aline Charles⁹, Gianluca Severi^{2,3}, Mathilde Touvier¹ & Marie Zins⁴

⁸Obstetrical, Perinatal and Pediatric Research Team (EPOPé), Centre for Research in Epidemiology and Statistics (CRESS), Inserm, INRAE, Université de Paris, Paris, France. ⁹Centre for Research in Epidemiology and Statistics (CRESS), Inserm, INRAE, Université de Paris, Paris, France. ¹⁰IRIS, UMR CNRS 8156, EHES, Inserm U997, Aubervilliers, France.

Constances cohort

Sofiane Kab⁴, Adeline Renuy⁴, Stéphane Le Got⁴, Céline Ribet⁴, Mireille Pellicer⁴, Emmanuel Wiernik⁴, Marcel Goldberg⁴ & Marie Zins⁴

E3N-E4N cohort

Fanny Artaud², Pascale Gerbouin-Rérolle², Mélody Enguix², Camille Laplanche², Roselyn Gomes-Rimav², Lyan Hoang², Emmanuelle Correia², Alpha Amadou Barry², Nadège Senina² & Gianluca Severi^{2,3}

NutriNet-Santé cohort

Fabien Szabo de Edelenyi¹, Julien Allègre¹, Nathalie Druesne-Pecollo¹, Younes Esseddik¹, Serge Hercberg¹ & Mathilde Touvier¹

Epipage2 and Elfe child cohorts

Marie-Aline Charles⁹, Pierre-Yves Ancel⁸, Valérie Benhammou¹¹, Anass Ritmi¹¹, Laetitia Marchand¹¹, Cecile Zaros¹², Elodie Lordmi¹², Adriana Candea¹², Sophie de Visme¹², Thierry Simeon¹², Xavier Thierry¹², Bertrand Geay¹², Marie-Noelle Dufourg¹² & Karen Milcent¹²

⁸Obstetrical, Perinatal and Pediatric Research Team (EPOPé), Centre for Research in Epidemiology and Statistics (CRESS), Inserm, INRAE, Université de Paris, Paris, France. ⁹Centre for Research in Epidemiology and Statistics (CRESS), Inserm, INRAE, Université de Paris, Paris, France. ¹¹EPIPAGE-2 Joint Unit, Paris, France. ¹²ELFE Joint Unit, Paris, France.

IPLESP – methodology and coordinating data center

Clovis Lusivika-Nzinga⁶, Gregory Pannetier⁶, Nathanael Lapidus⁶, Isabelle Goderel⁶, Céline Dorival⁶, Jérôme Nicol⁶, Olivier Robineau⁶ & Fabrice Carrat^{6,7}

Inserm

Cindy Lai¹³, Liza Belhadji¹³, Hélène Esperou¹³ & Sandrine Couffin-Cadiergues¹³

¹³Inserm, Paris, France.

Institut de Santé Publique

Jean-Marie Gagliolo¹⁴

¹⁴Aviesan, Inserm, Paris, France.

CEPH-Biobank

Hélène Blanché¹⁵, Jean-Marc Sébaoun¹⁵, Jean-Christophe Beaudoin¹⁵, Laetitia Gressin¹⁵, Valérie Morel¹⁵, Ouissam Ouili¹⁵ & Jean-François Deleuze¹⁵

¹⁵Fondation Jean Dausset-CEPH (Centre d'Etude du Polymorphisme Humain), CEPH-Biobank, Paris, France.

Unité des Virus Emergents

Stéphane Priet⁵, Paola Mariela Saba Villarroel⁵, Toscane Fourié⁵, Souand Mohamed Ali⁵, Abdenour Amroun⁵, Morgan Seston⁵, Nazli Ayhan⁵, Boris Pastorino⁵ & Xavier de Lamballerie⁵