



HAL
open science

Foraminifera and plastic pollution: Knowledge gaps and research opportunities

Vincent M.P. Bouchet, Laurent Seuront, Akira Tsujimoto, Julien Richirt, Fabrizio Frontalini, Masashi Tsuchiya, Misako Matsuba, Hidetaka Nomaki

► To cite this version:

Vincent M.P. Bouchet, Laurent Seuront, Akira Tsujimoto, Julien Richirt, Fabrizio Frontalini, et al.. Foraminifera and plastic pollution: Knowledge gaps and research opportunities. *Environmental Pollution*, 2023, 324, pp.121365. 10.1016/j.envpol.2023.121365 . hal-04061703

HAL Id: hal-04061703

<https://hal.science/hal-04061703>

Submitted on 7 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Foraminifera and plastic pollution: knowledge gaps and research opportunities**

2

3 Vincent M.P. Bouchet^{1,*}, Laurent Seuront^{1,2,3}, Akira Tsujimoto⁴, Julien Richirt,^{1,8} Fabrizio
4 Frontalini⁵, Masashi Tsuchiya⁶, Misako Matsuba⁷, Hidetaka Nomaki⁸

5 ¹ Univ. Lille, CNRS, Univ. Littoral Côte d'Opale, IRD, UMR8187, LOG, Laboratoire
6 d'Océanologie et de Géosciences, Station Marine de Wimereux, F 59000 Lille, France

7 ² Department of Marine Energy and Resource, Tokyo University of Marine Science and
8 Technology, 4-5-7 Konan, Minato-ku, Tokyo, 108-8477, Japan

9 ³ Department of Zoology and Entomology, Rhodes University, Grahamstown, 6140, South
10 Africa

11 ⁴ Faculty of Education, Shimane University, 1060 Nishikawatsu-cho, Matsue-
12 shi, Shimane, 690-8504, Japan

13 ⁵ Department of Pure and Applied Sciences, Università degli Studi di Urbino “Carlo Bo”,
14 61029, Urbino, Italy

15 ⁶ Research Institute for Global Change (RIGC), Japan Agency for Marine-Earth Science and
16 Technology (JAMSTEC), 2-15 Natsushima-cho, Yokosuka 237-0061, Japan

17 ⁷ Biodiversity Division, National Institute of Environmental Studies, 16-2 Onogawa, Tsukuba,
18 Ibaraki, 305-8506, Japan

19 ⁸ X-star, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15
20 Natsushima-cho, Yokosuka 237-0061, Japan

21

22 KEYWORDS. emerging pollutants, coastal habitats, biology, ecology, ecosystem functioning

23

24 ABSTRACT

25 Plastic has become one of the most ubiquitous and environmentally threatening sources of
26 pollution in the Anthropocene. Beyond the conspicuous visual impact and physical damages,
27 plastics both carry and release a cocktail of harmful chemicals, such as monomers, additives
28 and persistent organic pollutants. Here we show through a review of the scientific literature
29 dealing with both plastic pollution and benthic foraminifera (Rhizaria), that despite their
30 critical roles in the structure and function of benthic ecosystems, only 0.4% of studies have
31 investigated the effects of micro- and nano-plastics on this group. Consequently, we urge to
32 consider benthic foraminifera in plastic pollution studies *via* a tentative roadmap that includes
33 (i) the use of their biological, physiological and behavioral responses that may unveil the
34 effects of microplastics and nanoplastics and (ii) the evaluation of the indicative value of
35 foraminiferal species to serve as proxies for the degree of pollution. This appears particularly
36 timely in the context of the development of management strategies to restore coastal
37 ecosystems.

38

39 **Introduction**

40 Plastic pollution in marine environments has been reported as early as 1970 (Carpenter and
41 Smith Jr 1972) with an increasing interest until the mid-1980s (Bean 1987). It nevertheless
42 took nearly two additional decades to fully acknowledge the extent of the related
43 environmental issues (Thompson et al. 2004). Due to both the exponential increase in plastic
44 production (i.e., from 1.5 Mt in 1950 to 368 Mt in 2019, PlasticsEurope 2020) and the
45 mismanagement of plastic waste (one third of the plastic waste (32 million Mt) from 93% of
46 the world's population was classified as mismanaged in 2010, Borrelle et al. 2020), the 5 to 13

47 Mt of plastics discharged into the ocean every year, are expected to reach to up to 90 Mt/year
48 by 2030 under business-as-usual scenarios (Borrelle et al. 2020). As an inherently durable
49 material, plastic debris persist in the environment and are dispersed over long distances by
50 winds and water currents (Alimi et al. 2018). The potential threats related to plastic debris *via*
51 ingestion or leaching of contaminants (additives and persistent organic pollutants, see review
52 in Delaeter et al. 2022) have been identified as a major global conservation issue and a key
53 priority for research (Vegter et al. 2014, Harrison and Hester 2019, Seuront et al. 2022).

54 Plastic debris typically constantly degrade and fragment into smaller and smaller pieces
55 over time under UV radiation and wave actions to eventually form microplastics (<5 mm;
56 MPs) and nanoplastics (<100 nm; NPs). These small particles continuously sink towards the
57 seafloor where they are typically buried in unvegetated and vegetated sediments in subtidal
58 and intertidal coastal areas (Cozzolino et al. 2020). Noticeably, coastal ecosystems like
59 seagrass and macroalgal meadows (Cozzolino et al. 2020), mussel beds (Nicastro et al. 2022),
60 coral reefs (Reichert et al. 2022), tidal marshes and mangroves act as barriers against the
61 dispersal of MPs and NPs debris towards the marine environment (Martin et al. 2020). As
62 such, coastal habitats are major sinks for MPs and NPs, which enter long-term sequestration
63 in sediments (Martin et al. 2020). As a consequence, plastic pollution may interfere with the
64 key role played by living organisms in coastal ecosystems in mitigating the effects of climate
65 change by storing carbon dioxide (CO₂) in sediments (Taillardat et al. 2018). Yet, the effects
66 of plastic burial on the benthic biocoenosis of intertidal areas and coastal shallow-waters have
67 been largely overlooked.

68 Of the studies that have explicitly tested the effects of plastic on benthic species, relatively
69 few have attempted to consider MPs and NPs pollution and meiofaunal species which, for the
70 most part, are invisible to the naked eye, but are nevertheless crucial elements of coastal
71 benthic ecosystems (Schratzberger and Ingels 2018). In particular, meiobenthic foraminifera

72 are an important component of these coastal ecosystems. Amongst meiobenthic organisms
73 (<1 mm), benthic foraminifera (Rhizaria) may represent up to 50% of the benthic eukaryotic
74 biomass in coastal sediments (Moodley et al. 2000). They also contribute significantly to the
75 carbon and nitrogen biogeochemical cycles in coastal ecosystems (Jauffrais et al. 2019,
76 Choquel et al. 2021). They occupy a major trophic position through bacterivory, herbivory or
77 carnivory (Nomaki et al. 2008, Chronopoulou et al. 2019). They also actively participate in
78 the benthic ecosystem functioning *via* sediment mixing and bioirrigation, hence enhancing the
79 fluxes of particulate and dissolved elements at the sediment-water interface (Bouchet and
80 Seuront 2020, Deldicq et al. 2020, Deldicq et al. 2021a). In addition, foraminifera can
81 accumulate large quantities of nitrate in their cell and reduce it through denitrification under
82 low-oxygen conditions (Risgaard-Petersen et al. 2006, Piña-Ochoa et al. 2010, Choquel et al.
83 2021); their contribution to the nitrogen cycle being, for instance, up to 70% in the North Sea
84 (Piña-Ochoa et al. 2010) or ranging from 50% to 100% in Swedish fjords (Choquel et al.
85 2021). In the context of biomonitoring, benthic foraminifera have been shown to be good
86 sentinel of the health of benthic habitats (Armynot du Châtelet et al. 2004). Due to their
87 sensitivity to environmental changes, shifts in species composition of benthic foraminiferal
88 communities quickly reflect changes in environmental conditions (Schönfeld et al. 2012).
89 Hence, they are used as indicators of pollution (e.g. Alve 1995, Francescangeli et al. 2016,
90 Polovodova Asteman et al. 2015), such as oil spills (e.g. Morvan et al. 2004), drill cutting
91 (e.g. Mojtahid et al. 2006), element trace metals (e.g. Armynot du Châtelet et al. 2004), urban
92 sewage (e.g. Melis et al. 2019), and aquaculture (e.g. He et al. 2019). They are also considered
93 as accurate bioassay in ecotoxicological experiments which reported test (shell)
94 malformations (Le Cadre and Debenay 2006), alteration of the growth (Denoyelle et al.
95 2012), oxydative stress (Frontalini et al. 2018a), physiological changes (Losada Ros et al.
96 2020), proteom modifications (Ciacci et al. 2022) and changes in species composition of

97 foraminiferal communities in mesocosm (Frontalini et al. 2018b) in response to, respectively,
98 copper, oil-based drilling muds, element trace metals, cadmium and mercury exposure.

99 Because of the recognized fundamental role of foraminifera in the coastal benthic realm
100 (Moodley et al. 1998, LeKieffre et al. 2018), it is an important concern how MPs and NPs can
101 affect them. Specifically, plastic burial in benthic sediments may lead to the ingestion of MPs
102 or NPs and to long-term exposure of benthic foraminifera to a variety of additives and
103 chemicals leaching from plastic items in the sediment. Element trace metals, for instance,
104 which may accumulate onto and then leach from weathered plastic surface, are known to be
105 harmful for foraminifera (e.g. Le Cadre and Debenay 2006, Frontalini et al. 2018a, 2018b).
106 Noticeably, a wide range of physiological and ultrastructural impairments (e.g., involving
107 thickening of organic lining of test, mitochondrial degeneration, proliferation of degradation
108 vacuoles, lysosomes, residual bodies and the increased production of neutral lipids and of
109 reactive oxygen species) have been reported in benthic foraminifera exposed to element trace
110 metals (Frontalini et al. 2018a), as well as morphological abnormalities and decreased growth
111 rate (Le Cadre and Debenay 2006, Denoyelle et al. 2012). In addition, sediment anoxia
112 induced by plastic accumulation in the marine litter (Krause et al. 2020) may also contribute
113 to shape benthic foraminifera communities and promote low-oxygen-tolerant species. On the
114 basis of the above evidences, it seems legitimate to hypothesize that benthic foraminifera
115 would be highly affected by plastic pollution.

116 In this context, through a systematic review, this study quantifies the current state of
117 plastic pollution research on benthic foraminifera, presents key results, and summarizes
118 perspectives that warrant further study. Here, we highlight why foraminifera should be given
119 more emphasis in studies aiming to understand the effect of plastics on the benthic
120 biocoenosis of intertidal areas and coastal shallow-waters. We further suggest a way forward
121 toward consolidating research and policy efforts to better address the effect of plastic

122 pollution as a potential driver of benthic foraminiferal community structure and role in the
123 functioning of such key habitats.

124

125 **Methods**

126


127 The field of research dedicated to plastic pollution in benthic marine ecosystems is
128 currently one of the most prolific. A literature search was performed in the Web of Science
129 (last accessed on January, 13, 2023) using the words plastic (i.e. 'plastic*'), marine (i.e.
130 'marine') and benthic (i.e. 'benth*'). A total of 928 publications were returned. To evaluate
131 the research effort for the main taxa encountered in benthic coastal habitats, the literature
132 search was refined through the use of different combinations of keywords (i.e. 'bacteria*',
133 'crustacean*', 'diatoms*', 'macroalgae*', 'cnidaria*', 'mollusca*', 'echinodermata*',
134 'polychaeta', 'ciliata*', 'nematoda*', 'foraminifera', respectively, together with 'plastic*',
135 'marine' and/or 'intertidal' and/or 'coastal', and 'benth*'). The reference lists within these
136 studies were also scrutinized to identify additional relevant studies. For this opinion paper, we
137 only included English-written studies that assessed the impact of plastic pollution on the
138 above-mentioned taxa. A total of 925 studies were selected. This literature search allowed us
139 to evaluate the status of the research on benthic foraminifera and plastic pollution compared
140 to other biological components of benthic ecosystems.

141

142 **Plastic pollution and benthic foraminifera: the big void**

143 Among the studies that considered the effect of plastic pollution on marine organisms, a
144 vast majority (i.e., 81%) essentially targeted bacteria, algae and crustaceans (Fig. 1). Protists
145 were critically underrepresented (i.e., 2%) and amongst the 24 papers that considered them,
146 ten of these studies focused on benthic foraminifera; amongst which four observed benthic

147 foraminifera on floating and benthic plastic debris (Jorissen 2014, Finger 2017, Birarda et al.
148 2021, Subías-Baratau et al. 2022), four focused on the inclusion of plastic particles in
149 foraminiferal tests (Tsuchiya and Nomaki 2019, Birarda et al. 2021, Joppien et al. 2022a,
150 Romano et al. 2023) and only four *stricto sensu* monitored the effect of MPs or NPs (Ciacci et
151 al. 2019, Langlet et al. 2020, Birarda et al. 2021, Joppien et al. 2022b). A recent review
152 (Delaeter et al. 2022) showed that, in plastic leachates-related studies, only 0.64% studies
153 considered benthic foraminifera, while about 98% were on macro-invertebrates. Note that,
154 according to our literature survey, there is still no study considering the impact of plastic
155 pollution on planktic foraminifera. In short, regardless of coastal ecosystems, the assessment
156 of MPs and NPs impacts on foraminifera overwhelmingly lags behind other marine species.
157



158

159 **Figure 1.** Research effort (number of publications) considering the physical and toxic effects
160 of plastic pollution for the most relevant marine benthic taxa.

161

162 Drifting and benthic plastic debris may be colonized by benthic foraminifera (Jorissen
163 2014, Finger 2017, Birarda et al. 2021, Subías-Baratau et al. 2022). This may contribute to the
164 dispersion of benthic foraminifera, and their introduction outside their natural range as
165 observed on rafted plastic items with shallow-water foraminiferal fouling transported by
166 currents from Japan to Hawaii and North-America (Finger 2017).

167 Foraminifera have mainly four types of tests: those that assemble calcium carbonate crystals
168 constructed intracellularly and build tests (porcelaneous test), those that crystallized calcium
169 carbonate (mainly) extracellularly (hyaline), those that collect sediment particle in the habitat
170 for a test (agglutinated), and those with an organic membrane. Plastic particles were found to
171 be included in the test of hyaline (Joppien et al. 2022a) and agglutinated foraminifera
172 (Tsuchiya and Nomaki 2019, Birarda et al. 2021, Romano et al. 2023), and, as such, might
173 further be transferred to predators of foraminifera like gastropods, copepods, polychaetes,
174 decapods, polychaetes and fishes (Buzas and Carle 1979, Culver and Lipps 2003, Nomaki et
175 al. 2008). Inclusion of plastic particles into the test of benthic foraminifera may likely expose
176 them to the harmful effects of plastic additives; see Delaeter et al. (2022) for a specific review
177 on the impacts of plastic leachates on organisms.

178 Amongst the only four papers dealing with the *stricto sensu* influence of plastic on benthic
179 foraminifera (see details in Table 1), three dealt with the effect of particles (Ciacci et al. 2019,
180 Birarda et al. 2021, Joppien et al. 2022b) and two with the effects of additives (Langlet et al.
181 2020, Birarda et al. 2021). Noticeably, the exposition and ingestion of NPs of polystyrene (1
182 mg.L⁻¹) induced physiological stress by promoting the accumulation of neutral lipids and
183 enhanced reactive oxygen species production in *Ammonia parkinsoniana* (Ciacci et al. 2019).
184 In the presence of seawater-soaked polyethylene MPs (1 particle.mL⁻¹), the coral-reef species
185 *Amphistegina gibbosa* indifferently fed on MPs and on *Artemia* sp. nauplii (Joppien et al.

186 2022b). The heterotrophic behavior of this symbiont-bearing species was clearly altered;
187 hence, it had to be more reliant on autotrophy. Furthermore, *Rosalina bradyi*, *Textularia*
188 *bocki*, and *Lobatula lobatula* were found to colonize *in situ* a plastic bag (most likely
189 polyethylene) buried in the sediment (Birarda et al. 2021). The accumulation of protein beta-
190 sheets was observed in all three species, a typical feature observed during oxidative stress
191 cytotoxicity (Novak et al. 2019). The authors hypothesized that the accumulation of MPs
192 debris and additives both in the cytoplasm and agglutinated test (i.e., *T. bocki*) may alter the
193 biomineralization process in foraminifera. Laboratory experiments further showed that the
194 plastic additive bis-(2-ethylhexyl) phthalate (0.3 mg.L^{-1}) can be incorporated in the cytoplasm
195 of *R. globularis* (Birarda et al. 2021). However, the intertidal foraminifera *Haynesina*
196 *germanica* did not show any change in motion behavior nor in respiration rate when exposed
197 to leachates from virgin polypropylene pellets (20 and 200 mg of pellets.L⁻¹; Langlet et al.
198 2020).

199 These limited number of studies have partially explored the potential hazard of plastic
200 contamination of foraminifera in suggesting that MPs and NPs may interfere in the feeding
201 behavior of benthic foraminifera and generate physiological stress. However, they do not
202 allow to draw any robust conclusion, in particular because of the heterogeneity between
203 studies in terms of targeted foraminiferal species, type of plastic polymer, sizes and
204 concentrations of plastic particles, exposure time and measured parameters (Table 1), which
205 altogether further hampers any comparison among these works. Hence, we emphasize that our
206 knowledge is entirely insufficient for an integrated understanding of MPs and NPs effects on
207 foraminifera, as evidenced by the fact that the response is apparently dependent on the type of
208 plastic and foraminiferal species.

209 Given the ubiquity and extent of plastic pollution, the ever-growing number of studies
210 assessing the effect of plastics on a wide range of organisms and the aforementioned potential

211 ecological role of benthic foraminifera in key coastal habitats, it is striking that they have
212 been disregarded for so long, which further stresses the need to amend the *status quo*. The
213 lack of attention to a group as dominant in coastal sediments as foraminifera has received
214 from the scientific community is no longer justified neither on methodological nor ecological
215 grounds.

Table 1: Publications on the effects of plastic on benthic foraminifera: studied species, plastic types and methodological approaches.

Study	Species	Symbiont/ Chloroplast	Habitat	Type of study	Polymer/Additives	Particles size	Exposure	Concentration	Time of the exposure	Parameters
Birarda et al. 2021	<i>Rosalina globularis</i>	None	Temperate, Coastal, shallow-water	Experimental	Phtalate (DEHP)	/	Additive	0.3 mg.L ⁻¹	7 weeks	Oxydative stress
	<i>Rosalina bradyi</i> , <i>Textularia bocki</i> , <i>Cibicides lobatulus</i>	None	Temperate, Coastal, shallow-water	<i>In situ</i>	Polyethylene	Plastic bag	Particles and additive	/	/	Particle inclusion in the test, Oxydative stress
Ciacchi et al. 2019	<i>Ammonia parkinsoniana</i>	None	Temperate, Coastal, shallow-water	Experimental	Polystyrene	42 nm	Particles	1 mg.L ⁻¹	24h	Ultrastructural and physiological changes (oxydative stress)
Joppien et al. 2022b	<i>Amphistegina gibbosa</i>	Symbiont-bearing	Tropical, Coral reefs, shallow-water	Experimental	Polyethylene terephthalate	150-300 μ m	Particles	1 and 2 particles.mL ⁻¹	24h	Feeding behavior (<i>Artemia</i> sp. versus plastic particles)
Langlet et al. 2020	<i>Haynesina germanica</i>	Kleptoplastic	Temperate, Coastal, Intertidal	Experimental	Polypropylene	3.3-4.7 mm	Leachates (24h in seawater)	20 and 200 mL of pellets.L ⁻¹	10h	Motion behavior, Respiration

217

218 **Benthic foraminifera: culprits and victims of plastic pollution?**

219 It is generally recognized that MPs and NPs have a negative impact on organisms (e.g.
220 Delaeter et al. 2022), but this is not confirmed to be true for foraminifera. Ironically, benthic
221 foraminifera may not be only prime suspects but also guilty of sequestering both MPs and
222 NPs in sediment through their bioturbation activity that would actively transfer plastic
223 particles from the water-sediment interface to the sediment matrix. Noticeably, they are able
224 to burrow particles in the size range (10-125 μm) of MPs (Gross 2002). This may help to
225 isolate plastic particles from biologically active area (surface sediments) to less active area,
226 and promote fossilization of MPs and NPs. However, sub-sediments may also constitute a
227 reservoir where plastics are trapped and be further re-introduced in the food-web by benthic
228 organisms feeding in this compartment (Sandgaard et al. 2023). In addition, the inclusion of
229 plastic particles in the test of hyaline and agglutinated benthic foraminifera (Tsuchiya and
230 Nomaki 2019, Birarda et al. 2021, Joppien et al. 2022a, Romano et al. 2023) possibly
231 contributes to the transfer of plastic particles and additives through the food web, and to the
232 trapping of MPs and NPs in the sediment. In this specific context, benthic foraminifera
233 became their own victims as the burial of MPs and NPs is likely to expose them to ever
234 increasing concentrations of the countless additives and chemicals carried by plastic items.
235 We do not yet have evidence of whether they are culprits or victims, but at the very least,
236 benthic foraminifera that take up plastic provide an archive for the accumulation of MPs and
237 NPs in marine habitats, and this is *in fine* likely to provide additional evidence of how plastic
238 can be considered as a key stratigraphic indicator of the Anthropocene as a distinctive strata
239 component (Corcoran et al. 2015, Zalasiewicz et al. 2016, Dibley 2018). We hence advocate
240 for more comprehensive research programs (*i*) to understand the role of benthic foraminifera

241 in plastic trapping in sediments, and (ii) to improve our poor knowledge of the effect of
242 plastic particles and their leachates on benthic foraminifera.


243 The assessment of the pernicious non-lethal effects of MPs and NPs on organisms is a
244 challenge in the benthic coastal realm, particularly in minute organisms like benthic
245 foraminifera that represent an emerging model in coastal ecosystems. Despite their noticeable
246 small number relative to the available literature (see Fig. 1), the recent studies on the response
247 of benthic foraminifera to various aspects of plastic exposure (Ciacci et al. 2019, Langlet et al.
248 2020, Birarda et al. 2021, Joppien et al. 2022b) have stressed how critical it is to understand
249 why, how and under which environmental conditions MPs and NPs affect benthic
250 foraminifera, and we plea to address the following scientific questions (Fig. 2):

251 (1) What are the processes involved in the integration of plastic particles in foraminiferal test?
252 In particular, is this particle size- and/or concentration-dependent? Does this occur during
253 shell construction that includes calcification and agglutination or *via* ingestion? Does this
254 enhance the exposure of benthic foraminifera to plastic?

255 (2) Does sediment reworking by benthic foraminifera contribute to MPs and NPs
256 accumulation in coastal sediments?

257 (3) What are the potential harmful effects of plastic on benthic foraminifera at community
258 level, individual level, and cellular level?

259 (4) *In fine*, do the harmful effects of MPs and NPs alter the role of foraminifera in the benthic
260 ecosystem functioning?



261

262 Figure 2. Schematic of the questions to be tackled to understand (1) how much plastic
 263 particles are incorporated in foraminiferal test, (2) to what extent foraminifera may contribute
 264 to the trapping of MPs and NPs in coastal sediments and (3) the potential harmful effect of
 265 plastic on (a) species composition of foraminiferal communities (community level), (b)
 266 behavior (individual level), (c) cellular alterations and physiological changes (cellular level),
 267 and (4) the possible consequences on the benthic ecosystem functioning *i.e.* sediment mixing
 268 and bioirrigation. These questions are intended to shed light on the emerging opportunities of
 269 benthic foraminifera as proxies for the characterization of plastic pollution in coastal
 270 ecosystems. (Filled blue circles: MPs/NPs particles, Add: additives, POPs: persistent organic
 271 pollutants)

272

273 Addressing those questions can be achieved in future researches that should—and ideally
 274 concurrently—monitor the hazards of MPs and NPs exposure at different levels from cells to
 275 communities in the following roadmap:

276 (i) Cellular level: Underlying mechanisms behind the responses observed at the individual
277 level can be understood through the study of cellular alterations and physiological changes.
278 For instance, recent advances in omics technologies have provided new tools to understand
279 the effects of contaminants, which, in turn, may serve as biomarkers to be used in
280 environmental monitoring as early warning of pollution (Tomanek 2011; Ciacci et al. 2022).
281 In details, the response of benthic foraminifera to MPs and NPs particles and leachates are
282 now deeply accessible through the advance in transcriptomics (Titelboim et al. 2020),
283 proteomics (Stuhr et al. 2018) and microscopy (Frontalini et al. 2019). In addition, alterations
284 in the proteome can, for instance, have consequences on the fitness (i.e. reproductive success)
285 of an organism.

286 (ii) Individual level: While pernicious non-lethal effects of MPs and NPs may likely not be
287 easy to observe in communities, targeted experiments on foraminiferal individuals with
288 environmentally relevant concentrations and focusing on specific biological features known to
289 respond to contamination can address such limitations. Specifically, motion behavior is an
290 acknowledged keystone process behind the success of foraging, mate-finding and habitat
291 selection (Bowler and Benton 2005). Motion behavior is still, however, the forgotten child of
292 foraminiferal ecology, despite its increasingly recognized variability (Seuront and Bouchet
293 2015, Jauffrais et al. 2016) and role in bioturbation (Bouchet and Seuront 2020, Deldicq et al.
294 2020, Deldicq et al. 2021a). In addition, benthic foraminiferal behavior is affected in the
295 presence of a stressor (gravity, organic matter, temperature; Seuront and Bouchet 2015,
296 Jauffrais et al. 2016, Deldicq et al. 2021b) that may have further consequences on their search
297 for food and for the optimal microhabitat. In this context, investigating the behavioral changes
298 induced by MPs and NPs on benthic foraminifera will contribute to improve our
299 understanding of the effects of plastic pollution on their survival, may serve to predict
300 population-level effects.

301 We further stress that measurements suggested at points *i)* and *ii)* does not require long-term
302 incubation and recent studies confirmed they can be easily performed on foraminifera, which
303 may facilitate their implementation.

304 *(iii)* Community level: MPs and NPs may implicitly alter the quality of habitats inhabited by
305 benthic foraminifera, hence undermine the survival of some species, and ultimately constrain
306 the composition of foraminiferal communities. To reveal the response of community
307 composition to MPs and NPs, it is essential to monitor their spatial distribution through joint
308 *in situ* sampling surveys of MPs, NPs, and foraminiferal species. By applying spatial models,
309 *i.e.*, spatial varying coefficient model (Murakami et al. 2017, Matsuba et al. 2020), to these
310 spatial data, it may be possible to predict the distribution and composition of foraminiferal
311 communities in response to spatial patterns of MPs and NPs and their concentrations in
312 sediments, as well as ocean environmental factors (e.g. current velocity) that affect the
313 distribution of MPs and NPs. Such a modeling approach would be one way to determine how
314 MPs and NPs pollution, coupled with macro-environmental changes in the ocean, including
315 climate change, could affect foraminiferal communities. In addition, taking advantage of the
316 recent development of the use of benthic foraminifera to assess ecological quality statuses
317 (Alve et al. 2016, Bouchet et al. 2021, Parent et al. 2021), it may provide an opportunity to
318 develop an index of the degree of plastic pollution in coastal ecosystems sediment through the
319 identification of indicator species of the presence/absence of MPs and NPs. Finally, a better
320 understanding of benthic foraminiferal community response to plastic pollution will
321 eventually help to grasp their role in marine ecosystems functioning.

322

323 **Conclusion**

324 Benthic foraminifera offer a putatively endless sets of opportunities to tackle the question of
325 the potential harmful effects of plastic pollution, from the cellular to the community level. We
326 plea here to embrace the opportunity to see in these abundant but minute—though highly

327 neglected (see Fig. 1)—organisms a unique prospect to refine our understanding of the effects
328 of one of the main environmental threats of the “Anthropocene” on the structure and functions
329 at play behind the future of coastal ecosystems under climate change. *In fine*, foraminifera
330 will provide proxies of the history and the degree of plastic pollution on the benthic
331 ecosystem. Last but not the least, there is, to date, no study on the interaction between plastic
332 and planktic foraminifera, although they may be in contact with MPs and NPs in the water
333 column. This last aspect must also deserve more attention in the future.

334

335 AUTHOR INFORMATION

336 **Corresponding Author**

337 * Vincent M.P. Bouchet, vincent.bouchet@univ-lille.fr

338 **Author Contributions**

339 L.S., H.N. and V.M.P.B conceived the idea of the paper. L.S. and V.M.P.B. wrote the original
340 draft. The manuscript was written through contributions of all authors. All authors have given
341 approval to the final version of the manuscript.

342 **Funding Sources**

343 This work was financially supported by the Pierre-Hubert Curien Sakura program (project
344 number: 47041NL), implemented by the French Ministry for Europe and Foreign Affairs, the
345 French Ministry of Higher Education, Research and Innovation and the Japan Society for the
346 Promotion of Science (JPJSBP 120213211). C. Delaeter is funded by a PhD fellowship from
347 the Région Hauts-de-France and the University of Lille. J. Richirt is funded by a post-doctoral
348 fellowship from Japan Society for the Promotion of Science (P21729).

349 **Competing interests statement**

350 The authors have no competing interests to declare.

351 ACKNOWLEDGMENT

352 The authors are thankful to Nicolas Spilmont for his constructive comments on an earlier
353 version of this manuscript. This work was financially supported by the Pierre-Hubert Curien
354 Sakura program (project number: 47041NL), implemented by the French Ministry for Europe
355 and Foreign Affairs, the French Ministry of Higher Education, Research and Innovation and
356 the Japan Society for the Promotion of Science (JPJSBP 120213211). C. Delaeter is funded
357 by a PhD fellowship from the Région Hauts-de-France and the University of Lille. J. Richirt
358 is funded by a post-doctoral fellowship from Japan Society for the Promotion of Science
359 (P21729).

360 REFERENCES

- 361 Alimi, O. S., J. F. Budarz, L. M. Hernandez, and N. Tufenkji. 2018. Microplastics and
362 Nanoplastics in Aquatic Environments: Aggregation, Deposition, and Enhanced
363 Contaminant Transport. *Environmental Science and Technology* **52**:1704-1724.
- 364 Alve, E. 1995. Benthic foraminiferal responses to estuarine pollution: a review. *Journal of*
365 *Foraminiferal Research* **25**:190-203.
- 366 Alve, E., S. Korsun, J. Schönfeld, N. Dijkstra, E. Golikova, C. Hess, K. Husum, and G.
367 Panieri. 2016. Foram-AMBI: A sensitivity index based on benthic foraminiferal
368 faunas from North-East Atlantic and Arctic fjords, continental shelves and slopes.
369 *Marine Micropaleontology* **122**:1-12.
- 370 Armynot du Châtelet, E., J.-P. Debenay, and R. Soulard. 2004. Foraminiferal proxies for
371 pollution monitoring in moderately polluted harbors. *Environmental Pollution* **127**:27-
372 40.
- 373 Bean, M. J. 1987. Legal strategies for reducing persistent plastics in the marine environment.
374 *Marine Pollution Bulletin* **18**:357-360.
- 375 Birarda, G., C. Buosi, F. Caridi, M. A. Casu, G. De Giudici, L. Di Bella, D. Medas, C.
376 Meneghini, M. Pierdomenico, A. Sabbatini, A. Surowka, and L. Vaccari. 2021.
377 Plastics, (bio)polymers and their apparent biogeochemical cycle: An infrared
378 spectroscopy study on foraminifera. *Environmental Pollution* **279**:116912.

379 Borrelle, S. B., J. Ringma, K. Lavender Law, C. C. Monnahan, L. Lebreton, A. McGivern, E.
380 Murphy, J. Jambeck, G. H. Leonard, M. A. Hilleary, M. Eriksen, H. P. Possingham,
381 H. De Frond, L. R. Gerber, B. Polidoro, A. Tahir, M. Bernard, N. Mallos, M. Barnes,
382 and C. M. Rochman. 2020. Predicted growth in plastic waste exceeds efforts to
383 mitigate plastic pollution. *Science* **369**:1515-1518.

384 Bouchet, V. M. P., F. Frontalini, F. Francescangeli, P.-G. Sauriau, E. Geslin, M. V. A.
385 Martins, A. Almogi-Labin, S. Avnaim-Katav, L. Di Bella, A. Cearreta, R. Coccioni,
386 A. Costelloe, M. D. Dimiza, L. Ferraro, K. Haynert, M. Martinez-Colon, R. Melis, M.
387 Schweizer, M. Triantaphyllou, A. Tsujimoto, B. Wilson, and E. Armynot du Châtelet.
388 2021. Indicative value of benthic foraminifera for biomonitoring: assignment to
389 ecological groups of sensibility to total organic carbon of species from European
390 intertidal areas and transitional waters. *Marine Pollution Bulletin* **164**:112071.

391 Bouchet, V. M. P., and L. Seuront. 2020. Strength may lie in numbers: intertidal foraminifera
392 non-negligible contribution to surface sediment reworking. *Open Journal of Marine
393 Science* **10**:131-140.

394 Bowler, D., and T. Benton. 2005. Causes and consequences of animal dispersal strategies:
395 relating individual behaviour to spatial dynamics. *Biological Reviews* **80**:205-225.

396 Buzas, M. A., and K. J. Carle. 1979. Predators of foraminifera in the Indian River, Florida.
397 *Journal of Foraminiferal Research* **9**:336-340.

398 Carpenter, E. J., and K. L. Smith Jr. 1972. Plastics on the Sargasso sea surface. *Science*
399 **175**:1240-1241.

400 Choquel, C., E. Geslin, E. Metzger, H. Filipsson, N. Risgaard-Petersen, P. Launeau, M.
401 Giraud, T. Jauffrais, B. Jesus, and A. Mouret. 2021. Denitrification by benthic
402 foraminifera and their contribution to N-loss from a fjord environment.
403 *Biogeosciences* **18**:327-341.

404 Chronopoulou, P.-M., I. Salonen, C. Bird, G.-J. Reichart, and K. A. Koho. 2019.
405 Metabarcoding Insights Into the Trophic Behavior and Identity of Intertidal Benthic
406 Foraminifera. *Frontiers in Microbiology* **10**:1169.

407 Ciacci, C., M. V. Grimmelpont, I. Corsi, E. Bergami, D. Curzi, D. Burini, V. M. P. Bouchet,
408 P. Ambrogini, P. Gobi, Y. Ujiié, Y. Yshitani, R. Coccioni, J. M. Bernhard, and F.
409 Frontalini. 2019. Nanoparticle-Biological Interactions in a Marine Benthic
410 Foraminifer. *Scientific Reports* **9**:19441.

- 411 Ciacci, C., M. Betti, S. Abramovich, M. Cavaliere, and F. Frontalini. 2022. Mercury-induced
412 oxidative stress response in benthic foraminifera: an *in vivo* experiment on
413 *Amphistegina lessonii*. *Biology* **11**:960.
- 414 Corcoran, P. L., T. Norris, T. Ceccanese, M. J. Walzak, P. A. Helm, and C. H. Marvin. 2015.
415 Hidden plastics of Lake Ontario, Canada and their potential preservation in the
416 sediment record. *Environmental Pollution* **204**:17-25.
- 417 Cozzolino, L., K. R. Nicastro, G. I. Zardi, and C. B. de Los Santos. 2020. Species-specific
418 plastic accumulation in the sediment and canopy of coastal vegetated habitats. *Science*
419 *of The Total Environment* **723**:138018.
- 420 Culver, S. J., and J. H. Lipps. 2003. Predation on and by foraminifera. Pages 7-32 *in* P. H.
421 Kelley, M. Kowalewski, and T. H. Hansen, editors. *Predator-Prey Interaction in the*
422 *Fossil Record*. Kluwer Academic/Plenum Publisher, New York.
- 423 Delaeter, C., N. Spilmont, V. M. P. Bouchet, and L. Seuront. 2022. Plastic leachates: Bridging
424 the gap between a conspicuous pollution and its pernicious effects on marine life.
425 *Science of The Total Environment* **826**:154091.
- 426 Deldicq, N., L. Seuront, and V. M. P. Bouchet. 2021a. Inter-specific and inter-individual trait
427 variability matter in surface sediment reworking rates of intertidal benthic
428 foraminifera. *Marine Biology* **168**:101.
- 429 Deldicq, N., D. Langlet, C. Delaeter, G. Beaugrand, L. Seuront, V.M.P. Bouchet.
430 2021b. Effects of temperature on the behaviour and metabolism of an intertidal
431 foraminifera and consequences for benthic ecosystem functioning. *Scientific Reports*
432 **11**:4013.
- 433 Deldicq, N., L. Seuront, D. Langlet, and V. M. P. Bouchet. 2020. Behavioural traits of
434 intertidal foraminifera: understanding their role in bioturbation. *Marine Ecology*
435 *Progress Series* **643**:21-31.
- 436 Denoyelle, M., E. Geslin, F. J. Jorissen, L. Cazes, and F. Galgani. 2012. Innovative use of
437 foraminifera in ecotoxicology: A marine chronic bioassay for testing potential toxicity
438 of drilling muds. *Ecological Indicators* **12**:17-25.
- 439 Dibley, B. 2018. The technofossil : a memento mori. *Journal Of Contemporary Archaeology*
440 **5**:44-52.
- 441 Finger, K. 2017. Tsunami-generated rafting of foraminifera across the North Pacific Ocean.
442 *Aquatic Invasions* **13**:17-30.
- 443 Francescangeli, F., E. Armynot du Chatelet, G. Billon, A. Trentesaux, and V. M. P. Bouchet.
444 2016. Palaeo-ecological quality status based on foraminifera of Boulogne- sur-Mer

445 harbour (Pas-de-Calais, Northeastern France) over the last 200 years. *Marine*
446 *Environmental Research* **117**:32-43.

447 Frontalini, F., M.P. Nardelli, D. Curzi, A. Martin-Gonzalez, A. Sabbatini, A. Negri, M.T.
448 Losada, P. Gobbi, R. Coccioni, and J.M. Bernhard. 2018a. Benthic foraminiferal
449 ultrastructural alteration induces by heavy metals. *Marine micropaleontology* **138**:83-
450 89.

451 Frontalini, F., M. Greco, L. Di Bella, F. Lejzerowicz, E. Reo, A. Caruso, C. Cosentino, A.
452 Maccotta, G. Scopelliti, M. P. Nardelli, M. T. Losada, E. Armynot du Châtelet, R.
453 Coccioni, and J. Pawlowski. 2018b. Assessing the effect of mercury pollution on
454 cultured benthic foraminifera community using morphological and eDNA
455 metabarcoding approaches. *Marine Pollution Bulletin* **129**:512-524.

456 Frontalini, F., M. T. Losada, T. Toyofuku, J. Tyszka, J. Goleń, L. de Nooijer, B. Canonico, E.
457 Cesarini, Y. Nagai, U. Bickmeyer, T. Ikuta, R. Tsubaki, C. Besteiro Rodriguez, E.
458 Al-Enezi, S. Papa, R. Coccioni, J. Bijma, and J. M. Bernhard. 2019. Foraminiferal
459 ultrastructure: A perspective from fluorescent and fluorogenic probes. *Journal of*
460 *Geophysical Research: Biogeosciences* **124**:2823-2850.

461 Gross, O. 2002. Sediment interactions of foraminifera: implications for food degradation and
462 bioturbation processes. *Journal of Foraminiferal Research* **32**:414-424.

463 Harrison, R. M., and R. E. Hester. 2019. *Plastics and the environment*. Royal Society of
464 Chemistry, Cambridge UK.

465 He, X., T. F. Sutherland, J. Pawlowski, and C. L. Abbott. 2019. Responses of foraminifera
466 communities to aquaculture-derived organic enrichment as revealed by environmental
467 DNA metabarcoding. *Molecular Ecology* **28**:1138-1153.

468 Jauffrais, T., B. Jesus, E. Geslin, F. Briand, and V. Martin Jézéquel. 2016. Locomotion speed
469 of the benthic foraminifer *Ammonia tepida* exposed to different nitrogen and carbon
470 sources. *Journal of Sea Research* **118**:52-58.

471 Jauffrais, T., C. LeKieffre, M. Schweizer, E. Geslin, E. Metzger, J. M. Bernhard, B. Jesus, H.
472 Filipsson, O. Maire, and A. Meibom. 2019. Kleptoplastidic benthic foraminifera from
473 aphotic habitats: insights into assimilation of inorganic C, N and S studied with sub-
474 cellular resolution. *Environ Microbiol.* **21**:125-141.

475 Joppien, M., H. Westphal, A. Chandrapavan, M. Stuhr, and S. S. Doo. 2022a. Nanoplastic
476 incorporation into an organismal skeleton. *Scientific Reports* **12**:14771.

477 Joppien, M., H. Westphal, M. Stuhr, and S. S. Doo. 2022b. Microplastics alter feeding
478 strategies of a coral reef organism. *Limnology and Oceanography Letters* **7**:131-139.

- 479 Jorissen, F. J. 2014. Colonization by the benthic foraminifer *Rosalina (Tretomphalus)*
480 *concinna* of Mediterranean drifting plastics. In *Marine litter in the Mediterranean and*
481 *Black Seas*, Briand, F. Ed.; CIESM Workshop Monograph, Vol. 46; CIESM
482 Publisher, 2014; pp 87-95.
- 483 Krause, S., M. Molari, E. V. Gorb, E. Kossel, and M. Haeckel. 2020. Persistence of plastic
484 debris and its colonization by bacterial communities after two decades on the abyssal
485 seafloor. *Scientific Reports* **10**:9484.
- 486 Langlet, D., V. M. P. Bouchet, C. Delaeter, and L. Seuront. 2020. Motion behavior and
487 metabolic response to microplastic leachates in the benthic foraminifera *Haynesina*
488 *germanica*. *Journal of Experimental Marine Biology and Ecology* **529**:151395.
- 489 Le Cadre, V., and J.-P. Debenay. 2006. Morphological and cytological responses of *Ammonia*
490 (foraminifera) to copper contamination: implication for the use of foraminifera as
491 bioindicators of pollution. *Environmental Pollution* **143**:304-317.
- 492 Losada Ros, M.T., E. Al-Enezi, E. Cesarini, B. Canonico, C. Bucci, M.V. Alves Martins, S.
493 Papa, and F. Frontalini. 2020. Assessing the Cadmium Effects on the Benthic
494 Foraminifer *Ammonia cf. parkinsoniana*: An Acute Toxicity Test. *Water* **12**:1018.
495 <https://doi.org/10.3390/w12041018>
- 496 Martin, C., F. Baalkhuyur, L. Valluzzi, V. Saderne, M. Cusack, H. Almahasheer, P. K.
497 Krishnakumar, L. Rabaoui, M. A. Qurban, A. Arias-Ortiz, P. Masqué, and C. M.
498 Duarte. 2020. Exponential increase of plastic burial in mangrove sediments as a major
499 plastic sink. *Science advances* **6**:eaaz5593.
- 500 Matsuba, M., Y. Tanaka, T. Yamakita, Y. Ishikawa, and K. Fujikura. 2020. Estimation of
501 tsunami debris on seafloors towards future disaster preparedness: Unveiling spatial
502 varying effects of combined land use and oceanographic factors. *Marine Pollution*
503 *Bulletin* **157**:111289.
- 504 Melis, R., M. Celio, V. M. P. Bouchet, G. Varagona, M. Bazzaro, M. Crosera, and N.
505 Pugliese. 2019. Seasonal Response of benthic foraminifera to anthropogenic pressure
506 in two stations of the Gulf of Trieste (northern Adriatic Sea): the marine protected area
507 of Miramare versus the Servola water sewage outfall. *Mediterranean Marine Science*
508 **20**:120-141.
- 509 Mojtahid, M., F. Jorissen, J. Durrieu, F. Galgani, H. Howa, F. Redois, and R. Camps. 2006.
510 Benthic foraminifera as bio-indicators of drill cutting disposal in tropical east Atlantic
511 outer shelf environments. *Marine Micropaleontology* **61**:58-75.

512 Morvan, J., V. Le Cadre, F. Jorissen, and J.-P. Debenay. 2004. Foraminifera as potential bio-
513 indicators of the "Erika" oil spill in the Bay of Bourgneuf: field and experimental
514 studies. *Aquatic Living Resources* **17**:317-322.

515 Moodley, L., H. T. S. Boschker, J. J. Middelburg, R. Pel, P. M. J. Herman, E. de Deckere, and
516 C. H. R. Heip. 2000. Ecological significance of benthic foraminifera: ¹³C labelling
517 experiments. *Marine Ecology Progress Series* **202**:289-295.

518 Murakami, D., T. Yoshida, H. Seya, D.A. Griffith, and Y. Yamagata. 2017. A Moran
519 coefficient-based mixed effects approach to investigate spatially varying relationships.
520 *Spatial Statistics* **19**:68–89.

521 Nicastro, K. R., L. Seuront, C. D. McQuaid, and G. I. Zardi. 2022. Symbiont-induced
522 intraspecific phenotypic variation enhances plastic trapping and ingestion in biogenic
523 habitats. *Science of The Total Environment* **826**:153922.

524 Nomaki, H., N. O. Ogawa, N. Ohkouchi, H. Suga, T. Toyofuku, M. Shimanaga, T.
525 Nakatsuka, and H. Kitazato. 2008. Benthic foraminifera as trophic links between
526 phytodetritus and benthic metazoans: carbon and nitrogen isotopic evidence. *Marine*
527 *Ecology Progress Series* **357**:153-164.

528 Novak, S., T. Romih, B. Drasler, G. Birarda, L. Vaccari, P. Ferraris, S. Sorieul, M. Zieba, V.
529 Sebastian, M. Arruebo, S. B. Hocevar, A. Jemec Kokalj, and D. Drobne. 2019. The *in*
530 *vivo* effects of silver nanoparticles on terrestrial isopods, *Porcellio scaber*, depend on
531 a dynamic interplay between shape, size and nanoparticle dissolution properties.
532 *Analyst* **144**:488e497.

533 Parent, B., C. Barras, E. Bicchi, L. M. Charrieau, C. Choquel, É. Bénétteau, G. M. Maillet, and
534 F. J. Jorissen. 2021. Comparison of Four Foraminiferal Biotic Indices Assessing the
535 Environmental Quality of Coastal Mediterranean Soft Bottoms. *Water* **13**:3193.

536 Piña-Ochoa, E., S. Høgslund, E. Geslin, T. Cedhagen, N. P. Revsbech, L. P. Nielsen, M.
537 Schweizer, F. J. Jorissen, S. Rysgaard, and N. Risgaard-Petersen. 2010. Widespread
538 occurrence of nitrate storage and denitrification among Foraminifera and Gromiida.
539 *Proceedings of the National Academy of Science USA* **107**:1148-1153.

540 PlasticsEurope. 2020. *Plastics – The Facts 2020: An Analysis of European Plastics*
541 *Production, Demand and Waste Data*.

542 Polovodova Asteman, I., D. Hanslik, and K. Nordberg. 2015. An almost completed pollution-
543 recovery cycle reflected by sediment geochemistry and benthic foraminiferal
544 assemblages in a Swedish–Norwegian Skagerrak fjord. *Marine Pollution Bulletin*
545 **95**:126-140.

546 Reichert, J., A. L. Arnold, N. Hammer, I. B. Miller, M. Rades, P. Schubert, M. Ziegler, and T.
547 Wilke. 2022. Reef-building corals act as long-term sink for microplastic. *Global*
548 *Change Biology* **28**:33-45.

549 Risgaard-Petersen, N., A. M. Langezaal, S. Ingvaridsen, M. C. Schmid, M. S. M. Jetten, H. J.
550 M. Op den Camp, J. W. M. Derksen, E. Pina-Ochoa, S. P. Eriksson, L. P. Nielsen, N.
551 P. Revsbech, T. Cedhagen, and G. J. van der Zwaan. 2006. Evidence for complete
552 denitrification in a benthic foraminifera. *Nature* **443**:93-96.

553 Romano, E., E. Bergami, L. Di Bella, M. Bains, D. Berto, A. D'Ambrosi, M. DI Fazio, M.
554 Galli, L. Medegheni, C. Panti, C. Provenzani, F. Rampazzo, and M. C. Fossi. 2023.
555 First record of microplastic in the environmental matrices of a Mediterranean marine
556 cave (Bue Marino, Sardinia, Italy). *Marine Pollution Bulletin* **186**:114452.

557 Sandgaard, M.H., A. Palmqvist, A. Bour, S.N. Grønland, A. Hooge, H. Selck, A. Thit, and K.
558 Syberg, K. 2023. Sediment matters as a route of microplastic exposure: A call for
559 more research on the benthic compartment. *Frontiers in Marine Science* **9**:1100567.
560 doi: 10.3389/fmars.2022.1100567

561 Schönfeld, J., E. Alve, E. Geslin, F. Jorissen, S. Korsun, S. Spezzaferri, S. Abramovich, A.
562 Almogi-Labin, E. Armynot du Chatelet, C. Barras, L. Bergamin, E. Bicchi, V. M. P.
563 Bouchet, A. Cearreta, L. Di Bella, N. Dijkstra, S. Trevisan Disaro, L. Ferraro, F.
564 Frontalini, G. Gennari, E. Golikova, K. Haynert, S. Hess, K. Husum, V. Martins, M.
565 McGann, S. Oron, E. Romano, S. Mello Sousa, and A. Tsujimoto. 2012. The
566 FOBIMO (FORaminiferal BIo-MONitoring) initiative—Towards a standardised
567 protocol for soft-bottom benthic foraminiferal monitoring studies. *Marine*
568 *Micropaleontology* **94-95**:1-13.

569 Schratzberger, M., and J. Ingels. 2018. Meiofauna matters: The roles of meiofauna in benthic
570 ecosystems. *Journal of Experimental Marine Biology and Ecology* **502**.

571 Seuront, L., and V. M. P. Bouchet. 2015. The devil lies in details: new insights into the
572 behavioural ecology of benthic foraminifera. *Journal of Foraminiferal Research*
573 **45**:390-401.

574 Seuront, L., G. Zardi, M. Uguen, V. M. P. Bouchet, C. Delaeter, S. Henry, N. Spilmont, and
575 K. Nicastro. 2022. A whale of a plastic tale: a plea for interdisciplinary studies to
576 tackle micro- and nanoplastic pollution in the marine realm. *Science of The Total*
577 *Environment* **846**:157187.

578 Stuhr, M., B. Blank-Landeshammer, C. E. Reymond, L. Kollipara, A. Sickmann, M. Kucera,
579 and H. Westphal. 2018. Disentangling thermal stress responses in a reef-calcifier and
580 its photosymbionts by shotgun proteomics. *Scientific Reports* **8**:3524.

581 Subías-Baratau, A., Sanchez-Vidal, A., Di Martino, E., Figuerola, B. 2022. Marine biofouling
582 organisms on beached, buoyant and benthic plastic debris in the Catalan Sea. *Marine*
583 *Pollution Bulletin* **175**:113405.

584 Taillardat, P., D. A. Friess, and M. Lupascu. 2018. Mangrove blue carbon strategies for
585 climate change mitigation are most effective at the national scale. *Biology Letters*
586 **14**:20180251.

587 Thompson, R. C., Y. Olsen, R. P. Mitchell, A. Davis, S. J. Rowland, A. W. G. John, D.
588 McGonigle, and A. E. Russel. 2004. Lost at sea: where is all the plastic? *Science*
589 **304**:838.

590 Titelboim, D., S. Thangarjan, D. Raposo, R. Morard, M. Kucera, S. Ashckenazi-Polivoda, A.
591 Almogi-Labin, B. Herut, S. Manda, S. Abramovich, D. A. Gold, and U. Abdu. 2020.
592 The transcriptomic signature of cold and heat stress in benthic foraminifera—
593 Implications for range expansions of marine calcifiers. **35**:2679-2690.

594 Tomanek, L. 2011. *Environmental Proteomics: Changes in the Proteome of Marine*
595 *Organisms in Response to Environmental Stress, Pollutants, Infection, Symbiosis, and*
596 *Development. Annual review of Marine Science* **3**:373-399.

597 Tsuchiya, M., and H. Nomaki. 2019. Microplastic as a "micro"indicator for Anthropocene:
598 understanding the actual condition of microplastic pollution by using agglutinated
599 foraminifera. *in* TMS 2019, the Micropaleontological Society's Joint Foraminifera and
600 Nannofossil Meeting, Fribourg, Switzerland.

601 Vegter, A. C., M. Barletta, C. Beck, J. Borrero, H. Burton, M. L. Campbell, M. F. Costa, M.
602 Eriksen, C. Eriksson, A. Estrades, K. V. K. Gilardi, B. D. Hardesty, J. A. Ivar do Sul,
603 J. L. Lavers, B. Lazar, L. Lebreton, W. J. Nichols, C. A. Ribic, P. G. Ryan, Q. A.
604 Schuyler, S. D. A. Smith, H. Takada, K. A. Townsend, C. C. C. Wabnitz, C. Wilcox,
605 L. C. Young, and M. Hamann. 2014. Global research priorities to mitigate plastic
606 pollution impacts on marine wildlife. *Endanger. Species Res.* **25**:225-247.

607 Zalasiewicz, J., C. Waters, J. I. D. Sul, P. Corcoran, A. Barnosky, A. Cearreta, M. Edgeworth,
608 A. Gałuszka, C. Jeandel, R. Leinfelder, J. McNeill, W. Steffen, C. Summerhayes, M.
609 Wagreich, M. Williams, A. P. Wolfe, and Y. Yonan. 2016. The geological cycle of
610 plastics and their use as a stratigraphic indicator of the Anthropocene. *Anthropocene*
611 **13**:4-17.

612

613