



HAL
open science

Artificial light at night triggers slight transcriptomic effects on melatonin signaling but not synthesis in tadpoles of two anuran species

Morgane Touzot, Adeline Dumet, Jean Secondi, Thierry Lengagne, H el ene Henri, Emmanuel Desouhant, Claude Duchamp, Nathalie Mondy

► To cite this version:

Morgane Touzot, Adeline Dumet, Jean Secondi, Thierry Lengagne, H el ene Henri, et al.. Artificial light at night triggers slight transcriptomic effects on melatonin signaling but not synthesis in tadpoles of two anuran species. *Comparative Biochemistry and Physiology - Part A: Molecular and Integrative Physiology*, 2023, 280, pp.111386. 10.1016/j.cbpa.2023.111386 . hal-04062709

HAL Id: hal-04062709

<https://hal.science/hal-04062709>

Submitted on 7 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 **Artificial light at night triggers slight transcriptomic effects on melatonin signalling but**
2 **not synthesis in tadpoles of two anuran species**

3

4 Morgane Touzot^{1, 2, *, a}, Adeline Dumet^{2, a}, Jean Secondi^{2, 3}, Thierry Lengagne², H el ene Henri⁴,
5 Emmanuel Desouhant⁴, Claude Duchamp² and Nathalie Mondy²

6

7 ¹ Department of Ecoscience, Aarhus University, C.F. M ollers All e 4, Building 1120, DK-
8 8000 Aarhus C, Denmark.

9 ² Univ Lyon, Universit e Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-
10 69622, Villeurbanne, France.

11 ³ Univ Angers, 49045 Angers, France.

12 ⁴ Univ Lyon, Universit e Claude Bernard Lyon 1, CNRS, UMR5558 LBBE, Villeurbanne, F-
13 69622, France.

14 * Corresponding author: Department of Ecoscience, Aarhus University, C.F. M ollers All e 4,
15 Building 1120, DK-8000 Aarhus C, Denmark. Email: morgane.touzot@ecos.au.dk.

16 ^a contributed equally to the work.

17

18 Email addresses of each authors: morgane.touzot@ecos.au.dk, adeline.dumet@univ-lyon1.fr,
19 jean.secondi@univ-angers.fr, thierry.lengagne@univ-lyon1.fr, helene.henri-manijean@univ-lyon1.fr,
20 Emmanuel.Desouhant@univ-lyon1.fr, claudeduchamp@univ-lyon1.fr,
21 nathalie.mondy@univ-lyon1.fr.

22

23 ORCID

24 morgane.touzot@ecos.au.dk 0000-0002-0927-2817

25 adeline.dumet@univ-lyon1.fr 0000-0003-2249-1085

26 jean.secondi@univ-angers.fr 0000-0001-8130-1195

27 thierry.lengagne@univ-lyon1.fr 0000-0001-7840-6068

28 helene.henri-manijean@univ-lyon1.fr 0000-0001-9609-1952

29 Emmanuel.Desouhant@univ-lyon1.fr 0000-0003-0317-4463

30 claudio.duchamp@univ-lyon1.fr 0000-0001-9853-7110

31 nathalie.mondy@univ-lyon1.fr 0000-0002-4479-7053

32

33

34 **Highlights:**

- 35 • Melatonin-related gene expression tested at night under 2 light levels in 2 anurans
- 36 • Gene analysis based on respective species transcriptome and valid reference genes
- 37 • Preliminary description of a **nocturnal** rhythm in gene expression in tadpoles
- 38 • In both species, light pollution did not affect melatonin synthesis at gene level
- 39 • Melatonin signaling was slightly altered by light pollution during the night

40 **Abstract:**

41 The worldwide expansion of artificial light at night (ALAN) is acknowledged as a
42 threat to biodiversity through alterations of the natural photoperiod triggering the disruption
43 of physiological functions. In vertebrates, melatonin production during the dark phase can be
44 decreased or suppressed by nocturnal light as shown in many taxa. But the impact of **ALAN**
45 **at low intensity mimicking light pollution in periurban area** has never been investigated in
46 amphibians. We filled this gap by studying the impact of low ALAN levels on the expression
47 of genes related to melatonin synthesis and signaling in two anurans (agile frog, *Rana*
48 *dalmatina*, and common toad, *Bufo bufo*). Circadian expression of genes encoding enzymes
49 catalyzing melatonin synthesis (aralkylamine N-acetyltransferase, AANAT and
50 acetylserotonin O-methyltransferase, ASMT) or melatonin receptors (Mel1a, Mel1b and
51 Mel1c) was investigated using RT-qPCR after 23 days of exposure to control (< 0.01 lx) or
52 low ALAN level (3 lx) at night. We showed that the relative abundance of most transcripts
53 was low in late afternoon and early evening (06 pm and 08 pm) and increased throughout the
54 night in *R. dalmatina*. However, a clear and ample **nocturnal** pattern of target gene expression
55 was not detected in control tadpoles of both species. Surprisingly, a low ALAN level had little
56 influence on the relative expression of most melatonin-related genes. Only Mel1c expression
57 in *R. dalmatina* and Mel1b expression in *B. bufo* were affected by ALAN. This target gene
58 approach therefore provides experimental evidence that the melatonin signaling pathway was
59 slightly affected by low ALAN level in anuran tadpoles.

60

61

62 **Keywords:**

63 Amphibian; anthropogenic light pollution; gene expression; melatonin receptors; night-time

64 circadian rhythm

65 **1. Introduction**

66 The natural light/dark cycle leads to the rhythmic development of biological processes,
67 from molecules to organisms (Gaston et al., 2013). However, for the last decades, due to the
68 worldwide use and spread of artificial light at night (ALAN), a dramatic alteration of the
69 photoperiod regime beyond its natural range has been observed especially in urbanised areas
70 (Bennie et al., 2016; Cinzano et al., 2001; Gaston et al., 2014, 2012). In both terrestrial and
71 aquatic ecosystems (Davies and Smyth, 2018; Desouhant et al., 2019; Gaston et al., 2013;
72 Jechow and Hölker, 2019; Longcore and Rich, 2004; Secondi et al., 2017), ALAN was shown
73 to impact numerous behavioural and physiological mechanisms altering reproduction and
74 survival (Gaston et al., 2017, 2014; Longcore and Rich, 2004; Schroer and Hölker, 2017;
75 Touzot et al., 2020, 2019). ALAN is therefore likely to threaten population dynamic of wild
76 species (Sanders et al., 2015) and community composition (Davies et al., 2015, 2012; Sanders
77 et al., 2021).

78 Most of these alterations could be linked to a disruption of circadian timekeeping,
79 mainly regulated by melatonin. In most vertebrates, this hormone, which is only synthesized
80 during nocturnal periods in the absence of light, is defined as the chemical expression of
81 darkness (Reiter, 1991; Zhao et al., 2019). In addition to its function of relaying information
82 about changes in the photoperiod, melatonin plays important physiological roles in the
83 regulation of oxidative stress (Reiter et al., 2016; Tan et al., 2010), immune system (Calvo et
84 al., 2013; Carrillo-Vico et al., 2013; Pohanka, 2013) or sexual hormones (Mayer et al., 1997;
85 Tan et al., 2010). Melatonin is synthesized from tryptophan through four successive
86 enzymatic steps, two of them involving rate-limiting enzymes (Hardeland, 2008). Indeed,
87 melatonin production is limited by the availability of 1) aralkylamine N-acetyltransferase
88 (AANAT), that catalyses the transfer of the acetyl group of Acetyl-CoA to the primary amine
89 of serotonin, thereby producing CoA and N-acetylserotonin, and 2) N acetylserotonin O-

90 methyltransferase (ASMT) that catalyses the methylation of N-acetylserotonin to generate
91 melatonin. The rate of both enzymatic steps depends on the light/dark cycle and therefore
92 impacts melatonin synthesis, but differences in the circadian expression of these enzymes
93 exist between species (Borjigin et al., 1995; Coon et al., 2002; Ribelayga et al., 1999).
94 Melatonin actions are mediated through transmembrane G-protein-linked receptors, with two
95 subtypes identified in mammals (Mel1a and Mel1b) and three subtypes characterized in
96 amphibians, fishes, and birds (Mel1a, Mel1b and Mel1c). In vertebrates, these receptors exist
97 in many structures within the brains and in peripheral tissues, including kidney, liver,
98 intestine, blood, gills and skin (Confente et al., 2010; Hardeland, 2008; Jones et al., 2012).
99 The expression of these receptors also varies along the light/dark cycle (Li et al., 2013).
100 However, slight differences in circadian gene expression pattern exist depending on subtypes,
101 tissues and species (Confente et al., 2010; Jones et al., 2012; Maugars et al., 2020; Rada and
102 Wiechmann, 2006), which could partly be explained by the fact that to exert full physiological
103 action, melatonin receptors may form functional homo or heterodimers (Ayoub et al., 2002).

104 Amphibian species, which are among the most endangered vertebrates today
105 (International Union for Conservation of Nature, IUCN), are particularly exposed to ALAN
106 due to the fact that wetlands are subjected to light pollution (Secondi et al., 2017). Among
107 amphibians, anurans are the most likely to be affected by ALAN (Buchanan, 2006). In
108 anurans, ALAN was shown to increase stress level (Forsburg et al., 2021), alter larval
109 development (Dananay and Benard, 2018), reduce nocturnal activity affecting daily energy
110 allocation (Touzot et al., 2019), impact breeding behaviour and depress male reproductive
111 success (Touzot et al., 2020). Moreover, in adult cane toads, *Rhinella marina*, ALAN reduced
112 and delayed nocturnal activity peak and decreased body mass gain (Secondi et al., 2021).
113 When ALAN intensity is sufficient to alter photoperiod perception, in early stages of
114 development, tadpoles with a relatively immature circadian system and incomplete

115 differentiation of organs and tissues (Gosner, 1960) might be particularly sensitive to rhythm
116 disruption through ALAN (Fonken and Nelson, 2016).

117 Although amphibians are an ecological relevant model group to focus on the effects of light
118 pollution studies, consequences of such pollution on melatonin synthesis and signalling have
119 not been investigated yet (Grubisic et al., 2019). The lack of data in amphibians may be
120 explained by the difficulty in measuring melatonin production in non-model animal species.
121 Among the classical techniques for measurements, radio-immuno assay (RIA) and more
122 recently enzyme immunoassay (EIA) can be used (e.g., Delgado and Vivien-Roels, 1989;
123 Edwards and Pivorun, 1991; Jessop et al., 2014; Wright et al., 2006; Wright and Alves, 2001
124 Bayarri et al., 2004; Rebollar et al., 1999), but they consume a large amount of plasma
125 samples. Conversely, nowadays such invasive blood sampling on adults is not feasible
126 because it involves injury risk on protected species, and the small size of tadpoles precludes
127 RIA analysis. Since proteins, such as enzymes and receptors, are encoded by genes,
128 assessment of gene expression by quantitative reverse transcription polymerase chain reaction
129 (RT-qPCR) appears as a relevant method, with the advantages of being high throughput,
130 repeatable and allowing the assay of multiple genes per sample, thus limiting animal sample
131 size (Ginzinger, 2002).

132 Thus, this study focused on the influence of ALAN on the expression of genes encoding
133 enzymes catalyzing melatonin synthesis or its receptors in two species of amphibians.
134 Because *Ranidae* and *Bufo* are two of the most widespread and species families of
135 amphibians worldwide (<https://amphibiaweb.org/lists/>), we selected *Rana dalmatina* and *Bufo*
136 *bufo* to investigate whether a pattern of common consequences exists in two species of
137 anurans. During the breeding season, adults of these species are frequently found in urban and
138 peri-urban areas with wetlands (Beebee, 1979), subjected to light pollution, and laying eggs,
139 themselves exposed to ALAN. In this study, we aimed at (a) validating a RT-qPCR method

140 valuable on both anuran species, and (b) assessing by RT-qPCR the nocturnal expression of
141 five target melatonin-related genes (AANAT, ASMT, Mel1a, Mel1b and Mel1c) on tadpoles
142 of both species exposed during one third of their aquatic life to low light illuminances: <0.01
143 lx, which correspond to a control light level, and 3 lx, representing levels encountered in areas
144 hosting amphibians (Secondi et al., 2017).

145 **2. Methods**

146 *2.1. Animal collection, housing conditions*

147 Six fragments of approximately 20 eggs each were randomly sampled from a clutch
148 of both species, *R. dalmatina* and *B. bufo*, in March during the breeding season *in natura* in
149 Saint André de Corcy (46°N, 5°E) and Sainte-Croix (45°N, 5°E), France, respectively. Those
150 sites were chosen as they are located approximately 20 km from Lyon, reducing ALAN
151 exposure *in natura*. Upon arrival at the animal care facility (EcoAquatron, University of
152 Lyon), the 6 fragments per species were individually placed in an aquarium (32 cm X 17.5 cm
153 X 18.5 cm) containing 7.4 L of dechlorinated and oxygenated water. Each fragment was
154 immediately randomly assigned to one of the two experimental treatments (control, *i.e.*, <0.01
155 lx, and ALAN night light treatment, *i.e.*, 3 lx), resulting in 3 aquariums per experimental
156 treatment and per species. We used the classification of Gosner (Gosner, 1960) to distinguish
157 the different stages of tadpoles. From stage 25 of Gosner, the stage at which independent
158 feeding is possible, to the end of the experiment, tadpoles were fed every two days with
159 boiled organic green salad. Each week, one-third of the water of the aquarium was renewed
160 with dechlorinated and oxygenated water. Ambient temperature and relative humidity were
161 kept constant during the whole experiment at 16.0 ± 0.3 °C and $55.3 \pm 0.3\%$, respectively
162 (mean \pm SEM). After egg hatching, tadpole mortality was monitored and was equal to zero
163 during our experiments.

164 2.2. *Light treatments*

165 To mimic natural photoperiod, all aquariums were kept under a photoperiod
166 corresponding to the latitude of Lyon (45°N) at the date of the experiment (at the beginning of
167 the experiment photoperiod of 12.5L/11.5D and at the end of the experiment photoperiod of
168 14L/10D). During the daytime, illuminance provided by light tubes (Philips Master TL-D 58
169 W/865 and Exo Terra Repti Glo 2.0, 40 W T8) for all aquariums was 1906 ± 159 lx (mean \pm
170 SEM). The spectrum of our daylight is characteristic of fluorescent lamps (Touzot et al.,
171 2021) and is recommended for the housing of amphibians in animal care facility. Until
172 hatching, during night-time, individuals remained under dark conditions (<0.01 lx). From the
173 day of hatching, *i.e.*, 5 days since the start of our study under our conditions, the aquariums
174 were exposed to their respective light treatment at night. During night-time, to simulate
175 ALAN, we used white light-emitting diode (LED) ribbons (white cold Light Plus, 6000-6500
176 K, 14 W, 60 LED/m) whose spectrum is different from our daylight (Touzot et al., 2021).
177 White LEDs were chosen because they are increasingly used for street lighting worldwide
178 (Falchi et al., 2016). A LED ribbon of 95 cm (57 LEDs) combined with a light diffuser was
179 suspended horizontally 41.5 cm above the bottom of the aquariums. Each LED ribbon was
180 connected to a dimmer (manual dimmer, 12 V max, 8 A) and a laboratory power supply (15
181 V/DC max, 3 A), which allowed illuminance to be set for each light treatment. The aquariums
182 belonging to one light treatment were isolated from the others with tarpaulins to avoid light
183 spilling over from another treatment. At night, the control group was exposed to dark
184 conditions, *i.e.*, <0.01 lx, hereafter called “control-group”, corresponding to the illuminance
185 of a sky under clear conditions with a quarter moon (Gaston et al., 2013). The experimental
186 group was exposed to 3 lx, hereafter called “ALAN-group”, which corresponds to levels
187 encountered in areas hosting amphibians (Secondi et al., 2017) and in urban parks (Segrestin
188 et al., 2021). The daylight and 3 lx illuminances were measured with a luxmeter (Illuminance

189 meter T-10A, Konica Minolta). Illuminances for the control-group were set using a highly
190 sensitive light meter (Sky Quality Meter SQM-L, Unihedron). Consequently, to compare
191 illuminances, SQM-measured values were converted into lx according to the relationship
192 curve between SQM and lx measurements (Touzot *et al.*, 2020), as lx is the main unit used in
193 ALAN studies (Longcore and Rich, 2004). Illuminances were measured at the water surface
194 of each aquarium (24 cm from the LED) and checked every week (0.005 ± 0.001 lx for the
195 control-group and 3.26 ± 0.75 lx for the ALAN-group, mean \pm SEM).

196 2.3. *Experiment*

197 When tadpoles reached development stage 31 of Gosner, corresponding to 23 days of
198 experiment, for each species, 5 tadpoles of each treatment were randomly sampled every two
199 hours between 06 pm and 08 am, resulting in 8 timepoints (random sampling on the 3
200 aquariums of each light treatment). For each species, this resulted in 80 tadpoles sampled.
201 Development stage 31 of Gosner was chosen, because it is an easily identifiable stage as the
202 foot is paddle-shaped at this stage only Gosner. Development stage 31 also allowed long-term
203 exposure to ALAN, while being distant enough from metamorphosis, *i.e.*, stage 46 of Gosner,
204 the ultimate stages of development during which thyroid hormones are strongly produced to
205 trigger many functional and morphological changes in the organisms (Galton, 1992;
206 Kikuyama *et al.*, 1993). Tadpoles were individually sampled and immediately placed in liquid
207 nitrogen and then stored at -80°C until molecular analyses. As the main source of melatonin
208 synthesis is located in the pineal gland and eyes in amphibians (Delgado and Vivien-Roels,
209 1989), RNA extractions were realised on tadpole head, after tissues collection using a section
210 behind the eyes.

211 2.4. *Molecular analyses*

212 2.4.1. *Target and housekeeping gene selection*

213 We examined the relative abundance of the transcripts of five target genes related to
214 melatonin; two enzymes: AANAT and ASMT, and three receptors: Mel1a, Mel1b and Mel1c.
215 Finding appropriate housekeeping genes was a prerequisite to limit the biases related to the
216 daily variation in their expression. The majority of studies in amphibians have focused on the
217 stability of housekeeping genes during development (Dhorne-Pollet et al., 2013; Lou et al.,
218 2014), but not at the circadian scale. Since we were interested in temporal monitoring of gene
219 expression, housekeeping genes selected for normalization should have a constant level of
220 expression over 24h (Hellemans et al., 2007). In a first step, using genes selected from
221 developmental studies in amphibians (Dhorne-Pollet et al., 2013; Lou et al., 2014), *i.e.*, rpl8
222 (60S ribosomal protein L8) and tbp (TATA-binding protein) or circadian rhythm studies in
223 rodents (Kamphuis et al., 2005), *i.e.*, β -actin and GluR2 (AMPA-type glutamate receptor), we
224 performed an analysis of gene expression stability overnight. For each target and
225 housekeeping gene, existing amphibian nucleotide sequences have been searched in the NCBI
226 Nucleotide database (Appendix Table A) and blasted on the respective *de novo* transcriptome
227 of each species (see Touzot et al., 2021 for *B. bufo* transcriptome and see Appendix Text A
228 for *R. dalmatina* transcriptome). After validation of the gene sequences (identity percentages
229 greater than 70% and minimum length of 200 bp), specific primer pairs of 20 bp were created
230 for both species based on their respective *de novo* transcriptome, except for *R. dalmatina* for
231 which GluR2 has not been tested, as it had not been chosen as housekeeping gene in *B. bufo*
232 (Table 1).

233

234 **Table 1**

235

236 2.4.2. RNA extraction and cDNA synthesis

237

238 Total RNA was extracted by adding 500 μ L of TRI Reagent (Molecular Research
239 Center MRC, TR118) to the 160 samples and by homogenising tissues with a tissue lyser
240 during 2 minutes at 30Hz (Retsch, MM200) and then placing them immediately on ice. All
241 remaining steps were carried out according to the manufacturer's protocol (Molecular
242 Research Center MRC, TR118). Then, the RNA was treated with Turbo DNase enzyme
243 (Turbo DNA free kit, Invitrogen, AM1907) and assayed by fluorescence using a Qubit
244 fluorometer (Qubit® RNA HS Assay Kits Molecular Probes, Invitrogen, Q32855). RNA
245 integrity was assessed using a Bioanalyzer 2100 with Agilent RNA 6000 Nano kit (Agilent,
246 5067-1511). Total RNA was reverse transcribed into cDNA using the transcriptase
247 SuperScript® III (Invitrogen, 18080-044) according to the manufacturer' protocol and using
248 oligo(dt) to isolate mRNAs. cDNA obtained were then treated with Ribonuclease H (RNase
249 H, Invitrogen, AM1907).

250 2.4.3. RT-qPCR realisation

251 RT-qPCR was performed on the 160 samples with Universal SYBR Green Supermix
252 (BIORAD, 1725274) according to the manufacturer's protocol. Samples were run in
253 duplicates on a thermocycler CFX96 (BIORAD) with the following program: initial
254 denaturation and enzyme activation for 30 s at 95 °C, followed by 40 cycles of 10 s at 95 °C
255 and 30 s at 60 °C. A melting curve was then created for validation of the PCR products.
256 Quantification cycle (Cq) values were obtained and analyzed with Biorad CFX Manager 3.0.
257 software, using the regression method. Cq values were considered valid if duplicate
258 measurements (Δ Cq) differed by less than 1 cycle, otherwise, both Cq values were removed
259 from the analysis. For each pair of primers, the amplification efficiency was calculated from a
260 standard curve, generated from 6 pre-dosed amplicon range points with a dilution factor of 10,
261 and was comprised between 90 and 110%. For technical reasons, mainly related to an assay

262 plate effect, some samples had a ΔCq higher than 1 and had to be removed from the analysis,
263 reducing the sampling size from 80 to 77 samples for AANAT, ASMT and Mel1b and to 75
264 samples for Mel1c in *R. dalmatina*, and from 80 samples to 65 samples for AANAT, ASMT,
265 Mel1a and Mel1c and to 63 samples for Mel1b in *B. bufo*.

266 2.4.4. Selection of housekeeping genes, and relative mRNA quantification and analysis

267 The stability of the four housekeeping genes was analysed using software programs
268 GeNorm v3.5 (Vandesompele et al., 2002) and Normfinder v5 (Andersen et al., 2004), in
269 order to select genes showing the most stable expression necessary for normalization
270 (Hellemans et al., 2007). The two software programs evaluate stability slightly differently.
271 GeNorm estimates the expression stability of a gene, considered as a housekeeping, by
272 calculating the average variation of the expression levels ratios between this gene and another
273 one, also considered as housekeeping gene, in all samples (Vandesompele et al., 2002). Genes
274 with the lowest average variation (*i.e.*, lowest calculated stability value) have the most stable
275 expression. Stability in GeNorm is estimated for the entire dataset without distinction between
276 factors (*i.e.*, light treatment or timepoint). On the other hand, Normfinder estimated the
277 expression stability of a target gene using the Cq values and allows to consider factors effect
278 (*i.e.*, light treatment or timepoint). Target genes were normalised with the selected
279 housekeeping genes using *easyqPCR* v1.21.0 package (Hellemans et al., 2007; Le Pape,
280 2013), which allows the target genes to be normalized by several housekeeping genes. The
281 normalisation of each target gene was done with the three selected housekeeping genes
282 measured at the same timepoint. Specifically, it converts the Cq values of each sample into a
283 normalized relative quantity, considering the variation in Cq between genes, the efficiency of
284 PCR and the number of housekeeping genes. Relative target gene expression was analysed
285 with a generalised additive model (GAM) as it is the standard and most relevant model for

286 analysing data with cyclicity (Faraway, 2006; Zuur et al., 2009). The GAM analysis was
287 conducted using the *mgcv* v1.8-28 package (Wood, 2006) with a gamma distribution and
288 inverse function, with timepoint, light treatment and their interaction as explanatory factors.
289 We selected the best models based on the lowest Akaike's Information Criterion (AIC) when
290 the difference in AIC score between two models was higher than 2. A significance threshold
291 of 0.05 was adopted for all statistical analyses. All analyses and graphics were performed with
292 R software v3.5.3 ("R Core Team," 2018).

293 2.5. Ethical note

294 The sampling of *R. dalmatina* and *B. bufo* clutches was authorized by the Préfecture
295 de l'Ain (DDPP01-20-369 and 16-271) and by the French government in accordance with the
296 ethical committee of the University of Lyon (APAFIS#27760-2020102115402767 v2). The
297 animal care structure "EcoAquatron" (University of Lyon) received an agreement of
298 veterinary services (approval DSV 692661201). At the end of the experiment, all the
299 remaining tadpoles were released on the capture sites.

300 3. Results

301 3.1. Housekeeping gene selection

302 The analysis of housekeeping gene expression stability with GeNorm and Normfinder
303 recommended to select three housekeeping genes in *B. bufo* (Table 2). β -actin, rpl8 and tbp
304 presented the lowest variation in relative expression (*i.e.*, lowest stability value) for both
305 software while GluR2 always showed the highest variability (see Table 2 and Appendices
306 Table B and Table C for details). Therefore, GluR2 has been excluded from our housekeeping
307 gene selection. The three housekeeping genes selected for *B. bufo* also showed a low variation
308 in expression in *R. dalmatina* (Table 2). Therefore, for both species we selected β -actin, rpl8
309 and tbp as housekeeping genes.

310

311 **Table 2**

312

313 *3.2. Influence of timepoint and ALAN on melatonin-related gene expression*

314 *3.2.1. In R. dalmatina*

315 In *R. dalmatina*, the GAM analyses revealed that the relative abundance of transcripts
316 encoding AANAT ($F_{1,76} = 3.64, p = 0.028$) and ASMT ($F_1 = 13.32, p < 0.001$) was affected
317 by timepoint (see Table 3 and Appendix Table D for details). A low relative abundance of
318 transcripts encoding AANAT and ASMT was observed in the late afternoon and early
319 evening (06 pm and 08 pm, respectively), while it increased throughout the night, nearly
320 doubling in the early morning and 2 hours after the beginning of the day period (06 am and 08
321 am, respectively) (Fig. 1a and Fig. 1b). Although all melatonin receptor subtypes have been
322 assayed in *R. dalmatina*, only the relative abundance of transcripts encoding Mel1b and
323 Mel1c have been analysed, as the relative abundance of transcripts encoding Mel1a was too
324 low to be exploited (Cq values between 30 and 35, *i.e.*, within the detection limit of RT-
325 qPCR). Timepoint ($F_1 = 21.24, p < 0.001$) had a significant effect on the relative abundance
326 of transcripts encoding Mel1b (see Table 3 and Appendix Table D for details). We observed a
327 progressive increase throughout the night to reach its highest value 2 hours after the beginning
328 of the day period (08 am) (Fig. 1c). Light treatment significantly affected the relative
329 abundance of transcripts encoding for melatonin-related gene through an interaction with
330 timepoint in Mel1c (see Table 3 and Appendix Table D for details). In control conditions, the
331 relative abundance of transcripts encoding Mel1c varied with timepoint ($F_{1,32} = 7.72, p =$
332 0.002). Indeed, the relative abundance of transcripts encoding Mel1c was weak in the late
333 afternoon and early evening (06 pm and 08 pm) and increased throughout the night, more than
334 doubling in the early morning and 2 hours after the beginning of the day period (06 am and 08

335 am) (Fig. 1d). Exposure to ALAN inhibited night-time variation of the relative abundance of
336 transcripts encoding for Mel1c ($F_1 = 0.36, p = 0.549$), as it remained statistically stable
337 throughout the night (Fig. 1d).

338

339 **Table 3**

340

341 **Figure 1**

342

343 *3.2.2. In B. bufo*

344 The GAM analyses showed that the relative abundance of transcripts encoding for
345 Mel1a was significantly affected by timepoint ($F_1 = 6.53, p = 0.013$) (see Table 4 and
346 Appendix Table E for details). The relative abundance of transcripts encoding for Mel1a was
347 low in the late afternoon and early evening (06 pm and 08 pm) and progressively increased
348 throughout the night, to its highest value 2 hours after the beginning of the day period (08 am)
349 (Fig. 2c). Light treatment significantly affected the relative abundance of transcripts encoding
350 for melatonin-related gene through an interaction with timepoint in Mel1b in *B. bufo* (see
351 Table 4 and Appendix Table E for details). Exposure to ALAN induced night-time increase
352 ($F_1 = 7.15, p = 0.009$) in the relative abundance of transcripts encoding for Mel1b (Fig. 2d).
353 While in control conditions, this parameter remained statistically stable throughout the night
354 ($F_{3,51} = 1.06, p = 0.387$). However, the relative abundance of transcripts encoding for
355 AANAT, ASMT and Mel1c (Fig. 2a, 2b and 2e, respectively) was not affected by timepoint
356 (see Table 4 and Appendix Table E for details).

357

358 **Table 4**

359

360 **Figure 2**

361

362 **4. Discussion**

363 In this study, we developed a RT-qPCR method to investigate the nocturnal changes in
364 the relative expression of two melatonin-synthesis related enzymes (AANAT and ASMT) and
365 three melatonin-related receptors (Mel1a, Mel1b and Mel1c) in tadpoles of two non-model
366 amphibian species under low levels of artificial light at night. Despite variations in relative
367 abundance of most transcripts during the night in *R. dalmatina*, a marked night-time circadian
368 pattern could not be demonstrated in control tadpoles of both species. Contrary to our working
369 hypotheses and surprisingly, a low level of ALAN had little influence on most melatonin-
370 related genes. Only Mel1c expression in *R. dalmatina* and Mel1b expression in *B. bufo* were
371 affected by low levels of ALAN at night.

372 *4.1. Robust molecular methodology based on use of respective transcriptomes and* 373 *validated housekeeping gene stability*

374 For our RT-qPCR analysis, we gave special attention to the methodological issue to
375 ascertain the correct assessment of relative gene expression in tadpoles. Firstly, considering
376 that our study was conducted on two non-model species, but which are representative of the
377 most widespread and species families of amphibians worldwide, primers were designed with
378 the respective transcriptome of each species, thus favoring sequence identity and detection of
379 the target mRNAs. Each target gene sequences were validated by identity percentages greater
380 than 70% when compared to sequences obtain in different taxa (amphibian, fish, birds, and

381 mammals) and when the minimum length of the sequence was of 200 bp. In addition, to
382 confirm the correct amplification of the sequences corresponding to our target genes, gene
383 amplification was checked by Sanger sequencing and melting curves confirmed the specificity
384 of amplification. Secondly, finding appropriate housekeeping genes for normalization was a
385 prerequisite to limit the biases related to the daily variation in gene expression. Indeed, the
386 normalization of relative quantities of target genes with housekeeping genes relies on the
387 assumption that these housekeeping genes are stably expressed across all tested samples,
388 stability which is most of the time lacking in the studies (Hellemans et al., 2007). In our study,
389 we tested and verified the stable expression across circadian scale and light treatments of
390 genes commonly used for amphibians, such as *rpl8* and *tbp* (Dhorne-Pollet et al., 2013; Lou et
391 al., 2014) or used in circadian rhythm studies in rodents, such as β -actin and *GluR2*
392 (*Kamphuis et al., 2005*). Among the genes tested, we selected those with the most stable
393 expression across circadian scale and light treatments and opted for a normalization based on
394 multiple housekeeping genes, known to produce more reliable data (Hellemans et al., 2007).
395 Our approach highlights the need to validate housekeeping genes for each biological model
396 and specific experimental conditions. This step is crucial especially when the expression of
397 the targeted genes varies with time. Considering all these methodological parameters, our
398 robust molecular biology methodology gave confidence about the results obtained in this
399 study.

400 *4.2. Lack of marked effects of low ALAN levels on melatonin synthesis*

401 In a recent study, we showed that tadpole global gene expression, through
402 transcriptomic approach, was largely impacted by low levels of ALAN illuminance (Touzot et
403 al., 2021). In the present study, we used target-specific approaches to decipher potential
404 effects of ALAN on melatonin-related gene expression, but these effects were weak and
405 complex. Indeed, we did not detect a marked impact of ALAN on the relative expression of

406 genes encoding melatonin synthesis enzymes. This unexpected result weakens the hypothesis
407 that in tadpoles the negative impact of ALAN could result from an alteration in the level of
408 melatonin synthesis and/or induce a shift in the rhythm of melatonin synthesis as shown in
409 fishes (Brüning et al., 2016, 2015; Kupprat et al., 2020; Vera et al., 2005), in birds (de Jong et
410 al., 2016; Dominoni et al., 2013; Jiang et al., 2020) and in mammals (Le Tallec et al., 2016;
411 Rahman et al., 2019). Several hypotheses could be put forward to explain our results. First,
412 tadpoles could have been exposed to a lower light illuminance than expected (as 3 lx was the
413 illuminance measured at the upper water level). However, melatonin production can be
414 suppressed by very low light illuminances (reviewed in Grubisic et al., 2019). Melatonin
415 production, measured in water tank or measured on individual plasma, is inhibited in fishes
416 and rodents, respectively, as early as 0.01 – 0.03 lx, and in human saliva from 6 lx. Because
417 in this study we only tested one level of ALAN (3 lx), we could also have missed a non-
418 monotone dose-response relationship (inverted U-shaped curve), where the effect of ALAN
419 on relative expression of genes increases up to a certain light level, and then decreases past
420 this threshold. Such pattern have already been observed for physiological parameters (Yang et
421 al., 2018, 2016) and behavioural variables (Gomes et al., 2021; Touzot et al., 2020), whose
422 responses to light exposure is non-linear. Still, our results support the possibility that
423 ecologically relevant and low levels of ALAN may have a masking effect on molecular
424 pacemakers, *i.e.*, circadian clock genes, in vertebrates, as it has been suggested in the diurnal
425 songbird, *Zebra finch* (Alaasam et al., 2021). The masking effect, which appears to occur at
426 low ALAN levels, corresponds to a situation in which the behavior and the physiology are
427 altered by the external environment (*i.e.*, low light intensity) without any change in the central
428 circadian clock genes (Mrosovsky, 1999; Rietveld et al., 1993). Second, the faint nocturnal
429 pattern in the expression of genes encoding melatonin synthesis enzymes in control tadpoles
430 could suggest that at this early stage of development, mechanisms regulating circadian rhythm

431 may not be fully matured. Indeed, it is known that in addition to a circadian rhythm, the
432 activity of AANAT varies during the development of tadpoles, with a low AANAT activity in
433 the first development stages (Alonzo-Gómez et al., 1994). For instance, in the Perez's frog,
434 *Pelophylax perezi*, an increase of 26% of AANAT activity was observed between the Gosner
435 development stages 26/27 and 40 (Alonzo-Gómez et al., 1994). However, many studies
436 showed that the circadian rhythm of melatonin synthesis is strongly dependent on the
437 light/dark cycle in tadpole (Green et al., 1999) and appears very early in the development of
438 amphibians (Alonzo-Gómez et al., 1994). Indeed, in *P. perezi*, a significant day/night rhythm
439 of AANAT was detected in the eyes at development stage 25 of Gosner and at development
440 stage 43/44 of Nieuwkoop and Faber in the *Xenopus laevis* (Alonzo-Gómez et al., 1994). The
441 small differences in overnight gene expression in tadpoles in the present study could also be
442 partly related to the use of the whole head for the RNA extraction which could contain other
443 melatonin-synthesizing tissues. Indeed, melatonin production was detected in the gut of adult
444 frogs, *Rana pipiens* and *Lithobates catesbeianus* (Bubenik and Pang, 1997) and in various
445 tissues and glands in *L. catesbeianus* tadpoles (Wright et al., 2001). These extra pineal and
446 extra retinal sites of melatonin production could be independent of the light/dark cycle and be
447 rather regulated by tissue demands, such as the local defense against oxidative stress (Acuña-
448 Castroviejo et al., 2014). Alternatively, although a positive correlation between retinal
449 AANAT activity and retina melatonin level has been demonstrated during the light/dark cycle
450 in adult green frogs, *P. esculenta* (d'Istria et al., 1994; Serino et al., 1993), this was not the
451 case in adult European green toads, *Bufo viridis* (Serino et al., 1993). Similarly, in adult
452 Perez's frogs, *P. perezi*, high retina AANAT activity during winter at any time of the day was
453 associated with low retina melatonin synthesis (Delgado et al., 1993). This could also be the
454 case in our species, *i.e.*, enzyme activity did not reflect melatonin synthesis. Indeed, even if

455 proteins come from the translation of mRNAs, it would be approximate to consider the
456 proteome as the exact reflect of the transcriptome.

457 4.3. Slight effects of low ALAN levels on melatonin signalling pathway

458 Present data also showed that the expression of genes encoding melatonin receptor
459 subtypes could be detected in tadpoles of both *R. dalmatina* and *B. bufo*, in accordance with
460 the ontogenic profile of melatonin binding sites in tadpoles of *P. perezi* (Isorna et al., 2005).
461 For both species, we observed weak changes in the relative expression of melatonin receptor
462 subtypes over time, with a global trend to overexpress at the end of the night/early morning. A
463 recent comparison of transcriptomic levels during the day and the night in a freshwater
464 teleost, *Oryzias latipes*, mainly revealed an up-regulation of melatonin receptors during the
465 night, however, photoperiod has also been shown to act in a gene-specific and tissue-specific
466 manner (Maugars et al., 2020). While it is at night that Mel1a increased in the pituitary and
467 the heart, and Mel1c increased in the eye and the skin, Mel1b was significantly higher during
468 the day in the adipose tissue of *O. latipes* (Maugars et al., 2020). Our results also showed for
469 the first time to our knowledge that the expression of genes encoding for membrane melatonin
470 receptors was impacted by ALAN, but without a clear pattern. In *R. dalmatina*, ALAN
471 inhibited the nocturnal increase of Mel1c, a receptor of the quinone reductase 2 family that
472 has only been identified in fish, amphibians, and birds (Li et al., 2013). Sparse studies in
473 amphibians revealed that the Mel1c receptor is located in the retina, brain and peripheral
474 tissues in amphibians (Ebisawa et al., 1994; Rada and Wiechmann, 2006; Wiechmann et al.,
475 2009, 1999) and a reduction of its expression could contribute to an alteration of the overall
476 regulation of melatonin signal transduction (Maugars et al., 2020). Further studies are
477 required to confirm this statement. In *B. bufo*, Mel1b, a receptor which is present in all
478 vertebrates (Dubocovich and Markowska, 2005; Reppert et al., 1994; Slominski et al., 2005)
479 and whose expression is stable over time in the control individuals of our study, was increased

480 when tadpoles were exposed to ALAN. It therefore appears that low levels of ALAN act on
481 the expression of genes encoding melatonin receptor in a gene-specific and a species-specific
482 manner in amphibians. Such an observation is consistent with previous studies showing that
483 the expression of melatonin receptors varies according to species, geographical area, lighting
484 regime, time of day, stage of neuronal development and endocrine status (Jones et al., 2012;
485 Vanecek, 1998; Witt-Enderby et al., 2003). Most of these studies concern adult organisms and
486 present results suggest that studies on juvenile are necessary to better understand
487 physiological impact of ALAN especially at early stage of organism development.

488 **5. Conclusion**

489 Through the development of a rigorous RT-qPCR method, we have described the
490 nocturnal changes in the relative expression of two melatonin-synthesis related enzymes and
491 three melatonin-related receptors in tadpoles of two non-model amphibian species, *R.*
492 *dalmatina* and *B. bufo*. For the first time, the results provided experimental evidence that a 3
493 lx ALAN illuminance, which is encountered in urban and peri urban areas, had weak effects
494 on melatonin-related genes and melatonin signaling at this stage of development in contrast to
495 the deleterious effects of such illuminance on adult physiology and behavior.

496 **6. Acknowledgements**

497 We thank A. Clair, J. Ulmann and L. Averty for their technical assistance in the EcoAquatron.
498 We also thanks T. Lefebure, P. Veber and V. Navratil for their help with *de novo*
499 transcriptomes assembly.

500 **7. Funding**

501 This work was supported by the French Government [PhD grants 2017–2020], by the
502 ‘LABEX IMU Laboratoire d’excellence Intelligence des Mondes Urbains’, by IDEX

503 Initiative d'excellence Université Lyon and by the project One Health approach of the
504 artificial light at night. The authors declare no competing or financial interests.

505 **8. Data availability**

506 *R. dalmatina de novo* transcriptome reads are in the process of being deposited on the of the
507 Sequence Read Archive (SRA) and *de novo* transcriptome assembly is in the process of being
508 deposited on the Transcriptome Shotgun Assembly (TSA) database of the SRA. All data will
509 be available on SRA BioProject.

510 **9. References**

- 511 Acuña-Castroviejo, D., Escames, G., Venegas, C., Díaz-Casado, M.E., Lima-Cabello, E., López, L.C.,
512 Rosales-Corral, S., Tan, D.-X., Reiter, R.J., 2014. Extrapineal melatonin: sources, regulation,
513 and potential functions. *Cell. Mol. Life Sci.* 71, 2997–3025. [https://doi.org/10.1007/s00018-](https://doi.org/10.1007/s00018-014-1579-2)
514 [014-1579-2](https://doi.org/10.1007/s00018-014-1579-2)
- 515 Alaasam, V.J., Liu, X., Niu, Y., Habibian, J.S., Pieraut, S., Ferguson, B.S., Zhang, Y., Ouyang, J.Q., 2021.
516 Effects of dim artificial light at night on locomotor activity, cardiovascular physiology, and
517 circadian clock genes in a diurnal songbird. *Environmental Pollution* 282, 117036.
518 <https://doi.org/10.1016/j.envpol.2021.117036>
- 519 Alonzo-Gómez, A.L., de Pedro, N., Gancedo, B., Alonso-Bedate, M., Valenciano, A.I., Delgado, M.J.,
520 1994. Ontogeny of Ocular Serotonin N-Acetyltransferase Activity Daily Rhythm in Four
521 Anuran Species. *General and Comparative Endocrinology* 94, 357–365.
522 <https://doi.org/10.1006/gcen.1994.1091>
- 523 Andersen, C.L., Jensen, J.L., Ørntoft, T.F., 2004. Normalization of Real-Time Quantitative Reverse
524 Transcription-PCR Data: A Model-Based Variance Estimation Approach to Identify Genes
525 Suited for Normalization, Applied to Bladder and Colon Cancer Data Sets. *Cancer Res* 64,
526 5245–5250. <https://doi.org/10.1158/0008-5472.CAN-04-0496>
- 527 Ayoub, M.A., Couturier, C., Lucas-Meunier, E., Angers, S., Fossier, P., Bouvier, M., Jockers, R., 2002.
528 Monitoring of Ligand-independent Dimerization and Ligand-induced Conformational Changes
529 of Melatonin Receptors in Living Cells by Bioluminescence Resonance Energy Transfer.
530 *Journal of Biological Chemistry* 277, 21522–21528. <https://doi.org/10.1074/jbc.M200729200>
- 531 Bayarri, M.J., Muñoz-Cueto, J.A., López-Olmeda, J.F., Vera, L.M., Rol de Lama, M.A., Madrid, J.A.,
532 Sánchez-Vázquez, F.J., 2004. Daily locomotor activity and melatonin rhythms in Senegal sole
533 (*Solea senegalensis*). *Physiology & Behavior* 81, 577–583.
534 <https://doi.org/10.1016/j.physbeh.2004.02.001>
- 535 Beebee, T.J.C., 1979. Habitats of the British amphibians (2): suburban parks and gardens. *Biological*
536 *Conservation* 15, 241–257. [https://doi.org/10.1016/0006-3207\(79\)90046-6](https://doi.org/10.1016/0006-3207(79)90046-6)
- 537 Bennie, J., Davies, T.W., Cruse, D., Gaston, K.J., Swenson, N., 2016. Ecological effects of artificial light
538 at night on wild plants. *Journal of Ecology* 104, 611–620. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2745.12551)
539 [2745.12551](https://doi.org/10.1111/1365-2745.12551)
- 540 Borjigin, J., Wang, M.M., Snyder, S.H., 1995. Diurnal variation in mRNA encoding serotonin N-
541 acetyltransferase in pineal gland. *Nature* 378, 783–785. <https://doi.org/10.1038/378783a0>

542 Brüning, A., Hölker, F., Franke, S., Kleiner, W., Kloas, W., 2016. Impact of different colours of artificial
543 light at night on melatonin rhythm and gene expression of gonadotropins in European perch.
544 Science of The Total Environment 543, 214–222.
545 <https://doi.org/10.1016/j.scitotenv.2015.11.023>

546 Brüning, A., Hölker, F., Franke, S., Preuer, T., Kloas, W., 2015. Spotlight on fish: Light pollution affects
547 circadian rhythms of European perch but does not cause stress. Science of The Total
548 Environment 511, 516–522. <https://doi.org/10.1016/j.scitotenv.2014.12.094>

549 Bubenik, G.A., Pang, S.F., 1997. Melatonin Levels in the Gastrointestinal Tissues of Fish, Amphibians,
550 and a Reptile. General and Comparative Endocrinology 106, 415–419.
551 <https://doi.org/10.1006/gcen.1997.6889>

552 Buchanan, B.W., 2006. Observed and potential effects of artificial night lighting on anuran
553 amphibians. Ecological consequences of artificial night lighting 192–220.

554 Calvo, J.R., González-Yanes, C., Maldonado, M.D., 2013. The role of melatonin in the cells of the
555 innate immunity: a review. J. Pineal Res. 55, 103–120. <https://doi.org/10.1111/jpi.12075>

556 Carrillo-Vico, A., Lardone, P., Álvarez-Sánchez, N., Rodríguez-Rodríguez, A., Guerrero, J., 2013.
557 Melatonin: Buffering the Immune System. IJMS 14, 8638–8683.
558 <https://doi.org/10.3390/ijms14048638>

559 Cinzano, P., Falchi, F., Elvidge, C.D., 2001. The first World Atlas of the artificial night sky brightness.
560 Monthly Notices of the Royal Astronomical Society 328, 689–707.
561 <https://doi.org/10.1046/j.1365-8711.2001.04882.x>

562 Confente, F., Rendón, M.C., Besseau, L., Falcón, J., Muñoz-Cueto, J.A., 2010. Melatonin receptors in a
563 pleuronectiform species, *Solea senegalensis*: Cloning, tissue expression, day–night and
564 seasonal variations. General and Comparative Endocrinology 167, 202–214.
565 <https://doi.org/10.1016/j.ygcen.2010.03.006>

566 Coon, S.L., del Olmo, E., Young, W.S., Klein, D.C., 2002. Melatonin Synthesis Enzymes in *Macaca*
567 *mulatta* : Focus on Arylalkylamine *N*-Acetyltransferase (EC 2.3.1.87). The Journal of Clinical
568 Endocrinology & Metabolism 87, 4699–4706. <https://doi.org/10.1210/jc.2002-020683>

569 d'Istria, M., Monteleone, P., Serino, I., Chieffi, G., 1994. Seasonal variations in the daily rhythm of
570 melatonin and NAT activity in the Harderian gland, retina, pineal gland, and serum of the
571 green frog, *Rana esculenta*. General and comparative endocrinology 96, 6–11.

572 Dananay, K.L., Benard, M.F., 2018. Artificial light at night decreases metamorphic duration and
573 juvenile growth in a widespread amphibian. Proc. R. Soc. B 285, 20180367.
574 <https://doi.org/10.1098/rspb.2018.0367>

575 Davies, T.W., Bennie, J., Gaston, K.J., 2012. Street lighting changes the composition of invertebrate
576 communities. Biol. Lett. 8, 764–767. <https://doi.org/10.1098/rsbl.2012.0216>

577 Davies, T.W., Coleman, M., Griffith, K.M., Jenkins, S.R., 2015. Night-time lighting alters the
578 composition of marine epifaunal communities. Biol. Lett. 11, 20150080.
579 <https://doi.org/10.1098/rsbl.2015.0080>

580 Davies, T.W., Smyth, T., 2018. Why artificial light at night should be a focus for global change
581 research in the 21st century. Glob Change Biol 24, 872–882.
582 <https://doi.org/10.1111/gcb.13927>

583 de Jong, M., Jeninga, L., Ouyang, J.Q., van Oers, K., Spoelstra, K., Visser, M.E., 2016. Dose-dependent
584 responses of avian daily rhythms to artificial light at night. Physiology & Behavior 155, 172–
585 179. <https://doi.org/10.1016/j.physbeh.2015.12.012>

586 Delgado, M.J., Aeonso-Gómez, A.L., Gancedo, B., De Pedro, N., Valenciano, A.I., Alonso-Bedate, M.,
587 1993. Serotonin *N*-acetyltransferase (NAT) activity and melatonin levels in the frog retina are
588 not correlated during the seasonal cycle. General and Comparative Endocrinology 92, 143–
589 150.

590 Delgado, M.J., Vivien-Roels, B., 1989. Effect of environmental temperature and photoperiod on the
591 melatonin levels in the pineal, lateral eye, and plasma of the frog, *Rana perezi*: Importance of
592 ocular melatonin. General and Comparative Endocrinology 75, 46–53.
593 [http://dx.doi.org/10.1016/0016-6480\(89\)90006-3](http://dx.doi.org/10.1016/0016-6480(89)90006-3)

594 Desouhant, E., Gomes, E., Mondy, N., Amat, I., 2019. Mechanistic, ecological, and evolutionary
595 consequences of artificial light at night for insects: review and prospective. *Entomol Exp Appl*
596 167, 37–58. <https://doi.org/10.1111/eea.12754>

597 Dhorne-Pollet, S., Th  lie, A., Pollet, N., 2013. Validation of novel reference genes for RT-qPCR studies
598 of gene expression in *Xenopus tropicalis* during embryonic and post-embryonic
599 development: *X. Tropicalis* Rt-Qpcr Reference Genes. *Dev. Dyn.* 242, 709–717.
600 <https://doi.org/10.1002/dvdy.23972>

601 Dominoni, D.M., Goymann, W., Helm, B., Partecke, J., 2013. Urban-like night illumination reduces
602 melatonin release in European blackbirds (*Turdus merula*): implications of city life for
603 biological time-keeping of songbirds. *Front Zool* 10, 60. [https://doi.org/10.1186/1742-9994-](https://doi.org/10.1186/1742-9994-10-60)
604 10-60

605 Dubocovich, M.L., Markowska, M., 2005. Functional MT1 and MT2 Melatonin Receptors in Mammals.
606 *ENDO* 27, 101–110. <https://doi.org/10.1385/ENDO:27:2:101>

607 Ebisawa, T., Karne, S., Lerner, M.R., Reppert, S.M., 1994. Expression cloning of a high-affinity
608 melatonin receptor from *Xenopus* dermal melanophores. *Proc. Natl. Acad. Sci. U.S.A.* 91,
609 6133–6137. <https://doi.org/10.1073/pnas.91.13.6133>

610 Edwards, M.L.O., Pivorun, E.B., 1991. The effects of photoperiod and different dosages of melatonin
611 on metamorphic rate and weight gain in *Xenopus laevis* tadpoles. *General and Comparative*
612 *Endocrinology* 81, 28–38. [http://dx.doi.org/10.1016/0016-6480\(91\)90122-M](http://dx.doi.org/10.1016/0016-6480(91)90122-M)

613 Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C.C.M., Elvidge, C.D., Baugh, K., Portnov, B.A., Rybnikova,
614 N.A., Furgoni, R., 2016. The new world atlas of artificial night sky brightness. *Sci. Adv.* 2,
615 e1600377. <https://doi.org/10.1126/sciadv.1600377>

616 Faraway, J.J., 2006. Extending the linear model with R: generalized linear, mixed effects and
617 nonparametric regression models. Chapman & Hall/CRC, Boca Raton.

618 Fonken, L.K., Nelson, R.J., 2016. Effects of light exposure at night during development. *Current*
619 *Opinion in Behavioral Sciences* 7, 33–39. <https://doi.org/10.1016/j.cobeha.2015.10.008>

620 Forsburg, Z.R., Guzman, A., Gabor, C.R., 2021. Artificial light at night (ALAN) affects the stress
621 physiology but not the behavior or growth of *Rana berlandieri* and *Bufo valliceps*.
622 *Environmental Pollution* 277, 116775. <https://doi.org/10.1016/j.envpol.2021.116775>

623 Galton, V.A., 1992. The role of thyroid hormone in amphibian metamorphosis. *Trends in*
624 *Endocrinology & Metabolism* 3, 96–100. [https://doi.org/10.1016/1043-2760\(92\)90020-2](https://doi.org/10.1016/1043-2760(92)90020-2)

625 Gaston, K.J., Bennie, J., Davies, T.W., Hopkins, J., 2013. The ecological impacts of nighttime light
626 pollution: a mechanistic appraisal: Nighttime light pollution. *Biol Rev* 88, 912–927.
627 <https://doi.org/10.1111/brv.12036>

628 Gaston, K.J., Davies, T.W., Bennie, J., Hopkins, J., 2012. REVIEW: Reducing the ecological
629 consequences of night-time light pollution: options and developments. *J Appl Ecol* 49, 1256–
630 1266. <https://doi.org/10.1111/j.1365-2664.2012.02212.x>

631 Gaston, K.J., Davies, T.W., Nedelec, S.L., Holt, L.A., 2017. Impacts of Artificial Light at Night on
632 Biological Timings. *Annu. Rev. Ecol. Evol. Syst.* 48, 49–68. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-ecolsys-110316-022745)
633 ecolsys-110316-022745

634 Gaston, K.J., Duffy, J.P., Gaston, S., Bennie, J., Davies, T.W., 2014. Human alteration of natural light
635 cycles: causes and ecological consequences. *Oecologia* 176, 917–931.
636 <https://doi.org/10.1007/s00442-014-3088-2>

637 Ginzinger, D.G., 2002. Gene quantification using real-time quantitative PCR. *Experimental*
638 *Hematology* 30, 503–512. [https://doi.org/10.1016/S0301-472X\(02\)00806-8](https://doi.org/10.1016/S0301-472X(02)00806-8)

639 Gomes, E., Rey, B., D  bias, F., Amat, I., Desouhant, E., 2021. Dealing with host and food searching in a
640 diurnal parasitoid: consequences of light at night at intra- and trans-generational levels.
641 *Insect Conserv Divers* 14, 235–246. <https://doi.org/10.1111/icad.12477>

642 Gosner, K.L., 1960. A Simplified Table for Staging Anuran Embryos and Larvae with Notes on
643 Identification. *Herpetologica* 16, 183–190.

644 Green, C.B., Liang, M.-Y., Steenhard, B.M., Besharse, J.C., 1999. Ontogeny of circadian and light
645 regulation of melatonin release in *Xenopus laevis* embryos. *Developmental Brain Research*
646 117, 109–116. [https://doi.org/10.1016/S0165-3806\(99\)00109-1](https://doi.org/10.1016/S0165-3806(99)00109-1)

647 Grubisic, M., Haim, A., Bhusal, P., Dominoni, D.M., Gabriel, K.M.A., Jechow, A., Kupprat, F., Lerner, A.,
648 Marchant, P., Riley, W., Stebelova, K., van Grunsven, R.H.A., Zeman, M., Zubidat, A.E., Hölker,
649 F., 2019. Light Pollution, Circadian Photoreception, and Melatonin in Vertebrates.
650 *Sustainability* 11, 6400. <https://doi.org/10.3390/su11226400>

651 Hardeland, R., 2008. Melatonin, hormone of darkness and more – occurrence, control mechanisms,
652 actions and bioactive metabolites. *Cell. Mol. Life Sci.* 65, 2001–2018.
653 <https://doi.org/10.1007/s00018-008-8001-x>

654 Hellemans, J., Mortier, G., De Paepe, A., Speleman, F., Vandesompele, J., 2007. qBase relative
655 quantification framework and software for management and automated analysis of real-time
656 quantitative PCR data. *Genome Biol* 8, R19. <https://doi.org/10.1186/gb-2007-8-2-r19>

657 International Union for Conservation of Nature, IUCN, The IUCN red list of threatened species, n.d.
658 International Union for Conservation of Nature, IUCN, <https://www.iucnredlist.org/> [WWW
659 Document]. URL <https://www.iucnredlist.org/> (accessed 5.18.20).

660 Isorna, E., Guijarro, A.I., Delgado, M.J., López-Patiño, M.A., Pedro, N. de, Luis Alonso-Gómez, Á.,
661 2005. Ontogeny of central melatonin receptors in tadpoles of the anuran *Rana perezi*:
662 modulation of dopamine release. *J Comp Physiol A* 191, 1099–1105.
663 <https://doi.org/10.1007/s00359-005-0032-2>

664 Jechow, A., Hölker, F., 2019. How dark is a river? Artificial light at night in aquatic systems and the
665 need for comprehensive night-time light measurements. *WIREs Water* 6.
666 <https://doi.org/10.1002/wat2.1388>

667 Jessop, T.S., Dempster, T., Letnic, M., Webb, J.K., 2014. Interplay among nocturnal activity,
668 melatonin, corticosterone and performance in the invasive cane toad (*Rhinella marinus*).
669 *General and Comparative Endocrinology* 206, 43–50.
670 <https://doi.org/10.1016/j.ygcen.2014.07.013>

671 Jiang, J., He, Y., Kou, H., Ju, Z., Gao, X., Zhao, H., 2020. The effects of artificial light at night on
672 Eurasian tree sparrow (*Passer montanus*): Behavioral rhythm disruption, melatonin
673 suppression and intestinal microbiota alterations. *Ecological Indicators* 108, 105702.
674 <https://doi.org/10.1016/j.ecolind.2019.105702>

675 Jones, C., Helfer, G., Brandstätter, R., 2012. Melatonin Receptor Expression in the Zebra Finch Brain
676 and Peripheral Tissues. *Chronobiology International* 29, 189–202.
677 <https://doi.org/10.3109/07420528.2011.642912>

678 Kamphuis, W., Cailotto, C., Dijk, F., Bergen, A., Buijs, R.M., 2005. Circadian expression of clock genes
679 and clock-controlled genes in the rat retina. *Biochemical and Biophysical Research*
680 *Communications* 330, 18–26. <https://doi.org/10.1016/j.bbrc.2005.02.118>

681 Kikuyama, S., Kawamura, K., Tanaka, S., Yamamoto, K., 1993. Aspects of Amphibian Metamorphosis:
682 Hormonal Control, in: *International Review of Cytology*. Elsevier, pp. 105–148.
683 [https://doi.org/10.1016/S0074-7696\(08\)60426-X](https://doi.org/10.1016/S0074-7696(08)60426-X)

684 Kupprat, F., Hölker, F., Kloas, W., 2020. Can skyglow reduce nocturnal melatonin concentrations in
685 Eurasian perch? *Environmental Pollution* 262, 114324.
686 <https://doi.org/10.1016/j.envpol.2020.114324>

687 Le Pape, S., 2013. EasyqpcR: low-throughput real-time quantitative PCR data analysis.

688 Le Tallec, T., Théry, M., Perret, M., 2016. Melatonin concentrations and timing of seasonal
689 reproduction in male mouse lemurs (*Microcebus murinus*) exposed to light pollution.
690 *JMAMMAL* 97, 753–760. <https://doi.org/10.1093/jmammal/gyw003>

691 Li, D., Smith, D., Hardeland, R., Yang, M., Xu, H., Zhang, L., Yin, H., Zhu, Q., 2013. Melatonin Receptor
692 Genes in Vertebrates. *IJMS* 14, 11208–11223. <https://doi.org/10.3390/ijms140611208>

693 Longcore, T., Rich, C., 2004. Ecological light pollution. *Frontiers in Ecology and the Environment* 2,
694 191–198. [https://doi.org/10.1890/1540-9295\(2004\)002\[0191:elp\]2.0.co;2](https://doi.org/10.1890/1540-9295(2004)002[0191:elp]2.0.co;2)

695 Lou, Q., Cao, S., Xu, W., Zhang, Y., Qin, Z., Wei, W., 2014. Molecular characterization and mRNA
696 expression of ribosomal protein L8 in *Rana nigromaculata* during development and under
697 exposure to hormones. *Journal of Environmental Sciences* 26, 2331–2339.
698 <https://doi.org/10.1016/j.jes.2014.09.018>

699 Maugars, G., Nourizadeh-Lillabadi, R., Weltzien, F.-A., 2020. New Insights Into the Evolutionary
700 History of Melatonin Receptors in Vertebrates, With Particular Focus on Teleosts. *Front.*
701 *Endocrinol.* 11, 538196. <https://doi.org/10.3389/fendo.2020.538196>

702 Mayer, I., Bornestaf, C., Borg, B., 1997. Melatonin in non-mammalian vertebrates: Physiological role
703 in reproduction? *Comparative Biochemistry and Physiology Part A: Physiology* 118, 515–531.
704 [https://doi.org/10.1016/S0300-9629\(96\)00468-9](https://doi.org/10.1016/S0300-9629(96)00468-9)

705 Mrosovsky, N., 1999. Masking: History, Definitions, and Measurement. *Chronobiology International*
706 16, 415–429. <https://doi.org/10.3109/07420529908998717>

707 Pohanka, M., 2013. Impact of melatonin on immunity: a review. *Open Medicine* 8.
708 <https://doi.org/10.2478/s11536-013-0177-2>

709 R Core Team, 2018. . R: A Language and Environment for Statistical Computing. R foundation for
710 statistical computing.

711 Rada, J.A.S., Wiechmann, A.F., 2006. Melatonin Receptors in Chick Ocular Tissues: Implications for a
712 Role of Melatonin in Ocular Growth Regulation. *Invest. Ophthalmol. Vis. Sci.* 47, 25.
713 <https://doi.org/10.1167/iovs.05-0195>

714 Rahman, S.A., Wright, K.P., Lockley, S.W., Czeisler, C.A., Gronfier, C., 2019. Characterizing the
715 temporal Dynamics of Melatonin and Cortisol Changes in Response to Nocturnal Light
716 Exposure. *Sci Rep* 9, 19720. <https://doi.org/10.1038/s41598-019-54806-7>

717 Rebollar, P.G., Ubilla, E., Peleteiro, J.B., Agapito, M.T., Alvariño, J.M., 1999. Determination of plasma
718 melatonin levels by enzyme-linked immunosorbent assay (EIA) in turbot (*Scophthalmus*
719 *maximus* L.) and tench (*Tinca tinca* L.). *J Physiol Biochem* 55, 341–347.

720 Reiter, R.J., 1991. Melatonin: The chemical expression of darkness. *Molecular and Cellular*
721 *Endocrinology* 79, C153–C158. [https://doi.org/10.1016/0303-7207\(91\)90087-9](https://doi.org/10.1016/0303-7207(91)90087-9)

722 Reiter, R.J., Mayo, J.C., Tan, D.-X., Sainz, R.M., Alatorre-Jimenez, M., Qin, L., 2016. Melatonin as an
723 antioxidant: under promises but over delivers. *J. Pineal Res.* 61, 253–278.
724 <https://doi.org/10.1111/jpi.12360>

725 Reppert, S.M., Weaver, D.R., Ebisawa, T., 1994. Cloning and characterization of a mammalian
726 melatonin receptor that mediates reproductive and circadian responses. *Neuron* 13, 1177–
727 1185. [https://doi.org/10.1016/0896-6273\(94\)90055-8](https://doi.org/10.1016/0896-6273(94)90055-8)

728 Ribelayga, C., Gauer, F., Calgari, C., Pevet, P., Simonneaux, V., 1999. Photoneural Regulation of Rat
729 Pineal Hydroxyindole- *O* -Methyltransferase (HIOMT) Messenger Ribonucleic Acid Expression:
730 An Analysis of Its Complex Relationship with HIOMT Activity ¹. *Endocrinology* 140, 1375–
731 1384. <https://doi.org/10.1210/endo.140.3.6552>

732 Rietveld, W.J., Minors, D.S., Waterhouse, J.M., 1993. Circadian Rhythms and Masking: An Overview.
733 *Chronobiology International* 10, 306–312. <https://doi.org/10.1080/07420529309059713>

734 Sanders, D., Frago, E., Kehoe, R., Patterson, C., Gaston, K.J., 2021. A meta-analysis of biological
735 impacts of artificial light at night. *Nat Ecol Evol* 5, 74–81. <https://doi.org/10.1038/s41559-020-01322-x>

736

737 Sanders, D., Kehoe, R., Tiley, K., Bennie, J., Cruse, D., Davies, T.W., Frank van Veen, F.J., Gaston, K.J.,
738 2015. Artificial nighttime light changes aphid-parasitoid population dynamics. *Sci Rep* 5,
739 15232. <https://doi.org/10.1038/srep15232>

740 Schroer, S., Hölker, F., 2017. Impact of Lighting on Flora and Fauna, in: Karlicek, R., Sun, C.-C., Zissis,
741 G., Ma, R. (Eds.), *Handbook of Advanced Lighting Technology*. Springer International
742 Publishing, Cham, pp. 1–33. https://doi.org/10.1007/978-3-319-00295-8_42-1

743 Secondi, J., Dupont, V., Davranche, A., Mondy, N., Lengagne, T., Théry, M., 2017. Variability of surface
744 and underwater nocturnal spectral irradiance with the presence of clouds in urban and peri-
745 urban wetlands. *PLoS ONE* 12, e0186808. <https://doi.org/10.1371/journal.pone.0186808>

746 Secondi, J., Mondy, N., Gippet, J.M.W., Touzot, M., Gardette, V., Guillard, L., Lengagne, T., 2021.
747 Artificial light at night alters activity, body mass, and corticosterone level in a tropical anuran.
748 Behavioral Ecology arab044. <https://doi.org/10.1093/beheco/arab044>
749 Segrestin, J., Mondy, N., Boisselet, C., Guigard, L., Lengagne, T., Poussineau, S., Secondi, J., Puijalon,
750 S., 2021. Effects of artificial light at night on the leaf functional traits of freshwater plants.
751 Freshw Biol 66, 2264–2271. <https://doi.org/10.1111/fwb.13830>
752 Serino, I., D'Istria, M., Monteleone, P., 1993. A comparative study of melatonin production in the
753 retina, pineal gland and Harderian gland of *Bufo viridis* and *Rana esculenta*. Comparative
754 Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology 106, 189–
755 193. [https://doi.org/10.1016/0742-8413\(93\)90271-L](https://doi.org/10.1016/0742-8413(93)90271-L)
756 Slominski, A., Fischer, T.W., Zmijewski, M.A., Wortsman, J., Semak, I., Zbytek, B., Slominski, R.M.,
757 Tobin, D.J., 2005. On the Role of Melatonin in Skin Physiology and Pathology. ENDO 27, 137–
758 148. <https://doi.org/10.1385/ENDO:27:2:137>
759 Tan, D.-X., Hardeland, R., Manchester, L.C., Paredes, S.D., Korkmaz, A., Sainz, R.M., Mayo, J.C.,
760 Fuentes-Broto, L., Reiter, R.J., 2010. The changing biological roles of melatonin during
761 evolution: from an antioxidant to signals of darkness, sexual selection and fitness. Biological
762 Reviews 85, 607–623. <https://doi.org/10.1111/j.1469-185X.2009.00118.x>
763 Touzot, M., Lefebure, T., Lengagne, T., Secondi, J., Dumet, A., Konecny-Dupre, L., Veber, P., Navratil,
764 V., Duchamp, C., Mondy, N., 2021. Transcriptome-wide deregulation of gene expression by
765 artificial light at night in tadpoles of common toads. Science of The Total Environment
766 151734. <https://doi.org/10.1016/j.scitotenv.2021.151734>
767 Touzot, M., Lengagne, T., Secondi, J., Desouhant, E., Théry, M., Dumet, A., Duchamp, C., Mondy, N.,
768 2020. Artificial light at night alters the sexual behaviour and fertilisation success of the
769 common toad. Environmental Pollution 259, 113883.
770 <https://doi.org/10.1016/j.envpol.2019.113883>
771 Touzot, M., Teulier, L., Lengagne, T., Secondi, J., Théry, M., Libourel, P.-A., Guillard, L., Mondy, N.,
772 2019. Artificial light at night disturbs the activity and energy allocation of the common toad
773 during the breeding period. Conservation Physiology 7.
774 <https://doi.org/10.1093/conphys/coz002>
775 Vandesompele, J., Preter, K.D., Roy, N.V., Paepe, A.D., 2002. Accurate normalization of real-time
776 quantitative RT-PCR data by geometric averaging of multiple internal control genes 3.
777 Vanecek, J., 1998. Cellular Mechanisms of Melatonin Action. Physiological Reviews 78, 687–721.
778 <https://doi.org/10.1152/physrev.1998.78.3.687>
779 Vera, L.M., López-Olmeda, J.F., Bayarri, M.J., Madrid, J.A., Sánchez-Vázquez, F.J., 2005. Influence of
780 Light Intensity on Plasma Melatonin and Locomotor Activity Rhythms in Tench.
781 Chronobiology International 22, 67–78. <https://doi.org/10.1081/CBI-200038157>
782 Wiechmann, A.F., Campbell, L.D., Defoe, D.M., 1999. Melatonin receptor RNA expression in *Xenopus*
783 retina. Molecular Brain Research 63, 297–303. [https://doi.org/10.1016/S0169-328X\(98\)00292-7](https://doi.org/10.1016/S0169-328X(98)00292-7)
784 Wiechmann, A.F., Hollaway, L.R., Rada, J.A.S., 2009. Melatonin receptor expression in *Xenopus laevis*
785 surface corneal epithelium: Diurnal rhythm of lateral membrane localization. Molecular
786 Vision 20.
787 Witt-Enderby, P.A., Bennett, J., Jarzynka, M.J., Firestone, S., Melan, M.A., 2003. Melatonin receptors
788 and their regulation: biochemical and structural mechanisms. Life Sciences 72, 2183–2198.
789 [https://doi.org/10.1016/S0024-3205\(03\)00098-5](https://doi.org/10.1016/S0024-3205(03)00098-5)
790 Wood, S.N., 2006. On confidence intervals for generalized additive models based on penalized
791 regression splines. Aust N Z J Stat 48, 445–464. <https://doi.org/10.1111/j.1467-842X.2006.00450.x>
792 Wright, M., Alves, C., Francisco, L., Shea, T., Visconti, R., Bruni, N., 2001. Regulation of the plasma
793 melatonin level in anuran tadpoles. Perspectives in Comparative Endocrinology. Monduzzi
794 Editore, Bologna 669–674.
795
796

797 Wright, M.L., Alves, C.D., 2001. The decrease in plasma melatonin at metamorphic climax in *Rana*
798 *catesbeiana* bullfrog tadpoles is induced by thyroxine 129, 653–663.

799 Wright, M.L., Francisco, L.L., Scott, J.L., Richardson, S.E., Carr, J.A., King, A.B., Noyes, A.G., Visconti,
800 R.F., 2006. Effects of bilateral and unilateral ophthalmectomy on plasma melatonin in *Rana*
801 tadpoles and froglets under various experimental conditions. *General and Comparative*
802 *Endocrinology* 147, 158–166. <https://doi.org/10.1016/j.ygcen.2005.12.014>

803 Yang, Y., Pan, C., Zhong, R., Pan, J., 2018. Artificial light and biological responses of broiler chickens:
804 dose-response2. *Journal of Animal Science* 96, 98–107. <https://doi.org/10.1093/jas/skx044>

805 Yang, Y., Yu, Y., Yang, B., Zhou, H., Pan, J., 2016. Physiological responses to daily light exposure. *Sci*
806 *Rep* 6, 24808. <https://doi.org/10.1038/srep24808>

807 Zhao, D., Yu, Y., Shen, Y., Liu, Q., Zhao, Z., Sharma, R., Reiter, R.J., 2019. Melatonin Synthesis and
808 Function: Evolutionary History in Animals and Plants. *Front. Endocrinol.* 10, 249.
809 <https://doi.org/10.3389/fendo.2019.00249>

810 Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A., Smith, G.M., 2009. Mixed effects models and
811 extensions in ecology with R, *Statistics for Biology and Health*. Springer New York, New York,
812 NY. <https://doi.org/10.1007/978-0-387-87458-6>

813

814

815 **Tables and Figures**

816 **Table 1:** List of primers used for RT-qPCR for each target and housekeeping gene according
817 to species. Target genes are AANAT: aralkylamine N-acetyltransferase, ASMT: N
818 acetylserotonin O-methyltransferase, Mel1a, Mel1b and Mel1c: melatonin receptors.
819 Housekeeping genes are β -actin, rpl8: 60S ribosomal protein L8, tbp: TATA-binding protein,
820 and GluR2: AMPA-type glutamate receptor.

821

822 **Table 2:** Housekeeping gene stability analysis with GeNorm and Normfinder for *B. bufo* and
823 *R. dalmatina*. GeNorm stability values are given for the entire dataset. Normfinder stability
824 was analysed according to light treatment or timepoint. A low stability value indicates a low
825 variation in gene expression either globally or in each of the tested conditions of light
826 treatment or timepoint.

827

828 **Table 3:** GAM results for each target gene in *R. dalmatina* with the AIC scores for the
829 complete model (Timepoint * Light treatment) and the best model. The « * » character
830 indicates the interaction and the additive effects between the explanatory terms. Bold P-values
831 are significant.

832

833 **Figure 1:** Relative abundance of transcripts encoding for target gene in *R. dalmatina* exposed
834 to two light treatments of different night-time illuminance: control-group (black) or ALAN-
835 group (orange) and sampled over 8 timepoints. Each panel corresponds to one melatonin-related
836 target gene with a) AANAT, b) ASMT, c) Mel1b, and d) Mel1c. When the light treatment is
837 significant, both treatments are distinguished (solid dots and lines), while when the light
838 treatment is not significant, the treatments are not distinct (one dashed line). Darkblue shaded
839 areas comprised between 20 and 06 hours correspond to night-time period. Dots represent
840 relative abundance of transcripts of each sample, lines represent the regression from the GAM,
841 and shaded areas correspond to the 95% confidence interval. Relative abundance of transcripts
842 was normalised based on the transcription of β -actin, rpl8 and tbp housekeeping genes.

843

844 **Table 4:** GAM results for each target gene in *B. bufo* with the AIC scores for the complete
845 model (Timepoint * Light treatment) and the best model. The « * » character indicates the
846 interaction and the additive effects between the explanatory terms. Bold P-values were
847 significant.

848

849 **Figure 2:** Relative abundance of transcripts encoding for target gene in *B. bufo* exposed to
850 two light treatments of different night-time illuminance: control-group (black) or ALAN-
851 group (orange) and sampled over 8 timepoints. Each panel corresponds to one melatonin-
852 related target gene with a) AANAT, b) ASMT, c) Mel1a, d) Mel1b and e) Mel1c. When the
853 light treatment is significant, both treatments are distinguished (solid dots and lines), while
854 when the light treatment is not significant, the treatments are not distinct (one dashed line).
855 Darkblue shaded areas comprised between 20 and 06 hours correspond to night-time period.
856 Dots represent relative abundance of transcripts of each sample, lines represent the regression

857 from the GAM, and shaded areas correspond to the 95% confidence interval. Relative
858 abundance of transcripts was normalised based on the transcription of β -actin, rpl8 and tbp
859 housekeeping genes.


Gene	Species	Forward 5'-3' primer sequence	Reverse 5'-3' primer sequence	Amplicon size (bp)
<i>Target genes</i>				
AANAT	<i>R. dalmatina</i>	CCTGCCAGTGAATTCCGTTG	CCTGACTTCATCCAGATGGA	116
	<i>B. bufo</i>	CCCGCCAGTGAATTCCGCTG	TCTAACTTCATCCAGGTGGA	116
ASMT	<i>R. dalmatina</i>	TTTCTTCAGTATATCTGATC	TAGCTTGAGGTAGGAATTG	104
	<i>B. bufo</i>	TTTCCACCACGTACTTGGTC	CAGCCTGCGGTAAGAAGTG	104
Mel1a	<i>R. dalmatina</i>	TTAATGGGAATAAGTGTCAT	ATCCAACAAATAAATTTGGC	184
	<i>B. bufo</i>	TTAATGGGAATAAGTGTCAT	ATCCAACAAATAAATTTGGC	184
Mel1b	<i>R. dalmatina</i>	TGCTGTTATATACGGATTGC	GCTCTTTGGGCATTCTGTTC	126
	<i>B. bufo</i>	TGCTGTGATTTATGGACTTC	GCTTTTAGGGCACTCTGTTCG	126
Mel1c	<i>R. dalmatina</i>	CAAGCACAGAGTAAGACAAG	GCCACCGCTAGGCCAATAAA	139
	<i>B. bufo</i>	CAAGTGAAGCACAGAGTAAG	GCTACAGCAAGGCCAATGAA	139
<i>Housekeeping genes</i>				
β-actin	<i>R. dalmatina</i>	GAAAATTCTGACTGAGAGGG	TGACCTGCCCATCAGGCAAC	140
	<i>B. bufo</i>	GAAGATTCTTACTGAGAGAG	TGACCTGCCCATCAGGCAAC	140
rpl8	<i>R. dalmatina</i>	GCTAAGCTCAGGGCTGTCTGA	AAGGCAACTTTAGCAAGTGG	109
	<i>B. bufo</i>	GCTAAGCTCAGGGCTGTCTGA	AAGGCAACCTTGGCAAGAGG	109
thp	<i>R. dalmatina</i>	TGAAGAGCAGTCGCGCTTGG	TTACATCACAGCTCCCTACC	115
	<i>B. bufo</i>	AACAATCTCGGTTAGCAGCC	GATGCGTGAGCACGAGACCC	149
GluR2	<i>B. bufo</i>	ACCTTGATTATCATTTTCATC	GAATTCCTTTGTTGACCCTG	149

Species	Gene	GeNorm stability value	Normfinder stability value	
			Light treatment effect	Timepoint effect
<i>B. bufo</i>	β -actin	0.806	0.04	0.23
	rpl8	0.938	0.07	0.26
	tbp	0.993	0.07	0.28
	GluR2	1.006	0.09	0.36
<i>R. dalmatina</i>	β -actin	0.424	0.03	0.10
	rpl8	0.446	0.03	0.10
	tbp	0.439	0.04	0.11

862 **Table 3**

Gene	Complete model AIC	Best model AIC	Explanatory term in the fixed part of the best model	Explained deviance (%)	Factor in the fixed part of the model	F-value	edf	P-value
AANAT	142.3	139.6	Timepoint	12.0	Timepoint	3.64	1.76	0.028
ASMT	168.6	165.0	Timepoint	16.2	Timepoint	13.32	1	<0.001
Mel1b	76.0	77.8	Timepoint	20.3	Timepoint	21.24	1	<0.001
Mel1c	37.5	37.5	Timepoint * Light treatment	19.6	Control-group * Timepoint ALAN-group * Timepoint	7.72 0.36	1.32 1	0.002 0.549

- Control-group
- ALAN-group
- Non-distinct treatment



Gene	Complete model AIC	Best model AIC	Explanatory term in the fixed part of the best model	Explained deviance (%)	Factor in the fixed part of the model	F-value	edf	P-value
AANAT	54.2	55.0	Timepoint	5.16	Timepoint	1.48	1.77	0.241
ASMT	115.4	111.5	Timepoint	2.05	Timepoint	0.47	1.42	0.684
Mel1a	136.9	133.3	Timepoint	10.10	Timepoint	6.53	1.00	0.013
Mel1b	132.7	132.7	Timepoint * Light	22.80	Control-group * Timepoint ALAN-group * Timepoint	1.06 7.15	3.51 1	0.387 0.009
Mel1c	101.7	102.7	Timepoint	0.03	Timepoint	0.016	1	0.900

Control-group
 ALAN-group
 Non-distinct treatment

