



HAL
open science

Sec22b is a critical and non-redundant regulator of plasma cell maintenance

Amélie Bonaud, Laetitia Gargowitsch, Simon M Gilbert, Elanchezhian Rajan, Pablo Canales-Herrerias, Daniel Stockholm, Nabila F Rahman, Mark O Collins, Hakan Taskiran, Danika L Hill, et al.

► **To cite this version:**

Amélie Bonaud, Laetitia Gargowitsch, Simon M Gilbert, Elanchezhian Rajan, Pablo Canales-Herrerias, et al.. Sec22b is a critical and non-redundant regulator of plasma cell maintenance. Proceedings of the National Academy of Sciences of the United States of America, 2023, 120 (2), pp.e2213056120. 10.1073/pnas.2213056120 . hal-04064289v1

HAL Id: hal-04064289

<https://hal.science/hal-04064289v1>

Submitted on 11 Apr 2023 (v1), last revised 18 Jul 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Sec22b is a critical and non-redundant regulator of plasma cell maintenance.**

2 Amélie Bonaud^{1,2,3}, Laetitia Gargowitsch⁴, Simon M. Gilbert⁵, Elanchezhian Rajan⁶, Pablo Canales-
3 Herrerias⁷, Daniel Stockholm^{8,9}, Nabila F. Rahman¹⁰, Mark O. Collins⁶, Hakan Taskiran^{11,12,13}, Danika
4 L. Hill^{14,15}, Andres Alloatti^{16,17}, Nagham Alouche⁴, Stéphanie Balor¹⁸, Vanessa Soldan¹⁸, Daniel
5 Gillet¹⁹, Julien Barbier¹⁹, Françoise Bachelier⁴, Kenneth G.C. Smith^{5,20}, Julia Jellusova^{11,12}, Pierre
6 Bruhns⁷, Sebastian Amigorena¹⁶, Karl Balabanian^{1,2,3}, Michelle A. Linterman¹⁴, Andrew A. Peden⁶ and
7 Marion Espéli^{1,2,3*}

9 *1- Université Paris-Cité, Institut de Recherche Saint-Louis, INSERM U1160, F-75010 Paris, France,*

10 *2- CNRS, GDR3697 “Microenvironment of tumor niches”, Micronit, France,*

11 *3- OPALE Carnot Institute, The Organization for Partnerships in Leukemia, Hôpital Saint-Louis,*
12 *Paris, France*

13 *4- Université Paris-Saclay, INSERM, Inflammation, Microbiome and Immunosurveillance, Clamart,*
14 *France.*

15 *5- Department of Medicine, University of Cambridge, Cambridge Biomedical Campus, Addenbrooke’s*
16 *Hospital, Cambridge, UK.*

17 *6- School of Bioscience, University of Sheffield, Western Bank, Sheffield, S102TN, UK*

18 *7- Institut Pasteur, Université de Paris, Unit of Antibodies in Therapy and Pathology, Inserm*
19 *UMR1222, F-75015 Paris*

20 *8- PSL Research University, EPHE, Paris, France*

21 *9- Sorbonne Université, INSERM, Centre de Recherche Saint-Antoine, CRSA, F-75012, Paris, France*

22 *10- Dementia Research Institute, University of Cardiff, Hadyn Ellis Building, Maindy Road, Cardiff,*
23 *CF24 4HQ*

24 *11- Institute of Clinical Chemistry and Pathobiochemistry, School of Medicine, Klinikum rechts der*
25 *Isar, Technical University of Munich, 81675 Munich, Germany*

26 *12- TranslaTUM, Center for Translational Cancer Research, Technical University of Munich, 81675*
27 *Munich, Germany*

28 *13- Max Planck Institute of Immunobiology and Epigenetics, 79108 Freiburg im Breisgau, Germany*

29 *14- Lymphocyte Signalling and Development, Babraham Institute, Babraham Research Campus,*
30 *Cambridge CB22 3AT, UK*

31 *15- Department of Immunology and Pathology, Monash University, Melbourne, Victoria, 3004,*
32 *Australia*

33 *16- PSL Research University, Institut Curie Research Center, INSERM U932, Paris, France*

34 *17- Facultad de Ciencias Médicas, Instituto de Inmunología Clínica y Experimental de Rosario*
35 *(IDICER)-CONICET/Universidad Nacional de Rosario, Rosario, Argentina.*

36 *18- METi, Centre de Biologie Intégrative, Université de Toulouse, CNRS, UPS, 31062, Toulouse,*
37 *France*

38 *19- Université Paris-Saclay, CEA, INRAE, Département Médicaments et Technologies pour la Santé*
39 *(DMTS), SIMoS, 91191 Gif-sur-Yvette, France*

40 *20- Cambridge Institute of Therapeutic Immunology & Infectious Disease, Jeffrey Cheah Biomedical*
41 *Centre Cambridge Biomedical, University of Cambridge, Cambridge, United Kingdom*

42

43 * Corresponding author: marion.espeli@inserm.fr

44 Classification: Biological Science – Immunology and Inflammation

45 Short title: **Sec22b controls plasma cell maintenance**

46 *Keywords:* Plasma cell – antibody – SNARE - Endoplasmic reticulum – Mitochondria.

47 **Abstract**

48 Despite the essential role of plasma cells in health and disease, the cellular mechanisms controlling
49 their survival and secretory capacity are still poorly understood. Here, we identified the Soluble N-
50 ethylmaleimide-sensitive factor Attachment Protein Receptor (SNARE) Sec22b as a unique and
51 critical regulator of plasma cell maintenance and function. In absence of Sec22b, plasma cells were
52 hardly detectable and serum antibody titres were dramatically reduced. Accordingly, *Sec22b* deficient
53 mice fail to mount a protective immune response. At the mechanistic level, we demonstrated that
54 Sec22b contributes to efficient antibody secretion and is a central regulator of plasma cell maintenance
55 through the regulation of their transcriptional identity and of the morphology of the endoplasmic
56 reticulum and mitochondria. Altogether, our results unveil an essential and non-redundant role for
57 Sec22b as a regulator of plasma cell fitness and of the humoral immune response.

58

59

60 **Significance Statement:**

61 Despite their central role in health and disease, the cellular mechanisms underlying how plasma cells
62 persist while producing large quantities of antibodies are still poorly understood. In this paper we
63 describe that plasma cell survival is under the strict dependency of the Soluble N-ethylmaleimide-
64 sensitive factor Attachment Protein Receptor (SNARE) molecule Sec22b. We showed that this
65 molecule is a key regulator of endoplasmic reticulum and mitochondrial structure in plasma cells and
66 in its absence the humoral immune response is abrogated. Our findings have important implications
67 for our understanding of antibody mediated immunity but also open avenues for targeting plasma cells
68 in pathological contexts.

69

70

71 **Introduction:**

72 Plasma cells (PCs) are the cellular source of humoral immunity via the long-term secretion of large
73 quantities of antibodies that provide protection against reinfection. These cells can also contribute to
74 diseases including plasmacytomas as well as antibody-mediated autoimmune and inflammatory
75 pathologies. However, the therapeutic arsenal to target PCs is still very limited. Despite the essential
76 role of these cells in health and disease, the cellular mechanisms controlling their secretory function
77 and their survival are poorly understood. Closing this knowledge gap is thus of paramount importance
78 for designing new approaches to target this cell type.

79 During the transition from B cell to PC, the cell is reprogrammed to produce and secrete around 10^2 -
80 10^3 antibodies per second (1). To accommodate this large protein load, PCs expand their endoplasmic
81 reticulum (ER) and adapt to tolerate the extra stress induced via upregulation of the Ire1 α /Xbp1
82 branch of the unfolded protein response (UPR) (2–4). Early works from the 70s report that antibody
83 secretion happens via the conventional constitutive exocytosis pathway meaning that they are not pre-
84 stocked in granules but secreted as they are produced and trafficked from the ER to the Golgi
85 apparatus where they are glycosylated (5–7).

86 Here, through a proteomics analysis, we identified the SNARE Sec22b as an interesting molecule
87 upregulated in mouse and human PCs. Members of the Soluble N-ethylmaleimide-sensitive factor
88 Attachment Protein Receptor (SNARE) family are essential for the intracellular transport and fusion of
89 protein cargoes between organelles. They are involved in both regulated and constitutive exocytosis
90 and form a large family composed of different subtypes (Qa-, Qb-, Qc- and R-SNAREs) that interact
91 to form a ternary SNARE complex allowing membrane fusion between vesicles and organelles (8).
92 Distinct sets of SNAREs are expressed on the different organelles and can also be cell type specific
93 (9). On top of vesicular transport, Sec22b have other non-canonical functions, some of which could be
94 highly relevant for PC biology, including ER branching, nuclear shuttling of transcription factors as
95 well as plasma membrane expansion (10–13). Using a conditional KO mouse model, we demonstrated
96 that Sec22b is essential for PC maintenance and effective humoral response after vaccination and
97 infection. At the mechanistic level we showed that Sec22b governs expression of genes involved in
98 cell-cycle but also mitochondrial function and ER structure. Accordingly, Sec22b-deficient PCs

99 displayed a dramatically altered ER network, reduced contact between the ER and the mitochondria
100 and hyperfused mitochondria associated with poor survival. To date, little is known about how
101 organelle homeostasis is controlled in PCs and our results highlight Sec22b as a critical regulator of
102 this process with clear relevance for normal immune response regulation and for PC-mediated
103 diseases.
104

105 **Results:**

106 *The Sec22b SNARE and its partner are overexpressed in plasma cells:*

107 To identify new molecular actors that contribute to PC biology we performed a mass-spectrometry
108 analysis of sorted splenic B cells and *in vitro* generated PCs. All the data generated are available
109 through a user-friendly interface (<https://plasmacytomics.shinyapps.io/home/>). We first confirmed that
110 the expression of well-known factors of B cell and PC differentiation behaved as one would predict.
111 As expected, the B cell transcription factor Pax5 was significantly downregulated in PCs compared to
112 B cells while the transcription factor Xbp1 was significantly upregulated in PCs (Figure 1A).
113 Molecules involved in ER to Golgi trafficking, including SNAREs (Syntaxin 5 (Stx5), Sec22b and
114 Ykt6), the syntaxin regulator Sec1/Munc-18 protein (SCFD1/Sly1) and tethering factors
115 (USO1/p115), were also upregulated at the protein level in PCs compared to B cells (Figure 1A and
116 Supplementary Figure 1). We confirmed these results by western blotting and observed that the
117 SNAREs Stx5, Sec22b and Ykt6 were twice more expressed in PCs compared to B cells (Figure 1B).
118 Published microarray and RNAseq datasets were also included to the Plasmacytomics interface and in
119 support of our proteomic data, we also observed a significant increase in the levels of expression of
120 the genes encoding these three SNAREs in sorted murine PCs compared to B cells (Figure 1C). Data
121 mining of human RNAseq datasets also showed significantly increased expression of *SEC22B* and
122 *STX5A* in human PCs compared to B cells (Figure 1D) suggesting that this is a conserved feature in
123 PCs across species.

124 Previous works have shown that the ER-localised SNARE Sec22b forms a complex with the Golgi-
125 localised SNARE Stx5 to allow fusion of transport vesicles between these two organelles (9, 14, 15).
126 In absence of a suitable animal model, we first tested the functional relevance of Stx5 expression in
127 PC *in vitro* using a shRNA specific for Stx5 together with GFP reporter expression (Supplementary
128 Figure 2A-D). From day 3, the frequency of total PCs in the *Stx5* knock down (KD) samples
129 progressively diminished while it was constant in the control samples (Supplementary Figure 2C-D
130 left panel) and this decrease was due entirely to the loss of GFP⁺ PCs in the *Stx5* KD samples
131 (Supplementary Figure 2D central panel). The frequency of GFP⁺ B cells was not modified by *Stx5*
132 KD (Supplementary Figure 2D, right panel), confirming the specific requirement of *Stx5* expression

133 for PC persistence *in vitro*. We next took advantage of a small molecule, Retro-2, reported to block
134 Stx5 function by mislocalizing it and blocking its recycling (16, 17). When *in vitro* differentiated WT
135 PCs were cultured for 5 hours in presence of non-toxic doses of Retro-2 we observed a significant
136 reduction of antibody secretion (Supplementary Figure 2E). Moreover, injection of a single dose of
137 Retro-2 intraperitoneally to WT mice led to a reduction of the antibodies secreted by BM PCs *ex vivo*
138 (Supplementary Figure 2F), supporting a role for Stx5 in antibody secretion on top of PC survival.
139 Altogether our data suggest that the machinery required for ER to Golgi transport including
140 Stx5/Sec22b is upregulated during PC differentiation and that disruption of Stx5 leads to defects in
141 both PC viability and function.

142

143 *Dramatic loss of plasma cells and circulating antibodies in absence of Sec22b:*

144 Taking in consideration the effect of Stx5 on PC maintenance and antibody secretion, we next
145 explored the implication of its partner Sec22b. We crossed a *Sec22b* floxed mouse model (10) with the
146 mb1-cre strain (18) to generate a mouse model lacking *Sec22b* expression specifically in the B cell
147 lineage (*Sec22b*^{fllox/fllox} x mb1-cre, hereafter referred to as *Sec22b*^{B-KO}) (Supplementary Figure 3A-B).
148 Data mining of publicly available RNAseq datasets (Immgen) suggest that *Sec22b* is expressed at a
149 constant low level throughout B cell differentiation (Supplementary Figure 3C) so we investigated
150 whether *Sec22b* deficiency may alter this process. In absence of *Sec22b*, B cell development was
151 roughly normal with only a mild reduction of the number of BM mature B cells and splenic follicular
152 B cells compared to WT mice, whereas BM B cell precursors, splenic immature, marginal zone and
153 CD93⁺/CD21⁺/CD23⁻ B cells were unaffected (Supplementary Figure 3D-E). Despite this roughly
154 normal B cell development, *Sec22b*^{B-KO} mice had almost no circulating antibody at steady state for all
155 the isotypes tested (Figure 2A). Igκ representing around 90% of secreted antibodies in the mouse were
156 reduced over 40 times. IgG1 titres were 55 times lower in absence of *Sec22b*. IgM, IgA and IgG3
157 were 40, 30 and 20 times lower in absence of *Sec22b*, respectively (Figure 2A).

158 This was associated with a ten-fold reduction of the frequency and absolute number of PCs in the
159 spleen and the BM with the mature CD19⁺/B220⁺ PCs being the most affected (Figure 2B and C).
160 Accordingly, we observed that the few remaining splenic PCs in *Sec22b*^{B-KO} mice display a less

161 mature profile than their WT counterparts with reduced expression of the PC markers *Cd93* and
162 *Tnfrsf13b* and of the PC master regulators *Prdm1*, *Xbp1* and *Irf4*. In contrast enhanced expression of
163 the B cell master regulators *Pax5* and *Bach2* were detected at the transcriptional level in *Sec22b*^{B-KO}
164 PCs compared to WT (Figure 2D). Thus, our results demonstrate that *Sec22b* plays a critical and non-
165 redundant role in PC maintenance and in the control of antibody circulating titres at steady state.

166

167 *Sec22b regulates plasma cell maintenance and molecular identity:*

168 To unravel at which step of PC differentiation *Sec22b* is required, we performed *in vitro*
169 differentiation assays of splenic B cells. In control cultures PC frequency doubles between day 2 and
170 day 4, whereas it remains constant between these two time points in *Sec22b*^{B-KO} cultures (Figure 3A-
171 B). This was associated with an enhanced apoptosis of *in vitro* derived PCs lacking *Sec22b* as detected
172 by the frequency of active Caspase3⁺ cells (Figure 3C). In addition, cell cycle was impaired in
173 *Sec22b*^{B-KO} PCs generated *in vitro* with a progressive exit from the cell cycle particularly clear at day 4
174 (Figure 3D). These results indicate that *Sec22b* is not required for the initiation of PC differentiation
175 but is important for the maintenance and the expansion of this cell subset.

176 To gain further insight into how *Sec22b* controls PC fate we performed RNAseq analysis on *in vitro*
177 generated PCs. We chose an early time point, 2 days after LPS stimulation, to detect defective
178 mechanisms before the loss of PCs observed rapidly thereafter. Despite being numerically normal
179 (Figure 3E-F), *Sec22b* deficient PCs at day 2 post stimulation were transcriptionally very different
180 from their WT counterparts as shown by unsupervised analyses (Figure 3E-F and Supplementary
181 Tables 1 and 2). Over 6000 genes were differentially regulated between both genotypes with 3388
182 genes up-regulated and 3184 genes down-regulated in *Sec22b*^{B-KO} PCs compared to WT PCs (14026
183 genes total, fold change >1.2 and q-value < 0.05) (Figure 3G). Although the cell purity was equivalent
184 between sorted *Sec22b*^{WT} and *Sec22b*^{B-KO} PCs (>80%), we cannot formally exclude that the
185 differences observed result from heterogeneity in the cells generated *in vitro* at this early step in both
186 conditions. Transcriptional expressions of *Ptprc* (encoding B220) and *Sdc1* (encoding CD138) were
187 equivalent between samples from *Sec22b*^{WT} and *Sec22b*^{B-KO} PCs, suggesting that, at least based on
188 these two markers, the sorted cells were homogeneous between the two groups (Supplementary figure

189 4A-B). Sec22b was recently proposed to contribute to transcription factor nuclear shuttling and this
190 might account for the important transcriptional differences observed in our datasets(13). Gene set
191 enrichment analyses (GSEA) revealed that several pathways were significantly different between WT
192 and Sec22b^{B-KO} PCs. In line with our experimental results (Figure 3H), Sec22b seems essential for
193 promoting cell cycle possibly via control of the Myc/E2F signalling axes. Pathways pertaining to
194 “Mitochondria” were also significantly downregulated in Sec22b^{B-KO} compared to WT PCs, whereas
195 “UPR”, “ER-Golgi transport” and “protein secretion” pathways were upregulated (Figure 3H and
196 Supplementary Figures 5 and 6). We thus explored more precisely how *Sec22b* deficiency may affect
197 these processes. We first observed that the expression of the second R-SNARE normally involved in
198 ER to Golgi trafficking, Ykt6, was conserved in absence of Sec22b. Moreover, expression of all the
199 Qa, Qb and Qc SNAREs with whom Sec22b or Ykt6 interact to form a trans-SNARE complex was
200 significantly increased in Sec22b^{B-KO} PCs suggesting the induction of compensatory mechanisms to
201 deal with cargo transport in absence of Sec22b (Figure 3I).

202 To unravel whether Sec22b intrinsically regulates antibody secretion we took advantage of a droplet
203 microfluidic-based technique to assess the secretion rate at the single cell level (1). Between 6000 and
204 10000 cells were individually encapsulated into droplets and analysed for their IgM secretion rate after
205 2- and 4-days of *in vitro* culture. Up to 12% of encapsulated cells secreted detectable amount of IgM
206 over the 40 minutes of imaging with less secreting cells in the Sec22b^{B-KO} condition. We observed that
207 *in vitro* generated WT PCs secreted on average 12 IgM/sec at day 2 and 20 IgM/sec at day 4. In
208 contrast, *Sec22b* deficient PCs secreted on average 2 to 4 times less IgM/sec at day 2 and 4,
209 respectively (Figure 3J). Hence, *Sec22b* deficiency leads to a significant reduction, albeit not a total
210 block, in antibody secretion. The massive drop in antibody titre in the serum of *Sec22b* deficient mice
211 is thus probably caused by the combination of reduced secretion rate together with severely altered PC
212 maintenance.

213

214 *Sec22b* is essential for ER expansion and structure in PCs:

215 Our transcriptional data indicate that in absence of Sec22b, all three branches of the UPR (i.e. *Ern1*,
216 *Atf4/Ddit3*, *Perk*, *Atf6*) (Figure 4A-B) were upregulated including the Atf4/Chop pathway normally

217 actively repressed in PCs to protect from ER-stress induced cell death (4). Our results suggest that in
218 absence of Sec22b this protective mechanism is deficient and could lead to Chop-mediated cell death
219 and exit from the cell cycle. Exacerbated UPR may be a sign of enhanced ER stress and we thus
220 assessed whether ER biogenesis might be altered. Using a permeable cell tracker, we showed that the
221 ER membrane expansion normally observed as PC differentiate was significantly reduced in absence
222 of Sec22b (Figure 4C). The structure of the ER was also altered in Sec22b^{B-KO} PCs. Whereas the
223 perinuclear envelope was roughly normal, peripheral ER appeared poorly branched with dilated
224 cisternae (Figure 4D), suggestive of accumulation of hyper-dilated ER and loss of the parallel rough
225 ER typically seen in WT PCs. Electron microscopy confirmed that the stacked ER sheets characteristic
226 of PCs were lost in absence of Sec22b with a hyper-dilatation of the ER cisternae and a defective
227 stacking that was even more evident at day 4 (Figure 4E). Dilated ER was observed in all but one PC
228 in Sec22b^{B-KO} PCs while it was observed in less than 11% of Sec22b^{WT} PCs. Sec22b expression is thus
229 crucial for ER spatial expansion and organisation during PC differentiation.

230

231 *Sec22b deficiency affects PC fitness via altered mitochondrial dynamics:*

232 In addition to protein folding and export, an important function of the ER is the control of organelle
233 dynamics and in particular of mitochondria. Indeed, ER-mitochondria membrane contact sites are
234 pivotal for ER-mitochondria exchange but also for mitochondrial fission and thus function (19).
235 Through our RNAseq analysis we observed a downregulation of several genes involved in
236 mitochondrial function and dynamics (Figure 3H and Supplementary Figure 5) including *Drp1*, *Inf2*
237 and *Spire1* that encode proteins involved in mitochondrial fission at the ER-mitochondria contact site
238 (Figure 5A). We thus quantified ER-mitochondria contact sites by proximity ligation assay (PLA -
239 measuring the distance between the mitochondrial protein VDAC1 and the ER protein IP3R1) and
240 revealed that they were significantly reduced in Sec22b^{B-KO} PCs (mean 1.233+/-0.06910 for Sec22b^{B-}
241 ^{WT} vs. 0.8571+/-0.05122 for Sec22b^{B-KO}) (Figure 5B and Supplementary Figure 7). Moreover,
242 mitochondrial content and potential were increased in PCs after 4 days in culture in Sec22b^{B-KO} PCs
243 compared to WT PCs (Figure 5C). Accordingly, we detected more mitochondria, increased total
244 mitochondrial area but with fewer fragments in Sec22b^{B-KO} PCs compared to WT PCs suggesting an

245 hyperfused mitochondrial phenotype (Figure 5D-E). We wondered whether this defective
246 mitochondrial conformation may contribute to the PC loss observed in absence of Sec22b. Treatment
247 of WT PCs with M1, a Drp1 antagonist, together with Mdivi, an agonist of mitochondrial fusion,
248 promoted the generation of a hyperfused mitochondria phenotype comparable to that of Sec22b^{B-KO}
249 PCs (Figure 5F). Moreover, M1+Mdivi-treated WT PCs numbers were significantly reduced in culture
250 compared to control conditions, confirming that mitochondrial dynamics affect PC fitness (Figure
251 5G). In addition to this hyperfused phenotype Sec22b^{B-KO} PCs presented several abnormalities in
252 mitochondrial gene expression (Figure 5H) including genes implicated in the respiratory chain (*Atp5*
253 family), mitochondrial fission (*Mtfp1* and *Mtfr2*), structure (*Timm* and *Tomm* families) and metabolite
254 transport (*Slc25* family) (Figure 5H) suggesting a global alteration of mitochondrial size and function
255 in absence of Sec22b. Altogether, these results demonstrate that *Sec22b* expression regulates ER-
256 mitochondria contact sites and consequently mitochondrial dynamics in PCs with important
257 consequences for cell fitness and survival.

258

259 *Sec22b is indispensable for the generation of a protective humoral immune response:*

260 Considering the importance of PCs and circulating antibodies for the humoral immune response, we
261 finally assessed the impact of *Sec22b* deficiency on these processes. Following T-dependent
262 immunization with sheep red blood cells (SRBC) (Figure 6A-C), the frequency and number of PCs in
263 the spleen remained extremely low in Sec22b^{B-KO} mice, being reduced over 100 times compared to
264 controls (Figure 6B). A similar observation was made for antibody titres after SRBC immunization
265 (Figure 6C). We also investigated antigen specific humoral immune response by immunizing ip with
266 the T-dependent antigen NP-KLH in alum and boosting with NP-KLH only (Figure 6D). Seven days
267 after the boost, the frequency and numbers of splenic and BM PCs were again dramatically reduced in
268 absence of Sec22b (Figure 6E). Moreover, we barely detected NP-specific IgM and IgG1 antibodies in
269 the serum of Sec22b^{B-KO} mice throughout the immunization while a potent antibody immune response
270 with the expected kinetics was observed in control animals (Figure 6F). We next infected WT and
271 *Sec22b* deficient mice with influenza A virus (Figure 6G) and observed a profound defect in the
272 frequency and number of PCs in the draining mediastinal lymph nodes (Figure 6H). Flu-specific

273 antibodies were undetectable in the serum of *Sec22b* deficient mice (Figure 6I). These defects were
274 associated with exacerbated weight loss in *Sec22b*^{B-KO} mice compared to their WT littermates
275 suggesting a poorer control of the infection in absence of *Sec22b* (Figure 6J). Altogether, these results
276 establish that *Sec22b* is essential and play a non-redundant role for the establishment of a potent and
277 efficient humoral immune response after both vaccination and infection.

278

279

280 **Discussion**

281 Despite their essential role in health and disease, how PCs maintain a high rate of antibody secretion
282 while persisting on the long term is unclear. Here we demonstrate that the SNARE Sec22b is essential
283 for PC maintenance. Accordingly, in absence of *Sec22b* expression in the B cell lineage, the humoral
284 immune response was severely impaired after both vaccination and infection.

285 The reduced antibody secretion rate observed could be caused by a defective ER to Golgi transport
286 due to altered formation of the Stx5/Sec22b SNARE complex. Supporting a role for this complex in
287 this process, PC treatment with Retro-2, a drug known to alter Stx5 cellular localisation (16), also
288 leads to decreased antibody secretion. However, after Retro-2 treatment but also in absence of Sec22b,
289 antibody secretion was reduced but not totally abrogated. Like Sec22b, the R-SNARE Ykt6 has been
290 shown to form a trans SNARE complex with the Qa, Qb and Qc SNAREs Gosr1/2, Stx5 and Bet1 and
291 could partially compensate for Sec22b loss (20, 21). All these SNAREs are strongly expressed in PCs
292 and the four Q-SNAREs are twice more expressed in Sec22b^{B-KO} PCs thus suggesting that redundant
293 mechanisms may exist to export antibodies at a suboptimal level via the classical constitutive or via
294 unconventional secretion pathways (22, 23).

295 Importantly, the altered PC maintenance in absence of Sec22b cannot uniquely be linked to the
296 defective secretory capacity of the cells as several studies report that PC survival and antibody
297 secretion are uncoupled. For example, mice lacking Xbp1 or the ligase Rctb, important for Xbp1
298 activation, display reduced antibody secretion but normal PC numbers (24–27). Moreover, PC
299 producing pathogenic antibodies often fail to secrete them while the PC compartment is not altered in
300 these mouse models (28). Another example is PCs generated in the LMP2A model that can survive
301 without secreting any antibody reinforcing the independence between antibody secretion and PC
302 survival (29). Additive defects must thus explain the poor persistence of PCs in absence of Sec22b.
303 We report here that *Sec22b* deficiency is associated with a profound alteration of the PC
304 transcriptional profile and with deregulation of the UPR as well as morphological alterations of the ER
305 and mitochondria.

306 In PCs, the Ire1 α /Xbp1 axis is normally highly expressed and favours antibody production and folding
307 without causing cell death (2, 30) whereas Atf6 is weakly expressed and the Perk-dependent branch of

308 the UPR is repressed (4, 31). Activation of the Perk/Atf4/Chop pathway has been associated with
309 increased susceptibility to cell death in PCs and is normally suppressed by the Ire1 α /Xbp1-dependent
310 Ufbp1 protein (32). In absence of Sec22b the upregulation of Perk may contribute to the inhibition of
311 the cell cycle we observed (33) and eventually to PC cell death (34) hence explaining, in part, the poor
312 PC maintenance *in vitro* and *in vivo* in absence of Sec22b.

313 An important observation made in absence of Sec22b was the disturbed ER structure with poorly
314 stacked and hyper-dilated cisternae. This could be the consequence of the reduced transport of
315 antibodies from the ER to the Golgi apparatus but could also be caused directly by the absence of
316 Sec22b as was shown in yeast and drosophila (35–37). In yeast, Sec22p interacts with Sey1p to
317 regulate sterol biosynthesis and with atlastins to support ER homotypic fusion (35, 36). In the
318 drosophila, Sec22 was identified as an important regulator of ER structure with a dilatation of the ER
319 when *Sec22* was mutated or knocked-down (37). More recently, *Sec22b* knock-down in HUVECs was
320 also associated with dilated ER but the mechanism at play remains unclear (38). Interestingly, the long
321 form of Stx5 was also shown to be important for ER branching via interaction with Climp63 that links
322 the ER membrane to the microtubules (11). Thus, Sec22b could directly control ER morphology in
323 PCs either by interacting with the long form of Stx5 or via another partner still to identify.

324 Another striking defect observed in *Sec22b* deficient PCs was the accumulation of hyperfused
325 mitochondria and the reduction of ER/mitochondria contact sites that regulate mitochondrial fission
326 (19). Sec22b, through its longin domain, has already been shown to regulate ER/plasma membrane
327 contact sites and consequently plasma membrane expansion in neurites (12, 39). Whether Sec22b
328 contributes to ER/mitochondria contact sites directly or indirectly remains to be established. In
329 addition, we demonstrated that hyperfused mitochondria were associated with poor PC survival hence
330 suggesting that altered mitochondrial dynamics contribute to the decreased PC number in absence of
331 Sec22b perhaps through metabolic rewiring.

332 To conclude, we demonstrated that Sec22b is a crucial regulator of PC survival. Altogether, our results
333 demonstrate that Sec22b-mediated regulation of organelle dynamics is indispensable for PC biology
334 and for the establishment of a protective humoral immune response (Supplementary Figure 8).
335 Discovery of pharmacological regulators of SNAREs is an active field of research notably for

336 neurological disorders and toxin antagonisms (40–42). Specifically targeting this molecule in a tissue
337 and time-controlled manner may thus constitute an attractive avenue for PC-mediated diseases.

338

339

340 Acknowledgments

341 We thank Dr. N. Setterblad, C. Doliger and S. Duchez (Plateforme technologique IRSL, Paris,
342 France), Dr. V. Parietti (Mouse facility IRSL, Paris, France), S. Guibert (Integragen), M. Khamyath
343 and V. Gourhand for their technical assistance. We are grateful to Pr. A. Toubert for his comments on
344 this manuscript. The study was supported by the Laboratory of Excellence in Research on Medication
345 and Innovative Therapeutics (LabEx LERMIT) (ME and KB), an ANR JCJC grant (ANR-19-CE15-
346 0019-01), an ANR @RAction grant (ANR-14-ACHN-0008), a “Fondation ARC pour la recherche sur
347 le cancer” grant (P JA20181208173) and a grant from IdEx Université Paris-Cité (ANR-18-IDEX-
348 0001) to ME, an ANR PRC grant (ANR-17-CE14-0019) and an INCa grant (PRT-K 2017) to KB.
349 P.B. acknowledges funding from the French National Research Agency grant ANR-18-CE15-0001
350 project Autoimmuni-B, by the Institut Carnot Pasteur Microbes et Santé grant ANR-11-CARN-0017-
351 01, the Institut Pasteur and the Institut National de la Santé et de la Recherche Médicale (INSERM).
352 M.A.L is supported by Biotechnology and Biological Sciences Research Council
353 (BBS/E/B/000C0427, BBS/E/B/000C0428, and the Campus Capability Core Grant to the Babraham
354 Institute). AP and MC were supported from a grant from the Biotechnology and Biological Sciences
355 Research Council (BB/L022389/1). DLH is supported by a National Health and Medical Research
356 Council Australia Early-Career Fellowship (APP1139911). NA was supported by a PhD fellowship
357 from the French Ministry for education and by a 4th year PhD fellowship from the “Fondation ARC
358 pour la recherche sur le cancer”. P.C-H. was supported partly by a stipend from the Pasteur - Paris
359 University (PPU) International PhD program, and by a fellowship from the French *Fondation pour la*
360 *Recherche Médicale* (FRM). KGCS was supported by the Wellcome Trust (Programme Grant Number
361 083650/Z/07/Z). J.J’s research is supported by the German Research Foundation project number:
362 419193696 and through the CRC1335. H.T’s is supported through the graduate school of the Max
363 Planck Institute for Immunobiology and Epigenetics (IMPRS-IE) and through the CRC1335. The
364 "EMiLy" U1160 INSERM unit is a member of the OPALE Carnot institute, The Organization for
365 Partnerships in Leukemia (Institut Carnot OPALE, Institut de Recherche Saint-Louis, Hôpital Saint-
366 Louis, Paris, France. Web : www.opale.org. Email : contact@opale.org).

367

368 **Author Contributions**

369 AB designed and performed experiments, analyzed data, and wrote the manuscript. LG, SMG, ER,
370 PCH, DS, DLH, NR, MC and MAL performed experiments and analyzed data. NA helped with
371 experiments. SB and VS performed electron microscopy experiments. AA, DG, JB, PB and SA
372 provided essential reagents and technologies. J.J. and H.T. planned, performed and analyzed the PLA
373 assays. KGCS, FB, KB, MAL contributed to the project design and to the manuscript redaction. AAP
374 contributed to the project design and to the manuscript redaction and set up the PlasmacytOMICs
375 platform. ME designed the project, analyzed data, and wrote the manuscript. All authors had the
376 opportunity to review and edit the manuscript.

377

378

379 **Declaration of Interests**

380 The authors declare that no conflict of interest exists.

381

382

383 **Methods**

384 *Mouse model, immunisation, and infection*

385 The Sec22b^{fl/fl} mice (C57Bl/6J background) were obtained from Dr. Sebastian Amigorena and crossed
386 with the mb1-Cre mice (18) (C57Bl/6J background) purchased from The Jackson Laboratory. All
387 experiments were conducted in compliance with the European Union guide for the care and use of
388 laboratory animals and has been reviewed and approved by appropriate institutional review
389 committees (C2EA-26, Animal Care and Use Committee, Villejuif, France and Comité d'éthique
390 Paris-Nord N°121, Paris, France). In all experiments Sec22b^{WT} controls were mb1cre⁺ or
391 Sec22b^{fl/fl} / Sec22b^{fl/+} mice in equal proportions. Immunizations/infections were performed
392 intraperitoneally (ip) with 100µg of 4-hydroxy-3-nitrophenylacetyl-keyhole limpet hemocyanine (NP-
393 KLH) (Biosearch Technologies) adjuvanted with alum (Inject Alum, Thermo Scientific) or with
394 200µL of Sheep Red Blood Cells (SRBC) (Eurobio) or intranasally with 10⁴ plaque-forming units of
395 influenza A/HK/x31 virus (H3N2) under inhalation anaesthesia with isoflurane. Influenza infections
396 were performed on chimeric mice. CD45.1 Recipient mice were lethally irradiated with 11gy before
397 reconstitution with bone marrow cells from Sec22b^{B-KO} or Sec22b^{B-WT} mice. Chimera were infected 8
398 to 12 weeks later once reconstitution was complete.

399

400 *Proteomic analysis*

401 Splenic B cells were purified using a mouse B cell isolation kit following the manufacturer's
402 instructions (Miltenyi Biotec) and cultured for 3 days with 10µg/ml lipopolysaccharide (Sigma). The
403 differentiated cells were enriched using anti-CD138 antibodies coupled to PE and anti-PE magnetic
404 beads (Miltenyi Biotec). Purified cells were lysed into FASP buffer. The protein concentration of the
405 samples was normalised, digested using the FASP method (43) using trypsin and 100µg was
406 fractionated into 36 fractions by high pH reverse phase chromatography, pooled into 12 fractions and
407 peptides were analysed using LC-MS/MS on an Orbitrap Elite (Thermo Fisher) hybrid mass
408 spectrometer equipped with a nanospray source, coupled with an Ultimate RSLCnano LC System
409 (Dionex). MS data was analysed using MaxQuant (44) version 1.5.8.3. Data was searched against
410 mouse UniProt sequence databases. Full technical details can be found in the supplementary material.

411

412 *Cell preparation and flow cytometry*

413 Spleen and bone marrow cells were isolated as previously described (45, 46). Briefly, single cell
414 suspensions were stained with appropriate antibodies (Supplementary table 3) in PBS supplemented
415 with 2% BSA and 2mM EDTA for cell surface staining. Staining with ER tracker (Invitrogen),
416 mitoTracker Green (Invitrogen) and mitoTracker Orange CMTMRos (Invitrogen) were performed as
417 recommended by the supplier. Intracellular staining was performed using the FoxP3/Transcription
418 factor staining buffer set (eBioscience) according to the provider recommendation. Flow cytometry
419 analyses were performed on a BD LSR Fortessa cytometer and cell sorting experiments for RNAseq
420 and Biomark analysis were performed using a BD FACS AriaIII cell sorter. Data were analysed with
421 the Flowjo software (TreeStar, Ashland, OR).

422

423 *In vitro cell differentiation and infection of primary cells*

424 1×10^6 splenocytes were stimulated with $5 \mu\text{g}/\text{mL}$ of lipopolysaccharide (LPS) (InVivoGen) for 2 or 4
425 days in complete culture medium composed of RPMI supplemented with 10% foetal calf serum
426 (Sigma), 0.05mM 2-mercaptoethanol, 100U/mL penicillin-streptomycin, 1mM sodium pyruvate and
427 0.1mM non-essential amino acids (Gibco). Where indicated, cells were treated with $20 \mu\text{M}$ M1 and
428 $10 \mu\text{M}$ Mdivi (Sigma) for 2 days from day 2 to day 4.

429 *Stx5a* specific shRNAs were designed thanks to the RNAi consortium
430 (<https://portals.broadinstitute.org/gpp/public>) and cloned in the pLKO.3G (addgene #14748) vectors.
431 Lentiviral particles were produced in the Human Embryonic Kidney 293T (HEK293T) cell line with
432 the psPAX2 (Addgene #12260) and pMD2.G (Addgene #12259) vectors. For primary cell
433 transduction, splenocytes were put in culture in complete culture medium in presence of 80ng/ml
434 CD40L (Thermoscientific) and 1U/mL IL-4 (Miltenyi) for 24 hours to promote B cell entry into cycle.
435 The cells were then washed and transduced with lentiviral particles together with polybrene. After
436 24hours cells were washed and differentiated into PCs by addition of LPS as indicated above.

437

438 *Western blot*

439 Cells were resuspended in RIPA lysis buffer supplemented with Protease and Phosphatase Inhibitor
440 (Thermofisher). Total proteins were quantified with Bradford buffer (Thermofisher). 15µg of proteins
441 were separated on a NuPAGE™ 4-12% Bis-Tris Gel (Invitrogen) and transferred to a PVDF
442 membrane. Primary antibodies (Supplementary Table 3) or β-actin (Cell Signaling) were incubated
443 overnight at 4°C. Secondary antibodies (Supplementary Table 3), specific for the primary antibody
444 species, conjugated to horseradish peroxidase (HRP) were incubated 2h at room temperature and
445 detected using Pierce ECL (Thermofisher) and signal was quantified by ChemiDoc™ Touch Gel
446 Imaging System (BIO RAD).

447

448 *Confocal microscopy*

449 Cells were loaded on poly-Lysine (Sigma) coated slides and fixed in PBS/4% PFA prior to analysis.
450 Cells were permeabilized with PBS/0.3% Triton, washed in PBS and incubated with primary
451 antibodies coupled or not overnight at 4°C and then with secondary antibodies 1 hour at room
452 temperature, when necessary (Supplementary Table 3). Cells were incubated with Hoescht
453 (Thermofisher) for 1 hour at room temperature. Images were acquired using a LSM800 confocal
454 microscope equipped with the Airyscan system (Zeiss) using a 63x objective and z-stacks. Images
455 were analysed with Fiji and Zen.

456

457 *Electron microscopy*

458 Cells were pre-fixed with 2% glutaraldehyde/2% PFA in Sorensen phosphate buffer 0.1M pH7.2 for
459 15 minutes before being fixed with 2.5% glutaraldehyde/2% PFA in Sorensen phosphate buffer 0.1M
460 pH7.2 for 2 hours at room temperature. Cells were then washed in Sorensen phosphate buffer,
461 resuspended in PFA 1% and sent to the METi platform (Toulouse, France). Cells were then post-fixed
462 with 1% OsO4 in Cacodylate buffer (0.1 M, pH 7.2, EMS, Hatfield, PA), rinsed in the same buffer and
463 pelleted, concentrated in agarose, and treated for 1 h with 2% aqueous uranyl acetate. The samples
464 were then dehydrated in a graded ethanol series and embedded in Epon. After 48 h of polymerization
465 at 60 °C, ultrathin sections (80 nm thick) were mounted on 200 mesh Formvar-carbon-coated copper
466 grids. Finally, sections were stained with Uranylless and lead citrate. Grids were examined with a TEM

467 (Jeol JEM-1400, JEOL Inc, Peabody, MA, USA) at 80 kV. Images were acquired using a digital
468 camera (Gatan Orius, Gatan Inc, Pleasanton, CA, USA).

469

470 *ELISA*

471 ELISA assays were performed as previously described (47) for the determination of the different
472 isotype of Ig or NP-specific antibody titres in sera or culture supernatants. Briefly, plates were pre-
473 coated with goat anti-mouse IgM (Southern Biotech) or with NP(15)-BSA (Biosearch Technologies).
474 After a step of saturation in PBS/2%BSA, diluted sera/supernatant were added before incubation with
475 HRP-conjugated secondary antibody. Enzymatic revelation was performed with the TMB substrate
476 reagent set (BD OptEIA). All Antibodies used are indicated in Supplementary table 3.

477

478 *Proximity Ligation Assay (PLA)*

479 Anti-VDAC1 (20B12AF2) (abcam) and Anti-IP3R1 (E-8) (Santa Cruz Biotechnology) antibodies
480 were used to generate PLA probes using Duolink in Situ Probemaker (Sigma). B cells were stimulated
481 with Lipopolysaccharide (LPS) (InVivoGen) (5µg/ml) for 2 days. Stimulated cells were settled on
482 PTFE diagnostic slides (Thermo Fisher Scientific) for 30 min at 37°C, fixed with 4 %
483 paraformaldehyde (Thermo Fisher Scientific) in PBS for 15 min, permeabilized with 0.5% saponin
484 (quillaja bark, Sigma) in PBS for 30 min and blocked with Duolink Blocking Solution (Sigma) for 1
485 hour. Cells were incubated with Anti-VDAC1 (1/100) and Anti-IP3R1 (1/100) PLA probes overnight
486 at 4°C. PLA was performed according to the manufacturer's instructions (Duolink, Sigma). Later,
487 cells were further incubated with anti-mouse IgM-FITC antibody (1/100) (Jackson Immuno Research)
488 for 30 min at 4°C. Resulting samples were mounted in Fluoromount-G with DAPI (Invitrogen).
489 Fluorescence was detected by using an inverted confocal microscope, Leica TCS SP8 (Leica). PCs
490 were detected based on their high IgM expression. Quantification of detected signal per nucleus in
491 these cells was performed manually.

492

493 *Hemagglutination inhibition (HAI) assay*

494 Antibody titers pre and post vaccination were determined using the hemagglutination inhibition (HAI)
495 assay using the standard WHO protocol, as previously described (48). Sera were treated overnight
496 with receptor-destroying enzyme (Denka Seiken Co.) and were subsequently tested by standard
497 methods using 4 HA units of virus and a 0.5% suspension of turkey red blood cells. HAI titers were
498 recorded as the reciprocal of the highest dilution of the serum that completely inhibited agglutination
499 of erythrocytes by 4 HA units of the virus.

500

501 *Biomark-based transcriptomic analysis*

502 Multiplex qPCR analysis was performed using the Biomark system (Fluidigm). Cells were sorted at
503 100 cells/well directly into PCR tubes containing 5 μ L of reverse transcription/pre-amplification mix
504 as previously described (45). Briefly, the mix contained 2X Reaction mix and SuperscriptIII
505 (CellDirect One-Step qRT-PCR kit, Invitrogen) and 0,2X Taqman assay (Life technologies)
506 (Supplementary Table 4). Targeted cDNA pre-amplification was performed for 19 cycles before
507 processing with Dynamic Array protocol according to the manufacturer's instructions (Fluidigm).
508 Wells positive for *Gapdh*, *Actb* and control gene (*Prdm1*, *Irf4* and *Xpb1* for PCs and *Pax5* for B cells)
509 expression and negative for expression of a control gene (*Cd3e*) were considered for further analysis.
510 Mean expression of *Actb* and *Gapdh* was used for normalization. Unsupervised clustering (Spearman
511 rank correlation test) and heatmap representation were generated with (<http://www.heatmapper.ca>)
512 using the Z scores.

513

514 *RNAseq and RNAseq-based transcriptomic analysis*

515 For RNAseq, *in vitro* generated PCs were sorted at day 2 and RNA were extracted with "RNeasy plus
516 micro kit" (Qiagen) as recommended. RNAs were proceeded by the Integragen company for
517 sequencing. Libraries were prepared with NEBNext® Ultra™ II Directional RNA Library Prep Kit for
518 Illumina protocol according supplier recommendations. Briefly the key stages of this protocol were
519 successively, the purification of PolyA containing mRNA molecules using poly-T oligo attached
520 magnetic beads from 100ng total RNA (with the Magnetic mRNA Isolation Kit from NEB), a
521 fragmentation using divalent cations under elevated temperature to obtain approximately 300bp

522 pieces, double strand cDNA synthesis and finally Illumina adapters ligation and cDNA library
523 amplification by PCR for sequencing. Sequencing was then carried out on Paired End 100b reads of
524 Illumina NovaSeq. Image analysis and base calling was performed using Illumina Real Time Analysis
525 (3.4.4) with default parameters.

526 For analysis STAR was used to obtain the number of reads associated to each gene in the Gencode
527 vM24 annotation (restricted to protein-coding genes, antisense and lincRNAs). Raw counts for each
528 sample were imported into R statistical software. Extracted count matrix was normalized for library
529 size and coding length of genes to compute FPKM expression levels. The Bioconductor edgeR
530 package was used to import raw counts into R statistical software, and compute normalized log2 CPM
531 (counts per millions of mapped reads) using the TMM (weighted trimmed mean of M-values) as
532 normalization procedure. The normalized expression matrix from the 500 most variant genes (based
533 on standard deviation) was used to classify the samples according to their gene expression patterns
534 using principal component analysis (PCA), hierarchical clustering and consensus clustering. PCA was
535 performed by FactoMineR::PCA function with “ncp = 10, scale.unit = FALSE” parameters.
536 Hierarchical clustering was performed by stats::hclust function (with euclidean distance and ward.D
537 method). Differential expression analysis was performed using the Bioconductor limma package and
538 the voom transformation. To improve the statistical power of the analysis, only genes expressed in at
539 least one sample (FPKM \geq 0.1) were considered. A q value threshold of \leq 0.05 and a minimum fold
540 change of 1.2 were used to define differentially expressed genes. Gene list from the differential
541 analysis was ordered by decreasing log2 fold change. Gene set enrichment analysis was performed by
542 clusterProfiler::GSEA function using the fgsea algorithm. Gene sets from MSigDB v7.2 database were
543 selected among the C2_curated and Hallmark classes, keeping only gene sets defined by 10-500
544 genes.

545

546 *DropMap-based analysis of IgM secretion*

547 Dropmap experiment were performed as described(1), modified to detect IgM secretion from single
548 cell. Cells from *in vitro* cultures at day2 or day 4 after LPS stimulation were centrifuged and
549 resuspended in DropMap medium (RPMI without phenol red, supplemented with 0.1% Pluronic F68,

550 25 mM HEPES pH 7.4, 10% KO serum replacement (all ThermoFisher) and 0.5% recombinant human
551 serum albumin (Sigma Aldrich). Microfluidic droplets were generated as water-in-oil emulsions using
552 a co-flow of aqueous phases, one containing bioassay reagents (bioassay phase) and the other one
553 containing *in vitro*-generated PCs (cell phase). *Bioassay phase*: Streptavidin-coated paramagnetic
554 beads (300nm, Ademtech) were washed with PBS using a magnet, and then resuspended in a 1 μ M
555 solution of CaptureSelect biotin anti-mouse Ig κ conjugate (ThermoFisher) and incubated at room
556 temperature (RT) for 20 min. The beads were washed with PBS and resuspended in 5% pluronic F127
557 (ThermoFisher) and incubated for 20 min at room temperature. Following a PBS wash, the beads were
558 resuspended in DropMap buffer and incubated at RT for 20 min. Beads were washed for the last time
559 with PBS and resuspended in a solution of 150nM Goat anti-mouse IgM (μ chain specific) F(ab')₂
560 (Alexa647, Jackson ImmunoResearch) in DropMap buffer. *Cell phase*: To produce single-cell
561 droplets, cell concentration was adjusted to achieve 0.3 cells per droplet in DropMap buffer. For
562 calibration purposes with cells, purified monoclonal IgM was diluted in DropMap buffer.
563 Droplets were produced by hydrodynamic flow-focusing on a custom-made microfluidic device as in
564 (1), by co-flowing the two aqueous phases. Immediately after generation, droplets were injected into
565 the 2D observation chamber (1) until filled, which was then closed for image acquisition. The droplet
566 array in the 2D chamber was imaged using a Nikon Ti-2 Eclipse inverted microscope with motorized
567 stage and excitation light source (Lumencor Spectra X). Fluorescence was captured using a 10x
568 objective and a Cy5 filter, and images were recorded by a digital CMOS camera (ORCA-flash 4.0,
569 Hamamatsu). For each time point, an array of 10x10 images was acquired, and a total of 6 acquisitions
570 were recorded through 37.5 min. The images were analyzed with a custom Matlab script as described
571 (1). In brief, the ratio of fluorescent signal between the beadline and the background was estimated for
572 every droplet at every time point. The fluorescence ratio was then used to estimate the concentration
573 of IgM in the droplets by using a calibration curve, which was generated by measuring the fluorescent
574 ratio from different concentrations of purified monoclonal IgM antibody.
575
576 *Statistical analysis*

577 The p-values were determined as indicated in the figure legends using the Prism GraphPad software
578 with the two-tailed unpaired Mann-Whitney non-parametric test for the WT vs Sec22b^{B-KO} comparison
579 ($*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$, “ns” = non-significant p -value), or with the 2way
580 ANOVA with Sidak correction for multiple comparisons (**££** $p < 0.01$, **£££** $p < 0.001$).

581

582

583 **References**

- 584 1. K. Eyer, *et al.*, Single-cell deep phenotyping of IgG-secreting cells for high-resolution immune
585 monitoring. *Nat. Biotechnol.* **35**, 977–982 (2017).
- 586 2. N. N. Iwakoshi, *et al.*, Plasma cell differentiation and the unfolded protein response intersect at
587 the transcription factor XBP-1. *Nat Immunol* **4**, 321–329 (2003).
- 588 3. S. J. Kirk, J. M. Cliff, J. A. Thomas, T. H. Ward, Biogenesis of secretory organelles during B
589 cell differentiation. *J Leukoc Biol* **87**, 245–255 (2010).
- 590 4. Y. Ma, Y. Shimizu, M. J. Mann, Y. Jin, L. M. Hendershot, Plasma cell differentiation initiates a
591 limited ER stress response by specifically suppressing the PERK-dependent branch of the
592 unfolded protein response. *Cell Stress and Chaperones* **15**, 281–293 (2010).
- 593 5. F. Melchers, Biosynthesis, transport and secretion of immunoglobulin in plasma cells.
594 *Histochem. J.* **3**, 389–397 (1971).
- 595 6. E. Reales, *et al.*, Identification of Soluble N -Ethylmaleimide-Sensitive Factor Attachment
596 Protein Receptor Exocytotic Machinery in Human Plasma Cells: SNAP-23 Is Essential for
597 Antibody Secretion. *J Immunol* **175**, 6686–6693 (2005).
- 598 7. A. Tartakoff, P. Vassali, M. Detraz, Plasma cell immunoglobulin secretion. Arrest is
599 accompanied by alterations the golgi complex. *Journal of Experimental Medicine* **146**, 1332–
600 1345 (1977).
- 601 8. R. Jahn, R. H. Scheller, SNAREs--engines for membrane fusion. *Nat Rev Mol Cell Biol* **7**, 631–
602 643 (2006).
- 603 9. D. E. Gordon, L. M. Bond, D. A. Sahlender, A. A. Peden, A targeted siRNA screen to identify
604 SNAREs required for constitutive secretion in mammalian cells. *Traffic* **11**, 1191–1204 (2010).
- 605 10. A. Alloatti, *et al.*, Critical role for Sec22b-dependent antigen cross-presentation in antitumor
606 immunity. *J. Exp. Med.* **214**, 2231–2241 (2017).
- 607 11. K. Miyazaki, *et al.*, Contribution of the long form of syntaxin 5 to the organization of the
608 endoplasmic reticulum. *J Cell Sci* **125**, 5658–5666 (2012).
- 609 12. M. Petkovic, *et al.*, The SNARE Sec22b has a non-fusogenic function in plasma membrane
610 expansion. *Nat Cell Biol* **16**, 434–444 (2014).
- 611 13. G. A. Duque, *et al.*, Sec22b Regulates Inflammatory Responses by Controlling the Nuclear
612 Translocation of NF- κ B and the Secretion of Inflammatory Mediators. *The Journal of*
613 *Immunology* **207**, 2297–2309 (2021).
- 614 14. L. C. Gonzalez, W. I. Weis, R. H. Scheller, A novel snare N-terminal domain revealed by the
615 crystal structure of Sec22b. *J Biol Chem* **276**, 24203–24211 (2001).
- 616 15. J. C. Hay, D. S. Chao, C. S. Kuo, R. H. Scheller, Protein interactions regulating vesicle transport
617 between the endoplasmic reticulum and Golgi apparatus in mammalian cells. *Cell* **89**, 149–158
618 (1997).
- 619 16. A. Forrester, *et al.*, Functional dissection of the retrograde Shiga toxin trafficking inhibitor
620 Retro-2. *Nat Chem Biol* **16**, 327–336 (2020).

- 621 17. B. Stechmann, *et al.*, Inhibition of Retrograde Transport Protects Mice from Lethal Ricin
622 Challenge. *Cell* **141**, 231–242 (2010).
- 623 18. E. Hobeika, *et al.*, Testing gene function early in the B cell lineage in mb1-cre mice. *Proc. Natl.*
624 *Acad. Sci. U.S.A.* **103**, 13789–13794 (2006).
- 625 19. M. Giacomello, A. Pyakurel, C. Glytsou, L. Scorrano, The cell biology of mitochondrial
626 membrane dynamics. *Nat Rev Mol Cell Biol* **21**, 204–224 (2020).
- 627 20. T. Zhang, W. Hong, Ykt6 forms a SNARE complex with syntaxin 5, GS28, and Bet1 and
628 participates in a late stage in endoplasmic reticulum-Golgi transport. *J Biol Chem* **276**, 27480–
629 27487 (2001).
- 630 21. H. Hasegawa, *et al.*, Mammalian ykt6 is a neuronal SNARE targeted to a specialized
631 compartment by its profilin-like amino terminal domain. *Mol Biol Cell* **14**, 698–720 (2003).
- 632 22. D. E. Gordon, *et al.*, VAMP3/Syb and YKT6 are required for the fusion of constitutive secretory
633 carriers with the plasma membrane. *PLoS Genet* **13**, e1006698 (2017).
- 634 23. Y. Liu, C. Barlowe, Analysis of Sec22p in endoplasmic reticulum/Golgi transport reveals
635 cellular redundancy in SNARE protein function. *Mol Biol Cell* **13**, 3314–3324 (2002).
- 636 24. J. Jurkin, *et al.*, The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and
637 controls antibody secretion in plasma cells. *EMBO J* **33**, 2922–2936 (2014).
- 638 25. A. M. McGehee, *et al.*, XBP-1-Deficient Plasmablasts Show Normal Protein Folding but Altered
639 Glycosylation and Lipid Synthesis. *The Journal of Immunology* **183**, 3690–3699 (2009).
- 640 26. A. M. Reimold, *et al.*, Plasma cell differentiation requires the transcription factor XBP-1. *Nature*
641 **412**, 300–307 (2001).
- 642 27. N. Taubenheim, *et al.*, High rate of antibody secretion is not integral to plasma cell
643 differentiation as revealed by XBP-1 deficiency. *J. Immunol.* **189**, 3328–3338 (2012).
- 644 28. A. Bonaud, *et al.*, A mouse model recapitulating human monoclonal heavy chain deposition
645 disease evidences the relevance of proteasome inhibitor therapy. *Blood* **126**, 757–765 (2015).
- 646 29. S. Casola, *et al.*, B cell receptor signal strength determines B cell fate. *Nat. Immunol.* **5**, 317–327
647 (2004).
- 648 30. J. Tellier, *et al.*, Blimp-1 controls plasma cell function through the regulation of immunoglobulin
649 secretion and the unfolded protein response. *Nat Immunol* **17**, 323–330 (2016).
- 650 31. B. T. Gaudette, N. N. Iwakoshi, L. H. Boise, Bcl-xL protein protects from C/EBP homologous
651 protein (CHOP)-dependent apoptosis during plasma cell differentiation. *J Biol Chem* **289**,
652 23629–23640 (2014).
- 653 32. H. Zhu, *et al.*, Ufbp1 promotes plasma cell development and ER expansion by modulating
654 distinct branches of UPR. *Nat Commun* **10**, 1084 (2019).
- 655 33. J. W. Brewer, J. A. Diehl, PERK mediates cell-cycle exit during the mammalian unfolded
656 protein response. *Proc Natl Acad Sci U S A* **97**, 12625–12630 (2000).
- 657 34. S. Bustany, J. Cahu, P. Guardiola, B. Sola, Cyclin D1 sensitizes myeloma cells to endoplasmic
658 reticulum stress-mediated apoptosis by activating the unfolded protein response pathway. *BMC*
659 *Cancer* **15**, 262 (2015).

- 660 35. M. Lee, *et al.*, SNAREs support atlastin-mediated homotypic ER fusion in *Saccharomyces*
661 *cerevisiae*. *J Cell Biol* **210**, 451–470 (2015).
- 662 36. M. Lee, Y. Moon, S. Lee, C. Lee, Y. Jun, Ergosterol interacts with Sey1p to promote atlastin-
663 mediated endoplasmic reticulum membrane fusion in *Saccharomyces cerevisiae*. *FASEB J* **33**,
664 3590–3600 (2019).
- 665 37. X. Zhao, *et al.*, Sec22 regulates endoplasmic reticulum morphology but not autophagy and is
666 required for eye development in *Drosophila*. *J Biol Chem* **290**, 7943–7951 (2015).
- 667 38. E. Karampini, *et al.*, Sec22b determines Weibel-Palade body length by controlling anterograde
668 ER-Golgi transport. *Haematologica* **106**, 1138–1147 (2021).
- 669 39. A. Gallo, *et al.*, Role of the Sec22b-E-Syt complex in neurite growth and ramification. *J Cell Sci*
670 **133**, jcs247148 (2020).
- 671 40. G. Yoo, S. Yeou, J. B. Son, Y.-K. Shin, N. K. Lee, Cooperative inhibition of SNARE-mediated
672 vesicle fusion by α -synuclein monomers and oligomers. *Sci Rep* **11**, 10955 (2021).
- 673 41. C. H. Jung, *et al.*, Inhibition of SNARE-driven neuroexocytosis by plant extracts. *Biotechnol*
674 *Lett* **31**, 361–369 (2009).
- 675 42. R. Noel, *et al.*, N-methyldihydroquinazolinone derivatives of Retro-2 with enhanced efficacy
676 against Shiga toxin. *J Med Chem* **56**, 3404–3413 (2013).
- 677 43. J. R. Wiśniewski, A. Zougman, N. Nagaraj, M. Mann, Universal sample preparation method for
678 proteome analysis. *Nat Methods* **6**, 359–362 (2009).
- 679 44. J. Cox, M. Mann, MaxQuant enables high peptide identification rates, individualized p.p.b.-
680 range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**, 1367–1372
681 (2008).
- 682 45. N. Alouche, *et al.*, Hematologic disorder-associated Cxcr4 gain-of-function mutation leads to
683 uncontrolled extrafollicular immune response. *Blood* **137**, 3050–3063 (2021).
- 684 46. A. Bonaud, K. Balabanian, M. Espéli, Immunophenotyping of the Medullary B Cell
685 Compartment In Mouse Models. *Methods Mol Biol* **2308**, 95–105 (2021).
- 686 47. A. Bonaud, *et al.*, Leupaxin Expression Is Dispensable for B Cell Immune Responses. *Front.*
687 *Immunol.* **11**, 466 (2020).
- 688 48. G. L. Chen, E. W. Lamirande, H. Jin, G. Kemble, K. Subbarao, Safety, immunogenicity, and
689 efficacy of a cold-adapted A/Ann Arbor/6/60 (H2N2) vaccine in mice and ferrets. *Virology* **398**,
690 109–114 (2010).

691
692

693 **Figure Legends:**

694 **Figure 1: The Syntaxin-5-Sec22b SNARE complex is overexpressed in plasma cells:**

695 A) Changes in protein abundance between murine Naive B cells and CD138 enriched PCs
696 generated *in vitro* were measured using label free LC-MS/MS analysis and plotted using the
697 PlasmacytOMICs interface. The Log₂ Fold change between naïve B cells and the indicated
698 cell subset is shown for each protein. Error bars show SEM. * - indicates False Discovery
699 Adjusted p-value < 0.05. We were unable to consistently measure sufficient peptides for
700 quantification of CD138 in the PC samples possibly due to its high level of glycosylation. The
701 fold change shown for XBP1 is an underestimate, as the ratio plotted is calculated using an
702 imputed value for the B-cell samples as the protein was not detected in these samples (see
703 Supplementary Figure 1). B) Representative immunoblots for Xbp1, IgM, CD138, Scfd1,
704 Uso1, Stx5, Ytk6, Sec22b and b-actin (top to bottom respectively) from samples prepared
705 from splenic B cells (left panel) or CD138 enriched *in vitro* differentiated PCs (right panel).
706 The band shown for CD138 is the non-glycosylated form of the protein. For Stx5 the two
707 bands correspond to the short and the long isoforms of the protein. The fold change between
708 B cells and PCs normalized to b-actin is indicated on the right for each protein. C) The
709 PlasmacytOMICs interface was used to perform a meta-analysis of gene expression changes
710 between murine Naive B cells and a range of antibody secreting cell types (mSplPC.Array =
711 microarray / mouse splenic PCs; mBMPC.Array = microarray / mouse BM PCs; mPB.rSeq =
712 RNAseq / mouse plasmablasts generated *in vitro*; mSplPC.rSeq = RNAseq / mouse splenic
713 PCs; mBMPC.rSeq = RNAseq / mouse BM PCs) for the indicated genes. The Log₂ Fold
714 change between naïve B cells and the indicated cell subset is shown for each gene. Error bars
715 show SEM . * - indicates False Discovery Adjusted p-value < 0.05. B) D) Changes in gene
716 expression between human Naive B cells and selection of human antibody secreting cell types
717 (hPB.rSeq = RNAseq / human blood PC; hBMPC.rSeq = RNAseq / human BM PC) were

718 calculated for the indicated genes using the PlasmacytOMICcs platform. The Log₂ Fold
719 change between naïve B cells and the indicated cell subset is shown for each gene. Error bars
720 show SEM . * - indicates False Discovery Adjusted p-value < 0.05. Sec22b^{WT} controls were
721 mb1cre⁺ and Sec22b^{fl/fl} / Sec22b^{fl/+} mice.

722

723 **Figure 2: Strong reduction of plasma cells and circulating antibodies in absence of**
724 **Sec22b:**

725 A) ELISA quantification of Ig titres in sera of unimmunized Sec22b^{WT} or Sec22b^{B-KO} mice.
726 B-C) Representative dot plots (top) and quantification (bottom) of absolute number of total
727 PCs (CD138⁺TACI⁺) and percentage of each PC subsets: PBs (CD138⁺TACI⁺B220⁺CD19⁺),
728 early PCs (CD138⁺TACI⁺B220^{low}CD19⁺) and late PCs (CD138⁺TACI⁺B220⁻CD19⁻) in
729 Sec22b^{WT} and Sec22b^{B-KO} mice in spleen (B) and bone marrow (C). Cells were first gated on
730 their size, structure and viability. Dead cells and doublets were excluded. D) Unsupervised
731 clustering based on the relative expression of *Prdm1*, *Xbp1*, *Irf4*, *Tnfrsf13b*, *CD93*, *Pax5* and
732 *bach2* of sorted Sec22b^{WT} and Sec22b^{B-KO} PCs determined by Biomark multiplex qPCRs at
733 steady state. The heatmap was generated using the heatmapper.ca website and row Z score
734 based on (2^{-DC_t}) values. n= 7-19 mice from 2-5 independent experiments. The p-values were
735 determined with the two-tailed Mann-Whitney non-parametric test. **p* < 0.05; ***p* < 0.01;
736 ****p*<0.001 *****p*<0.0001. “ns” = non-significant *p*-value. Sec22b^{WT} controls were mb1cre⁺
737 and Sec22b^{fl/fl} / Sec22b^{fl/+} mice.

738

739 **Figure 3: Sec22b is essential for plasma cell maintenance and molecular identity**

740 A-B) Representative dot plots (A) and quantification (B) of PCs (CD138⁺B220^{+/-}) generated
741 *in vitro* from Sec22b^{WT} and Sec22b^{B-KO} splenocytes after 2 or 4 days of stimulation with LPS.
742 Cells were gated on their size and structure, on their viability and doublets were excluded. C)

743 Frequency of Caspase 3⁺ PCs at day 2 and 4 post LPS stimulation determined by flow
744 cytometry. D) Flow cytometry analysis of the cell cycle phases based on DAPI and Ki-67
745 staining in *in vitro* generated Sec22b^{WT} and Sec22b^{B-KO} PCs at day 2 (left) and day 4 (right)
746 post LPS stimulation. E-I) RNAseq analysis of Sec22b^{WT} and Sec22b^{B-KO} PCs generated *in*
747 *vitro* at day2 post LPS stimulation. E) Principal component analysis of the RNAseq data of
748 Sec22b^{WT} and Sec22b^{B-KO} PCs. The proportion of variance is indicated for PC1 and PC2. F)
749 Unsupervised clustering of Sec22b^{WT} and Sec22b^{B-KO} PCs based on the 500 most
750 differentially expressed genes. G) Volcano-plot showing the differentially expressed genes
751 between Sec22b^{WT} and Sec22b^{B-KO} PCs. Genes significantly downregulated and upregulated
752 in Sec22b^{B-KO} PCs are shown in green and red, respectively (FC>1.2; q value<0.05). H)
753 Normalized Enrichment Scores (NES) of representative gene sets significantly enriched in
754 Sec22b^{B-KO} vs Sec22b^{WT} PCs and characteristic of selected cellular pathways. Gene sets
755 significantly downregulated and upregulated in Sec22b^{B-KO} PCs are shown in green and red,
756 respectively. I) Expression of *Ykt6*, *Gosr1*, *Gosr2*, *Stx5* and *Bet1* in fragments per kilobase of
757 exon per million reads mapped (FPKM) determined by RNAseq in Sec22b^{WT} and Sec22b^{B-KO}
758 PCs. J) Quantification of the IgM secretion rate from *in vitro* generated PCs from Sec22b^{WT}
759 and Sec22b^{B-KO} splenocytes after 2 or 4 days of LPS stimulation analysed by DropMap. Each
760 point represents one cell.

761 (A-B) n=12-16 mice from at least 3 pooled independent experiments. (C-D) n=4 of 1
762 representative experiment out of 2. (E-I) n=4 in 1 experiment; (J) n=4 in 2 independent
763 experiments. The p-values were determined with the two-tailed Mann-Whitney non-
764 parametric test **p* < 0.05; ***p* < 0.01; ****p*<0.001, “ns” = non-significant *p*-value. Sec22b^{WT}
765 controls were mb1cre⁺ and Sec22b^{fl/fl} / Sec22b^{fl/+} mice.

766

767 **Figure 4: Sec22b is essential for ER expansion and structure in PCs**

768 A) Expression of *Ern1* (encoding Ire1a), *Atf4*, *Atf6*, *Ddit3* (encoding Chop) in fragments per
769 kilobase of exon per million reads mapped (FPKM) determined by RNAseq in *Sec22b*^{WT} and
770 *Sec22b*^{B-KO} PCs. B) Heatmap showing the relative expression of selected ER stress genes
771 from PCs generated from *Sec22b*^{WT} and *Sec22b*^{B-KO} splenocytes after 2 days of LPS
772 stimulation, determined by Biomark multiplex qPCRs at steady state. C) Flow cytometric
773 quantification of ER-Tracker MFI (geometrical mean) on PCs generated from *Sec22b*^{WT} and
774 *Sec22b*^{B-KO} splenocytes after 2 or 4 days of LPS stimulation. n=4, one representative
775 experiment shown out of 3. D) Confocal microscopy images of *Sec22b*^{WT} and *Sec22b*^{B-KO}
776 PCs obtained from splenocytes stimulated with LPS for 2 days. Cells were stained with anti-
777 IgM to detect antibodies, anti-calnexin to detect the ER and anti-IRF4 antibodies. Nuclei were
778 counterstained with Hoechst. Images are representative of 60 *Sec22b*^{WT} PC and 71 *Sec22b*^{B-}
779 ^{KO} PC from 2 mice per group. E) Electron microscopy images of *Sec22b*^{WT} (left) and
780 *Sec22b*^{B-KO} (right) PCs obtained from splenocytes stimulated with LPS for 2 (top) or 4
781 (bottom) days. Red arrowheads indicate ER sheets. Scale bar = 1mm or 0.5mm. Images are
782 representative of 64 WT PCs (9 at day 2 and 55 at day 4), and 39 KO PCs (21 at day 2 and 18
783 at day 4). Data are from 5 mice per genotype out of 2 experiments.

784 The heatmap were generated using the heatmapper.ca website, row Z score based on $(2^{-\Delta Ct})$
785 values. The p-values were determined with the two-tailed Mann-Whitney non-parametric test
786 $*p < 0.05$. *Sec22b*^{WT} controls were mb1cre⁺ and *Sec22b*^{fl/fl} / *Sec22b*^{fl/+} mice.

787

788 **Figure 5: Sec22b deficiency affects PC fitness via altered mitochondrial dynamics**

789 A) Expression of *Dnm1l*, *Inf2* and *Spire1* in FPKM determined by RNAseq in *Sec22b*^{WT} and
790 *Sec22b*^{B-KO} PCs obtained from splenocytes stimulated with LPS for 2 days. n=4 and data are
791 representative of 1 experiment. B) PLA performed on splenocytes stimulated 2 days with
792 LPS. Representative images are shown (left), PC stained with anti-IgM (green), PLA (red dot)

793 and nuclei counterstained with DAPI. Quantification of dot per PC (middle) (one
794 representative experiment) and mean of dot number per PC in each experiment (right) are
795 represented. Scale bar = 30 μ m. n= 1271 cells Sec22b^{WT} and 1893 cells Sec22b^{B-KO} of 3-4
796 mice in 3 independent experiment. C) Flow cytometric quantification of Mito-tracker (mitoT)
797 green (left) and mitoT orange CMTMRos (right) MFI (geometrical mean) on PCs generated
798 from Sec22^{WT} and Sec22b^{B-KO} splenocytes after 2 or 4 days of LPS stimulation. n= 4 mice in
799 1 representative experiment out of 3. D-E) Confocal microscopy images of PCs obtained from
800 Sec22b^{WT} and Sec22b^{B-KO} splenocytes stimulated with LPS for 4 days. Cells were stained
801 with anti-IgM to detect antibodies, anti-cytochrome c to detect mitochondria, and nuclei were
802 counterstained with Hoechst. Scale bar = 10 μ m. Representative images (D) and quantification
803 (E) of the mean area (top), mean intensity (mean per z-stack) (middle) and number of
804 fragments (over the full cell volume quantified through 10 z-stacks) (bottom) per cell of the
805 cytochrome c staining in PCs obtained from Sec22^{WT} and Sec22b^{B-KO} splenocytes stimulated
806 with LPS for 4 days. n=304 for the Sec22b^{WT} and n=231 for the Sec22b^{B-KO} from 3
807 independent mice per genotype. F-G) 3D projection of confocal microscopy images of PCs
808 obtained from Sec22^{WT} and Sec22b^{B-KO} splenocytes stimulated with LPS for 4 days and after
809 2 days of treatment with DMSO or M1/Mdivi for Sec22^{WT}. Cells were stained with anti-IgM,
810 anti-cytochrome c antibodies and nuclei were counterstained with Hoechst. Scale bar = 6mm.
811 Representative images (F) and quantification (G) of fold change of PC frequency normalized
812 to control Sec22b^{WT} PCs are shown. n=10 from 2 pooled experiments. H) Supervised
813 heatmap based on RNAseq expression data of 9 mitochondrial gene families differentially
814 expressed between Sec22b^{WT} and Sec22b^{B-KO} PCs. For flow cytometry experiment cells were
815 gated on their size and structure, on their viability and doublets were excluded. The p-values
816 were determined with the two-tailed unpaired Mann-Whitney non-parametric test * $p < 0.05$;
817 **** $p < 0.0001$, “ns” = non-significant p -value except for PLA experiment, determined with

818 Welch's t test or paired t-test. Sec22b^{WT} controls were mb1cre⁺ and Sec22b^{fl/fl} / Sec22b^{fl/+}
819 mice.

820

821 **Figure 6: Sec22b is indispensable for the generation of a protective humoral immune**
822 **response:**

823 A) Schematic representation of the SRBC immunisation protocol for Sec22b^{WT} and Sec22b^{B-}
824 ^{KO} mice. B) Representative dot plots (top), frequency (bottom left) and absolute number
825 (bottom right) of splenic PCs (CD138⁺B220^{+/-}) determined by flow cytometry 7 days after
826 SRBC immunization. C) ELISA quantification of IgM (top) and IgG1 (bottom) serum titres 7
827 days after SRBC immunized. A-C one representative experiment of 2 is shown. D) Schematic
828 representation of the NP-KLH immunization/ boost protocol. Sec22b^{WT} or Sec22b^{B-KO} mice
829 were immunized, rechallenged 28 days later and analysed 7 days later. Sera were collected at
830 days 7-, 14-, 21- and 7-days post boost. E) Frequency (left) and absolute number (right) of
831 splenic PCs (CD138⁺B220^{+/-}) (top) and bone marrow (BM) PCs (CD138⁺B220^{+/-}) (bottom)
832 determined by flow cytometry 7 days post boost with NP-KLH (n=9 mice from 2 pooled
833 independent experiments). F) ELISA quantification of NP-IgM (top) and NP-IgG1 (bottom)
834 serum titres at days 7-, 14-, 21- post primary immunization and 7-days post boost (n=5 mice.
835 One experiment representative of 2 is shown). G) Schematic representation of the Flu
836 infection protocol for Sec22b^{WT} and Sec22b^{B-KO} mice. Mice were analysed 14 days after
837 infection with influenza A virus. H) Frequency (left) and absolute number (right) of
838 mediastinal PCs (CD138⁺B220^{+/-}) determined by flow cytometry 14 days after infection with
839 influenza A virus. I) Quantification of HAI titres 14 days post flu infection. J) Weight change
840 of Sec22b^{WT} or Sec22b^{B-KO} mice over 12 days after flu infection. (H-I; n=5, one
841 representative experiment of 2 is shown, J; n=15 from 2 pooled experiments). For flow
842 cytometry experiment cells were first gated on their size and structure, then their viability.

843 Dead cells and doublets were excluded. The p-values were determined with the two-tailed
844 Mann-Whitney non-parametric test * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$ or
845 with the 2way ANOVA with Sidak correction for multiple comparisons (**¶¶** $p < 0.01$, **¶¶¶**
846 $p < 0.001$). Sec22b^{WT} controls were mb1cre⁺ and Sec22b^{fl/fl} / Sec22b^{fl/+} mice.

847

848

849

850