

Emergence of exopolysaccharides overproducers is linked to environmental spatial structure via redox state

Maxime Ardré, Maxime Deforet

▶ To cite this version:

Maxime Ardré, Maxime Deforet. Emergence of exopoly
saccharides overproducers is linked to environmental spatial structure via redox state. MSphere, in
Press, $10.1128/{\rm msphere.00123-23}$. hal-04065104

HAL Id: hal-04065104 https://hal.science/hal-04065104

Submitted on 11 Apr 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Emergence of exopolysaccharides overproducers is linked to environmental spatial structure via redox state

Maxime Ardré¹, Maxime Deforet^{2,*}

1. Laboratoire Biophysique et Evolution, CBI, ESPCI Paris, Université PSL, CNRS, 75005 Paris, France

2. Sorbonne Université, CNRS, Institut de Biologie Paris-Seine (IBPS), Laboratoire Jean Perrin (LJP), F-75005, Paris

* maxime.deforet@sorbonne-universite.fr

Abstract:

The formation of biofilm at the air-liquid interface of a still flask is related to the emergence of exopolysaccharides (EPS) overproducers. These mutants have the ability to remain near the surface, where oxygen is abundant. Yet, it is still unclear what role oxygen plays in cellular metabolism in this condition. A. Besse, M.-C. Groleau and E. Déziel (mSphere e00057-23, 2023 <u>https://doi.org/10.1128/msphere.00057-23</u>) explains that the redox state of cells is key in understanding the emergence of EPS overproducers. They found that the spatial distribution of oxidizing agent (not oxygen specifically) controls the advantage of remaining near the air-liquid interface, and hence the advantage that EPS-overproduction confers. All together this research paves the way for a deeper comprehension of the relationship between the environment's spatial structure and population dynamics.

A bacterial culture flask left on the bench for many days will result in the production of biofilm at the air-liquid interface (ALI). A study by Besse and colleagues reveals with clarity that the seemingly simple observation has profound implications.

The formation of bacterial biofilm at the ALI is frequently triggered by the colonization of the surface by a mutant derived from the ancestral population. This mutant evolved to overproduce a polymer, which provides a competitive advantage at the ALI. Several mutants of exopolysaccharides (EPS) overproduction have been identified across species (Spiers 2002, Kovack 2019, Besse 2022), as well as the metabolic pathways and mutations associated with them. They are mediated by diguanylate cyclases (DGCs), which regulate the secondary signaling molecule cyclic di-GMP (c-di-GMP) (Filloux 2016). Specifically, the mutant develops a mutation in the DGCs that regulate the amount of c-di-GMP, leading to an over-expression of c-di-GMP (Spiers 2002, Yan 2017) and subsequent polymer production. In a dynamic and spatially organized environment, these mutations are essential for survival.

The abundance of oxygen at the ALI is generally accepted as the reason for colonization of this environment (Rainey 1998, Kobayashi 2007). Fast use of oxygen by the bacterial population leads to an oxygen depletion in the bulk of the liquid: a significant gradient of oxygen rapidly develops in a still flask with virtually no oxygen 1 mm below the ALI (Koza 2010) (Figure A).

In this context, the ability for bacterial cells to remain at the liquid surface is critical. The insightful research undertaken by Besse et al. provides a clear description of the cellular metabolic state in relation to the oxygen availability.

The authors went beyond the notion that dissolved oxygen is the driving force behind biofilm development and demonstrated that, instead, any oxidizing agent could play a role in biofilm formation. While oxygen is the most common oxidizing agent, an alternate agent, such as nitrate, can be used to maintain a healthy metabolism. To prove their point, they measured the cells' metabolic activity in a flask by measuring their redox state through the NADH/NAD+ ratio. When nitrate is supplied to the bacterial population in place of oxygen for oxidation, much fewer EPS-overproducing mutants were found to invade the ancestral population (Figure B).

On the other hand, the authors ensured that a modified strain Δanr that is incapable of metabolizing nitrate (due to the shutdown of the denitrification pathway by ANR) resulted in the selection of EPS-overproducing mutants even when nitrate was supplied. Moreover, they calculated the quantitative threshold of the NADH/NAD+ ratio below which the redox state of the population promotes the invasion of EPS-overproducing mutants.

A feedback loop between how the oxidizing agent is spread and how the ALI biofilm is made

Their research emphasizes that the spatial heterogeneity of an oxidizing agent is responsible for the selection of overproducer mutants at the ALI. Further evidence is shown when cultures are treated with pyocyanin, a diffuse metabolite that can transport electrons from the cell to a distant oxidant. Addition of pyocyanin to the flask makes the oxidizing agent's activity available across a greater distance. Therefore, the oxidizing environment becomes more homogeneous (less spatially structured, Figure B), decreasing the advantage of remaining at the ALI. Mutants that produce excessive EPS are less likely to outperform the ancestral strain. Conversely, adding DNase reduced extracellular DNA, a known electron shuttle in biofilms, reinforced the spatial structure and promoted emergence of EPS-overproducing mutants.

The spatial structure of the environment determines the invasion of mutations responsible for the excessive synthesis of EPS. To understand the feedback loop between ecology and evolution, however, one must understand the mechanisms underlying the structuring of the environment. Recent research (Ardré 2019) indicates that biofilm formation at the ALI influences the spatial distribution of diffusible metabolites produced at the ALI, in part through hydrodynamic instability-induced mixing. For instance, this mixing from the top to the bottom of the flask may explain why oxidized resazurin (pink dye) is visible so deeply under the ALI in Fig. 2C, below the typical penetration length of oxygen from the ALI as observed by Koza (Koza 2010).

This research shows that mutants who overproduce EPS benefit from a spatially heterogeneous and more concentrated oxygen distribution at the ALI because their redox state is well balanced there. Yet, it is still unclear why these mutants make up only a fraction of 30% of the population within a few millimeters of the ALI (Figure 2C of the article). The mechanisms that allow the stable coexistence of mutants and ancestors at the ALI are still unknown. Addressing this complex topic will undoubtedly require a collaborative effort from experts in microbial physiology, evolutionary biology, cell mechanics, hydrodynamics, and statistical physics.

References:

Ardré M, Dufour D, Rainey PB. 2019. Causes and Biophysical Consequences of Cellulose Production by Pseudomonas fluorescens SBW25 at the Air-Liquid Interface. Journal of Bacteriology 201.

Besse A, Groleau M-C, Trottier M, Vincent AT, Déziel E. 2022. Pseudomonas aeruginosa Strains from Both Clinical and Environmental Origins Readily Adopt a Stable Small-Colony-Variant Phenotype Resulting from Single Mutations in c-di-GMP Pathways. Journal of Bacteriology 204.

Kobayashi K. 2007. Bacillus subtilis Pellicle Formation Proceeds through Genetically Defined Morphological Changes. Journal of Bacteriology 189:4920–4931.

Kovács ÁT, Dragoš A. 2019. Evolved Biofilm: Review on the Experimental Evolution Studies of Bacillus subtilis Pellicles. Journal of Molecular Biology 431:4749–4759.

Koza A, Moshynets O, Otten W, Spiers AJ. 2010. Environmental modification and niche construction: developing O2 gradients drive the evolution of the Wrinkly Spreader. The ISME Journal 5:665–673.

O'Toole G, Kaplan HB, Kolter R. 2000. Biofilm Formation as Microbial Development. Annual Review of Microbiology 54:49–79.

Rainey PB, Travisano M. 1998. Adaptive radiation in a heterogeneous environment - Nature. Nature. https://www.nature.com/articles/27900.

Spiers AJ, Kahn SG, Bohannon J, Travisano M, Rainey PB. 2002. Adaptive Divergence in Experimental Populations of Pseudomonas fluorescens. I. Genetic and Phenotypic Bases of Wrinkly Spreader Fitness. Genetics 161:33–46.

Valentini M, Filloux A. 2016. Biofilms and Cyclic di-GMP (c-di-GMP) Signaling: Lessons from Pseudomonas aeruginosa and Other Bacteria. Journal of Biological Chemistry 291:12547–12555.

Yan J, Deforet M, Boyle KE, Rahman R, Liang R, Okegbe C, P. Dietrich LE, Qiu W, Xavier JB. 2017. Bow-tie signaling in c-di-GMP: Machine learning in a simple biochemical network. Bow-tie signaling in c-di-GMP: Machine learning in a simple biochemical network | PLOS Computational Biology.

https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1005677.

Wong LS, Johnson MS, Zhulin IB, Taylor BL. 1995. Role of methylation in aerotaxis in Bacillus subtilis. Journal of Bacteriology 177:3985–3991.



A. Still flask

B. Stirred flask Still flask with e⁻ transporters Still flask supplemented with oxidants

Figure: A still flask induces a spatial structure of O_2 (the only oxidizing agent present in the media), which promotes the emergence of EPS-overproducing mutants and the formation of a biofilm at the air-liquid interface. When the flask is well stirred, or when the oxidizing agent

is dispersed in the media (nitrate or pyocyanin), the spatial structure is suppressed and no biofilm forms at the air-liquid interface.