



**HAL**  
open science

# Bilinear Strichartz estimates and almost sure global solutions for the nonlinear Schrödinger equation

Nicolas Burq, Aurélien Poiret, Laurent Thomann

► **To cite this version:**

Nicolas Burq, Aurélien Poiret, Laurent Thomann. Bilinear Strichartz estimates and almost sure global solutions for the nonlinear Schrödinger equation. 2023. hal-04066398

**HAL Id: hal-04066398**

**<https://hal.science/hal-04066398v1>**

Preprint submitted on 19 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

---

# BILINEAR STRICHARTZ ESTIMATES AND ALMOST SURE GLOBAL SOLUTIONS FOR THE NONLINEAR SCHRÖDINGER EQUATION

by

Nicolas Burq, Aurélien Poiret & Laurent Thomann

---

**Abstract.** — The purpose of this article is to construct global solutions, in a probabilistic sense, for the nonlinear Schrödinger equation posed on  $\mathbb{R}^d$ , in a supercritical regime. Firstly, we establish Bourgain type bilinear estimates for the harmonic oscillator which yields a gain of half a derivative in space for the local theory with randomised initial conditions, for the cubic equation in  $\mathbb{R}^3$ . Then, thanks to the lens transform, we are able to obtain global in time solutions for the nonlinear Schrödinger equation without harmonic potential. Secondly, we prove a Kato type smoothing estimate for the linear Schrödinger equation with harmonic potential. This allows us to consider the Schrödinger equation with a nonlinearity of odd degree in a supercritical regime, in any dimension  $d \geq 2$ .

**Résumé.** — L'objectif de cet article est de construire des solutions globales, en un sens probabiliste, pour l'équation de Schrödinger non linéaire posée sur  $\mathbb{R}^d$ , en régime surcritique. Tout d'abord, nous établissons des estimations bilinéaires de type Bourgain pour l'oscillateur harmonique. Celles-ci permettent un gain d'une demi-dérivée en l'espace pour la théorie locale avec des conditions initiales aléatoires, en ce qui concerne l'équation cubique dans  $\mathbb{R}^3$ . Puis, grâce à la transformée de lentille, nous sommes en mesure d'obtenir des solutions globales en temps pour l'équation de Schrödinger non linéaire sans potentiel harmonique. Dans un second temps, nous prouvons un effet régularisant de type Kato pour l'équation de Schrödinger linéaire avec potentiel harmonique. Ceci nous permet de considérer l'équation de Schrödinger avec une non-linéarité de degré impair dans un régime surcritique, dans toute dimension  $d \geq 2$ .

---

**2000 Mathematics Subject Classification.** — 35Q55 ; 35R60 ; 35P05.

**Key words and phrases.** — Bourgain type bilinear estimates, global solutions, harmonic oscillator, random data, super-critical non linear Schrödinger equations, smoothing effect.

N. Burq and L. Thomann were partially supported by the ANR project "SMOOTH" ANR-22-CE40-0017.

## Contents

1. Introduction and results.....	2
2. Preliminary results.....	10
3. The bilinear estimate for the harmonic oscillator.....	19
4. The smoothing effect for the harmonic oscillator.....	32
5. Random initial data and Sobolev spaces.....	38
6. The fixed point argument in dimension $d = 3$ , global existence, and scattering.....	45
7. Estimation of the regularity of the random initial data and proof of Theorem 1.2.....	56
8. The fixed point argument for equation (1.11).....	66
9. Estimation of the regularity of the initial random data and proof of Theorem 1.4.....	73
Appendix A. Tools of pseudo-differential operators and applications to eigenfunctions.....	77
References.....	83

## 1. Introduction and results

**1.1. Introduction.** — In this article, by the means of random initial conditions, we construct global solutions to the nonlinear Schrödinger equation in a supercritical regime. Namely, in the following we will consider the equation

$$(1.1) \quad \begin{cases} i \frac{\partial U}{\partial t} + \Delta U = \kappa |U|^{p-1} U, & (t, x) \in \mathbb{R} \times \mathbb{R}^d \\ U(0, x) = u_0(x), \end{cases}$$

where  $\kappa \in \{-1, 1\}$ , where  $p \geq 3$  is an odd integer, and  $d \geq 1$ .

The starting points of this article were two unpublished papers from the PhD thesis of the second author (in French), see [60, 61]. The first result of the present paper concerns the cubic Schrödinger equation in dimension  $d = 3$ . In this case the well-posedness is obtained using regularizing properties of random series combined with bilinear estimates for the harmonic oscillator. Then, in the second part of the article we prove global existence results for the nonlinear Schrödinger equation in dimension  $d \geq 2$ , with a nonlinearity of degree  $p \geq 5$ . This result is also obtained by taking benefit from stochastic properties of random series, but combined with the Kato smoothing effect which is established here for the Schrödinger equation with harmonic potential.

Since the pioneering works of Bourgain [5, 6] and the works of Tzvetkov [72, 73] on the Schrödinger equation, the papers of Burq-Tzvetkov [21, 22, 23] on the wave equation, there have been many contributions to the study of dispersive equations with random initial conditions. Among them, there are results on the Schrödinger equation by Colliander-Oh [32], Benyi-Oh-Pocovnicu [1, 3], Bourgain-Bulut [9, 10], Nahmod-Oh-Rey Bellet-Staffilani [54], Nahmod-Staffilani [55], Dodson-Lührman-Mendelson [42], Kilip-Murphy-Visan [50], Oh-Tzvetkov-Zhang [58], Deng-Nahmod-Yue [39, 40], on the wave equation by Oh-Pocovnicu-Tzvetkov [57], Burq-Lebeau [15], de Suzzoni [34, 35], Bourgain-Bulut [10], Bringmann [11, 12, 13], Sun-Tzvetkov [66], on the Benjamin-Ono equation by Tzvetkov-Visciglia [74, 75], Deng-Tzvetkov-Visciglia [36], Deng [38], and many others.

In some of the more recent works on the topic, some of the material from [60, 61] was applied, extended or generalised, see *e.g.* [62, 59]. Therefore it appeared relevant to rework the material from these papers and to gather them in a reader accessible version, which is the aim of the present article.

The strategy of the proof, already used in [69, 20, 37] and more recently in [19, 52], is the following. We first study a nonlinear Schrödinger equation with harmonic potential, for which we are able to prove local existence results. Then thanks to the lens transform, this latter equation is conjugated to the equation (1.1) for which we are able to deduce global existence results.

For  $d \geq 1$ , we define the harmonic oscillator by

$$H = -\Delta_d + |x|^2 = \sum_{j=1}^d \left( -\frac{\partial^2}{dx_j^2} + x_j^2 \right),$$

and we denote by  $\{h_n, n \geq 0\}$  an orthonormal basis of  $L^2(\mathbb{R}^d)$  of eigenvectors of  $H$  (the Hermite functions). The eigenvalues of  $H$  are the  $\{2(\ell_1 + \dots + \ell_d) + d, \ell \in \mathbb{N}^d\}$ , and we can order them in a non decreasing sequence  $\{\lambda_n^2, n \geq 0\}$ , repeated according to their multiplicities, and so that

$$Hh_n = \lambda_n^2 h_n, \quad \forall n \in \mathbb{N}.$$

It is then easy to observe that  $\lambda_n \sim n^{1/(2d)}$ , when  $n \rightarrow +\infty$ .

In the following,  $H^s(\mathbb{R}^d)$  and  $W^{s,p}(\mathbb{R}^d)$  denote the usual Sobolev spaces. We also define the harmonic Sobolev spaces, associated to the harmonic oscillator. The harmonic Sobolev space  $\mathcal{H}^s(\mathbb{R}^d)$  is defined as the closure of the Schwartz space for the norm

$$\|u\|_{\mathcal{H}^s(\mathbb{R}^d)} = \|H^{s/2}u\|_{L^2(\mathbb{R}^d)}.$$

Similarly, the space  $\mathcal{W}^{s,p}(\mathbb{R}^d)$  is defined as the closure of the Schwartz space for the norm

$$\|u\|_{\mathcal{W}^{s,p}(\mathbb{R}^d)} = \|H^{s/2}u\|_{L^p(\mathbb{R}^d)}.$$

In fact, this latter norm is a weighted Sobolev norm, because from [45] : for all  $1 < p < +\infty$  and  $s \geq 0$ , there exists a constant  $C > 0$  such that

$$(1.2) \quad \frac{1}{C} \|u\|_{\mathcal{W}^{s,p}(\mathbb{R}^d)} \leq \|(-\Delta)^{s/2}u\|_{L^p(\mathbb{R}^d)} + \|\langle x \rangle^s u\|_{L^p(\mathbb{R}^d)} \leq C \|u\|_{\mathcal{W}^{s,p}(\mathbb{R}^d)}.$$

Since the family  $\{h_n, n \geq 0\}$  forms a Hilbertian basis of  $L^2(\mathbb{R}^d)$ , any  $u \in \mathcal{H}^\sigma(\mathbb{R}^d)$  can be written

$$u = \sum_{n \in \mathbb{N}} c_n h_n \quad \text{where} \quad \|u\|_{\mathcal{H}^\sigma(\mathbb{R}^d)}^2 = \sum_{n \in \mathbb{N}} \lambda_n^{2\sigma} |c_n|^2 < +\infty.$$

Next, assume that  $(\Omega, \mathcal{T}, P)$  is a given probability space and that  $(g_n(\omega))_{n \in \mathbb{N}}$  is an identically distributed sequence of independent complex Gaussian random variables

$$g_n \sim \mathcal{N}_{\mathbb{C}}(0, 1),$$

namely, the density of  $g_n$  is given by  $\frac{1}{\pi} e^{-|z|^2} dL$  where  $dL$  denotes the Lebesgue measure on  $\mathbb{C}$ . Then, for  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^d)$ , which can be expanded as

$$u_0(x) = \sum_{n \in \mathbb{N}} c_n h_n(x),$$

we can consider the application  $\omega \mapsto u_0(\omega, \cdot)$ , from  $(\Omega, \mathcal{T})$  to  $\mathcal{H}^\sigma(\mathbb{R}^d)$  that we equip with its Borelian  $\sigma$ -algebra, defined by

$$u_0^\omega(x) = \sum_{n \in \mathbb{N}} c_n g_n(\omega) h_n(x).$$

We can check that the application  $\omega \mapsto u_0^\omega$  is in  $L^2(\Omega, \mathcal{H}^\sigma(\mathbb{R}^d))$  and we define the probability measure  $\mu$  as the distribution of this random variable. By definition, we have the following equality:

$$P(\omega \in \Omega : \Psi(u_0^\omega) \in A) = \mu(f \in \mathcal{H}^\sigma(\mathbb{R}^d) : \Psi(f) \in A),$$

for any measurable  $\Psi : (\mathcal{H}^\sigma(\mathbb{R}^d), \mathcal{B}(\mathcal{H}^\sigma(\mathbb{R}^d))) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$  and any set  $A \in \mathcal{B}(\mathbb{R})$ .

Let us first state a result which shows that the randomisation does not improve the initial condition in the Sobolev scale, and that it does not improve its spatial localisation.

**Theorem 1.1.** — *For all  $s \geq 0$ , if  $u_0 \notin \mathcal{H}^s(\mathbb{R}^d)$  then*

$$u_0^\omega \notin H^s(\mathbb{R}^d) \quad \text{and} \quad \langle x \rangle^s u_0^\omega \notin L^2(\mathbb{R}^d), \quad \omega - a.s.$$

By (1.2) it follows that one therefore has  $u_0^\omega \notin \mathcal{H}^s(\mathbb{R}^d)$ ,  $\omega - a.s.$

**1.2. Almost sure global existence for the cubic Schrödinger equation in dimension 3.** — Our first results concern the cubic equation

$$(1.3) \quad \begin{cases} i \frac{\partial U}{\partial t} + \Delta U = \kappa |U|^2 U, & (t, x) \in \mathbb{R} \times \mathbb{R}^3 \\ U(0, x) = u_0(x), \end{cases}$$

where  $\kappa \in \{-1, 1\}$ . Let us first recall some deterministic results concerning (1.3). There exists  $T > 0$  and a space  $X_T^s$  continuously embedded into  $C^0([-T, T]; H^s(\mathbb{R}^3))$  such that:

- If  $s > 1/2$  (subcritical case) then for all  $R > 0$ , there exists  $T_R > 0$  such that if  $\|u_0\|_{H^s(\mathbb{R}^3)} \leq R$ , then there exists a unique local solution  $u \in X_{T_R}^s$  to the equation (1.3), see [47, 26]. Moreover, the mapping  $u_0 \in B_{H^s}(0, R) \rightarrow u \in X_{T_R}^s$  is continuous. This means that the problem is locally well-posed. If  $T$  can be chosen equal to  $+\infty$ , we say that the problem is globally well-posed.

In the case where the problem is globally well-posed, it is natural to study the behavior of the solution in  $+\infty$ : if for all  $u_0 \in H^s(\mathbb{R}^3)$ , there exists  $u_+ \in H^s(\mathbb{R}^3)$  such that  $\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u_+\|_{H^s(\mathbb{R}^3)} = 0$ , which is known as scattering. According

to [31] it is the case, for  $\kappa = 1$ , as soon as  $s > \frac{4}{5}$ .

- If  $s = 1/2$  (critical case), one can prove the local existence of a unique solution for each initial data as in the subcritical case, but the existence time of the solution depends on  $u_0$  and not only on  $\|u_0\|_{H^s(\mathbb{R}^3)}$ . Therefore the globalization problem is a complex problem. See [26].

- If  $s < 1/2$  (supercritical case), according to [27], we know that there exists a sequence of real numbers  $t_n \in \mathbb{R}$  which tends to 0 and a sequence of functions  $u_n \in H^s(\mathbb{R}^3)$  such that  $\lim_{n \rightarrow +\infty} \|u_n\|_{H^s(\mathbb{R}^3)} = 0$  and  $\lim_{n \rightarrow +\infty} \|u_n(t_n)\|_{H^s(\mathbb{R}^3)} = +\infty$ , where  $u_n(t)$  denotes a solution of the equation (1.3) with initial data  $u_n$ . Therefore, the flow of the equation (1.3) is not continuous in 0, which implies that the problem is not well posed and the usual methods do not allow to study the equation in this situation.

We assume in this section that the eigenfunctions of the harmonic oscillator are given by the tensor eigenfunctions, *i.e.* for  $n \geq 0$ , there exist  $(n_1, n_2, n_3) \in \mathbb{N}^3$  such that for  $x = (x_1, x_2, x_3) \in \mathbb{R}^3$

$$(1.4) \quad h_n(x) = e_{n_1}(x_1)e_{n_2}(x_2)e_{n_3}(x_3),$$

where  $(-\partial_{x_j}^2 + x_j^2)e_{n_k} = (2n_k + 1)e_{n_k}$  and  $\lambda_n^2 = 2(n_1 + n_2 + n_3) + 3$ . Here  $(e_n)_{n \in \mathbb{N}}$  denotes the basis of the eigenfunctions of the harmonic oscillator in dimension 1 and  $(\mu_n^2)_{n \in \mathbb{N}}$  the associated eigenfunction sequence, so that we have  $\mu_n^2 = 2n + 1$ .

Indeed,  $e_n$  are given by the formula

$$e_n(x) = (-1)^n c_n e^{x^2/2} \frac{d^n}{dx^n} (e^{-x^2}), \quad \text{with} \quad \frac{1}{c_n} = (n!)^{\frac{1}{2}} 2^{\frac{n}{2}} \pi^{\frac{1}{4}}.$$

Our main result then reads :

**Theorem 1.2.** — *Let  $\sigma \in ]0, \frac{1}{2}[$  with  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$  and  $s \in ]\frac{1}{2}, \frac{1}{2} + \sigma[$ , then there exists a set  $\Omega' \subset \Omega$  satisfying the following conditions:*

- (i)  $P(\Omega') > 0$ .
- (ii) *For any element  $\omega \in \Omega'$ , there exists a unique global solution  $U$  to the equation (1.3) with initial data  $u_0^\omega$  such that*

$$U(t) - e^{it\Delta} u_0^\omega \in \mathcal{C}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^3)).$$

Moreover, for any element  $\omega \in \Omega'$ , there exist  $L_+, L_- \in \mathcal{H}^s(\mathbb{R}^3)$  such that

$$(1.5) \quad \begin{aligned} \lim_{t \rightarrow +\infty} \|U(t) - e^{it\Delta} (u_0^\omega + L_+)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \\ \lim_{t \rightarrow -\infty} \|U(t) - e^{it\Delta} (u_0^\omega + L_-)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \end{aligned}$$

and

$$(1.6) \quad \begin{aligned} \lim_{t \rightarrow +\infty} \|e^{-it\Delta} U(t) - (u_0^\omega + L_+)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \\ \lim_{t \rightarrow -\infty} \|e^{-it\Delta} U(t) - (u_0^\omega + L_-)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0. \end{aligned}$$

Finally, when  $\eta \rightarrow 0$  we have

$$(1.7) \quad \mu(u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) : \text{we have global existence and scattering} \mid \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \eta) \rightarrow 1.$$

Notice that since  $e^{\pm it\Delta}$  does not act on  $\mathcal{H}^\sigma(\mathbb{R}^3)$ , the properties (1.5) and (1.6) are different. It is interesting to notice that Theorem 1.2 is not a small data result in the critical  $H^{1/2}(\mathbb{R}^3)$  space. More precisely, for  $0 < \sigma < 1/2$ , let  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) \setminus H^{1/2}(\mathbb{R}^3)$ . Then for  $\omega \in \Omega'$ ,  $u_0^\omega$  satisfies the conclusion of Theorem 1.2 and by Theorem 1.1 we have  $u_0^\omega \notin H^{1/2}(\mathbb{R}^3)$ . Similarly, for  $K \geq 1$  we can define  $[u_0]_K = \sum_{\lambda_n < K} c_n h_n \in \bigcap_{s \geq 0} \mathcal{H}^s(\mathbb{R}^3)$ . Then Theorem 1.2 applies to this initial condition and we have  $\|[u_0^\omega]_K\|_{H^{1/2}(\mathbb{R}^3)} \gg 1$  for  $K \gg 1$  (we refer to Proposition 5.4 for a precise statement). This is the counterpart with some global existence results obtained for the Navier-Stokes equation, see [28, 29, 30].

We can obtain a more quantitative statement in the case of small initial conditions  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$ . For  $0 < \alpha < 1$ , there exists a set  $\Omega_\alpha \subset \Omega'$  satisfying  $P(\Omega_\alpha) \geq 1 - \alpha$ ,

under the condition

$$(1.8) \quad \alpha = C_1 \exp\left(-\frac{C_2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}\right),$$

for some universal constants  $C_1, C_2 > 0$ .

It is likely that our approach can be extended to any dimension  $d \geq 2$ . In this case we also gain a half-derivative compared to the deterministic problem. We will not give more details.

Our result can be extended to more general i.i.d. random variables  $(g_n)_{n \in \mathbb{N}}$ . Actually, we need the  $g_n$  to be centered, that they admit moment at any order and satisfy multilinear chaos estimates as in Proposition 7.2.

To establish our results, we use the ideas of N. Burq and N. Tzvetkov developed in [21, 22] by randomizing the initial data. This allows us to gain derivatives in  $L^p$  spaces, for  $p > 2$ .

The proof essentially relies on two intermediate results:

- (i) The lens transformation (introduced in [56, 25] and used in [20, 68]) which allows us to reduce to proving a local existence theorem on  $] -\frac{\pi}{4}; \frac{\pi}{4}[$  for the Schrödinger equation with harmonic potential (see Section 2.1).
- (ii) The existence of a bilinear estimate for the harmonic oscillator (see Section 3) which allows to gain the half-derivative on the first order terms in  $u_0$ . This estimate is the analogue of the bilinear estimate for the usual Laplacian proved by Bourgain in [7].

The results of Theorem 1.2 extend the results of [20, 37] obtained in space dimension  $d = 1$  and  $d = 2$ . In [62] similar results were obtained in space dimension  $d = 3$ , but the randomisation of the initial condition was different, which implied scattering results in stronger harmonic Sobolev norms. In the papers [11, 12, 2, 24] some scattering results are obtained using a randomisation based on the Littlewood-Paley decomposition of the initial condition.

The existence of a bilinear estimate for the harmonic oscillator being of interest in itself, we state it below in any dimension  $d \geq 2$ . Let us first define the dyadic localisation operators. Let  $\eta \in C_0^\infty(\mathbb{R})$  such that  $\eta_{[0,1]} = 1$  and  $\eta_{[2,+\infty[} = 0$ . Set  $\psi(x) = \eta(x) - \eta(4x)$ . For dyadic numbers  $N = 2^j$ , we define the following sequence of operators:

$$(1.9) \quad \Delta_N(u) = \begin{cases} \psi\left(\frac{H}{N^2}\right)u & \text{for } N \geq 1, \\ 0 & \text{else.} \end{cases}$$



Notice that if  $\lambda_n \notin [\frac{N}{2}, \sqrt{2}N]$  then  $\Delta_N(h_n) = 0$  and that  $\sum_{N \text{ dyadic}} \Delta_N(u) = u$ .

**Theorem 1.3.** — *Let  $d \geq 2$ . Then for any  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u, v \in \mathcal{S}'(\mathbb{R}^d)$ ,*

$$(1.10) \quad \begin{aligned} & \|e^{itH} \Delta_N(v) e^{itH} \Delta_M(u)\|_{L^2([-1,1]; L^2(\mathbb{R}^d))} \leq \\ & \leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|\Delta_N(v)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

The first bilinear estimate has been obtained by Bourgain [7] for the Schrödinger group on  $\mathbb{R}^d$ . For bilinear estimates on compact manifolds and application to the well-posedness of nonlinear Schrödinger equation, we refer to [17].

We refer to the work [59] of F. Planchon, N. Tzvetkov, and N. Visiglia for an alternative proof of Theorem 1.3 in the case  $d = 2$ , with the improvement  $\delta = 0$ . This bilinear estimate has been used in [59] in order to obtain bounds on the growth of norms for the nonlinear Schrödinger equation with harmonic potential.

**1.3. Almost sure global existence for the nonlinear Schrödinger equation in any dimension  $d \geq 2$ .** — We now consider the following Schrödinger equation with polynomial nonlinearity

$$(1.11) \quad \begin{cases} i \frac{\partial U}{\partial t} + \Delta U = \kappa |U|^{p-1} U, & (t, x) \in \mathbb{R} \times \mathbb{R}^d \\ U(0, x) = u_0(x), \end{cases}$$

where  $\kappa \in \{-1, 1\}$  and  $p \geq 5$  denotes an odd integer, and  $d \geq 2$ .

We prove the following results:

**Theorem 1.4.** — *Let  $u_0 \in \mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)$  then there exist  $s \in ]\frac{d}{2} - \frac{2}{p-1}, \frac{d}{2}[$  and a set of  $\Omega' \subset \Omega$  such that the following conditions are met:*

- (i)  $P(\Omega') > 0$ .
- (ii) For all element  $\omega \in \Omega'$ , there is a unique global solution  $U$  to the equation (1.11) in the space  $e^{it\Delta} u_0^\omega + X^s$  with initial data  $u_0^\omega$ .
- (iii) For all element  $\omega \in \Omega'$ , there exists  $L_+, L_- \in \mathcal{H}^s(\mathbb{R}^d)$  such that

$$(1.12) \quad \begin{aligned} & \lim_{t \rightarrow +\infty} \|U(t) - e^{it\Delta} (u_0^\omega + L_+)\|_{H^s(\mathbb{R}^d)} = 0, \\ & \lim_{t \rightarrow -\infty} \|U(t) - e^{it\Delta} (u_0^\omega + L_-)\|_{H^s(\mathbb{R}^d)} = 0, \end{aligned}$$

and

$$(1.13) \quad \begin{aligned} \lim_{t \rightarrow +\infty} \|e^{-it\Delta}U(t) - (u_0^\omega + L_+)\|_{\mathcal{H}^s(\mathbb{R}^d)} &= 0, \\ \lim_{t \rightarrow -\infty} \|e^{-it\Delta}U(t) - (u_0^\omega + L_-)\|_{\mathcal{H}^s(\mathbb{R}^d)} &= 0. \end{aligned}$$

Finally, when  $\eta \rightarrow 0$  we have

$$(1.14) \quad \mu(u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) : \text{we have global existence and scattering} \mid \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \eta) \rightarrow 1.$$

The proof of this result relies on the Kato smoothing effect for the linear Schrödinger equation with harmonic potential. In dimension 1, the smoothing effect was used in [20] in order to obtain local existence results. We observe here that this approach can be extended in high dimension, in the case of a nonlinearity of degree  $p \geq 5$  and an initial condition  $u_0 \in \mathcal{H}^{(d-1)/2}(\mathbb{R}^d)$ .

We refer to [42, 50] for almost sure scattering results for NLS.

Let us state the smoothing effect for the harmonic oscillator, which has his own interest.

**Theorem 1.5.** — *Let  $\epsilon \in ]0, \frac{1}{2}[$  then there exists a constant  $C > 0$  such that for all  $u_0 \in L^2(\mathbb{R}^d)$ ,*

$$(1.15) \quad \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{H}^{1/2-2\epsilon} e^{itH} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq C \|u_0\|_{L^2(\mathbb{R}^d)},$$

and for all  $u_0 \in \mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)$ ,

$$(1.16) \quad \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{-\Delta}^{d/2-2\epsilon} e^{itH} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq C \|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)}.$$

The first results on the local smoothing effect were obtained in [49, 64, 76, 33]. Here the proof follows the Doi method [43, 44]. See also [77, 78, 70] for results on the smoothing effect for the Schrödinger equation with confining potentials.

The result of Theorem 1.5 is an extension to the well-known smoothing effect for the linear Schrödinger group

$$\left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{-\Delta}^{1/2-2\epsilon} e^{it\Delta} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq C \|u_0\|_{L^2(\mathbb{R}^d)},$$

and for all  $u_0 \in H^{\frac{d-1}{2}}(\mathbb{R}^d)$ ,

$$\left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{-\Delta}^{d/2-2\epsilon} e^{it\Delta} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq C \|u_0\|_{H^{\frac{d-1}{2}}(\mathbb{R}^d)}.$$

**1.4. Notations.** — In this paper  $c, C > 0$  denote constants the value of which may change from line to line. These constants will always be universal, or uniformly bounded with respect to the other parameters. We denote by  $H = -\Delta + |x|^2$  the harmonic oscillator on  $\mathbb{R}^d$ , and for  $\sigma \in \mathbb{R}$  we define the Sobolev space  $\mathcal{H}^\sigma(\mathbb{R}^d)$  by the norm  $\|u\|_{\mathcal{H}^\sigma(\mathbb{R}^d)} = \|H^{\sigma/2}u\|_{L^2(\mathbb{R}^d)}$ . More generally, we define the spaces  $\mathcal{W}^{\sigma,p}(\mathbb{R}^d)$  by the norm  $\|u\|_{\mathcal{W}^{\sigma,p}(\mathbb{R}^d)} = \|H^{\sigma/2}u\|_{L^p(\mathbb{R}^d)}$ . The Fourier transform is defined by  $\mathcal{F}f(\xi) = \int_{\mathbb{R}^d} e^{-ix\xi} f(x) dx$ , for  $f \in \mathcal{S}(\mathbb{R}^d)$ . We denote by  $\langle x \rangle = \sqrt{1 + |x|^2}$ .

**1.5. Organization of the rest of the article.** — In Section 2, we present some harmonic analysis results, including study of the lens transform, Strichartz estimates for the Schrödinger group with harmonic potential, properties of the Bourgain space, and Hermite functions estimates. Section 3 is devoted to the proof of the bilinear estimates for the harmonic oscillator. Then, in Section 4, we prove the local smoothing effect. In Section 5 we prove that the randomized initial data does not allow to gain derivatives in the  $L^2$  scale and that it does not enjoy better spatial localisation properties (Theorem 1.1). Section 6 is devoted to the fixed point argument for the cubic NLS in dimension 3. In Section 7 we study the regularity of the randomised initial conditions which allows to complete the proof of Theorem 1.2. In Section 8 and in Section 9, using a fixed point argument, we prove the results on the NLS in dimension  $d \geq 2$ .

## 2. Preliminary results

In this section, except in the fourth part, the dimension of space is assumed to be any  $d \geq 1$ .

**2.1. The lens transformation.** — As in [69, 20], we use the lens transform which allows to work with the Schrödinger equation with harmonic potential. More precisely, suppose that  $U(s, y)$  is a solution of the problem (1.1). Then the function  $u(t, x)$  defined for  $|t| < \frac{\pi}{4}$  and  $x \in \mathbb{R}$  by

$$(2.1) \quad u(t, x) = \mathcal{L}(U)(t, x) := \left(\frac{1}{\cos 2t}\right)^{d/2} U\left(\frac{\tan(2t)}{2}, \frac{x}{\cos(2t)}\right) e^{-\frac{i|x|^2 \tan(2t)}{2}} \\ := \mathcal{L}_t(U \big|_{s=\frac{\tan(2t)}{2}})(x)$$

where

$$(2.2) \quad \mathcal{L}_t(G)(x) = \left(\frac{1}{\cos(2t)}\right)^{d/2} G\left(\frac{x}{\cos(2t)}\right) e^{-\frac{i|x|^2 \tan(2t)}{2}},$$

solves the problem

$$(2.3) \quad \begin{cases} i\partial_t u - Hu = \kappa \cos^{\frac{d}{2}(p-1)-2}(2t)|u|^{p-1}u, & |t| < \frac{\pi}{4}, x \in \mathbb{R}^d, \\ u(0, \cdot) = U_0, \end{cases}$$

where  $H = -\partial_x^2 + |x|^2$ . Similarly, if  $U = e^{is\Delta_y}U_0$  is a solution of the linear Schrödinger equation, then

$$(2.4) \quad u = e^{-itH}U_0 = \mathcal{L}(U)$$

is the solution of the linear harmonic Schrödinger equation with the same initial data. In other words, if we denote by  $\Psi(s, s')$  the map which sends the data at time  $s'$  to the solution at time  $s$  of (2.3), and by  $\Phi(t, t')$  the map which sends the data at time  $t'$  to the solution at time  $t$  of (1.11), the family  $(\mathcal{L}_t)_{|t| < \frac{\pi}{4}}$  conjugates the linear and the nonlinear flows: with  $t(s) = \frac{\arctan(2s)}{2}$ ,  $s(t) = \frac{\tan(2t)}{2}$ ,

$$\mathcal{L}_{t(s)} \circ e^{i(s-s')\Delta_y} = e^{-i(t(s)-t(s'))H} \circ \mathcal{L}_{t(s')}.$$

and

$$\mathcal{L}_{t(s)} \circ \Psi(s, s') = \Phi(t(s), t(s')) \circ \mathcal{L}_{t(s')}.$$

By (2.4) we have

$$(2.5) \quad e^{-itH}u(t, x) = \left(\frac{1}{\cos(2t)}\right)^{d/2} e^{it\Delta}u\left(\frac{\tan(2t)}{2}, \frac{x}{\cos(2t)}\right) e^{-\frac{i|x|^2 \tan(2t)}{2}}$$

and equivalently

$$(2.6) \quad e^{it\Delta}u(t, x) = \left(\frac{1}{\sqrt{1+4t^2}}\right)^{d/2} e^{-itH}u\left(\frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}}\right) e^{\frac{i|x|^2 t}{1+4t^2}}.$$

**2.2. The Strichartz estimates for the harmonic oscillator.** — In the following, we will say that a pair  $(q, r) \in [2, +\infty]^2$  is admissible if and only if

$$(q, r, d) \neq (2, +\infty, 2) \quad \text{and} \quad \frac{2}{q} = \frac{d}{2} - \frac{d}{r}.$$

We define, for  $s \in \mathbb{R}$  and  $T > 0$  the Strichartz spaces for the Schrödinger equation

$$\begin{aligned} X^s &= \bigcap_{(q,r) \text{ admissible}} L^q(\mathbb{R}; W^{s,r}(\mathbb{R}^d)), \\ X_T^s &= \bigcap_{(q,r) \text{ admissible}} L^q([-T, T]; W^{s,r}(\mathbb{R}^d)). \end{aligned}$$

Similarly, we define the Strichartz spaces for the harmonic oscillator

$$\begin{aligned}\bar{X}^s &= \bigcap_{(q,r) \text{ admissible}} L^q\left(\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]; \mathcal{W}^{s,r}(\mathbb{R}^d)\right), \\ \bar{X}_T^s &= \bigcap_{(q,r) \text{ admissible}} L^q([-T, T]; \mathcal{W}^{s,r}(\mathbb{R}^d)).\end{aligned}$$

The Strichartz estimates for the usual Schrödinger equation can be found in [67]. Let us state the Strichartz inequalities for the harmonic oscillator.

**Proposition 2.1.** — *For any time  $T \geq 0$ , there exists a constant  $C_T > 0$  such that for any function  $u \in \mathcal{H}^s(\mathbb{R}^d)$  and any function  $F \in L^{q'}([-T, T]; \mathcal{W}^{s,r'}(\mathbb{R}^d))$  with  $(q, r)$  admissible*

$$(2.7) \quad \|e^{-itH}u\|_{\bar{X}_T^s} \leq C_T \|u\|_{\mathcal{H}^s(\mathbb{R}^d)},$$

$$(2.8) \quad \left| \int_0^t e^{-i(t-s)H} F(s) ds \right|_{\bar{X}_T^s} \leq C \|F\|_{L^{q'}([-T, T]; \mathcal{W}^{s,r'}(\mathbb{R}^d))}.$$

*Proof.* — The Strichartz estimates follow from the study of the kernel of  $e^{itH}$ , and we refer to [20, Section 5] for a proof. Alternatively, let us show here how (2.7) also follows from the Strichartz estimates for the Schrödinger flow with the free Laplacian. By replacing  $u$  by  $e^{iT H}u$ , we see that it is enough to prove the estimate for some  $T > 0$ , for example for  $T = \frac{\pi}{4}$ . Similarly, by replacing  $u$  by  $H^{\frac{s}{2}}u$ , we can restrain the proof to the case where  $s = 0$ . Thus, if the couple  $(q, r)$  is admissible, by (2.5) we obtain

$$\begin{aligned}\|e^{-itH}u\|_{L^q\left(\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]; L^r(\mathbb{R}^d)\right)} &= \\ &= \left\| \left(\frac{1}{\cos(2t)}\right)^{d/2} e^{it\Delta} u \left(\frac{\tan(2t)}{2}, \frac{x}{\cos(2t)}\right) e^{-\frac{i|x|^2 \tan(2t)}{2}} \right\|_{L^q\left(\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]; L^r(\mathbb{R}^d)\right)} \\ &= \|e^{it\Delta}u\|_{L^q(\mathbb{R}; L^r(\mathbb{R}^d))}.\end{aligned}$$

Then, we can use the Strichartz estimates for the Laplacian to conclude.

The proof of (2.8), which relies on a duality argument, is the same as the proof of the Strichartz estimates for the Laplacian that can be found in [67, Section 2.3].  $\square$

**Proposition 2.2.** — *If  $s \geq 0$  then there exists a constant  $C > 0$  such that for all  $p \in [1, +\infty]$  and  $q \in [1, +\infty]$  satisfying  $\frac{2}{p} + \frac{d}{q} - \frac{d}{2} \leq 0$ , we have for any  $u \in L^p\left(\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]; \mathcal{W}^{s,q}(\mathbb{R}^d)\right)$ ,*

$$\|U\|_{L^p(\mathbb{R}; \mathcal{W}^{s,q}(\mathbb{R}^d))} \leq C \|u\|_{L^p\left(\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]; \mathcal{W}^{s,q}(\mathbb{R}^d)\right)}.$$

As a consequence, let  $s \geq 0$  and  $u \in \overline{X}^s$  then  $U \in X^s$  and there exists a constant  $C > 0$  such that for all  $u \in \overline{X}^s$ ,

$$\|U\|_{X^s} \leq C\|u\|_{\overline{X}^s}.$$

*Proof.* — By interpolation, it is sufficient to prove the result for  $s = n \in \mathbb{N}$ . Let  $\alpha \in \mathbb{N}^d$  with  $|\alpha| \leq n$ , then thanks to the Leibniz formula, we obtain

$$\begin{aligned} \partial_x^\alpha U(t, x) &= \partial_x^\alpha \left( \left( \frac{1}{\sqrt{1+4t^2}} \right)^{d/2} u \left( \frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) e^{\frac{i|x|^2 t}{1+4t^2}} \right) \\ &= \left( \frac{1}{\sqrt{1+4t^2}} \right)^{d/2} \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} \partial_x^\beta \left( u \left( \frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) \right) \partial_x^{\alpha-\beta} \left( e^{\frac{i|x|^2 t}{1+4t^2}} \right). \end{aligned}$$

Then, as

$$|\partial_x^{\alpha-\beta} (e^{\frac{i|x|^2 t}{1+4t^2}})| \leq C_{\alpha,\beta} \left( 1 + \left| \frac{x}{\sqrt{1+4t^2}} \right|^{|\alpha-\beta|} \right),$$

we establish

$$\begin{aligned} |\partial_x^\alpha U(t, x)| &\leq \\ \sum_{0 \leq \beta \leq \alpha} C_{\alpha,\beta} \left( \frac{1}{\sqrt{1+4t^2}} \right)^{d/2+|\beta|} |\partial_x^\beta u| \left( \frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) \left( 1 + \left| \frac{x}{\sqrt{1+4t^2}} \right|^{|\alpha-\beta|} \right). \end{aligned}$$

Therefore,

$$\begin{aligned} \|\partial_x^\alpha U\|_{L^p(\mathbb{R}; L^q(\mathbb{R}^d))} &\leq \\ &\leq \sum_{0 \leq \beta \leq \alpha} C_{\alpha,\beta} \left\| \left( \frac{1}{\sqrt{1+4t^2}} \right)^{d/2+d/q+|\beta|} \left\| u \left( \frac{\arctan(2t)}{2}, x \right) \right\|_{\mathcal{W}^{|\alpha|,q}(\mathbb{R}^d)} \right\|_{L^p(\mathbb{R})} \\ &\leq \sum_{0 \leq \beta \leq \alpha} C_{\alpha,\beta} \left\| \left( \frac{1}{\sqrt{1+\tan^2 2t}} \right)^{d/2+d/q-2/p+|\beta|} u(t, x) \right\|_{L^p([-\frac{\pi}{4}, \frac{\pi}{4}]; \mathcal{W}^{|\alpha|,q}(\mathbb{R}^d))}. \end{aligned}$$

To conclude, it is enough to notice that  $\frac{d}{2} + \frac{d}{q} - \frac{2}{p} + |\beta| \geq 0$  for  $\beta \in \mathbb{N}^d$ . □

**2.3. Some properties of Bourgain spaces.** — We now define the Bourgain spaces and recall some of their different properties.

We define the space  $\overline{X}^{s,b} = \overline{X}^{s,b}(\mathbb{R} \times \mathbb{R}^d)$  as the closure of  $C_0^\infty(\mathbb{R} \times \mathbb{R}^d)$  for the norm

$$\begin{aligned} \|u\|_{\overline{X}^{s,b}}^2 &= \sum_{n \in \mathbb{N}} \|\langle t + \lambda_n^2 \rangle^b \lambda_n^s \widehat{P_n u}(t)\|_{L_t^2(\mathbb{R}; L_x^2(\mathbb{R}^d))}^2 \\ &= \sum_{n \in \mathbb{N}} \|H^{s/2} e^{itH} P_n u(t, \cdot)\|_{L_x^2(\mathbb{R}^d; H_t^b(\mathbb{R}))}^2, \end{aligned}$$

where  $\widehat{P_n u}(t)$  denotes the Fourier transform, with respect to the time variable, of  $P_n u := \langle u, h_n \rangle_{L^2(\mathbb{R}^d) \times L^2(\mathbb{R}^d)} h_n$ .

In [67, Corollary 2.10], we can find the following proposition:

**Proposition 2.3.** — *For any real  $b > \frac{1}{2}$ , there exists a constant  $C > 0$  such that for any real  $s \in \mathbb{R}$  and for any admissible pair  $(q, r)$ , if  $u \in \overline{X}^{s,b}$  then  $u \in L^q(\mathbb{R}; \mathcal{W}^{s,r}(\mathbb{R}^d))$  and*

$$\|u\|_{L^q(\mathbb{R}; \mathcal{W}^{s,r}(\mathbb{R}^d))} \leq C \|u\|_{\overline{X}^{s,b}}.$$

Thanks to [17, Lemma 2.4], we can establish the following proposition:

**Proposition 2.4.** — *For any  $\theta \in [0, 1]$ , if  $b > \frac{1-\theta}{2}$  then there exists a constant  $C > 0$  such that for any  $s \in \mathbb{R}$  and any function  $u \in \overline{X}^{s,b}$ ,*

$$\|u\|_{L^{\frac{2}{\theta}}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^d))} \leq C \|u\|_{\overline{X}^{s,b}}.$$

*Proof.* — It is enough to consider the case  $s = 0$ . Using the inverse Fourier transform, we have

$$P_n u(t) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{it\tau} \widehat{P_n u}(\tau) d\tau = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{\langle \tau + \lambda_n^2 \rangle^b}{\langle \tau + \lambda_n^2 \rangle^b} e^{it\tau} \widehat{P_n u}(\tau) d\tau.$$

Then, for  $b > \frac{1}{2}$ , we deduce from the Cauchy-Schwarz inequality that

$$|P_n u(t)| \leq C \left( \int_{\mathbb{R}} \langle \tau + \lambda_n^2 \rangle^{2b} |\widehat{P_n u}(\tau)|^2 d\tau \right)^{1/2}.$$

Thus, by squaring and summing for  $n \in \mathbb{N}$ , we obtain that for any  $b > \frac{1}{2}$  and function  $u \in \overline{X}^{0,b}$ ,

$$\|u\|_{L^\infty(\mathbb{R}; L^2(\mathbb{R}^d))} \leq C \|u\|_{\overline{X}^{0,b}}.$$

On the other hand,

$$\|u\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} = \|u\|_{\overline{X}^{0,0}}$$

thus the result follows by interpolation.  $\square$

**Proposition 2.5.** — For any constant  $0 < \delta < 1$ , there exist two constants  $b' < \frac{1}{2}$  and  $C > 0$  such that for any  $s \in \mathbb{R}$  and any function  $u \in L^{1+\delta}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^d))$ ,

$$\|u\|_{\overline{X}^{s, -b'}} \leq C \|u\|_{L^{1+\delta}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^d))}.$$

*Proof.* — From Proposition 2.4, we have by duality that for any  $\theta \in [0, 1]$  and  $b > \frac{1-\theta}{2}$ , there exists  $C > 0$  such that for any function  $u \in L^{\frac{2}{2-\theta}}(\mathbb{R}; L^2(\mathbb{R}^d))$

$$\|u\|_{\overline{X}^{0, -b}} \leq C \|u\|_{L^{\frac{2}{2-\theta}}(\mathbb{R}; L^2(\mathbb{R}^d))}.$$

Then we choose  $\theta = \frac{2\delta}{1+\delta}$  and  $b = \frac{1-\theta+\delta}{2} < \frac{1}{2}$  to obtain the result. □

In [67, Lemma 2.11], we can obtain the following proposition:

**Proposition 2.6.** — If  $\psi \in C_0^\infty(\mathbb{R})$  then for any  $b \geq 0$ , there exists a constant  $C > 0$  such that for any  $s \in \mathbb{R}$  and any function  $u \in \overline{X}^{s, b}$ ,

$$\|\psi(t)u\|_{\overline{X}^{s, b}} \leq C \|u\|_{\overline{X}^{s, b}}.$$

Finally, we give a last proposition whose proof can be found in [46, Lemma 3.2] (taking  $b' = 1 - b$  and  $T = 1$ ).

**Proposition 2.7.** — Let  $\psi \in C_0^\infty(\mathbb{R})$  then for any  $1 \geq b > \frac{1}{2}$ , there exists a constant  $C > 0$  such that for any  $s \in \mathbb{R}$  and any function  $F \in \overline{X}^{s, b-1}$ ,

$$\left\| \psi(t) \int_0^t e^{-i(t-s)H} F(s) ds \right\|_{\overline{X}^{s, b}} \leq C \|F\|_{\overline{X}^{s, b-1}}.$$

From now on, we set  $T = \frac{\pi}{4}$ , then we define the new Bourgain space which will interest us: we define the space  $\overline{X}_T^{s, b} = \overline{X}^{s, b}([-T; T] \times \mathbb{R}^d)$  as the subset of  $\overline{X}^{s, b}$  for which the following norm

$$\|u\|_{\overline{X}_T^{s, b}} = \inf_{w \in \overline{X}^{s, b}} \left\{ \|w\|_{\overline{X}^{s, b}} \text{ with } w|_{[-T, T]} = u \right\}$$

is finite. Proposition 2.4 allows us to obtain the following result:

**Proposition 2.8.** — Let  $b > \frac{1}{2}$  and  $s \in \mathbb{R}$ , then  $\overline{X}_T^{s, b} \hookrightarrow C^0([-T, T]; \mathcal{H}^s(\mathbb{R}^d))$ .

And finally, thanks to Proposition 2.2 and Proposition 2.3, we obtain the following result:

**Proposition 2.9.** — Let  $b > \frac{1}{2}$ ,  $s \geq 0$  and  $u \in \overline{X}_{\pi/4}^{s, b}$  then  $U \in X^s$  and there exists a constant  $c > 0$  such that for any  $u \in \overline{X}_{\pi/4}^{s, b}$ ,

$$\|U\|_{X^s} \leq c \|u\|_{\overline{X}_{\pi/4}^{s, b}}.$$



**2.4. Basic properties of the eigenfunctions of the harmonic oscillator in dimension  $d = 3$ .** — In this fourth, we give some estimates of the tensor eigenfunctions of the harmonic oscillator.

**Proposition 2.10.** — *For any  $\delta > 0$ , there exists a constant  $C_\delta > 0$  such that for all  $n, m, k \in \mathbb{N}$ ,*

$$(2.9) \quad \|h_n\|_{L^4(\mathbb{R}^3)} \leq C\lambda_n^{-1/4}(\log \lambda_n)^3,$$

$$(2.10) \quad \|h_n\|_{L^\infty(\mathbb{R}^3)} \leq C\lambda_n^{-1/6},$$

*Proof.* — Denote by  $\mu_n = \sqrt{2n+1}$ , so that  $(-\partial_x^2 + x^2)e_n = \mu_n^2 e_n$ . In dimension 1, the following estimates are known by the work [51], namely

$$\|e_n\|_{L^4(\mathbb{R})} \leq C\mu_n^{-1/4} \log \mu_n, \quad \|e_n\|_{L^\infty(\mathbb{R})} \leq C\mu_n^{-1/6}.$$

Let us prove (2.9). In dimension 3, as  $\lambda_n^2 = \mu_{n_1}^2 + \mu_{n_2}^2 + \mu_{n_3}^2$  then there exists  $j \in \{1, 2, 3\}$  such that  $\mu_{n_j}^2 \geq \frac{\lambda_n^2}{3}$ . Thus, we obtain

$$\begin{aligned} \|h_n\|_{L^4(\mathbb{R}^3)} &= \|e_{n_1}\|_{L^4(\mathbb{R})} \|e_{n_2}\|_{L^4(\mathbb{R})} \|e_{n_3}\|_{L^4(\mathbb{R})} \\ &\leq C\mu_{n_1}^{-1/4} \mu_{n_2}^{-1/4} \mu_{n_3}^{-1/4} (\log \mu_{n_1})(\log \mu_{n_2})(\log \mu_{n_3}) \\ &\leq C\lambda_n^{-1/4} (\log \lambda_n)^3, \end{aligned}$$

because  $\mu_{n_1} \leq \lambda_n$ ,  $\mu_{n_2} \leq \lambda_n$  and  $\mu_{n_3} \leq \lambda_n$ . The estimate (2.10) is obtained in a similar manner.  $\square$

**Lemma 2.11.** — *For any  $s \in \mathbb{R}$ , there exists a constant  $C > 0$  such that for all  $f, g, h \in \mathcal{H}^s(\mathbb{R})$ ,*

$$\begin{aligned} \|f(x_1)g(x_2)h(x_3)\|_{\mathcal{H}^s(\mathbb{R}^3)} &\leq \\ &\leq C(\|f\|_{\mathcal{H}^s(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \|h\|_{L^2(\mathbb{R})} + \|f\|_{L^2(\mathbb{R})} \|g\|_{\mathcal{H}^s(\mathbb{R})} \|h\|_{L^2(\mathbb{R})} + \\ &\quad + \|f\|_{L^2(\mathbb{R})} \|g\|_{L^2(\mathbb{R})} \|h\|_{\mathcal{H}^s(\mathbb{R})}). \end{aligned}$$

*Proof.* — We write

$$f(x_1) = \sum_{n=0}^{+\infty} a_n e_n(x_1), \quad g(x_2) = \sum_{m=0}^{+\infty} b_m e_m(x_2), \quad h(x_3) = \sum_{k=0}^{+\infty} c_k e_k(x_3).$$

Then, since the family  $(e_n(x_1)e_m(x_2)e_k(x_3))_{(n,m,k) \in \mathbb{N}^3}$  forms a Hilbertian basis of  $L^2(\mathbb{R}^3)$  composed of eigenfunctions for  $H$  associated to the eigenvalue  $\mu_n^2 + \mu_m^2 + \mu_k^2$ ,

we obtain

$$\begin{aligned}
 \|f(x_1)g(x_2)h(x_3)\|_{\mathcal{H}^s(\mathbb{R}^3)}^2 &= \sum_{n,m,k} |a_n|^2 |b_m|^2 |c_k|^2 (\mu_n^2 + \mu_m^2 + \mu_k^2)^s \leq \\
 &\leq C \sum_{n,m,k \geq 0} |a_n|^2 |b_m|^2 |c_k|^2 \mu_n^{2s} + C \sum_{n,m,k \geq 0} |a_n|^2 |b_m|^2 |c_k|^2 \mu_m^{2s} + \\
 &\quad + C \sum_{n,m,k \geq 0} |a_n|^2 |b_m|^2 |c_k|^2 \mu_k^{2s} \\
 &\leq C (\|f\|_{\mathcal{H}^s(\mathbb{R})}^2 \|g\|_{L^2(\mathbb{R})}^2 \|h\|_{L^2(\mathbb{R})}^2 + \|f\|_{L^2(\mathbb{R})}^2 \|g\|_{\mathcal{H}^s(\mathbb{R})}^2 \|h\|_{L^2(\mathbb{R})}^2 + \\
 &\quad + C \|f\|_{L^2(\mathbb{R})}^2 \|g\|_{L^2(\mathbb{R})}^2 \|h\|_{\mathcal{H}^s(\mathbb{R})}^2),
 \end{aligned}$$

which was the claim. □

**Proposition 2.12.** — *For all  $\delta > 0$  and all  $s \in [0, 1]$ , there exists a constant  $C > 0$  such that for all  $n, m, k \in \mathbb{N}$ ,*

$$(2.11) \quad \|h_n h_m\|_{\mathcal{H}^s(\mathbb{R}^3)} \leq C \max(\lambda_n, \lambda_m)^{s-1/2+\delta},$$

$$(2.12) \quad \|h_n h_m h_k\|_{\mathcal{H}^s(\mathbb{R}^3)} \leq C \max(\lambda_n, \lambda_m, \lambda_k)^{s-1/2+\delta}.$$

*Proof.* — To begin with, recall the following bilinear estimate which is proven in [20, Lemma A.8]: for all  $0 \leq \theta \leq 1$ , we have

$$(2.13) \quad \|e_n e_m\|_{\mathcal{H}^\theta(\mathbb{R})} \leq C \max(\mu_n, \mu_m)^{-1/2+\theta} \min(\log \mu_n, \log \mu_m)^{1/2},$$

with the notation  $\mu_n = \sqrt{2n+1}$ . We first prove (2.11) in the case  $s = 0$ . We can suppose that  $\max(\lambda_n, \lambda_m) = \lambda_n$  and  $\max(\mu_{n_1}, \mu_{n_2}, \mu_{n_3}) = \mu_{n_1}$ , then  $\mu_{n_1}^2 \geq \frac{\lambda_n^2}{3} \geq \frac{\lambda_m^2}{3} \geq \frac{\mu_{m_1}^2}{3}$ . Thus, thanks to (2.13), we obtain that for all  $\delta > 0$

$$\begin{aligned}
 \|h_n h_m\|_{L^2(\mathbb{R}^3)} &= \|e_{n_1} e_{m_1}\|_{L^2(\mathbb{R})} \|e_{n_2} e_{m_2}\|_{L^2(\mathbb{R})} \|e_{n_3} e_{m_3}\|_{L^2(\mathbb{R})} \\
 &\leq \|e_{n_1} e_{m_1}\|_{L^2(\mathbb{R})} \|e_{n_2}\|_{L^4(\mathbb{R})} \|e_{m_2}\|_{L^4(\mathbb{R})} \|e_{n_3}\|_{L^4(\mathbb{R})} \|e_{m_3}\|_{L^4(\mathbb{R})} \\
 &\leq C_\delta \mu_{n_1}^{-1/2+\delta} \\
 &\leq C_\delta \lambda_n^{-1/2+\delta}.
 \end{aligned}$$

We now consider the case  $s = 1$ . Then the general case  $0 \leq s \leq 1$  will follow by interpolation. Using Lemma 2.11, (2.9) and (2.13) with  $\theta = 1$ , we obtain

$$\begin{aligned} \|h_n h_m\|_{\mathcal{H}^1(\mathbb{R}^3)} &\leq \\ &\leq C(\|e_{n_1} e_{m_1}\|_{\mathcal{H}^1(\mathbb{R})} + \|e_{n_2} e_{m_2}\|_{\mathcal{H}^1(\mathbb{R})} + \|e_{n_3} e_{m_3}\|_{\mathcal{H}^1(\mathbb{R})}) \\ &\leq C(\max(\mu_{n_1}, \mu_{m_1})^{1/2+\delta} + \max(\mu_{n_2}, \mu_{m_2})^{1/2+\delta} + \max(\mu_{n_3}, \mu_{m_3})^{1/2+\delta}) \\ &\leq C \max(\lambda_n, \lambda_m)^{1/2+\delta}, \end{aligned}$$

which was the claim.

We turn to the proof of (2.12) with  $s = 0$ . Suppose that  $\max(\lambda_n, \lambda_m, \lambda_k) = \lambda_n$ . Then, by (2.10) and (2.11), we obtain

$$\begin{aligned} \|h_n h_m h_k\|_{L^2(\mathbb{R}^3)} &\leq \|h_n h_m\|_{L^2(\mathbb{R}^3)} \|h_k\|_{L^\infty(\mathbb{R}^3)} \\ &\leq C_\delta \lambda_n^{-1/2+\delta} \\ &\leq C_\delta \max(\lambda_n, \lambda_m, \lambda_k)^{-1/2+\delta}, \end{aligned}$$

hence the result. Next, assume that  $s = 1$ . Thanks to the previous inequality,

$$\begin{aligned} \|h_n h_m h_k\|_{\mathcal{H}^1(\mathbb{R}^3)} &\leq \|h_n h_m\|_{\mathcal{H}^1(\mathbb{R}^3)} \|h_k\|_{L^\infty(\mathbb{R}^3)} + \|h_n h_m\|_{L^2(\mathbb{R}^3)} \|h_k\|_{W^{1,\infty}(\mathbb{R}^3)} \\ &\leq C \max(\lambda_n, \lambda_m)^{1/2+\delta} \lambda_k^{-1/6} + C \max(\lambda_n, \lambda_m)^{-1/2+\delta} \lambda_k^{5/6} \\ &\leq C \max(\lambda_n, \lambda_m, \lambda_k)^{1/2+\delta}. \end{aligned}$$

The general case  $0 \leq s \leq 1$  follows by interpolation.  $\square$

**Lemma 2.13.** — *Let  $\delta > 0$ ,  $\ell \geq 4$  and  $N \geq 1$ , then there exists a constant  $C_N > 0$  such that if we assume  $\lambda_{n_1} \geq \lambda_{n_2}^{1+\delta}$  and  $\lambda_{n_2} \geq \lambda_{n_3} \geq \dots \geq \lambda_{n_\ell}$ , then*

$$\left| \int_{\mathbb{R}^3} \prod_{j=1}^{\ell} h_{n_j}(x) dx \right| \leq C_N \lambda_{n_1}^{-N}.$$

*Proof.* — Using (2.9) and (2.10), we obtain for all  $k \geq 1$

$$\begin{aligned}
 \left| \int_{\mathbb{R}^3} \prod_{j=1}^{\ell} h_{n_j}(x) dx \right| &\leq \lambda_{n_1}^{-2k} \|h_{n_1} H^k(\prod_{j=2}^{\ell} h_{n_j})\|_{L^1(\mathbb{R}^3)} \\
 &\leq \lambda_{n_1}^{-2k} \|h_{n_1}\|_{L^2(\mathbb{R}^3)} \prod_{j=2}^{\ell} \|h_{n_j}\|_{H^{2k}(\mathbb{R}^3)} \\
 &\leq C_k \lambda_{n_1}^{-2k} \lambda_{n_2}^{2k} \prod_{j=2}^{\ell} \|h_{n_j}\|_{L^{2(\ell-1)}(\mathbb{R}^3)} \\
 &\leq C_k \left( \frac{\lambda_{n_2}}{\lambda_{n_1}} \right)^{2k} \\
 &\leq C_k \lambda_{n_1}^{-\frac{2k\delta}{1+\delta}},
 \end{aligned}$$

which was the claim. □

### 3. The bilinear estimate for the harmonic oscillator

The aim of this section is to prove the bilinear estimate of Theorem 1.3. We notice that the result will follow from:

**Theorem 3.1.** — *Let  $d \geq 2$ . For any  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C > 0$  and  $\epsilon > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u, v \in \mathcal{S}'(\mathbb{R}^d)$ ,*

$$\begin{aligned}
 &\|e^{itH} \Delta_N(v) e^{itH} \Delta_M(u)\|_{L^2([-\epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq \\
 &\leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|\Delta_N(v)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{L^2(\mathbb{R}^d)}.
 \end{aligned}$$

Indeed, we can replace  $u$  by  $e^{i\epsilon H} u$  and  $v$  by  $e^{i\epsilon H} v$  to obtain

$$\begin{aligned}
 &\|e^{i(t+\epsilon)H} \Delta_N(v) e^{i(t+\epsilon)H} \Delta_M(u)\|_{L^2([-\epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq \\
 &\leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|\Delta_N(v)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{L^2(\mathbb{R}^d)}.
 \end{aligned}$$

Then, we use the change of variables  $t \longleftrightarrow t + \epsilon$  and Theorem 3.1 to obtain that

$$\begin{aligned}
 &\|e^{itH} \Delta_N(v) e^{itH} \Delta_M(u)\|_{L^2([-\epsilon, 2\epsilon]; L^2(\mathbb{R}^d))} \leq \\
 &\leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|\Delta_N(v)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{L^2(\mathbb{R}^d)}.
 \end{aligned}$$

We can thus iterate the procedure  $2E(\frac{1}{\epsilon})$  times to establish Theorem 1.3 and we are thus reduced to show Theorem 3.1.

Recall that  $\eta \in C_0^\infty(\mathbb{R})$  is such that  $\eta_{[0,1]} = 1$  and  $\eta_{[2,+\infty[} = 0$ , and was used in the definition (1.9) of  $\Delta_N$ . Let  $r \ll 1$  and  $\phi \in C_0^\infty(\mathbb{R})$  satisfy

$$\phi(x) = \begin{cases} 1 & \text{for } x \in [1/4; 2], \\ 0 & \text{for } x \in [0, 1/4 - r] \cup [2 + r, +\infty[, \end{cases}$$

and set  $\Delta'_N = \phi(\frac{H}{N^2})$ . Then using that  $\phi(x)(\eta(x) - \eta(4x)) = \eta(x) - \eta(4x)$  for all  $x \in \mathbb{R}$ , we notice that for all  $N \geq 1$

$$(3.1) \quad \Delta'_N \circ \Delta_N = \Delta_N.$$

Let us then observe that to prove Theorem 3.1, it is sufficient to show

$$(3.2) \quad \|e^{itH} \Delta'_N(v) e^{itH} \Delta'_M(u)\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq \\ \leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|v\|_{L^2(\mathbb{R}^d)} \|u\|_{L^2(\mathbb{R}^d)}.$$

Indeed, if the previous inequality is satisfied, we can apply it to  $v$  replaced by  $\Delta_N(v)$  and  $u$  replaced by  $\Delta_M(u)$  then we can use (3.1) to obtain Theorem 3.1.

• **Case  $M \sim N$  with  $M \geq N$ .** For  $d = 2$ , we can use the Strichartz inequalities (Proposition 2.1) to obtain that

$$\|e^{itH} \Delta'_N(v) e^{itH} \Delta'_M(u)\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq \\ \leq \|e^{itH} \Delta'_N(v)\|_{L^4([- \epsilon, \epsilon]; L^4(\mathbb{R}^d))} \|e^{itH} \Delta'_N(v)\|_{L^4([- \epsilon, \epsilon]; L^4(\mathbb{R}^d))} \\ \leq \|e^{itH} \Delta'_N(v)\|_{L^4([- \pi, \pi]; L^4(\mathbb{R}^d))} \|e^{itH} \Delta'_N(v)\|_{L^4([- \pi, \pi]; L^4(\mathbb{R}^d))} \\ \leq C \|\Delta'_N(v)\|_{L^2(\mathbb{R}^d)} \|\Delta'_N(v)\|_{L^2(\mathbb{R}^d)} \\ \leq C \|v\|_{L^2(\mathbb{R}^d)} \|u\|_{L^2(\mathbb{R}^d)}.$$

For  $d \geq 3$ , using again Proposition 2.1 and the Sobolev embeddings, we establish that

$$\begin{aligned}
 & \|e^{itH} \Delta'_N(v) e^{itH} \Delta'_M(u)\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq \\
 & \leq \|e^{itH} \Delta'_N(v)\|_{L^\infty([- \epsilon, \epsilon]; L^d(\mathbb{R}^d))} \|e^{itH} \Delta'_N(v)\|_{L^2([- \epsilon, \epsilon]; L^{\frac{2d}{d-2}}(\mathbb{R}^d))} \\
 & \leq \|e^{itH} \Delta'_N(v)\|_{L^\infty([- \pi, \pi]; \mathcal{W}^{\frac{d-2}{2}, 2}(\mathbb{R}^d))} \|e^{itH} \Delta'_N(v)\|_{L^2([- \pi, \pi]; L^{\frac{2d}{d-2}}(\mathbb{R}^d))} \\
 & \leq C \|\Delta'_N(v)\|_{\mathcal{H}^{\frac{d-2}{2}}(\mathbb{R}^d)} \|\Delta'_N(v)\|_{L^2(\mathbb{R}^d)} \\
 & \leq CN^{\frac{d-2}{2}} \|v\|_{L^2(\mathbb{R}^d)} \|u\|_{L^2(\mathbb{R}^d)}.
 \end{aligned}$$

Therefore (3.2) is proved in the case  $M \sim N$ .

• **Case  $M \geq 10N$ .** We now have to prove that

$$(3.3) \quad \|e^{itH} v_N e^{itH} u_M\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))} \leq CN^{\frac{d-2}{2}} \left(\frac{N}{M}\right)^{1/2-\delta} \|v\|_{L^2(\mathbb{R}^d)} \|u\|_{L^2(\mathbb{R}^d)}.$$

We can write

$$u_M = \chi\left(\frac{4|x|^2}{M^2}\right)u_M + (1 - \chi)\left(\frac{4|x|^2}{M^2}\right)u_M$$

with  $\chi \in C_0^\infty(\mathbb{R})$ ,  $0 \leq \chi \leq 1$ , satisfying

$$\chi(x) = \begin{cases} 1 & \text{if } |x| \leq 15/32, \\ 0 & \text{if } |x| \geq 1/2. \end{cases}$$

By the triangle inequality, we have to estimate the following two terms:

$$(3.4) \quad \left\| \left( e^{itH} \chi\left(\frac{4|x|^2}{M^2}\right) u_M \right) (e^{itH} v_N) \right\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))}$$

and

$$(3.5) \quad \left\| \left( e^{itH} (1 - \chi)\left(\frac{4|x|^2}{M^2}\right) u_M \right) (e^{itH} v_N) \right\|_{L^2([- \epsilon, \epsilon]; L^2(\mathbb{R}^d))}.$$

**3.1. Estimate of the term (3.4).** — We have the following result.

**Proposition 3.2.** — *For all  $\delta > 0$ , there exists a constant  $C_\delta > 0$  such that for all  $u \in H^{-1/2+\delta}(\mathbb{R}^d)$  and  $v \in \mathcal{H}^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)$ ,*

$$\|e^{itH} u e^{itH} v\|_{L^2([- \frac{\pi}{8}, \frac{\pi}{8}]; L^2(\mathbb{R}^d))} \leq C_\delta \|u\|_{H^{-1/2+\delta}(\mathbb{R}^d)} \|v\|_{\mathcal{H}^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)}.$$

*Proof.* — From [65, Theorem 2.4] (coming from [7]) we have that for all  $\delta > 0$ , there exists a constant  $C_\delta > 0$  such that for all  $u \in H^{-1/2+\delta}(\mathbb{R}^d)$  and  $v \in H^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)$ ,

$$\|e^{it\Delta}u e^{it\Delta}v\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} \leq C_\delta \|u\|_{H^{-1/2+\delta}(\mathbb{R}^d)} \|v\|_{H^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)}.$$

Using the lens transformation (see Section 2.1) and the previous inequality, we obtain that

$$\begin{aligned} & \|e^{itH}u e^{itH}v\|_{L^2([-\frac{\pi}{8}, \frac{\pi}{8}]; L^2(\mathbb{R}^d))}^2 = \\ &= \|e^{-itH}u e^{-itH}v\|_{L^2([-\frac{\pi}{8}, \frac{\pi}{8}]; L^2(\mathbb{R}^d))}^2 \\ &= \int_{]-\frac{\pi}{8}, \frac{\pi}{8}[} \int_{\mathbb{R}^d} \frac{1}{|\cos(2t)|^{2d}} |e^{it\Delta}u e^{it\Delta}v|^2 \left( \frac{\tan(2t)}{2}, \frac{x}{\cos(2t)} \right) dx dt \\ &= \int_{]-\frac{\pi}{8}, \frac{\pi}{8}[} \int_{\mathbb{R}^d} \frac{1}{|\cos(2t)|^d} |e^{it\Delta}u e^{it\Delta}v|^2 \left( \frac{\tan(2t)}{2}, x \right) dx dt \\ &= \int_{-1/2}^{1/2} \int_{\mathbb{R}^d} (1 + (2t)^2)^{d/2-1} |e^{it\Delta}u e^{it\Delta}v|^2(t, x) dx dt \\ &\leq C \int_{\mathbb{R}} \int_{\mathbb{R}^d} |e^{it\Delta}u e^{it\Delta}v|^2(t, x) dx dt \\ &\leq C \|u\|_{H^{-1/2+\delta}(\mathbb{R}^d)}^2 \|v\|_{H^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)}^2 \\ &\leq C \|u\|_{H^{-1/2+\delta}(\mathbb{R}^d)}^2 \|v\|_{\mathcal{H}^{\frac{d-1}{2}-\delta}(\mathbb{R}^d)}^2, \end{aligned}$$

where in the last inequality we used (1.2).  $\square$

We will show that for all  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C_\delta > 0$  such that for all dyadic  $N, M \geq 1$ , and  $u, v \in L^2(\mathbb{R}^d)$ , if  $M \geq N$  then

$$\begin{aligned} & \|(e^{itH}v_N)(e^{itH}\chi\left(\frac{4|x|^2}{M^2}\right)u_M)\|_{L^2([-\frac{\pi}{8}, \frac{\pi}{8}]; L^2(\mathbb{R}^d))} \leq \\ & \leq C_\delta N^{\frac{d-2}{2}} \left(\frac{N}{M}\right)^{1/2-\delta} \|u\|_{L^2(\mathbb{R}^d)} \|v\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

To do this, using Proposition 3.2, it suffices to prove that for any  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C_\delta > 0$  such that for all dyadic  $M \geq 1$  and  $u \in L^2(\mathbb{R}^d)$

$$\|\chi\left(\frac{4|x|^2}{M^2}\right)u_M\|_{H^{-1/2+\delta}(\mathbb{R}^d)} \leq C_\delta M^{-1/2+\delta} \|u\|_{L^2(\mathbb{R}^d)}.$$

We have

$$\|\chi\left(\frac{4|x|^2}{M^2}\right)u_M\|_{L^2(\mathbb{R}^d)} \leq \|u_M\|_{L^2(\mathbb{R}^d)} \leq C\|u\|_{L^2(\mathbb{R}^d)}.$$

Thus, by interpolation, it is sufficient to show that there exists a constant  $C > 0$  such that for all dyadic  $M \geq 1$  and  $u \in L^2(\mathbb{R}^d)$ ,

$$(3.6) \quad \|\chi\left(\frac{4|x|^2}{M^2}\right)u_M\|_{H^{-1}(\mathbb{R}^d)} \leq CM^{-1}\|u\|_{L^2(\mathbb{R}^d)}.$$

We then use the semi-classical calculus. For a function  $u$ , we define  $\mathbf{u} : x \mapsto u\left(\frac{x}{\sqrt{h}}\right)$  where  $h = \frac{1}{M^2}$ .

Observe that

$$(3.7) \quad \chi\left(\frac{4|x|^2}{M^2}\right)\left[\frac{H}{M^2}\right](u)(x) = [\chi(4|x|^2)(-h^2\Delta + |x|^2)](\mathbf{u})(\sqrt{h}x)$$

and that

$$(3.8) \quad \chi\left(\frac{4|x|^2}{M^2}\right)\left[\phi\left(\frac{H}{M^2}\right)\right](u)(x) = [\chi(4|x|^2)\phi(-h^2\Delta + |x|^2)](\mathbf{u})(\sqrt{h}x).$$

Thus, to prove (3.6), it is sufficient to show that there exists a constant  $C > 0$  such that for all  $h \in ]0, 1]$  and  $u \in L^2(\mathbb{R}^d)$ ,

$$(3.9) \quad \|\chi(4|x|^2)\phi(|x|^2 + |h\nabla|^2)u\|_{H^{-1}(\mathbb{R}^d)} \leq Ch\|u\|_{L^2(\mathbb{R}^d)}.$$

Indeed,

$$\begin{aligned} \|\chi\left(\frac{4|x|^2}{M^2}\right)u_M\|_{H^{-1}(\mathbb{R}^d)} &\leq \|[\chi(4|x|^2)\phi(-h^2\Delta + |x|^2)](\mathbf{u})(\sqrt{h}\cdot)\|_{H^{-1}(\mathbb{R}^d)} \\ &\leq h^{-\frac{1}{2}-\frac{d}{4}}\|[\chi(4|x|^2)\phi(-h^2\Delta + |x|^2)](\mathbf{u})\|_{H^{-1}(\mathbb{R}^d)} \\ &\leq Ch^{\frac{1}{2}-\frac{d}{4}}\|\mathbf{u}\|_{L^2(\mathbb{R}^d)} \\ &\leq Ch^{1/2}\|u\|_{L^2(\mathbb{R}^d)} \\ &\leq CM^{-1}\|u\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

Let us then prove (3.9). Thanks to Proposition A.5, we have

$$\begin{aligned} &\|\chi(4|x|^2)\phi(|x|^2 + |h\nabla|^2)u\|_{H^{-1}(\mathbb{R}^d)} \leq \\ &\leq \|\chi(4|x|^2)[\phi(|x|^2 + |h\nabla|^2) - \text{Op}_h(\phi(|x|^2 + |\xi|^2))\chi_2]u\|_{H^{-1}(\mathbb{R}^d)} + \\ &\quad + \|\chi(4|x|^2)\text{Op}_h(\phi(|x|^2 + |\xi|^2))\chi_2u\|_{H^{-1}(\mathbb{R}^d)} \\ &\leq h\|u\|_{L^2(\mathbb{R}^d)} + \|\chi(4|x|^2)\text{Op}_h(\phi(|x|^2 + |\xi|^2))\chi_2u\|_{H^{-1}(\mathbb{R}^d)}. \end{aligned}$$



Thus, it is sufficient to evaluate  $\|\chi(4|x|^2)Op_h(\phi(|x|^2 + |\xi|^2))\chi_2u\|_{H^{-1}(\mathbb{R}^d)}$ . We have

$$\begin{aligned}
& \chi(4|x|^2)Op_h(\phi(|x|^2 + |\xi|^2))\chi_2u = \\
&= \frac{\chi(4|x|^2)}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \phi(|x|^2 + (h\xi)^2) \mathcal{F}(\chi_2u)(\xi) d\xi \\
&= \frac{h}{(2\pi)^d} \int_{\mathbb{R}^d} (h\xi) \cdot \nabla(e^{ix \cdot \xi}) \frac{\chi(4|x|^2)}{i|h\xi|^2} \phi(|x|^2 + (h\xi)^2) \mathcal{F}(\chi_2u)(\xi) d\xi \\
&= \frac{h}{(2\pi)^d} \left( \int_{\mathbb{R}^d} (h\xi) \cdot \nabla \left( e^{ix \cdot \xi} \frac{\chi(4|x|^2)}{i|h\xi|^2} \phi(|x|^2 + |h\xi|^2) \mathcal{F}(\chi_2u)(\xi) \right) d\xi \right. \\
&\quad \left. - \int_{\mathbb{R}^d} e^{ix \cdot \xi} (h\xi) \cdot \nabla \left( \frac{\chi(4|x|^2)}{i|h\xi|^2} \phi(|x|^2 + |h\xi|^2) \right) \mathcal{F}(\chi_2u)(\xi) d\xi \right) \\
&= \frac{h}{(2\pi)^d} \left( \operatorname{div}_x \left( \int_{\mathbb{R}^d} (h\xi) e^{ix \cdot \xi} \frac{\chi(4|x|^2)}{i|h\xi|^2} \phi(|x|^2 + |h\xi|^2) \mathcal{F}(\chi_2u)(\xi) d\xi \right) \right. \\
&\quad \left. - \int_{\mathbb{R}^d} e^{ix \cdot \xi} (h\xi) \cdot \nabla \left( \frac{\chi(4|x|^2)}{i|h\xi|^2} \phi(|x|^2 + |h\xi|^2) \right) \mathcal{F}(\chi_2u)(\xi) d\xi \right).
\end{aligned}$$

Then, since  $4|x|^2 \leq \frac{1}{2}$  and  $\frac{1}{4}^- \leq |x|^2 + |\xi|^2 \leq 2^+$  implies  $\frac{1}{8}^- \leq |\xi|^2$ , we deduce that

$$(x, \xi) \longrightarrow \xi \frac{\chi(4|x|^2)}{|\xi|^2} \phi(|x|^2 + |\xi|^2) \in S^0$$

and

$$(x, \xi) \longrightarrow \xi \cdot \nabla \left( \frac{\chi(4|x|^2)}{|\xi|^2} \phi(|x|^2 + |\xi|^2) \right) \in S^0.$$

Thus, by Theorem A.2, we have

$$\begin{aligned}
& \left\| \frac{h}{(2\pi)^d} \operatorname{div}_x \left( \int_{\mathbb{R}^d} (h\xi) e^{ix \cdot \xi} \frac{\chi(4|x|^2)}{i(h\xi)^2} \phi(|x|^2 + (h\xi)^2) \mathcal{F}(\chi_2u)(\xi) d\xi \right) \right\|_{H^{-1}(\mathbb{R}^d)} \leq \\
& \leq \frac{h}{(2\pi)^d} \left\| \int_{\mathbb{R}^d} (h\xi) e^{ix \cdot \xi} \frac{\chi(4|x|^2)}{i(h\xi)^2} \phi(|x|^2 + (h\xi)^2) \mathcal{F}(\chi_2u)(\xi) d\xi \right\|_{L^2(\mathbb{R}^d)} \\
& \leq Ch \|\chi_2u\|_{L^2(\mathbb{R}^d)} \\
& \leq Ch \|u\|_{L^2(\mathbb{R}^d)}
\end{aligned}$$

and

$$\begin{aligned}
& \left\| \frac{h}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} (h\xi) \cdot \nabla \left( \frac{\chi(4|x|^2)}{i(h\xi)^2} \phi(|x|^2 + (h\xi)^2) \right) \mathcal{F}(\chi_2 u)(\xi) d\xi \right\|_{H^{-1}(\mathbb{R}^d)} \leq \\
& \leq \left\| \frac{h}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} (h\xi) \cdot \nabla \left( \frac{\chi(4|x|^2)}{i(h\xi)^2} \phi(|x|^2 + (h\xi)^2) \right) \mathcal{F}(\chi_2 u)(\xi) d\xi \right\|_{L^2(\mathbb{R}^d)} \\
& \leq Ch \|\chi_2 u\|_{L^2(\mathbb{R}^d)} \\
& \leq Ch \|u\|_{L^2(\mathbb{R}^d)}.
\end{aligned}$$

This proves (3.9) and the estimate (3.4) follows.

### 3.2. Estimate of the term (3.5). —

**Proposition 3.3.** — *There exists a time  $T \in ]0, \frac{\pi}{4}[$  such that for all  $K \geq 0$ , there exists a constant  $C_K > 0$  such that for all  $M \geq 1$  and  $u \in L^2(\mathbb{R}^d)$ ,*

$$\left\| \chi \left( \frac{8|x|^2}{M^2} \right) e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq C_K M^{-K} \|u\|_{L^2(\mathbb{R}^d)}.$$

Let us show that Proposition 3.3 implies the bound (3.5). From the Sobolev embedding

$$\|u\|_{L^\infty([a, b]; L^\infty(\mathbb{R}^d))} \leq C \|u\|_{L^\infty([a, b]; \mathcal{H}^{\frac{d}{2}+1}(\mathbb{R}^d))}$$

and Proposition 3.3, we deduce for  $M \geq 10N$  that

$$\begin{aligned}
& \left\| \chi \left( \frac{8|x|^2}{M^2} \right) \left( e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right) (e^{itH} v_N) \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq \\
& \leq \left\| \chi \left( \frac{8|x|^2}{M^2} \right) e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \|e^{itH} v_N\|_{L^\infty([-T, T]; L^\infty(\mathbb{R}^d))} \\
& \leq C \left\| \chi \left( \frac{8|x|^2}{M^2} \right) e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \|v_N\|_{\mathcal{H}^{\frac{d}{2}+1}(\mathbb{R}^d)} \\
& \leq C_K M^{-K} N^{\frac{d}{2}+1} \|u\|_{L^2(\mathbb{R}^d)} \|v\|_{L^2(\mathbb{R}^d)} \\
& \leq C_K M^{-K + \frac{d}{2}+1} \|u\|_{L^2(\mathbb{R}^d)} \|v\|_{L^2(\mathbb{R}^d)}.
\end{aligned}$$

And so, it is enough to estimate, for  $M \geq 10N$ , the following term:

$$\left\| \left( (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \right) \left( e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right) (e^{itH} v_N) \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))}.$$

Thanks to Proposition 2.1, we obtain that for all  $R \geq 1$ ,

$$\begin{aligned}
& \left\| (e^{itH} v_N)(1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \left( e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right) \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} = \\
& = \left\| \langle x \rangle^R (e^{itH} v_N) \langle x \rangle^{-R} (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \left( e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right) \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\
& \leq M^{-R} \left\| \langle x \rangle^R (e^{itH} v_N) (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \left( e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right) \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\
& \leq M^{-R} \left\| (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \langle x \rangle^R (e^{itH} v_N) \right\|_{L^4([-T, T]; L^4(\mathbb{R}^d))} \\
& \quad \cdot \left\| e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right\|_{L^4([-T, T]; L^4(\mathbb{R}^d))} \\
& \leq M^{-R} \left\| (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \langle x \rangle^R (e^{itH} v_N) \right\|_{L^4([-T, T]; L^4(\mathbb{R}^d))} \\
& \quad \cdot \left\| e^{itH} (1 - \chi) \left( \frac{4|x|^2}{M^2} \right) u_M \right\|_{L^4([-T, T]; \mathcal{W}^{\frac{d-2}{4}, \frac{2d}{d-1}}(\mathbb{R}^d))} \\
& \leq M^{-R} \left\| \langle x \rangle^R (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) (e^{itH} v_N) \right\|_{L^\infty([-T, T]; L^4(\mathbb{R}^d))} M^{\frac{d-2}{4}} \|u_M\|_{L^2(\mathbb{R}^d)}.
\end{aligned}$$

Then, since

$$\text{Supp} \left\{ (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \right\} \subset \left\{ |x|^2 \geq \frac{M^2}{8} \frac{15}{16} \right\} \subset \{|x|^2 \geq 5N^2\}$$

from Proposition A.4, we deduce

$$\left\| (1 - \chi) \left( \frac{8|x|^2}{M^2} \right) \langle x \rangle^R e^{itH} v_N \right\|_{L^4([-T, T]; L^4(\mathbb{R}^d))} \leq C \|v\|_{L^2(\mathbb{R}^d)}.$$

*Proof of Proposition 3.3.* — Using semi-classical analysis as in (3.8), it suffices to prove the existence of a time  $T \in ]0, \frac{\pi}{4}[$  such that for all  $N \geq 1$ , there exists a constant  $C_N > 0$  such that for all  $u \in L^2(\mathbb{R}^d)$  and  $h \in ]0, 1]$ ,

$$\left\| \chi(8|x|^2) e^{itH_h/h} (1 - \chi) (4|x|^2) \phi(|x|^2 + |h\nabla|^2) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq C_N h^N \|u\|_{L^2(\mathbb{R}^d)}.$$

Using Proposition A.5, it is sufficient to establish the following result: for any function  $g(x, \xi) \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  satisfying  $\text{Supp}(g) \subset \{\frac{1}{8} \leq |\xi|^2 + |x|^2 \leq 4\}$  and all integer  $N \geq 1$ , there exists a constant  $C_N > 0$  such that for all  $u \in L^2(\mathbb{R}^d)$  and  $h \in ]0, 1]$ ,

$$\left\| \chi(8|x|^2) e^{itH_h/h} (1 - \chi) (4|x|^2) \text{Op}_h(g) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq C_N h^N \|u\|_{L^2(\mathbb{R}^d)}.$$

Indeed, by Proposition A.5,

$$\begin{aligned}
 & \left\| \chi(8|x|^2) e^{itH_h/h} (1-\chi)(4|x|^2) \phi(|x|^2 + |h\nabla|^2) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq \\
 & \leq \left\| \chi(8|x|^2) e^{itH_h/h} (1-\chi)(4|x|^2) [\phi(|x|^2 + |h\nabla|^2) u - \sum_{j=0}^{N-1} h^j \text{Op}_h(\Psi_j) u] \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\
 & \quad + \sum_{j=0}^{N-1} h^j \left\| \chi(8|x|^2) e^{itH_h/h} (1-\chi)(4|x|^2) \text{Op}_h(\Psi_j(x, \xi)) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\
 & \leq C_N h^N \|u\|_{L^2(\mathbb{R}^d)} + \sum_{j=0}^{N-1} h^j \left\| \chi(8|x|^2) e^{itH_h/h} (1-\chi)(4|x|^2) \text{Op}_h(\Psi_j) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\
 & \leq C_N h^N \|u\|_{L^2(\mathbb{R}^d)}.
 \end{aligned}$$

□

**Lemma 3.4.** — *There exists a time  $T \in ]0, \frac{\pi}{4}[$  such that if  $g \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  satisfying  $\text{Supp}(g) \subset \{\frac{1}{8} \leq |\xi|^2 + |x|^2 \leq 4\}$  then for any integer  $K \geq 1$ , there exists a constant  $C_K > 0$  such that for all  $u \in L^2(\mathbb{R}^d)$  and  $h \in ]0, 1]$ ,*

$$\left\| \chi(8|x|^2) e^{itH_h/h} (1-\chi)(4|x|^2) \text{Op}_h(g) u \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq C_K h^K \|u\|_{L^2(\mathbb{R}^d)}.$$

*Proof.* — We define

$$w(s, x) = \int_{\mathbb{R}^d} e^{\frac{i}{h}\Phi(s, x, \xi)} a(s, x, \xi, h) \hat{u}\left(\frac{\xi}{h}\right) \frac{d\xi}{(2\pi h)^d}$$

where

$$a(s, x, \xi, h) = \sum_{j=0}^K h^j a_j(s, x, \xi).$$

Assume that

$$\begin{cases} \Phi(0, x, \xi) = x \cdot \xi, \\ \partial_s \Phi - |\nabla \Phi|^2 - |x|^2 = 0, \\ a_0(0, x, \xi) = (1-\chi)(4|x|^2)g(x, \xi), \\ \partial_s a_0 - 2\nabla a_0 \cdot \nabla \Phi - a_0 \Delta \Phi = 0, \end{cases}$$

and for  $1 \leq j \leq K$

$$\begin{cases} a_j(0, x, \xi) = 0, \\ \partial_s a_j - 2\nabla a_j \cdot \nabla \Phi - a_j \Delta \Phi = -i\Delta a_{j-1}. \end{cases}$$

Then

$$\begin{aligned} ih\partial_s w + (-h^2\Delta + |x|^2)w &= -h^{K+2} \int_{\mathbb{R}^d} e^{\frac{i}{h}\Phi(s,x,\xi)} \Delta(a_K(s,x,\xi)) \hat{u}\left(\frac{\xi}{h}\right) \frac{d\xi}{(2\pi h)^d} \\ &:= h^{K+2} f. \end{aligned}$$

Since  $w_0 = (1 - \chi)(4|x|^2)Op_h(g(x, \xi))u$ , hence

$$w = e^{itH_h/h}(1 - \chi)(4|x|^2)Op_h(g(x, \xi))u - ih^{K+1} \int_0^t e^{i(t-s)H_h/h} f(s) ds,$$

which in turn implies

$$\begin{aligned} \chi(8|x|^2)e^{itH_h/h}(1 - \chi)(4|x|^2)Op_h(g(x, \xi))u &= \\ &= \chi(8|x|^2)w(t, x) - ih^{K+1}\chi(8|x|^2) \int_0^t e^{i(t-s)H_h/h} f(s) ds. \end{aligned}$$

Let us notice that if  $\Psi$  is a solution of the equation  $\partial_t \Psi + |\nabla \Psi|^2 = 0$  with initial data  $\Psi(0, x, \xi) = x \cdot \xi$  then  $\Phi(t, x, \xi) = \Psi\left(-\frac{\tan 2t}{2}, \frac{x}{\cos 2t}, \xi\right) + \frac{|x|^2 \tan 2t}{2}$  is a solution of the equation  $\partial_t \Phi - |\nabla \Phi|^2 - |x|^2 = 0$  with the same initial data.

By the method of characteristics, we obtain

$$\Psi(t, x, \xi) = -t|\xi|^2 + x \cdot \xi,$$

then we deduce that

$$\Phi(t, x, \xi) = \frac{\tan(2t)}{2}(|\xi|^2 + |x|^2) + \frac{x \cdot \xi}{\cos(2t)}.$$

Hence

$$\nabla \Phi = \frac{\xi}{\cos(2t)} + x \tan(2t) \quad \text{and} \quad \Delta \Phi = d \tan(2t).$$

Using the method of characteristics, we obtain

$$(3.10) \quad a_0(t, x - 2 \int_0^t \nabla \Phi, \xi) = \frac{a_0(0, x, \xi)}{|\cos 2t|^{\frac{d}{2}}},$$

and

$$(3.11) \quad a_j(t, x - 2 \int_0^t \nabla \Phi, \xi) = -i \int_0^t \left| \frac{\cos(2s)}{\cos(2t)} \right|^{\frac{d}{2}} \Delta a_{j-1}(s, x - 2 \int_0^s \nabla \Phi, \xi) ds.$$

Now, for any  $\xi \in \mathbb{R}^d$  and  $|x| \leq \frac{1}{2}$ ,

$$a_0(0, x, \xi) = 0.$$

As a consequence, for all  $\xi \in \mathbb{R}^d$ ,  $t \in ]-\frac{\pi}{4}, \frac{\pi}{4}[$ ,  $|x| \leq \frac{\sqrt{15}}{8\sqrt{2}}$  and  $j \in \mathbb{N}$

$$a_j(t, x - 2 \int_0^t \nabla \Phi, \xi) = 0.$$

We have  $\int_0^t \nabla \Phi = \xi F(t) - \frac{x \log \cos(2t)}{2}$  with  $F$  a continuous function satisfying  $F(0) = 0$ . Thus, for all  $\epsilon > 0$ , there exists a time  $T \in ]0, \frac{\pi}{4}]$  such that if  $|t| \leq T$  then  $|F(t)| \leq \epsilon$  and  $|\log \cos 2t| \leq \epsilon$ .

Observe that

$$y = x - 2 \int_0^t \nabla \Phi = x(1 + \log \cos 2t) - 2\xi F(t)$$

can be solved by

$$x = \frac{y + 2\xi F(t)}{1 + \log \cos 2t}.$$

So if  $|y| \leq \frac{1}{4}$ ,  $|\xi|^2 \leq 4$  and  $|t| \leq T$  then  $|x| \leq \frac{1/4+4\epsilon}{1-\epsilon} \leq \frac{\sqrt{15}}{8\sqrt{2}}$ , if  $\epsilon < 1$  is small enough.

This implies that for any  $j \in \mathbb{N}$  and  $|t| \leq T$

$$\text{Supp}(a_j(t)) \subset B_x(0, \frac{1}{4})^c \times B_\xi(0, 2).$$

Therefore, if  $|t| \leq T$ , since  $\text{Supp}(\chi(8|x|^2)) \subset B_x(0, \frac{1}{4})$ , we deduce that

$$\chi(8|x|^2)e^{itH_h/h}(1 - \chi)(4|x|^2)\mathcal{O}p_h(g(x, \xi))u = -ih^{K+1}\chi(8|x|^2) \int_0^t e^{i(t-s)H_h/h} f(s) ds.$$

Then, by Theorem 2.1, we obtain

$$\begin{aligned} & \|\chi(8|x|^2)e^{itH_h/h}(1 - \chi)(4|x|^2)\mathcal{O}p_h(g(x, \xi))u\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \leq \\ & \leq h^{K+1} \left\| \chi(8|x|^2) \int_0^t e^{i(t-s)H_h/h} f(s) ds \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \\ & \leq h^{K+1} \|f\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \\ & \leq h^{K+1} \|\Delta a_K\|_{L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d)))} \|u\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

The lemma is therefore proved if  $\Delta a_K \in L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d)))$ .

We prove by induction on  $K \in \mathbb{N}$ , that for all  $\alpha \in \mathbb{N}^d$ ,

$$\partial_x^\alpha a_K \in L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d))).$$

For  $K = 0$ , using (3.10), we see by change of variables that

$$\partial_x^\alpha a_0 \in L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d)))$$

if  $\partial_x^\alpha a_0(0) \in L_x^2(\mathbb{R}^d, L_\xi^2(\mathbb{R}^d))$ . But  $a_0(0) \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  with  $\text{Supp}(a_0(0)) \subset \{(x, \xi) : |x|^2 \leq 1, |\xi|^2 \leq 4\}$  and the case  $K = 0$  is obvious.

Let us assume the result established at rank  $K - 1$  and show it at rank  $K$ . Using (3.11), we note that  $\partial_x^\alpha a_K \in L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d)))$  if  $\partial_x^{\alpha+2} a_{K-1} \in L_t^1([-T, T]; L_x^2(\mathbb{R}^d; L_\xi^2(\mathbb{R}^d)))$ . This last statement being clear by induction hypothesis.  $\square$

**3.3. Bilinear estimates and Bourgain spaces.** — The aim of this section is to write the bilinear estimate of Theorem 1.3 in Bourgain spaces. More precisely, we establish the following result:

**Proposition 3.5.** — *Let  $\psi \in C_0^\infty(\mathbb{R})$ . There exists  $\delta_0 \in ]0, \frac{1}{2}]$  such that for all  $\delta \in ]0, \delta_0]$ , there exist  $b' < \frac{1}{2}$  and a constant  $C > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u_0, u, v \in \mathcal{S}'(\mathbb{R}^d)$ ,*

$$(3.12) \quad \|\Delta_N(v)\Delta_M(u)\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} \leq \\ \leq C \min(N, M)^{\frac{d-2}{2} + \delta} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2 - \delta} \|\Delta_N(v)\|_{\overline{X}^{0, b'}} \|\Delta_M(u)\|_{\overline{X}^{0, b'}}.$$

and

$$(3.13) \quad \|\Delta_N(\psi(t)e^{-itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} \leq \\ \leq C \min(N, M)^{\frac{d-2}{2} + \delta} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2 - \delta} \|\Delta_N(u_0)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{\overline{X}^{0, b'}}.$$

To prove these results, we adapt the proof of [18, Lemma 4.4]. Let us begin by noting that it is sufficient to prove the following two propositions:

**Proposition 3.6.** — *For all  $b \in ]\frac{1}{2}, 1]$  and  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u, v \in \mathcal{S}'(\mathbb{R}^d)$ ,*

$$\|\Delta_N(v)\Delta_M(u)\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} \leq \\ \leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2 - \delta} \|\Delta_N(v)\|_{\overline{X}^{0, b}} \|\Delta_M(u)\|_{\overline{X}^{0, b}}.$$

**Proposition 3.7.** — *Let  $\psi \in C_0^\infty(\mathbb{R})$  then for all  $b \in ]\frac{1}{2}, 1]$  and  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u_0, v \in \mathcal{S}'(\mathbb{R}^d)$*

$$\|\Delta_N(\psi(t)e^{-itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^d))} \leq \\ \leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2 - \delta} \|\Delta_N(u_0)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{\overline{X}^{0, b}}.$$

Indeed, for any  $\epsilon > 0$ , according to Proposition 2.4 (with  $\theta = \frac{1}{2}$ ), we obtain

$$\begin{aligned} \|\Delta_N(v)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} &\leq \|\Delta_N(v)\|_{L^4(\mathbb{R};L^2(\mathbb{R}^d))}\|\Delta_M(u)\|_{L^4(\mathbb{R};L^\infty(\mathbb{R}^d))} \\ &\leq C\|\Delta_N(v)\|_{\overline{X}^{0,1/4+\epsilon}}\|\Delta_M(u)\|_{\overline{X}^{d/2+\epsilon,1/4+\epsilon}}, \end{aligned}$$

$$\begin{aligned} \|\Delta_N(\psi(t)e^{itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} &\leq \\ &\leq \|\Delta_N(\psi(t)e^{itH}u_0)\|_{L^4(\mathbb{R};L^\infty(\mathbb{R}^d))}\|\Delta_M(u)\|_{L^4(\mathbb{R};L^2(\mathbb{R}^d))} \\ &\leq C\|\Delta_N(u_0)\|_{\mathcal{H}^{d/2-1/2+\epsilon}(\mathbb{R}^d)}\|\Delta_M(u)\|_{\overline{X}^{0,1/4+\epsilon}} \\ &\leq C\|\Delta_N(u_0)\|_{\mathcal{H}^{d/2+\epsilon}(\mathbb{R}^d)}\|\Delta_M(u)\|_{\overline{X}^{0,1/4+\epsilon}}, \end{aligned}$$

and

$$\begin{aligned} \|\Delta_N(\psi(t)e^{itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} &\leq \\ &\|\Delta_N(\psi(t)e^{itH}u_0)\|_{L^4(\mathbb{R};L^2(\mathbb{R}^d))}\|\Delta_M(u)\|_{L^4(\mathbb{R};L^\infty(\mathbb{R}^d))} \\ &\leq C\|\Delta_N(\psi(t)e^{itH}u_0)\|_{L^\infty(\mathbb{R};L^2(\mathbb{R}^d))}\|\Delta_M(u)\|_{\overline{X}^{d/2+\epsilon,1/4+\epsilon}} \\ &\leq C\|\Delta_N(u_0)\|_{L^2(\mathbb{R}^d)}\|\Delta_M(u)\|_{\overline{X}^{d/2+\epsilon,1/4+\epsilon}}. \end{aligned}$$

Therefore, by interpolation, for all  $\theta \in [0, 1]$ , we obtain

$$\begin{aligned} \|\Delta_N(v)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} &\leq \\ &\leq C\min(M, N)^{\frac{d-2}{2}+\theta(1+\epsilon)}\left(\frac{\min(M, N)}{\max(M, N)}\right)^{(1/2-\delta)(1-\theta)} \\ &\quad \cdot \|\Delta_N(v)\|_{\overline{X}^{0,b(1-\theta)+\theta(1/4+\epsilon)}}\|\Delta_M(u)\|_{\overline{X}^{0,b(1-\theta)+\theta(1/4+\epsilon)}} \end{aligned}$$

and

$$\begin{aligned} \|\Delta_N(\psi(t)e^{itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} &\leq \\ &\leq C\min(M, N)^{\frac{d-2}{2}+\theta(1+\epsilon)}\left(\frac{\min(M, N)}{\max(M, N)}\right)^{(1/2-\delta)(1-\theta)} \\ &\quad \cdot \|\Delta_N(u_0)\|_{L^2(\mathbb{R}^d)}\|\Delta_M(u)\|_{\overline{X}^{0,b(1-\theta)+\theta(1/4+\epsilon)}}. \end{aligned}$$

Choose  $\delta = \frac{\epsilon}{2}$  and  $\theta = \frac{\epsilon}{4}$  then

$$\begin{aligned} b(1-\theta) + \theta\left(\frac{1}{4} + \epsilon\right) &= b - \frac{b\epsilon}{4} + \frac{\epsilon}{4}\left(\frac{1}{4} + \epsilon\right) \\ &\leq b - \frac{\epsilon}{8} + \frac{\epsilon}{16} + \frac{\epsilon^2}{4} \leq b - \frac{\epsilon}{17}. \end{aligned}$$



It is then enough to take  $b = \frac{1}{2} + \frac{\epsilon}{34}$  and to set  $b' = b(1 - \theta) + \theta(\frac{1}{4} + \epsilon) < \frac{1}{2}$  to obtain

$$\begin{aligned} & \|\Delta_N(v)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} \leq \\ & \leq C \min(N, M)^{\frac{d-2}{2}+\epsilon} \left( \frac{\min(M, N)}{\max(N, M)} \right)^{(1/2-\delta)(1-\theta)} \|\Delta_N(v)\|_{\overline{X}^{0,b'}} \|\Delta_M(u)\|_{\overline{X}^{0,b'}} \end{aligned}$$

and

$$\begin{aligned} & \|\Delta_N(\psi(t)e^{-itH}u_0)\Delta_M(u)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^d))} \leq \\ & \leq C \min(N, M)^{\frac{d-2}{2}+\epsilon} \left( \frac{\min(M, N)}{\max(M, N)} \right)^{(1/2-\delta)(1-\theta)} \|\Delta_N(u_0)\|_{L^2(\mathbb{R}^d)} \|\Delta_M(u)\|_{\overline{X}^{0,b'}}. \end{aligned}$$

To conclude, it is sufficient to note that

$$\left( \frac{1}{2} - \delta \right) (1 - \theta) = \frac{1}{2} - \frac{5\epsilon}{8} + \frac{\epsilon^2}{8} \geq \frac{1}{2} - \epsilon$$

and the inequalities (3.12) and (3.13) follow with  $\delta_0 = \epsilon$ .

Then, as for [18, Lemma 4.4], to prove Proposition 3.6, it is sufficient to establish the following proposition:

**Proposition 3.8.** — *For all  $b \in ]\frac{1}{2}, 1]$  and  $\delta \in ]0, \frac{1}{2}]$ , there exists a constant  $C > 0$  such that for all dyadic numbers  $N, M \geq 1$  and  $u, v \in \mathcal{S}'(\mathbb{R}^d)$*

$$\begin{aligned} & \|\Delta_N(v)\Delta_M(u)\|_{L^2([0,1];L^2(\mathbb{R}^d))} \leq \\ & \leq C \min(N, M)^{\frac{d-2}{2}} \left( \frac{\min(N, M)}{\max(N, M)} \right)^{1/2-\delta} \|\Delta_N(v)\|_{\overline{X}^{0,b}} \|\Delta_M(u)\|_{\overline{X}^{0,b}}. \end{aligned}$$

Finally, to obtain Proposition 3.7 and Proposition 3.8, it is enough to use [17, Lemma 2.1] and Theorem 1.3.

## 4. The smoothing effect for the harmonic oscillator

**4.1. Some preliminary results.** — We start by establishing two preliminary lemmas.

**Lemma 4.1.** — *Let  $s_1$  and  $s_2$  be two real numbers.*

(i) *If  $\max(s_2, s_1 + s_2) \leq 1$  then there exists a constant  $C > 0$  such that for all  $u \in L^2(\mathbb{R}^d)$ ,*

$$\| [\sqrt{H}^{s_1+s_2}, \langle x \rangle^{-s_1}] u \|_{L^2(\mathbb{R}^d)} \leq C \|u\|_{L^2(\mathbb{R}^d)},$$

(ii) *If  $s_2 \geq -1$  then there exists a constant  $C > 0$  such that for all  $u \in H^{s_1-1}(\mathbb{R}^d)$ ,*

$$\| [\sqrt{-\Delta}^{s_1}, \langle x \rangle^{-s_2}] u \|_{L^2(\mathbb{R}^d)} \leq C \|u\|_{H^{s_1-1}(\mathbb{R}^d)},$$

(iii) If  $s_2 \leq 1$  then there exists a constant  $C > 0$  such that for all  $u \in \mathcal{H}^{s_1-s_2}(\mathbb{R}^d)$ ,

$$\|[\sqrt{H}^{s_1}, \langle x \rangle^{-s_2}]u\|_{L^2(\mathbb{R}^d)} \leq C\|u\|_{\mathcal{H}^{s_1-s_2}(\mathbb{R}^d)}.$$

*Proof.* — To evaluate the regularity of the previous commutators, we use the Weyl-Hörmander pseudo-differential calculus associated with the metric  $\frac{dx^2}{1+|x|^2} + \frac{d\xi^2}{1+|\xi|^2}$ .

The class of symbols  $S(\mu, m)$  associated to the previous metric is the space of regular functions on  $\mathbb{R}^d \times \mathbb{R}^d$  which satisfy  $|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha, \beta} \langle x \rangle^{\mu-\alpha} \langle \xi \rangle^{m-\beta}$ .

Thus, we have (see [48, Section 18.5], [63] or [4]) that if  $a_1 \in S(\mu_1, m_1)$  and  $a_2 \in S(\mu_2, m_2)$  then the commutator  $[Op(a_1), Op(a_2)]$  is a pseudo-differential operator with a symbol in the class  $S(\mu_1 + \mu_2 - 1, m_2 + m_2 - 1)$ .

Here we will use that  $a(x, \xi) = (|x|^2 + |\xi|^2)^{\alpha/2} \in S(\alpha, \alpha)$ .

(i) As a consequence, if  $\max(s_2, s_1 + s_2) \leq 1$

$$[\sqrt{H}^{s_1+s_2}, \langle x \rangle^{-s_1}] \in S(s_2 - 1, s_1 + s_2 - 1) \subset S(0, s_1 + s_2 - 1).$$

Moreover, as recalled in [53], if  $q \in S(0, \mu)$  then for any  $s \in \mathbb{R}$ , there exists a constant  $C > 0$  such that

$$\|Op(q)u\|_{H^{s-\mu}(\mathbb{R}^d)} \leq C\|u\|_{H^s(\mathbb{R}^d)}.$$

Thus, we can take  $s = \mu = s_1 + s_2 - 1$  to obtain that

$$\|[\sqrt{H}^{s_1+s_2}, \langle x \rangle^{-s_1}]u\|_{L^2(\mathbb{R}^d)} \leq C\|u\|_{H^{s_1+s_2-1}(\mathbb{R}^d)} \leq \|u\|_{L^2(\mathbb{R}^d)}.$$

(ii) Similarly

$$[\sqrt{-\Delta}^{s_1}, \langle x \rangle^{-s_2}] \in S(-s_2 - 1, s_1 - 1) \subset S(0, s_1 - 1),$$

and we can conclude as previously.

(iii) We have

$$[\sqrt{H}^{s_1}, \langle x \rangle^{-s_2}]\sqrt{H}^{s_2-s_1} \in S(-1, s_2 - 1) \subset S(0, s_2 - 1).$$

Then

$$\|[\sqrt{H}^{s_1}, \langle x \rangle^{-s_2}]\sqrt{H}^{s_2-s_1}u\|_{L^2(\mathbb{R}^d)} \leq C\|u\|_{L^2(\mathbb{R}^d)},$$

and it is enough to replace  $u$  by  $\sqrt{H}^{s_1-s_2}u$  to obtain the desired result.  $\square$

**Lemma 4.2.** — *Let  $a \in S(2\epsilon, 0)$ . Then for all  $u \in \mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)$ ,*

$$\left| \int_{\mathbb{R}^d} a(x) \cdot \nabla u(x) \bar{u}(x) dx \right| \leq C\|u\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2.$$

*Proof.* — By duality, we have

$$(4.1) \quad \left| \int_{\mathbb{R}^d} a(x) \cdot \nabla u(x) \bar{u}(x) dx \right| \leq \|au\|_{\mathcal{H}^{1/2-\epsilon}(\mathbb{R}^d)} \|\nabla u\|_{\mathcal{H}^{-1/2+\epsilon}(\mathbb{R}^d)}.$$

We use again the Weyl-Hörmander pseudo-differential calculus introduced in the proof of Lemma 4.1. Since  $a \in S(2\epsilon, 0)$ , we deduce that  $\|au\|_{\mathcal{H}^{1/2-\epsilon}(\mathbb{R}^d)} \leq C\|u\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}$ . Then we observe that  $\|\nabla u\|_{\mathcal{H}^{-1/2+\epsilon}(\mathbb{R}^d)} \leq C\|u\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}$ , which implies the result by (4.1).  $\square$

With these different lemmas established, we can proceed to the proof of the smoothing effect.

#### 4.2. Proof of (1.15). —

**Step 1:** Let us show that for any  $\alpha < 1$ , there exists a constant  $C > 0$  such that for any  $u \in \mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)$

$$(4.2) \quad 2(1-\alpha) \left\| \frac{1}{\langle x \rangle^{\alpha/2}} \nabla u \right\|_{L^2(\mathbb{R}^d)}^2 \leq C \|u\|_{\mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)}^2 - \operatorname{Re} \left( \int_{\mathbb{R}^d} \left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, H \right] u(x) \bar{u}(x) dx \right).$$

By Theorem A.1 with  $h = 1$ , we obtain, since  $\nabla = iOp(\xi)$ ,

$$(4.3) \quad \begin{aligned} \left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, \Delta \right] &= \\ &= \left( \frac{x \cdot \nabla}{\langle x \rangle^\alpha} \right) \Delta - \Delta \left( \frac{x \cdot \nabla}{\langle x \rangle^\alpha} \right) \\ &= iOp \left( \frac{x \cdot \xi}{\langle x \rangle^\alpha} \right) Op(-|\xi|^2) - iOp(-|\xi|^2) Op \left( \frac{x \cdot \xi}{\langle x \rangle^\alpha} \right) \\ &= 2Op \left( \left( \frac{|\xi|^2}{\langle x \rangle^\alpha} - \alpha \frac{(x \cdot \xi)^2}{\langle x \rangle^{\alpha+2}} \right) + i\alpha(d+2) \frac{x \cdot \xi}{\langle x \rangle^{\alpha+2}} - i\alpha(\alpha+2) \frac{(x \cdot \xi)|x|^2}{\langle x \rangle^{\alpha+4}} \right). \end{aligned}$$

Then, using that  $\alpha < 1$ , we obtain

$$\begin{aligned}
& \operatorname{Re} \left( \int_{\mathbb{R}^d} \operatorname{Op} \left( \left( \frac{|\xi|^2}{\langle x \rangle^\alpha} - \alpha \frac{(x \cdot \xi)^2}{\langle x \rangle^{\alpha+2}} \right) u(x) \bar{u}(x) dx \right) \right) = \\
& = \operatorname{Re} \int_{\mathbb{R}^d} \left( \frac{-\Delta u}{\langle x \rangle^\alpha} \bar{u} + \alpha \frac{(x \cdot \nabla)^2 u - (x \cdot \nabla) u}{\langle x \rangle^{\alpha+2}} \bar{u} \right) dx \\
& = \operatorname{Re} \int_{\mathbb{R}^d} \left( \nabla u \cdot \nabla \left( \frac{\bar{u}}{\langle x \rangle^\alpha} \right) - \alpha (x \cdot \nabla u) \operatorname{div} \left( \frac{x \bar{u}}{\langle x \rangle^{\alpha+2}} \right) - \alpha \frac{(x \cdot \nabla u)}{\langle x \rangle^{\alpha+2}} \bar{u} \right) dx \\
& = \int_{\mathbb{R}^d} \left( \frac{|\nabla u|^2}{\langle x \rangle^\alpha} - \alpha \frac{|x \cdot \nabla u|^2}{\langle x \rangle^{\alpha+2}} \right) dx + \\
& \quad + \alpha \operatorname{Re} \int_{\mathbb{R}^d} \left( (\alpha + 2) \frac{|x|^2 (x \cdot \nabla u)}{\langle x \rangle^{\alpha+4}} \bar{u} - (d + 2) \frac{(x \cdot \nabla u)}{\langle x \rangle^{\alpha+2}} \bar{u} \right) dx \\
& \geq (1 - \alpha) \left\| \frac{1}{\langle x \rangle^{\alpha/2}} \nabla u \right\|_{L^2(\mathbb{R}^d)}^2 + \\
& \quad + \alpha \operatorname{Re} \int_{\mathbb{R}^d} \left( (\alpha + 2) \frac{|x|^2 (x \cdot \nabla u)}{\langle x \rangle^{\alpha+4}} \bar{u} - (d + 2) \frac{(x \cdot \nabla u)}{\langle x \rangle^{\alpha+2}} \bar{u} \right) dx.
\end{aligned}$$

Thanks to Lemma 4.2 with  $\epsilon = 0$ , we establish

$$\left| \int_{\mathbb{R}^d} \operatorname{Op} \left( \frac{x \cdot \xi}{\langle x \rangle^{\alpha+2}} \right) u \bar{u} dx \right|, \left| \int_{\mathbb{R}^d} \operatorname{Op} \left( \frac{(x \cdot \xi) |x|^2}{\langle x \rangle^{\alpha+4}} \right) u \bar{u} dx \right| \leq C \|u\|_{\mathcal{H}^{1/2}(\mathbb{R}^d)}^2,$$

and

$$\left| \int_{\mathbb{R}^d} \left( \frac{|x|^2 (x \cdot \nabla u)}{\langle x \rangle^{\alpha+4}} \bar{u} \right) dx \right|, \left| \int_{\mathbb{R}^d} \left( \frac{x \cdot \nabla u}{\langle x \rangle^{\alpha+2}} \bar{u} \right) dx \right| \leq C \|u\|_{\mathcal{H}^{1/2}(\mathbb{R}^d)}^2.$$

Thus by (4.3), for any  $\alpha < 1$ , there exists a constant  $C > 0$  such that for any  $u \in H^{1/2}(\mathbb{R}^d)$ ,

$$\operatorname{Re} \left( \int_{\mathbb{R}^d} \left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, \Delta \right] u(x) \bar{u}(x) dx \right) \geq 2(1 - \alpha) \left\| \frac{1}{\langle x \rangle^{\alpha/2}} \nabla u \right\|_{L^2(\mathbb{R}^d)}^2 - C \|u\|_{\mathcal{H}^{1/2}(\mathbb{R}^d)}^2.$$

In a similar way, we have

$$\left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, -|x|^2 \right] = -2 \operatorname{Op} \left( \frac{|x|^2}{\langle x \rangle^\alpha} \right),$$

and therefore

$$\begin{aligned}
\operatorname{Re} \left( \int_{\mathbb{R}^d} \left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, -|x|^2 \right] u(x) \overline{u(x)} dx \right) &= -2 \int_{\mathbb{R}^d} \frac{|x|^2}{\langle x \rangle^\alpha} |u(x)|^2 dx \\
&\geq -C \|u\|_{\mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)}^2.
\end{aligned}$$

This implies (4.2).

**Step 2:** We now prove that for any  $\alpha \in ]0, 1[$ , there exists a constant  $C > 0$  such that for any  $T \geq 0$  and  $u_0 \in \mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)$ ,

$$(4.4) \quad \int_0^T \left\| \frac{1}{\langle x \rangle^{\alpha/2}} \nabla (e^{-itH} u_0) \right\|_{L^2(\mathbb{R}^d)}^2 dt \leq CT \|u_0\|_{\mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)}^2.$$

Let us set  $u = e^{-itH} u_0$ , then we get

$$\begin{aligned} -i \int_{\mathbb{R}^d} \left[ \frac{x \cdot \nabla}{\langle x \rangle^\alpha}, H \right] u(t, x) \overline{u(t, x)} dx &= \\ &= \int_{\mathbb{R}^d} \frac{x \cdot \nabla \partial_t u(t, x)}{\langle x \rangle^\alpha} \overline{u(t, x)} dx + i \int_{\mathbb{R}^d} \frac{x \cdot \nabla}{\langle x \rangle^\alpha} u(t, x) \overline{H u(t, x)} dx \\ &= \int_{\mathbb{R}^d} \frac{x \cdot \nabla \partial_t u(t, x)}{\langle x \rangle^\alpha} \overline{u(t, x)} dx + \int_{\mathbb{R}^d} \frac{x \cdot \nabla}{\langle x \rangle^\alpha} u(t, x) \overline{\partial_t u(t, x)} dx \\ &= \partial_t \left( \int_{\mathbb{R}^d} \frac{x \cdot \nabla}{\langle x \rangle^\alpha} u(t, x) \overline{u(t, x)} dx \right). \end{aligned}$$

Thus, thanks to (4.2), we obtain for  $T \geq 0$ ,

$$\begin{aligned} 2(1-\alpha) \int_0^T \left\| \frac{1}{\langle x \rangle^{\alpha/2}} \nabla (e^{-itH} u_0) \right\|_{L^2(\mathbb{R}^d)}^2 dt &\leq \\ &\leq CT \|u_0\|_{\mathcal{H}^{(2-\alpha)/2}(\mathbb{R}^d)}^2 + \operatorname{Re} \left( i \int_{\mathbb{R}^d} \frac{x \cdot \nabla u_0}{\langle x \rangle^\alpha} \overline{u_0} - \frac{x \cdot \nabla (e^{-iTH} u_0)}{\langle x \rangle^\alpha} \overline{e^{-iTH} u_0} dx \right). \end{aligned}$$

Then, we can apply Lemma 4.2 with  $\epsilon = 1 - \alpha/2$ , to deduce (4.4).

**Step 3 :** In (4.4), we choose  $\alpha = 1 - 2\epsilon$  with  $\epsilon \in ]0, \frac{1}{2}[$  so that we have

$$(4.5) \quad \int_0^T \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \nabla (e^{-itH} u_0) \right\|_{L^2(\mathbb{R}^d)}^2 dt \leq CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2.$$

Using Lemma 4.1, we get

$$\begin{aligned} &\int_0^T \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} H^{1/2} e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt \leq \\ &\leq 2 \int_0^T \left\| H^{1/2} \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt + 2 \int_0^T \left\| \left[ \frac{1}{\langle x \rangle^{1/2-\epsilon}}, H^{1/2} \right] e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt \\ &\leq 2 \int_0^T \left\| H^{1/2} \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt + CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2. \end{aligned}$$

Then, using (1.2) and (4.5), we obtain

$$\begin{aligned}
& \int_0^T \left\| H^{1/2} \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 \leq \\
& \leq C \int_0^T \left\| \langle x \rangle^{1/2+\epsilon} e^{itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt + C \int_0^T \left\| (-\Delta)^{1/2} \left( \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right) \right\|_{L^2(\mathbb{R}^d)}^2 dt \\
& \leq CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2 + C \int_0^T \left\| \nabla \left( \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right) \right\|_{L^2(\mathbb{R}^d)}^2 dt \\
& \leq CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2 + C \int_0^T \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \nabla (e^{itH} u_0) \right\|_{L^2(\mathbb{R}^d)}^2 dt \\
& \leq CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2.
\end{aligned}$$

Therefore, we obtain

$$\int_0^T \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} H^{1/2} e^{-itH} u_0 \right\|_{L^2(\mathbb{R}^d)}^2 dt \leq CT \|u_0\|_{\mathcal{H}^{1/2+\epsilon}(\mathbb{R}^d)}^2.$$

And we can replace  $u_0$  by  $H^{-1/4-\epsilon/2} u_0$  to deduce (1.15).

**4.3. Proof of (1.16).** — Using Lemma 4.1 and (1.15), we get

$$\begin{aligned}
& \left\| \sqrt{H}^{d/2-2\epsilon} \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq \\
& \leq \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{H}^{d/2-2\epsilon} e^{itH} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} + \\
& \quad + \left\| \left[ \sqrt{H}^{d/2-2\epsilon}, \frac{1}{\langle x \rangle^{1/2-\epsilon}} \right] e^{itH} u_0 \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \\
& \leq C \|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}.
\end{aligned}$$

Then, using (1.2), we establish

$$\left\| \sqrt{-\Delta}^{d/2-2\epsilon} \left( \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right) \right\|_{L^2([-\pi, \pi] \times \mathbb{R}^d)} \leq C \|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}.$$

And finally, using Lemma 4.1, we can conclude that

$$\begin{aligned}
& \left\| \frac{1}{\langle x \rangle^{1/2-\epsilon}} \sqrt{-\Delta}^{d/2-2\epsilon} (e^{itH} u_0) \right\|_{L^2([- \pi, \pi] \times \mathbb{R}^d)} \leq \\
& \leq \left\| \left[ \frac{1}{\langle x \rangle^{1/2-\epsilon}}, \sqrt{-\Delta}^{d/2-2\epsilon} \right] e^{itH} u_0 \right\|_{L^2([- \pi, \pi] \times \mathbb{R}^d)} + \\
& \quad + \left\| \sqrt{-\Delta}^{d/2-2\epsilon} \left( \frac{1}{\langle x \rangle^{1/2-\epsilon}} e^{itH} u_0 \right) \right\|_{L^2([- \pi, \pi] \times \mathbb{R}^d)} \\
& \leq C \|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}.
\end{aligned}$$

### 5. Random initial data and Sobolev spaces

This section is devoted to the proof of Theorem 1.1, which shows that the initial randomized data does not gain any derivative in  $L^2(\mathbb{R}^d)$  and does not enjoy any better localisation property.

The proof of this result will rely on micro-analysis tools. We first step is to establish the following statement which gives a precise description of the phase-space localisation of the Hermite functions.

**Proposition 5.1.** — *For all  $s \geq 0$ , there exist two constants  $C_1, C_2 > 0$  such that for all  $n \in \mathbb{N}$ ,*

$$(5.1) \quad C_1 \lambda_n^s \leq \|(-\Delta)^{s/2} h_n\|_{L^2(\mathbb{R}^d)} \leq C_2 \lambda_n^s,$$

and

$$(5.2) \quad C_1 \lambda_n^s \leq \|\langle x \rangle^{s/2} h_n\|_{L^2(\mathbb{R}^d)} \leq C_2 \lambda_n^s.$$

*Proof.* — Let us first prove this result in the particular case where  $(h_n)_{n \geq 0}$  is the basis of the tensor eigenfunctions, since the argument is then particularly easy. Using (1.2), we have

$$\begin{aligned}
c(\|(-\Delta)^{s/2} h_n\|_{L^2(\mathbb{R}^d)} + \|\langle x \rangle^s h_n\|_{L^2(\mathbb{R}^d)}) & \leq \lambda_n^s = \|h_n\|_{\mathcal{H}^s(\mathbb{R}^d)} \leq \\
& \leq C(\|(-\Delta)^{s/2} h_n\|_{L^2(\mathbb{R}^d)} + \|\langle x \rangle^s h_n\|_{L^2(\mathbb{R}^d)}),
\end{aligned}$$

then

$$\begin{aligned}
C' \|(-\Delta)^{s/2} h_n\|_{L^2(\mathbb{R}^d)} & \leq \lambda_n^s \leq \\
& \leq C(\|(-\Delta)^{s/2} h_n\|_{L^2(\mathbb{R}^d)} + \|(-\Delta)^{s/2} (\widehat{h}_n)\|_{L^2(\mathbb{R}^d)} + \|h_n\|_{L^2(\mathbb{R}^d)}).
\end{aligned}$$

But since the eigenfunctions are tensor functions, then  $h_n(x) = e^{i\theta_n} \widehat{h}_n(x)$ , for some  $\theta_n \in \mathbb{R}$ , since this equality is true in dimension 1, and (5.1) follows. The proof of (5.2) is similar.

Let us now prove (5.1) in the general case. We set  $h = \frac{1}{\lambda_n^2}$  and  $\Phi_h(x) = \frac{1}{h^{d/4}} h_n(\lambda_n x)$  so that  $(-h^2 \Delta + |x|^2 - 1)\Phi_h = 0$  and  $\|\Phi_h\|_{L^2(\mathbb{R}^d)} = 1$ . To prove (5.1), it is sufficient to establish that there exists a constant  $C_1 > 0$  such that for all  $h > 0$ ,

$$h^s \|(-\Delta)^{s/2} \Phi_h\|_{L^2(\mathbb{R}^d)} \geq C_1.$$

Let us proceed by contradiction and suppose that

$$(5.3) \quad \liminf_{h \rightarrow 0} h^s \|(-\Delta)^{s/2} \Phi_h\|_{L^2(\mathbb{R}^d)} = 0.$$

By [14, Theorem 2], there exists a positive measure  $\mu \in \mathcal{M}_+(\mathbb{R}^d \times \mathbb{R}^d)$  such that for any function  $a \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ ,

$$\lim_{h \rightarrow 0} \langle a(x, h|\nabla|) \Phi_h, \Phi_h \rangle_{L^2(\mathbb{R}^d) \times L^2(\mathbb{R}^d)} = \int_{\mathbb{R}^d \times \mathbb{R}^d} \text{tr}(a(x, \xi)) \mu(dx d\xi).$$

Let us recall that  $(x, \xi) \in \text{Supp}(\mu)^c$  if and only if there exists  $r > 0$  such that for all  $\phi \in C_0^\infty(B(x, r) \times B(\xi, r))$ ,

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \phi(x, \xi) \mu(dx, d\xi) = 0.$$

By Proposition A.3, if  $a \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  with  $\text{Supp}(a) \cap \{(x, \xi) : |x|^2 + |\xi|^2 = 1\} = \emptyset$  then for all  $N \in \mathbb{N}$ , there exists  $E_N \in \text{Op}(T^{-2})$  and  $R_N \in \text{Op}(T^{-(N+1)})$  such that

$$E_N \circ (-h^2 \Delta + |x|^2 - 1) = a(x, h|\nabla|) - h^{N+1} R_N.$$

Therefore

$$\langle a(x, h|\nabla|) \Phi_h, \Phi_h \rangle_{L^2(\mathbb{R}^d) \times L^2(\mathbb{R}^d)} = h^{N+1} \langle R_N \Phi_h, \Phi_h \rangle_{L^2(\mathbb{R}^d) \times L^2(\mathbb{R}^d)},$$

then

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} a(x, \xi) \mu(dx d\xi) = 0.$$

And finally, we establish that

$$\text{Supp}(\mu) \subset \{(x, \xi) : |x|^2 + |\xi|^2 = 1\}.$$

Again, by Proposition A.3, if  $a \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  with  $\text{Supp}(a) \cap \{(x, \xi) : \xi^2 = 0\} = \emptyset$  then for all  $N \in \mathbb{N}$ , there exists  $E_N \in \text{Op}(S^{-s})$  and  $R_N \in \text{Op}(S^{-(N+1)})$  such that

$$E_N \circ \sum_{j=1}^d |h \partial_{x_j}|^s = a(x, h|\nabla|) - h^{N+1} R_N.$$



But according to [53] and (5.3), we get

$$\begin{aligned} \lim_{h \rightarrow 0} \left| \langle E_N \circ \sum_{j=1}^d |h \partial_{x_j}|^s \Phi_h, \Phi_h \rangle_{L^2(\mathbb{R}^d) \times L^2(\mathbb{R}^d)} \right| &\leq \lim_{h \rightarrow 0} \left\| E_N \circ \sum_{j=1}^d |h \partial_{x_j}|^s \Phi_h \right\|_{L^2(\mathbb{R}^d)} \\ &\leq \lim_{h \rightarrow 0} \left\| \sum_{j=1}^d |h \partial_{x_j}|^s \Phi_h \right\|_{L^2(\mathbb{R}^d)} = 0. \end{aligned}$$

Therefore

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} a(x, \xi) \mu(dx d\xi) = 0,$$

and we establish that

$$\text{Supp}(\mu) \subset \{(x, \xi) : |\xi|^2 = 0\}.$$

Then for  $a \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  we have

$$\begin{aligned} 0 &= \int_{\mathbb{R}^d} [-h^2 \Delta + |x|^2 - 1, h^{-1} Op_h(a)] \Phi_h \overline{\Phi_h} \\ &= \frac{1}{i} \int_{\mathbb{R}^d} \{-h^2 \Delta + |x|^2 - 1, Op_h(a)\} \Phi_h \overline{\Phi_h} + h \int_{\mathbb{R}^d} Op_h(R) \Phi_h \overline{\Phi_h}. \end{aligned}$$

Thus, we deduce that for any function  $a \in C_0^\infty(\mathbb{R}^d)$ ,

$$(5.4) \quad \int_{\mathbb{R}^d \times \mathbb{R}^d} (\xi \partial_x a - x \partial_\xi a) d\mu(x, \xi) = 0.$$

Let  $(x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d$  and set for  $t \in \mathbb{R}$ ,

$$\begin{cases} x(t) = x \cos(t) + \xi \sin(t), \\ \xi(t) = \xi \cos(t) - x \sin(t), \end{cases}$$

*i.e.*

$$\begin{cases} \dot{x}(t) = \xi(t) \text{ with } x(0) = x, \\ \dot{\xi}(t) = -x(t) \text{ with } \xi(0) = \xi. \end{cases}$$

Using (5.4), we get that for all  $t \in \mathbb{R}$  and  $a \in C_0^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ ,

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} a(x \cos(t) + \xi \sin(t), \xi \cos(t) - x \sin(t)) d\mu(x, \xi) = \int_{\mathbb{R}^d \times \mathbb{R}^d} a(x, \xi) d\mu(x, \xi).$$

Therefore if  $(x_0, \xi_0) \in \text{Supp}(\mu)$  then for all  $r > 0$ , there exists  $a \in C_0^\infty(B((x_0, \xi_0), r))$  such that for all  $t \in \mathbb{R}$ ,

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} a(x \cos(t) + \xi \sin(t), \xi \cos(t) - x \sin(t)) d\mu(x, \xi) \neq 0.$$

However

$$\begin{aligned} & \text{Supp}(a(x \cos(t) + \xi \sin(t), \xi \cos(t) - x \sin(t))) \\ & \subset \left\{ (x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d / (x \cos(t) + \xi \sin(t), \xi \cos(t) - x \sin(t)) \in B((x_0, \xi_0), r) \right\} \\ & \subset B(x_0 \cos(t) - \xi_0 \sin(t), 2r) \times B(x_0 \sin(t) + \xi_0 \cos(t), 2r), \end{aligned}$$

and therefore, for all  $t \in \mathbb{R}$ ,  $(x_0 \cos(t) - \xi_0 \sin(t), x_0 \sin(t) + \xi_0 \cos(t)) \in \text{Supp}(\mu)$ .

But for  $\xi_0 = 0$ ,  $x_0^2 = 1$  then  $x_0 \sin(t) + \xi_0 \cos(t) = x_0 \sin(t) = 0$  is impossible and therefore the estimate (5.1) is proved by contradiction. The proof of (5.2) is analogous.  $\square$

Recall the Paley-Zygmund inequality: Let  $X$  be a random variable in  $L^2(\Omega)$ , then for all  $0 \leq \lambda \leq 1$ ,

$$(5.5) \quad P(X \geq \lambda E(X)) \geq (1 - \lambda)^2 \frac{E(X)^2}{E(X^2)}.$$

For a function  $\chi \in C_0^\infty(\mathbb{R}^d)$  such that  $0 \leq \chi \leq 1$ , with  $\chi(x) = 1$  if  $|x| \leq 1$  and  $\chi(x) = 0$  if  $|x| \geq 2$ , we define

$$\begin{aligned} \sigma_N^2 &= \sum_{n \in \mathbb{N}} \chi^2 \left( \frac{\lambda_n^2}{N^2} \right) |c_n|^2 \lambda_n^{2s} \xrightarrow{N \rightarrow +\infty} +\infty, \\ S_N &= \left\| \chi \left( \frac{H}{N^2} \right) u_0 \right\|_{H^s(\mathbb{R}^d)}, \\ M &= \sup_{N \in \mathbb{N}^*} S_N. \end{aligned}$$

Let us turn to the proof of Theorem 1.1.

**Lemma 5.2.** — *For all  $s \geq 0$ , we have*

$$P\left(\omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) u_0^\omega \right\|_{H^s(\mathbb{R}^d)} = +\infty\right) \in \{0, 1\}.$$

*Proof.* — The random initial condition can be written  $u_0^\omega = \sum_{j \geq 0} X_j(\omega)$ , where

$(X_j)_{j \in \mathbb{N}}$  is a sequence of independent random variables satisfying  $X_j(\omega) \in H^s(\mathbb{R}^d)$ ,  $\omega$ -almost surely. Therefore, we have

$$\sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) u_0^\omega \right\|_{H^s(\mathbb{R}^d)} = +\infty$$

if and only if for all  $K \in \mathbb{N}$ ,

$$\sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) \left( \sum_{j \geq K} X_j(\omega) \right) \right\|_{H^s(\mathbb{R}^d)} = +\infty,$$

so if we put  $F_j = \sigma(X_j, X_{j+1}, \dots)$  we have that

$$\left\{ \omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) u_0^\omega \right\|_{H^s(\mathbb{R}^d)} = +\infty \right\} \in \bigcap_{K \in \mathbb{N}^*} F_K.$$

As a consequence, the set  $\left\{ \omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) u_0^\omega \right\|_{H^s(\mathbb{R}^d)} = +\infty \right\}$  is an element of the asymptotic  $\sigma$ -algebra and the lemma is proved by the 0-1 law.  $\square$

**Lemma 5.3.** — For  $s \geq 0$ , if  $\sum_{n \in \mathbb{N}} |c_n|^2 \lambda_n^{2s} = +\infty$  then

$$P \left( \omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\| \chi \left( \frac{H}{N^2} \right) u_0^\omega \right\|_{H^s(\mathbb{R}^d)} = +\infty \right) = 1.$$

*Proof.* — Recall the notation  $M = \sup_{N \in \mathbb{N}^*} S_N$ . Thus by Lemma 5.2 it is sufficient to establish that

$$(5.6) \quad P(M = +\infty) > 0.$$

Thanks to (5.1), we get

$$\begin{aligned} E(S_N^2) &= E \left( \left\| \chi \left( \frac{H}{N^2} \right) u_0 \right\|_{H^s(\mathbb{R}^d)}^2 \right) \\ &\geq E \left( \sum_{n,m} \chi \left( \frac{\lambda_n^2}{N^2} \right) \chi \left( \frac{\lambda_m^2}{N^2} \right) c_n \overline{c_m} g_n(\omega) \overline{g_m(\omega)} \int_{\mathbb{R}^d} (-\Delta)^s(h_n) (-\Delta)^s(h_m) dx \right) \\ &= E \left( \sum_{n \in \mathbb{N}} \chi^2 \left( \frac{\lambda_n^2}{N^2} \right) |c_n|^2 |g_n(\omega)|^2 \|(-\Delta)^s(h_n)\|_{L^2(\mathbb{R}^d)}^2 \right) \\ &\geq C_1 \sigma_N^2. \end{aligned}$$

Therefore, thanks to the Zygmund inequality (5.5) with  $X = S_N^2$ , we establish

$$(5.7) \quad \begin{aligned} P \left( M^2 \geq \frac{C_1 \sigma_N^2}{2} \right) &\geq P \left( S_N^2 \geq \frac{C_1 \sigma_N^2}{2} \right) \geq P \left( S_N^2 \geq \frac{E(S_N^2)}{2} \right) \\ &\geq \frac{1}{4} \frac{E \left( \left\| \chi \left( \frac{H}{N^2} \right) u_0 \right\|_{H^s(\mathbb{R}^d)}^2 \right)^2}{E \left( \left\| \chi \left( \frac{H}{N^2} \right) u_0 \right\|_{H^s(\mathbb{R}^d)}^4 \right)}. \end{aligned}$$

Then, thanks to (5.1), we have

$$\begin{aligned}
 & E\left(\left\|\chi\left(\frac{H}{N^2}\right)u_0\right\|_{H^s(\mathbb{R}^d)}^4\right) \leq \\
 & \leq E\left(\sum_{n,m} \chi\left(\frac{\lambda_n^2}{N^2}\right)\chi\left(\frac{\lambda_m^2}{N^2}\right)c_n\overline{c_m} g_n(\omega)\overline{g_m(\omega)} \int_{\mathbb{R}^d} (-\Delta)^s(h_n)(-\Delta)^s(h_m)dx\right)^2 \\
 & \quad + E\left(\sum_{n,m} \chi\left(\frac{\lambda_n^2}{N^2}\right)\chi\left(\frac{\lambda_m^2}{N^2}\right)c_n\overline{c_m} g_n(\omega)\overline{g_m(\omega)}\right)^2 \\
 & \leq CE\left(\sum_{n_1,n_2,n_3,n_4} \chi\left(\frac{\lambda_{n_1}^2}{N^2}\right)\chi\left(\frac{\lambda_{n_2}^2}{N^2}\right)\chi\left(\frac{\lambda_{n_3}^2}{N^2}\right)\chi\left(\frac{\lambda_{n_4}^2}{N^2}\right)c_{n_1}\overline{c_{n_2}}c_{n_3}\overline{c_{n_4}}\right. \\
 & \quad \cdot \|(-\Delta)^s(h_{n_1})\|_{L^2(\mathbb{R}^d)}\|(-\Delta)^s(h_{n_2})\|_{L^2(\mathbb{R}^d)}\|(-\Delta)^s(h_{n_3})\|_{L^2(\mathbb{R}^d)}\|(-\Delta)^s(h_{n_4})\|_{L^2(\mathbb{R}^d)}\Big) \\
 & \quad + CE\left(\sum_n \chi^2\left(\frac{\lambda_n^2}{N^2}\right)|c_n|^2\right)^2 \\
 & \leq C_2\sigma_N^4.
 \end{aligned}$$

Therefore from (5.7) we deduce  $P\left(M^2 \geq \frac{C_1\sigma_N^2}{2}\right) \geq \frac{1}{4}\frac{C_1^2}{C_2}$ . Finally, using the monotone convergence theorem, we obtain  $P(M = +\infty) \geq \frac{1}{4}\frac{C_1^2}{C_2}$ , which implies (5.6).  $\square$

We are now able to complete the proof of Theorem 1.1. Indeed, let us prove it by contradiction and assume that  $P(\omega \in \Omega : u_0^\omega \in H^s(\mathbb{R}^d)) > 0$ . According to Proposition A.6,

$$\forall u \in H^s(\mathbb{R}^d), \quad \sup_{N \in \mathbb{N}^*} \left\|\chi\left(\frac{H}{N^2}\right)u\right\|_{H^s(\mathbb{R}^d)} \leq C\|u\|_{H^s(\mathbb{R}^d)},$$

and from the latter inequality we deduce that

$$P\left(\omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\|\chi\left(\frac{H}{N^2}\right)u_0^\omega\right\|_{H^s(\mathbb{R}^d)} < +\infty\right) > 0.$$

Therefore

$$P\left(\omega \in \Omega : \sup_{N \in \mathbb{N}^*} \left\|\chi\left(\frac{H}{N^2}\right)u_0^\omega\right\|_{H^s(\mathbb{R}^d)} = +\infty\right) < 1$$

which contradicts Proposition 5.3.

As a consequence, we have proven that if  $u_0 \notin \mathcal{H}^s(\mathbb{R}^d)$  then  $u_0^\omega \notin H^s(\mathbb{R}^d)$ . The fact that  $\langle x \rangle^s u_0^\omega \notin L^2(\mathbb{R}^d)$  is obtained similarly, and this concludes the proof of Theorem 1.1.

To finish this part, we evaluate the Sobolev norm of the initial data. This will establish that Theorem 1.2 will hold true for supercritical equations with large initial data.

**Proposition 5.4.** — *Let  $\sigma \geq 0$ ,  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$  and  $s \geq \sigma$ . Assume that for all  $n \in \mathbb{N}$ ,*

$$\lambda_n^{2s} |c_n|^2 \leq 1$$

then for all  $t \geq 0$ ,

$$\mu\left(u : \left\| \chi\left(\frac{H}{N^2}\right)u \right\|_{\mathcal{H}^s(\mathbb{R}^3)} \leq t\right) \leq e^{t^2 - \frac{1}{2}\|\chi\left(\frac{H}{N^2}\right)u_0\|_{\mathcal{H}^s(\mathbb{R}^3)}^2}$$

*Proof.* — Using that  $-\ln(1+u) \leq -\frac{u}{2}$  for all  $u \in [0, 1]$  and the Markov inequality, we obtain

$$\begin{aligned} \mu\left(u \in \mathcal{H}^\sigma(\mathbb{R}^3) : \left\| \chi\left(\frac{H}{N^2}\right)u \right\|_{\mathcal{H}^s(\mathbb{R}^3)} \leq t\right) &= \\ &= P\left(\omega \in \Omega : e^{-\|\chi\left(\frac{H}{N^2}\right)u_0^\omega\|_{\mathcal{H}^s(\mathbb{R}^3)}^2} \geq e^{-t^2}\right) \\ &\leq e^{t^2} E\left(e^{-\|\chi\left(\frac{H}{N^2}\right)u_0^\omega\|_{\mathcal{H}^s(\mathbb{R}^3)}^2}\right) \\ &\leq e^{t^2} \prod_{n \in \mathbb{N}} E\left(e^{-\chi^2\left(\frac{\lambda_n^2}{N^2}\right)\lambda_n^{2s}|c_n|^2|X|^2}\right) \\ &\leq e^{t^2} \prod_{n \in \mathbb{N}} \left(\frac{1}{1 + \chi^2\left(\frac{\lambda_n^2}{N^2}\right)\lambda_n^{2s}|c_n|^2}\right) \\ &\leq e^{t^2} \prod_{n \in \mathbb{N}} \left(e^{-\frac{1}{2}\chi^2\left(\frac{\lambda_n^2}{N^2}\right)\lambda_n^{2s}|c_n|^2}\right) \\ &\leq e^{t^2 - \frac{1}{2}\|\chi\left(\frac{H}{N^2}\right)u_0\|_{\mathcal{H}^s(\mathbb{R}^3)}^2}. \end{aligned}$$

□

*Remarks :* (i) For example, if  $u_0 \notin \mathcal{H}^s(\mathbb{R}^3)$  and  $\lambda_n^{2s}|c_n|^2 \leq 1$ ,  $\forall n \in \mathbb{N}$ , we obtain that for all  $t \geq 0$ ,

$$\lim_{N \rightarrow +\infty} \mu\left(u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) : \left\| \chi\left(\frac{H}{N^2}\right)u_0 \right\|_{\mathcal{H}^s(\mathbb{R}^d)} \leq t\right) = 0.$$

This does mean that the Sobolev norm of the initial data is not small.

(ii) For example, for  $\epsilon \ll 1$ , we can choose  $c_n = \frac{\epsilon}{\lambda_n^s} \frac{1}{\sqrt{n}}$  and obtain for  $t \geq 0$ ,

$$\mu\left(u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) : \left\| \chi\left(\frac{H}{N^2}\right)u_0 \right\|_{\mathcal{H}^s(\mathbb{R}^d)} \leq t\right) \leq \exp\left(t^2 - C'\epsilon^2 \ln^2 N\right) \xrightarrow{N \rightarrow +\infty} 0,$$

but  $\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} = C''\epsilon \ll 1$ .

## 6. The fixed point argument in dimension $d = 3$ , global existence, and scattering

Let us introduce the following equation:

$$(6.1) \quad \begin{cases} i \frac{\partial u}{\partial t} - Hu = \kappa \cos(2t)|u|^2u, & (t, x) \in \mathbb{R} \times \mathbb{R}^3 \\ u(0, x) = u_0(x). \end{cases}$$

We will show that this equation is locally well posed. Then, thanks to the lens transform we will be able to show that (1.3) is globally well-posed.

**6.1. Some nonlinear analysis.** — In this part, we establish the estimates which will be used to apply a fixed point theorem.

**Lemma 6.1.** — *There exists  $b' < \frac{1}{2}$  such that for all  $\delta > 0$  and  $K \geq 1$ , there exists a constant  $C_K > 0$  such that if  $N_1 \geq N_2^{1+\delta}$  and  $N_2 \geq N_3 \geq N_4$  then for all  $u_1, u_2, u_3, u_4 \in \bar{X}^{0, b'}$ ,*

$$\left| \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(u_1) \Delta_{N_2}(u_2) \Delta_{N_3}(u_3) \Delta_{N_4}(u_4) \right| \leq C_K N_1^{-K} \prod_{j=1}^4 \|\Delta_{N_j}(u_j)\|_{\bar{X}^{0, b'}}.$$

*Proof.* — We begin with the case  $u_j(t, x) = c_j(t)h_{n_j}(x)$ . We have

$$\begin{aligned} \left| \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(u_1) \Delta_{N_2}(u_2) \Delta_{N_3}(u_3) \Delta_{N_4}(u_4) \right| &= \\ &= \left| \int_{\mathbb{R} \times \mathbb{R}^3} \prod_{j=1}^4 \psi\left(\frac{\lambda_{n_j}^2}{N_j^2}\right) c_j(t) h_{n_j}(x) dt dx \right| \\ &\leq \prod_{j=1}^4 \psi\left(\frac{\lambda_{n_j}^2}{N_j^2}\right) \int_{\mathbb{R}} |c_1(t) \cdots c_4(t)| dt \left| \int_{\mathbb{R}^3} h_{n_1}(x) \cdots h_{n_4}(x) dx \right| \end{aligned}$$

Then by Lemma 2.13 and Proposition 2.4, we deduce

$$\begin{aligned}
(6.2) \quad \left| \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(u_1) \Delta_{N_2}(u_2) \Delta_{N_3}(u_3) \Delta_{N_4}(u_4) \right| &\leq \\
&\leq C_K N_1^{-K} \prod_{j=1}^4 \psi\left(\frac{\lambda_{n_j}^2}{N_j^2}\right) \prod_{j=1}^4 \|c_j\|_{L_t^4(\mathbb{R})} \\
&\leq C_K N_1^{-K} \prod_{j=1}^4 \|\Delta_{N_j}(u_j)\|_{L^4(\mathbb{R}; L^2(\mathbb{R}^3))} \\
&\leq C_K N_1^{-K} \prod_{j=1}^4 \|\Delta_{N_j}(u_j)\|_{\overline{X}^{0, b'}}.
\end{aligned}$$

For the general case, let  $u_j(t, x) = \sum_{k \geq 0} c_{j,k}(t) h_k(x)$ , then by (6.2)

$$\begin{aligned}
&\left| \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(u_1) \Delta_{N_2}(u_2) \Delta_{N_3}(u_3) \Delta_{N_4}(u_4) \right| \leq \\
&\leq \sum_{k_1, k_2, k_3, k_4 \geq 0} \left| \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(c_{1, k_1} h_{k_1}) \Delta_{N_2}(c_{2, k_2} h_{k_2}) \Delta_{N_3}(c_{3, k_3} h_{k_3}) \Delta_{N_4}(c_{4, k_4} h_{k_4}) \right| \\
&\leq C_K N_1^{-K} \sum_{k_1, k_2, k_3, k_4 \geq 0} \prod_{j=1}^4 \|\Delta_{N_j}(c_{j, k_j} h_{k_j})\|_{\overline{X}^{0, b'}} \\
&\leq C_K N_1^{-K+12} \sqrt{\sum_{k_1, k_2, k_3, k_4 \geq 0} \prod_{j=1}^4 \|\Delta_{N_j}(c_{j, k_j} h_{k_j})\|_{\overline{X}^{0, b'}}^2}.
\end{aligned}$$

Then we have

$$\begin{aligned}
\sum_{k_j \geq 0} \|\Delta_{N_j}(c_{j, k_j} h_{k_j})\|_{\overline{X}^{0, b'}}^2 &= \sum_{k_j \geq 0} \sum_{n \geq 0} \|\langle \tau + \lambda_n \rangle^{b'} P_n(\widehat{c_{j, k_j} h_{k_j}})\|_{L_t^2(\mathbb{R}; L_x^2(\mathbb{R}^3))}^2 \\
&= \sum_{n \geq 0} \|\langle \tau + \lambda_n \rangle^{b'} \widehat{P_n}(u_j)\|_{L_t^2(\mathbb{R}; L_x^2(\mathbb{R}^3))}^2 \\
&= \|u_j\|_{\overline{X}^{0, b'}}^2,
\end{aligned}$$

which concludes the proof.  $\square$

**Proposition 6.2.** — *There exists  $b' < \frac{1}{2}$  such that for all  $b > \frac{1}{2}$  and  $s > \frac{1}{2}$ , there exist two constants  $C > 0$  and  $\kappa > 0$  such that for any  $v \in \overline{X}^{s, b}$  and all*

$$N_3 \leq N_2 \leq N_1,$$

$$(6.3) \quad \|\Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(v)\|_{\overline{X}^{s,-b'}} \leq CN_1^{-\kappa}\|v\|_{\overline{X}^{s,b}}^3.$$

*Proof.* — By duality, it is sufficient to show that there exists a constant  $\delta > 0$  such that

$$\int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(v)\Delta_M(w) \leq CN_1^{-\kappa}M^{-\delta}\|v\|_{\overline{X}^{s,b}}^3\|w\|_{\overline{X}^{-s,b'}}.$$

Thanks to Lemma 6.1, we only need to treat the case where  $M \leq N_1^{1+\delta}$ .

- Case  $N_3 \leq M \leq N_1^{1+\delta}$ . Using (3.12), we obtain

$$\begin{aligned} & \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(v)\Delta_M(w) \leq \\ & \leq \|\Delta_{N_1}(v)\Delta_{N_2}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \|\Delta_M(w)\Delta_{N_3}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \\ & \leq (N_2N_3)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{N_3}{M}\right)^{1/2-\delta} \|\Delta_{N_1}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{0,b}} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\ & \leq \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{N_3}{M}\right)^{1/2-\delta} (N_2N_3)^{1/2+\delta-s} \left(\frac{M}{N_1}\right)^s \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}} \\ & \leq (N_2N_3)^{1-s} M^{-1/2+\delta} N_1^{-1/2+(1+s)\delta} \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}} \\ & \leq M^{-\delta} M^{1/2-s+(1+s)\delta} N_1^{1/2-s+2\delta} \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}} \\ & \leq M^{-\delta} N_1^{1-2s+(3+s)\delta} \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}}. \end{aligned}$$

- Case  $M \leq N_3 \leq N_1^{1+\delta}$ . Using (3.12), we get

$$\begin{aligned} & \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(v)\Delta_M(w) \leq \\ & \leq \|\Delta_{N_1}(v)\Delta_{N_2}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \|\Delta_M(w)\Delta_{N_3}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \\ & \leq (N_2M)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{M}{N_3}\right)^{1/2-\delta} \|\Delta_{N_1}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{0,b}} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\ & \leq N_2M \left(\frac{1}{N_3N_1}\right)^{1/2-\delta} \left(\frac{M}{N_1N_2N_3}\right)^s \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}} \\ & \leq M^{-\delta} N_2^{1-s} N_3^{1/2+2\delta} \left(\frac{1}{N_1}\right)^{1/2+s-\delta} \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}} \\ & \leq M^{-\delta} N_1^{1-2s+3\delta} \|v\|_{\overline{X}^{s,b}}^3 \|w\|_{\overline{X}^{-s,b'}}, \end{aligned}$$

which concludes the proof.  $\square$



**Proposition 6.3.** — *There exists  $b' < \frac{1}{2}$  such that for all  $b > \frac{1}{2}$  and  $s > \frac{1}{2}$ , there exist two constants  $C, \kappa > 0$  such that if for some  $\lambda > 0$ , we have for all  $N$ ,*

$$\|\Delta_N(e^{itH}u_0)\|_{L^2([-\pi,\pi];L^\infty(\mathbb{R}^3))} \leq \lambda N^{-1/6}$$

then for all  $v \in \overline{X}^{s,b}$  and all  $N_3 \leq N_2 \leq N_1$ ,

$$(6.4) \quad \|\Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(\psi(t)e^{-itH}u_0)\|_{\overline{X}^{s,-b'}} \leq CN_1^{-\kappa}(\|v\|_{\overline{X}^{s,b}}^3 + \lambda^3),$$

$$(6.5) \quad \|\Delta_{N_1}(v)\Delta_{N_2}(\psi(t)e^{-itH}u_0)\Delta_{N_3}(v)\|_{\overline{X}^{s,-b'}} \leq CN_1^{-\kappa}(\|v\|_{\overline{X}^{s,b}}^3 + \lambda^3).$$

*Proof.* — We show (6.4), the proof of (6.5) being similar. By duality, it is sufficient to show that there exists a constant  $\delta > 0$  such that

$$\begin{aligned} \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(\psi(t)e^{-itH}u_0)\Delta_M(w) &\leq \\ &\leq CN_1^{-\kappa}M^{-\delta}(\|v\|_{\overline{X}^{s,b}}^3 + \lambda^3)\|w\|_{\overline{X}^{-s,b'}}. \end{aligned}$$

Thanks to Lemma 6.1, we only need to treat the case where  $M \leq N_1^{1+\delta}$ .

- Case  $N_2 \leq M \leq N_1^{1+\delta}$ . Using (3.12) and Proposition 2.3, we get

$$\begin{aligned} &\int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v)\Delta_{N_2}(v)\Delta_{N_3}(\psi(t)e^{-itH}u_0)\Delta_M(w) \leq \\ &\leq \|\Delta_{N_2}(v)\Delta_M(w)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \|\Delta_{N_3}(\psi(t)e^{-itH}u_0)\|_{L^2([-\pi,\pi];L^\infty(\mathbb{R}^3))} \|\Delta_{N_1}(v)\|_{L^\infty([-\pi,\pi];L^2(\mathbb{R}^3))} \\ &\leq N_2^{1/2+\delta} \left(\frac{N_2}{M}\right)^{1/2-\delta} \|\Delta_{N_1}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(e^{itH}u_0)\|_{L^2([-\pi,\pi];L^\infty(\mathbb{R}^3))} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\ &\leq N_2^{1/2+\delta} \left(\frac{N_2}{M}\right)^{1/2-\delta} \left(\frac{M}{N_1N_2}\right)^s N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\ &\leq N_2^{1-s} M^{s-1/2+\delta} N_1^{-s} N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\ &\leq M^{-\delta} N_2^{1-s} N_1^{-1/2+(3+s)\delta} N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\ &\leq N_1^{1/2-s+(3+s)\delta} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}}. \end{aligned}$$

- Case  $M \leq N_2 \leq N_1^{1+\delta}$ . Using (3.12) and Proposition 2.3, we get

$$\begin{aligned}
 & \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(v) \Delta_{N_2}(v) \Delta_{N_3}(\psi(t)e^{-itH}u_0) \Delta_M(w) \leq \\
 & \leq \|\Delta_{N_2}(v) \Delta_M(w)\|_{L^2(\mathbb{R}; L^2(\mathbb{R}^3))} \|\Delta_{N_3}(\psi(t)e^{-itH}u_0)\|_{L^2([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \|\Delta_{N_1}(v)\|_{L^\infty([-\pi, \pi]; L^2(\mathbb{R}^3))} \\
 & \leq M^{1/2+\delta} \left(\frac{M}{N_2}\right)^{1/2-\delta} \|\Delta_{N_1}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(e^{itH}u_0)\|_{L^2([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\
 & \leq M^{1/2+\delta} \left(\frac{M}{N_2}\right)^{1/2-\delta} \left(\frac{M}{N_1 N_2}\right)^s N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\
 & \leq N_2^{-1/2-s+\delta} M^{1+s} N_1^{-s} N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\
 & \leq N_2^{1/2+\delta} N_1^{-s} N_3^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\
 & \leq M^{-\delta} N_1^{1/2-s+2\delta} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}},
 \end{aligned}$$

whiv □

**Proposition 6.4.** — *There exists  $b' < \frac{1}{2}$  such that for all  $b > \frac{1}{2}$  and  $s > \frac{1}{2}$ , there exists a constant  $C > 0$  such that if for some  $\lambda > 0$ , we have for all  $N$ ,*

$$\|\Delta_N(e^{itH}u_0)\|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \lambda N^{-1/6} \quad \text{and} \quad \| [e^{itH}u_0]^2 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \lambda^2$$

then for all  $v \in \overline{X}^{s,b}$ ,

$$(6.6) \quad \|v(\psi(t)e^{-itH}u_0)(\psi(t)e^{-itH}u_0)\|_{\overline{X}^{s,-b'}} \leq C(\|v\|_{\overline{X}^{s,b}}^3 + \lambda^3).$$

*Proof.* — Using Proposition 2.5 and Proposition 2.3, we get

$$\begin{aligned}
 & \|v(\psi(t)e^{-itH}u_0)(\psi(t)e^{-itH}u_0)\|_{\overline{X}^{s,-b'}} \leq \\
 & \leq \|v(\psi(t)e^{-itH}u_0)(\psi(t)e^{-itH}u_0)\|_{L^{1+\delta}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^3))} \\
 & \leq \|v(\psi(t)e^{-itH}u_0)(\psi(t)e^{-itH}u_0)\|_{L^{1+\delta}([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \\
 & \leq \|v\|_{L^\infty([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \|e^{itH}u_0\|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))}^2 + \\
 & \quad + \|v\|_{L^2([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \| [e^{itH}u_0]^2 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \\
 & \leq \lambda^2 \|v\|_{\overline{X}^{s,b}} + \lambda^2 \|v\|_{L^2(\mathbb{R}; \mathcal{W}^{s,6}(\mathbb{R}^3))} \\
 & \leq C(\|v\|_{\overline{X}^{s,b}}^3 + \lambda^3),
 \end{aligned}$$

hence the result. □

**Proposition 6.5.** — *There exists  $b' < \frac{1}{2}$  such that for all  $b > \frac{1}{2}$  and  $s > \frac{1}{2}$ , there exists a constant  $C > 0$  such that if for some  $\lambda > 0$ , we have*

$$\| [e^{itH} u_0]^3 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \lambda^3$$

then for all  $v \in \overline{X}^{s,b}$ ,

$$(6.7) \quad \| (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) \|_{\overline{X}^{s,-b'}} \leq C\lambda^3.$$

*Proof.* — Using Proposition 2.5, we get

$$\begin{aligned} \| (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) \|_{\overline{X}^{s,-b'}} &\leq \\ &\leq \| (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) (\psi(t)e^{-itH} u_0) \|_{L^{1+\delta}(\mathbb{R}; \mathcal{H}^s(\mathbb{R}^3))} \\ &\leq C \| [e^{itH} u_0]^3 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \\ &\leq C\lambda^3, \end{aligned}$$

hence the result.  $\square$

**Proposition 6.6.** — *There exists  $b' < \frac{1}{2}$  such that for all  $b > \frac{1}{2}$  and  $s \in ]\frac{1}{2}, 1[$ , there exist  $C > 0, \kappa > 0$  and  $R \in [2, +\infty[$  such that if for some  $\lambda > 0$ , we have for all  $N \geq 1$ ,*

$$\begin{cases} \| u_0 \|_{L^2(\mathbb{R}^3)} \leq \lambda, \\ \| \Delta_N(e^{itH} u_0) \|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \lambda N^{-1/6}, \\ \| \Delta_N(e^{itH} u_0) \|_{L^R([- \pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \lambda N^{s-1/4}, \end{cases}$$

then for all  $v \in \overline{X}^{s,b}$  and all  $N_3 \leq N_2 \leq N_1$ ,

$$(6.8) \quad \| \Delta_{N_1}(\psi(t)e^{-itH} u_0) \Delta_{N_2}(v) \Delta_{N_3}(v) \|_{\overline{X}^{s,-b'}} \leq CN_1^{-\kappa} (\lambda^3 + \|v\|_{\overline{X}^{s,b}}^3).$$

*Proof.* — Let  $\delta > 0$  be small enough, to be fixed later.

**Case**  $N_1 \geq (N_2 N_3)^{\frac{1-s}{1-s-4\delta}}$ . By duality, it is sufficient to establish

$$\int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(\psi(t)e^{itH} u_0) \Delta_{N_2}(v) \Delta_{N_3}(v) \Delta_M(w) \leq CN_1^{-\kappa} M^{-\delta} \|w\|_{\overline{X}^{-s,b'}} (\lambda^3 + \|v\|_{\overline{X}^{s,b}}^3).$$

Thanks to Lemma 6.1, we only need to treat the case where  $M \leq N_1^{1+\delta}$ . If  $N_3 \leq M$  then using (3.12) and (3.13), we obtain

$$\begin{aligned}
& \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\Delta_{N_3}(v)\Delta_M(w) \leq \\
& \leq \|\Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \|\Delta_{N_3}(v)\Delta_M(w)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \\
& \leq (N_2N_3)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{N_3}{M}\right)^{1/2-\delta} \|\Delta_{N_1}(u_0)\|_{L^2} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{0,b}} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\
& \leq (N_2N_3)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{N_3}{M}\right)^{1/2-\delta} \left(\frac{M}{N_2N_3}\right)^s \lambda \|\Delta_{N_2}(v)\|_{\overline{X}^{s,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{s,b}} \|\Delta_M(w)\|_{\overline{X}^{-s,b'}} \\
& \leq M^{-\delta} N_1^{-\delta} N_1^{-1+s+4\delta} (N_2N_3)^{1-s} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\
& \leq M^{-\delta} N_1^{-\delta} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}}.
\end{aligned}$$

Then, if  $N_3 \geq M$ , using (3.12) and (3.13), we obtain

$$\begin{aligned}
& \int_{\mathbb{R} \times \mathbb{R}^3} \Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\Delta_{N_3}(v)\Delta_M(w) \leq \\
& \leq \|\Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \|\Delta_{N_3}(v)\Delta_M(w)\|_{L^2(\mathbb{R};L^2(\mathbb{R}^3))} \\
& \leq (N_2M)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{M}{N_3}\right)^{1/2-\delta} \|\Delta_{N_1}(u_0)\|_{L^2} \|\Delta_{N_2}(v)\|_{\overline{X}^{0,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{0,b}} \|\Delta_M(w)\|_{\overline{X}^{0,b'}} \\
& \leq (N_2M)^{1/2+\delta} \left(\frac{N_2}{N_1}\right)^{1/2-\delta} \left(\frac{M}{N_3}\right)^{1/2-\delta} \left(\frac{M}{N_2N_3}\right)^s \lambda \|\Delta_{N_2}(v)\|_{\overline{X}^{s,b}} \|\Delta_{N_3}(v)\|_{\overline{X}^{s,b}} \|\Delta_M(w)\|_{\overline{X}^{-s,b'}} \\
& \leq M^{-\delta} N_1^{-\delta} N_1^{-1+s+4\delta} (N_2N_3)^{1-s} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}} \\
& \leq M^{-\delta} N_1^{-\delta} \lambda \|v\|_{\overline{X}^{s,b}}^2 \|w\|_{\overline{X}^{-s,b'}}.
\end{aligned}$$

**Case**  $N_1 \leq (N_2 N_3)^{\frac{1-s}{1-s-4\delta}}$ . Using Proposition 2.5 and Proposition 2.3, we establish

$$\begin{aligned}
& \|\Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\Delta_{N_3}(v)\|_{\overline{X}^{s,-b'}} \leq \\
& \leq \|\Delta_{N_1}(\psi(t)e^{-itH}u_0)\Delta_{N_2}(v)\Delta_{N_3}(v)\|_{L^{1+\delta}(\mathbb{R};\mathcal{H}^s(\mathbb{R}^3))} \\
& \leq \|\Delta_{N_1}(e^{-itH}u_0)\Delta_{N_2}(v)\Delta_{N_3}(v)\|_{L^{1+\delta}([- \pi, \pi];\mathcal{H}^s(\mathbb{R}^3))} \\
& \leq \|\Delta_{N_1}(e^{itH}u_0)\|_{L^{\frac{(1+\delta)(1+2\delta)}{\delta}}([- \pi, \pi];\mathcal{W}^{s,4}(\mathbb{R}^3))} \prod_{j=1}^2 \|\Delta_{N_j}(v)\|_{L^{2(1+2\delta)}(\mathbb{R};L^8(\mathbb{R}^3))} \\
& \quad + \|\Delta_{N_1}(e^{itH}u_0)\|_{L^4([- \pi, \pi];L^\infty(\mathbb{R}^3))} \|\Delta_{N_2}(v)\|_{L^\infty([- \pi, \pi];\mathcal{H}^s(\mathbb{R}^3))} \|\Delta_{N_3}(v)\|_{L^2([- \pi, \pi];\mathcal{W}^{s,6}(\mathbb{R}^3))} \\
& \quad + \|\Delta_{N_1}(e^{itH}u_0)\|_{L^4([- \pi, \pi];L^\infty(\mathbb{R}^3))} \|\Delta_{N_2}(v)\|_{L^2([- \pi, \pi];\mathcal{W}^{s,6}(\mathbb{R}^3))} \|\Delta_{N_3}(v)\|_{L^\infty([- \pi, \pi];\mathcal{H}^s(\mathbb{R}^3))} \\
& \leq N_1^{s-1/4} (N_2 N_3)^{\frac{9}{8}-\frac{1}{1+2\delta}-s} \lambda \|v\|_{\overline{X}^{s,b}}^2 + N_1^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2 \\
& \leq (N_2 N_3)^{\frac{(1-s)(s-1/4+\delta)}{1-s-4\delta}} (N_2 N_3)^{\frac{9}{8}-\frac{1}{1+\delta}-s} N_1^{-\delta} \lambda \|v\|_{\overline{X}^{s,b}}^2 + N_1^{-1/6} \lambda \|v\|_{\overline{X}^{s,b}}^2,
\end{aligned}$$

with

$$\begin{aligned}
& \frac{(1-s)(s-1/4+\delta)}{1-s-4\delta} + \frac{9}{8} - \frac{1}{1+2\delta} - s = \\
& = s - \frac{1}{4} + \delta + \frac{4\delta(s-\frac{1}{4}+\delta)}{1-s-4\delta} + \frac{9}{8} - \frac{1}{1+2\delta} - s \\
& = \frac{7}{8} + \frac{4\delta(s-\frac{1}{4}+\delta)}{1-s-4\delta} - \frac{1}{1+2\delta} \\
& = -\frac{1}{8} + o(\delta) < 0.
\end{aligned}$$

And finally, the proposition is proved with  $R = \frac{(1+\delta)(1+2\delta)}{\delta}$ .  $\square$

**6.2. Local well-posedness of equation (6.1).** — Let  $\lambda > 0$  and define  $E_0(\lambda)$  as the set of functions  $u_0 \in L^2(\mathbb{R}^3)$  which satisfy

$$(6.9) \quad \begin{cases} \|u_0\|_{L^2(\mathbb{R}^3)} \leq \lambda \\ \| [e^{itH}u_0]^2 \|_{L^4([- \pi, \pi];\mathcal{H}^s(\mathbb{R}^3))} \leq \lambda^2 \\ \| [e^{itH}u_0]^3 \|_{L^4([- \pi, \pi];\mathcal{H}^s(\mathbb{R}^3))} \leq \lambda^3 \\ \|\Delta_N(e^{itH}u_0)\|_{L^4([- \pi, \pi];L^\infty(\mathbb{R}^3))} \leq \lambda N^{-1/6}, \quad \forall N \geq 1 \\ \|\Delta_N(e^{itH}u_0)\|_{L^R([- \pi, \pi];\mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \lambda N^{s-1/4}, \quad \forall N \geq 1 \end{cases}$$

where  $R > 2$  is fixed by Proposition 6.6.

**Proposition 6.7.** — Let  $\frac{1}{2} < s < 1$  then there exist  $C > 0$  and  $b > 1/2$  such that if  $u_0 \in E_0(\lambda)$  with  $\lambda > 0$ , then for all  $v \in \overline{X}^{s,b}$

$$\begin{aligned} \left\| \psi(t) \int_0^t e^{-i(t-s)H} \psi(s) \cos(2s) |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) ds \right\|_{\overline{X}^{s,b}} &\leq \\ &\leq C(\lambda^3 + \|v\|_{\overline{X}^{s,b}}^3). \end{aligned}$$

*Proof.* — For all  $b > \frac{1}{2}$ , using Proposition 2.6 and Proposition 2.7, we obtain

$$\begin{aligned} \left\| \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) ds \right\|_{\overline{X}^{s,b}} &\leq \\ &\leq C \left\| \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) \right\|_{\overline{X}^{s,b-1}} \\ &\leq C \left\| |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) \right\|_{\overline{X}^{s,b-1}}. \end{aligned}$$

Then using (6.3), (6.4), (6.5), (6.6), (6.7), (6.8), we establish the existence of an integer  $b' < \frac{1}{2}$  such that for all  $u_0 \in E_0(\lambda)$ ,

$$\left\| |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) \right\|_{\overline{X}^{s,-b'}} \leq C(\lambda^3 + \|v\|_{\overline{X}^{s,b}}^3).$$

It is then sufficient to choose  $b = 1 - b' > \frac{1}{2}$  and the proposition is proved.  $\square$

**Proposition 6.8.** — Let  $\frac{1}{2} < s < 1$  then there exists a constant  $C > 0$  and  $b > 1/2$  such that if  $u_0 \in E_0(\lambda)$  with  $\lambda > 0$ , then for all  $v \in \overline{X}_T^{s,b}$

$$\begin{aligned} \left\| \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) ds \right\|_{\overline{X}_T^{s,b}} &\leq \\ &\leq C(\lambda^3 + \|v\|_{\overline{X}_T^{s,b}}^3). \end{aligned}$$

*Proof.* — Let  $w \in \overline{X}^{s,b}$  such that  $w|_{[-T,T]} = v$  then

$$\begin{aligned} \left\| \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v|^2 (\psi(s) e^{-isH} u_0 + v) ds \right\|_{\overline{X}_T^{s,b}} &\leq \\ &\leq \left\| \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + w|^2 (\psi(s) e^{-isH} u_0 + w) ds \right\|_{\overline{X}^{s,b}} \\ &\leq C(\lambda^3 + \|w\|_{\overline{X}^{s,b}}^3) \end{aligned}$$

for all  $w \in \overline{X}^{s,b}$ .  $\square$

In a similar way, one could prove the following result:

**Proposition 6.9.** — *Let  $\frac{1}{2} < s < 1$  then there exists a constant  $C > 0$  and a real  $b > 1/2$  such that if  $u_0 \in E_0(\lambda)$  with  $\lambda > 0$ , then for all  $v_1, v_2 \in \overline{X}_T^{s,b}$*

$$\begin{aligned} & \left\| \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s)\psi(s) |\psi(s)e^{-isH}u_0 + v_1|^2 (\psi(s)e^{-isH}u_0 + v_1) ds \right. \\ & \quad \left. - \psi(t) \int_0^t e^{-i(t-s)H} \cos(2s)\psi(s) |\psi(s)e^{-isH}u_0 + v_2|^2 (\psi(s)e^{-isH}u_0 + v_2) ds \right\|_{\overline{X}_T^{s,b}} \\ & \leq C \|v_1 - v_2\|_{\overline{X}_T^{s,b}} (\lambda^2 + \|v_1\|_{\overline{X}_T^{s,b}}^2 + \|v_2\|_{\overline{X}_T^{s,b}}^2). \end{aligned}$$

**Theorem 6.10.** — *Let  $\frac{1}{2} < s < 1$  then there exists a constant  $C > 0$  and a real  $b > 1/2$  such that if  $u_0 \in E_0(\lambda)$  with  $\lambda < \frac{1}{2\sqrt{C}}$  then there is a unique solution to the equation (6.1) with initial data  $u_0$  in the space  $e^{-itH}u_0 + B_{\overline{X}_T^{s,b}}(0, \frac{1}{2}\sqrt{\frac{1}{C}})$ .*

*Proof.* — We define

$$\begin{aligned} L(v) := & \\ & - i\kappa\psi(t) \int_0^t e^{-i(t-s)H} \cos(2s)\psi(s) |\psi(s)e^{-isH}u_0 + v(s)|^2 (\psi(s)e^{-isH}u_0 + v(s)) ds, \end{aligned}$$

$u = e^{-itH}u_0 + v$  is the unique solution to (6.1) in the space  $e^{-itH}u_0 + B_{\overline{X}_T^{s,b}}(0, R)$  if and only if  $v$  is the unique fixed point of  $L$  in space  $B_{\overline{X}_T^{s,b}}(0, R)$ . According to Proposition 6.8 and Proposition 6.9, there exists a constant  $C > 0$  such that

$$\begin{aligned} \|L(v)\|_{\overline{X}_T^{s,b}} & \leq C(\lambda^3 + \|v\|_{\overline{X}_T^{s,b}}^3), \\ \|L(v_1) - L(v_2)\|_{\overline{X}_T^{s,b}} & \leq C\|v_1 - v_2\|_{\overline{X}_T^{s,b}} (\lambda^2 + \|v_1\|_{\overline{X}_T^{s,b}}^2 + \|v_2\|_{\overline{X}_T^{s,b}}^2). \end{aligned}$$

Thus, if  $\lambda < \frac{1}{2\sqrt{C}}$  then  $L$  is a contraction of the complete space  $B_{\overline{X}_T^{s,b}}(0, \frac{1}{2}\sqrt{\frac{1}{C}})$  and therefore has a unique fixed point.  $\square$

**6.3. Global solutions and scattering for the equation (1.3).** — Thanks to the lens transform and the result of Theorem 6.10, we are now able to establish the existence of global solutions for the equation (1.3). We also prove the uniqueness of the solution and then show that this solution scatters at  $t \rightarrow \pm\infty$ . Recall that the set  $E_0(\lambda)$  is defined in (6.9).

**Theorem 6.11.** — *If  $\frac{1}{2} < s < 1$  then there exist two constants  $C, c > 0$  such that if  $u_0 \in E_0(\lambda)$  with  $\lambda < \frac{1}{2\sqrt{C}}$  then there exists a unique global solution to the equation (1.3) with initial data  $u_0$  in the space  $e^{it\Delta}u_0 + B_{X^s}(0, \frac{c}{2}\sqrt{\frac{1}{C}})$ .*

*Proof.* — Let  $u$  be given by Theorem 6.10 and define

$$U(t, x) = \left( \frac{1}{\sqrt{1+4t^2}} \right)^{3/2} u \left( \frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) e^{\frac{i|x|^2 t}{1+4t^2}}.$$

According to Section 2.1, since  $u$  is a solution of (6.1) on  $[-\pi/4, \pi/4]$  then  $U$  is a global solution of (1.3).

Thus, to obtain the theorem, it suffices to notice that

$$(e^{it\Delta} u_0)(t, x) = \left( \frac{1}{\sqrt{1+4t^2}} \right)^{3/2} (e^{-itH} u_0) \left( \frac{\arctan(2t)}{2}, \frac{x}{\sqrt{1+4t^2}} \right) e^{\frac{i|x|^2 t}{1+4t^2}},$$

and to use Proposition 2.9. □

The existence of the solutions being proved, one with analogous estimates, one can show that the solutions are unique.

We then prove that the constructed solutions scatter for  $t \rightarrow +\infty$  and  $t \rightarrow -\infty$ .

**Theorem 6.12.** — *Let  $U$  be the unique solution of (1.3) constructed in Theorem 6.11, then there exist  $L_+, L_- \in \mathcal{H}^s(\mathbb{R}^3)$  such that*

$$(6.10) \quad \begin{aligned} \lim_{t \rightarrow +\infty} \|U(t) - e^{it\Delta}(u_0 + L_+)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \\ \lim_{t \rightarrow -\infty} \|U(t) - e^{it\Delta}(u_0 + L_-)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \end{aligned}$$

and

$$(6.11) \quad \begin{aligned} \lim_{t \rightarrow +\infty} \|e^{-it\Delta}U(t) - (u_0 + L_+)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0, \\ \lim_{t \rightarrow -\infty} \|e^{-it\Delta}U(t) - (u_0 + L_-)\|_{\mathcal{H}^s(\mathbb{R}^3)} &= 0. \end{aligned}$$

*Proof.* — Let us treat the case  $t \rightarrow +\infty$ . In the following, we set  $T = \frac{\pi}{4}$ . We have shown that

$$-i\psi(t) \int_0^t e^{-i(t-s)H} \cos(2s)\psi(s)|\psi(s)e^{-isH}u_0 + v(s)|^2(\psi(s)e^{-isH}u_0 + v(s))ds \in \overline{X}_T^{s,b}.$$

hence by Lemma 2.8,

$$\begin{aligned} -ie^{-itH} \int_0^t e^{isH} \cos(2s)\psi(s)|\psi(s)e^{-isH}u_0 + v(s)|^2(\psi(s)e^{-isH}u_0 + v(s))ds \\ \in C^0([-T, T]; \mathcal{H}^s(\mathbb{R}^3)). \end{aligned}$$



And therefore, there exists a function  $L \in \mathcal{H}^s(\mathbb{R}^3)$  such that

$$\begin{aligned} & \lim_{t \rightarrow T} \left\| L - i\kappa e^{-itH} \int_0^t e^{isH} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v(s)|^2 (\psi(s) e^{-isH} u_0 + v(s)) ds \right\|_{\mathcal{H}^s(\mathbb{R}^3)} \\ &= \lim_{t \rightarrow T} \left\| e^{itH} L - i\kappa \int_0^t e^{isH} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v(s)|^2 (\psi(s) e^{-isH} u_0 + v(s)) ds \right\|_{\mathcal{H}^s(\mathbb{R}^3)} \\ &= \lim_{t \rightarrow T} \left\| e^{iT H} L - i\kappa \int_0^t e^{isH} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v(s)|^2 (\psi(s) e^{-isH} u_0 + v(s)) ds \right\|_{\mathcal{H}^s(\mathbb{R}^3)} \\ &= 0. \end{aligned}$$

But for  $t \in [-T, T]$ ,

$$u(t) = e^{-itH} u_0 - i\kappa e^{-itH} \int_0^t e^{isH} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v(s)|^2 (\psi(s) e^{-isH} u_0 + v(s)) ds.$$

So, by (2.6), we obtain

$$\begin{aligned} U(t) &= e^{it\Delta} u_0 + \\ &+ e^{it\Delta} \left[ -i\kappa \int_0^{\frac{\arctan(2t)}{2}} e^{isH} \cos(2s) \psi(s) |\psi(s) e^{-isH} u_0 + v(s)|^2 (\psi(s) e^{-isH} u_0 + v(s)) ds \right] \\ &:= e^{it\Delta} u_0 + e^{it\Delta} F(t). \end{aligned}$$

Then

$$\lim_{t \rightarrow +\infty} \|F(t) - L_+\|_{\mathcal{H}^s(\mathbb{R}^3)} = 0,$$

with  $L_+ = e^{iT H} L \in \mathcal{H}^s(\mathbb{R}^3)$ , which proves (6.11). Then we have

$$\begin{aligned} \lim_{t \rightarrow +\infty} \|e^{it\Delta} F(t) - e^{it\Delta} L_+\|_{H^s(\mathbb{R}^3)} &= \lim_{t \rightarrow +\infty} \|F(t) - L_+\|_{H^s(\mathbb{R}^3)} \\ &\leq C \lim_{t \rightarrow +\infty} \|F(t) - L_+\|_{\mathcal{H}^s(\mathbb{R}^3)} = 0, \end{aligned}$$

hence (6.10) follows.  $\square$

## 7. Estimation of the regularity of the random initial data and proof of Theorem 1.2

**7.1. Estimation of the regularity of the random initial data.** — In this section, we estimate the regularity of the random data by proving large deviation type estimates. In particular, we establish that  $u_0^\omega \in \bigcup_{\lambda \geq 1} E_0(\lambda)$ , for almost any  $\omega \in \Omega$

(recall that  $E_0(\lambda)$  is defined in (6.9)).

For  $\Lambda > 0$  we define,

$$\begin{aligned}
 \Omega_\Lambda = & \left\{ \omega \in \Omega : \left\{ \|u_0^\omega\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \Lambda \right\} \cap \left\{ \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \Lambda^2 \right\} \right. \\
 & \cap \left\{ \| [e^{itH} u_0^\omega]^3 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \Lambda^3 \right\} \\
 (7.1) \quad & \cap \bigcap_{N \text{ dyadic}} \left\{ \|\Delta_N(e^{itH} u_0^\omega)\|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \Lambda N^{-1/6} \right\} \\
 & \left. \cap \bigcap_{N \text{ dyadic}} \left\{ \|\Delta_N(e^{itH} u_0^\omega)\|_{L^R([- \pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \Lambda N^{s-1/4} \right\} \right\}
 \end{aligned}$$

and the aim of this part is to prove the following theorem:

**Theorem 7.1.** — *There are two constants  $C, c > 0$  such that for any  $\Lambda \geq 0$  and all  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$ ,*

$$(7.2) \quad P(\Omega_\Lambda^c) \leq C \exp\left(-\frac{c\Lambda^2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}\right).$$

We start by establishing Wiener chaos type inequalities for complex Gaussian random variables.

**Proposition 7.2.** — *Assume that  $g_n \sim \mathcal{N}_\mathbb{C}(0, 1)$  are independent random variables, then there exists a constant  $C > 0$  such that for all  $q \geq 2$  and all sequences  $(c_n)_n \in \ell^2(\mathbb{N})$ ,  $(c_{n,m})_{n,m} \in \ell^2(\mathbb{N} \times \mathbb{N})$  and  $(c_{n,m,k})_{n,m,k} \in \ell^2(\mathbb{N} \times \mathbb{N} \times \mathbb{N})$ ,*

$$(7.3) \quad \left\| \sum_{n \in \mathbb{N}} c_n g_n(\omega) \right\|_{L^q(\Omega)} \leq C q^{\frac{1}{2}} \sqrt{\sum_{n \in \mathbb{N}} |c_n|^2},$$

$$(7.4) \quad \left\| \sum_{n,m \in \mathbb{N}} c_{n,m} g_n(\omega) g_m(\omega) \right\|_{L^q(\Omega)} \leq C q \sqrt{\sum_{n,m \in \mathbb{N}} |c_{n,m}|^2},$$

$$(7.5) \quad \left\| \sum_{n,m,k \in \mathbb{N}} c_{n,m,k} g_n(\omega) g_m(\omega) g_k(\omega) \right\|_{L^q(\Omega)} \leq C q^{\frac{3}{2}} \sqrt{\sum_{n,m,k \in \mathbb{N}} |c_{n,m,k}|^2}.$$

*Proof.* — The bound (7.3) is the Khintchine inequality, and we refer to [21] for a proof.

Since the random variables  $g_n$  are independent complex Gaussians, by [71, Proposition 2.4] (Wiener chaos estimates) there exists a constant  $C > 0$  such that for any  $q \geq 2$ ,

$$(7.6) \quad \left\| \sum_{n,m \in \mathbb{N}} c_{n,m} g_n(\omega) g_m(\omega) \right\|_{L^q(\Omega)} \leq C q \left\| \sum_{n,m \in \mathbb{N}} c_{n,m} g_n(\omega) g_m(\omega) \right\|_{L^2(\Omega)}.$$

Next,

$$\begin{aligned}
& \left\| \sum_{n,m \in \mathbb{N}} c_{n,m} g_n(\omega) g_m(\omega) \right\|_{L^2(\Omega)}^2 = \\
& = \sum_{n,n',m,m' \in \mathbb{N}} c_{n,m} \overline{c_{n',m'}} E \left( g_n(\omega) g_m(\omega) \overline{g_{n'}(\omega) g_{m'}(\omega)} \right) \\
& \leq \sum_{n=n'=m=m' \in \mathbb{N}} || + \sum_{n=n',m=m' \in \mathbb{N}} || + \sum_{n=m,n'=m' \in \mathbb{N}} || + \sum_{n=m',n'=m \in \mathbb{N}} ||.
\end{aligned}$$

Then

$$\begin{aligned}
\sum_{n=n'=m=m' \in \mathbb{N}} \left| c_{n,m} \overline{c_{n',m'}} E \left( g_n(\omega) g_m(\omega) \overline{g_{n'}(\omega) g_{m'}(\omega)} \right) \right| &= \sum_{n \in \mathbb{N}} |c_{n,n}|^2 E(|g_n(\omega)|^4) \\
&\leq \sum_{n,m \in \mathbb{N}} |c_{n,m}|^2,
\end{aligned}$$

and using that  $E(g_n(\omega)^2) = 0$ , we obtain

$$\begin{aligned}
\sum_{n=m,n'=m' \in \mathbb{N}} \left| c_{n,m} \overline{c_{n',m'}} E \left( g_n(\omega) g_m(\omega) \overline{g_{n'}(\omega) g_{m'}(\omega)} \right) \right| &= \\
&= \sum_{n,m \in \mathbb{N}} |c_{n,n}| |c_{m,m}| \left| E \left( g_n(\omega)^2 \overline{g_m(\omega)^2} \right) \right| \\
&= \sum_{n \in \mathbb{N}} |c_{n,n}|^2 E(|g_n(\omega)|^4) \\
&\leq \sum_{n,m \in \mathbb{N}} |c_{n,m}|^2,
\end{aligned}$$

thus (7.4) is proved. We can proceed in the same way to obtain (7.5) since the inequality (7.6) is true for any product of random variables.  $\square$

We now prove Theorem 7.1. By definition (7.1) of the set  $\Omega_\Lambda$  we have

$$\begin{aligned}
 P(\Omega_\Lambda^c) &\leq P(\omega \in \Omega : \|u_0^\omega\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \geq \Lambda) \\
 &\quad + P(\omega \in \Omega : \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \geq \Lambda^2) \\
 &\quad + P(\omega \in \Omega : \| [e^{itH} u_0^\omega]^3 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \geq \Lambda^3) \\
 &\quad + P\left( \bigcup_{N \text{ dyadic}} \left\{ \omega \in \Omega : \|\Delta_N(e^{itH} u_0^\omega)\|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \geq \Lambda N^{-1/6} \right\} \right) \\
 &\quad + P\left( \bigcup_{N \text{ dyadic}} \left\{ \omega \in \Omega : \|\Delta_N(e^{itH} u_0^\omega)\|_{L^R([- \pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \geq \Lambda N^{s-1/4} \right\} \right)
 \end{aligned}$$

so it is enough to establish the bound (7.2) for each of the terms. Let us carry out the proof for the second and the fourth term (for the other terms, the approach is similar).

• **Case**  $\| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \geq \Lambda^2$ . Since  $s \in [\frac{1}{2}, \frac{1}{2} + \sigma[$ , it is enough to prove the following lemma:

**Lemma 7.3.** — *For any  $\epsilon > 0$ , there exist two constants  $C, c > 0$  such that for any  $\Lambda > 0$  and any function  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$ ,*

$$P\left(\omega \in \Omega : \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^{\sigma+1/2-2\epsilon}(\mathbb{R}^3))} \geq \Lambda^2\right) \leq C e^{-\frac{c\Lambda^2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}}.$$

*Proof.* — Note that it is sufficient to prove the estimate for  $\Lambda \geq C\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}$ . Thanks to the Markov and Minkowski inequalities, we obtain for  $q \geq 4$ ,

$$\begin{aligned}
 P\left(\omega \in \Omega : \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^{\sigma+1/2-2\epsilon}(\mathbb{R}^3))} \geq \Lambda^2\right) &= \\
 &= P\left(\omega \in \Omega : \| H^{\sigma/2+1/4-\epsilon} [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; L^2(\mathbb{R}^3))}^q \geq \Lambda^{2q}\right) \\
 &\leq \Lambda^{-2q} E_\omega \left( \| H^{\sigma/2+1/4-\epsilon} [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; L^2(\mathbb{R}^3))}^q \right) \\
 &= \Lambda^{-2q} \| H^{\sigma/2+1/4-\epsilon} [e^{itH} u_0^\omega]^2 \|_{L^q(\Omega; L^4([- \pi, \pi]; L^2(\mathbb{R}^3)))}^q \\
 &\leq \Lambda^{-2q} \| H^{\sigma/2+1/4-\epsilon} [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; L^2(\mathbb{R}^3; L^q(\Omega)))}^q.
 \end{aligned}$$

Then with (7.4) we get

$$\begin{aligned}
& \| H^{\sigma/2+1/4-\epsilon} [e^{itH} u_0^\omega]^2 \|_{L^q(\Omega)} = \\
& = \left\| H^{\sigma/2+1/4-\epsilon} \left[ \sum_{n,m \in \mathbb{N}} e^{it(\lambda_n^2 + \lambda_m^2)} c_n c_m h_n(x) h_m(x) g_n(\omega) g_m(\omega) \right] \right\|_{L^q(\Omega)} \\
& \leq \left\| \sum_{n,m \in \mathbb{N}} e^{it(\lambda_n^2 + \lambda_m^2)} c_n c_m H^{\sigma/2+1/4-\epsilon} [h_n(x) h_m(x)] g_n(\omega) g_m(\omega) \right\|_{L^q(\Omega)} \\
& \leq Cq \sqrt{\sum_{n,m \in \mathbb{N}} |c_n|^2 |c_m|^2 H^{\sigma/2+1/4-\epsilon} [h_n(x) h_m(x)]^2}.
\end{aligned}$$

By the triangle inequality, we obtain

$$\begin{aligned}
& P \left( \omega \in \Omega : \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^{\sigma+1/2-2\epsilon}(\mathbb{R}^3))} \geq \Lambda^2 \right) \\
& \leq \left( \frac{Cq}{\Lambda^2} \right)^q \left\| \sum_{n,m \in \mathbb{N}} |c_n|^2 |c_m|^2 H^{\sigma/2+1/4-\epsilon} [h_n(x) h_m(x)] \right\|_{L^2([- \pi, \pi]; L^1(\mathbb{R}^3))}^{q/2} \\
& = C \left( \frac{Cq}{\Lambda^2} \right)^q \left( \sum_{n,m \in \mathbb{N}} |c_n|^2 |c_m|^2 \| H^{\sigma/2+1/4-\epsilon} [h_n(x) h_m(x)] \|_{L^2(\mathbb{R}^3)}^2 \right)^{q/2}.
\end{aligned}$$

Then thanks to Proposition 2.12 with  $\delta = \epsilon$ , we deduce that

$$\begin{aligned}
\| H^{\sigma/2+1/4-\epsilon} [h_n(x) h_m(x)] \|_{L^2(\mathbb{R}^3)}^2 & \leq C_\epsilon \max(\lambda_n, \lambda_m)^{2(\sigma-\epsilon)} \\
& \leq C \max(\lambda_n, \lambda_m)^{2\sigma}.
\end{aligned}$$

And finally, we have

$$\begin{aligned}
& P \left( \omega \in \Omega : \| [e^{itH} u_0^\omega]^2 \|_{L^4([- \pi, \pi]; \mathcal{H}^{\sigma+1/2-2\epsilon}(\mathbb{R}^3))} \geq \Lambda^2 \right) \leq \\
& \leq \left( \frac{Cq}{\Lambda^2} \right)^q \left( \sum_{n,m \in \mathbb{N}} |c_n|^2 |c_m|^2 \max(\lambda_n, \lambda_m)^{2\sigma} \right)^{q/2} \\
& \leq \left( \frac{Cq \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}{\Lambda^2} \right)^q.
\end{aligned}$$

Then it is enough to choose  $q = \frac{t^2}{2C \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2} \geq 4$  to conclude.  $\square$

• **Case**  $\| \Delta_N [e^{itH} u_0^\omega] \|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \geq N^{-1/6} \Lambda$ . Let us start by establishing the following lemma:

**Lemma 7.4.** — For all  $p_1, p_2 \in [2, +\infty[$  and  $\epsilon > 0$ , there exist two constants  $C, c > 0$  such that for all  $\Lambda > 0$ ,  $N \geq 1$  and  $u_0 \in \mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)$ ,

$$\begin{aligned} P\left(\omega \in \Omega : \|\Delta_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]; W^{\epsilon, p_2}(\mathbb{R}^3))} \geq \Lambda N^{-1/6-\sigma+2\epsilon}\right) &\leq \\ &\leq C e^{-\frac{c\Lambda^2}{\|\Delta_N(u_0)\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}^2}}. \end{aligned}$$

*Proof.* — We replace  $u_0$  by  $\Delta_N(u_0)$ , therefore we are led to prove that

$$P\left(\omega \in \Omega : \|\Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]; W^{\epsilon, p_2}(\mathbb{R}^3))} \geq \Lambda N^{-1/6-\sigma+2\epsilon}\right) \leq C e^{-\frac{c\Lambda^2}{\|u_0\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}^2}}.$$

As for Lemma 7.3, it is sufficient to prove the estimate for  $\Lambda \geq C\|u_0\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}$ . Thanks to the Markov and Minkowski inequalities, we obtain for  $q \geq p_1, p_2$ ,

$$\begin{aligned} P\left(\omega \in \Omega : \|\Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]; W^{\epsilon, p_2}(\mathbb{R}^3))} \geq \Lambda N^{-1/6-\sigma+2\epsilon}\right) &\leq \\ &\leq \left(\frac{N^{1/6+\sigma-2\epsilon} E_\omega(\|\Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]; W^{\epsilon, p_2}(\mathbb{R}^3))})}{\Lambda}\right)^q \\ &\leq \left(\frac{N^{1/6+\sigma-2\epsilon}}{\Lambda}\right)^q \|H^{\epsilon/2} \Delta'_N(e^{itH}u_0^\omega)\|_{L^q(\Omega), L^{p_1}([- \pi, \pi]); L^{p_2}(\mathbb{R}^3)}^q \\ &\leq \left(\frac{N^{1/6+\sigma-2\epsilon}}{\Lambda}\right)^q \|H^{\epsilon/2} \Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]); L^{p_2}(\mathbb{R}^3); L^q(\Omega)}^q. \end{aligned}$$

But using (7.3),

$$\begin{aligned} \|H^{\epsilon/2} \Delta'_N(e^{itH}u_0^\omega)\|_{L^q(\Omega)} &\leq \left\| \sum_{n \in \mathbb{N}} \phi\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^\epsilon e^{it\lambda_n^2} c_n h_n(x) g_n(\omega) \right\|_{L^q(\Omega)} \\ &\leq \left\| \sum_{\lambda_n \sim N} \phi\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^\epsilon e^{it\lambda_n^2} c_n h_n(x) g_n(\omega) \right\|_{L^q(\Omega)} \\ &\leq C\sqrt{q} \sqrt{\sum_{\lambda_n \sim N} \phi^2\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^{2\epsilon} |c_n|^2 |h_n(x)|^2}. \end{aligned}$$

Then, using (2.9) and (2.10), we obtain

$$\begin{aligned}
& \|H^{\epsilon/2}\Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]); L^{p_2}(\mathbb{R}^3); L^q(\Omega)} \leq \\
& \leq C\sqrt{q} \left\| \sum_{\lambda_n \sim N} \phi^2\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^{2\epsilon} |c_n|^2 |h_n(x)|^2 \right\|_{L^{p_1/2}([- \pi, \pi]); L^{p_2/2}(\mathbb{R}^3)}^{1/2} \\
& \leq C\sqrt{q} \sqrt{\sum_{\lambda_n \sim N} \phi^2\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^{2\epsilon} |c_n|^2 \|h_n(x)\|_{L^{p_2}(\mathbb{R}^3)}^2} \\
& \leq C\sqrt{q} \sqrt{\sum_{\lambda_n \sim N} \phi^2\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^{2\epsilon-1/3} |c_n|^2} \\
& \leq C\sqrt{q} N^{-\sigma-1/6+2\epsilon} \sqrt{\sum_{\lambda_n \sim N} \phi^2\left(\frac{\lambda_n^2}{N^2}\right) \lambda_n^{2(\sigma-\epsilon)} |c_n|^2} \\
& \leq C\sqrt{q} N^{-\sigma-1/6+2\epsilon} \|\Delta'_N(u_0)\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)} \\
& \leq C\sqrt{q} N^{-\sigma-1/6+2\epsilon} \|u_0\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}.
\end{aligned}$$

Finally, we have for all  $q \geq p_1, p_2$ ,

$$\begin{aligned}
P\left(\omega \in \Omega : \|\Delta'_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]); W^{\epsilon, p_2}(\mathbb{R}^3)} \geq \Lambda N^{-1/6-\sigma+2\epsilon}\right) & \leq \\
& \leq \left(\frac{C\sqrt{q}\|u_0\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}}{\Lambda}\right)^q.
\end{aligned}$$

It is then sufficient to choose  $q = \left(\frac{\Lambda}{4C\|u_0\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}}\right)^2 \geq p_1, p_2$  to prove Lemma 7.4.  $\square$

Then for  $p_2 = \frac{3}{\epsilon} + \epsilon$ , we have

$$W^{\epsilon, p_2}(\mathbb{R}^3) \hookrightarrow L^\infty(\mathbb{R}^3).$$

Hence for all  $p_1 \in [2, +\infty[$  and  $\epsilon > 0$ , there exist two constants  $C, c > 0$  such that for all  $\Lambda > 0$ ,  $N \geq 1$  and  $u_0 \in \mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)$ ,

$$P\left(\omega \in \Omega : \|\Delta_N(e^{itH}u_0^\omega)\|_{L^{p_1}([- \pi, \pi]); L^\infty(\mathbb{R}^3)} \geq \Lambda N^{-1/6-\sigma+2\epsilon}\right) \leq C e^{-\frac{c\Lambda^2}{\|\Delta_N(u_0)\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}^2}}.$$

Then we can choose  $p_1 = 4$  and use that  $\|\Delta_N(u_0)\|_{\mathcal{H}^{\sigma-\epsilon}(\mathbb{R}^3)}^2 \leq N^{-2\epsilon}\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2$  to obtain for any  $\epsilon > 0$  the existence of two constants  $C, c > 0$  such that for all  $\Lambda > 0$ ,

$N \geq 1$  and  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$ ,

$$(7.7) \quad P\left(\omega \in \Omega : \|\Delta_N(e^{itH}u_0^\omega)\|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \geq \Lambda N^{-1/6}\right) \leq C e^{-\frac{cN^{2(\sigma-\epsilon)}\Lambda^2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}}.$$

We have to prove that there exist two constants  $C, c > 0$  such that for all  $\Lambda > 0$ ,  $N \geq 1$  and  $u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3)$ ,

$$(7.8) \quad P\left(\bigcup_N \left\{\omega \in \Omega : \|\Delta_N(e^{itH}u_0^\omega)\|_{L^4([- \pi, \pi]; L^\infty(\mathbb{R}^3))} \geq \Lambda N^{-1/6}\right\}\right) \leq C e^{-\frac{c\Lambda^2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}}.$$

We can assume that  $\Lambda \geq C\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}$ , and we set  $\alpha = \left(\frac{\Lambda}{C\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}}\right)^2$  and choosing  $\epsilon < \sigma$  in (7.7), it is enough to show that

$$\forall \delta > 0, \exists C, c > 0 / \forall \alpha \geq 1, \sum_{N \text{ dyadic}} e^{-\alpha N^\delta} \leq C e^{-c\alpha}.$$

Using that there exists  $c > 0$ , such that  $ck \leq 2^{\delta k} - 1$

$$\sum_{N \text{ dyadic}} e^{-\alpha N^\delta} = e^{-\alpha} \sum_{k \geq 0} e^{-\alpha(2^{\delta k} - 1)} \leq e^{-\alpha} \sum_{k \geq 0} e^{-c\alpha k} \leq C e^{-\alpha},$$

and (7.8) is proved. This completes the proof of Theorem 7.1.

**7.2. Proof of Theorem 1.2.** — Recall the definition (7.1) of the set  $\Omega_\Lambda$ . To show Theorem 1.2, it is sufficient to show that for any  $\Lambda > 0$ ,  $P(\Omega_\Lambda) > 0$ . To do this, we first establish that it is sufficient to show the result for a finite number of terms in the initial data. For  $u_0 = \sum_{n \in \mathbb{N}} c_n h_n \in L^2(\mathbb{R}^3)$ , we define for  $K \in \mathbb{N}^*$ ,

$$[u_0]_K = \sum_{\lambda_n < K} c_n h_n, \quad [u_0]^K = \sum_{\lambda_n \geq K} c_n h_n.$$



By independence of the random variables, we obtain

$$\begin{aligned}
P(\Omega_\Lambda) \geq & \mu \left( \left\| [u_0]^K \right\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \frac{\Lambda}{2}, \left\| (e^{itH}[u_0]^K)^2 \right\|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^2}{2}, \right. \\
& \left. \left\| (e^{itH}[u_0]^K)^3 \right\|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^3}{2}, \right. \\
& \left. \bigcap_N \left\{ \left\| \Delta_N(e^{itH}[u_0]^K) \right\|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \frac{\Lambda N^{-1/6}}{2} \right\}, \right. \\
& \left. \bigcap_N \left\{ \left\| \Delta_N(e^{itH}[u_0]^K) \right\|_{L^R([-\pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \frac{\Lambda N^{s-1/4}}{2} \right\} \right). \\
& \cdot \mu \left( \left\| [u_0]_K \right\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \frac{\Lambda}{2}, \left\| (e^{itH}[u_0]_K)^2 \right\|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^2}{2}, \right. \\
& \left. \left\| (e^{itH}[u_0]_K)^3 \right\|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^3}{2}, \right. \\
& \left. \bigcap_N \left\{ \left\| \Delta_N(e^{itH}[u_0]_K) \right\|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \frac{\Lambda N^{-1/6}}{2} \right\}, \right. \\
& \left. \bigcap_N \left\{ \left\| \Delta_N(e^{itH}[u_0]_K) \right\|_{L^R([-\pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \frac{\Lambda N^{s-1/4}}{2} \right\} \right).
\end{aligned}$$

Let us denote by  $P_{\Lambda, K}$  the first probabilistic term of this inequality. Then by Theorem 7.1, for all  $\Lambda \geq 0$  and  $K \in \mathbb{N}^*$ ,

$$P_{\Lambda, K} \geq 1 - C \exp \left( - \frac{c\Lambda^2}{\left\| [u_0]^K \right\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2} \right),$$

with

$$\lim_{K \rightarrow +\infty} \left\| [u_0]^K \right\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2 = 0.$$

Therefore, there is an integer  $K \geq 1$  such that  $C \exp(-\frac{c\Lambda^2}{\| [u_0]_K \|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}) \leq \alpha$ , and we deduce that

$$(7.9) \quad \frac{P(\Omega_\Lambda)}{1-\alpha} \geq \mu \left( u_0 \in \mathcal{H}^\sigma(\mathbb{R}^3) : \| [u_0]_K \|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \frac{\Lambda}{2}, \| (e^{itH} [u_0]_K)^2 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^2}{2}, \right. \\ \left. \| (e^{itH} [u_0]_K)^3 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^3}{2}, \right. \\ \left. \bigcap_N \left\{ \| \Delta_N(e^{itH} [u_0]_K) \|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \frac{\Lambda N^{-1/6}}{2} \right\}, \right. \\ \left. \bigcap_N \left\{ \| \Delta_N(e^{itH} [u_0]_K) \|_{L^R([-\pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \frac{\Lambda N^{s-1/4}}{2} \right\} \right).$$

As a result, we are reduced to prove the result for a finite number of terms in the initial data. We notice that there are two constants  $C_1, C_2 > 0$  such that  $C_1 n^{\frac{1}{3}} \leq \lambda_n^2 \leq C_2 n^{\frac{1}{3}}$  then that  $|\{\lambda_n \leq K\}| \leq CK^6$ . Thus by Proposition 2.10 and by the Cauchy-Schwarz inequality

$$(7.10) \quad \left\{ \omega \in \Omega : \| [u_0^\omega]_K \|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \frac{\Lambda}{2} \right\} = \left\{ \omega \in \Omega : \sum_{\lambda_n \leq K} \lambda_n^{2\sigma} |c_n|^2 |g_n(\omega)|^2 \leq \frac{\Lambda^2}{CK^6} \right\} \subset A_\Lambda$$

where

$$A_\Lambda = \bigcap \left\{ \omega \in \Omega : \| (e^{itH} [u_0^\omega]_K)^2 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^2}{2} \right\}, \\ \bigcap \left\{ \omega \in \Omega : \| (e^{itH} [u_0^\omega]_K)^3 \|_{L^4([-\pi, \pi]; \mathcal{H}^s(\mathbb{R}^3))} \leq \frac{\Lambda^3}{2} \right\}, \\ \bigcap \bigcap_N \left\{ \omega \in \Omega : \| \Delta_N(e^{itH} [u_0^\omega]_K) \|_{L^4([-\pi, \pi]; L^\infty(\mathbb{R}^3))} \leq \frac{\Lambda}{2} N^{-1/6} \right\}, \\ \bigcap \bigcap_N \left\{ \omega \in \Omega : \| \Delta_N(e^{itH} [u_0^\omega]_K) \|_{L^R([-\pi, \pi]; \mathcal{W}^{s,4}(\mathbb{R}^3))} \leq \frac{\Lambda}{2} N^{s-1/4} \right\}.$$

We are now able to complete the proof of Theorem 1.2. By (7.9) and (7.10), it is sufficient to show that for any integer  $K \geq 1$  and any  $\Lambda > 0$ ,

$$P\left(\omega \in \Omega : \sum_{\lambda_n \leq K} \lambda_n^{2\sigma} |c_n|^2 |g_n(\omega)|^2 \leq \frac{\Lambda^2}{CK^6}\right) > 0.$$

But, by independence,

$$\begin{aligned} P\left(\omega \in \Omega : \sum_{\lambda_n \leq K} \lambda_n^{2\sigma} |c_n|^2 |g_n(\omega)|^2 \leq \frac{\Lambda^2}{CK^6}\right) &\geq \\ &\geq P\left(\bigcap_{\lambda_n \leq K} \left\{\omega \in \Omega : |g_n(\omega)|^2 \leq \frac{\Lambda^2}{CK^{12} \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}\right\}\right) \\ &= \prod_{\lambda_n \leq K} P\left(\omega \in \Omega : |g_n(\omega)|^2 \leq \frac{\Lambda^2}{CK^{12} \|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}\right) > 0, \end{aligned}$$

because for all  $R > 0$  and all  $n \in \mathbb{N}$ ,  $P(\omega \in \Omega : |g_n(\omega)| \leq R) > 0$ .

It remains to show (1.7). By Theorem 6.11 and Theorem 6.12, it is sufficient to establish that for any  $\Lambda > 0$ ,

$$\lim_{\eta \rightarrow 0} P(\omega \in \Omega_\Lambda : \|u_0^\omega\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \eta) = 1.$$

Actually, by adapting the proof of [23, Appendix A.2], we can obtain that

$$P(\omega \in \Omega_\Lambda^c : \|u_0^\omega\|_{\mathcal{H}^\sigma(\mathbb{R}^3)} \leq \eta) \leq C e^{-c \frac{\Lambda^2}{\eta^2}},$$

and (1.7) is proven.

**7.3. Proof of (1.8).** — To prove (1.8), it is enough to show that for any  $\Lambda > 0$  and any  $\alpha \in ]0, 1]$ ,  $P(\Omega_\Lambda) \geq 1 - \alpha$ . This result is clear from Theorem 7.1 since

$$P(\Omega_\Lambda) \geq 1 - C_1 \exp\left(-\frac{C_2}{\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}^2}\right) \geq 1 - \alpha$$

if  $\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^3)}$  is small enough.

## 8. The fixed point argument for equation (1.11)

We introduce the following equation:

$$(8.1) \quad \begin{cases} i \frac{\partial u}{\partial t} - Hu = \kappa \cos(2t)^{\frac{d(p-1)}{2}-2} |u|^{p-1} u, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ u(0, x) = u_0(x), \end{cases}$$

where  $p \geq 5$  denotes an odd integer and  $\kappa \in \{-1, 1\}$ . We will first show that this equation is almost surely locally well-posed, and then deduce global well-posedness of (1.11) using the lens transform.

### 8.1. Some nonlinear estimates and local well-posedness of equation (8.1).

— In this section, we establish estimates that will be useful to apply the Picard fixed point theorem. We start by showing two preliminary lemmas.

**Lemma 8.1.** — *Let  $(q, r) \in [2, +\infty[ \times [2, +\infty]$ ,  $s, s_0 \geq 0$  and assume that  $s - s_0 > \frac{d}{2} - \frac{2}{q} - \frac{d}{r}$ , then there are two constants  $\kappa, C > 0$  such that for all  $T \geq 0$  and  $u \in \overline{X}_T^s$ ,*

$$\|u\|_{L^q([-T, T]; \mathcal{W}^{s_0, r}(\mathbb{R}^d))} \leq CT^\kappa \|u\|_{\overline{X}_T^s}.$$

*Proof.* — Let  $\epsilon > 0$  then there exists  $\kappa_\epsilon > 0$  such that

$$\|u\|_{L^q([-T, T]; \mathcal{W}^{s_0, r}(\mathbb{R}^d))} \leq T^{\kappa_\epsilon} \|u\|_{L^{q+\epsilon}([-T, T]; \mathcal{W}^{s_0, r}(\mathbb{R}^d))}.$$

Now the couple  $(q + \epsilon, \frac{2d(q+\epsilon)}{dq+d\epsilon-4})$  is admissible with

$$\mathcal{W}^{s, \frac{2d(q+\epsilon)}{dq+d\epsilon-4}}(\mathbb{R}^d) \hookrightarrow \mathcal{W}^{s_0, r}(\mathbb{R}^d) \quad \text{if} \quad s - s_0 \geq \frac{d}{2} - \frac{2}{q + \epsilon} - \frac{d}{r}.$$

But, as  $s - s_0 > \frac{d}{2} - \frac{2}{q} - \frac{d}{r}$  then there is  $0 < \epsilon \ll 1$  such that  $s - s_0 \geq \frac{d}{2} - \frac{2}{q+\epsilon} - \frac{d}{r}$ .  $\square$

**Lemma 8.2.** — *Let  $s \geq 0$  then there exists a constant  $C > 0$  such that for all functions  $f$  and  $g$  in  $\mathcal{S}(\mathbb{R}^d)$ ,*

$$\|(-\Delta)^{s/2}(fg)\|_{L^2(\mathbb{R}^d)} \leq C(\|g(-\Delta)^{s/2}f\|_{L^2(\mathbb{R}^d)} + \|f(-\Delta)^{s/2}g\|_{L^2(\mathbb{R}^d)}).$$

*Proof.* — Using the Fourier transform we obtain

$$\begin{aligned} \|(-\Delta)^{s/2}(fg)\|_{L^2(\mathbb{R}^d)} &\leq C\|\xi|^s \mathcal{F}(fg)\|_{L^2(\mathbb{R}^d)} \\ &= C\|\xi|^s \mathcal{F}(f) \star \mathcal{F}(g)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

But for all  $\xi, \eta \in \mathbb{R}^d$ ,

$$|\xi|^s \leq (|\eta| + |\xi - \eta|)^s \leq C_s(|\eta|^s + |\xi - \eta|^s),$$

therefore

$$\begin{aligned} \|(-\Delta)^{s/2}(fg)\|_{L^2(\mathbb{R}^d)} &\leq C_s(\|(|\cdot|^s \mathcal{F}(f)) \star \mathcal{F}(g)\|_{L^2(\mathbb{R}^d)} + \|\mathcal{F}(f) \star (|\cdot|^s \mathcal{F}(g))\|_{L^2(\mathbb{R}^d)}) \\ &= C_s(\|\mathcal{F}(g(-\Delta)^{s/2}f)\|_{L^2(\mathbb{R}^d)} + \|\mathcal{F}(f(-\Delta)^{s/2}g)\|_{L^2(\mathbb{R}^d)}) \\ &= C_s(\|g(-\Delta)^{s/2}f\|_{L^2(\mathbb{R}^d)} + \|f(-\Delta)^{s/2}g\|_{L^2(\mathbb{R}^d)}), \end{aligned}$$

which was the claim.  $\square$

Then, the expected estimates are established.

**Proposition 8.3.** — *Let  $s > \frac{d}{2} - \frac{2}{p-1}$  then there exist two constants  $C > 0$  and  $\kappa > 0$  such that if we assume*

$$\|e^{-itH}u_0\|_{L^p([- \pi, \pi]; L^\infty(\mathbb{R}^d))} \leq \lambda$$

for some  $\lambda > 0$ , then for any  $0 < T \leq 1$ ,  $v \in \overline{X}_T^s$  and  $f_j = v$  or  $f_j = e^{-itH}u_0$ ,

$$\|(\sqrt{-\Delta}^s v) \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa(\lambda^p + \|v\|_{\overline{X}_T^s}^p),$$

and

$$\|\langle x \rangle^s v \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa(\lambda^p + \|v\|_{\overline{X}_T^s}^p).$$

*Proof.* — According to the Hölder inequality and (1.2),

$$\begin{aligned} \|(\sqrt{-\Delta}^s v) \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} &\leq \\ &\leq \|\sqrt{-\Delta}^s v\|_{L^\infty([-T, T]; L^2(\mathbb{R}^d))} \prod_{j=2}^p \|f_j\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))} \\ &\leq C\|v\|_{L^\infty([-T, T]; \mathcal{H}^s(\mathbb{R}^d))} \prod_{j=2}^p \|f_j\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))}, \end{aligned}$$

and

$$\begin{aligned} \|\langle x \rangle^s v \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} &\leq \|\langle x \rangle^s v\|_{L^\infty([-T, T]; L^2(\mathbb{R}^d))} \prod_{j=2}^p \|f_j\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))} \\ &\leq \|v\|_{L^\infty([-T, T]; \mathcal{H}^s(\mathbb{R}^d))} \prod_{j=2}^p \|f_j\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))}. \end{aligned}$$

If  $f_j = v$  then as  $s > \frac{d}{2} - \frac{2}{p-1}$ , we can use Lemma 8.1 to get

$$\|v\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))} \leq CT^\kappa \|v\|_{\overline{X}_T^s}.$$

If  $f_j = e^{-itH}u_0$  then according to the Hölder inequality,

$$\begin{aligned} \|e^{-itH}u_0\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))} &\leq T^{\frac{1}{p-1} - \frac{1}{p}} \|e^{-itH}u_0\|_{L^p([- \pi, \pi]; L^\infty(\mathbb{R}^d))} \\ &\leq T^{\frac{1}{p(p-1)}} \lambda, \end{aligned}$$

which completes the proof.  $\square$

**Lemma 8.4.** — *If  $0 < s < \frac{d}{2}$  then there exist two constants  $C > 0$  and  $\kappa > 0$  such that if we assume that*

$$\|e^{-itH}u_0\|_{L^p([- \pi, \pi]; \mathcal{W}^{\frac{1}{8}, \infty}(\mathbb{R}^d))} \leq \lambda,$$

and

$$\|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)} \leq \lambda$$

for some  $\lambda > 0$ , then for all  $0 < T \leq 1$ ,

$$\|\langle x \rangle^s (e^{-itH}u_0)^p\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa \lambda^p.$$

*Proof.* — According to the Hölder inequality and (1.2), we obtain

$$\begin{aligned} & \|\langle x \rangle^s (e^{-itH}u_0)^p\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq \\ & \leq \|\langle x \rangle^{\frac{d}{2}} (e^{-itH}u_0)^p\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \\ & \leq \|\langle x \rangle^{\frac{d-1}{2}} e^{-itH}u_0\|_{L^\infty([-T, T]; L^2(\mathbb{R}^d))} \|\langle x \rangle^{\frac{1}{2(p-1)}} e^{-itH}u_0\|_{L^{p-1}([-T, T]; L^\infty(\mathbb{R}^d))}^{p-1} \\ & \leq CT^{1/p} \|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)} \|e^{-itH}u_0\|_{L^p([-T, T]; \mathcal{W}^{\frac{1}{8}, \infty}(\mathbb{R}^d))}^{p-1} \\ & \leq CT^{1/p} \lambda^p. \end{aligned}$$

□

**Lemma 8.5.** — *There exist  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$  and constants  $C, \kappa > 0$  such that if we assume that*

$$\|e^{-itH}u_0\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \lambda,$$

and

$$\|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)} \leq \lambda$$

for some  $\lambda > 0$ , then for all  $0 < T \leq 1$ ,  $v \in \overline{X}_T^s$  and  $f_j = v$  or  $f_j = e^{-itH}u_0$ ,

$$\|\sqrt{-\Delta}^s (e^{-itH}u_0) \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa (\lambda^p + \|v\|_{\overline{X}_T^s}^p).$$

*Proof.* — For all  $\epsilon \in ]0, \frac{1}{2}[$ , according to the Hölder inequality, we have

$$\begin{aligned} & \|\sqrt{-\Delta}^s (e^{-itH} u_0) \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq \\ & \leq \left\| \frac{\sqrt{-\Delta}^s (e^{-itH} u_0)}{\langle x \rangle^{1/2-\epsilon}} \right\|_{L^2([-T, T]; L^2(\mathbb{R}^d))} \prod_{j=2}^p \|\langle x \rangle^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)} f_j\|_{L^{2(p-1)}([-T, T]; L^\infty(\mathbb{R}^d))}. \end{aligned}$$

Then, we choose  $s = \frac{d}{2} - 2\epsilon$  with  $\epsilon \ll 1$  to obtain by using (1.16) that

$$\begin{aligned} & \|\sqrt{-\Delta}^s (e^{-itH} u_0) \prod_{j=2}^p f_j\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq \\ & \leq \lambda \prod_{j=2}^p \|\langle x \rangle^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)} f_j\|_{L^{2(p-1)}([-T, T]; L^\infty(\mathbb{R}^d))} \\ & \leq \lambda \prod_{j=2}^p \|f_j\|_{L^{2(p-1)}([-T, T]; \mathcal{W}^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)+\epsilon, \frac{d}{\epsilon}+1}(\mathbb{R}^d))}. \end{aligned}$$

If  $f_j = e^{-itH} u_0$ , by interpolation, we there exists a constant  $\kappa > 0$  such that

$$\begin{aligned} & \|e^{-itH} u_0\|_{L^{2(p-1)}([-T, T]; \mathcal{W}^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)+\epsilon, \frac{d}{\epsilon}+1}(\mathbb{R}^d))} \leq \\ & \leq CT^\kappa \|e^{-itH} u_0\|_{L^{2p}([-T, T]; \mathcal{W}^{s_0, \infty}(\mathbb{R}^d))} \|e^{-itH} u_0\|_{L^\infty([-T, T]; \mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d))}^{1-\theta} \end{aligned}$$

where  $\theta = \frac{d-\epsilon}{d+\epsilon}$  and  $s_0 = (\frac{1-\theta}{\theta})(\frac{d-1}{2}) + \frac{1}{\theta(p-1)}(\frac{1}{2}-\epsilon) + \frac{\epsilon}{\theta}$ .

Firstly,  $\|e^{-itH} u_0\|_{L^\infty([-T, T]; \mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d))} = \|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)} \leq \lambda$ . Then as

$$s_0 = \frac{1}{2(p-1)} + C\epsilon + o(\epsilon) \leq \frac{1}{7}$$

we deduce  $\|e^{-itH} u_0\|_{L^{2p}([-T, T]; \mathcal{W}^{s_0, \infty}(\mathbb{R}^d))} \leq \lambda$  and

$$\|e^{-itH} u_0\|_{L^{2(p-1)}([-T, T]; \mathcal{W}^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)+\epsilon, \frac{d}{\epsilon}+1}(\mathbb{R}^d))} \leq \lambda.$$

If  $f_j = v$ , since  $s - \frac{1}{p-1}(\frac{1}{2}-\epsilon) > \frac{d}{2} - \frac{1}{p-1} - \frac{d\epsilon}{d+\epsilon}$  (if  $\epsilon \ll \frac{1}{2(p-2)}$ ) then by Lemma 8.1, we obtain

$$\|v\|_{L^{2(p-1)}([-T, T]; \mathcal{W}^{\frac{1}{p-1}(\frac{1}{2}-\epsilon)+\epsilon, \frac{d}{\epsilon}+1}(\mathbb{R}^d))} \leq CT^\kappa \|v\|_{\overline{X}_T^s},$$

hence the result.  $\square$

We therefore introduce the natural set of initial conditions. Let  $\lambda \geq 0$  and define  $F_0(\lambda)$  as the set of functions  $u_0 \in \mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)$  which satisfy

$$(8.2) \quad \begin{cases} \|u_0\|_{\mathcal{H}^{\frac{d-1}{2}}(\mathbb{R}^d)} & \leq \lambda, \\ \|e^{-itH}u_0\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} & \leq \lambda. \end{cases}$$

Then, we can establish the two main results of this section.

**Proposition 8.6.** — *There exists  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$  and constants  $C, \kappa > 0$  such that if  $u_0 \in F_0(\lambda)$  for some  $\lambda > 0$  then for all  $v, v_1, v_2 \in \overline{X}_T^s$  and  $0 < T \leq 1$ ,*

$$\begin{aligned} \left\| \int_0^t e^{-i(t-s)H} \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) ds \right\|_{\overline{X}_T^s} &\leq \\ &\leq CT^\kappa (\lambda^p + \|v\|_{\overline{X}_T^s}^p), \end{aligned}$$

and

$$\begin{aligned} &\left\| \int_0^t e^{-i(t-s)H} \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v_1|^{p-1} (e^{-isH}u_0 + v_1) ds \right. \\ &\quad \left. - \int_0^t e^{-i(t-s)H} \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v_2|^{p-1} (e^{-isH}u_0 + v_2) ds \right\|_{\overline{X}_T^s} \\ &\leq CT^\kappa \|v_1 - v_2\|_{\overline{X}_T^s} (\lambda^{p-1} + \|v_1\|_{\overline{X}_T^s}^{p-1} + \|v_2\|_{\overline{X}_T^s}^{p-1}). \end{aligned}$$

*Proof.* — We only prove the first estimate. Using Proposition 2.1 and (1.2), we obtain

$$\begin{aligned} &\left\| \int_0^t e^{-i(t-s)H} \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) ds \right\|_{\overline{X}_T^s} \leq \\ &\leq C \left\| \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) \right\|_{L^1([-T, T]; \mathcal{H}^s(\mathbb{R}^d))} \\ &\leq C \left\| |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) \right\|_{L^1([-T, T]; \mathcal{H}^s(\mathbb{R}^d))} \\ &\leq C \left\| (-\Delta)^s (|e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v)) \right\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \\ &\quad + C \left\| \langle x \rangle^s |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) \right\|_{L^1([-T, T]; L^2(\mathbb{R}^d))}. \end{aligned}$$

Then, using Lemma 8.2 and Proposition 8.3, Lemma 8.4, and Lemma 8.5, we can obtain a constant  $\kappa > 0$  such that for all  $u_0 \in F_0(\lambda)$ ,  $0 < T \leq 1$  and  $v \in \overline{X}_T^s$ ,

$$\left\| (-\Delta)^s (|e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v)) \right\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa (\lambda^p + \|v\|_{\overline{X}_T^s}^p),$$

and

$$\left\| \langle x \rangle^s |e^{-isH}u_0 + v|^{p-1} (e^{-isH}u_0 + v) \right\|_{L^1([-T, T]; L^2(\mathbb{R}^d))} \leq CT^\kappa (\lambda^p + \|v\|_{\overline{X}_T^s}^p),$$



which was to prove.  $\square$

We are now able to state the local well-posedness result for equation (8.1).

**Theorem 8.7.** — *There exist  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$ ,  $C > 0$  and  $\delta > 0$  such that for any  $0 < T \leq 1$ , if  $u_0 \in F_0(\lambda)$  with  $\lambda < CT^{-\delta}$  then there is a unique solution to the equation (8.1) on  $[-T, T]$  in the space  $e^{-itH}u_0 + B_{\overline{X}_T^s}(0, \lambda)$ .*

*Proof.* — Let us define

$$L(v) = -i\kappa \int_0^t e^{-i(t-s)H} \cos(2s)^{\frac{d(p-1)}{2}-2} |e^{-isH}u_0 + v(s)|^{p-1} (e^{-isH}u_0 + v(s)) ds,$$

and note that  $u = e^{-itH}u_0 + v$  is the unique solution of (8.1) on  $[-T, T]$  in the space  $e^{-itH}u_0 + B_{\overline{X}_T^s}(0, R)$  if and only if  $v$  is the unique fixed point of  $L$  on  $B_{\overline{X}_T^s}(0, R)$ .

According to Proposition 8.6, there are two constants  $C > 0$  and  $\kappa > 0$  such that

$$\begin{aligned} \|L(v)\|_{\overline{X}_T^s} &\leq CT^\kappa (\lambda^p + \|v\|_{\overline{X}_T^s}^p) \\ \|L(v_1) - L(v_2)\|_{\overline{X}_T^s} &\leq CT^\kappa \|v_1 - v_2\|_{\overline{X}_T^s} (\lambda^{p-1} + \|v_1\|_{\overline{X}_T^s}^{p-1} + \|v_2\|_{\overline{X}_T^s}^{p-1}). \end{aligned}$$

Therefore if  $\lambda < (\frac{1}{8CT^\kappa})^{\frac{1}{p-1}}$  then  $L$  is a contraction of  $B_{\overline{X}_T^s}(0, \lambda)$  and the theorem follows.  $\square$

**8.2. Global solutions and scattering for the equation (1.11).** — Thanks to the lens transform and the results of the previous section, we are able obtain the existence of global solutions for the equation (1.11).

**Theorem 8.8.** — *There exist  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$  and  $C_1, C_2 > 0$  such that if  $u_0 \in F_0(\lambda)$  with  $\lambda < C_1$  then there is a global solution to (1.11) in the space  $e^{it\Delta}u_0 + B_{X^s}(0, C_2)$ .*

*Proof.* — Let  $u$  be given by Theorem 8.7 with  $T = \frac{\pi}{4}$ . We apply to  $u$  the lens transformation defined in Section 2.1 to obtain a function  $U$  which, according to Proposition 2.2, satisfies the conditions of the theorem.  $\square$

**Theorem 8.9.** — *There exist  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$ ,  $C_1, C_2 > 0$  and  $\delta > 0$  such that for any  $0 < T \leq 1$ , if  $u_0 \in F_0(\lambda)$  with  $\lambda < C_1(\arctan 2T)^{-\delta}$  then there is a solution to (1.11) on  $[-T, T]$  in the space  $e^{it\Delta}u_0 + B_{X_T^s}(0, C_2\lambda^p)$ .*

*Proof.* — Let  $u$  be given by Theorem 8.7 at  $T$  replaced by  $\frac{1}{2} \arctan 2T$ . Then, as for the previous proof, we apply to  $u$  the lens transformation, which yields the result.  $\square$

We then prove the uniqueness of the constructed solutions.

**Proposition 8.10.** — Let  $s \in ]\frac{d}{2} - \frac{2}{p-1}; \frac{d}{2}[$ ,  $u_0 \in F_0(\lambda)$  and  $T \in ]0, 1]$ . Let  $U_1$  and  $U_2$  be two solutions of (1.11) on  $[-T, T]$  on the space  $e^{it\Delta}u_0 + X_T^s$  then,

$$U_1(t) = U_2(t) \text{ in } L^2(\mathbb{R}^d), \forall t \in [-T, T].$$

*Proof.* — By reversibility of the equation, is enough to consider the case  $t \in [0, T]$ . For all  $t \in \mathbb{R}$ , we have

$$\begin{aligned} \frac{d}{dt} \|U_1(t) - U_2(t)\|_{L^2(\mathbb{R}^d)}^2 &= \\ &= 2\operatorname{Re}(\langle \partial_t(U_1(t) - U_2(t)), U_1(t) - U_2(t) \rangle_{L^2(\mathbb{R}^d)L^2(\mathbb{R}^d)}) \\ &= 2|\langle |U_1(t)|^{p-1}U_1(t) - |U_2(t)|^{p-1}U_2(t), U_1(t) - U_2(t) \rangle_{L^2(\mathbb{R}^d)L^2(\mathbb{R}^d)}| \\ &\leq 2\|U_1(t) - U_2(t)\|_{L^2(\mathbb{R}^d)} \| |U_1(t)|^{p-1}U_1(t) - |U_2(t)|^{p-1}U_2(t) \|_{L^2(\mathbb{R}^d)} \\ &\leq C\|U_1(t) - U_2(t)\|_{L^2(\mathbb{R}^d)}^2 (\|U_1(t)\|_{L^\infty(\mathbb{R}^d)}^{p-1} + \|U_2(t)\|_{L^\infty(\mathbb{R}^d)}^{p-1}). \end{aligned}$$

Then, by the Grönwall Lemma, the result is proved if  $\|U_j(t)\|_{L^\infty(\mathbb{R}^d)}^{p-1} \in L_{loc}^1$ , since  $\|U_1(0) - U_2(0)\|_{L^2(\mathbb{R}^d)}^2 = 0$ . By using Proposition 2.2, we get

$$\begin{aligned} \|U_j\|_{L^{p-1}([0, T], L^\infty(\mathbb{R}^d))} &\leq \|e^{it\Delta}u_0\|_{L^{p-1}([0, T]; L^\infty(\mathbb{R}^d))} + \|V_j\|_{L^{p-1}([0, T]; L^\infty(\mathbb{R}^d))} \\ &\leq C_T(\|e^{-itH}u_0\|_{L^{p-1}([-\pi, \pi]; L^\infty(\mathbb{R}^d))} + \|V_j\|_{X_T^s}) \\ &\leq C_T(\lambda + \|V_j\|_{X_T^s}), \end{aligned}$$

hence the result. □

Finally, we prove that the global solutions constructed scatter at  $+\infty$  and at  $-\infty$ , namely (1.12) and (1.13). The proof is the same as the proof of Theorem 6.12 and is left here.

## 9. Estimation of the regularity of the initial random data and proof of Theorem 1.4

**9.1. Estimation of the regularity of the initial random data.** — For  $\Lambda > 0$ , recall the definition (8.2) of the set  $F_0(\Lambda)$ , and define

$$\Omega_\Lambda = \{\omega \in \Omega : u_0^\omega \in F_0(\Lambda)\}.$$

The purpose of this part is to establish the following theorem:

**Theorem 9.1.** — *There exist constants  $C, c > 0$  such that for any  $\Lambda > 0$ ,*

$$P(\Omega_\Lambda^c) \leq C \exp\left(-c \frac{\Lambda^2}{\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}^2}\right).$$

*Proof.* — By the triangle inequality, we can write

$$(9.1) \quad P(\Omega_\Lambda^c) \leq P(\omega \in \Omega : \|u_0^\omega\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \geq \Lambda) + P(\omega \in \Omega : \|e^{-itH}u_0\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \geq \Lambda)$$

and it suffices to bound each of the previous two terms.

We first estimate the first term of (9.1). It is sufficient to establish the estimate for  $\Lambda \geq C\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}$ . Let  $q \geq 1$  then according to the Markov inequality, we get

$$\begin{aligned} P\left(\omega \in \Omega : \|u_0^\omega\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \geq \Lambda\right) &= P\left(\omega \in \Omega : \sum_{n \in \mathbb{N}} \lambda_n^{d-1} |c_n|^2 |g_n(\omega)|^2 \geq \Lambda^2\right) \\ &\leq \Lambda^{-2q} \left\| \sum_{n \in \mathbb{N}} \lambda_n^{d-1} |c_n|^2 |g_n|^2 \right\|_{L^q(\Omega)}^q \\ &\leq \Lambda^{-2q} \left( \sum_{n \in \mathbb{N}} \lambda_n^{d-1} |c_n|^2 \|g_n\|_{L^{2q}(\Omega)}^2 \right)^q \\ &\leq \left( C \left( \frac{q}{c} \right)^{\frac{1}{2}} \frac{\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}}{\Lambda} \right)^{2q}. \end{aligned}$$

Then we can choose  $q = c \left( \frac{\Lambda}{2C\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}} \right)^2 \geq 1$  to get

$$P\left(\omega \in \Omega : \|u_0^\omega\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \geq \Lambda\right) \leq \frac{1}{2^{2q}} = e^{-2 \ln(2)q} \leq \exp\left(-\frac{c\Lambda^2}{\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}^2}\right).$$

From then on, the second term of (9.1) remains to be estimated. For this, let us recall the estimates of the eigenfunctions of the harmonic oscillator whose proof can be found in [51, Corollary 3.2]. For all  $p \in [4, +\infty]$ , there exists a constant  $C > 0$  such that for all  $n \in \mathbb{N}$ ,

$$(9.2) \quad \|h_n\|_{L^p(\mathbb{R}^d)} \leq C\lambda_n^{-1+\frac{d}{2}} \quad \text{if } d \geq 2.$$

Since  $\mathcal{W}^{\frac{1}{6}, r}(\mathbb{R}^d) \hookrightarrow \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d)$  if  $\frac{1}{6} - \frac{1}{7} > \frac{d}{r}$ , we have to prove that there exist two constants  $C, c > 0$  such that for all  $\Lambda \geq 0$  and  $r \geq 2$ ,

$$(9.3) \quad P\left(\omega \in \Omega : \|e^{-itH}u_0^\omega\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{6}, r}(\mathbb{R}^d))} \geq \Lambda\right) \leq C \exp\left(-c \frac{\Lambda^2}{\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}^2}\right).$$

It is sufficient to show the estimate for  $t \geq C\|u_0\|_{\mathcal{H}^\sigma(\mathbb{R}^d)}$ . According to the Markov and Minkowski inequalities, we obtain for  $q \geq \max(2p, r, 2)$ ,

$$\begin{aligned} P\left(\omega \in \Omega : \|e^{-itH}u_0^\omega\|_{L^{2p}([-\pi, \pi]; \mathcal{W}^{\frac{1}{6}, r}(\mathbb{R}^d))} \geq \Lambda\right) &\leq \\ &\leq \Lambda^{-q} \|e^{-itH}u_0^\omega\|_{L^q(\Omega, L^{2p}([-\pi, \pi]; \mathcal{W}^{\frac{1}{6}, r}(\mathbb{R}^d)))}^q \\ &\leq \Lambda^{-q} \|H^{\frac{1}{12}}e^{-itH}u_0^\omega\|_{L^{2p}([-\pi, \pi]; L^r(\mathbb{R}^d, L^q(\Omega)))}^q. \end{aligned}$$

Then, thanks to (7.3), we obtain

$$\begin{aligned} \|H^{\frac{1}{12}}e^{-itH}u_0^\omega\|_{L^q(\Omega)} &= \left\| \sum_{n \in \mathbb{N}} \lambda_n^{\frac{1}{6}} c_n e^{-it\lambda_n^2} h_n(x) g_n(\omega) \right\|_{L^q(\Omega)} \leq \\ &\leq Cq^{\frac{1}{2}} \sqrt{\sum_{n \in \mathbb{N}} \lambda_n^{\frac{1}{3}} |c_n|^2 |h_n(x)|^2}. \end{aligned}$$

And finally, by the triangle inequality and (9.2), we have

$$\begin{aligned} P\left(\omega \in \Omega : \|e^{-itH}u_0^\omega\|_{L^{2p}([-\pi, \pi]; \mathcal{W}^{\frac{1}{6}, r}(\mathbb{R}^d))} \geq \Lambda\right) &\leq \\ &\leq \left(\frac{Cq^{\frac{1}{2}}}{\Lambda}\right)^q \left\| \sqrt{\sum_{n \in \mathbb{N}} \lambda_n^{\frac{1}{3}} |c_n|^2 |h_n(x)|^2} \right\|_{L^r(\mathbb{R}^d)}^q \\ &\leq \left(\frac{Cq^{\frac{1}{2}}}{\Lambda}\right)^q \left\| \sum_{n \in \mathbb{N}} \lambda_n^{\frac{1}{3}} |c_n|^2 |h_n(x)|^2 \right\|_{L^{r/2}(\mathbb{R}^d)}^{q/2} \\ &\leq \left(\frac{Cq^{\frac{1}{2}}}{\Lambda}\right)^q \left( \sum_{n \in \mathbb{N}} \lambda_n^{\frac{1}{3}} |c_n|^2 \|h_n(x)\|_{L^r(\mathbb{R}^d)}^2 \right)^{q/2} \\ &\leq \left( \frac{Cq^{\frac{1}{2}} \|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}}{\Lambda} \right)^q. \end{aligned}$$

Thus, it is enough to choose  $q = \left( \frac{\Lambda}{2C\|u_0\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)}} \right)^2$  to get (9.3). □

**9.2. Proof of Theorem 1.4.** — To complete the proof of this result, it is sufficient to establish that for all  $\Lambda > 0$ ,

$$(9.4) \quad P(\Omega_\Lambda) > 0.$$

For  $u_0 = \sum_{n \in \mathbb{N}} c_n h_n \in L^2(\mathbb{R}^d)$ , we define for  $N \in \mathbb{N}^*$ ,

$$[u_0]_N = \sum_{\lambda_n < N} c_n h_n, \quad [u_0]^N = \sum_{\lambda_n \geq N} c_n h_n.$$

**Lemma 9.2.** — *There exists  $N \in \mathbb{N}^*$  such that*

$$P(\Omega_\Lambda) \geq \frac{1}{2} P(\omega \in \Omega : \|[u_0^\omega]_N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \frac{\Lambda}{2}, \|e^{-itH}[u_0^\omega]_N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \frac{\Lambda}{2}).$$

*Proof.* — By independence, using Theorem 9.1, we obtain

$$\begin{aligned} P(\Omega_\Lambda) &= P\left(\omega \in \Omega : \|[u_0^\omega]_N + [u_0^\omega]^N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \Lambda, \right. \\ &\quad \left. \|e^{-itH}[u_0^\omega]_N + e^{-itH}[u_0^\omega]^N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \Lambda\right) \\ &\geq P\left(\omega \in \Omega : \|[u_0^\omega]_N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \frac{\Lambda}{2}, \|e^{-itH}[u_0^\omega]_N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \frac{\Lambda}{2}\right) \\ &\quad \cdot P\left(\omega \in \Omega : \|[u_0^\omega]^N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \frac{\Lambda}{2}, \|e^{-itH}[u_0^\omega]^N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \frac{\Lambda}{2}\right) \\ &\geq P\left(\omega \in \Omega : \|[u_0^\omega]_N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \frac{\Lambda}{2}, \|e^{-itH}[u_0^\omega]_N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \frac{\Lambda}{2}\right) \\ &\quad \cdot \left(1 - C \exp\left(-\frac{c\Lambda^2}{\sum_{\lambda_n \geq N} \lambda_n^{d-1} |c_n|^2}\right)\right). \end{aligned}$$

But we have  $\lim_{N \rightarrow +\infty} 1 - C \exp\left(-\frac{c\Lambda^2}{\sum_{\lambda_n \geq N} \lambda_n^{d-1} |c_n|^2}\right) = 1$ , thus there is  $N \in \mathbb{N}^*$  such

$$\text{that } 1 - C \exp\left(-\frac{c\Lambda^2}{\sum_{\lambda_n \geq N} \lambda_n^{d-1} |c_n|^2}\right) \geq 1/2. \quad \square$$

Therefore, to prove (9.4), it is sufficient to prove the following proposition:

**Lemma 9.3.** — *For all  $\Lambda > 0$  and  $N \in \mathbb{N}^*$ ,*

$$P\left(\omega \in \Omega : \|[u_0^\omega]_N\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \Lambda, \|e^{-itH}[u_0^\omega]_N\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{7}, \infty}(\mathbb{R}^d))} \leq \Lambda\right) > 0.$$

*Proof.* — We get

$$\begin{aligned}
 P\left(\omega \in \Omega : \left\| [u_0^\omega]_N \right\|_{\mathcal{H}^{(d-1)/2}(\mathbb{R}^d)} \leq \Lambda, \left\| e^{-itH} [u_0^\omega]_N \right\|_{L^{2p}([- \pi, \pi]; \mathcal{W}^{\frac{1}{p}, \infty}(\mathbb{R}^d))} \leq \Lambda \right) &\geq \\
 &\geq P\left(\omega \in \Omega : \sum_{\lambda_n < N} \lambda_n^{d-1} |c_n|^2 |g_n(\omega)|^2 \leq \frac{C\Lambda^2}{N^{2d}}\right) \\
 &\geq P\left(\omega \in \Omega : \sum_{\lambda_n < N} |g_n(\omega)|^2 \leq \frac{C\Lambda^2}{N^{2d} \|u_0\|_{\mathcal{H}^{d-1}(\mathbb{R}^d)}^2}\right) \\
 &\geq P\left(\bigcap_{\lambda_n < N} \left\{ \omega \in \Omega : |g_n(\omega)|^2 \leq \frac{C\Lambda^2}{N^{4d} \|u_0\|_{\mathcal{H}^{d-1}(\mathbb{R}^d)}^2} \right\}\right) \\
 &= \prod_{\lambda_n < N} P\left(\omega \in \Omega : |g_n(\omega)|^2 \leq \frac{C\Lambda^2}{N^{4d} \|u_0\|_{\mathcal{H}^{d-1}(\mathbb{R}^d)}^2}\right) > 0.
 \end{aligned}$$

□

### Appendix A. Tools of pseudo-differential operators and applications to eigenfunctions

For  $m \in \mathbb{R}$ , we define  $T^m$  as the vector space of symbols  $q(x, \xi) \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$  such that for all  $\alpha \in \mathbb{N}^d$  and  $\beta \in \mathbb{N}^d$ , there exists a constant  $C_{\alpha, \beta} > 0$  such that for all  $(x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d$ , we have

$$|\partial_x^\alpha \partial_\xi^\beta q(x, \xi)| \leq C_{\alpha, \beta} (1 + |x| + |\xi|)^{m-\beta}.$$

Similarly, let  $S^m$  be the vector space of symbols satisfying

$$|\partial_x^\alpha \partial_\xi^\beta q(x, \xi)| \leq C_{\alpha, \beta} (1 + |\xi|)^{m-\beta}.$$

For  $q \in S^m \cup T^m$  and  $h > 0$ , let  $Op_h(q)$  be the operator defined by

$$\begin{aligned}
 Op_h(q)f(x) &= (2\pi h)^{-d} \int_{\mathbb{R}^d \times \mathbb{R}^d} e^{i(x-y)\cdot\xi/h} q(x, \xi) f(y) dy d\xi \\
 &= (2\pi h)^{-d} \int_{\mathbb{R}^d} e^{ix\cdot\xi/h} q(x, \xi) \hat{f}(\xi/h) d\xi \\
 &= (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix\cdot\xi} q(x, h\xi) \hat{f}(\xi) d\xi.
 \end{aligned}$$

Let  $p(x, \xi) = |\xi|^2 + |x|^2 - 1$  and define  $H_h = Op_h(p) \in Op_h(T^2)$ .

**A.1. Microlocal analysis for the harmonic oscillator.** — In [53], we can then obtain the following two results:

**Theorem A.1.** — *If  $q_1 \in S^{m_1}$  (respectively  $T^{m_1}$ ) and  $q_2 \in S^{m_2}$  (respectively  $T^{m_2}$ ) then there exists a symbol  $q \in S^{m_1+m_2}$  (respectively  $T^{m_1+m_2}$ ) such that*

$$Op_h(q_1) \circ Op_h(q_2) = Op_h(q)$$

with

$$q = \sum_{|\alpha| \leq N} \frac{h^{|\alpha|}}{i^{|\alpha|}} \partial_\xi^\alpha q_1 \partial_x^\alpha q_2 + h^{N+1} r_N$$

where  $r_N \in S^{m_1+m_2-(N+1)}$  (respectively  $T^{m_1+m_2-(N+1)}$ ).

**Theorem A.2.** — *If  $q(x, \xi) \in S^0$  then for any  $s \in \mathbb{R}$ , there exists a constant  $C > 0$  such that for any  $h \in ]0, 1]$  and any  $u \in H^s(\mathbb{R}^d)$ ,*

$$\|Op_h(q)u\|_{H^s(\mathbb{R}^d)} \leq C \|u\|_{H^s(\mathbb{R}^d)}.$$

We then state the following property which will allow us to invert the harmonic oscillator modulo a very regular remainder term.

Let  $\delta > 0$  and define a function  $\eta \in C^\infty(\mathbb{R}^d)$  such that

$$\eta(x) = \begin{cases} 0 & \text{if } |x| \leq 1 + \delta, \\ 1 & \text{if } |x| \geq 1 + 2\delta. \end{cases}$$

**Proposition A.3.** — *For any  $N \in \mathbb{N}^*$ , there exist two pseudo-differential operators  $E_N \in Op_h(T^{-2})$  and  $R_N \in Op_h(T^{-(N+1)})$  such that*

$$E_N \circ H_h = \eta + h^{N+1} R_N.$$

*Proof.* — Recall that  $p(x, \xi) = |\xi|^2 + |x|^2 - 1$  and set

$$f_0 = \frac{\eta}{p} \in T^{-2}.$$

For  $n \geq 1$ , we define  $f_n$  by induction in the following way:

$$f_n = -\frac{1}{p} \sum_{|\alpha|+j=n, j \neq n} \frac{1}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p \in T^{-2-n}.$$

Finally we set

$$E_N = Op_h\left(\sum_{0 \leq j \leq N} h^j f_j\right).$$

Then, by Proposition A.1,

$$\begin{aligned}
 E_N \circ H_h &= Op_h \left( \sum_{0 \leq j \leq N} h^j f_j \right) \circ Op_h(p) \\
 &= Op_h \left( \sum_{|\alpha|+j \leq N} \frac{h^{|\alpha|+j}}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p + h^{N+1} r_N \right) \\
 &= Op_h \left( \sum_{|\alpha|+j \leq N} \frac{h^{|\alpha|+j}}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p \right) + h^{N+1} R_N
 \end{aligned}$$

with  $R_N = Op_h(r_N) \in Op_h(T^{-(N+1)})$ . Therefore

$$\begin{aligned}
 \sum_{|\alpha|+j \leq N} \frac{h^{|\alpha|+j}}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p &= f_0 p + \sum_{1 \leq \ell \leq N} \sum_{|\alpha|+j=\ell} \frac{h^\ell}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p \\
 &= \eta + \sum_{1 \leq \ell \leq N} h^\ell \left( \sum_{|\alpha|+j=\ell, j \neq \ell} \frac{1}{i^{|\alpha|}} \partial_\xi^\alpha f_j \partial_x^\alpha p + f_{\ell \cdot} p \right) \\
 &= \eta,
 \end{aligned}$$

which was the claim.  $\square$

**A.2. A localisation property of the eigenfunctions of the harmonic oscillator in dimension  $d \geq 1$ .** — We can now establish a localisation property of the eigenfunctions of the harmonic oscillator.

**Proposition A.4.** — *Let  $d \geq 1$ . For all integers  $K, N \geq 1$ , all  $c > 1$  and all  $1 \leq p \leq +\infty$ , there exists a constant  $C > 0$  such that for all  $n \geq 0$ ,*

$$\| \langle x \rangle^K h_n \|_{L^p(|x| \geq c\lambda_n)} \leq C \lambda_n^{-N}.$$

*Proof.* — By definition, we have  $(-\Delta + |x|^2 - \lambda_n^2)h_n = 0$ . Therefore, if we set  $h = \frac{1}{\lambda_n^2}$  and  $\Phi(x) = h_n(\lambda_n x)$  then

$$(-h^2 \Delta + |x|^2 - 1)\Phi = 0.$$

Let  $\delta \ll 1$  and set  $\chi \in C^\infty(\mathbb{R}^d)$  such that

$$\chi(x) = \begin{cases} 0 & \text{if } |x| \leq 1, \\ 1 & \text{if } |x| \geq 1 + \delta, \end{cases}$$

and set also  $\tilde{\chi} \in C^\infty(\mathbb{R}^d)$  such that

$$\tilde{\chi}(x) = \begin{cases} 0 & \text{if } |x| \leq 1 + 2\delta, \\ 1 & \text{if } |x| \geq 1 + 3\delta. \end{cases}$$



Then

$$H_h(\chi\Phi) = -h^2\Phi\Delta\chi - 2h^2\nabla\chi \cdot \nabla\Phi.$$

Next, thanks to Proposition A.3, we obtain

$$\eta\chi\Phi = -E_N(h^2\Phi\Delta\chi + 2h^2\nabla\chi \cdot \nabla\Phi) - h^{N+1}R_N(\chi\Phi).$$

And finally

$$\langle x \rangle^K \tilde{\chi}\eta\chi\Phi = -\langle x \rangle^K \tilde{\chi}E_N(h^2\Phi\Delta\chi + 2h^2\nabla\chi \cdot \nabla\Phi) - h^{N+1}\langle x \rangle^K \tilde{\chi}R_N(\chi\Phi).$$

- Estimate of  $\langle x \rangle^K \tilde{\chi}E_N(\Phi\Delta\chi)$ : We have

$$\tilde{\chi}(x)E_N(\Phi\Delta\chi)(x) = \frac{\tilde{\chi}(x)}{(2\pi h)^d} \int_{\xi, 1 \leq |y| \leq 1+\delta} e^{i(x-y)\cdot\xi/h} E_N(x, \xi)(\Delta\chi\Phi)(y) dy d\xi.$$

Since  $|x| > 1 + 2\delta$ , then  $|x - y| > \delta$ . Next, observe that we have

$$\int_{|x-y|>\delta} \leq \int_{|x_1-y_1|>\delta} + \int_{|x_2-y_2|>\delta} + \cdots + \int_{|x_d-y_d|>\delta},$$

so that we are reduced to treat the term where  $|x_1 - y_1| > \delta$ .

From  $\frac{h^M}{i^M(x_1-y_1)^M} \partial_{\xi_1}^M e^{i(x-y)\cdot\xi/h} = e^{i(x-y)\cdot\xi/h}$  and from an integration by parts, we deduce

$$\begin{aligned} \langle x \rangle^K \tilde{\chi}(x)E_N(\Phi\Delta\chi)(x) &= \\ &= (-1)^M \frac{\tilde{\chi}(x)}{(2\pi h)^d} \int_{\substack{\xi, 1 \leq |y| \leq 1+\delta \\ |x_1-y_1|>\delta}} \frac{h^M e^{i(x-y)\cdot\xi/h} \langle x \rangle^K \partial_{\xi_1}^M E_N(x, \xi)(\Phi\Delta\chi)(y)}{i^M(x_1 - y_1)^M} dy d\xi. \end{aligned}$$

Therefore, as  $E_N \in T^{-2}$ , we obtain

$$|\langle x \rangle^K \tilde{\chi}(x)E_N(\Phi\Delta\chi)(x)| \leq Ch^{M-d} |\tilde{\chi}(x)| \int_{\xi, 1 \leq |y| \leq 1+\delta} \frac{|(\Phi\Delta\chi)(y)|}{(1 + |x| + |\xi|)^{2+M-K}} d\xi dy.$$

We can then assume that  $M \geq 1$  is such that  $M > 2d + K - 1$ , and therefore

$$\begin{aligned} |\langle x \rangle^K \tilde{\chi}(x)E_N(\Phi\Delta\chi)(x)| &\leq \\ &\leq Ch^{M-d} \|\Delta\chi\Phi\|_{L^1(\mathbb{R}^d)} \frac{|\tilde{\chi}(x)|}{(1 + |x|)^{1+M-K-d}} \int_{\mathbb{R}^d} \frac{d\xi}{(1 + |\xi|)^{d+1}} \\ &\leq Ch^{M-d} \|\Phi\|_{L^2(\mathbb{R}^d)} \frac{1}{(1 + |x|)^{1+M-K-d}}. \end{aligned}$$

And finally, for all integers  $M, K \geq 1$ , there exists a constant  $C > 0$  such that for all  $h \in ]0, 1]$ ,

$$\|\langle x \rangle^K \tilde{\chi} E_N(\Phi \Delta \chi)\|_{L^p(\mathbb{R}^d)} \leq Ch^{M-d} \|\Phi\|_{L^2(\mathbb{R}^d)}.$$

- Estimate of  $\langle x \rangle^K \tilde{\chi} E_N(\nabla \chi \cdot \nabla \Phi)$ : Since  $\Phi$  satisfies  $(-h^2 \Delta + |x|^2)\Phi = \Phi$ , then  $h \|\nabla \Phi\|_{L^2(\mathbb{R}^d)} \leq \|\Phi\|_{L^2(\mathbb{R}^d)}$ . As a consequence we can proceed as the first term to get the same kind of estimate.

- Estimate of  $h^{N+1} \langle x \rangle^K \tilde{\chi} R_N(\chi \Phi)$ : We have

$$h^{N+1} \langle x \rangle^K \tilde{\chi}(x) R_N(\chi \Phi)(x) = h^{N+1} \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{i(x-y) \cdot \xi} \langle x \rangle^K \tilde{\chi}(x) r_N(x, h\xi) \mathcal{F}(\chi \Phi)(\xi) d\xi$$

with

$$\langle x \rangle^K \tilde{\chi}(x) r_N(x, \xi) \in T^{-N-1+K} \subset T^0 \subset S^0 \quad \text{for } N \geq K - 1.$$

Using Theorem A.2 and the Sobolev embeddings, we obtain

$$\begin{aligned} \|h^{N+1} \langle x \rangle^K \tilde{\chi} R_N(\chi \Phi)\|_{L^p(\mathbb{R}^d)} &\leq h^{N+1} \|\langle x \rangle^K \tilde{\chi} R_N(\chi \Phi)\|_{H^{d/2+1}(\mathbb{R}^d)} \\ &\leq Ch^{N+1} \|\chi \Phi\|_{H^{d/2+1}(\mathbb{R}^d)} \\ &\leq Ch^{N+1} \|\Phi\|_{H^{d/2+1}(\mathbb{R}^d)}. \end{aligned}$$

Finally, we obtain that for all integers  $K, N \geq 1$ , for all  $p \in [1, +\infty]$  and  $c > 1$ , that there exists a constant  $C > 0$  such that for all  $0 < h \leq 1$ , we have

$$\|\langle x \rangle^K \Phi\|_{L^p(|x| \geq c)} \leq Ch^N \|\Phi\|_{H^{d/2+1}(\mathbb{R}^d)}.$$

Then, returning to the initial variable, we get that for all integers  $K, N \geq 1$ , for all  $p \in [1, +\infty]$  and  $c > 1$ , there exists a constant  $C > 0$  such that for all  $0 < h \leq 1$  and  $n \in \mathbb{N}$  with  $h = \frac{1}{\lambda_n^2}$ , we have

$$\begin{aligned} \|\langle \sqrt{h}x \rangle^K h_n\|_{L^p(|x| \geq c\lambda_n)} &\leq Ch^{N-d/(2p)-d/4-1/2} \|h_n\|_{H^{d/2+1}(\mathbb{R}^d)} \\ &\leq Ch^{N-d/(2p)-d/4-1/2} \|h_n\|_{\mathcal{H}^{d/2+1}(\mathbb{R}^d)} \\ &\leq Ch^{N-d/(2p)-d/2-1} \|h_n\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

But

$$\begin{aligned} \|\langle x \rangle^K h_n\|_{L^p(|x| \geq c\lambda_n)} &\leq \|h_n\|_{L^p(|x| \geq c\lambda_n)} + h^{-K/2} \|\langle \sqrt{h}x \rangle^K h_n\|_{L^p(|x| \geq c\lambda_n)} \\ &\leq Ch^{N-d/(2p)-d/2-1-K/2} \|h_n\|_{L^2(\mathbb{R}^d)} \\ &\leq C\lambda_n^{-2N+d/p+d+K+2} \|h_n\|_{L^2(\mathbb{R}^d)} \end{aligned}$$

which concludes the proof. □

**A.3. Functional and pseudo-differential calculus.** — The following property of functional calculus explains that some operators can be approximated by pseudo-differential operators.

**Proposition A.5.** — *Let  $\Phi \in C_0^\infty(\mathbb{R})$  and  $\chi_2 \in C_0^\infty(\mathbb{R}^d)$  with  $\chi_2(x) = 1$  for  $x \in B(0, 1^+)$ . Then for all  $N \in \mathbb{N}^*$  and  $s \geq 0$ , there is a constant  $C_{N,s} > 0$  such that for all  $h \in ]0, 1]$  and  $u \in L^2(\mathbb{R}^d)$ ,*

$$\|\Phi(|x|^2 + |h\nabla|^2)u - \sum_{j=0}^{N-1} h^j Op_h(\Psi_j)\chi_2 u\|_{H^s(\mathbb{R}^d)} \leq C_{N,s} h^{N-s} \|u\|_{L^2(\mathbb{R}^d)},$$

where  $\Psi_0(x, \xi) = \Phi(|x|^2 + |\xi|^2)$ ,  $Supp(\Psi_j) \subset \{(x, \xi) : |x|^2 + |\xi|^2 \in Supp(\Phi)\}$  and  $\Psi_j \in T^{-j} \subset S^0$ .

*Proof.* — We use [16, Proposition 2.1] (notice that *stricto sensu* [16, Proposition 2.1] is only for a semi-classical Laplace operator, but the proof given applies also here). If  $\chi_1 \in C_0^\infty(\mathbb{R}^d)$  is such that  $\chi_1 \chi_2 = \chi_1$  then

$$\|\Phi(|x|^2 + |h\nabla|^2)\chi_1 u - \sum_{j=0}^{N-1} h^j Op_h(\Psi_j(x, \xi))\chi_2 u\|_{H^s(\mathbb{R}^d)} \leq C_{N,s} h^{N-s} \|u\|_{L^2(\mathbb{R}^d)},$$

with  $\Psi_0(x, \xi) = \Phi(|x|^2 + |\xi|^2)$ ,  $Supp(\Psi_j) \subset \{(x, \xi) : |x|^2 + |\xi|^2 \in Supp(\Phi)\}$  and  $\Psi_j \in T^{-j}$ . Then, it is enough to show that

$$\|\Phi(|x|^2 + |h\nabla|^2)(1 - \chi_1)u\|_{H^s(\mathbb{R}^d)} = \mathcal{O}(h^\infty) \|u\|_{L^2(\mathbb{R}^d)}.$$

We choose  $\chi_1 = 1$  on  $B(0, R)$ ,  $R > 1$ , and the result then follows from Proposition A.4.  $\square$

**A.4. A continuity result.** — Thanks to the functional calculus, it is clear that  $\|\chi(\frac{H}{N^2})u\|_{\mathcal{H}^s(\mathbb{R}^d)} \leq C \|u\|_{\mathcal{H}^s(\mathbb{R}^d)}$ , however the continuity result stated in the next proposition is not straightforward and relies on Proposition A.5.

**Proposition A.6.** — *For any  $s \geq 0$ , there exists a constant  $C > 0$  such that for any  $N \in \mathbb{N}^*$  and any function  $u \in H^s(\mathbb{R}^d)$ ,*

$$\|\chi\left(\frac{H}{N^2}\right)u\|_{H^s(\mathbb{R}^d)} \leq C \|u\|_{H^s(\mathbb{R}^d)}.$$

*Proof.* — Using (3.7) and (3.8), it is sufficient to show that:  $\forall s \geq 0, \exists C > 0$  and  $h_0$  such that  $\forall 0 < h \leq h_0, \forall u \in H^s(\mathbb{R}^d)$

$$(A.1) \quad \|\chi(|x|^2 + |h\nabla|^2)u\|_{H^s(\mathbb{R}^d)} \leq C \|u\|_{H^s(\mathbb{R}^d)}.$$

Indeed, recalling that  $\mathbf{u}(x) = u(\frac{x}{\sqrt{h}})$ , then

$$\begin{aligned}
 \|\chi\left(\frac{H}{N^2}\right)u\|_{H^s(\mathbb{R}^d)} &\leq \\
 &\leq \|[\chi(|x|^2 + |h\nabla|^2)]\mathbf{u}(\sqrt{h}\cdot)\|_{H^s(\mathbb{R}^d)} \\
 &\leq \|[\chi(|x|^2 + |h\nabla|^2)]\mathbf{u}(\sqrt{h}\cdot)\|_{L^2(\mathbb{R}^d)} + \|\sqrt{-\Delta}^s[\chi(|x|^2 + |h\nabla|^2)]\mathbf{u}(\sqrt{h}\cdot)\|_{L^2(\mathbb{R}^d)} \\
 &\leq h^{-d/4}\|u\|_{L^2(\mathbb{R}^d)} + h^{s/2-d/4}\|[\chi(|x|^2 + |h\nabla|^2)]\mathbf{u}\|_{H^s(\mathbb{R}^d)} \\
 &\leq \|u\|_{L^2(\mathbb{R}^d)} + h^{s/2-d/4}\|u\|_{H^s(\mathbb{R}^d)} \\
 &\leq \|u\|_{H^s(\mathbb{R}^d)}.
 \end{aligned}$$

By interpolation, we can limit the proof to the case where  $s$  is an integer. Thanks to Proposition A.5 (with  $N = s$ ), there exists a constant  $C > 0$  such that for any  $h \in ]0, 1]$  and any function  $u \in L^2(\mathbb{R}^d)$ , we have

$$\|\chi(|x|^2 + |h\nabla|^2)u - \sum_{j=0}^N h^j Op_h(\Psi_j)\chi_2 u\|_{H^s(\mathbb{R}^d)} \leq C\|u\|_{L^2(\mathbb{R}^d)},$$

with  $Supp(\Psi_j(x, \xi)) \subset \{(x, \xi) : |x|^2 + |\xi|^2 \in Supp(\chi)\}$ .

Thus, to obtain (A.1), it suffices to obtain that for any  $s \geq 0$ , there exist two constants  $C > 0$  and  $h_0 \geq 1$  such that for any  $h \in ]0, h_0]$  and any function  $u \in H^s(\mathbb{R}^d)$ ,

$$(A.2) \quad \|Op_h(\Psi_j)u\|_{H^s(\mathbb{R}^d)} \leq C\|u\|_{H^s(\mathbb{R}^d)}.$$

Finally, to establish (A.2), it is enough to use Theorem A.2 and to notice that

$$(x, \xi) \longrightarrow \chi(|x|^2 + |\xi|^2) \in S^0 \quad \text{and} \quad (x, \xi) \longrightarrow \Psi_j(x, \xi) \in S^0.$$

□

## References

- [1] A. Bényi, T. Oh, and O. Pocovnicu. Wiener randomization on unbounded domains and an application to almost sure well-posedness of NLS. Excursions in harmonic analysis, Volume 4. The February Fourier talks at the Norbert Wiener Center, College Park, MD, USA, 2002-2013. Cham: Birkhäuser/Springer. Applied and Numerical Harmonic Analysis, 3–25 (2015).
- [2] A. Bényi, T. Oh, and O. Pocovnicu. On the probabilistic Cauchy theory of the cubic nonlinear Schrödinger equation on  $\mathbb{R}^d$ ,  $d \geq 3$ . *Trans. Am. Math. Soc.*, Ser. B 2, 1–50 (2015).

- [3] A. Bényi, T. Oh, and O. Pocovnicu. On the probabilistic Cauchy theory for nonlinear dispersive PDEs. Landscapes of time-frequency analysis. Based on talks given at the inaugural conference on aspects of time-frequency analysis, Turin, Italy, July 5-7, 2018. Cham: Birkhäuser. *Appl. Numer. Harmon. Anal.*, 1–32 (2019).
- [4] J.-M. Bouclet. Distributions spectrales pour des opérateurs perturbés. PhD thesis, Université de Nantes, 2000.
- [5] J. Bourgain. Periodic nonlinear Schrödinger equation and invariant measures. *Comm. Math. Phys.* 166 (1994), no. 1, 1–26.
- [6] J. Bourgain. Invariant measures for the 2D-defocusing nonlinear Schrödinger equation. *Comm. Math. Phys.* 176 (1996), no. 2, 421–445.
- [7] J. Bourgain. Refinements of Strichartz’ inequality and applications to 2D-NLS with critical nonlinearity. *Internat. Math. Res. Notices* 1998, no. 5, 253–283.
- [8] J. Bourgain and A. Bulut. Almost sure global well posedness for the radial nonlinear Schrödinger equation on the unit ball I: the 2D case. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 31 (2014), no. 6, 1267–1288.
- [9] J. Bourgain and A. Bulut. Almost sure global well-posedness for the radial nonlinear Schrödinger equation on the unit ball II: the 3d case. *J. Eur. Math. Soc. (JEMS)* 16 (2014), no. 6, 1289–1325.
- [10] J. Bourgain and A. Bulut. Invariant Gibbs measure evolution for the radial nonlinear wave equation on the 3d ball. *J. Funct. Anal.* 266 (2014), no. 4, 2319–2340.
- [11] B. Bringmann. Almost-sure scattering for the radial energy-critical nonlinear wave equation in three dimensions. *Anal. PDE* 13 (2020), no. 4, 1011–1050.
- [12] B. Bringmann. Almost sure scattering for the energy critical nonlinear wave equation. *Amer. J. Math.* 143 (2021), no. 6, 1931–1982.
- [13] B. Bringmann. Almost sure local well-posedness for a derivative nonlinear wave equation. *Int. Math. Res. Not. IMRN* 2021, no. 11, 8657–8697.
- [14] N. Burq. Mesures semi-classiques et mesures de défaut. (French) [Semiclassical measures and defect measures] *Séminaire Bourbaki, Vol. 1996/97. Astérisque* 245 (1997), Exp. No. 826, 4, 167–195.
- [15] N. Burq and G. Lebeau. Injections de Sobolev probabilistes et applications. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(6):917–962, 2013.
- [16] N. Burq, P. Gérard, and N. Tzvetkov. Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds. *Amer. J. Math.* 126 (2004), no. 3, 569–605.
- [17] N. Burq, P. Gérard, and N. Tzvetkov. Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces. *Invent. Math.* 159 (2005), no. 1, 187–223.
- [18] N. Burq, P. Gérard, and N. Tzvetkov. Multilinear eigenfunction estimates and global existence for the three dimensional nonlinear Schrödinger equations. *Ann. Sci. École Norm. Sup. (4)* 38 (2005), no. 2, 255–301.
- [19] N. Burq and L. Thomann. Almost sure scattering for the one dimensional nonlinear Schrödinger equation. *Mem. Amer. Math. Soc.*, to appear.

- [20] N. Burq, L. Thomann, and N. Tzvetkov. Long time dynamics for the one dimensional non linear Schrödinger equation. *Ann. Inst. Fourier (Grenoble)* 63 (2013), no. 6, 2137–2198.
- [21] N. Burq and N. Tzvetkov. Random data Cauchy theory for supercritical wave equations. I. Local theory. *Invent. Math.* 173 (2008), no. 3, 449–475.
- [22] N. Burq and N. Tzvetkov. Random data Cauchy theory for supercritical wave equations. II. A global existence result. *Invent. Math.* 173 (2008), no. 3, 477–496.
- [23] N. Burq and N. Tzvetkov. Probabilistic well-posedness for the cubic wave equation. *J. Eur. Math. Soc. (JEMS)* 16 (2014), no. 1, 1–30.
- [24] N. Camps. Scattering for the cubic Schrödinger equation in 3D with randomized radial initial data. *Trans. Amer. Math. Soc.* 376 (2023), no. 1, 285–333.
- [25] R. Carles. Critical nonlinear Schrödinger equations with and without harmonic potential. *Math. Models Methods Appl. Sci.* 12 (2002), no. 10, 1513–1523.
- [26] T. Cazenave and F. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in  $H^s$ . *Nonlinear Anal., Theory Methods Appl.* 14, No. 10, 807–836 (1990).
- [27] M. Christ, J. Colliander, and T. Tao. Ill-posedness for nonlinear Schrödinger and wave equations. *Preprint : arXiv : 0311048*.
- [28] J.-Y. Chemin and I. Gallagher. Wellposedness and stability results for the Navier-Stokes equations in  $\mathbb{R}^3$ . *Ann. Inst. H. Poincaré Anal. Non Linéaire* 26 (2009), no. 2, 599–624.
- [29] J.-Y. Chemin and I. Gallagher. Large, global solutions to the Navier-Stokes equations, slowly varying in one direction. *Trans. Amer. Math. Soc.* 362 (2010), no. 6, 2859–2873.
- [30] J.-Y. Chemin, I. Gallagher, and M. Paicu. Global regularity for some classes of large solutions to the Navier-Stokes equations. *Ann. of Math. (2)* 173 (2011), no. 2, 983–1012.
- [31] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on  $\mathbb{R}^3$ . *Comm. Pure Appl. Math.* 57 (2004), no. 8, 987–1014.
- [32] J. Colliander and T. Oh. Almost sure well-posedness of the cubic nonlinear Schrödinger equation below  $L^2(\mathbb{T})$ . *Duke Math. J.* 161 (2012), no. 3, 367–414.
- [33] P. Constantin and J.-C. Saut. Local smoothing properties of Schrödinger equations. *Indiana Univ. Math. J.* 38, No. 3, 791–810 (1989).
- [34] A.-S. de Suzzoni. Invariant measure for the cubic wave equation on the unit ball of  $R^3$ . *Dyn. Partial Differ. Equ.* 8 (2011), no. 2, 127–47.
- [35] A.-S. de Suzzoni. Large data low regularity scattering results for the wave equation on the Euclidean space. *Comm. Partial Differential Equations* 38 (2013), no. 1, 1–49.
- [36] Y. Deng, N. Tzvetkov, and N. Visciglia. Invariant measures and long time behaviour for the Benjamin-Ono equation III. *Comm. Math. Phys.* 339 (2015), no. 3, 815–857.
- [37] Y. Deng. Two-dimensional nonlinear Schrödinger equation with random radial data. *Anal. PDE* 5 (2012), no. 5, 913–960.
- [38] Y. Deng. Invariance of the Gibbs measure for the Benjamin-Ono equation. *J. Eur. Math. Soc. (JEMS)* 17 (2015), no. 5, 1107–1198.

- [39] Y. Deng, A. Nahmod, and H. Yue. Invariant Gibbs measures and global strong solutions for nonlinear Schrödinger equations in dimension two. *Preprint : arXiv: 1910.98492*.
- [40] Y. Deng, A. Nahmod, and H. Yue. Random tensors, propagation of randomness, and nonlinear dispersive equations. *Invent. Math.* (2021), no. 2, 539–686.
- [41] S. A. Denisov. Infinite superlinear growth of the gradient for the two-dimensional Euler equation. *Discrete Contin. Dyn. Syst.* 23 (2009), no. 3, 755–764.
- [42] B. Dodson, J. Lührmann, and D. Mendelson. Almost sure local well-posedness and scattering for the 4D cubic nonlinear Schrödinger equation. *Adv. Math.* 347, 619–676 (2019).
- [43] S.-I. Doi. Smoothing effects of Schrödinger evolution groups on Riemannian manifolds. *Duke Math. J.* 82, No. 3, 679–706 (1996).
- [44] S.-I. Doi. Commutator algebra and abstract smoothing effect. *J. Funct. Anal.* 168, No. 2, 428–469 (1999).
- [45] J. Dziubanski and P. Glowacki. Sobolev spaces related to Schrödinger operators with polynomial potentials. *Math. Z.* 262 (2009), no. 4, 881–894.
- [46] J. Ginibre. Le problème de Cauchy pour des EDP semi-linéaires périodiques en variables d’espace (d’après Bourgain). (French) [The Cauchy problem for periodic semilinear PDE in space variables (after Bourgain)] Séminaire Bourbaki, Vol. 1994/95. *Astérisque* No. 237 (1996), Exp. No. 796, 4, 163–187.
- [47] J. Ginibre and G. Velo. On a class of nonlinear Schrödinger equations. I. The Cauchy problem, general case. *J. Funct. Anal.* 32, 1–32 (1979).
- [48] L. Hörmander. The analysis of linear partial differential operators. III. Pseudo-differential operators. Reprint of the 1994 edition. *Classics in Mathematics*. Springer, Berlin, 2007. viii+525 pp.
- [49] T. Kato. On the Cauchy problem for the (generalized) Korteweg-de Vries equation. *Adv. Math., Suppl. Stud.* 8, 93–128 (1983).
- [50] R. Killip, J. Murphy, and M. Visan. Almost sure scattering for the energy-critical NLS with radial data below  $H^1(\mathbb{R}^4)$ . *Commun. Partial Differ. Equations* 44, No. 1, 51–71 (2019).
- [51] H. Koch and D. Tataru.  $L^p$  eigenfunction bounds for the Hermite operator. *Duke Math. J.* 128 (2005), no. 2, 369–392.
- [52] M. Latocca. Almost sure scattering at mass regularity for radial Schrödinger equations. *Nonlinearity*, 35 (2022), no.10.
- [53] A. Martinez. An introduction to semiclassical and microlocal analysis. Universitext. Springer-Verlag, New York, 2002. viii+190 pp.
- [54] A. Nahmod, T. Oh, Tadahiro, L. Rey-Bellet, and G. Staffilani. Invariant weighted Wiener measures and almost sure global well-posedness for the periodic derivative NLS. *J. Eur. Math. Soc. (JEMS)* 14 (2012), no. 4, 1275–1330.
- [55] A. Nahmod and G. Staffilani. Almost sure well-posedness for the periodic 3D quintic nonlinear Schrödinger equation below the energy space. *J. Eur. Math. Soc. (JEMS)* 17 (2015), no. 7, 1687–1759.

- [56] U. Niederer. Maximal kinematical invariance group of the harmonic oscillator. *Helv. Phys. Acta*, v. 46 (1973), no. 2, 191–200.
- [57] T. Oh, O. Pocovnicu, and N. Tzvetkov. Probabilistic local Cauchy theory of the cubic nonlinear wave equation in negative Sobolev spaces. *Ann. Inst. Fourier (Grenoble)* 72 (2022), no. 2, 771–830.
- [58] T. Oh, N. Tzvetkov, and Y. Wang. Solving the 4NLS with white noise initial data. *Forum Math. Sigma* 8 (2020), Paper No. e48, 63 pp.
- [59] F. Planchon, N. Tzvetkov, and N. Visciglia. Growth of Sobolev norms for 2d NLS with harmonic potential. *Rev. Mat. Iberoam.*, (2022). DOI 10.4171/RMI/1371.
- [60] A. Poiret. Solutions globales pour des équations de Schrödinger sur-critiques en toutes dimensions. *Preprint : arXiv:1207.3519*.
- [61] A. Poiret. Solutions globales pour l'équation de Schrödinger cubique en dimension 3. *Preprint : arXiv:1207.1578*.
- [62] A. Poiret, D. Robert, and L. Thomann. Probabilistic global well-posedness for the supercritical nonlinear harmonic oscillator. *Anal. PDE* 7 (2014), no. 4, 997–1026.
- [63] D. Robert. Autour de l'approximation semi-classique. (French) [On semiclassical approximation] Progress in Mathematics, 68. Birkhäuser Boston, Inc., Boston, MA, 1987. x+329 pp.
- [64] P. Sjölin. Regularity of solutions to the Schrödinger equation. *Duke Math. J.* 55, 699–715 (1987).
- [65] G. Staffilani. The theory of nonlinear Schrödinger equations : part1.
- [66] C. Sun and N. Tzvetkov. New examples of probabilistic well-posedness for nonlinear wave equations. *J. Funct. Anal.* (2020), no. 2, 108322, 47 pp.
- [67] T. Tao. Nonlinear dispersive equations. Local and global analysis. CBMS Regional Conference Series in Mathematics, 106. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2006. xvi+373 pp.
- [68] T. Tao. A pseudoconformal compactification of the nonlinear Schrödinger equation and applications. *New York J. Math.* 15 (2009), 265–282.
- [69] L. Thomann. Random data Cauchy problem for supercritical Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 26 (2009), no. 6, 2385–2402.
- [70] L. Thomann. A remark on the Schrödinger smoothing effect. *Asymptotic Anal.* 69, No. 1-2, 117–123 (2010).
- [71] L. Thomann and N. Tzvetkov. Gibbs measure for the periodic derivative nonlinear Schrödinger equation. *Nonlinearity* 23 (2010), no. 11, 2771–2791.
- [72] N. Tzvetkov. Invariant measures for the nonlinear Schrödinger equation on the disc. *Dyn. Partial Differ. Equ.* 3 (2006), no. 2, 111–160.
- [73] N. Tzvetkov. Invariant measures for the defocusing nonlinear Schrödinger equation. *Ann. Inst. Fourier (Grenoble)* 58 (2008), no. 7, 2543–2604.
- [74] N. Tzvetkov and N. Visciglia. Invariant measures and long-time behavior for the Benjamin-Ono equation. *Int. Math. Res. Not. IMRN* 2014, no. 17, 4679–4714.



- [75] N. Tzvetkov and N. Visciglia. Invariant measures and long time behaviour for the Benjamin-Ono equation II. *J. Math. Pures Appl.* (9) 103 (2015), no. 1, 102–141.
- [76] L. Vega. Schrödinger equations: Pointwise convergence to the initial data. *Proc. Am. Math. Soc.* 102, No. 4, 874–878 (1988).
- [77] K. Yajima and G. Zhang. Smoothing property for Schrödinger equations with potential superquadratic at infinity. *Comm. Math. Phys.* 221 (2001), no. 3, 573–590.
- [78] K. Yajima and G. Zhang. Local smoothing property and Strichartz inequality for Schrödinger equations with potentials superquadratic at infinity. *J. Differential Equations* 202 (2004), no. 1, 81–110.

---

NICOLAS BURQ, Laboratoire de Mathématiques d’Orsay, CNRS, Université Paris–Saclay, Bâtiment 307, F-91405 Orsay Cedex, and Institut Universitaire de France

*E-mail* : `nicolas.burq@math.u-psud.fr`

AURÉLIEN POIRET, Lycée Jeanne d’Albret, 6 rue Giraud Teulon, 78100 Saint-Germain-en-Laye • *E-mail* : `aurelien.poiret@gmail.com`

LAURENT THOMANN, Université de Lorraine, CNRS, IECL, F-54000 Nancy, France

*E-mail* : `laurent.thomann@univ-lorraine.fr`