



HAL
open science

The Structure of Plant–Pollinator Networks is Affected by Crop Type in a Highly Intensive Agricultural Landscape

Claire Gay, Sabrina Gaba, Vincent Bretagnolle

► **To cite this version:**

Claire Gay, Sabrina Gaba, Vincent Bretagnolle. The Structure of Plant–Pollinator Networks is Affected by Crop Type in a Highly Intensive Agricultural Landscape. SSRN Electronic Journal, 2023, pp.4378048. 10.2139/ssrn.4378048 . hal-04071662

HAL Id: hal-04071662

<https://hal.science/hal-04071662v1>

Submitted on 19 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The structure of plant–pollinator networks is affected by crop**
2 **type in a highly intensive agricultural landscape**

3

4

5 **Claire Gay^{a,b,*}, Sabrina Gaba^{a,b,c}, Vincent Bretagnolle^{a,b,c}**

6 ^a UMR 7372 Centre d'Etudes Biologiques de Chizé, CNRS & La Rochelle Université – 79360

7 Villiers-En-Bois (France)

8 ^b USC 1339 Centre d'Etudes Biologiques de Chizé, INRAE, CNRS & La Rochelle Université – 79360

9 Villiers-En-Bois (France)

10 ^c LTSER « Zone Atelier Plaine & Val de Sèvre » – 79360 Villiers-En-Bois (France)

11 *Author to whom correspondence should be addressed: claire.gay@cebc.cnrs.fr

12

13 **Abstract**

14 In agricultural landscapes, bipartite networks formed by pollinators and the flowers they
15 forage on, are characterised by the presence of species managed by humans, whether
16 honeybees (*Apis mellifera*) or crop plants. These managed species can affect the structural
17 properties of these networks because of spatial and temporal variation in the availability of
18 resources, and competition for these resources; for example, crop plants such as oilseed rape
19 and sunflower produce a large number of flowers during a short blooming period. Here, we
20 examined the structure of plant–pollinator networks in an intensive agricultural landscape, the
21 LTSER Zone Atelier Plaine & Val de Sèvre (West of France). We compiled a six-year
22 monitoring dataset of plant–pollinator interactions, sampling by sweep-nets along transects in
23 the main six crop types (811 fields in total). To describe networks, we computed six metrics:
24 connectance, nodes number, modularity, specialisation and two nestedness measures. Strong
25 differences between crops were observed in nestedness and specialisation: in oilseed rape,

26 networks were much more nested and much less specialised than in sunflower. In addition,
27 the link 'honeybee–crop flower' differed between the two mass-flowering crops. Honeybees
28 and sunflowers appeared as specialist species in sunflower crops, interacting strongly and
29 almost exclusively with each other, whereas they are usually considered highly generalist
30 species. Indeed, sunflower pollination may be almost entirely driven by honeybees,
31 conversely to oilseed rape crops, where the presence of wild bees and other insects tended to
32 produce a more diversified network.

33

34 **Key words**

35 Bipartite networks, Mass-flowering crops, Weeds, Honeybees, Spatio-temporal heterogeneity

36

37 **Highlights**

- 38 • Honeybees and crop flowers drive the network structure in agricultural environment
- 39 • Main mass-flowering crops got very distinct network properties
- 40 • Specialisation of managed species is highly variable among mass-flowering crops

41 **1. Introduction**

42 Insect biodiversity is declining worldwide, and numerous local insect extinctions have already
43 been documented (Habel et al., 2019). This decline has high consequences for the ecosystem
44 services that rely upon insects, not least pollination (Frankie et al., 2009). In temperate
45 regions, 88% of plants depend on zoogamous pollination for their reproduction (Ollerton et
46 al., 2011) and 70% of crops depend to some extent on insect pollination (Klein et al., 2007),
47 which improves yields of insect-pollinated crops and ensure agricultural production, as in
48 oilseed rape (OSR; *Brassica sp.*) and sunflower (SF; *Helianthus sp.*) (Perrot et al., 2019,
49 2018). In arable farming systems, insect pollinators forage partly – and sometimes mainly –
50 on mass-flowering crops, but availability of these crops does not cover the entire season of the
51 pollinators' activity (Odoux et al., 2014). Between OSR and SF blooms (the so-called 'dearth
52 period'), the resources provided by weeds are essential for maintaining pollinators, including
53 the honeybee *Apis mellifera* (Carvalho et al., 2011; Requier et al., 2015). Weeds and other
54 wild plants are therefore critical for pollinators (Bretagnolle and Gaba, 2015), and contribute
55 to their diversity and abundance at both field and landscape scales (Carvalho et al., 2011).
56 Among insect pollinators, bees and hoverflies contain generalist species that can interact with
57 many wild flowers (Dunn et al., 2020) or crop plants, and both improve pollination service
58 (Garibaldi et al., 2011). Hoverflies are more resilient to agricultural modifications than bees in
59 agricultural landscapes (Jauker et al., 2009); conversely, butterflies which also pollinate many
60 wild plants, have been dramatically affected by agricultural intensification (Johst et al., 2006).
61 Overall, weeds and pollinators are closely intertwined in agricultural farmland, and form
62 ecological interaction networks, defined by a set of nodes (species) interconnected by one or
63 more links (interactions). There is evidence that such networks have changed in structure over
64 the last century, due to pollinator and plant community changes, with many interactions
65 involving specialist species being lost (Mathiasson and Rehan, 2020), mostly as a result of

66 land use intensification (Ferreira et al., 2013). Indeed, the persistence of pollinators – together
67 with the services they provide – depends on stability, hence the structure of plant–pollinator
68 ecological networks.

69 However, plant–pollinator ecological networks in agricultural landscapes present unique
70 properties: first, they harbour managed and over-abundant species; second, they are highly
71 dynamic in space and time due to crop rotations. Such characteristics are expected to impact
72 the nature and structure of plant–pollinator networks. First, farmland ecological networks are
73 structured around managed honeybees in addition to wild pollinators, and managed crops in
74 addition to weeds (Geslin et al., 2017). The effect of mass-flowering crops on one hand, and
75 the presence of the honeybee on the other hand, can be related to ‘invasive species’ in arable
76 systems (Geslin et al., 2017; Stanley and Stout, 2014), i.e. super-generalist species that
77 interact with multiple groups of species (Giannini et al., 2015), leading to asymmetry and
78 nestedness of the network (Ferreira et al., 2013). Perhaps counterintuitively, this asymmetry
79 and nestedness contributes to the stability and the resistance of networks to land use change
80 (Memmott et al., 2004). Links that emerge from these super-abundant species may, however,
81 come at the expense of others, leading to species role shifts (Albrecht et al., 2014), or even
82 loss of species in both plants or pollinators, which should affect the structure and functioning
83 of networks (Valido et al., 2019). For instance, OSR flowers number c. 500,000/ha (Hoyle
84 and Cresswell, 2007), are brightly coloured and attractive, and produce both pollen and
85 nectar. OSR can thus create a distortion in mutualistic networks and has already been shown
86 to be related to the decline of specialist long-tongued bumblebees in favour of short-tongued
87 species better adapted to its flower traits (Diekötter et al., 2010). Honeybees in return may
88 decrease wild bee richness, leading to networks with fewer links (Lázaro et al., 2021).

89 However, a decrease or even a removal of supergeneralist (key) species may destabilise the
90 network (Kovács-Hostyánszki et al., 2019). Secondly, agricultural landscapes display very

91 high spatio-temporal unpredictability (due to crop rotations), and seasonal heterogeneity
92 (Rollin et al., 2016): honeybees are present from April to September (Odoux et al., 2014),
93 while OSR and SF, the main super-generalist plants, have very short flowering peaks
94 (Bretagnolle and Gaba, 2015), and wild bees show generally have a short presence within the
95 season. Crop rotation and spatial heterogeneity is therefore expected to affect the temporal
96 and spatial dynamics of plant–pollinator networks (Chakraborty et al., 2021).

97 There is currently a lack of descriptive information on plant–pollinator networks in farmland
98 habitats, especially at large spatial scale or over longer time-scales (Olesen et al., 2011).

99 Indeed, most studies investigated the crop compartment in relation to the semi-natural
100 elements or its plants (e.g. Magrath et al., 2018; Stanley and Stout, 2014), while few studies
101 aimed at investigating differences among crops (but see Kovács-Hostyánszki et al., 2019) and
102 even fewer the effect of crop diversity. This is unfortunate, since crop effect was found to be
103 the most statistically important effect in a vast study of predation–prey networks in
104 agricultural habitats (Ma et al., 2019). Indeed, descriptive information is critical to
105 understanding community stability and robustness, which are linked to the number of species
106 and the number of links between them (Carpentier et al., 2021), and therefore network
107 resilience, which relies on redundancy in the number of pollinator species per plant species
108 (Memmott et al., 2004) and network nested structure (Bascompte et al., 2003). Network
109 stability is improved by pollinator diversity (Bendel et al., 2019), nestedness and low
110 modularity, though possibly at the expense of persistence (Thébault and Fontaine, 2010).

111 Along environmental gradients from less- to more- impacted by human footprint, e.g. from
112 natural sites to monoculture farming, an increase in connectance and nestedness but a
113 decrease in the number of nodes as well as modularity were shown (Morrison et al., 2020).

114 Increased connectance is expected to decrease secondary extinctions (Blüthgen et al., 2007)

115 and improves robustness at the community level as well as stability (Thébaud and Fontaine,
116 2010).

117 In this study, we used a descriptive approach to qualify plant–pollinator networks in a highly
118 intensive farming system, at a large spatial scale (study site of 450km²) with data collected on
119 six different crop types (811 fields in total). Because our sampling effort was high, we first
120 paid attention to two generally overlooked patterns, the effect of sample size and the degree
121 distribution. Sampling effort has been shown to affect many network metrics, through
122 sampling intensity and sampling bias, but also underlying species abundances (Blüthgen et
123 al., 2007). Degree distribution is highly skewed, resulting in part from many ‘forbidden links’
124 (Jordano, 2016). The frequency distribution of links is claimed to be best described by power-
125 law distributions (Blüthgen, 2010), which have been shown to fit most natural networks. We
126 also describe the overall visual structure of networks by seasons and crop types. Then, we
127 investigated major features of the network, from simple network descriptors (number of
128 nodes, connectance) to more complex ones (nestedness, modularity, specialisation),
129 quantifying their level of variation among crop types and seasons, while accounting for
130 sample size. We expected plant–pollinator network structure to vary seasonally because of the
131 presence of mass-flowering crops, and the spillover or dilution effects that result from such
132 presence. Indeed, we expected strong variations in network structure among crop types, in
133 particular between mass-flowering and other crops, but also between arable crops and
134 grasslands. We further predicted a clear dominance of the managed honeybee, especially
135 during the flower blooms of OSR and SF, which dominate landscapes when flowering. In
136 particular, we predicted that networks would be more specialised during OSR and SF blooms
137 than in other periods and other crops. We also predicted that networks would be more nested
138 during the blooming periods of mass-flowering crops than in the dearth periods, and thus in

139 these crops – as a result of super-generalist and abundant species presence – more able to
140 include the same partners as the specialists in their spectrum of interactions.

141

142 **2. Materials and methods**

143 **2.1. Study site**

144 Our study site is the Long Term Socio-Ecological Research (LTSER) site ‘Zone Atelier
145 Plaine & Val de Sèvre’, located in the Nouvelle-Aquitaine region of southwest France, south
146 of the city of Niort (Bretagnolle et al., 2018). It covers 450 km², harbouring c. 13,000 fields.
147 We focused on six years of survey, 2015–2020, which is one of the largest time intervals of
148 plant–pollinator network studies (but see Chacoff et al., 2018; Petanidou et al., 2008; Ponisio
149 et al., 2017). Each year, 100 to 220 different fields were surveyed (some fields were studied
150 for more than one year) with a standardised protocol. Fields were selected in a stratified
151 random process (see Bretagnolle et al. (2018) for details). From 2015–2020, the number of
152 fields sampled per respective year were 174, 217, 195, 160, 117 and 115 (see ESM1a and 1b
153 in Appendix A). Crop surveyed included 7% corn (*Zea mays*) and sorghum (*Sorghum*
154 *bicolor*), 8% alfalfa (*Medicago sativa*), 12% meadows (temporary or permanent), 18% oilseed
155 rape (*Brassica napus*), 22% sunflower (*Helianthus annuus*), and 33% cereal crops. Cereals
156 included barley (*Hordeum vulgare*), durum wheat (*Triticum turgidum*), einkorn (*Triticum*
157 *monococcum*), oat (*Avena sativa*), rye (*Secale cereale*), triticale (\times *Triticosecale*) and winter
158 wheat (*Triticum aestivum*). Sampling was performed from April to August, i.e. all crops were
159 surveyed at least once during their respective flowering periods: OSR from April–May;
160 sunflower from July–August; alfalfa, meadows and cereals from May–June to the end of
161 August; and corn and sorghum from July–August.

162

163 **2.2. Sampling of plant–pollinator interactions**

164 Insects surveyed included bees (both honeybees and wild bees), butterflies and hoverflies
165 (Syrphidae), i.e. the three main pollinator groups in the LTSER. This study site hosts a large
166 number of pollinator species (Rollin et al., 2016). Each year, two (2015) or three (2016–2020)
167 50 m long transects were performed per field: one in the field centre, one in the field margin
168 (section between the field edge and the first seeding line) and the last in between, at c. 25 m
169 from the margin (except in 2015). Field margins generally harbour a more diverse fauna
170 compared to field cores. Transects were parallel to tram lines. For each transect, the surveyed
171 area was a theoretical volume, 2.5 m each side, 2.5 m in front of the observer and c. 1 m
172 above the observer's head. Surveys were conducted under calm weather conditions, i.e. sunny
173 days with air temperatures in full sun mostly above 15°C (95% T >16°C; 12° to 45°C, in
174 sunlight), and between 7:30 a.m. and 6:30 p.m. In 27.9% of the fields (226 of 811), two
175 sampling sessions were conducted (early- and late-season). Each plant–pollinator interaction
176 (i.e. an insect landing on a flower) was tallied, with both plant and pollinator being identified
177 (pollinators being caught with a sweep-net for later identification by a professional
178 entomologist if necessary). Transects were walked along very slowly (c. 12 s/m), for a total
179 duration of 10 min each. The observer stopped the timer each time an insect had to be
180 identified or captured. Butterflies were always identified by sight; however, given the time
181 spent in flight in butterflies, all individuals crossing the sampled theoretical volume were
182 tallied, even if not on a flower. Almost all pollinators and plants (94% and 96% respectively)
183 were identified to species level.

184

185 **2.3. Statistical analyses**

186 *2.3.1. Sampling completeness and frequency distribution of degrees*

187 We estimated sampling completeness of plants and pollinators using the Chao estimator of
188 asymptotic species richness (Chacoff et al., 2012), which is based on rare species (those that

189 occur only once or twice). We calculated the Chao 2 estimator using the *vegan* R package
190 (v2.5.7; Oksanen et al., 2020) and evaluated the percentage of completion (ratio between the
191 observed value and estimated value) of our sampling of plants and pollinators.
192 We analysed the frequency distribution of degrees in total as well as in subnetworks (per crop
193 and per period), testing for best-fit with *a priori* distributions, i.e. Exponential, Poisson,
194 Weibull, Lognormal, Negative binomial, Power law and Fisher's logseries. We used the
195 *fitdistrplus* package (Delignette-Muller and Dutang, 2015), except for the Fisher's logseries
196 (which is not implemented in this package), which we computed. Power law was tested with
197 the *powerLaw* package (Gillespie, 2015). Best fit was checked using the Akaike Information
198 Criterion (AIC; Burnham et al., 2011). We fixed a threshold of 2 units' difference in AIC to
199 select the best model among the candidate models.

200

201 2.3.2. Number of nodes and number of interactions: crop and temporal effects

202 From floral visitation events, we constructed bipartite networks with plant and bee nodes,
203 using a visitation event as a plant–pollinator interaction. To analyse how the network structure
204 varied among crops and seasons, we built interaction subnetworks. Six crop categories were
205 retained: OSR, cereals, alfalfa, meadows, SF and corn–sorghum. We also split the data into
206 three equal-length seasons, each 45 days long: April 1st–May 20th (OSR flowering peak,
207 characterised by high resource availability for insects), May 21st–July 9th, characterised by a
208 general lack of flowering crop resources available (i.e. the dearth period) – when honeybee
209 colony sizes are, perhaps surprisingly, at their highest (Odoux et al., 2014) – and finally, July
210 10th–August 22nd, corresponding to the SF flowering peak. We also divided our dataset into
211 six main taxonomic groups: honeybees, bumblebees, other social bees (including some
212 *Halictidae*, e.g. *Halictus rubicundus*, *Lasioglossum pauxillum*), solitary bees, hoverflies and
213 butterflies. To describe seasonal patterns in the number of interactions, we standardised our

214 data, since the number of transects per field and the number of fields sampled per crop and
215 per year varied. We thus converted the number of interactions per transect per field into a
216 number per hectare (ha), since each transect surveyed an area of 250 m². Next, we grouped
217 samples per fortnight by averaging values (with standard-error, se) of interactions per ha per
218 fortnight for each crop. Obtained values were compared with ANOVA and post-hoc Tukey
219 tests. Data were log-transformed.

220

221 2.3.3. *Network structure descriptors*

222 A large number of indices is available to describe network structure (see ESM2, which
223 presents their respective properties in Appendix A). We selected the following ones: number
224 of nodes (i.e. the number of species in interaction), connectance (the ratio between the number
225 of links observed and the total number of possible links), modularity – which gives insights
226 into the presence of subnetworks among the network (called modules, i.e. weakly interlinked
227 subsets of species that are constituted of strongly interacting species) and which was
228 calculated using the greedy Louvain algorithm method (Blondel et al., 2008) – and
229 nestedness, described by NODF (Nestedness metrics based on Overlap and Decreasing Fill;
230 Almeida-Neto et al., 2008). The latter is a property describing to what extent the partners of
231 specialists are a subset of the partners of generalists. We also used WNODF (Weighted
232 Nestedness metrics based on Overlap and Decreasing Fill; Almeida-Neto & Ulrich, 2011), the
233 quantitative alternative to NODF that accounts for the number of similar interactions. Finally,
234 we focused on the specialisation H_2' (standardised two-dimensional Shannon entropy) – the
235 degree of specialisation between the two levels of the entire network (Blüthgen et al., 2006) –
236 and the d' species-level specialisation measure (standardised Kullback–Leibler distance,
237 Blüthgen et al., 2006), corrected by de Manincor et al. (2020), representing the degree of
238 interaction specialisation at the species level. Network descriptors were obtained using the R

239 packages *bipartite* (Dormann et al., 2009), *igraph* (Csardi and Nepusz, 2006), *NetIndices*
240 (Kones et al., 2009) and *vegan* (Oksanen et al., 2020).
241 Some of these metrics (i.e. connectance, nestedness) are sensitive to sample or network size
242 (Dormann et al., 2009). We thus checked how indices varied with increasing numbers of
243 samples (according to periods and crops) by calculating all metrics (excluding d') with
244 increasing numbers of sampled fields. Five to 195 fields (every 10 field steps) were randomly
245 sampled (with replacement) from the overall sample. At each step (i.e. field number), 100
246 groups of fields were constituted, the six network metrics were computed and the mean and
247 the 95% confidence interval (CI) were calculated. The maximum bootstrap threshold to
248 compare the three seasons was 175 field groups, when OSR period samples stopped, but we
249 also went up to 195 fields to compare only dearth period and sunflower period. When
250 comparing the six different crop types, we had to stop at a threshold of 35 fields per group
251 (100 groups) in the bootstrap because we had very few fields of corn–sorghum and alfalfa. To
252 increase the sampling effort, we had to lose crops, and were left with only four crops with 100
253 groups of 85 fields. However, when focusing on the two mass-flowering crops, OSR and SF,
254 we were able to analyse 100 groups of 165 fields. Therefore, the number of fields
255 bootstrapped varied according to the crops to be compared.

256

257 *2.3.4. Field selection*

258 To calculate Chao 2 estimates and the number of interactions per hectare, we included the
259 fields in which no insects had been observed (138 fields, 17.0%). However, to represent
260 interaction networks and calculate metrics on aggregated fields, we only retained the fields on
261 which at least one interaction was observed in the dataset.

262 All analyses were run with *R software v.4.1.0* (R Core Team, 2021).

263

264 **3. Results**

265 **3.1. Sampling completeness and frequency distribution of degrees**

266 A total of 37,133 plant–pollinator interactions were tallied over 811 fields (127 of which were
267 studied in at least two years between 2015 and 2020), occurring between 200 species of
268 pollinators foraging on 165 species of plants (including individuals that were only identified
269 to genus level). We found 96 species of solitary bees, 35 species of social bees (including the
270 honeybee *A. mellifera* and 14 species of bumblebees), 38 species of butterflies and 31 species
271 of hoverflies. A total of 758 interactions were obtained between one plant or one pollinator
272 occurring only once. No interaction was found between one plant and one pollinator occurring
273 both only once. In regard to pollinators, over the full network, 80 species (40.0%) interacted
274 with only one plant (crop or weed), 29 species (14.5%) interacted with two, while 91 species
275 (45.5%) were more generalist (>2 plants, up to 92 plants). Similar values were obtained for
276 plants, with 38 weed species interacting with only one insect species (23%), and 16 with two
277 insect species (9.7%). There were 138 fields without any observed interaction (17.0%). We
278 caught 64.5% of pollinators, and recorded 78.6% of the plant community, based on Chao 2
279 estimators.

280 Overall, the frequency distribution of degrees (i.e. interacting plants/pollinators) fitted better
281 to the Fisher’s log-series (see ESM3 in Appendix A) than to Lognormal ($\Delta_{AIC} = 73$) or Power
282 laws ($\Delta_{AIC} = 77$). When splitting by crop or per season, similar results were found.

283 Differences in the goodness-of-fit of Fisher’s log-series and Lognormal were, however, much
284 lower in alfalfa ($\Delta_{AIC} = 12$) and in corn and sorghum crops ($\Delta_{AIC} = 3$) than in other crops.

285

286 **3.2. Number of nodes and number of interactions: crop and temporal effects**

287 The total number of nodes varied highly among crops (165, 114, 186, 178, 161 and 75 in
288 meadows, alfalfa, cereals, SF, OSR and corn–sorghum, respectively). Accumulation curves

289 per crop indicated that these differences among crops (Fig. 1) could be detected even from
290 low sample sizes (c. $n = 50$). Thus, independently of the number of sampled fields, meadows
291 showed the highest values of node numbers, followed by alfalfa, cereals, OSR and SF (Fig.
292 1). Year did not affect the pattern of the accumulation curves (see ESM4 in Appendix A),
293 although 2017 was a species-rich year regardless of the sampling effort.

294 The number of plant–pollinator interactions, here standardised per hectare of sampling effort
295 (which can be considered as an instantaneous visitation rate per hectare), was on average
296 338.43 ± 23.19 interactions/ha (mean \pm se). This value, however, showed strong seasonal
297 variations, with most interactions occurring in early July when considering all crops, or mid-
298 June when excluding SF (Fig. 2). There were also strong variations among crops, with
299 interactions in SF being overwhelmingly dominant and driving seasonal variation. In early
300 July, we found significant differences in the average number of interactions among crops
301 ($F_{5,62} = 20.16$, p -value < 0.0001), with significantly higher values for SF than in alfalfa (T_{HSD}
302 p -adj < 0.001), meadows (T_{HSD} p -adj < 0.001), cereals (T_{HSD} p -adj < 0.001) and OSR (T_{HSD} p -
303 adj = 0.0015).

304 Network structure differed strongly among crops in relation to variation of the total number of
305 nodes, either in the repartition of interactions between the main groups of plants or between
306 the main groups of pollinators (Fig. 3). In flowering crops, the crop flower was generally the
307 most visited flower in the crop considered. For instance, 95.27% of pollinators visited SF
308 plant in SF fields, and 86.07% OSR plants in OSR fields. However, this was true only for
309 mass-flowering crops: alfalfa represented approximately a third of all visited flowers in alfalfa
310 fields (36.12%) and corn was poorly visited in corn–sorghum fields (7.55%). In those crops,
311 pollinators preferentially foraged on wild flowers (89.62%). Moreover, honeybees’
312 dominance was not of the same magnitude in all crops: they were overrepresented in SF
313 (91.95%) and OSR (76.60%), while hoverflies rather than honeybees were the most-

314 interacting group in other crops (32.55% in corn and sorghum crops, 56.42% in cereal crops,
315 36.02% in alfalfa crops and 53.38% in meadows). Interestingly, when found in other crops,
316 OSR and SF flowers showed the same interaction pattern, i.e. interacting mainly with
317 honeybees. Moreover, hoverflies foraged more on wild flowers than did any other insect
318 group, potentially explaining their high presence in weed-rich crops such as meadows or
319 alfalfa. For instance, whereas the total percentage of weed visits by all pollinator groups was
320 63.78% in alfalfa crops, it reached 77.94% when considering only hoverfly visits.

321 The overall pattern of the network is depicted on Fig. 4, which represents standardised
322 interactions between pollinators and plants according to their degree (i.e. number of
323 interaction partners). Species with the highest degrees were not necessarily the most frequent,
324 although species' frequency of occurrence and degree were strongly related (i.e. decreasing
325 pattern of species degrees were concomitant with decreasing species frequencies) (Fig. 4). In
326 terms of the number of interactions, honeybees and bumblebees were most prevalent (83.72%
327 of the total interactions number, $n = 31,089$), followed by hoverflies (8.21%, $n = 3,050$),
328 solitary bees (2.69%, $n = 998$) and butterflies (1.36%, $n = 506$). Honeybees were by far the
329 dominant insect species (80.80% of all interactions, $n = 30,007$), the next two being
330 *Sphaerophoria scripta* (3.80%, $n = 1,412$) and *Episyrphus balteatus* (2.19%, $n = 812$), two
331 hoverflies. However, although *S. scripta* was found 20 fewer times than honeybees in total, its
332 number of plant partners (92) was higher than that of honeybees (75). Butterflies were far less
333 often observed, with only 81 interactions for the most common butterfly, *Pieris rapae*.

334 Concerning plants, SF represented 72.36% ($n = 26,871$) of the whole sample, seven times
335 more than OSR (10.69%, $n = 3,971$), and 60 times more than the most-interacting weed,
336 *Torilis arvensis* (1.20%, $n = 446$). Therefore, the complete network was dominated by
337 honeybees and mass-flowering crops: honeybees interacting with OSR or SF represented
338 77.44% of all interactions.

339

340 **3.3. Network structure**

341 *3.3.1. Seasonal effect on network structure*

342 The average number of nodes during the dearth period (mid-May to early July) was similar to
343 the average number of nodes for all periods, regardless of the sampling effort (Fig. 5a),
344 reaching 214 (CI_{95%}: 195–230) species when considering a sampling effort of 175 fields. The
345 average number of species was nearly equal during SF and OSR flowering blooms (154 (142–
346 165) and 148 (146–151), respectively, in 175 fields); that is to say, 1.4-times lower than
347 during the dearth period, but with a weaker variability among fields during OSR flowering.
348 Mean connectance did not vary a lot with period, being 3% (2.6–3.4) during bloom periods
349 and 2.5% (2.2–2.8) during the dearth period in 175 fields. Mean modularity was slightly
350 higher for the OSR blooming network (0.42 (0.41–0.43)) than for other periods (0.39 (0.36–
351 0.41) in SF blooming and 0.37 (0.35–0.38) in dearth period)). Specialisation of the network
352 remained constant during dearth and SF periods when considering network sizes above 100
353 fields (see ESM2 for details in Appendix A), but decreased much more quickly in the OSR
354 blooming period, with no overlapping 95% in CIs. This highlights that OSR blooming period
355 was represented by more generalist species than the other periods. Specialisation was higher
356 during the SF bloom and dearth periods, with average values between 0.49 (0.43–0.55) and
357 0.50 (0.46–0.52), respectively, compared to the OSR flowering bloom (0.30 (0.29–0.33)). The
358 OSR blooming period was characterised by a higher nestedness (37.2 (35.6–38.6) in 175
359 fields) than the SF blooming period (24.0 (21.3–26.9)) and dearth period (27.0 (24.4–29.9)).
360 Similar patterns were observed when using weighted NODF: 14.8 (12.7–16.8) during the
361 dearth period, 13.3 (11.8–15.0) during the SF blooming period, and 21.9 (20.6–23.1) during
362 the OSR blooming period. Thus, it seems that the flowering period of the OSR was the most
363 prone to host specialist species that had partners similar to generalist species.

364

365 3.3.2. Crop effect on network structure

366 All crops' subnetworks were similarly poorly connected (c. 3% at a threshold of 85 fields)
367 (Fig. 5b). Although OSR blooming period was the most modular period (see above), mean
368 modularity was almost identical whatever the crop considered, at about 0.5. At an equal
369 sampling survey effort of 85 fields, modularity was 0.43 (CI_{95%}: 0.39–0.47) for meadows,
370 0.45 (0.40–0.49) for SF and 0.48 (0.45–0.52) for OSR. With a higher sampling rate (n = 165
371 fields), but at the cost of losing crop types in the comparisons, OSR showed significant higher
372 modularity values than SF. The two mass-flowering crops' networks also had very different
373 specialisation (H2'), showing the two most extreme values among our six crops. OSR was far
374 less specialised (0.31 (0.29–0.33) at 165 fields) than all other crops, while SF (0.58 (0.55–
375 0.61)) showed the most specialised network; the other crops showed intermediate values, e.g.
376 0.45 (0.43–0.47) for meadows and 0.40 (0.34–0.46) for cereals at a survey effort of 85 fields.
377 Low specialisation in OSR could be partially explained by the low specialisation of the most
378 abundant insect, the honeybee ($d' = 0.27$) and the low specialisation of the crop flower, OSR
379 ($d' = 0.21$). Indeed, in OSR, the average d' of all species was 0.26 for insects (n = 97, median
380 = 0.20) and 0.23 for plants (n = 68, median = 0.44), meaning that the network was lowly
381 specialised at both trophic levels and highly driven by the honeybee and OSR flower
382 dominance. In SF, specialisation was 0.67 for SF (more than three times higher than in OSR)
383 and 0.42 for the honeybee (1.5-times higher than in OSR). SF was thus highly connected to
384 honeybees despite honeybees being slightly less specialised than SF, indicating a larger
385 partners' spectrum. The average d' of all insect species in SF was 0.41 (n = 109, median =
386 0.50) and 0.67 for plants (n = 65, median = 0.51), again reflecting the preponderant roles of
387 honeybees and SF. Nestedness (NODF) increased with sampling effort, being higher in OSR
388 than in any other crops (mean value of 29.8 (22.5–36.8) when sampling effort was low, i.e. 35

389 fields), especially in meadows (16.1 (11.0–21.3)), cereals (21.3 (15.1–28.4)) and SF (21.6
390 (16.8–29.2)), rather than in alfalfa (25.5 (21.0–30.8)). When increasing sampling effort (85-
391 field threshold), nestedness reached 33.0 (27.7–38.2) in OSR networks, c. 25% more than in
392 cereals (23.9 (19.9–28.5)) and in SF (23.4 (19.5–28.1)). Weighted NODF average values
393 showed similar patterns: SF reached a WNODF mean value of 14.5 (13.1–15.7) when
394 considering a threshold of 165 fields, whereas OSR networks peaked on average at 21.6
395 (20.3–22.5).

396

397 **4. Discussion**

398 The detection of several hundreds of species in the plant–pollinator network indicates that the
399 LTSER ZAPVS is characterised by a high diversity of species despite its intensive farming
400 system. While the honeybee, SF and OSR are by far the species that establish most
401 interactions, they are not necessarily those with the greatest number of interacting partners.
402 Although OSR flower is the species with the highest degree (number of interaction partners),
403 confirming its structuring role in the network (Diekötter et al., 2010; Stanley and Stout, 2014),
404 it is closely followed by wild plant species, generally hosted in less-intensively managed but
405 more diverse crops such as alfalfa or meadows. Moreover, the hoverfly *Sphaerophoria scripta*
406 showed far more partners than honeybee, including OSR and SF flowers, suggesting that its
407 role in the network has to be explored in more detail. *Bombus* species are also frequent
408 visitors of SF, as Syrphidae of OSR (Jauker and Wolters, 2008). Although these are rarely
409 studied as key species in networks, they might be strongly involved in OSR and SF
410 pollination (Breeze et al., 2011), and could act as pollination insurance if honeybees were to
411 decline dramatically (Sánchez-Bayo and Wyckhuys, 2019).

412 Completeness of survey effort was, however, moderate: 65% for insects but 80% for plants
413 according to the Chao 2 estimator, despite our dataset covering six months (from April to

414 September) and six years. The phenology of some plants and insects being very short may
415 explain why not all species were detected, in addition to the classic and inherently imperfect
416 detection method. In addition, the low connectance within crops (around 2–3%) suggests that
417 few interactions are carried out among all possible ones. These forbidden links (Jordano,
418 2016) could result from spatial or temporal constraints (Valdovinos, 2019). Indeed, one of our
419 possible explanations is that each crop network corresponds to a set of different fields that
420 belongs to different landscape entities and different sampling weeks, which generates a high
421 community variability in terms of phenology and location. But a low connectance, i.e. high
422 proportion of forbidden links, is in certain cases one of the indicators of destabilised networks
423 (Landi et al., 2018).

424 The hypothesis of the dominance of honeybee interactions with crop flowers during OSR
425 flowering and SF flowering, and not in other periods, was largely verified. Overdominance of
426 managed species may destabilise networks (Valido et al., 2019). The stability of a network
427 can be evaluated through the values of some network metrics (Thébault and Fontaine, 2010).
428 A stable bipartite mutualistic network is often a network with a high connectance and many
429 species, as well as a nested – but not very modular – network (Thébault and Fontaine, 2010).
430 In our case, connectance was always very low (<5%), whether we considered the total
431 network or the subnetworks per season or per crop. When the connectance is so low,
432 modularity and nestedness are usually positively correlated (Fortuna et al., 2010), hence a
433 situation with a nested but not modular network is unlikely, therefore possibly suggesting
434 instability in our networks. In particular, the corn–sorghum network, which is highly modular
435 and comprises few species, may show particularly low stability, though we had relatively
436 small sample sizes. Meadows were a special case, endowed with a large number of species
437 but not very nested and not very connected compared to other crops. Thus, although meadow
438 networks appear to be resilient networks thanks to the number of species they host (Okuyama

439 and Holland, 2008), they display a certain vulnerability because the sharing of partners is
440 weak, like in the studied SF crops, but contrary to the OSR crops.

441 Indeed, the most striking differences in networks were found between OSR and SF, both
442 being mass-flowering crops. As expected, these two crops drove the overall behaviour of
443 networks during their blooming season (approximately the same metrics of behaviour
444 between the OSR crop and OSR period, and between the SF crop and SF period), although the
445 meadows, weeds in cereals, and alfalfa were flowering at the same time. However, the OSR
446 and SF crops' networks were somewhat opposed in some of the key metrics, hence possibly
447 showing different network stability (Duchenne et al., 2022). The OSR network did not
448 harbour many species, but was highly nested compared to the other crop networks, which
449 therefore suggests it was a resilient network. Its nestedness may result from the fact that the
450 OSR network was mostly composed of species with a high degree of generalisation (i.e. the
451 average species specialisation was quite low in OSR crops), leading to a relatively nested
452 structure (Fontaine et al., 2009). Thus, as the OSR network was the most nested network in
453 our study, it was expected that the general core of species it included were generalists rather
454 than specialists, which was indeed the case. The interaction between OSR and honeybees has
455 a structuring effect for the network, corroborating studies that highlight OSR flower and its
456 partners as a hub that positively influences the visitation and pollination of neighbouring wild
457 plants, especially those with similar morphology (Thompson et al., 2021). OSR is also known
458 to host diverse insect communities (Stanley et al., 2013), while honeybees remain one of the
459 most efficient pollinators and, as demonstrated here, the most important partner of OSR.

460 Conversely, SF appears less stable than OSR because it is much less nested for an equivalent
461 diversity of species, and is composed of more specialised species, including the honeybee–SF
462 duo, which both qualify as specialist species, interacting strongly and almost entirely with
463 each other – a pattern that was not necessarily found in other studies (e.g. Valido et al., 2019).

464 Such a quasi-exclusive relationship between honeybees and SF as we found here may reflect
465 either competition within pollinators or plant communities in SF crops, or a strong niche
466 differentiation between wild and managed species. Conversely, OSR may host more
467 generalist species that interact with specialist species and therefore enhance the persistence of
468 the network (Liao et al., 2022).

469

470 **5. Conclusion**

471 Our results suggest that the honeybee and crop flower pair may differ between OSR and SF,
472 being an almost exclusive relationship in SF that prevents other partners by competition,
473 while allowing other modules in OSR hence increasing nestedness and consequently
474 improving network stability. This divergence may possibly result from phenological
475 difference, or as a consequence of flower shape – a divergence that can be further explored
476 with species trait approaches.

477

478 **Author Contributions**

479 Conceptualisation, V.B. and S.G.; Methodology, C.G., V.B. and S.G.; Formal Analysis, C.G.;
480 Writing-Original Draft Preparation, C.G., S.G. and V.B; Writing-Review and Editing, C.G.,
481 S.G. and V.B. All authors have read and agreed to the published version of the manuscript.

482

483 **Declaration of Competing Interest**

484 The authors declare that they have no known competing financial interests or personal
485 relationships that could have appeared to influence the work reported in this paper.

486

487 **Acknowledgements**

488 We thank in particular Jean-Luc Gautier, Alexis Saintilan and Marilyn Roncoroni for the
489 monitoring, sampling and identification of the pollinator' data. We also thank the farmers of
490 the LTSER 'Zone Atelier Plaine & Val de Sèvre' involved in the research program. CG
491 received funding for her PhD grant from the project SHOWCASE (SHOWCASing synergies
492 between agriculture, biodiversity and ecosystem services to help farmers capitalising on
493 native biodiversity) within the European Union's Horizon 2020 Research and Innovation
494 Programme under grant agreement N° 862480. This study was supported by the 'Programme
495 Pollinisateurs' (2015–2020) funded by the Ministère de l'Ecologie et de l'Environnement.

496

497 **Appendix A. Supporting information**

498 Supplementary data associated with this article can be found in the online version at [...]

499

500 **References**

- 501 Albrecht, M., Padrón, B., Bartomeus, I., Traveset, A., 2014. Consequences of plant invasions on
502 compartmentalization and species' roles in plant–pollinator networks. *Proc. R. Soc. B.* 281,
503 20140773. <https://doi.org/10.1098/rspb.2014.0773>
- 504 Almeida-Neto, M., Guimarães, P., Guimarães, P.R., Loyola, R.D., Ulrich, W., 2008. A consistent
505 metric for nestedness analysis in ecological systems: reconciling concept and measurement.
506 *Oikos* 117, 1227–1239. <https://doi.org/10.1111/j.0030-1299.2008.16644.x>
- 507 Almeida-Neto, M., Ulrich, W., 2011. A straightforward computational approach for measuring
508 nestedness using quantitative matrices. *Environmental Modelling & Software* 26, 173–178.
509 <https://doi.org/10.1016/j.envsoft.2010.08.003>
- 510 Bascompte, J., Jordano, P., Melián, C.J., Olesen, J.M., 2003. The Nested Assembly of Plant-Animal
511 Mutualistic Networks. *Proceedings of the National Academy of Sciences of the United States*
512 *of America* 100, 9383–9387.

513 Bendel, C.R., Kral-O'Brien, K.C., Hovick, T.J., Limb, R.F., Harmon, J.P., 2019. Plant–pollinator
514 networks in grassland working landscapes reveal seasonal shifts in network structure and
515 composition. *Ecosphere* 10. <https://doi.org/10.1002/ecs2.2569>

516 Blondel, V.D., Guillaume, J.-L., Lambiotte, R., Lefebvre, E., 2008. Fast unfolding of communities in
517 large networks. *Journal of Statistical Mechanics: Theory and Experiment* 2008, P10008.
518 <https://doi.org/10.1088/1742-5468/2008/10/P10008>

519 Blüthgen, N., 2010. Why network analysis is often disconnected from community ecology: A critique
520 and an ecologist's guide. *Basic and Applied Ecology* 11, 185–195.
521 <https://doi.org/10.1016/j.baae.2010.01.001>

522 Blüthgen, Nico, Menzel, F., Blüthgen, Nils, 2006. Measuring specialization in species interaction
523 networks. *BMC Ecology* 13.

524 Blüthgen, Nico, Menzel, F., Hovestadt, T., Fiala, B., Blüthgen, Nils, 2007. Specialization, Constraints,
525 and Conflicting Interests in Mutualistic Networks. *Current Biology* 17, 341–346.
526 <https://doi.org/10.1016/j.cub.2006.12.039>

527 Breeze, T.D., Bailey, A.P., Balcombe, K.G., Potts, S.G., 2011. Pollination services in the UK: How
528 important are honeybees? *Agriculture, Ecosystems & Environment* 142, 137–143.
529 <https://doi.org/10.1016/j.agee.2011.03.020>

530 Bretagnolle, V., Berthet, E., Gross, N., Gauffre, B., Plumejeaud, C., Houte, S., Badenhauer, I.,
531 Monceau, K., Allier, F., Monestiez, P., Gaba, S., 2018. Towards sustainable and
532 multifunctional agriculture in farmland landscapes: Lessons from the integrative approach of a
533 French LTSER platform. *Science of The Total Environment* 627, 822–834.
534 <https://doi.org/10.1016/j.scitotenv.2018.01.142>

535 Bretagnolle, V., Gaba, S., 2015. Weeds for bees? A review. *Agron. Sustain. Dev.* 35, 891–909.
536 <https://doi.org/10.1007/s13593-015-0302-5>

537 Burnham, K.P., Anderson, D.R., Huyvaert, K.P., 2011. AIC model selection and multimodel inference
538 in behavioral ecology: some background, observations, and comparisons. *Behav Ecol*
539 *Sociobiol* 65, 23–35. <https://doi.org/10.1007/s00265-010-1029-6>

540 Carpentier, C., Barabás, G., Spaak, J.W., De Laender, F., 2021. Reinterpreting the relationship
541 between number of species and number of links connects community structure and stability.
542 *Nature Ecology & Evolution* 5, 1102–1109. <https://doi.org/10.1038/s41559-021-01468-2>

543 Carvalho, L.G., Veldtman, R., Shenkute, A.G., Tesfay, G.B., Pirk, C.W.W., Donaldson, J.S.,
544 Nicolson, S.W., 2011. Natural and within-farmland biodiversity enhances crop productivity:
545 Weeds maximize nature benefits to crops. *Ecology Letters* 14, 251–259.
546 <https://doi.org/10.1111/j.1461-0248.2010.01579.x>

547 Chacoff, N.P., Resasco, J., Vázquez, D.P., 2018. Interaction frequency, network position, and the
548 temporal persistence of interactions in a plant–pollinator network. *Ecology* 99, 21–28.
549 <https://doi.org/10.1002/ecy.2063>

550 Chacoff, N.P., Vázquez, D.P., Lomáscolo, S.B., Stevani, E.L., Dorado, J., Padrón, B., 2012.
551 Evaluating sampling completeness in a desert plant-pollinator network: Sampling a plant-
552 pollinator network. *Journal of Animal Ecology* 81, 190–200. <https://doi.org/10.1111/j.1365-2656.2011.01883.x>

553

554 Chakraborty, P., Chatterjee, S., Smith, B.M., Basu, P., 2021. Seasonal dynamics of plant pollinator
555 networks in agricultural landscapes: how important is connector species identity in the
556 network? *Oecologia* 196, 825–837. <https://doi.org/10.1007/s00442-021-04975-y>

557 Csardi, G., Nepusz, T., 2006. The Igraph Software Package for Complex Network Research.
558 *InterJournal. Complex Systems* 1695.

559 de Manincor, N., Hautekèete, N., Mazoyer, C., Moreau, P., Piquot, Y., Schatz, B., Schmitt, E.,
560 Zélazny, M., Massol, F., 2020. How biased is our perception of plant-pollinator networks? A
561 comparison of visit- and pollen-based representations of the same networks. *Acta Oecologica*
562 105, 103551. <https://doi.org/10.1016/j.actao.2020.103551>

563 Delignette-Muller, M.L., Dutang, C., 2015. fitdistrplus: An R Package for Fitting Distributions.
564 *Journal of Statistical Software* 64. <https://doi.org/10.18637/jss.v064.i04>

565 Diekötter, T., Kadoya, T., Peter, F., Wolters, V., Jauker, F., 2010. Oilseed rape crops distort plant-
566 pollinator interactions. *Journal of Applied Ecology* 47, 209–214.
567 <https://doi.org/10.1111/j.1365-2664.2009.01759.x>

568 Dormann, C.F., Frund, J., Bluthgen, N., Gruber, B., 2009. Indices, Graphs and Null Models:
569 Analyzing Bipartite Ecological Networks. *TOECOLJ* 2, 7–24.
570 <https://doi.org/10.2174/1874213000902010007>

571 Duchenne, F., Wüest, R.O., Graham, C.H., 2022. Seasonal structure of interactions enhances
572 multidimensional stability of mutualistic networks. *Proceedings of the Royal Society B:
573 Biological Sciences* 289. <https://doi.org/10.1098/rspb.2022.0064>

574 Dunn, L., Lequerica, M., Reid, C.R., Latty, T., 2020. Dual ecosystem services of syrphid flies
575 (Diptera: Syrphidae): pollinators and biological control agents. *Pest Manag Sci* 76, 1973–
576 1979. <https://doi.org/10.1002/ps.5807>

577 Ferreira, P.A., Boscolo, D., Viana, B.F., 2013. What do we know about the effects of landscape
578 changes on plant–pollinator interaction networks? *Ecological Indicators* 31, 35–40.
579 <https://doi.org/10.1016/j.ecolind.2012.07.025>

580 Fontaine, C., Thébault, E., Dajoz, I., 2009. Are insect pollinators more generalist than insect
581 herbivores? *Proceedings of the Royal Society B: Biological Sciences* 276, 3027–3033.
582 <https://doi.org/10.1098/rspb.2009.0635>

583 Fortuna, M.A., Stouffer, D.B., Olesen, J.M., Jordano, P., Mouillot, D., Krasnov, B.R., Poulin, R.,
584 Bascompte, J., 2010. Nestedness versus modularity in ecological networks: two sides of the
585 same coin? *Journal of Animal Ecology*. <https://doi.org/10.1111/j.1365-2656.2010.01688.x>

586 Frankie, G.W., Thorp, R.W., Hernandez, J., Rizzardi, M., Ertter, B., Pawelek, J.C., Witt, S.L.,
587 Schindler, M., Coville, R., Wojcik, V.A., 2009. Native bees are a rich natural resource in
588 urban California gardens. *California Agriculture* 63, 113–120.
589 <https://doi.org/10.3733/ca.v063n03p113>

590 Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R., Cunningham, S.A.,
591 Carvalho, L.G., Chacoff, N.P., Dudenhöffer, J.H., Greenleaf, S.S., Holzschuh, A., Isaacs,
592 R., Krewenka, K., Mandelik, Y., Mayfield, M.M., Morandin, L.A., Potts, S.G., Ricketts, T.H.,
593 Szentgyörgyi, H., Viana, B.F., Westphal, C., Winfree, R., Klein, A.M., 2011. Stability of
594 pollination services decreases with isolation from natural areas despite honey bee visits:

595 Habitat isolation and pollination stability. *Ecology Letters* 14, 1062–1072.
596 <https://doi.org/10.1111/j.1461-0248.2011.01669.x>

597 Geslin, B., Gauzens, B., Baude, M., Dajoz, I., Fontaine, C., Henry, M., Ropars, L., Rollin, O.,
598 Thébault, E., Vereecken, N.J., 2017. Massively Introduced Managed Species and Their
599 Consequences for Plant–Pollinator Interactions. *Advances in Ecological Research* 57, 147–
600 199. <https://doi.org/10.1016/bs.aecr.2016.10.007>

601 Giannini, T.C., Garibaldi, L.A., Acosta, A.L., Silva, J.S., Maia, K.P., Saraiva, A.M., Guimarães, P.R.,
602 Kleinert, A.M.P., 2015. Native and Non-Native Supergeneralist Bee Species Have Different
603 Effects on Plant-Bee Networks. *PLoS ONE* 10, e0137198.
604 <https://doi.org/10.1371/journal.pone.0137198>

605 Gillespie, C.S., 2015. Fitting Heavy Tailed Distributions: The *powerLaw* Package. *J. Stat. Soft.* 64.
606 <https://doi.org/10.18637/jss.v064.i02>

607 Habel, J.C., Ulrich, W., Biburger, N., Seibold, S., Schmitt, T., 2019. Agricultural intensification drives
608 butterfly decline. *Insect Conserv Divers* icad.12343. <https://doi.org/10.1111/icad.12343>

609 Hoyle, M., Cresswell, J.E., 2007. The effect of wind direction on cross-pollination in wind-pollinated
610 gm crops. *Ecological Applications* 17, 1234–1243. <https://doi.org/10.1890/06-0569>

611 Jauker, F., Diekötter, T., Schwarzbach, F., Wolters, V., 2009. Pollinator dispersal in an agricultural
612 matrix: opposing responses of wild bees and hoverflies to landscape structure and distance
613 from main habitat. *Landscape Ecol* 24, 547–555. <https://doi.org/10.1007/s10980-009-9331-2>

614 Jauker, F., Wolters, V., 2008. Hover flies are efficient pollinators of oilseed rape. *Oecologia* 156, 819–
615 823. <https://doi.org/10.1007/s00442-008-1034-x>

616 Johst, K., Drechsler, M., Thomas, J., Settele, J., 2006. Influence of mowing on the persistence of two
617 endangered large blue butterfly species: Influence of mowing on butterfly persistence. *Journal*
618 *of Applied Ecology* 43, 333–342. <https://doi.org/10.1111/j.1365-2664.2006.01125.x>

619 Jordano, P., 2016. Sampling networks of ecological interactions. *Functional Ecology* 30, 1883–1893.
620 <https://doi.org/10.1111/1365-2435.12763>

621 Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C.,
622 Tschamtko, T., 2007. Importance of pollinators in changing landscapes for world crops. *Proc.*
623 *R. Soc. B.* 274, 303–313. <https://doi.org/10.1098/rspb.2006.3721>

624 Kones, J.K., Soetaert, K., Van Oevelen, D., Owino, J., 2009. Are network indices robust indicators of
625 food web functioning? a Monte Carlo approach. *Ecological Modelling* 370–382.

626 Kovács-Hostyánszki, A., Földesi, R., Báldi, A., Endrédi, A., Jordán, F., 2019. The vulnerability of
627 plant-pollinator communities to honeybee decline: A comparative network analysis in
628 different habitat types. *Ecological Indicators* 97, 35–50.
629 <https://doi.org/10.1016/j.ecolind.2018.09.047>

630 Landi, P., Minoarivelo, H.O., Brännström, Å., Hui, C., Dieckmann, U., 2018. Complexity and stability
631 of ecological networks: a review of the theory. *Population Ecology* 60, 319–345.
632 <https://doi.org/10.1007/s10144-018-0628-3>

633 Lázaro, A., Müller, A., Ebmer, A.W., Dathe, H.H., Scheuchl, E., Schwarz, M., Risch, S., Pauly, A.,
634 Devalez, J., Tschulin, T., Gómez-Martínez, C., Papas, E., Pickering, J., Waser, N.M.,
635 Petanidou, T., 2021. Impacts of beekeeping on wild bee diversity and pollination networks in
636 the Aegean Archipelago. *Ecography* 44, 1353–1365. <https://doi.org/10.1111/ecog.05553>

637 Liao, J., Bearup, D., Strona, G., 2022. A patch-dynamic metacommunity perspective on the
638 persistence of mutualistic and antagonistic. *Ecology* 103. <https://doi.org/10.1002/ecy.3686>

639 Ma, A., Lu, X., Gray, C., Raybould, A., Tamaddoni-Nezhad, A., Woodward, G., Bohan, D.A., 2019.
640 Ecological networks reveal resilience of agro-ecosystems to changes in farming management.
641 *Nature Ecology & Evolution* 3, 260–264. <https://doi.org/10.1038/s41559-018-0757-2>

642 Magrach, A., Holzschuh, A., Bartomeus, I., Riedinger, V., Roberts, S.P.M., Rundlof, M., Vujic, A.,
643 Wickens, J.B., Wickens, V.J., Bommarco, R., Gonzalez-Varo, J.P., Potts, S.G., Smith, H.G.,
644 Steffan-Dewenter, I., Vila, M., 2018. Plant-pollinator networks in semi-natural grasslands are
645 resistant to the loss of pollinators during blooming of mass-flowering crops. *Ecography* 41,
646 62–74. <https://doi.org/10.1111/ecog.02847>

647 Mathiasson, M.E., Rehan, S.M., 2020. Wild bee declines linked to plant-pollinator network changes
648 and plant species introductions. *Insect Conservation and Diversity* 13, 595–605.
649 <https://doi.org/10.1111/icad.12429>

650 Memmott, J., Waser, N.M., Price, M.V., 2004. Tolerance of pollination networks to species
651 extinctions. *Proc. R. Soc. Lond. B* 271, 2605–2611. <https://doi.org/10.1098/rspb.2004.2909>

652 Morrison, B.M.L., Brosi, B.J., Dirzo, R., 2020. Agricultural intensification drives changes in hybrid
653 network robustness by modifying network structure. *Ecology Letters* 23, 359–369.
654 <https://doi.org/10.1111/ele.13440>

655 Odoux, J.-F., Aupinel, P., Gateff, S., Requier, F., Henry, M., Bretagnolle, V., 2014. ECOBEE: a tool
656 for long-term honey bee colony monitoring at the landscape scale in West European intensive
657 agroecosystems. *Journal of Apicultural Research* 53, 57–66.
658 <https://doi.org/10.3896/IBRA.1.53.1.05>

659 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
660 O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2020.
661 *vegan: Community Ecology Package*. R package version 2.5-7. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
662 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).

663 Okuyama, T., Holland, J.N., 2008. Network structural properties mediate the stability of mutualistic
664 communities. *Ecology Letters* 11, 208–216. <https://doi.org/10.1111/j.1461-0248.2007.01137.x>

665 Olesen, J.M., Stefanescu, C., Traveset, A., 2011. Strong, Long-Term Temporal Dynamics of an
666 Ecological Network. *PLoS ONE* 6, e26455. <https://doi.org/10.1371/journal.pone.0026455>

667 Ollerton, J., Winfree, R., Tarrant, S., 2011. How many flowering plants are pollinated by animals?
668 *Oikos* 120, 321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>

669 Perrot, T., Gaba, S., Roncoroni, M., Gautier, J.-L., Bretagnolle, V., 2018. Bees increase oilseed rape
670 yield under real field conditions. *Agriculture, Ecosystems & Environment* 266, 39–48.
671 <https://doi.org/10.1016/j.agee.2018.07.020>

672 Perrot, T., Gaba, S., Roncoroni, M., Gautier, J.-L., Saintilan, A., Bretagnolle, V., 2019. Experimental
673 quantification of insect pollination on sunflower yield, reconciling plant and field scale
674 estimates. *Basic and Applied Ecology* 34, 75–84. <https://doi.org/10.1016/j.baae.2018.09.005>

675 Petanidou, T., Kallimanis, A.S., Tzanopoulos, J., Sgardelis, S.P., Pantis, J.D., 2008. Long-term
676 observation of a pollination network: fluctuation in species and interactions, relative
677 invariance of network structure and implications for estimates of specialization: High
678 plasticity in plant-pollinator networks. *Ecology Letters* 11, 564–575.
679 <https://doi.org/10.1111/j.1461-0248.2008.01170.x>

680 Ponisio, L.C., Gaiarsa, M.P., Kremen, C., 2017. Opportunistic attachment assembles plant–pollinator
681 networks. *Ecol Lett* 20, 1261–1272. <https://doi.org/10.1111/ele.12821>

682 R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for
683 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

684 Requier, F., Odoux, J.-F., Tamic, T., Moreau, N., Henry, M., Decourtye, A., Bretagnolle, V., 2015.
685 Honey bee diet in intensive farmland habitats reveals an unexpectedly high flower richness
686 and a major role of weeds. *Ecological Applications* 25, 881–890. <https://doi.org/10.1890/14-1011.1>

687

688 Rollin, O., Benelli, G., Benvenuti, S., Decourtye, A., Wratten, S.D., Canale, A., Desneux, N., 2016.
689 Weed-insect pollinator networks as bio-indicators of ecological sustainability in agriculture. A
690 review. *Agron. Sustain. Dev.* 36, 8. <https://doi.org/10.1007/s13593-015-0342-x>

691 Sánchez-Bayo, F., Wyckhuys, K.A.G., 2019. Worldwide decline of the entomofauna: A review of its
692 drivers. *Biological Conservation* 232, 8–27. <https://doi.org/10.1016/j.biocon.2019.01.020>

693 Stanley, D.A., Gunning, D., Stout, J.C., 2013. Pollinators and pollination of oilseed rape crops
694 (*Brassica napus* L.) in Ireland: ecological and economic incentives for pollinator conservation.
695 *Journal of Insect Conservation* 17, 1181–1189. <https://doi.org/10.1007/s10841-013-9599-z>

696 Stanley, D.A., Stout, J.C., 2014. Pollinator sharing between mass-flowering oilseed rape and co-
697 flowering wild plants: implications for wild plant pollination. *Plant Ecol* 215, 315–325.
698 <https://doi.org/10.1007/s11258-014-0301-7>

699 Thébault, E., Fontaine, C., 2010. Stability of Ecological Communities and the Architecture of
700 Mutualistic and Trophic Networks. *Science* 329, 853–856.
701 <https://doi.org/10.1126/science.1188321>

- 702 Thompson, A., Ștefan, V., Knight, T.M., 2021. Oilseed Rape Shares Abundant and Generalized
703 Pollinators with Its Co-Flowering Plant Species. *Insects* 12, 1096.
704 <https://doi.org/10.3390/insects12121096>
- 705 Valdovinos, F.S., 2019. Mutualistic networks: moving closer to a predictive theory. *Ecology Letters*
706 22, 1517–1534. <https://doi.org/10.1111/ele.13279>
- 707 Valido, A., Rodríguez-Rodríguez, M.C., Jordano, P., 2019. Honeybees disrupt the structure and
708 functionality of plant-pollinator networks. *Scientific Reports* 9, 4711.
709 <https://doi.org/10.1038/s41598-019-41271-5>