



HAL
open science

Support and distribution inference from noisy data

Jeremie Capitaio-Miniconi, Élisabeth Gassiat, Luc Lehéricy

► **To cite this version:**

Jeremie Capitaio-Miniconi, Élisabeth Gassiat, Luc Lehéricy. Support and distribution inference from noisy data. 2025. hal-04073724v3

HAL Id: hal-04073724

<https://hal.science/hal-04073724v3>

Preprint submitted on 11 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Support and distribution inference from noisy data

J eremie Capitaou-Miniconi*,  elisabeth Gassiat* and Luc Lehericy**

* Universit  Paris-Saclay, CNRS, Laboratoire de math matiques d'Orsay, 91405, Orsay, France,

** Universit  C te d'Azur, CNRS, LJAD, France.

Abstract

Understanding what kind of noise in the observations allows to recover low dimensional structures of a signal is of interest in statistical learning, as a first step to build efficient dimension reduction procedures. In this work we give a new contribution on the type of noise which can affect the data without preventing to build consistent estimators of the support and distribution of the signal. We focus on the situation where the observations are corrupted with additive and independent noise. We prove that for general classes of supports, it is possible to recover both the support and the distribution of the signal without knowing the noise distribution and with no sample of the noise. We exhibit classes of distributions over which we prove adaptive minimax rates (up to a log log factor) for the estimation of the support in Hausdorff distance. Moreover, for the class of distributions with compact support, we provide estimators of the unknown (in general singular) distribution and prove maximum rates in Wasserstein distance. We also prove an almost matching lower bound on the associated minimax risk.

1 Introduction

1.1 Context and aim

It is often observed that high dimensional data has a low intrinsic dimension. The computational geometry point of view gave rise to a number of interesting algorithms (see [6] and references therein) for the reconstruction of a non linear shape from a point cloud, and the case of non noisy data is by now relatively well understood. See for instance [1] for an overview of the subject and references when the observations are sampled on an unknown manifold, and [14] and [24] for the estimation of the distribution using the Wasserstein metric as loss function. Some of the geometric ideas that have been developed to handle non noisy data can be applied, or adapted, to handle noisy data and build estimators with controlled risk. However, the upper bound on the risk is meaningful only when the bound on the noise is small, or even when the noise tends to 0 with the amount of data tending to infinity. See [2], [1], [12], [16], [25], [3]. The estimation of the distribution with unknown but small (and orthogonal to the unknown manifold) noise is handled in [14] with a kernel estimator. Non parametric Bayesian methods have been explored in [5] for observations on a tubular neighborhood of the unknown manifold, that is for bounded noise.

In this paper, we aim at giving a new contribution on the type of *unknown* noise which can affect the data without preventing to build consistent estimators of the support and of the law of the signal. We are interested in the estimation of possibly low dimensional supports, and of distributions supported on such supports, when the observations are corrupted with additive noise: this is called the deconvolution model. In contrast with previous literature, we consider the setting where the distribution of the noise is unknown and where we do not have access to auxiliary data to estimate it.

When we accept to consider noise with known distribution, usual estimation procedures are roughly based on the fact that it is possible to get an estimator of the characteristic function of the non noisy data by dividing an estimator of the characteristic function of the noisy data by that of the noise. The authors of [20] consider data corrupted with Gaussian noise, and

propose as estimator of the manifold an upper level set of an estimator of a kernel smoothed density of the unknown distribution. With the truncated Hausdorff loss, the authors prove that their estimator achieves a maximum risk (over some class of distributions) upper bounded by $(\sqrt{\log n})^{-1+\delta}$ for any positive δ , and prove a lower bound of order $(\log n)^{-1+\delta}$ for the minimax risk. Taking an upper level set of an estimated density had been earlier proposed to estimate a support based on non noisy data in [9]. In the context of full dimensional convex support and with additive Gaussian noise, [7] proposes an estimation procedure using convexity ideas. The authors prove an upper bound of order $\log \log n / \sqrt{\log n}$ and a lower bound of order $(\log n)^{-2/\tau}$ for the minimax Hausdorff risk, for any $\tau \in (0, 1)$. Earlier work with known noise and with full dimensional support is [22], where the author first builds an estimator of the unknown density using deconvolution ideas, then samples from this estimated density and takes a union of balls centered on the sampled points, such as in [13]. Focusing on the estimation of the distribution, with known Gaussian noise, the authors of [11] prove matching upper and lower bounds for the minimax risk using the Wasserstein distance. Results for other known noises, but limited to one dimensional observations, can be found in [10].

1.2 Contribution and main results

The intuition behind our work is that certain geometrical properties of the support of the signal induce sufficient structure for the deconvolution problem to be solvable even without knowledge of the noise distribution. The main original contributions of our work are twofold.

- We exhibit simple geometric properties of a support so that, whatever the distribution on such a support—provided it does not have too heavy tails—, the deconvolution problem can be solved without any knowledge regarding the noise.
- We propose classes of distributions over which we prove adaptive minimax rates (up to a $\log \log$ factor) for the estimation of the support in Hausdorff distance. Moreover, for the class of distributions with compact support, we provide estimators of the unknown (in general singular) distribution and prove rates for the maximum risk in Wasserstein distance. We also prove an almost matching lower bound on the associated minimax risk.

In Section 2, we precise the setting and the identifiability of the deconvolution problem of [17], which our paper builds upon. The main assumption needed in [17] to ensure that the recovery of the distribution of the signal is possible, (Adep), comes from a complex analysis argument and does not have a clear probabilistic or geometric interpretation; as such it may seem difficult to verify outside of specific cases. We exhibit in Section 2.3 simple geometric properties of a support so that, whatever the distribution on such a support, (Adep) holds and therefore the deconvolution problem can be solved without any knowledge regarding the noise, see Corollaries 1 and 2. The genericity of these geometric conditions is considered in Section 2.4.

Section 3 provides an overview of our estimation procedures and the estimation of the characteristic function of the signal. We focus on support estimation in Section 4. We are able to propose estimators achieving adaptive minimax rates (up to a $\log \log$ factor) for the estimation of the support in Hausdorff distance. The minimax rates are investigated over well chosen classes of distributions, see Theorems 3, 4 and 5. Specifically, the minimax risk for the Hausdorff distance is upper bounded by $(\log \log n)^L / (\log n)^\kappa$ for some L , where $\kappa \in (1/2, 1]$ is a parameter characterizing the heaviness of the tail of the distribution of the signal ($\kappa = 1$ corresponds to compactly supported distributions, and $\kappa = 1/2$ to sub-Gaussian distributions). On the other hand, the minimax risk is lower bounded by $1/(\log n)^\kappa$ if $\kappa \in (1/2, 1)$ and $1/(\log n)^{1-\delta}$ if $\kappa = 1$, δ being any (small) positive number. Adaptivity is with respect to κ . In some sense, exhibiting these classes of distributions allows to fill the gap between the upper and lower bounds in [20] and [7], together with the extension to totally unknown noise.

Section 5 is devoted to the estimation of the distribution when it is compactly supported. We consider the estimation of the unknown (in general singular) distribution of the hidden non noisy data itself when it has a compact support. We prove almost matching upper and lower

bounds of order $1/\log n$ for the estimation risk of the distribution in Wasserstein distance, see Theorem 6 and Theorem 7.

Appendix A presents an index of the notations defined along the paper. Detailed proofs for Sections 2, 3, 4 and 5 are given in the remaining appendices.

1.3 Notations

The Euclidean norm (in any dimension) will be denoted $\|\cdot\|_2$. If A is a subset of \mathbb{R}^D , we write $d(x, A) = \inf\{\|x - y\|_2 \mid y \in A\}$. For any $r > 0$, we write $B_r = (-r, r)$. For any measurable function f on B_r^D , we write $\|f\|_{\infty, r}$ the essential supremum of f over B_r^D and $\|f\|_{2, r}$ the norm of f in $L^2(B_r^D)$.

2 Geometric conditions for identifiability

In this section, we recall the general identifiability Theorem proved in [17]. We provide geometric conditions on the support of the signal that ensure that this theorem can be applied.

2.1 Setting

We consider independent and identically distributed observations Y_i , $i = 1, \dots, n$ coming from the model

$$Y = X + \varepsilon, \quad (1)$$

in which the signal X and the noise ε are independent random variables. We assume that the observation has dimension at least two, and that its coordinates can be partitioned in such a way that the corresponding blocks of noise variables are independently distributed, that is

$$Y = \begin{pmatrix} Y^{(1)} \\ Y^{(2)} \end{pmatrix} = \begin{pmatrix} X^{(1)} \\ X^{(2)} \end{pmatrix} + \begin{pmatrix} \varepsilon^{(1)} \\ \varepsilon^{(2)} \end{pmatrix} = X + \varepsilon \quad (2)$$

in which $Y^{(1)}, X^{(1)}, \varepsilon^{(1)} \in \mathbb{R}^{d_1}$ and $Y^{(2)}, X^{(2)}, \varepsilon^{(2)} \in \mathbb{R}^{d_2}$, for $d_1, d_2 \geq 1$ with $d_1 + d_2 = D$, and we assume that the noise components $\varepsilon^{(1)}$ and $\varepsilon^{(2)}$ are independent random variables. We write G the distribution of X and \mathcal{M}_G its support. For $i \in \{1, 2\}$, we write $\mathbb{Q}^{(i)}$ the distribution of $\varepsilon^{(i)}$, so that $\mathbb{Q} = \mathbb{Q}^{(1)} \otimes \mathbb{Q}^{(2)}$ is the distribution of ε .

We shall not make any more assumption on the distribution of the noise ε , and we shall not assume that its distribution is known. Indeed in [17], it is proved that under very mild conditions on the distribution of the signal X , model (2) is fully identifiable, that is one can recover G , and thus its support, and \mathbb{Q} from the convolution $G * \mathbb{Q}$.

2.2 Identifiability Theorem

Let us introduce the assumptions on the distribution of the signal we shall use. To state the first assumption, we let κ be a positive real number.

A(κ) There exist $a, b > 0$ such that for all $\lambda \in \mathbb{R}^D$, $\mathbb{E}[\exp(\lambda^\top X)] \leq a \exp(b\|\lambda\|_2^{1/\kappa})$.

Assumption A(κ) is about the tail of G as the following proposition shows.

Proposition 1. • *A random variable X satisfies A(1) if and only if its support is compact.*

- *A random variable X satisfies A(κ) for $\kappa \in [0, 1)$ if and only if there exist constants $c, d > 0$ such that for any $t \geq 0$,*

$$\mathbb{P}(\|X\|_2 \geq t) \leq c \exp(-dt^{1/(1-\kappa)}).$$

The proof of Proposition 1 is detailed in Appendix B.1.

Under $A(\kappa)$, the characteristic function of the signal can be extended into the multivariate analytic function

$$\begin{aligned} \Phi_X : \mathbb{C}^{d_1} \times \mathbb{C}^{d_2} &\longrightarrow \mathbb{C} \\ (z_1, z_2) &\longmapsto \mathbb{E} \left[\exp \left(iz_1^\top X^{(1)} + iz_2^\top X^{(2)} \right) \right]. \end{aligned}$$

The second assumption is the following.

(Adep) For any $z_0 \in \mathbb{C}^{d_1}$, $z \mapsto \Phi_X(z_0, z)$ is not the null function and for any $z_0 \in \mathbb{C}^{d_2}$, $z \mapsto \Phi_X(z, z_0)$ is not the null function.

It has been shown in [17] that several models satisfy this assumption, such as the repeated measurements submodel (see Corollary 2.3 in [17]) or the noisy independent component analysis submodel (see Corollary 2.2 in [17]). In particular, it follows directly from Corollary 2.4 of [17] and the inverse function Theorem that the errors in variable regression model (in which $X^{(2)} = g(X^{(1)})$ for some function g) satisfies assumption (Adep) when the regression function is a non-constant differentiable function. In this paper, we are interested in identifying supports of distributions and we will provide geometrical conditions so that Assumption (Adep) is verified, see Section 2.3.

Obviously, if no centering constraint is put on the signal or on the noise, it is possible to translate the signal by a fixed vector $m \in \mathbb{R}^D$ and the noise by $-m$ without changing the observation. The model can thus be identifiable only up to translation.

Theorem 1 (from [17]). *If the distribution of the signal satisfies $A(\kappa)$ for $\kappa \in (1/2, 1]$ and (Adep), then the distribution of the signal and the distribution of the noise can be recovered up to translation from the distribution of the observations.*

The proof of this theorem is based on recovering Φ_X for which the structure provided by (Adep) is crucial. The arguments show that knowing the characteristic function of the observations in a neighborhood of the origin allows to recover Φ_X in a neighborhood of the origin, and then over the whole multidimensional complex plane.

2.3 Sufficient geometrical conditions for (Adep) to hold

In [8], the authors prove that (Adep) holds for random variables supported on a sphere. In this section, we provide very general conditions on the support of a random variable which are sufficient for (Adep) to hold. At the end of this section, we show concrete examples where these conditions are satisfied.

We define the following assumptions (H1) and (H2). Here, if A is a subset of \mathbb{R}^D , we write $\text{Diam}(A)$ its diameter $\sup\{\|x - y\|_2 \mid x, y \in A\}$.

(H1) There exist $(A_\Delta)_{\Delta>0}$ and $(B_\Delta)_{\Delta>0}$ such that $A_\Delta \subset \mathbb{R}^{d_2}$, $B_\Delta \subset \mathbb{R}^{d_1}$, $\mathbb{P}(X^{(2)} \in A_\Delta) > 0$, $\lim_{\Delta \rightarrow 0} \text{Diam}(B_\Delta) = 0$ and $\mathbb{P}(X^{(1)} \in B_\Delta \mid X^{(2)} \in A_\Delta) = 1$.

(H2) There exist $(A_\Delta)_{\Delta>0}$ and $(B_\Delta)_{\Delta>0}$ such that $A_\Delta \subset \mathbb{R}^{d_1}$, $B_\Delta \subset \mathbb{R}^{d_2}$, $\mathbb{P}(X^{(1)} \in A_\Delta) > 0$, $\lim_{\Delta \rightarrow 0} \text{Diam}(B_\Delta) = 0$ and $\mathbb{P}(X^{(2)} \in B_\Delta \mid X^{(1)} \in A_\Delta) = 1$.

We prove that these assumptions are sufficient to ensure identifiability provided that $A(\kappa)$ is satisfied. In what follows we shall say that a random variable satisfies (Adep) when its characteristic function does.

Theorem 2. *Assume that the distribution of X satisfies $A(\kappa)$, (H1) and (H2). Then X satisfies $A(\kappa)$ and (Adep).*

The proof of Theorem 2 is detailed in Section B.2.

One can interpret the assumptions (H1) and (H2) geometrically as shown in Figure 1. In essence, it means that there exists a slice (along the first d_1 , resp. last d_2 , coordinates, with

base A_Δ) such that the random variable belongs to this slice with positive probability and such that on this slice, the support of the distribution is contained in an orthogonal slice (along the last d_2 , resp. first d_1 , coordinates) of diameter shrinking to zero with Δ .

Let us introduce sets of support that guarantee that (H1) and (H2) hold for any signal with support in these sets. Let $\pi^{(1:d_1)}$ (resp. $\pi^{(d_1+1:D)}$) be the projection which extracts the first d_1 (resp. last d_2) coordinates of vectors of \mathbb{R}^D , and let

$$\mathcal{A}_1(\Delta, \varepsilon) = \{\mathcal{M} \subset \mathbb{R}^D \mid \text{There exists } x = (x_1, x_2) \in \mathcal{M} \\ \text{such that } \text{Diam} \left(\pi^{(1:d_1)} \left[\mathcal{M} \cap (\mathbb{R}^{d_1} \times \bar{B}(x_2, \varepsilon)) \right] \right) < \Delta\}$$

and

$$\mathcal{A}_2(\Delta, \varepsilon) = \{\mathcal{M} \subset \mathbb{R}^D \mid \text{There exists } x = (x_1, x_2) \in \mathcal{M} \\ \text{such that } \text{Diam} \left(\pi^{(d_1+1:D)} \left[\mathcal{M} \cap (\bar{B}(x_1, \varepsilon) \times \mathbb{R}^{d_2}) \right] \right) < \Delta\}.$$

The following corollary is a direct consequence of Theorem 2.

Corollary 1. *Let $\mathcal{M} \in \mathcal{A} := (\cap_{\Delta>0} \cup_{\varepsilon>0} \mathcal{A}_1(\Delta, \varepsilon)) \cap (\cap_{\Delta>0} \cup_{\varepsilon>0} \mathcal{A}_2(\Delta, \varepsilon))$. Then any random variable with support \mathcal{M} satisfies (H1) and (H2).*

This is analog to (H1) and (H2) in the sense that if $\mathcal{M} \in \mathcal{A}$, there exists sequences $(A_\Delta)_{\Delta>0}$ (of the form $\bar{B}(x_1, \varepsilon)$) and $(B_\Delta)_{\Delta>0}$ with $\text{Diam}(B_\Delta) < \Delta$ and such that if $(X^{(1)}, X^{(2)})$ has support \mathcal{M} , then conditionally to $X^{(1)} \in A_\Delta$, $X^{(2)} \in B_\Delta$ a.s., and likewise for (H2). Note that this is not strictly equivalent since the A_Δ are not necessarily open in (H1) and (H2).

In the case of compact supports, these sets $(\cap_{\Delta>0} \cup_{\varepsilon>0} \mathcal{A}_1(\Delta, \varepsilon))$ admit a simple formulation. Define the sets \mathcal{B}_1 and \mathcal{B}_2 as

$$\mathcal{B}_1 = \{\mathcal{M} \subset \mathbb{R}^D \text{ compact} \mid \exists x_1 \in \mathbb{R}^{d_1}, \text{Card}(\{x_1\} \times \mathbb{R}^{d_2} \cap \mathcal{M}) = 1\},$$

and

$$\mathcal{B}_2 = \{\mathcal{M} \subset \mathbb{R}^D \text{ compact} \mid \exists x_2 \in \mathbb{R}^{d_2}, \text{Card}(\mathbb{R}^{d_1} \times \{x_2\} \cap \mathcal{M}) = 1\}.$$

Corollary 2. *Let \mathcal{M} be a subset of \mathbb{R}^D such that $\mathcal{M} \in \mathcal{B}_1 \cap \mathcal{B}_2$. If X is a random variable with support \mathcal{M} , then X satisfies A(1), (H1) and (H2).*

The proof of Corollary 2 is detailed in Section B.3.

We now give some illustrations of when Corollaries 1 and 2 hold.

First, any closed Euclidean ball, and more generally any strictly convex compact set in \mathbb{R}^D , is in $\mathcal{B}_1 \cap \mathcal{B}_2$. To see this, consider the points of the set with maximal first (resp. last) coordinate: they are unique by strict convexity, which ensures that the set is in \mathcal{B}_1 (resp. \mathcal{B}_2). The same holds for any compact set with strictly convex hull.

For non-compact supports, it is possible to satisfy Corollary 1 but not be in $\mathcal{B}_1 \cap \mathcal{B}_2$: Figure 2 shows a situation where the sets A_Δ and B_Δ cannot be nested, since there is no single “pinching point” as in \mathcal{B}_1 and \mathcal{B}_2 .

2.4 Genericity of (H1) and (H2)

Write $\mathcal{A} = (\cap_{\Delta>0} \cup_{\varepsilon>0} \mathcal{A}_1(\Delta, \varepsilon)) \cap (\cap_{\Delta>0} \cup_{\varepsilon>0} \mathcal{A}_2(\Delta, \varepsilon))$ the set of supports we introduced in Corollary 1.

We are interested in understanding whether signals “almost always” have a support in \mathcal{A} . Several notions of genericity exist when one wants to show that a property is “almost always” true.




Figure 1: Illustration of (H1) and \mathcal{A}_1 . As A_Δ narrows on the “tip” of the support, B_Δ can be taken smaller and smaller. This narrowing as A_Δ tends to the extremal point of the support is why \mathcal{B}_1 holds for compact supports with strictly convex convex hull.




Figure 2: Illustration of (H1) and \mathcal{A}_1 for a non-compact support. As A_Δ goes to infinity and becomes thinner, so does B_Δ .

One approach is to show that the property holds almost surely in a probabilistic sense. The issue with this is that we need a measure on the closed subsets of \mathbb{R}^D , of which there is no canonical example. As an illustration, we construct a small, random perturbation of the Euclidean space (based on partitioning it into arbitrarily small simplices which are then moved at random by a small amount). When applied to a compact set, this perturbation produces a random variable taking values in the compact subsets of \mathbb{R}^D . We show that almost surely, regardless of the initial support, this new support satisfies Corollary 2. The precise construction of this transformation and the statement of the probabilistic genericity of Corollary 2 are detailed in Section B.4.

Another approach of a more topological nature is to show that the set \mathcal{A} , seen as a subset of the metric space of all closed subsets of \mathbb{R}^D endowed with the Hausdorff distance, is a dense G_δ set (*i.e.* it is a dense, countable intersection of open sets), or in other words, is comeagre.

This property follows from the fact that the sets $\cup_{\varepsilon>0} \mathcal{A}_1(\Delta, \varepsilon)$ are open with respect to the Hausdorff distance, and are increasing in Δ . Therefore, \mathcal{A} is a countable intersection of open sets (*i.e.* a G_δ set) which is in addition dense (see Proposition 3 from Section B.5).

Equivalently, this means that the complement of \mathcal{A} is a meagre set, and therefore the set of all supports satisfying (H1) and (H2) is comeagre. Do note that while this notion ensures that \mathcal{A} is dense in the set of all closed subsets of \mathbb{R}^D , its complement is also dense, see Proposition 3 as well.

3 The estimation procedures

3.1 Overview

The identifiability Theorem 1 is the base upon which our estimators are built. A key part of its proof is the following result: if a multivariate analytic function ϕ satisfies $\phi(0) = 1$, $\phi(t) = \phi(-t)$ for all t , as well as assumptions $A(\kappa)$ for some $\kappa > 1/2$, (Adep) and

$$\phi(t_1, t_2)\Phi_X(t_1, 0)\Phi_X(0, t_2) = \Phi_X(t_1, t_2)\phi(t_1, 0)\phi(0, t_2)$$

for all (t_1, t_2) in a neighborhood of zero in $\mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$, then $\phi = \Phi_X$. In particular, fixing some $\nu_{\text{est}} > 0$, the only function ϕ such that

$$\int_{B_{\nu_{\text{est}}}^{d_1} \times B_{\nu_{\text{est}}}^{d_2}} |\phi(t_1, t_2)\Phi_X(t_1, 0)\Phi_X(0, t_2) - \Phi_X(t_1, t_2)\phi(t_1, 0)\phi(0, t_2)|^2 \times |\Phi_{\varepsilon^{(1)}}(t_1)\Phi_{\varepsilon^{(2)}}(t_2)|^2 dt_1 dt_2 = 0$$

is Φ_X . Indeed the characteristic functions of the noises $\Phi_{\varepsilon^{(i)}}$, $i \in \{1, 2\}$, are continuous and equal to 1 in zero. Moreover, the introduction of these characteristic functions allow to combine them with the characteristic functions Φ_X of the signal to get the characteristic functions of the observations, so that we are able to construct an empirical criterion M_n that estimates the above integral, see (4). Finally, following the usual M -estimation procedure, we then construct an estimator $\hat{\phi}$ of Φ_X by minimizing M_n over some set of multivariate analytic functions. Note that the resulting $\hat{\phi}$ is in general not the characteristic function of a probability distribution.

A difference between our estimator and the one from [17] is that we do not directly use $\hat{\phi}$ to estimate the distribution of the signal. Indeed, while there is always a one-to-one correspondence between the distribution G and its characteristic function Φ_X , the inverse Fourier transform approach to recover G from Φ_X breaks down when G is singular. However, it is always possible to recover the convolution $G * \Psi$ of G by Ψ for any properly chosen kernel Ψ . Moreover, when Ψ is an approximation of the unity, the distribution of Z will be close to the Dirac in zero, thus $X + Z$ will concentrate close to the support of X . In Section 4.1, we propose an estimator of the support as an upper-level set of an estimated density following ideas of [20], the main difference being with the smoothing kernel we choose. Indeed, with this kernel, no prior knowledge on the intrinsic dimension is needed to build the estimator. The precise choice of Ψ is non trivial and is given in Section 4.1.

Proposition 2 directly ensures that the estimator

$$\hat{g} := \mathcal{F}^{-1} [\hat{\phi} \cdot \mathcal{F}[\Psi]]$$

is close to $\bar{g} := \mathcal{F}^{-1} [\Phi_X \cdot \mathcal{F}[\Psi]] = G * \Psi$, whose upper level sets are close to \mathcal{M}_G . Here, for any integrable function f from \mathbb{R}^D to \mathbb{R} , we denote by $\mathcal{F}[f]$ (resp. $\mathcal{F}^{-1}[f]$) the (resp. inverse) Fourier transform of f defined, for all $y \in \mathbb{R}^d$, by

$$\mathcal{F}[f](y) = \int e^{it^\top y} f(t) dt \quad \text{and} \quad \mathcal{F}^{-1}[f](y) = \left(\frac{1}{2\pi}\right)^D \int e^{-it^\top y} f(t) dt.$$

Then, the estimator for the support of the signal is taken as the upper level set

$$\widehat{\mathcal{M}} = \{y : \hat{g}(y) > \lambda\},$$

where λ has to be well chosen. To obtain the best possible rates requires a further polynomial truncation of $\hat{\phi}$ before Fourier inversion, depending on the parameter κ . We then provide a model choice procedure to adapt in κ and get adaptive rates. Estimation of the distribution starts from the estimation of the support and follows similar ideas.

3.2 Estimation of the characteristic function

For any $S > 0$, let $\Upsilon_{\kappa,S}$ be the subset of multivariate analytic functions from \mathbb{C}^D to \mathbb{C} defined as follows.

$$\Upsilon_{\kappa,S} = \left\{ \phi \text{ analytic s.t. } \forall z \in \mathbb{R}^D, \overline{\phi(z)} = \phi(-z), \phi(0) = 1 \text{ and } \forall i \in \mathbb{N}^D \setminus \{0\}, \left| \frac{\partial^i \phi(0)}{\prod_{a=1}^d i_a!} \right| \leq \frac{S^{\|i\|_1}}{\|\dot{i}\|_1^{\kappa\|i\|_1}} \right\} \quad (3)$$

where $\|i\|_1 = \sum_{a=1}^D i_a$. If the distribution of X satisfies $A(\kappa)$, then there exists S such that $\Phi_X \in \Upsilon_{\kappa,S}$, and the converse also holds, see Lemma 3.1 in [17].

Let $\Phi_{\varepsilon^{(i)}}$ be the characteristic function of $\varepsilon^{(i)}$, $i = 1, 2$, and define for all $\phi \in \Upsilon_{\kappa,S}$ and any $\nu > 0$,

$$M(\phi; \nu | \Phi_X) = \int_{B_{\nu}^{d_1} \times B_{\nu}^{d_2}} |\phi(t_1, t_2) \Phi_X(t_1, 0) \Phi_X(0, t_2) - \Phi_X(t_1, t_2) \phi(t_1, 0) \phi(0, t_2)|^2 \times |\Phi_{\varepsilon^{(1)}}(t_1) \Phi_{\varepsilon^{(2)}}(t_2)|^2 dt_1 dt_2.$$

Fix some $\nu_{\text{est}} > 0$, and define M_n for any ϕ as follows

$$M_n(\phi) = \int_{B_{\nu_{\text{est}}}^{d_1} \times B_{\nu_{\text{est}}}^{d_2}} |\phi(t_1, t_2) \tilde{\phi}_n(t_1, 0) \tilde{\phi}_n(0, t_2) - \tilde{\phi}_n(t_1, t_2) \phi(t_1, 0) \phi(0, t_2)|^2 dt_1 dt_2, \quad (4)$$

where for all $(t_1, t_2) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$,

$$\tilde{\phi}_n(t_1, t_2) = \frac{1}{n} \sum_{\ell=1}^n \exp \left\{ it_1^\top Y_\ell^{(1)} + it_2^\top Y_\ell^{(2)} \right\}.$$

By the law of large numbers, $M_n(\phi)$ is a consistent estimator of $M(\phi; \nu_{\text{est}} | \Phi_X)$. As usual in M-estimation to get consistency of the estimator, we need to minimize M_n over a closed set of functions over which Φ_X is the only minimizer of $M(\cdot; \nu_{\text{est}} | \Phi_X)$. We thus introduce \mathcal{H} a subset of functions $\mathbb{C}^D \rightarrow \mathbb{C}^D$ such that all elements of \mathcal{H} satisfy (Adep) and such that the set of the restrictions to $B_{\nu_{\text{est}}}^D$ of functions in \mathcal{H} is closed in $L^2(B_{\nu_{\text{est}}}^D)$. Indeed we shall consider consistency of the estimator in $L^2(B_{\nu_{\text{est}}}^D)$, see Proposition 2 below. For instance, \mathcal{H} can be defined using submodels such as errors in variable regression with regression functions in a closed set of smooth functions when it is believed that the signal X belongs to this submodel. \mathcal{H} can also be defined using geometric constraints such as those described in Section 2.3, as soon as it is believed that the support of X satisfies those constraints since we need $\Phi_X \in \mathcal{H}$. We now introduce the following assumption.

(Amin) There exists $\alpha \in (0, 1)$ and $\nu_0 > 0$ such that for all $\nu \in (0, \nu_0]$, there exists $c(\nu, \alpha) > 0$ such that for all $\Phi_X \in \mathcal{H} \cap \Upsilon_{1/2,S}$ and for all $\phi \in \mathcal{H} \cap \Upsilon_{1/2,S}$ such that $\phi \neq \Phi_X$,

$$\left\| \frac{\phi}{\Phi_X} - \frac{\phi(\cdot, 0)}{\Phi_X(\cdot, 0)} - \frac{\phi(0, \cdot)}{\Phi_X(0, \cdot)} + 1 \right\|_{2,\nu} \geq c(\nu, \alpha) \max \left(\left\| \frac{\phi(\cdot, 0)}{\Phi_X(\cdot, 0)} - 1 \right\|_{2,\nu}, \left\| \frac{\phi(0, \cdot)}{\Phi_X(0, \cdot)} - 1 \right\|_{2,\nu} \right)^{1+\alpha},$$

where $f(\cdot, 0)$ denotes the function $t_1 \in \mathbb{R}^{d_1} \mapsto f(t_1, 0)$ and likewise $f(0, \cdot) : t_2 \in \mathbb{R}^{d_2} \mapsto f(0, t_2)$ for any function f on \mathbb{R}^D .

From now on, we assume that \mathcal{H} satisfies (Amin) for some fixed $\alpha \in (0, 1)$.

Let also $\mathbb{C}_m[X]$ be the set of polynomial functions of degree m in D indeterminates. For any integer m and any $\kappa < 1$, we define $\widehat{\Phi}_{n,m,\kappa}$ be a (up to $1/n$) measurable minimizer of the functional $\phi \mapsto M_n(\phi)$ over $\Upsilon_{\kappa,S} \cap \mathcal{H} \cap \mathbb{C}_m[X]$, that is

$$M_n(\widehat{\Phi}_{n,m,\kappa}) \leq \inf_{\phi \in \Upsilon_{\kappa,S} \cap \mathcal{H} \cap \mathbb{C}_m[X]} M_n(\phi) + \frac{1}{n}.$$

For good choices of m , $\widehat{\Phi}_{n,m,\kappa}$ is a consistent estimator of Φ_X in $L^2(B_\nu^D)$ at polynomial rate. The constants will depend on the signal through κ and S , and on the noise through its second moment and the following quantity:

$$c_\nu = \inf\{|\Phi_{\varepsilon^{(1)}}(t)|, t \in B_\nu^{d_1}\} \wedge \inf\{|\Phi_{\varepsilon^{(2)}}(t)|, t \in B_\nu^{d_2}\}. \quad (5)$$

Note that for any noise distribution, for small enough ν , c_ν is a positive real number. For any $\nu > 0$, $c(\nu) > 0$, $E > 0$, define $\mathcal{Q}^{(D)}(\nu, c(\nu), E)$ the set of distributions $\mathbb{Q} = \otimes_{j=1}^2 \mathbb{Q}_j$ on \mathbb{R}^D such that $c_\nu \geq c(\nu)$ and $\int_{\mathbb{R}^D} \|x\|^2 d\mathbb{Q}(x) \leq E$.

Proposition 2. *For all $\kappa_0 > 1/2$, $\nu \in (0, \nu_{est}]$, $S, c(\nu), E, C > 0$ and $\delta \in (0, 1)$, there exist positive constants c and n_0 such that the following holds: let $\kappa \in [\kappa_0, 1]$, for all $\Phi_X \in \Upsilon_{\kappa,S} \cap \mathcal{H}$ and $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$, for all $n \geq n_0$ and $s \geq 1$, with probability at least $1 - 2e^{-s}$,*

$$\sup_{\kappa' \in [\kappa_0, \kappa], m \in [\frac{2}{\kappa'} \frac{\log n}{\log \log n}, C \frac{\log n}{\log \log n}]} \int_{B_\nu^{d_1} \times B_\nu^{d_2}} |\widehat{\Phi}_{n,m,\kappa'}(t) - \Phi_X(t)|^2 dt \leq c \left(\frac{s}{n^{1/(1+2\alpha)}} \right)^{1-\delta}.$$

The proof of Proposition 2 is based on results in [8] and [17] and is detailed in Appendix C. For sake of simplicity, we denote $\widehat{\Phi}_{n,\kappa}$ the estimator $\widehat{\Phi}_{n,m,\kappa}$ in which $m = \lceil 4 \frac{\log n}{\log \log n} \rceil$. Note that with this value of m , we may apply the inequality in Proposition 2 uniformly for $\kappa' \in [\kappa_0, \kappa]$ whatever $\kappa_0 \in (1/2, 1]$ and $\kappa \in (\kappa_0, 1]$.

4 Estimation of the support

In Section 4.1, we provide an estimator of the support of the signal when κ is known, and prove an upper bound for the maximum risk in Hausdorff distance. Section 4.2 is devoted to the construction of an adaptive estimator of the support for unknown κ . In Section 4.3, we prove a lower bound which shows that our estimator is minimax up to some power of $\log \log n$ for all $\kappa \in (1/2, 1)$ and up to any small power of $\log n$ for $\kappa = 1$.

4.1 Upper bound

We shall consider the minimax risk in Hausdorff distance, which is defined, for A_1 and A_2 subsets of \mathbb{R}^D , as

$$d_H(A_1, A_2) = \sup_{x \in A_1 \cup A_2} |d(x, A_1) - d(x, A_2)|.$$

When $\kappa < 1$, the support of G is not compact. Since we allow the support to be a non-compact set, we define a truncated loss function as in [20]. We fix \mathcal{K} a compact subset of \mathbb{R}^D (which can be arbitrarily large) and for any S_1, S_2 subsets of \mathbb{R}^D , the truncated loss function is

$$H_{\mathcal{K}}(S_1, S_2) = d_H(S_1 \cap \mathcal{K}, S_2 \cap \mathcal{K}).$$

We now define the class over which we will prove an upper bound for the maximum risk in Hausdorff distance. Since consistency of $\widehat{\Phi}_{n,m,\kappa}$ is obtained for X such that $\Phi_X \in \mathcal{H} \cap \Upsilon_{\kappa,S}$ we summarize $\mathcal{L}(\kappa, S, \mathcal{H})$ as the set of distributions G such that, if X is a random variable with distribution G , then $\Phi_X \in \mathcal{H} \cap \Upsilon_{\kappa,S}$. Moreover, for any positive constants a, d and r_0 , we define $St_{\mathcal{K}}(a, d, r_0)$ as the set of positive measures G such that for all $x \in \mathcal{M}_G \cap \mathcal{K}$, for all $r \leq r_0$, $G(B(x, r)) \geq ar^d$. The distributions in $St_{\mathcal{K}}(a, d, r_0)$ are called (a, d) -standard. Such a class of distributions is commonly used for inferring topological information, see for instance [6].

Remark 1. • If a measure μ (for instance the d -dimensional Hausdorff measure on a manifold) is (a, d') -standard for some positive constants a and d' , and if G admits a density g with respect to μ such that g is lower bounded by $c > 0$, then G is (ac, d') -standard.

- We do not make any assumption on the reach of the support of G (see [15]) since it is not necessary here, although it provides a convenient way to check the (a, d) -standard assumption: if \mathcal{M}_G is a Riemannian manifold of dimension d with $\text{reach}(\mathcal{M}_G) \geq \tau_{\min} > 0$, then the d -dimensional Hausdorff measure restricted to \mathcal{M}_G is (a, d) -standard for some $a > 0$ (see Lemma 32 of [20]).

We now introduce the kernel we shall use for our construction. Fix $A > 0$ and define, for all $y \in \mathbb{R}$,

$$u_A(y) = \exp \left\{ -\frac{1}{(1-2y)^A} - \frac{1}{(1+2y)^A} \right\} 1_{[-\frac{1}{2}, \frac{1}{2}]}(y)$$

and for all $y \in \mathbb{R}^D$,

$$\psi_A(y) = I(A) \mathcal{F}^{-1}[u_A * u_A](\|y\|_2) \text{ with } I(A) = \frac{1}{\int \mathcal{F}^{-1}[u_A * u_A](\|x\|_2) dx}.$$

For $h > 0$ and $x \in \mathbb{R}^D$, we write $\psi_{A,h}(x) = h^{-D} \psi_A(x/h)$, hence $\mathcal{F}[\psi_{A,h}](t) = \mathcal{F}[\psi_A](th)$. Note that $\mathcal{F}[\psi_A]$ has compact support, so we may define the function \bar{g} by

$$\forall y \in \mathbb{R}^D, \bar{g}(y) = \left(\frac{1}{2\pi} \right)^D \int e^{-it^\top y} \mathcal{F}[\psi_A](ht) \Phi_X(t) dt,$$

which may be rewritten using usual Fourier calculus, for all $y \in \mathbb{R}^D$, as

$$\bar{g}(y) = (\psi_{A,h} * G)(y) = \frac{1}{h^D} \int_{\mathbb{R}^D} \psi_A \left(\frac{\|y-u\|_2}{h} \right) dG(u).$$

The density \bar{g} is a kernel smoothing of the distribution G . The bandwidth parameter h will be chosen appropriately in Theorem 3 below. We now construct an estimator of \bar{g} by truncating $\widehat{\Phi}_{n,\kappa}$ depending on κ . Adaptation with respect to κ is handled in Section 4.2. For some integer $m_\kappa > 0$ to be chosen later, let

$$\forall y \in \mathbb{R}^D, \widehat{g}_{n,\kappa}(y) = \left(\frac{1}{2\pi} \right)^D \int e^{-it^\top y} \mathcal{F}[\psi_A](ht) T_{m_\kappa} \widehat{\Phi}_{n,\kappa}(t) dt,$$

in which for any multivariate analytic function ϕ defined in a neighborhood of 0 in \mathbb{C}^D as $\phi : x \mapsto \sum_{(i_1, \dots, i_D) \in \mathbb{N}^D} c_i \prod_{a=1}^D x_a^{i_a}$, its truncation on $\mathbb{C}_m[X]$ for any integer m is

$$T_m \phi : x \mapsto \sum_{(i_1, \dots, i_D) \in \mathbb{N}^D, i_1 + \dots + i_D \leq m} c_i \prod_{a=1}^D x_a^{i_a}.$$

Since for all $t \in \mathbb{R}^D$, $T_{m_\kappa} \widehat{\Phi}_{n,\kappa}(-t) = T_{m_\kappa} \widehat{\Phi}_{n,\kappa}(t)$, the function $\widehat{g}_{n,\kappa}$ is real valued. Finally, define an estimator of the support of the signal as the upper level set

$$\widehat{\mathcal{M}}_\kappa = \{y \in \mathbb{R}^D \mid \widehat{g}_{n,\kappa}(y) > \lambda_{n,\kappa}\},$$

for some $\lambda_{n,\kappa}$. The main theorem of this section gives an upper bound of the maximum risk.

Theorem 3. Let $\kappa \in (1/2, 1]$, $a > 0$, $d \leq D$, $r_0 > 0$. For $c_h \geq \exp(2D+2)$ and $\ell \in (0, 1)$, define m_κ and h as

$$m_\kappa = \left\lfloor \frac{1}{6\kappa} \frac{\log(n)}{\log \log(n)} \right\rfloor, \quad h = c_h S m_\kappa^{-\kappa}$$

and $\lambda_{n,\kappa}$ depending whether $d < D$ or $d = D$ as

- if $d < D$,

$$\lambda_{n,\kappa} = \left(\frac{1}{h}\right)^\ell,$$

- if $d = D$,

$$\lambda_{n,\kappa} = \frac{1}{4}ac_A^D d_A.$$

Then for any $\kappa_0 \in (1/2, 1]$, $\nu \in (0, \nu_{est}]$, $c(\nu) > 0$, $E > 0$, $S > 0$, there exists n_0 and $C > 0$ such that for all $n \geq n_0$,

$$\sup_{\kappa \in [\kappa_0, 1]} \sup_{\substack{G \in \text{St}_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H}) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \frac{\log(n)^\kappa}{\log(\log(n))^{\kappa + \frac{A+1}{A}}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_\kappa)] \leq C.$$

Remark 2. • We prove in the next section a nearly matching lower bound. Thus, the minimax rate of convergence of the support in truncated Hausdorff distance depends on κ , that is on the way the distribution of the signal behaves at infinity. This rate deteriorates when the distribution of the signal has heavier tails. Indeed, since the distribution of the noise is unknown, taking into account distant observation points to build the estimator of the support becomes more difficult.

- When $d < D$, thanks to the use of the kernel ψ_A , our estimator does not require the knowledge of d , which has to be compared with the estimator in [20] where prior knowledge of d is needed.
- In [20], the upper bound on the rate is of order $1/\sqrt{\log n}$. Here we get a bound of order $1/(\log n)^\kappa$ depending on the tail of the distribution of the signal. We do not need to know the distribution of the noise, contrarily to [20] where the distribution of the noise is used in the construction of the estimator, as usual in the classical deconvolution literature.
- It may be seen from the proof of Theorem 3 that the choice $\lambda_{n,\kappa} = \frac{1}{4}ac_A^D d_A$ is valid for any d . However, this requires the knowledge of a .
- Note that there are two truncation steps: the first one is implicit in the construction of $\widehat{\Phi}_{n,\kappa}$ (chosen at the end of Section 3.2) and the second one appears in the definition of $\widehat{g}_{n,\kappa}$. This second truncation is necessary to control the error of $\widehat{\Phi}_{n,\kappa}$ on $B_{1/h}^D$ (instead of the error on B_ν^D in Proposition 2), and the degree m_κ in the second truncation is always smaller than the degree m used in the construction of $\widehat{\Phi}_{n,\kappa}$.
- The constant ℓ can be chosen arbitrarily between 0 and 1.

The proof of Theorem 3 is detailed in Appendix D.1. As in [20], the idea is to lower bound \bar{g} on the support \mathcal{M}_G when the bandwidth parameter h becomes small, and to upper bound it on every points further than a small distance (depending on h) from that support, see Lemmas 3 and 4 in Appendix D.1.

4.2 Adaptation to unknown κ

We now propose a data-driven model selection procedure to select κ such that the resulting estimator has the right rate of convergence. As usual, the idea is to perform a bias-variance trade off. Although we have an upper bound for the variance term, the bias is not easily accessible. We will use Goldenshluger and Lepski's method, see [21]. The variance bound is given as follows:

$$\sigma_n(\kappa) = c_\sigma \frac{(\log \log n)^{\kappa + \frac{A+1}{A}}}{(\log n)^\kappa}.$$

Fix some $\kappa_0 > 1/2$. The bias proxy is defined as

$$B_n(\kappa) = 0 \vee \sup_{\kappa' \in [\kappa_0, \kappa]} \left(H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\kappa'}) - \sigma_n(\kappa') \right).$$

The estimator of κ is now given by

$$\widehat{\kappa}_n \in \arg \min \{B_n(\kappa) + \sigma_n(\kappa), \kappa \in [\kappa_0, 1]\},$$

and the estimator of the support of the signal is $\widehat{\mathcal{M}}_{\widehat{\kappa}_n}$. The following theorem states that this estimator is rate adaptive.

Theorem 4. *For any $\kappa_0 \in (1/2, 1]$, $\nu \in (0, \nu_{est}]$, $c(\nu) > 0$, $E > 0$, $S > 0$, $a > 0$, $d \leq D$, there exists $c_\sigma > 0$ such that*

$$\limsup_{n \rightarrow +\infty} \sup_{\kappa \in [\kappa_0, 1]} \sup_{\substack{G \in St_{\mathcal{K}}(a, d) \cap \mathcal{L}(\kappa, S, \mathcal{H}) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \frac{\log(n)^\kappa}{\log(\log(n))^{\kappa + \frac{A+1}{A}}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\widehat{\kappa}_n})] < +\infty.$$

The proof of Theorem 4 is detailed in Appendix D.5.

4.3 Lower bound

The aim of this subsection is to prove a lower bound for the minimax risk of the estimation of \mathcal{M}_G using the distance $H_{\mathcal{K}}$ as loss function. The proof of Theorem 5 is based on Le Cam's two-points method, see [27], one of the most widespread technique to derive lower bounds. Note that we can not use the lower bound proved in [20] since the two distributions they use for the signal X in their two-points proof have Gaussian tails, for which $\kappa = 1/2$.

Theorem 5. *For any $\kappa \in (1/2, 1)$, there exists $S_\kappa > 0$, $a_\kappa > 0$ and \mathcal{H}_κ^* a set of complex functions satisfying (Adep) such that the set of the restrictions of its elements to $[-\nu, \nu]^D$ is closed in $L_2([-\nu, \nu]^D)$ for any $\nu > 0$, and such that for all $S \geq S_\kappa$, $a \leq a_\kappa$, $d \geq 1$, $0 < r_0 < 1$, $E > 0$ and $\nu \in (0, \nu_{est}]$ such that $c(\nu) > 0$, there exists $C > 0$ depending only on a , D , S , E and ν , and there exists n_0 , such that for all $n \geq n_0$,*

$$\inf_{\widehat{\mathcal{M}}} \sup_{\substack{G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H}_\kappa^*) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}})] \geq \frac{C}{\log(n)^\kappa}, \quad (6)$$

and for any $\delta \in (0, 1)$, there exists $C > 0$ depending only on a , D , S , E , ν and δ , and there exists n_0 , such that for all $n \geq n_0$,

$$\inf_{\widehat{\mathcal{M}}} \sup_{\substack{G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(1, S, \mathcal{H}_1^*) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}})] \geq \frac{C}{\log(n)^{1+\delta}}, \quad (7)$$

where the infimum in (6) and (7) is taken over all possible estimators $\widehat{\mathcal{M}}$ of \mathcal{M}_G .

Remark 3. • *The lower bound in Theorem 5 almost matches the upper bound for the maximum risk of our estimator in Theorem 3. Thus our work identifies the main factor in the minimax rate for the estimation of the support in Hausdorff loss. Notice that (7) is weaker than (6) with $\kappa = 1$, but we were not able to find a f_κ suitable to prove (6) with $\kappa = 1$, see Lemma 7 in Appendix D.6.*

- *In [20], the lower bound does not match the upper bound by a larger power in the rate (almost twice).*
- *The sets of supports we consider are not the same as that considered in [20]. In [20], the authors assume that the support is a regular manifold with lower bounded reach. We do not assume regularity, we only assume that the distribution of the signal is (a, d) -standard.*

The proof of Theorem 5 is detailed in Appendix D.6.

5 Estimation of the distribution of the signal

In this section, we assume that the support \mathcal{M}_G of G is a compact subset of \mathbb{R}^D . For any $\eta > 0$ and $A \subset \mathbb{R}^D$, A_η will denote the η -offset of A , that is the set of all points x in \mathbb{R}^D such that $d(x, A) \leq \eta$.

To estimate G , we shall consider the probability density \bar{g} defined in Section 4.1 and define the probability distribution $P_{\psi_{A, h_n}}$ on \mathbb{R}^D such that, for any \mathcal{O} borelian set of \mathbb{R}^D ,

$$P_{\psi_{A, h_n}}(\mathcal{O}) = \int_{\mathcal{O}} \bar{g}(y) dy.$$

We then estimate $P_{\psi_{A, h_n}}$ using the estimation of \bar{g} defined in Section 4.1 for $\kappa = 1$, $\hat{g}_n := \hat{g}_{n,1}$. Since \hat{g}_n can be non positive, we use $\hat{g}_n^+ = \max\{0, \hat{g}_n\}$ and renormalize it to get a probability distribution. We shall also estimate $P_{\psi_{A, h_n}}$ with a probability distribution having support on a (small) offset of the estimated support $\widehat{\mathcal{M}}$ restricted to the closed euclidean ball $\bar{B}(0, R_n)$, for some radius R_n that grows to infinity with n . Thus we fix some $\eta > 0$ and define $\hat{P}_{n, \eta}$ such that, for any \mathcal{O} borelian set of \mathbb{R}^D ,

$$\hat{P}_{n, \eta}(\mathcal{O}) = \frac{1}{\int_{(\widehat{\mathcal{M}} \cap \bar{B}(0, R_n))_\eta} \hat{g}_n^+(y) dy} \int_{\mathcal{O} \cap (\widehat{\mathcal{M}} \cap \bar{B}(0, R_n))_\eta} \hat{g}_n^+(y) dy = c_n \int_{\mathcal{O} \cap (\widehat{\mathcal{M}} \cap \bar{B}(0, R_n))_\eta} \hat{g}_n^+(y) dy.$$

5.1 Upper bound for the Wasserstein risk

The aim of this subsection is to give an upper bound of the Wasserstein maximum risk for the estimation of G . For any $p \in [1, +\infty)$ and any two probability measures μ and ν on \mathbb{R}^D , we write $W_p(\mu, \nu)$ the Wasserstein distance of order p between μ and ν , that is

$$W_p(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \left(\int_{\mathbb{R}^D \times \mathbb{R}^D} \|x - y\|_2^p d\pi(x, y) \right)^{1/p},$$

where $\Pi(\mu, \nu)$ is the set of probability measures on $\mathbb{R}^D \times \mathbb{R}^D$ that have marginals μ and ν .

Theorem 6. *For all $\nu \in (0, \nu_{est}]$, $c(\nu) > 0$, $E > 0$, $S > 0$, $\eta > 0$, $a > 0$, $r_0 > 0$, $d \leq D$, $p \in [1, +\infty)$, define m_n , h_n and λ_n as in Theorem 3 for $\kappa = 1$. Assume that $\lim_{n \rightarrow +\infty} R_n = +\infty$ and that there exists $\delta \in (0, \frac{1}{2})$ such that $R_n \leq \exp(n^{1/2-\delta})$. Then there exist n_0 and $C > 0$ such that for all $n \geq n_0$,*

$$\sup_{\substack{G \in \text{St}_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(1, S, \mathcal{H}) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [W_p(G, \hat{P}_{n, \eta})] \leq C \frac{\log \log(n)}{\log(n)}.$$

The proof of Theorem 6 is detailed in Appendix E.1. Note that the magnitude of η does not appear to be crucial when looking at the proof, at least in an asymptotic perspective. Considering an offset of size η of $\widehat{\mathcal{M}}$ is meant to control the variance of the estimator $\hat{P}_{n, \eta}$.

Remark 4. • *The lower bound in Theorem 7 almost matches the upper bound for the maximum risk of our estimator in Theorem 6. Thus our work identifies the main factor in the minimax rate for the estimation of the distribution in Wasserstein loss.*

- *Comparison with earlier results in the deconvolution setting [11] or [10] with known noise is not easy since the classes of signals they consider is much different than the ones we consider.*

5.2 Lower bound for the Wasserstein risk

The aim of this subsection is to establish a lower bound for the minimax Wasserstein risk of order p for any $p \geq 1$. Again, we can not use previous lower bounds proved in [11] or [10] since they use in the two-points method signals with distributions having too heavy tails.

Theorem 7. For any $p \geq 1$, there exists $S_1 > 0$, $a_1 > 0$ and \mathcal{H}_1^* a set of complex functions satisfying (Adep) such that the set of the restrictions of its elements to $[-\nu, \nu]^D$ is closed in $L_2([-\nu, \nu]^D)$ for any $\nu > 0$, and such that for all $S \geq S_1$, $a \leq a_1$, $d \geq 1$, $0 < r_0 < 1$, $E > 0$ and $\nu \in (0, \nu_{est}]$ such that $c(\nu) > 0$, there exists $C > 0$ depending only on a , D , S , E and ν , and there exists n_0 , such that for all $n \geq n_0$,

$$\inf_{\widehat{P}_n} \sup_{\substack{G \in \text{St}_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(1, S, \mathcal{H}_1^*) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \mathbb{E}_{(G^* \mathbb{Q})^{\otimes n}} [W_p(G, \widehat{P}_n)] \geq C \frac{1}{\log(n)^{1+\delta}},$$

where the infimum is taken on all possible estimate \widehat{P}_n of G .

As for Theorem 5, we use Le Cam's two-points method with the same two distributions $G_0(1)$ and $G_1(1)$. The proof essentially consists in showing that there exists a constant $C > 0$ independent of γ such that $W_p(G_0(1), G_1(1)) \geq CH_{\mathcal{K}}(M_0(\gamma), M_1(\gamma))$, that is $W_p(G_0(1), G_1(1)) \geq C\gamma$ for a constant $C > 0$. Once such an equality is established, the lower bound follows from taking γ as for Theorem 5.

The rest of the proof is detailed in Appendix E.2.

6 Acknowledgments

Jeremie Capitaio-Miniconi would like to acknowledge support from the UDOPIA-ANR-20-THIA-0013. Élisabeth Gassiat would like to acknowledge the Institut Universitaire de France and the ANR ASCAI: ANR-21-CE23-0035-02. This work was supported by the French government, through the UCA^{Jedi} and 3IA Côte d'Azur Investissements d'Avenir managed by the National Research Agency (ANR-15-IDEX-01 and ANR-19-P3IA-0002).

A Summary of notations

To help the reader, this section gathers the notations introduced in this paper, along with where they are defined.

- For any $r > 0$, we write $B_r = (-r, r)$ and for any measurable function f on B_r^D , we write $\|f\|_{\infty, r}$ the essential supremum of f over B_r^D and $\|f\|_{2, r} = (\int_{B_r^D} |f(u)|^2 du)^{1/2}$. Section 1.3.
- Φ_X : the characteristic function of X , $\Phi_{\varepsilon^{(i)}}$: the characteristic function of $\varepsilon^{(i)}$. Section 2.2.
- Assumption A(κ): There exist $a, b > 0$ such that for all $\lambda \in \mathbb{R}^D$, $\mathbb{E}[\exp(\lambda^\top X)] \leq a \exp(b \|\lambda\|_2^{1/\kappa})$. Section 2.2.
- Assumption (Adep): For any $z_0 \in \mathbb{C}^{d_1}$, $z \mapsto \Phi_X(z_0, z)$ is not the null function and for any $z_0 \in \mathbb{C}^{d_2}$, $z \mapsto \Phi_X(z, z_0)$ is not the null function. Section 2.2.
- \mathcal{M}_G : support of the distribution G . Section 2.2.
- For all $(t_1, t_2) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$, $\tilde{\phi}_n(t_1, t_2) = \frac{1}{n} \sum_{\ell=1}^n \exp\{it_1^\top Y_\ell^{(1)} + it_2^\top Y_\ell^{(2)}\}$. Section 3.2.
- $M(\phi; \nu | \Phi_X)$ and $M_n(\phi)$, Section 3.2:

$$M(\phi; \nu | \Phi_X) = \int_{B_\nu^{d_1} \times B_\nu^{d_2}} |\phi(t_1, t_2) \Phi_X(t_1, 0) \Phi_X(0, t_2) - \Phi_X(t_1, t_2) \phi(t_1, 0) \phi(0, t_2)|^2 \times |\Phi_{\varepsilon^{(1)}}(t_1) \Phi_{\varepsilon^{(2)}}(t_2)|^2 dt_1 dt_2,$$

$$M_n(\phi) = \int_{B_{\nu_{est}}^{d_1} \times B_{\nu_{est}}^{d_2}} |\phi(t_1, t_2) \tilde{\phi}_n(t_1, 0) \tilde{\phi}_n(0, t_2) - \tilde{\phi}_n(t_1, t_2) \phi(t_1, 0) \phi(0, t_2)|^2 dt_1 dt_2.$$

- For $S > 0$ and $\kappa > 0$, $\Upsilon_{\kappa,S} = \{\phi \text{ analytic s.t. } \forall z \in \mathbb{R}^D, \overline{\phi(z)} = \phi(-z), \phi(0) = 1 \text{ and } \forall i \in \mathbb{N}^D \setminus \{0\}, \left| \frac{\partial^i \phi(0)}{\prod_{a=1}^D i_a!} \right| \leq \frac{S^{\|i\|_1}}{\|i\|_1^{\kappa \|i\|_1}}\}$. Section 3.2.
- \mathcal{H} : a subset of functions $\mathbb{C}^D \rightarrow \mathbb{C}^D$ such that all elements of \mathcal{H} satisfy (Adep) and such that the set of the restrictions to $[-\nu_{\text{est}}, \nu_{\text{est}}]^D$ of functions in \mathcal{H} is closed in $L^2([-\nu_{\text{est}}, \nu_{\text{est}}]^D)$. Section 3.2.
- For any integer m , $\mathbb{C}_m[X]$ is the set of polynomial functions of degree m in D indeterminates with complex coefficients. Section 3.2.
- For any integer m and any $\kappa < 1$, $\widehat{\Phi}_{n,m,\kappa}$ is a (up to $1/n$) measurable minimizer of the functional $\phi \mapsto M_n(\phi)$ over $\Upsilon_{\kappa,S} \cap \mathcal{H} \cap \mathbb{C}_m[X]$. Section 3.2.
- $c_\nu = \inf\{|\Phi_{\varepsilon(1)}(t)|, t \in [-\nu, \nu]^{d_1}\} \wedge \inf\{|\Phi_{\varepsilon(2)}(t)|, t \in [-\nu, \nu]^{d_2}\}$. Section 3.2.
- For any $\nu > 0$, $c(\nu) > 0$, $E > 0$, $\mathcal{Q}^{(D)}(\nu, c(\nu), E)$ is the set of distributions $\mathbb{Q} = \otimes_{j=1}^2 \mathbb{Q}_j$ on \mathbb{R}^D such that $c_\nu \geq c(\nu)$ and $\int_{\mathbb{R}^D} \|x\|^2 d\mathbb{Q}(x) \leq E$. Section 3.2.
- $d_H(A_1, A_2)$ is the Hausdorff distance between A_1 and A_2 subsets of \mathbb{R}^D . Section 4.1.
- \mathcal{K} : a compact subset of \mathbb{R}^D (which can be arbitrarily large). Section 4.1.
- For all S_1, S_2 subsets of \mathbb{R}^D , $H_{\mathcal{K}}(S_1, S_2) = d_H(S_1 \cap \mathcal{K}, S_2 \cap \mathcal{K})$. Section 4.1.
- $St_{\mathcal{K}}(a, d, r_0)$: set of positive measures G such that for all $x \in \mathcal{M}_G \cap \mathcal{K}$, for all $r \leq r_0$, $G(B(x, r)) \geq ar^d$. The distributions in $St_{\mathcal{K}}(a, d, r_0)$ are called (a, d) -standard. Section 4.1.
- $\mathcal{L}(\kappa, S, \mathcal{H})$: set of distributions G such that, if X is a random variable with distribution G , then $\Phi_X \in \mathcal{H} \cap \Upsilon_{\kappa,S}$. Section 4.1.
- Fourier and inverse Fourier transform : when f is an integrable function from \mathbb{R}^D to \mathbb{R} , for all $y \in \mathbb{R}^D$, $\mathcal{F}[f](y) = \int e^{it^\top y} f(t) dt$ and $\mathcal{F}^{-1}[f](y) = (\frac{1}{2\pi})^D \int e^{-it^\top y} f(t) dt$. End of Section 3.1.
- For any $A > 0$, for all $y \in \mathbb{R}$, $u_A(y) = \exp\left\{-\frac{1}{(1-2y)^A} - \frac{1}{(1+2y)^A}\right\} 1_{[-\frac{1}{2}, \frac{1}{2}]}(y)$. Section 4.1.
- For any $A > 0$ and $y \in \mathbb{R}^D$, $\psi_A(y) = I(A) \mathcal{F}^{-1}[u_A * u_A](\|y\|_2)$, with $I(A) = \int \frac{1}{\mathcal{F}^{-1}[u_A * u_A](\|x\|_2)} dx$ and $\psi_{A,h}(x) = h^{-D} \psi_A(x/h)$. Section 4.1.
- The constants $c_A > 0$ and $d_A > 0$: for all $x \in \{y \in \mathbb{R}^D : \|y\|_2 \leq c_A\}$, $\psi_A(x) \geq d_A$. Section 4.1.
- For any multivariate analytic function ϕ defined in a neighborhood of 0 in \mathbb{C}^D as $\phi : x \mapsto \sum_{(i_1, \dots, i_D) \in \mathbb{N}^D} c_i \prod_{a=1}^D x_a^{i_a}$, for any integer m , its truncation on $\mathbb{C}_m[X]$ is $T_m \phi : x \mapsto \sum_{(i_1, \dots, i_D) \in \mathbb{N}^D, i_1 + \dots + i_D \leq m} c_i \prod_{a=1}^D x_a^{i_a}$. Section 4.1.
- For all $y \in \mathbb{R}^D$, $\bar{g}(y) = (\frac{1}{2\pi})^D \int e^{-it^\top y} \mathcal{F}[\psi_A](ht) \Phi_X(t) dt$. Section 4.1.
- For all $y \in \mathbb{R}^d$, $\widehat{g}_{n,\kappa}(y) = (\frac{1}{2\pi})^D \int e^{-it^\top y} \mathcal{F}[\psi_A](ht) T_{m,\kappa} \widehat{\Phi}_{n,\kappa}(t) dt$. Section 4.1.
- $\widehat{\mathcal{M}}_\kappa = \{y \in \mathbb{R}^D \mid \widehat{g}_{n,\kappa}(y) > \lambda_{n,\kappa}\}$. Section 4.1.
- $\Gamma_{n,\kappa} = \|\widehat{g}_{n,\kappa} - \bar{g}\|_\infty = \sup_{y \in \mathbb{R}^D} |\widehat{g}_{n,\kappa}(y) - \bar{g}(y)|$. Section 4.1.
- For any $\eta > 0$, A_η denotes the η -offset of A . Section 5.
- For any borelian set \mathcal{O} of \mathbb{R}^D , $P_{\psi_A, h_n}(\mathcal{O}) = \int_{\mathcal{O}} \bar{g}(y) dy$. Section 5.
- $\widehat{g}_n := \widehat{g}_{n,1}$ and $\widehat{g}_n^+ = \max\{0, \widehat{g}_n\}$. Section 5.

- For any borelian set \mathcal{O} of \mathbb{R}^D , $\widehat{P}_{n,\eta}(\mathcal{O}) = \frac{1}{\int_{(\widehat{\mathcal{M}} \cap \widehat{B}(0, R_n))_\eta} \widehat{g}_n^+(y) dy} \int_{\mathcal{O} \cap (\widehat{\mathcal{M}} \cap \widehat{B}(0, R_n))_\eta} \widehat{g}_n^+(y) dy = c_n \int_{\mathcal{O} \cap (\widehat{\mathcal{M}} \cap \widehat{B}(0, R_n))_\eta} \widehat{g}_n^+(y) dy$.
- For any $p \in [1, +\infty)$ and any two probability measures μ and ν on \mathbb{R}^D , $W_p(\mu, \nu)$ is the Wasserstein distance of order p between μ and ν . Section 5.1.
- For $k, l \in \{1, \dots, D\}$ with $k \leq l$, $\pi^{(k:l)} : (x_1, \dots, x_D) \in \mathbb{R}^D \mapsto (x_k, \dots, x_l) \in \mathbb{R}^{l-k+1}$ and $\pi^{(k)} = \pi^{(k:k)}$. Section 2.3.

B Proofs for Section 2

B.1 Proof of Proposition 1

Case $\kappa = 1$.

First, it is clear that any compactly supported distribution satisfies A(1). Conversely, if $\mathbb{E}[e^{\langle \lambda, X \rangle}] \leq a \exp(b \|\lambda\|_2)$, then for any $\mu > 0$, we get, for any $b' > b$, if we denote $(e_j)_{1 \leq j \leq D}$ the canonical basis of \mathbb{R}^D ,

$$\begin{aligned}
\mathbb{P}(\|X\|_2 \geq Db') &\leq \sum_{j=1}^D \mathbb{P}(|X_j| \geq b') \\
&= \sum_{j=1}^D \{\mathbb{P}(X_j \geq b') + \mathbb{P}(X_j \leq -b')\} \\
&= \sum_{j=1}^D \{\mathbb{P}(\langle \mu e_j, X \rangle \geq \mu b') + \mathbb{P}(-\langle \mu e_j, X \rangle \geq \mu b')\} \\
&\leq \sum_{j=1}^D \left\{ \frac{\mathbb{E}[\exp(\langle \mu e_j, X \rangle)]}{\exp(b'\mu)} + \frac{\mathbb{E}[\exp(-\langle \mu e_j, X \rangle)]}{\exp(b'\mu)} \right\} \quad \text{by Markov inequality} \\
&\leq 2D \frac{a \exp(b\mu)}{\exp(b'\mu)} \xrightarrow{\mu \rightarrow +\infty} 0,
\end{aligned}$$

and hence $\|X\|_2 \leq Db$ almost surely.

Case $\kappa < 1$.

Assume that for any $\lambda \in \mathbb{R}^D$, $\mathbb{E}[e^{\langle \lambda, X \rangle}] \leq a \exp(b \|\lambda\|_2^{1/\kappa})$ for some $a, b > 0$. Then by using the same directional method as for $\kappa = 1$, we get that for any $\mu, t \geq 0$,

$$\begin{aligned}
\mathbb{P}(\|X\|_2 \geq t) &\leq 2Da \exp(b\mu^{1/\kappa} - \mu t) \\
&= 2Da \exp\left(-\left(\frac{\kappa}{b}\right)^{\frac{\kappa}{1-\kappa}} (1-\kappa)t^{\frac{1}{1-\kappa}}\right) \quad \text{by taking } \mu = \left(\frac{\kappa t}{b}\right)^{\frac{\kappa}{1-\kappa}},
\end{aligned}$$

hence the result.

Now, assume that for any $t \geq 0$, $\mathbb{P}(\|X\|_2 \geq t) \leq c \exp(-dt^{1/(1-\kappa)})$ for some $c, d > 0$, then by the Cauchy-Schwarz inequality, for any $\lambda \in \mathbb{R}^D$,

$$\mathbb{E}[e^{\langle \lambda, X \rangle}] \leq \mathbb{E}[e^{\|\lambda\|_2 \|X\|_2}].$$

Then, using that for any nonnegative random variable Y , $\mathbb{E}[Y] = \int_{t \geq 0} \mathbb{P}(Y \geq t) dt$,

$$\begin{aligned} \mathbb{E}[e^{\langle \lambda, X \rangle}] &\leq 1 + \int_{t \geq 1} \mathbb{P}(e^{\|\lambda\|_2 \|X\|_2} \geq t) dt \\ &\leq 1 + \int_{s \geq 0} \mathbb{P}(\|X\|_2 \geq s) \|\lambda\|_2 \exp(\|\lambda\|_2 s) ds \quad \text{with } t = e^{\|\lambda\|_2 s} \\ &\leq 1 + c \|\lambda\|_2 \int_{s \geq 0} \exp(-ds^{\frac{1}{1-\kappa}} + \|\lambda\|_2 s) ds \\ &\leq 1 + c \|\lambda\|_2^{\frac{1}{\kappa}} \int_{s' \geq 0} \exp(\|\lambda\|_2^{\frac{1}{\kappa}} (-ds'^{\frac{1}{1-\kappa}} + s')) ds' \quad \text{with } s = s' \|\lambda\|_2^{\frac{1}{\kappa-1}}. \end{aligned}$$

Note that $-ds'^{\frac{1}{1-\kappa}} + s' \leq \kappa (\frac{1-\kappa}{d})^{1/\kappa-1}$ for any $s' \geq 0$, and $-ds'^{\frac{1}{1-\kappa}} + s' \leq -s'$ when $s' \geq (\frac{2}{d})^{1/\kappa-1}$. In particular,

$$\begin{aligned} \mathbb{E}[e^{\langle \lambda, X \rangle}] &\leq 1 + c \|\lambda\|_2^{\frac{1}{\kappa}} \int_{s'=0}^{(\frac{2}{d})^{1/\kappa-1}} \exp(\|\lambda\|_2^{\frac{1}{\kappa}} (-ds'^{\frac{1}{1-\kappa}} + s')) ds' \\ &\quad + c \|\lambda\|_2^{\frac{1}{\kappa}} \int_{s' \geq (\frac{2}{d})^{1/\kappa-1}} \exp(\|\lambda\|_2^{\frac{1}{\kappa}} (-ds'^{\frac{1}{1-\kappa}} + s')) ds' \\ &\leq 1 + c \|\lambda\|_2^{\frac{1}{\kappa}} \left(\frac{2}{d}\right)^{1/\kappa-1} \exp\left(\kappa \|\lambda\|_2^{\frac{1}{\kappa}} \left(\frac{1-\kappa}{d}\right)^{1/\kappa-1}\right) + c \exp\left(-\|\lambda\|_2^{\frac{1}{\kappa}} \left(\frac{2}{d}\right)^{1/\kappa-1}\right), \end{aligned}$$

which proves that $A(\kappa)$ holds.

B.2 Proof of Theorem 2

Consider a random variable X satisfying $A(\kappa)$. Theorem 2 is a direct consequence of the following Lemma 1. Indeed, for any $z_0 \in \mathbb{C}^{d_1}$ and $z \in \mathbb{C}^{d_2}$,

$$\mathbb{E} \left[\exp \left(iz_0^\top X^{(1)} + iz^\top X^{(2)} \right) \right] = \mathbb{E} \left[\mathbb{E} \left[\exp \left(iz_0^\top X^{(1)} \right) \mid X^{(2)} \right] \exp \left(iz^\top X^{(2)} \right) \right].$$

Let us show that $z \mapsto \mathbb{E}[\exp(iz_0^\top X^{(1)} + iz^\top X^{(2)})]$ is the null function if and only if the random variable $\mathbb{E}[\exp(iz_0^\top X^{(1)}) \mid X^{(2)}]$ is zero $\mathbb{P}_{X^{(2)}}$ -a.s. Define h the measurable function such that

$$\mathbb{E} \left[\exp \left(iz_0^\top X^{(1)} \right) \mid X^{(2)} \right] = h \left(X^{(2)} \right) \quad \mathbb{P}_{X^{(2)}} - a.s.,$$

and assume that for all $z \in \mathbb{C}^{d_2}$, $\mathbb{E}[h(X^{(2)}) \exp(iz^\top X^{(2)})] = 0$. Write $\Psi(h, iz)$ this expectation (which is zero for all $z \in \mathbb{C}^{d_2}$), and note that $\Psi(h, iz) = \Psi(\overline{h}, \overline{iz})$. As a consequence,

$$\mathbb{E} \left[\operatorname{Re}(h(X^{(2)})) \exp \left(iz^\top X^{(2)} \right) \right] = \frac{1}{2} \left(\Psi(h, iz) + \overline{\Psi(h, iz)} \right) = 0,$$

and likewise for $\operatorname{Im}(h)$. By replacing $h(X^{(2)})$ by its real or imaginary part, let us now assume that it is real-valued.

Denote for all $x \in \mathbb{R}^{d_2}$, $h_+(x) = \max\{h(x), 0\}$ and $h_-(x) = \max\{-h(x), 0\}$. We get for all $z \in \mathbb{C}^{d_2}$,

$$\mathbb{E} \left[h_+ \left(X^{(2)} \right) \exp \left(iz^\top X^{(2)} \right) \right] = \mathbb{E} \left[h_- \left(X^{(2)} \right) \exp \left(iz^\top X^{(2)} \right) \right].$$

If $h(X^{(2)})$ is not zero $\mathbb{P}_{X^{(2)}}$ -a.s., then the measures with density h_+ and h_- with respect to $\mathbb{P}_{X^{(2)}}$ have positive (and equal) total measure. After dividing by this total measure, we obtain two probability measures with equal characteristic functions, therefore they are the same probability distributions, so that $h_+(X^{(2)}) = h_-(X^{(2)})$, $\mathbb{P}_{X^{(2)}}$ -a.s., which implies $h(X^{(2)}) = 0$, $\mathbb{P}_{X^{(2)}}$ -a.s.

Likewise, for any $z_0 \in \mathbb{C}^{d_1}$, $z \mapsto \mathbb{E}[\exp(iz_0^\top X^{(1)} + iz^\top X^{(2)})]$ is the null function if and only if $\mathbb{E}[\exp(iz_0^\top X^{(1)}) \mid X^{(1)}]$ is zero $\mathbb{P}_{X^{(1)}}$ -a.s.

Lemma 1. *Assume (H1) and (H2). Then, for all $z \in \mathbb{C}^{d_1}$, $\mathbb{E}[\exp(iz^\top X^{(1)}) | X^{(2)}]$ is not $\mathbb{P}_{X^{(2)}}$ -a.s. the null random variable and for all $z \in \mathbb{C}^{d_2}$, $\mathbb{E}[\exp(iz^\top X^{(2)}) | X^{(1)}]$ is not $\mathbb{P}_{X^{(1)}}$ -a.s. the null random variable*

Proof of Lemma 1. Let $z \in \mathbb{C}^{d_1}$ be such that $\mathbb{E}[\exp(iz^\top X^{(1)}) | X^{(2)}]$ is $\mathbb{P}_{X^{(2)}}$ -a.s. the null random variable. Then for any $\Delta > 0$, if we denote A_Δ and B_Δ the sets given by (H1) and take $x_\Delta \in B_\Delta$,

$$\mathbb{E}[\exp(iz^\top (X^{(1)} - x_\Delta)) | X^{(2)}] 1_{X^{(2)} \in A_\Delta} = 0 \quad \mathbb{P}_{X^{(2)}}\text{-a.s.},$$

and taking the real part of this equation shows that

$$\mathbb{E}[\cos(\operatorname{Re}(z)^\top (X^{(1)} - x_\Delta)) \exp(-\operatorname{Im}(z)^\top (X^{(1)} - x_\Delta)) | X^{(2)}] 1_{X^{(2)} \in A_\Delta} = 0 \quad \mathbb{P}_{X^{(2)}}\text{ a.s.} \quad (8)$$

Using that $\operatorname{Diam}(B_\Delta) \rightarrow 0$ as $\Delta \rightarrow 0$ from (H1), we can fix $\Delta > 0$ small enough that if $x \in B_\Delta$, $\cos(\operatorname{Re}(z)^\top (x - x_\Delta)) > 0$. Since $\mathbb{P}(X^{(1)} \in B_\Delta | X^{(2)} \in A_\Delta) = 1$, this means that the expectation in (8) is positive, which is a contradiction. Thus $\mathbb{E}[\exp(iz^\top (X^{(1)} - x_\Delta)) | X^{(2)}]$ is not $\mathbb{P}_{X^{(2)}}$ -a.s. the null random variable.

The proof of the other part of Lemma 1 is analogous using (H2). □

B.3 Proof of Corollary 2

Let \mathcal{M} be a compact subset of \mathbb{R}^D . Let us first prove that the function $u \mapsto \operatorname{Diam}(\{u\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$ is upper semi-continuous.

Let $u \in \mathbb{R}^{d_1}$. Since \mathcal{M} is compact, there exist sequences $u_n \rightarrow u$ and (x_n, y_n) in $(\{u_n\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$ such that $\|x_n - y_n\|_2 = \operatorname{Diam}(\{u_n\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$ and $\lim_{n \rightarrow +\infty} \|x_n - y_n\|_2 = \limsup_{v \rightarrow u} \operatorname{Diam}(\{v\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$. Moreover, we may assume that there exists (x, y) in $(\{u\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$ such that $x_n \rightarrow x$ and $y_n \rightarrow y$. Taking the limit along those sequences shows that $\operatorname{Diam}(\{u\} \times \mathbb{R}^{d_2} \cap \mathcal{M}) \geq \|x - y\| = \limsup_{v \rightarrow u} \operatorname{Diam}(\{v\} \times \mathbb{R}^{d_2} \cap \mathcal{M})$, proving the claimed upper-semi continuity.

Now, since \mathcal{M} is compact, there exists $R > 0$ such that $\mathcal{M} \subset \bar{B}(0, R)$. If moreover $\mathcal{M} \in \mathcal{B}_1$, there exists $x_1 \in \mathbb{R}^{d_1}$ such that $\operatorname{Diam}(\{x_1\} \times \mathbb{R}^{d_2} \cap \mathcal{M}) = 0$. Using the upper semi-continuity shows that $\mathcal{M} \in \bigcap_{n \geq 1} \mathcal{A}_2(1/n, R)$. Likewise, if $\mathcal{M} \in \mathcal{B}_2$, there exists $x_2 \in \mathbb{R}^{d_2}$ such that $\operatorname{Diam}(\mathbb{R}^{d_1} \times \{x_2\} \cap \mathcal{M}) = 0$ and $\mathcal{M} \in \bigcap_{n \geq 1} \mathcal{A}_1(1/n, R)$.

The end of the proof follows from Corollary 1 and the fact that any random variable with compact support satisfies A(1).

B.4 Measure theoretical genericity and random perturbation of the space

In this section, we construct a random and small perturbation of \mathbb{R}^D and show that any compact set is almost surely transformed into a compact set in $\mathcal{B}_1 \cap \mathcal{B}_2$.

More precisely, for any $\varepsilon > 0$, we define a (random) continuous bijection $f : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that almost surely, $|f(x) - x| \leq \varepsilon$ for all $x \in \mathbb{R}^D$, and such that if \mathcal{M} is compact, then $f(\mathcal{M})$ is in $\mathcal{B}_1 \cap \mathcal{B}_2$ almost surely. This random bijection does not depend on which support \mathcal{M} is considered, and can for instance be seen as a modeling of the imperfections of “realistic” supports, or as a way to introduce a Bayesian prior on the support. In that sense, compact supports are almost surely in $\mathcal{B}_1 \cap \mathcal{B}_2$, and thus compactly supported random variables almost surely satisfy (Adep).

There is no canonical way to define a random perturbation of \mathbb{R}^D . Our approach is to tile the space with simplices, then add a small perturbation to each vertex of the tiling, keeping the transformation linear inside each simplex. Visually, this results in a small, random crumpling of the Euclidean space.

B.4.1 Simplicial tiling of \mathbb{R}^D

Let us recall a few definitions about simplicial complexes. For any $k \in \{0, \dots, D\}$, a k -simplex of \mathbb{R}^D is the convex hull of $(k+1)$ affinely independent points of \mathbb{R}^D . A simplicial complex \mathcal{P} is a set of simplices such that every face of a simplex from \mathcal{P} is also in \mathcal{P} , and the non-empty intersection of any two simplices $F_1, F_2 \in \mathcal{P}$ is a face of both F_1 and F_2 . \mathcal{P} is a homogeneous simplicial D -complex if each simplex of dimension less than D of \mathcal{P} is the face of a D -simplex of \mathcal{P} . For any simplex F , we write $\text{relint}(F)$ its relative interior. Finally, a homogeneous simplicial D -complex \mathcal{P} is called a simplicial tiling of $A \subset \mathbb{R}^D$ if the relative interior of its simplices form a partition of A . Note that the facets of \mathcal{P} , that is, its D -simplices, do not necessarily form a partition of A : two facets can have a non-empty intersection when they share a face.

First, consider a finite simplicial tiling of the hypercube $[0, 1]^D$, and extend it to \mathbb{R}^D by mirroring it along the hyperplanes orthogonal to the canonical axes crossing them at integer coordinates. Formally, for any $k = (k_1, \dots, k_D) \in \mathbb{Z}^D$, the hypercube $\prod_{i=1}^D [k_i, k_i + 1]$ contains the tiling of $[0, 1]^D$, mirrored along axis i if and only if k_i is odd. The faces of the hypercubes defined in this way match, as each pair of hypercubes sharing a face are mirrors of each other with respect to that face. Thus, the resulting tiling \mathcal{P} is a simplicial tiling of \mathbb{R}^D .

Let $(x_n)_{n \in \mathbb{N}}$ be the sequence of vertices of the simplicial tiling \mathcal{P} (i.e. its 0-simplices). We identify each simplex $F \in \mathcal{P}$ with the set of its 0-dimensional faces $\{x_i\}_{i \in I}$, and write F_I in that case. Note that the set I is unique for any given simplex F and characterizes F .

B.4.2 Perturbation of the tiling

Fix a small $r > 0$. Let $(\varepsilon_n)_{n \in \mathbb{N}}$ be a sequence of i.i.d. uniform variables on $[-r, r]^D$, and define \mathcal{P}^ε the simplicial complex defined by

$$\mathcal{P}^\varepsilon = \{\{x_i + \varepsilon_i\}_{i \in I} : \{x_i\}_{i \in I} \in \mathcal{P}\}.$$

Note that since the original tiling of $[0, 1]^D$ was finite, there exists $r_0 > 0$ such that for any $(\varepsilon_n)_{n \in \mathbb{N}} \in ([-r_0, r_0]^D)^{\mathbb{N}}$, the vertices of any simplex in \mathcal{P} are still affinely independent after being moved according to ε and any two simplices $F, F' \in \mathcal{P}$ sharing a face F'' (resp. with no intersection) are transformed into two simplices of \mathcal{P}^ε that share exactly the transformation of F'' (resp. with no intersection), so that \mathcal{P}^ε is indeed a simplicial complex. Finally, \mathcal{P}^ε still covers \mathbb{R}^D (as seen when moving each vertex in $[-1, 2]^D$ one after the other along a continuous path, showing that no hole is created in the covering of $[0, 1]^D$ at any point in time), so for any $r \in (0, r_0]$, \mathcal{P}^ε is almost surely a simplicial tiling of \mathbb{R}^D .

Since the relative interiors of the simplices of \mathcal{P} define a partition of \mathbb{R}^D , for each $z \in \mathbb{R}^D$, there exists exactly one face $F_I \in \mathcal{P}$ such that $z \in \text{relint}(F_I)$. Writing $z = \sum_{i \in I} \alpha_i x_i$ (for $\alpha \in (0, 1]^{|I|}$ such that $\sum_{i \in I} \alpha_i = 1$), we define the image of z by the perturbation as $f^\varepsilon(z) = \sum_{i \in I} \alpha_i (x_i + \varepsilon_i)$. In other words, each simplex is deformed according to the linear transformation given by the perturbation of its vertices.

The mapping f^ε is a (random) bijective and continuous transformation of \mathbb{R}^D that is “small”, in the sense that almost surely, $\sup_{z \in \mathbb{R}^D} \|z - f^\varepsilon(z)\| \leq r$.

Note that the transformation f^ε can be made with arbitrarily small granularity: the same approach works when considering tilings of $[0, \delta]^D$ for any $\delta > 0$ instead of $[0, 1]^D$ (up to changing r). We may also iterate several random independent transformations $f^{\varepsilon^{(1)}} \circ \dots \circ f^{\varepsilon^{(m)}}$ for $m \geq 1$, and the transformation of \mathcal{M} will still almost surely belong to $\mathcal{B}_1 \cap \mathcal{B}_2$.

Theorem 8. *Let $r \in (0, r_0]$ with r_0 as above, $\varepsilon = (\varepsilon_n)_{n \in \mathbb{N}}$ be a sequence of i.i.d. uniform r.v. on $[-r, r]^D$, $\delta > 0$, and f^ε be the bijective transformation of \mathbb{R}^D defined above.*

Then for any (random) continuous mapping $H : \mathbb{R}^D \rightarrow \mathbb{R}^D$ that is independent of ε , the mapping $F : z \mapsto \delta f^\varepsilon(\frac{H(z)}{\delta})$ satisfies: for any compact set $\mathcal{M} \subset \mathbb{R}^D$, $F(\mathcal{M}) \in \mathcal{B}_1 \cap \mathcal{B}_2$ a.s..

This shows that for any compact set $\mathcal{M} \in \mathbb{R}^D$, a small change into the set $F(\mathcal{M})$ where F is a transformation of \mathbb{R}^D of the type described in the Theorem almost surely results in a set in $\mathcal{B}_1 \cap \mathcal{B}_2$.

B.4.3 Proof of Theorem 8

Let \mathcal{M} be a compact set of \mathbb{R}^D . Since ε and H are independent, and thus f^ε and $H(\mathcal{M})$ are independent, writing μ_H the distribution of $H(\mathcal{M})$:

$$\mathbb{P}(F(\mathcal{M}) \in \mathcal{B}_1 \cap \mathcal{B}_2) = \int \mathbb{P}\left(f^\varepsilon\left(\frac{h}{\delta}\right) \in \mathcal{B}_1 \cap \mathcal{B}_2\right) d\mu_H(h) = 1,$$

provided that for any compact set $\mathcal{M}' \in \mathbb{R}^D$, $f^\varepsilon(\mathcal{M}') \in \mathcal{B}_1 \cap \mathcal{B}_2$ a.s..

Thus, it suffices to show that for any compact set $\mathcal{M} \in \mathbb{R}^D$, almost surely, $f^\varepsilon(\mathcal{M})$ is in the set \mathcal{B}_1 from Corollary 2. The proof for \mathcal{B}_2 is identical.

We will show that $\text{Card}(\arg \max_{z \in f^\varepsilon(\mathcal{M})} \pi^{(1)}(z)) = 1$, where $\pi^{(1)}(z)$ is the first coordinate of z . First, since \mathcal{M} is compact and f^ε is continuous, $f^\varepsilon(\mathcal{M})$ is compact, therefore the supremum of $\pi^{(1)}$ is reached at least at one point.

Lemma 2. *The two following properties hold almost surely.*

1. Let $F'_I, F'_J \in \mathcal{P}^\varepsilon$ be two different simplices, then at least one of the two following points holds:

- $\sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_I)} \pi^{(1)}(x) \neq \sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)} \pi^{(1)}(x)$
- $\pi^{(1)}$ does not reach its maximum on $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_I)$ or does not reach its maximum on $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)$.

2. Let $F'_I \in \mathcal{P}^\varepsilon$, then the supremum of $\pi^{(1)}$ on $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_I)$ is reached at at most one point of $\text{relint}(F'_I)$.

A consequence of this lemma is that almost surely, the maximizer of $\pi^{(1)}$ on $f^\varepsilon(\mathcal{M})$ is unique, as all maximizers of $\pi^{(1)}$ on $f^\varepsilon(\mathcal{M})$ belong to the relative interior of one simplex of \mathcal{P}^ε , which shows that $f^\varepsilon(\mathcal{M})$ is almost surely in \mathcal{B}_1 .

Proof of Lemma 2. The following functions will be of use in the proof. For any finite $J \subset \mathbb{N}$ such that $F'_J = \{x_i + \varepsilon_i\}_{i \in J} \in \mathcal{P}^\varepsilon$, for any $j \in J$ and $\alpha \in (0, 1]$, let

$$u_{\alpha, J} : e \in \mathbb{R} \mapsto \sup \left\{ \alpha \pi^{(1)}(x_j) + e + \sum_{k \in J \setminus \{j\}} \alpha_k \pi^{(1)}(x_k + \varepsilon_k), \text{ where} \right. \\ \left. z = \alpha x_j + \sum_{k \in J \setminus \{j\}} \alpha_k x_k \in \mathcal{M}, \alpha_k \in (0, 1] \text{ and } \alpha + \sum_k \alpha_k = 1 \right\}.$$

In other words, $u_{\alpha, J}$ is the supremum of $\pi^{(1)}$ on the slice of $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)$ that gives weight α to the vertex $(x_j + \varepsilon_j)$. To simplify the notations, let $w_k : z \mapsto \alpha_k$ be the “weight” functions. It is straightforward to check that

1. the function $u_{\alpha, J}$ is linear with slope α ,
2. $\sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)} \pi^{(1)}(x) = \sup_{\alpha \in (0, 1]} u_{\alpha, J}(\pi^{(1)}(\varepsilon_j))$,
3. the function $h : \pi^{(1)}(\varepsilon_j) \mapsto \sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)} \pi^{(1)}(x)$ (all coordinates of all ε_k other than $\pi^{(1)}(\varepsilon_j)$ being fixed) is convex,
4. if the supremum of $\pi^{(1)}$ on the closure of $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)$ is reached at some point $z \in F'_J$ when $\pi^{(1)}(\varepsilon_j) = e$, then $w_j(z)$ is a sub-gradient of h at e ,
5. since the number of points where the sub-gradient of a convex function on \mathbb{R} is not unique is at most countable, almost surely (whether all coordinates of all ε_k other than $\pi^{(1)}(\varepsilon_j)$ are fixed or not), h has a unique sub-gradient at $\pi^{(1)}(\varepsilon_j)$.

Let us now prove the first point of the lemma. Let $F'_I = \{x_i + \varepsilon_i\}_{i \in I}$ and $F'_J = \{x_i + \varepsilon_i\}_{i \in J}$ be two different simplices of \mathcal{P}^ε , and let $j \in J \setminus I$ (by exchanging the two simplices, we may assume without loss of generality that J is not a subset of I).

Consider the following, conditionally to $(\varepsilon_n)_{n \neq j}$ and $\pi^{(2:D)}(\varepsilon_j)$. Assume that $h(\pi^{(1)}(\varepsilon_j)) = \sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_I)} \pi^{(1)}(x)$ (otherwise we are in the first case of the first point of the lemma). We may assume without loss of generality (by point 5 above) that the sub-gradient of h at $\pi^{(1)}(\varepsilon_j)$ is unique. Two cases are possible:

- the sub-gradient of h at $\pi^{(1)}(\varepsilon_j)$ is 0. Then $\pi^{(1)}$ does not reach its maximum on $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)$, since if z is a maximizer of $\pi^{(1)}$, then $w_j(z) = 0$ by point 4,
- the sub-gradient of h at $\pi^{(1)}(\varepsilon_j)$ is positive, so there exists a single point e such that $h(e) = \sup_{x \in f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_I)} \pi^{(1)}(x)$. Since $\pi^{(1)}(\varepsilon_j)$ is uniform on $[-r, r]$ by construction, we almost surely have $\pi^{(1)}(\varepsilon_j) \neq e$, and thus this second case almost surely never happens.

For the second point of the lemma, by points 4 and 5, if the set of maximizers of $\pi^{(1)}$ on $f^\varepsilon(\mathcal{M}) \cap \text{relint}(F'_J)$ is a non-empty set \mathcal{Z} , then for any $j \in J$, almost surely, w_j is constant on \mathcal{Z} . Since every point $z \in F'_J$ is characterized by the vector $(w_j(z))_{j \in J}$, this shows that \mathcal{Z} contains a single point, which concludes the proof. \square

B.5 Density of \mathcal{A} and its complement

The proposition below shows that any support \mathcal{M} can be altered by a small perturbation to produce both supports that satisfy (H1) and (H2) and supports that satisfy neither. *A fortiori*, the same is true for (Adep), as on one hand (H1), (H2) and $A(\kappa)$ ensure (Adep) by Theorem 2 and on the other hand a small perturbation of the signal (by adding noise with independent components supported on a small neighborhood of zero) is enough to no longer satisfy (Adep).

Proposition 3. *The set $\mathcal{A} = (\cap_{\Delta > 0} \cup_{\varepsilon > 0} \mathcal{A}_1(\Delta, \varepsilon)) \cap (\cap_{\Delta > 0} \cup_{\varepsilon > 0} \mathcal{A}_2(\Delta, \varepsilon))$ of supports satisfying Corollary 1 and its complement are dense in the set of closed subsets of \mathbb{R}^D endowed with the Hausdorff distance.*

Proof. First, let us show that \mathcal{A} is dense.

Let $\delta > 0$ and let \mathcal{M} be a closed subset of \mathbb{R}^D , we show that there exists a closed \mathcal{M}' in $\cap_{\Delta > 0} (\mathcal{A}_1(\Delta, \delta) \cap \mathcal{A}_2(\Delta, \delta))$ (and thus in \mathcal{A}) such that $d_H(\mathcal{M}, \mathcal{M}') \leq 8\delta$.

For $k, l \in \{1, \dots, D\}$ with $k \leq l$, write $\pi^{(k:l)}$ the projection $\pi^{(k:l)} : (x_1, \dots, x_D) \in \mathbb{R}^D \mapsto (x_k, \dots, x_l) \in \mathbb{R}^{l-k+1}$ and $\pi^{(k)} = \pi^{(k:k)}$.

Let $z = (z_1, z_2) \in \mathcal{M}$ with $z_1 = \pi^{(1:d_1)}(z)$ and $z_2 = \pi^{(d_1+1:D)}(z)$, \mathcal{M}' is defined by cutting the space in half through z orthogonally to the space of the first d_1 coordinates and spreading the two halves apart, connecting them by a single segment to ensure it is in $\mathcal{A}_2(\Delta, \delta)$, then cut and connect again orthogonally to the $(d_1 + 1)$ -th axis to be in $\mathcal{A}_1(\Delta, \delta)$.

Formally, define \mathcal{M}' as the union of:

- $\{y \mid y = (y_1, y_2) \in \mathcal{M}, \pi^{(1)}(y_1) \leq \pi^{(1)}(z_1) \text{ and } \pi^{(1)}(y_2) \leq \pi^{(1)}(z_2)\}$,
- $\{(y_1 + 4\delta(1, 0, \dots, 0), y_2) \mid y = (y_1, y_2) \in \mathcal{M}, \pi^{(1)}(y_1) \geq \pi^{(1)}(z_1) \text{ and } \pi^{(1)}(y_2) \leq \pi^{(1)}(z_2)\}$,
- $\{(y_1, y_2 + 4\delta(1, 0, \dots, 0)) \mid y = (y_1, y_2) \in \mathcal{M}, \pi^{(1)}(y_1) \leq \pi^{(1)}(z_1) \text{ and } \pi^{(1)}(y_2) \geq \pi^{(1)}(z_2)\}$,
- $\{(y_1 + 4\delta(1, 0, \dots, 0), y_2 + 4\delta(1, 0, \dots, 0)) \mid y = (y_1, y_2) \in \mathcal{M}, \pi^{(1)}(y_1) \geq \pi^{(1)}(z_1) \text{ and } \pi^{(1)}(y_2) \geq \pi^{(1)}(z_2)\}$,
- the segment between z and $(z_1 + 4\delta(1, 0, \dots, 0), z_2)$ and the segment between z and $(z_1, z_2 + 4\delta(1, 0, \dots, 0))$.

An illustration of this construction is given in Figure 3.

By construction, the Hausdorff distance between this set \mathcal{M}' and \mathcal{M} is smaller than 8δ (the points in the first four sets have moved at most 8δ and the segments are at distance at most 8δ of z). \mathcal{M}' is also closed, and taking $x = (z_1 + 2\delta(1, 0, \dots, 0), z_2)$ and $x_2 = (z_1, z_2 + 2\delta(1, 0, \dots, 0))$




Figure 3: Transforming \mathcal{M} into a set $\mathcal{M}' \in \mathcal{B}_1 \cap \mathcal{B}_2$

in the definition of $\mathcal{A}_1(\Delta, \delta)$ and $\mathcal{A}_2(\Delta, \delta)$ is enough to check that $\mathcal{M}' \in \mathcal{A}_1(\Delta, \delta) \cap \mathcal{A}_2(\Delta, \delta)$ for any $\Delta > 0$.

To show that the complement of \mathcal{A} is dense, let \mathcal{M} be a closed subset of \mathbb{R}^D and $\eta > 0$, and let $\mathcal{M}' = \{x + y \mid x \in \mathcal{M}, y \in [-\eta, \eta]^D\}$. Then $d_H(\mathcal{M}, \mathcal{M}') \leq \eta\sqrt{D}$ by construction, and for any $\Delta \leq 2\eta$ and $\varepsilon > 0$, $\mathcal{M}' \notin \mathcal{A}_1(\Delta, \varepsilon)$, and thus $\mathcal{M}' \in \mathcal{A}^c$ where \mathcal{A}^c is the complement of the set \mathcal{A} .

Note that if \mathcal{M} is the support of a random variable X , then \mathcal{M}' is the support of $X + Y$, where Y is a uniform random variable on $[-\eta, \eta]^D$ that is independent of X . Since $\Phi_{X+Y} = \Phi_X \Phi_Y$ and Φ_Y does not satisfy (Adep), Φ_{X+Y} does not satisfy it either. \square

C Proof of Proposition 2

Let $\kappa_0 \in (1/2, 1]$.

From Lemma H.3 of [18], there exists a constant $c_T > 0$ such that for all $\kappa' \in [\kappa_0, 1]$, $m \geq D/\kappa'$ and $\phi \in \Upsilon_{\kappa', S}$,

$$\|\phi - T_m \phi\|_{\infty, \nu_{\text{est}}} \leq c_T (S \nu_{\text{est}})^m m^{-\kappa' m + D}.$$

Let $\kappa' \in [\kappa_0, 1]$ and assume that $m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}$, then we get $\|\phi - T_m \phi\|_{\infty, \nu_{\text{est}}} = O(n^{-2+o_n(1)})$, where $o_n(1)$ denotes a sequence tending to 0 when n tends to infinity. In particular, there exists n_0 such that for all $n \geq n_0$,

$$\sup_{\kappa' \in [\kappa_0, 1]} \sup_{\nu \in (0, \nu_{\text{est}}]} \sup_{m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}} \sup_{\phi \in \Upsilon_{\kappa', S}} \|\phi - T_m \phi\|_{2, \nu} \leq \frac{1}{n} \quad (9)$$

and

$$\sup_{\kappa' \in [\kappa_0, 1]} \sup_{m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}} \sup_{\phi \in \Upsilon_{\kappa', S}} |M_n(\phi) - M_n(T_m \phi)| \leq c \|\phi - T_m \phi\|_{\infty, \nu_{\text{est}}} \leq \frac{1}{n} \quad (10)$$

for some $c > 0$ that depends only on ν_{est} , κ_0 and S , using that $\sup_{\phi \in \Upsilon_{\kappa_0, S}} \|\phi\|_{\infty, \nu_{\text{est}}} < +\infty$.

Let now $\kappa \in [\kappa_0, 1]$ and $\Phi_X \in \Upsilon_{\kappa, S} \cap \mathcal{H}$. By definition, for any $m \geq 1$ and $\kappa' \in [\kappa_0, \kappa]$, $\widehat{\Phi}_{n, m, \kappa'}$ is such that $\widehat{\Phi}_{n, m, \kappa'} \in \Upsilon_{\kappa', S} \cap \mathcal{H}$ and

$$\begin{aligned} M_n(T_m \widehat{\Phi}_{n, m, \kappa'}) &\leq \inf_{\phi \in \Upsilon_{\kappa', S} \cap \mathcal{H}} M_n(T_m \phi) + \frac{1}{n} \\ &\leq M_n(T_m \Phi_X) + \frac{1}{n} \end{aligned}$$

and thus, by (10),

$$\sup_{\kappa' \in [\kappa_0, \kappa]} \sup_{m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}} M_n(\widehat{\Phi}_{n, m, \kappa'}) \leq M_n(\Phi_X) + \frac{2}{n}. \quad (11)$$

Straightforward computations prove that for some constant C , for all $\phi \in \Upsilon_{\kappa_0, S}$,

$$|M_n(\phi) - M(\phi; \nu_{\text{est}} | \Phi_X)| \leq C \frac{\|Z_n\|_{\infty, \nu_{\text{est}}}}{\sqrt{n}}$$

where $Z_n(t) = \sqrt{n}(\tilde{\phi}_n(t) - \Phi_X(t)\Phi_{\varepsilon(1)}(t_1)\Phi_{\varepsilon(2)}(t_2))$. Therefore, for any $\nu \in (0, \nu_{\text{est}}]$, writing $\zeta_{m, \kappa'} = \widehat{\Phi}_{n, m, \kappa'} - \Phi_X$, for some constant C ,

$$\begin{aligned} M(\widehat{\Phi}_{n, m, \kappa'}; \nu | \Phi_X) &\leq M(\widehat{\Phi}_{n, m, \kappa'}; \nu_{\text{est}} | \Phi_X) \\ &\leq 2C \frac{\|Z_n\|_{\infty, \nu_{\text{est}}}}{\sqrt{n}} + \frac{2}{n}. \end{aligned}$$

Combining this with the concentration properties of Z_n described in the proof of Lemma A.1 of [17] shows that for any $\delta > 0$, there exist $c_\delta > 0$ and n_0 (which do not depend on κ) such that for all $n \geq n_0$, with probability at least $1 - e^{-c_\delta n}$,

$$\sup_{\kappa' \in [\kappa_0, \kappa]} \sup_{m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}} M(\widehat{\Phi}_{n, m, \kappa'}; \nu_{\text{est}} | \Phi_X) \leq \delta.$$

In addition, since $\Upsilon_{\kappa_0, S} \cap \mathcal{H}$ is compact in $L^2([-\nu_{\text{est}}, \nu_{\text{est}}]^D)$, $\phi \mapsto M(\phi; \nu_{\text{est}} | \Phi_X)$ is continuous on $L^2([-\nu_{\text{est}}, \nu_{\text{est}}]^D)$, and $M(\phi; \nu_{\text{est}} | \Phi_X) = 0$ implies $\phi = \Phi_X$ for all $\phi \in \mathcal{H} \cap \Upsilon_{\kappa_0, S}$ by Theorem 1, and for all $\eta > 0$, there exists $\delta > 0$ such that

$$\inf_{\phi \in \Upsilon_{\kappa_0, S} \cap \mathcal{H} \text{ s.t. } \|\phi - \Phi_X\|_{2, \nu_{\text{est}}} \geq \eta} M(\phi; \nu_{\text{est}} | \Phi_X) > \delta.$$

Therefore, for all $\eta > 0$, there exist $c_\eta > 0$ and n_0 (which do not depend on κ) such that for all $n \geq n_0$, with probability at least $1 - e^{-c_\eta n}$,

$$\sup_{\kappa' \in [\kappa_0, \kappa]} \sup_{m \geq \frac{2}{\kappa'} \frac{\log n}{\log \log n}} \|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}} \leq \eta. \quad (12)$$

Now, using (20) in [8] (which is a consequence of direct computations and norm comparisons in [18] Section J) we get that there exists $\eta_1 > 0$ such that as soon as $\|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}} \leq \eta_1$,

$$M(\widehat{\Phi}_{n, m, \kappa'}; \nu | \Phi_X) \leq c_M \frac{\|Z_n\|_{\infty, \nu_{\text{est}}}}{\sqrt{n}} \|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}}^{1 - \epsilon(\|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}})} + \frac{2}{n}, \quad (13)$$

where for all $u > 0$, $\epsilon(u) = b/\log \log(1/u)$ for some $b > 0$. Fix now $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$ and $E > 0$ such that $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$. In particular, $c_\nu \geq c(\nu) > 0$. Assumption (Amin) and Proposition 1 in [19] imply that there exists $\eta_2 > 0$ and c depending only on ν , c_ν and $c(\nu, \alpha)$ such that as soon as $\|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}} \leq \eta_2$,

$$M(\widehat{\Phi}_{n, m, \kappa'}; \nu | \Phi_X) \geq c \|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}}^{2(1+\alpha)}. \quad (14)$$

Let $\eta = \min\{\eta_1, \eta_2\}$. Then, by (14) and (13), as soon as $\|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}} \leq \eta$,

$$\|\zeta_{m, \kappa'}\|_{2, \nu}^{2+2\alpha} \leq \frac{2}{c} \max \left(c_M \frac{\|Z_n\|_{\infty, \nu_{\text{est}}}}{\sqrt{n}} \|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}}^{1 - \epsilon(\|\zeta_{m, \kappa'}\|_{2, \nu_{\text{est}}})}, \frac{2}{n} \right). \quad (15)$$

Now, following the proof of (25) of Section A.3 of [17], for any $\nu' \geq \nu$, $m \geq 1$ and $\phi \in \mathbb{C}_m[X]$,

$$\|\phi\|_{2, \nu'} \leq m^{D/2} \left(4 \frac{\nu'}{\nu}\right)^{m+D/2} \|\phi\|_{2, \nu}. \quad (16)$$

Assuming $m \in [\frac{2}{\kappa'} \frac{\log n}{\log \log n}, C \frac{\log n}{\log \log n}]$ in the following series of inequalities (for some fixed $C >$

$2/\kappa')$,

$$\begin{aligned}
\|\zeta_{m,\kappa'}\|_{2,\nu_{\text{est}}} &\leq 2 \max(\|T_m \zeta_{m,\kappa'}\|_{2,\nu_{\text{est}}}, \frac{3}{n}) && \text{by (9)} \\
&\leq 2 \max\left(\|T_m \zeta_{m,\kappa'}\|_{2,\nu} m^{\frac{D}{2}} \left(4 \frac{\nu_{\text{est}}}{\nu}\right)^{m+\frac{D}{2}}, \frac{3}{n}\right) && \text{by (16)} \\
&\leq 4 \max\left(\|\zeta_{m,\kappa'}\|_{2,\nu} m^{\frac{D}{2}} \left(4 \frac{\nu_{\text{est}}}{\nu}\right)^{m+\frac{D}{2}}, \frac{3m^{\frac{D}{2}} \left(4 \frac{\nu_{\text{est}}}{\nu}\right)^{m+\frac{D}{2}}}{n}\right) && \text{by (9)} \\
&\leq n^{\epsilon(1/n)} \max\left(\|\zeta_{m,\kappa'}\|_{2,\nu}, \frac{1}{n}\right), && (17)
\end{aligned}$$

up to increasing the constant b in the definition of $u \mapsto \epsilon(u)$, which can be done without loss of generality. Together with (15) and (12), one gets for all $s \in [1, c_\eta n]$ (assuming $c_\eta \leq 1$ without loss of generality), with probability at least $1 - e^{-c_\eta n} - e^{-s} \geq 1 - 2e^{-s}$ (on the event where $\|Z_n\|_{\infty, \nu_{\text{est}}} \leq c_Z \sqrt{s}$ and (12) holds) that for all $\kappa' \in [\kappa_0, \kappa]$ and $m \in [\frac{2}{\kappa'} \frac{\log n}{\log \log n}, C \frac{\log n}{\log \log n}]$,

$$\|\zeta_{m,\kappa'}\|_{2,\nu}^{2+2\alpha} \leq c \max\left(\sqrt{\frac{s}{n}} n^{\epsilon(1/n)} \left(\|\zeta_{m,\kappa'}\|_{2,\nu} \vee \frac{1}{n}\right)^{1-\epsilon(\|\zeta_{m,\kappa'}\|_{2,\nu_{\text{est}}})}, \frac{1}{n}\right)$$

for some constant $c > 0$ that does not depend on κ, κ' or m , from which Proposition 2 follows.

D Proofs for Section 4

D.1 Proof of Theorem 3

The following properties of ψ_A and $\mathcal{F}[\psi_A]$ will be of use.

- (I) The support of $\mathcal{F}[\psi_A]$ is the unit ball $\{y \in \mathbb{R}^D : \|y\|_2 \leq 1\}$.
- (II) $\psi_A > 0$ and $\mathcal{F}[\psi_A] \geq 0$.
- (III) ψ_A is continuous so that there exist constants $c_A > 0$ and $d_A > 0$ such that for all $x \in \{y \in \mathbb{R}^D : \|y\|_2 \leq c_A\}$, $\psi_A(x) \geq d_A$.
- (IV) ψ_A and $\psi_{A,h}$ are probability densities on \mathbb{R}^D .
- (V) (Lemma in [26]) For all $A > 0$, there exists $\beta_A > 0$ such that

$$\lim_{\|t\|_2 \rightarrow \infty} \exp\{\beta_A \|t\|_2^{\frac{A}{A+1}}\} \psi_A(t) = 0 \tag{18}$$

- (VI) It holds

$$\|\psi_{A,h}\|_2 = \frac{I(A)}{h^{D/2}} \|u_A * u_A\|_2. \tag{19}$$

We shall also need the following lemmas.

Lemma 3. *Assume $G \in \text{St}_{\mathcal{K}}(a, d, r_0)$, then for any $h \leq r_0/c_A$,*

$$\inf_{y \in \mathcal{M}_G \cap \mathcal{K}} \bar{g}(y) \geq ac_A^d d_A \left(\frac{1}{h}\right)^{D-d}.$$

The proof of Lemma 3 is detailed in Section D.2.

Lemma 4. *For any $C_1 > 0$, there exists $h_0 > 0$ depending only on C_1, D and A such that for any $h \leq h_0$,*

$$\sup \left\{ \bar{g}(y) \mid y \in \mathcal{K}, d(y, \mathcal{M}_G) > h \left[\frac{D}{\beta_A} \log \left(\frac{1}{h} \right) \right]^{\frac{A+1}{A}} \right\} \leq C_1.$$

The proof of Lemma 4 is detailed in Section D.3.

The last ingredient is to control the difference between the convoluted density and its estimator, defined as $\Gamma_{n,\kappa} = \|\widehat{g}_{n,\kappa} - \bar{g}\|_\infty = \sup_{y \in \mathbb{R}^D} |\widehat{g}_{n,\kappa}(y) - \bar{g}(y)|$. The following Lemma uses Proposition 2 to show that it converges uniformly to zero with high probability.

Lemma 5. *Let $\kappa_0 \in (1/2, 1]$ and $S > 0$. For all $\kappa \in [\kappa_0, 1]$, take $m_\kappa \leq \frac{1}{6\kappa} \frac{\log n}{\log \log n}$ such that $\inf_{\kappa \in [\kappa_0, 1]} m_\kappa \rightarrow +\infty$ as $n \rightarrow +\infty$. Let $h = c_h S m_\kappa^{-\kappa}$ for some $c_h \geq \exp\{2D + 2\}$.*

Then for any $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$, $E > 0$, for any $\eta > 0$, there exists n_0 such that for all $n \geq n_0$,

$$\sup_{\kappa \in [\kappa_0, 1]} \sup_{G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})} \sup_{\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)} \mathbb{P}_{(G * \mathbb{Q})^{\otimes n}}(\Gamma_{n,\kappa} \geq \eta) \leq \frac{2}{n}.$$

The proof of Lemma 5 is detailed in Appendix D.4.

To prove Theorem 3, the idea is to lower bound \bar{g} on the support \mathcal{M}_G when the bandwidth parameter h becomes small, and to upper bound it on every points further than a small distance (depending on h) from that support.

Let now $\kappa_0 \in (1/2, 1]$, $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$, $E > 0$, $S > 0$ and $C > 0$. Let $\kappa \in [\kappa_0, 1]$, $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$ and $G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})$.

By Lemma 3, for any $h \leq (r_0/c_A) \wedge 1$,

$$\inf_{y \in \mathcal{M}_G \cap \mathcal{K}} \widehat{g}_{n,\kappa}(y) \geq \inf_{y \in \mathcal{M}_G \cap \mathcal{K}} \bar{g}(y) - \Gamma_{n,\kappa} \geq ac_A^d d_A \left(\frac{1}{h}\right)^{D-d} - \Gamma_{n,\kappa} \geq \frac{ac_A^d d_A}{2} \left(\frac{1}{h}\right)^{D-d}$$

as soon as $\Gamma_{n,\kappa} \leq \frac{ac_A^d d_A}{2}$, and this lower bound is strictly larger than $\lambda_{n,\kappa}$ for any d . This implies that on the event where $\Gamma_{n,\kappa} \leq \frac{ac_A^d d_A}{2}$, $\mathcal{M}_G \cap \mathcal{K} \subset \widehat{\mathcal{M}}_\kappa \cap \mathcal{K}$. Next,

$$\sup_{y \in \mathcal{K}: d(y, \mathcal{M}_G) \geq h \left[\frac{D}{\beta_A} \log\left(\frac{1}{h}\right)\right]^{\frac{A+1}{A}}} \widehat{g}_{n,\kappa}(y) \leq \sup_{y \in \mathcal{K}: d(y, \mathcal{M}_G) \geq h \left[\frac{D}{\beta_A} \log\left(\frac{1}{h}\right)\right]^{\frac{A+1}{A}}} \bar{g}(y) + \Gamma_{n,\kappa}.$$

Choosing $C_1 = \frac{ac_A^d d_A}{16}$ and applying Lemma 4 we get that, on the event where $\Gamma_{n,\kappa} \leq \frac{ac_A^d d_A}{16}$,

$$\sup_{y \in \mathcal{K}, d(y, \mathcal{M}_G) \geq h \left[\frac{D}{\beta_A} \log\left(\frac{1}{h}\right)\right]^{\frac{A+1}{A}}} \widehat{g}(y) \leq 2C_1,$$

and this upper bound is strictly less than $\lambda_{n,\kappa}$ for any d . This implies that

$$\left\{ y : y \in \mathcal{K}, d(y, \mathcal{M}_G) > h \left[\frac{D}{\beta_A} \log\left(\frac{1}{h}\right)\right]^{\frac{A+1}{A}} \right\} \cap \widehat{\mathcal{M}}_\kappa = \emptyset.$$

As a result, we have proved that for all $\kappa_0 \in (1/2, 1]$, $S > 0$, $a > 0$, $d \leq D$, $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$ and $E > 0$, there exist $c' > 0$ and n_0 such that for all $n \geq n_0$, for all $\kappa \in [\kappa_0, 1]$, $G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})$ and $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$, with $(G * \mathbb{Q})^{\otimes n}$ -probability at least $1 - \frac{2}{n}$,

$$\frac{\log(n)^\kappa}{\log(\log(n))^{\kappa + \frac{A+1}{A}}} H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_\kappa) \leq c'. \quad (20)$$

Using the fact that $H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_\kappa)$ is uniformly upper bounded, the theorem follows.

D.2 Proof of Lemma 3

Let $y \in \mathcal{M}_G \cap \mathcal{K}$. By property (III) of ψ_A ,

$$\begin{aligned}\bar{g}(y) &= \frac{1}{h^D} \int \psi_A \left(\frac{\|y-u\|}{h} \right) dG(u) \\ &\geq \frac{1}{h^D} \int_{\|u-y\|_2 \leq c_A h} \psi_A \left(\frac{\|y-u\|}{h} \right) dG(u) \\ &\geq \frac{1}{h^D} d_A G(B(y, c_A h)) \\ &\geq \frac{1}{h^D} d_A a(c_A h)^d.\end{aligned}$$

D.3 Proof of Lemma 4

Recall the definition of \bar{g} : for all $y \in \mathbb{R}^D$,

$$\bar{g}(y) = \frac{1}{h^D} \int \psi_A \left(\frac{\|y-u\|}{h} \right) dG(u).$$

Let $C_1 > 0$ and $\epsilon > 0$. By Property (V) of ψ_A , there exists $T > 0$ (depending on A and C_1) such that for any $t \geq T$, $\psi_A(t) \leq C_1 \exp(-\beta_A t^{A/(A+1)})$. Take $y \in \mathbb{R}^D$ such that $d(y, \mathcal{M}_G) > (\frac{\epsilon}{\beta_A})^{\frac{A+1}{A}} h \log(\frac{1}{h})^{\frac{A+1}{A}}$, then for all $u \in \mathcal{M}_G$, $\frac{\|y-u\|}{h} \geq (\beta_A^{-1} \log(\frac{1}{h\epsilon}))^{\frac{A+1}{A}}$, therefore there exists $h_0 > 0$ depending only on ϵ , D , A and T (thus C_1 such that $h \leq h_0$ implies $\frac{\|y-u\|}{h} \geq T$ and thus

$$\begin{aligned}\psi_A \left(\frac{\|y-u\|}{h} \right) &\leq C_1 \exp \left\{ -\beta_A \left(\frac{\|y-u\|}{h} \right)^{A/(A+1)} \right\} \\ &\leq C_1 \exp \left\{ -\log \left(\frac{1}{h^\epsilon} \right) \right\} \\ &= C_1 h^\epsilon,\end{aligned}$$

and finally $\bar{g}(y) \leq C_1 (\frac{1}{h})^{D-\epsilon}$ since G is a probability distribution. Lemma 4 follows by taking $\epsilon = D$.

D.4 Proof of Lemma 5

Recall $\Gamma_{n,\kappa} = \|\hat{g}_{n,\kappa} - \bar{g}\|_\infty$. We first relate it to $\|T_{m_\kappa} \hat{\Phi}_{n,\kappa} - \Phi_X\|_{2,1/h}$.

Lemma 6. *Let $\kappa \in (0, 1]$, $h > 0$, $m > 0$. For any $A > 0$,*

$$\Gamma_{n,\kappa} \leq I(A) \frac{\|u_A * u_A\|_2}{h^{D/2}} \|T_{m_\kappa} \hat{\Phi}_{n,\kappa} - \Phi_X\|_{2,1/h}.$$

Proof. For $y \in \mathbb{R}^D$,

$$\hat{g}_{n,\kappa}(y) - \bar{g}(y) = \left(\frac{1}{2\pi} \right)^D \int e^{-it^\top y} \mathcal{F}[\psi_A](th) (T_{m_\kappa} \hat{\Phi}_{n,\kappa}(t) - \Phi_X(t)) dt.$$

Since $\mathcal{F}[\psi_A](th)$ is 0 for $\|t\|_2 > 1/h$,

$$\begin{aligned}\hat{g}_{n,\kappa}(y) - \bar{g}(y) &= \left(\frac{1}{2\pi} \right)^D \int e^{-it^\top y} \mathcal{F}[\psi_A](th) (T_{m_\kappa} \hat{\Phi}_{n,\kappa}(t) - \Phi_X(t)) 1_{\|t\|_2 \leq 1/h} dt \\ &= \mathcal{F}^{-1}[\mathcal{F}[\psi_h] \{ (T_{m_\kappa} \hat{\Phi}_{n,\kappa} - \Phi_X) 1_{\|t\|_2 \leq 1/h} \}](y) \\ &= \mathcal{F}^{-1}[\mathcal{F}[\psi_{A,h}] * \mathcal{F}^{-1}[(T_{m_\kappa} \hat{\Phi}_{n,\kappa} - \Phi_X) 1_{\|t\|_2 \leq 1/h}]](y).\end{aligned}\tag{21}$$

By Young's convolution inequality,

$$\|\hat{g}_{n,\kappa} - \bar{g}\|_\infty \leq \|\mathcal{F}^{-1}[\mathcal{F}[\psi_{A,h}]]\|_2 \|\mathcal{F}^{-1}[(T_{m_\kappa} \hat{\Phi}_{n,\kappa} - \Phi_X) 1_{\|t\|_2 \leq 1/h}]\|_2.$$

Finally, using Parseval's equality and the fact that $\mathcal{F}^{-1}[\mathcal{F}[\psi_{A,h}]] = \psi_{A,h}$,

$$\|\widehat{g}_{n,\kappa} - \bar{g}\|_\infty \leq \|\psi_{A,h}\|_2 \|T_{m_\kappa} \widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,1/h},$$

and use that $\|\psi_{A,h}\|_2 = \frac{I(A)}{h^{D/2}} \|u_A * u_A\|_2$ to conclude the proof. \square

Let us now show that with proper choices for the parameters m_κ and h , $\Gamma_{n,\kappa}$ tends to 0 with high probability.

Let $\kappa_0 \in (1/2, 1]$, $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$, $E > 0$, $S > 0$ and $C > 0$. By Proposition 2, taking $s = \log n$ and δ small enough (to be chosen later), there exists c and n_0 such that for all $n \geq n_0$,

$$\sup_{\kappa \in [\kappa_0, 1]} \sup_{\substack{G \in \text{St}_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H}) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)}} \mathbb{P}_{(G * \mathbb{Q})^{\otimes n}} \left(\|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,1/h} \leq c(\log n) n^{-(1-\delta)/(1+2\alpha)} \right) \geq 1 - \frac{2}{n}. \quad (22)$$

Take these c and n_0 , and let $\kappa \in [\kappa_0, 1]$, $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$ and $G \in \text{St}_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})$.

Using inequalities analogous to (28)-(29) p.17 of [17], we get that for all integer m ,

$$\|T_m \widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,1/h}^2 \leq 4U(h) + 4m^D \left(2 + 2\frac{1}{h\nu}\right)^{2m+D} \left(2V(\nu) + \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2\right), \quad (23)$$

where

$$U(h) = ch^{-D-2m-2/\kappa} S^{2m} m^{-2\kappa m+2D} \exp(2\kappa(S/h)^{1/\kappa})$$

and $V(\nu) = c(S\nu)^{2m+2/\kappa} m^{-2\kappa m+2D}$.

Thus, applying Lemma 6 and using $h = c_h S m_\kappa^{-\kappa}$, there exists $C > 0$ (which does not depend on κ) such that on the event where (23) holds:

$$\begin{aligned} \Gamma_{n,\kappa}^2 &\leq C(c_h S)^{-2D-2m_\kappa-2/\kappa} m_\kappa^{2D(\kappa+1)+2} S^{2m_\kappa} \exp(2\kappa c_h^{-1/\kappa} m_\kappa) \\ &\quad + C m_\kappa^{D(1+\kappa)} \left(2 + 2\frac{m_\kappa^\kappa}{c_h S\nu}\right)^{2m_\kappa+D} \left((S\nu)^{2m_\kappa+2/\kappa} m_\kappa^{-2\kappa m_\kappa+2D} + \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right). \end{aligned} \quad (24)$$

The first term of the upper bound is upper bounded as follows.

$$\begin{aligned} &(c_h S)^{-2D-2m_\kappa-2/\kappa} m_\kappa^{2D(\kappa+1)+2} S^{2m_\kappa} \exp(2\kappa c_h^{-1/\kappa} m_\kappa) \\ &= S^{-2D-2/\kappa} \exp\{(-2D-2m_\kappa-2/\kappa)\log(c_h) + (2D(\kappa+1)+2)\log(m_\kappa) + 2\kappa c_h^{-1/\kappa} m_\kappa\} \\ &\leq C \exp\{(-2\log(c_h) + 1)m_\kappa + 2(D(\kappa+1)+1)\log(m_\kappa)\} \quad (25) \\ &\leq C \exp\{(-2\log(c_h) + 3 + 2D(\kappa+1))m_\kappa\}, \quad (26) \end{aligned}$$

for another constant $C > 0$, where inequality (25) holds because $2\kappa c_h^{1/\kappa} > 1$ and inequality (26) holds because $\log(m_\kappa) \leq m_\kappa$. The second term of the upper bound is upper bounded by

$$\begin{aligned} &m_\kappa^{D(1+\kappa)} \left(2 + 2\frac{m_\kappa^\kappa}{c_h S\nu}\right)^{2m_\kappa+D} \left((S\nu)^{2m_\kappa+2/\kappa} m_\kappa^{-2\kappa m_\kappa+2D} + \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right) \\ &\leq C m_\kappa^{D(1+2\kappa)} (2\kappa m_\kappa)^{2\kappa m_\kappa} (2\kappa)^{-2\kappa m_\kappa} (c_h S\nu)^{-2m_\kappa-D} \\ &\quad \times \left((S\nu)^{2m_\kappa+2/\kappa} m_\kappa^{-2\kappa m_\kappa+2D} + \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right) \\ &\leq C \left(\exp\left\{(-2\log(c_h) + (3D+2\kappa))m_\kappa\right\} \right. \\ &\quad \left. + (2\kappa m_\kappa)^{2\kappa m_\kappa} \exp\left\{(-2\log(c_h) + D(1+2\kappa))m_\kappa\right\} \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right) \end{aligned}$$

for another constant $C > 0$. Putting all together, we get that for yet another constant $C > 0$,

$$\Gamma_{n,\kappa}^2 \leq C \max \left(\exp \left\{ (-2 \log(c_h) + 3 + 2D(\kappa + 1))m_\kappa \right\}, \exp \left\{ (-2 \log(c_h) + (3D + 2\kappa))m_\kappa \right\}, \right. \\ \left. (2\kappa m_\kappa)^{2\kappa m_\kappa} \exp \left\{ (-2 \log(c_h) + D(1 + 2\kappa))m_\kappa \right\} \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right).$$

Choosing $c_h \geq \exp\{2D + 2\}$ and $m_\kappa \leq \frac{1}{6\kappa} \frac{\log n}{\log \log n}$, it follows that

$$\Gamma_{n,\kappa}^2 \leq C e^{-m_\kappa} \left[1 \vee n^{1/6} \|\widehat{\Phi}_{n,\kappa} - \Phi_X\|_{2,\nu}^2 \right]. \quad (27)$$

By (22), taking $\delta < 1 - \frac{1+2\alpha}{6}$ (which is possible since $\alpha < 1$), we obtain that on the event of (22), $\Gamma_{n,\kappa}^2 \leq C e^{-m_\kappa} \rightarrow 0$ (up to changing C), which concludes the proof.

D.5 Proof of Theorem 4

Fix $\kappa_0 \in (1/2, 1]$, $S > 0$, $a > 0$, $d \leq D$, $\nu \in (0, \nu_{\text{est}}]$, $c(\nu) > 0$, $E > 0$. Using the end of the proof of Theorem 3, there exist n_0 and c' such that for all $\kappa \in [\kappa_0, 1]$, all $G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})$ and all $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$, with $(G * \mathbb{Q})^{\otimes n}$ -probability at least $1 - \frac{2}{n}$,

$$\frac{\log(n)^\kappa}{\log(\log(n))^{\kappa + \frac{A+1}{A}}} H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_\kappa) \leq c'. \quad (28)$$

Let us now choose $c_\sigma = c'$ and consider the event where (20) holds. By the triangular inequality, for any $\kappa \in [\kappa_0, 1]$,

$$H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_\kappa) + H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \\ \leq \sigma_n(\kappa) + H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}).$$

Now, using the definition of $B_n(\cdot)$, if $\kappa \leq \widehat{\kappa}_n$, then

$$H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq B_n(\widehat{\kappa}_n) + \sigma_n(\kappa)$$

while if $\kappa \geq \widehat{\kappa}_n$, then

$$H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq B_n(\kappa) + \sigma_n(\widehat{\kappa}_n)$$

so that in all cases,

$$H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq B_n(\widehat{\kappa}_n) + \sigma_n(\kappa) + B_n(\kappa) + \sigma_n(\widehat{\kappa}_n) \\ \leq 2B_n(\kappa) + 2\sigma_n(\kappa)$$

using the definition of $\widehat{\kappa}_n$, and therefore

$$H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq 2B_n(\kappa) + 3\sigma_n(\kappa).$$

By the triangular inequality and the definition of $B_n(\cdot)$,

$$B_n(\kappa) \leq 0 \vee \sup_{\kappa' \in [\kappa_0, \kappa]} \left\{ H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \mathcal{M}_G) + H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\kappa'}) - \sigma_n(\kappa') \right\} \\ \leq H_{\mathcal{K}}(\widehat{\mathcal{M}}_\kappa, \mathcal{M}_G) + 0 \vee \sup_{\kappa' \in [\kappa_0, \kappa]} \left\{ H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\kappa'}) - \sigma_n(\kappa') \right\} \\ \leq \sigma_n(\kappa).$$

Thus, for all $\kappa \in [\kappa_0, 1]$, all $G \in St_{\mathcal{K}}(a, d, r_0) \cap \mathcal{L}(\kappa, S, \mathcal{H})$ and all $\mathbb{Q} \in \mathcal{Q}^{(D)}(\nu, c(\nu), E)$, with $(G * \mathbb{Q})^{\otimes n}$ -probability at least $1 - \frac{2}{n}$,

$$H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\widehat{\kappa}_n}) \leq 5\sigma_n(\kappa),$$

and using the fact that $H_{\mathcal{K}}(\mathcal{M}_G, \widehat{\mathcal{M}}_{\widehat{\kappa}}) \leq \sup_{x, x' \in \mathcal{K}} d(x, x')$ on the event of probability at most $2/n$ where this does not hold, Theorem 4 follows.

D.6 Proof of Theorem 5

As usual for the two-points method, the idea is to find two distributions having support as far as possible in $H_{\mathcal{K}}$ -distance, and a noise such that the joint distributions of the observations have total variation distance upper bounded by some $C < 1$. We shall consider the noise as in [17], with independent identically distributed coordinates having density q defined as

$$q : x \in \mathbb{R} \mapsto c_q \frac{1 + \cos(cx)}{(\pi^2 - (cx)^2)^2}$$

for some $c > 0$, where c_q is such that q is a probability density, and with characteristic function

$$\mathcal{F}[q] : t \mapsto c_q \left[\left(1 - \left|\frac{t}{c}\right|\right) \cos\left(\pi \frac{t}{c}\right) + \frac{1}{\pi} \sin\left(\pi \left|\frac{t}{c}\right|\right) \right] \mathbf{1}_{-c \leq t \leq c}.$$

Let us now define the two distributions to apply the two-points method. For any $\kappa \in (1/2, 1]$, we first choose a density function f_{κ} according to the following Lemma.

Lemma 7. *For any $\kappa \in (1/2, 1)$, $p \geq 1$, there exists a continuous density function $f_{\kappa} : \mathbb{R} \rightarrow \mathbb{R}$ in $L_p(\mathbb{R})$, positive everywhere, and positive constants A, B such that for all $u \in \mathbb{R}$,*

$$|\mathcal{F}[f_{\kappa}](u)| \leq A \exp(-B|u|^{\frac{1}{\kappa}}) \quad \text{and} \quad |\mathcal{F}[f_{\kappa}]'(u)| \leq A \exp(-B|u|^{\frac{1}{\kappa}}).$$

For any $\delta \in (0, 1)$, there exists a continuous compactly supported density function $f_1 : [-1, 1] \rightarrow \mathbb{R}$ positive everywhere such that

$$|\mathcal{F}[f_1](u)| \leq A \exp(-B|u|^{\delta}) \quad \text{and} \quad |\mathcal{F}[f_1]'(u)| \leq A \exp(-B|u|^{\delta}).$$

The proof of Lemma 7 is detailed in Section D.7.

Then, inspired by [20], for all $\gamma \in (0, 1]$, define $\tilde{g}_{\gamma} : \mathbb{R} \rightarrow \mathbb{R}$ and $g_{\gamma} : \mathbb{R} \rightarrow \mathbb{R}^{D-d}$ for all $x \in \mathbb{R}$ as

$$\tilde{g}_{\gamma}(x) = \cos\left(\frac{x}{\gamma}\right) \quad \text{and} \quad g_{\gamma}(x) = (\tilde{g}_{\gamma}(x), 0, \dots, 0)^{\top}.$$

Let $M_0(\gamma) = \{(u, \gamma g_{\gamma}(u)) : u \in \mathbb{R}\}$, $M_1(\gamma) = \{(u, -\gamma g_{\gamma}(u)) : u \in \mathbb{R}\}$, and let $J \in \mathbb{R}^{D \times D}$ be the matrix

$$J = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1/2 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}.$$

For any $\kappa \in (1/2, 1]$, let $U(\kappa)$ be the random variable in \mathbb{R} having density f_{κ} defined in Lemma 7 and let $S_0(\kappa) = (U(\kappa), \gamma g_{\gamma}(U(\kappa)))$, $S_1(\kappa) = (U(\kappa), -\gamma g_{\gamma}(U(\kappa)))$. For $i \in \{0, 1\}$, we shall denote $T_i(\kappa)$ the distribution of $S_i(\kappa)$. Finally we define $X_i(\kappa) = JS_i(\kappa)$, $i = 0, 1$ and $G_i(\kappa)$ the distribution of $X_i(\kappa)$. When $\kappa < 1$, the supports of $S_i(\kappa)$ and $X_i(\kappa)$ do not depend on κ . They are illustrated in Figure 4. The transformation J is such that the support of X_i is the graph of a bijective function, which makes it easy to satisfy (Adep) through for instance assumptions (H1), (H2) and Theorem 2 from Section 2.3.

Lemma 8. *For any $i \in \{0, 1\}$ and $\kappa \in (1/2, 1]$, $X_i(\kappa)$ satisfies $A(\kappa)$.*

The proof of Lemma 8 is detailed in Section D.8.

Lemma 9. *There exists $a_0 > 0$ such that for $i \in \{0, 1\}$, for any $d \geq 1$, $r_0 < 1$ and $a \leq a_0$, $G_i(\kappa) \in St_{\mathcal{K}}(a, d, r_0)$.*

The proof of Lemma 9 is detailed in Section D.9.

Lemma 10. *For any $i \in \{0, 1\}$ and $\kappa \in (1/2, 1]$, $\Phi_{X_i(\kappa)}$ satisfy (Adep)*




Figure 4: Left: support of S_0 and S_1 when $\gamma = 1$. Right: support of X_0 and X_1 when $\gamma = 1$.

The proof of Lemma 10 is detailed in Section D.10.

We finally set $\mathcal{H}_\kappa^* = \{\Phi_{X_0(\kappa)}, \Phi_{X_1(\kappa)}\}$. As a finite set, it is closed in $L_2([-\nu, \nu]^D)$.

Let us comment on dimensionality. The distributions used here are distributions with support of dimension 1. This is not an issue since the d in the definition of $St_{\mathcal{K}}(a, d, r_0)$ is an upper bound on the dimension of the support. We could also have used supports with dimension d by adding to $X_i(\kappa)$ an independent uniform distribution on a Euclidean ball of a linear space of dimension d .

The support of $G_i(\kappa)$ is $JM_i(\gamma)$, and the following lemma follows easily from the fact that for $i \in \{0, 1\}$, $JM_i(\gamma) = \gamma JM_i(1)$.

Lemma 11. *For any $\gamma > 0$,*

$$H_{\mathcal{K}}(JM_0(\gamma), JM_1(\gamma)) = \gamma H_{\mathcal{K}}(JM_0(1), JM_1(1)). \quad (29)$$

To obtain the lower bound, the parameter γ will be chosen as large as possible while making sure that the joint distributions of the observations have total variation distance smaller than some $C < 1$.

In the following, we will write A, B, C (with upper case letters) positive constants that can change from line to line. As in [17] and [20], we use the upper bound:

$$\|(G_0(\kappa) * Q)^{\otimes n} - (G_1(\kappa) * Q)^{\otimes n}\|_{TV} \leq 1 - (1 - \|(G_0(\kappa) * Q) - (G_1(\kappa) * Q)\|_{TV})^n,$$

where $\|\cdot\|_{TV}$ denotes the total variation distance. Using Le Cam's two-points method, the minimax rate will be lower bounded by $H(JM_0(\gamma), JM_1(\gamma))$, that is γ , (see Lemma 11) provided that there exists a constant $C > 0$ such that $\|(G_0(\kappa) * Q)^{\otimes n} - (G_1(\kappa) * Q)^{\otimes n}\|_{TV} \leq C < 1$, so that we only need to find $C > 0$ such that

$$\int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \leq \frac{C}{n}.$$

Since Q has a density q over \mathbb{R}^D , $G_0(\kappa) * Q$ and $G_1(\kappa) * Q$ also have a density over \mathbb{R}^D . We first prove that for $i \in \{0, 1\}$,

$$\int_{\mathbb{R}^D} \prod_{j=1}^D x_j^2 \left| \frac{d(G_i(\kappa) * Q)}{dx}(x) \right|^2 dx < +\infty. \quad (30)$$

Indeed,

$$\int_{\mathbb{R}^D} \prod_{j=1}^D x_j^2 \left| \frac{d(G_i(\kappa) * Q)}{dx}(x) \right|^2 dx \leq \left\| \frac{d(G_i(\kappa) * Q)}{dx} \right\|_{\infty} \int_{\mathbb{R}^D} \prod_{j=1}^D x_j^2 d(G_i(\kappa) * Q)(x).$$

First, $\left\| \frac{d(G_i(\kappa) * Q)}{dx} \right\|_\infty \leq \|q\|_\infty^D < \infty$. Moreover, for $k \in \{1, \dots, D\}$, writing $X_i(\kappa)^{[k]}$ and $\varepsilon^{[k]}$ for the k -th coordinate of $X_i(\kappa)$ and ε ,

$$\begin{aligned} \int_{\mathbb{R}^D} \prod_{j=1}^D x_j^2 |d(G_i(\kappa) * Q)(x)| &= \mathbb{E} \left[\prod_{k=1}^D (X_i(\kappa)^{[k]} + \varepsilon^{[k]})^2 \right] \\ &= \mathbb{E} [(X_i(\kappa)^{[1]} + \varepsilon^{[1]})^2 (X_i(\kappa)^{[2]} + \varepsilon^{[2]})^2] \prod_{k=3}^D \mathbb{E} [(\varepsilon^{[k]})^2]. \end{aligned} \quad (31)$$

We have that $(X_i(\kappa)^{[2]} + \varepsilon^{[2]})^2 \leq a^2 (X_i(\kappa)^{[1]})^2 + 2\gamma X_i(\kappa)^{[1]} + 2X_i(\kappa)^{[1]} \varepsilon^{[2]} + (1 + \gamma)(\varepsilon^{[2]})^2 + \gamma^2$, using (31) and the fact that $\varepsilon^{[2]}$ is independent of all other variables and that, for $k \in \{1, 2\}$, $X_i^{[1]}$ is independent of $\varepsilon^{[k]}$, we finally get that $\int \prod_{j=1}^D x_j^2 |d(G_i(\kappa) * Q)(x)|$ is upper bounded by product and sum of expectation of $((\varepsilon^{[j]})^2)_{j \in \{1, \dots, D\}}$, $(X_i(\kappa)^{[1]})^2$, $(X_i(\kappa)^{[1]})^3$ and $(X_i(\kappa)^{[1]})^4$ which are all finite thanks to Lemma 7.

By the Cauchy-Schwarz inequality,

$$\begin{aligned} \int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \\ \leq \pi^{D/2} \left(\int \prod_{j=1}^D (1 + x_j^2) \left| \frac{d((G_0(\kappa) - G_1(\kappa)) * Q)}{dx}(x) \right|^2 dx \right)^{1/2}. \end{aligned} \quad (32)$$

By Parseval's identity, for all $\eta \in \{0, 1\}^D$,

$$\begin{aligned} \int_{\mathbb{R}^D} \prod_{j=1}^D x_j^{2\eta_j} \left| \frac{d((G_0(\kappa) - G_1(\kappa)) * Q)}{dx}(x) \right|^2 dx \\ = \int_{\mathbb{R}^D} \left| \left(\prod_{j=1}^D \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[G_0(\kappa)] - \mathcal{F}[G_1(\kappa)])(t) \mathcal{F}[Q](t) \right|^2 dt \\ = \int_{[-c, c]^D} \left| \left(\prod_{j=1}^D \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[G_0(\kappa)] - \mathcal{F}[G_1(\kappa)])(t) \mathcal{F}[Q](t) \right|^2 dt, \end{aligned}$$

since $\mathcal{F}[Q]$ and for $\eta \in \{0, 1\}^D$, $\partial^\eta \mathcal{F}[Q]$ are supported on $[-c, c]^D$. Moreover, they are bounded functions, so that there exists a constant C (depending only on D) such that

$$\begin{aligned} \int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \\ \leq C \sum_{\eta \in \{0, 1\}^D} \int_{[-c, c]^D} \left| \left(\prod_{j=1}^D \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[G_0(\kappa)] - \mathcal{F}[G_1(\kappa)])(t) \right|^2 dt \\ = \sum_{\eta \in \{0, 1\}^D} \int_{[-c, c]^D} \left| \left(\prod_{j=1}^D \partial_{t_j}^{\eta_j} \right) (t \mapsto \mathcal{F}[S_0] - \mathcal{F}[S_1])(A_a^\top t) \right|^2 dt. \end{aligned}$$

Using the change of variable $u = A^\top t$, and noticing that $\{A^\top t ; t \in [-c, c]^D\} \subset [-2c, 2c]^D$, there exists a constant $C > 0$ depending on D and a such that

$$\begin{aligned} \int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \\ \leq C \sum_{\eta \in \{0, 1\}^D} \int_{[-2c, 2c]^D} \left| \left(\prod_{j=1}^D \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[S_0] - \mathcal{F}[S_1])(u) \right|^2 du. \end{aligned}$$

For all $t = (t_1, \dots, t_D) \in \mathbb{R}^D$, for $i \in \{0, 1\}$, $\mathcal{F}[T_i](t) = \mathcal{F}[\tilde{T}_i](t_1, t_2)$, where \tilde{T}_i is the distribution of the 2 first coordinates of $S_i(\kappa)$ under T_i . There exists a constant $C > 0$ such that

$$\begin{aligned} & \int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \\ & \leq C \sum_{\eta \in \{0,1\}^2} \int_{[-2c, 2c]^2} \left| \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \right|^2 dt. \end{aligned} \quad (33)$$

Following the same approach as [20], we get that for all $t = (t_1, t_2) \in \mathbb{R}^2$,

$$\begin{aligned} (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) &= \int_{\mathbb{R}} \{e^{it_1 u + i\gamma t_2 \tilde{g}_\gamma(u)} - e^{it_1 u - i\gamma t_2 \tilde{g}_\gamma(u)}\} f_\kappa(u) du \\ &= 2i \int_{\mathbb{R}} e^{it_1 u} \sin(t_2 \gamma \tilde{g}_\gamma(u)) f_\kappa(u) du \\ &= 2i \int_{\mathbb{R}} e^{it_1 u} \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k+1} \gamma^{2k+1}}{(2k+1)!} \tilde{g}_\gamma^{2k+1}(u) f_\kappa(u) du. \end{aligned}$$

Since $\sum_{k=0}^{\infty} \int_{\mathbb{R}} \frac{|t_2|^{2k+1} \gamma^{2k+1}}{(2k+1)!} |\tilde{g}_\gamma^{2k+1}(u)| f_\kappa(u) du$ is finite, we can switch integral and sum thanks to Fubini Theorem, so that

$$\begin{aligned} (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) &= 2i \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k+1} \gamma^{2k+1}}{(2k+1)!} \int_{\mathbb{R}} e^{it_1 u} \tilde{g}_\gamma^{2k+1}(u) f_\kappa(u) du \\ &= 2i \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k+1} \gamma^{2k+1}}{(2k+1)!} m_k(t_1), \end{aligned}$$

with for all $u \in \mathbb{R}$,

$$m_k(u) = \mathcal{F}[\tilde{g}^{2k+1} f_\kappa](u) = \underbrace{(\mathcal{F}[\tilde{g}] * \mathcal{F}[\tilde{g}] * \dots * \mathcal{F}[\tilde{g}] * \mathcal{F}[f_\kappa])}_{2k+1 \text{ times}}(u). \quad (34)$$

Since

$$\mathcal{F}[x \mapsto \cos(\frac{x}{\gamma})] = \frac{1}{2} \delta_{-\frac{1}{\gamma}} + \frac{1}{2} \delta_{\frac{1}{\gamma}},$$

for all $u \in \mathbb{R}$,

$$\begin{aligned} \underbrace{(\mathcal{F}[\tilde{g}] * \mathcal{F}[\tilde{g}] * \dots * \mathcal{F}[\tilde{g}])}_{2k+1 \text{ times}}(u) &= \underbrace{\mathcal{F}[\cos(\frac{\cdot}{\gamma}) * \dots * \mathcal{F}[\cos(\frac{\cdot}{\gamma})]}_{2k+1 \text{ times}} \\ &= \left(\frac{1}{2}\right)^{2k+1} \sum_{j=1}^{2k+1} \binom{2k+1}{j} \delta_{a_j}, \end{aligned}$$

where $a_j = (2j - 2k - 1)/\gamma$. By (34),

$$m_k(u) = \left(\frac{1}{2}\right)^{2k+1} \sum_{j=0}^{2k+1} \binom{2k+1}{j} \mathcal{F}[f_\kappa](u - a_j).$$

Therefore,

$$\sup_{|t| \leq c} |m_k(t)| \leq \sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_\kappa] \left(t - \frac{2j - 2k - 1}{\gamma} \right) \right|$$

and

$$\sup_{|t| \leq c} |m'_k(t)| \leq \sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_\kappa]' \left(t - \frac{2j - 2k - 1}{\gamma} \right) \right|.$$

Assume first that $\kappa \in (1/2, 1)$

For γ that satisfies $\gamma \leq \frac{1}{2c}$, by Lemma 7, there exist two constants A, B independent of γ and k such that

$$\sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_\kappa] \left(t - \frac{2j-2k-1}{\gamma} \right) \right| \leq A \exp(-B\gamma^{-\frac{1}{\kappa}})$$

and

$$\sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_\kappa]' \left(t - \frac{2j-2k-1}{\gamma} \right) \right| \leq A \exp(-B\gamma^{-\frac{1}{\kappa}}).$$

Thus,

$$\sup_{|t| \leq c} |m_k(t)| \leq A \exp(-B\gamma^{-\frac{1}{\kappa}}), \quad (35)$$

and

$$\sup_{|t| \leq c} |m'_k(t)| \leq A \exp(-B\gamma^{-\frac{1}{\kappa}}). \quad (36)$$

For all $\eta \in \{0, 1\}^2$, and $t \in [-c, c]^2$,

$$\begin{aligned} \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) &= \prod_{j=1}^2 \partial_{t_j}^{\eta_j} \left[2i \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k+1} \gamma^{2k+1}}{(2k+1)!} m_k(t_1) \right] \\ &= 2i\eta_2 \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k} \gamma^{2k+1}}{(2k)!} \partial_{t_1}^{\eta_1} m_k(t_1) + 2i(1-\eta_2) \sum_{k=0}^{\infty} \frac{(-1)^k t_2^{2k+1} \gamma^{2k+1}}{(2k+1)!} \partial_{t_1}^{\eta_1} m_k(t_1), \end{aligned}$$

so that

$$\begin{aligned} &\left| \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \right| \\ &\leq 2 \sum_{k=0}^{\infty} \frac{|t_2|^{2k} \gamma^{2k+1}}{(2k)!} |\partial_{t_1}^{\eta_1} m_k(t_1)| + 2 \sum_{k=0}^{\infty} \frac{|t_2|^{2k+1} \gamma^{2k+1}}{(2k+1)!} |\partial_{t_1}^{\eta_1} m_k(t_1)|. \end{aligned}$$

By (35) and (36), there exists a constant $C > 0$ which depends only on D and A such that

$$\left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{S}_0] - \mathcal{F}[\tilde{S}_1])(t) \leq C \exp(-B\gamma^{-\frac{1}{\kappa}}) \sup_{|t_2| \leq c} \left(\gamma \cosh(|t_2|\gamma) + \sinh(|t_2|\gamma) \right).$$

For γ small enough, there exists a constant $C_1 > 0$ which depends only on $D, A > 0$ and $C > 0$ such that

$$\left| \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \right| \leq C \exp(-B\gamma^{-\frac{1}{\kappa}}).$$

Finally, using (33), there exist constants $C > 0$ and $B > 0$ which depend only on D such that

$$\int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \leq C \exp(-B\gamma^{-\frac{1}{\kappa}}).$$

Taking $\gamma = c_\gamma (\log n)^{-\kappa}$ with $c_\gamma \leq B_1^\kappa$ shows that there exists $C > 0$ such that

$$\int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \leq \frac{C}{n}.$$

Let us now consider the case $\kappa = 1$

For γ that satisfies $\gamma \leq \frac{1}{2c}$, by Lemma 7, for all $\delta \in (0, 1)$, there exist two constants $A > 0$, $B > 0$ independent of γ and k such that

$$\sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_1] \left(t - \frac{2j-2k-1}{\gamma} \right) \right| \leq A \exp(-B\gamma^{-\delta}),$$

and

$$\sup_{|t| \leq c, 0 \leq j \leq 2k+1} \left| \mathcal{F}[f_1]' \left(t - \frac{2j-2k-1}{\gamma} \right) \right| \leq A \exp(-B\gamma^{-\delta}).$$

Thus, there exist constants $A > 0$ and $B > 0$ independent of γ and k such that

$$\sup_{|t| \leq c} |m_k(t)| \leq A \exp(-B\gamma^{-\delta}), \quad (37)$$

and

$$\sup_{|t| \leq c} |m'_k(t)| \leq A \exp(-B\gamma^{-\delta}). \quad (38)$$

Doing the same computation as in the case $\kappa \in (1/2, 1)$ shows that for all $\eta \in \{0, 1\}^2$ and $t \in [-c, c]^2$,

$$\begin{aligned} & \left| \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \right| \\ & \leq 2 \sum_{k=0}^{\infty} \frac{|t_2|^{2k} \gamma^{2k+1}}{(2k)!} |\partial_{t_1}^{\eta_1} m_k(t_1)| + 2 \sum_{k=0}^{\infty} \frac{|t_2|^{2k+1} \gamma^{2k+1}}{(2k+1)!} |\partial_{t_1}^{\eta_1} m_k(t_1)|. \end{aligned}$$

By (37) and (38), there exist constants $C > 0$ and $B > 0$ which depend only on D such that

$$\left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \leq C \exp(-B\gamma^{-\delta}) \sup_{|t_2| \leq c} \left(\gamma \cosh(|t_2|\gamma) + \sinh(|t_2|\gamma) \right).$$

For γ small enough, there exists a constant $C > 0$ which depends only on D such that

$$\left| \left(\prod_{j=1}^2 \partial_{t_j}^{\eta_j} \right) (\mathcal{F}[\tilde{T}_0] - \mathcal{F}[\tilde{T}_1])(t) \right| \leq C \exp(-B\gamma^{-\delta}).$$

Finally, using (33), there exists a constant $C > 0$ which depends only on D such that

$$\int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \leq C \exp(-B\gamma^{-\delta}).$$

Taking $\gamma = c_\gamma (\log n)^{-\frac{1}{\delta}}$ with $c_\gamma \leq B^{\frac{1}{\delta+1}}$ shows that there exists $C > 0$ such that

$$\int_{\mathbb{R}^D} |d(G_0(\kappa) * Q)(x) - d(G_1(\kappa) * Q)(x)| \leq \frac{C}{n}.$$

D.7 Proof of Lemma 7

Case $\kappa \neq 1$.

This case is based on [23]. In the following, we will note constants that can change with upper case A , B and C . In [23], Theorem 2, the author defines for any positive constants $\mu > 0$, $q > 1$ and $a > 0$ a function $\zeta_{q,\mu,a}$, such that for $x \in \mathbb{R}$,

$$\zeta_{q,\mu,a}(x) = -i \int_{\mathcal{C}} z^\mu \exp(z^q - qax^2z) dz,$$

where \mathcal{C} is a curve in the complex plane so that the maximum of $|z^\mu \exp(z^q - qax^2z)|$ for $z \in \mathcal{C}$ is attained on the positive real line. The author shows that $\zeta_{q,\mu,a}$ and $\zeta_{q,\mu,a}^2$ are integrable functions.

The author uses the saddle-point integration method to show that there exist $A > 0$ and $B > 0$ which depend on q, μ and a such that

$$|\mathcal{F}[\zeta_{q,\mu,a}](t)| \leq A \exp(-Bx^{\frac{2q}{q+1}}). \quad (39)$$

Finally, for $\kappa \in (1/2, 1)$, fix $\mu > 0, a > 0$, and define

$$f_\kappa = c_{f_\kappa} \operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]^2 * u_1,$$

where $u_1 : x \in \mathbb{R} \mapsto \exp(-\frac{1}{1-4x^2})1_{(-1/2,1/2)}(x)$ and c_{f_κ} is a constant that ensures that f_κ is a density.

Let us first prove that there exist A and B positive constants such that $|\mathcal{F}[\operatorname{Re}[\zeta_{q,\mu,a}]^2](t)| \leq A \exp(-B|t|^{1/\kappa})$.

$$\begin{aligned} |\mathcal{F}[\operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]^2](t)| &= |\mathcal{F}[\operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}] * \mathcal{F}[\operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]](t)| \\ &\leq A \int_{\mathbb{R}} \exp(-B|x-y|^{1/\kappa} - B|y|^{1/\kappa}) dy \\ &= \int_{|y-x| \geq |x|/2} \exp(-B|x-y|^{1/\kappa} - B|y|^{1/\kappa}) dy \\ &\quad + \int_{|y-x| < |x|/2} \exp(-B|x-y|^{1/\kappa} - B|y|^{1/\kappa}) dy \\ &\leq A \exp(-B|x|^{1/\kappa}). \end{aligned} \quad (40)$$

Finally, for all $t \in \mathbb{R}$, using that $|\mathcal{F}[u_1](t)| \leq 1$,

$$\begin{aligned} |\mathcal{F}[f_\kappa](t)| &= |\mathcal{F}[\operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]^2](t)| |\mathcal{F}[u_1](t)| \\ &\leq c_{f_\kappa} A \exp(-B|x|^{1/\kappa}). \end{aligned}$$

For $x \in \mathbb{R}$, $\mathcal{F}[f_\kappa(x)]' = \mathcal{F}[x \mapsto x f_\kappa(x)]$ and

$$x f_\kappa(x) = c_{f_\kappa} v * \operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]^2(x) + c_{f_\kappa} u_1 * \tilde{\zeta}(x),$$

where $v : x \in \mathbb{R} \mapsto x u_1(x)$ and $\tilde{\zeta} : x \in \mathbb{R} \mapsto x \operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]^2(x)$.

Following the same proof as Theorem 2 of [23], there exists $A > 0$ and $B > 0$ such that for all $t \in \mathbb{R}$, $|\mathcal{F}[x \mapsto x \operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]](t)| \leq A \exp(-B|t|^{1/\kappa})$, so that, following the proof of (40), $|\mathcal{F}[\tilde{\zeta}](t)| \leq A \exp(-B|t|^{1/\kappa})$. Hence, there exists $A > 0$ and $B > 0$ such that $|\mathcal{F}[f_\kappa]'(t)| \leq A \exp(-B|t|^{1/\kappa})$.

Finally, note that f_κ is continuous as a convolution of an integrable function with a smooth function, and that for all $x \in \mathbb{R}$, $f_\kappa(x) > 0$ since $\operatorname{Re}[\zeta_{\frac{1}{2\kappa-1},\mu,a}]$ and u are not the null function almost everywhere.

Case $\kappa = 1$.

Let $\delta \in (0, 1)$ and define $f_1 : x \in \mathbb{R} \mapsto c_{f_1} (u_{\frac{1}{1-\delta}} * u_{\frac{1}{1-\delta}})(x)$, where c_{f_1} is a constant that ensures that f_1 is a probability density.

There exist $A > 0$ and $B > 0$ such that $\mathcal{F}[f_1](x) \leq A \exp(-B|x|^\delta)$, see Lemma in [26]. Moreover, $\mathcal{F}[f_1]'(x) = 2c_{f_1} \mathcal{F}[u_{\frac{1}{1-\delta}}](x) \mathcal{F}[u_{\frac{1}{1-\delta}}]'(x) \leq A \|x \mapsto x u_{\frac{1}{1-\delta}}(x)\|_1 \exp(-B|x|^\delta)$.

Finally, note that f_1 is continuous and does not vanish on its support.

D.8 Proof of Lemma 8

First, by Lemma 7, for any $\kappa \in (1/2, 1]$, $U(\kappa)$ satisfies $A(\kappa)$.

Let $i \in \{0, 1\}$. For any $\lambda = (\lambda_1, \dots, \lambda_D) \in \mathbb{R}^D$,

$$\begin{aligned} \mathbb{E}[\exp(\lambda^\top X_i(\kappa))] &= \mathbb{E} \left[\exp \left(\left(\lambda_1 + \frac{1}{2} \lambda_2 \right) U(\kappa) + (-1)^i \gamma \lambda_2 \frac{1}{2} \cos \left(\frac{U(\kappa)}{\gamma} \right) \right) \right] \\ &\leq e^{\gamma \frac{1}{2} |\lambda_2|} \mathbb{E} \left[\exp \left(\left(\lambda_1 + \frac{1}{2} \lambda_2 \right) U(\kappa) \right) \right]. \end{aligned} \quad (41)$$

Since $U(\kappa)$ satisfies $A(\kappa)$, there exist positive constants $A > 0$ and $B > 0$ such that for all $\lambda = (\lambda_1, \dots, \lambda_D) \in \mathbb{R}^D$, $\mathbb{E}[\exp(\lambda^\top X_i(\kappa))] \leq A \exp(B|\lambda|^{1/\kappa})$. Applying this in (41),

$$\begin{aligned} \mathbb{E}[\exp(\lambda^\top X_i(\kappa))] &\leq A \exp \left(\gamma \frac{1}{2} |\lambda_2| + B \left| \lambda_1 + \frac{1}{2} \lambda_2 \right|^{1/\kappa} \right) \\ &\leq A' \exp(B' |\lambda|^{1/\kappa}) \end{aligned}$$

for some other constants A' and B' since $1 \leq \kappa$, so that $X_i(\kappa)$ satisfies $A(\kappa)$.

D.9 Proof of Lemma 9

The proof is done in five steps.

1. We show that γg_γ is 1-lipschitz.
2. For $i \in \{0, 1\}$ and $\kappa \in (\frac{1}{2}, 1]$, we compute the density p_i of $T_i(\kappa)$ with respect to the 1-dimensional Hausdorff measure μ_H and we show that for any compact set \mathcal{K} , there exists $b(\kappa, \mathcal{K}) > 0$ such that, for all $u \in M_i(\gamma) \cap \mathcal{K}$, $|p_i(u)| \geq b(\kappa, \mathcal{K})$.
3. We show that for $i \in \{0, 1\}$, $\mu_H(\cdot \cap M_i(\gamma))$ is in $St_{\mathcal{K}}(2, d, r_0)$.
4. We deduce that for $i \in \{0, 1\}$ and $d \geq 1$, T_i is in $St_{\mathcal{K}}(2b(\kappa, \mathcal{K}), d, r_0)$.
5. Finally, we show that for $i \in \{0, 1\}$, $d \geq 1$ and a small enough, $G_i(\kappa) \in St_{\mathcal{K}}(a, d, r_0)$.

Proof of 1 For all $x \in \mathbb{R}$, $|\gamma \tilde{g}'_\gamma(x)| = |\sin(\frac{x}{\gamma})| \leq 1$, which implies that γg_γ is 1-Lipschitz.

Proof of 2 Let us first compute the density p_i of $T_i(\kappa)$ with respect to μ_H . For $i \in \{0, 1\}$, denote $\zeta_i : x \in \mathbb{R} \mapsto (x, (-1)^i \gamma g_\gamma(x))$. Let \mathcal{B} be an open subset of \mathbb{R}^D . For any $\kappa \in (\frac{1}{2}, 1]$,

$$T_i(\kappa)(\mathcal{B}) = \mathbb{P}[\zeta_i(U(\kappa)) \in \mathcal{B}] = \mathbb{P}[U(\kappa) \in \zeta_i^{-1}(\mathcal{B})] = \int_{\zeta_i^{-1}(\mathcal{B})} f_\kappa(u) du.$$

Let $\text{Jac } \zeta_i : u \in \mathbb{R} \mapsto \sqrt{1 + \gamma^2 \tilde{g}_\gamma(u)^2}$ be the Jacobian of ζ_i . By the Area Formula (see equation (2.47) in [4]),

$$s_i(\kappa)(\mathcal{B}) = \int_{\zeta_i^{-1}(\mathcal{B})} \frac{f_\kappa(u)}{\text{Jac } \zeta_i(u)} \text{Jac } \zeta_i(u) du = \int_{\mathcal{B} \cap M_i(\gamma)} \frac{f_\kappa(\pi^{(1)}(u))}{\text{Jac } \zeta_i(\pi^{(1)}(u))} d\mu_H(u).$$

We then have that for all $x \in \mathbb{R}^D$,

$$p_i(x) = \frac{f_\kappa(\pi^{(1)}(x))}{\text{Jac } \zeta_i(\pi^{(1)}(x))} \mathbf{1}_{M_i(\gamma)}(x).$$

Since f_κ is continuous and does not vanish on its support, for any compact set \mathcal{K} , $M_i(\gamma) \cap \mathcal{K}$ is a compact subset of the support of f_κ . Thus, since f_κ is continuous and does not vanish on its support, for any compact set \mathcal{K} , there exists $c(\kappa, \mathcal{K}) > 0$ such that for all $u \in M_i(\gamma) \cap \mathcal{K}$, $f_\kappa(u) \geq c(\kappa, \mathcal{K})$. Moreover, for $i \in \{0, 1\}$, $\text{Jac } \zeta_i(u) \leq \sqrt{2}$. Therefore, for all $x \in M_i(\gamma) \cap \mathcal{K}$, $|p_i(x)| \geq \frac{c(\kappa, \mathcal{K})}{\sqrt{2}}$.

Proof of 3 Recall that the 1-dimensional Hausdorff measure μ_H is defined as the limit $\lim_{\eta \rightarrow 0} \mu_H^\eta$, where for any set Z

$$\mu_H^\eta(Z) = \inf \left\{ \sum_{i \in \mathbb{N}} \text{Diam}(A_i) : Z \subset \bigcup_i A_i \text{ and } \forall i, \text{Diam}(A_i) \leq \eta \right\}.$$

For any $z \in M_i(\gamma)$, there exists $x_0 \in \mathbb{R}$ such that $z = (x_0, (-1)^i \gamma g_\gamma(x_0))$ and, for any $r > 0$,

$$B(z, r) \cap M_i(\gamma) \supset \{(x, (-1)^i \gamma g_\gamma(x)), x \in B(x_0, r)\}$$

since $|x - x_0| \leq r$ implies $\|\gamma g_\gamma(x) - \gamma g_\gamma(x_0)\|_\infty \leq r$.

Let $(A_i)_{i \in \mathbb{N}}$ be a covering of $\{(x, (-1)^i \gamma g_\gamma(x)), x \in B(x_0, r)\}$, and $B_i = \pi^{(1)}(A_i)$, then B_i is a covering of $B(x_0, r)$. For all $\eta > 0$,

$$\mu_H^\eta(\{(x, (-1)^i \gamma g_\gamma(x)), x \in B(x_0, r)\}) \geq \mu_H^\eta(B(x_0, r)),$$

thus $\mu_H(B(z, r) \cap M_i(\gamma)) \geq \mu_H(B(x_0, r)) = 2r$. If $r_0 \leq 1$, then for any $r \leq r_0$,

$$\mu_H(B(z, r) \cap M_i(\gamma)) \geq 2r^d,$$

which proves 3.

Proof of 4 Let $x_i \in M_i(\gamma) \cap \mathcal{K}$ and $r_0 < 1$. Then for all $r \leq r_0$,

$$\begin{aligned} T_i(B(x_i, r) \cap M_i(\gamma)) &= \int_{B(x_i, r) \cap M_i(\gamma)} p_i(u) d\mu_H(u) \\ &\geq b(\kappa, \mathcal{K}) \mu_H(B(x_i, r) \cap M_i(\gamma)) \geq 2b(\kappa, \mathcal{K}) r^d. \end{aligned}$$

Proof of 5 For $i \in \{0, 1\}$, let $x_i \in JM_i(\gamma) \cap \mathcal{K}$, $r_0 < 1$, and take $\tilde{\mathcal{K}}$ such that $J^{-1}\mathcal{K} \subset \tilde{\mathcal{K}}$. For all $r \leq r_0$,

$$\begin{aligned} G_i(\kappa)(B(x_i, r)) &= \mathbb{P}[JS_i(\kappa) \in B(x_i, r)] \geq \mathbb{P}\left[S_i(\kappa) \in B\left(J^{-1}x_i, \frac{r}{\|J\|_{\text{op}}}\right)\right] \\ &\geq \frac{2b(\kappa, J^{-1}\mathcal{K})}{\|J\|_{\text{op}}} r \\ &\geq \frac{2b(\kappa, \tilde{\mathcal{K}})}{\|J\|_{\text{op}}} r, \end{aligned}$$

so that for some a_0 and all $a \leq a_0$, $G_i(\kappa)(B(x_i, r)) \geq ar^d$.

D.10 Proof of Lemma 10

Let us write $m_{i,\gamma}(x) = (x + (-1)^i \frac{\gamma}{2} \cos(\frac{x}{\gamma}), 0, \dots, 0)$, so that $X_i(\kappa) = (U(\kappa), m_{i,\gamma}(U(\kappa)))$. For $i \in \{0, 1\}$, let $w_{i,\kappa,\gamma}$ be the density of the first coordinate of $m_{i,\gamma}(U(\kappa))$, then

$$M_\kappa = \sup_{x \in \mathbb{R}, \gamma \in [0,1], i \in \{0,1\}} \{w_{i,\kappa,\gamma}(x) \vee f_\kappa(x)\}$$

is an upper bound of the density of $X_i(\kappa)^{(1)}$ and of the first coordinate of $X_i(\kappa)^{(2)}$ with respect to the Lebesgue measure. Let us show that M_κ is finite. First, note that $m_{i,\gamma}$ is one-to-one from \mathbb{R} to $\mathbb{R} \times \{0\}^{D-2}$ and $m_{i,\gamma}^{-1}$ is Lipschitz with Lipschitz constant upper bounded by 1/2. One can easily check that for all $x \in \mathbb{R}$,

$$w_{i,\kappa,\gamma}(x) = f_\kappa((m_{i,\gamma})_1(x, 0, \dots, 0)) \frac{1}{(m_{i,\gamma})'_1(m_{i,\gamma}^{-1}(x, 0, \dots, 0))},$$

where $(m_{i,\gamma})_1(x)$ is the first coordinate of $m_{i,\gamma}(x)$. Since $(m_{i,\gamma})'_1$ is lower bounded by 1/2, $M_\kappa \leq \sup_{x \in \mathbb{R}} f_\kappa(x)$, which is finite.

For any $\Delta > 0$, define the sets:

$$\begin{aligned} A_{\Delta}^{(1)} &= [-\Delta, \Delta] \quad \text{and} \quad B_{\Delta}^{(2)} = \bar{B}\left(0, \left(\frac{1}{2} + 2\right)\Delta\right) \cap (\mathbb{R} \times \{0\}^{D-2}), \\ A_{\Delta}^{(2)} &= [-\Delta, \Delta] \times \{0\}^{D-2} \quad \text{and} \quad B_{\Delta}^{(1)} = \bar{B}(0, \Delta) \cap \mathbb{R}. \end{aligned}$$

Define $c_{\Delta, \kappa} = \mathbb{P}[U(\kappa) \in A_{\Delta}^{(1)}] \wedge \inf_{\gamma \in [0, 1], i \in \{0, 1\}} \mathbb{P}[m_{i, \gamma}(U(\kappa)) \in A_{\Delta}^{(2)}]$, and let us prove that $c_{\Delta, \kappa} > 0$.

First, $\mathbb{P}[U(\kappa) \in A_{\Delta}^{(1)}] > 0$ since the density of $U(\kappa)$ is positive everywhere on its support. Then, for $i \in \{0, 1\}$,

$$\begin{aligned} \mathbb{P}[m_{i, \gamma}(U(\kappa)) \in A_{\Delta}^{(2)}] &= \mathbb{P}\left(U(\kappa) + (-1)^i \frac{1}{2} \gamma \cos\left(\frac{U(\kappa)}{\gamma}\right) \in [-\Delta, \Delta]\right) \\ &\geq \mathbb{P}\left(U(\kappa) \in [-\Delta/2, \Delta/2], (-1)^i \frac{1}{2} \gamma \cos\left(\frac{U(\kappa)}{\gamma}\right) \in [-\Delta/2, \Delta/2]\right) \\ &\geq \mathbb{P}\left(U(\kappa) \in [-\Delta/2, \Delta/2], \cos\left(\frac{U(\kappa)}{\gamma}\right) \in [-\Delta/\gamma, \Delta/\gamma]\right) \\ &\geq \mathbb{P}\left(U(\kappa) \in \left[-\frac{\Delta}{2}, \frac{\Delta}{2}\right] \cap [\arccos(\Delta), \pi - \arccos(\Delta)]\right), \end{aligned}$$

which is positive.

For any $\Delta > 0$ define $B_{\Delta, i, \gamma}^{(1)} = m_{i, \gamma}^{-1}(A_{\Delta}^{(2)})$. Then

$$\begin{aligned} \text{Diam}(B_{\Delta, i, \gamma}^{(1)}) &= \sup_{x, y \in B_{\Delta, i, \gamma}^{(1)}} |x - y| \\ &= \sup_{x, y \in A_{\Delta}^{(2)}} |m_{i, \gamma}^{-1}(x) - m_{i, \gamma}^{-1}(y)| \\ &\leq \frac{1}{2} \sup_{x, y \in A_{\Delta}^{(2)}} \|x - y\| \leq \Delta. \end{aligned}$$

Thus, $B_{\Delta, i, \gamma}^{(1)} \subset B_{\Delta}^{(1)}$, and

$$\mathbb{P}[(X_i(\kappa))^{(1)} \in B_{\Delta, i, \gamma}^{(1)} | (X_i(\kappa))^{(2)} \in A_{\Delta}^{(2)}] \geq \mathbb{P}[(X_i(\kappa))^{(1)} \in B_{\Delta, i, \gamma}^{(1)} | (X_i(\kappa))^{(2)} \in A_{\Delta}^{(2)}] = 1.$$

Similarly, define $B_{\Delta, i, \gamma}^{(2)} = m_{i, \gamma}(A_{\Delta}^{(1)}) = \{(x + (-1)^i \gamma \frac{1}{2} \cos(\frac{x}{\gamma}), 0, \dots, 0), x \in [-\Delta, \Delta]\}$, then

$$\begin{aligned} \text{Diam}(B_{\Delta, i, \gamma}^{(2)}) &= \sup_{x, y \in A_{\Delta}^{(1)}} \left| x + (-1)^i \gamma \frac{1}{2} \cos\left(\frac{x}{\gamma}\right) - y - (-1)^i \gamma \frac{1}{2} \cos\left(\frac{y}{\gamma}\right) \right| \\ &\leq 2\Delta + \gamma \frac{1}{2} \left| \cos\left(\frac{x}{\gamma}\right) - \cos\left(\frac{y}{\gamma}\right) \right| \\ &\leq \left(\frac{1}{2} + 2\right) \Delta. \end{aligned}$$

Thus, $B_{\Delta, i, \gamma}^{(2)} \subset B_{\Delta}^{(2)}$, and

$$\mathbb{P}[(X_i(\kappa))^{(2)} \in B_{\Delta, i, \gamma}^{(2)} | (X_i(\kappa))^{(1)} \in A_{\Delta}^{(1)}] \geq \mathbb{P}[(X_i(\kappa))^{(2)} \in B_{\Delta, i, \gamma}^{(2)} | (X_i(\kappa))^{(1)} \in A_{\Delta}^{(1)}] = 1.$$

Thus, the distribution of $X_i(\kappa)$ satisfies (H1) and (H2). Lemma 10 follows from Lemma 8 and Theorem 2.

E Proofs for Section 5

E.1 Proof of Theorem 6

We shall need two technical lemmas. The following one is easily proved following the arguments at the end of the proof of Theorem 3.

Lemma 12. Let G be a probability measure with compact support \mathcal{M}_G . Assume that $G \in St_{\mathcal{M}_G}(a, d, r_0)$ for some constants $a > 0$, $d > 0$ and $r_0 > 0$. Recall that $\Gamma_n := \Gamma_{n,1} = \|\hat{g}_n - \bar{g}\|_\infty$. Then

(1) For any $C_1 > 0$ and $c > 0$, there exists $h_0 > 0$ such that if $h_n \leq h_0$, on the event where

$$C_1 + \Gamma_n < \lambda_n < ac_A^d d_A \left(\frac{1}{h_n}\right)^{D-d} - \Gamma_n,$$

it holds

$$\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_c.$$

(2) For m_n, h_n, λ_n chosen as in Theorem 3, for all $C_1 \in (0, ac_A^d d_A)$ and $\delta' > 0$, there exist $C > 0$ and $n_0 \geq 0$ such that for all $n \geq n_0$, with probability at least $1 - 2\exp(-n^{1/2-\delta'})$,

$$\Gamma_n^2 \leq Ce^{-m_n} \quad \text{and} \quad C_1 + \Gamma_n < \lambda_n < ac_A^d d_A \left(\frac{1}{h_n}\right)^{D-d} - \Gamma_n.$$

and in particular, for all $c > 0$ and $\delta' > 0$, there exists $n'_0 \geq 0$ such that for all $n \geq n'_0$, with probability at least $1 - 2\exp(-n^{1/2-\delta'})$,

$$\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_c.$$

In particular, since $R_n \rightarrow +\infty$ and \mathcal{M}_G is compact, up to increasing n_0 , on this event,

$$\mathcal{M}_G \subset \widehat{\mathcal{M}} \cap \bar{B}(0, R_n) \subset (\mathcal{M}_G)_c.$$

In the rest of the proof of the Theorem, we lighten the notation $\widehat{\mathcal{M}} \cap \bar{B}(0, R_n)$ into $\widehat{\mathcal{M}}$ (equivalently, we redefine the estimator $\hat{\mathcal{M}}$ as the intersection of the estimator of Section 4.1 with the closed euclidean ball of radius R_n).

Lemma 13. Let G be a probability measure with compact support \mathcal{M}_G . Assume that $G \in St_{\mathcal{M}_G}(a, d, r_0)$ for some constants $a > 0$, $d > 0$ and $r_0 > 0$. Then for any $\alpha > 0$, $c > 0$ and $p \in [1, +\infty)$, there exists $C(\alpha, c) > 0$ such that, on the event where

$$\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_c,$$

it holds

$$\|\bar{g}\|_{L_1(\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c)} \leq C(\alpha, c) h_n^\alpha \quad \text{and} \quad \int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c} \|x\|^p |\bar{g}(x)| dx \leq C(\alpha, c) h_n^\alpha.$$

Proof. By definition,

$$\|\bar{g}\|_{L_1(\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c)} = \frac{1}{h_n^D} \int_{x \in \mathcal{M}_G} \int_{y \in \mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c} \psi_A \left(\frac{\|y - x\|_2}{h_n} \right) dy dG(x).$$

By Point (V) of the properties of ψ_A , for any $A > 0$, there exists $C > 0$ such that for any $x \in \mathcal{M}_G$ and $y \in \mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c$,

$$\psi_A \left(\frac{\|y - x\|_2}{h_n} \right) \leq C \exp \left(-\beta_A \frac{\|y - x\|_2^{A/(A+1)}}{h_n^{A/(A+1)}} \right) \leq C \exp \left(-\beta_A \frac{d(y, \mathcal{M}_G)^{A/(A+1)}}{h_n^{A/(A+1)}} \right).$$

Since $\mathcal{M}_G \subset \widehat{\mathcal{M}}$, for all $y \in \mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c$, $d(y, \mathcal{M}_G) \geq c$, so for any $\alpha > 0$, there exists a constant $\tilde{C} > 0$ such that

$$C \exp \left(-\beta_A \frac{d(y, \mathcal{M}_G)^{A/(A+1)}}{h_n^{A/(A+1)}} \right) \leq \tilde{C} \frac{h_n^{D+\alpha}}{d(y, \mathcal{M}_G)^{D+\alpha}}.$$

Moreover, since \mathcal{M}_G is compact, $\text{Diam}(\mathcal{M}_G)$ is finite, so that on the event where $\mathcal{M}_G \subset \widehat{\mathcal{M}}$,

$$\begin{aligned} \int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c} \frac{1}{d(y, \mathcal{M}_G)^{D+\alpha}} dy &\leq \int_{\mathbb{R}^D \setminus (\mathcal{M}_G)_c} \frac{1}{d(y, \mathcal{M}_G)^{D+\alpha}} dy < \infty. \\ &\leq \int_{\mathbb{R}^D} \left(\frac{1}{c \vee (\|y\| - \text{Diam}(\mathcal{M}_G)/2)} \right)^{D+\alpha} dy < \infty. \end{aligned}$$

Therefore, for all $c > 0$ and $\alpha > 0$, there exists C depending on A, D, c, α and $\text{Diam}(\mathcal{M}_G)$ such that

$$\|\bar{g}\|_{L^1(\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c)} \leq Ch_n^\alpha.$$

The proof that the same holds for $\int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_c} \|x\|^p \bar{g}(x) dx$ is similar. \square

Let $G \in St_{\mathcal{M}_G}(a, d, r_0)$ be such that if $X \sim G$, then $\Phi_X \in \mathcal{H} \cap \Upsilon_{1,S}$. Fix $p \in [1, +\infty)$. We use a bias-variance decomposition of $W_p(G, \widehat{P}_{n,\eta})$ through the triangle inequality

$$W_p(G, \widehat{P}_{n,\eta}) \leq W_p(G, P_{\psi_{A,h}}) + W_p(P_{\psi_{A,h}}, \widehat{P}_{n,\eta}).$$

The proof is done in several steps :

- (1) We first show that there exists $C > 0$ depending only on A and D such that the bias satisfies

$$W_p(G, P_{\psi_{A,h_n}}) \leq Ch_n.$$

- (2) We prove that for any $\alpha \geq 1$, on the event where

$$\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_c,$$

there exists $C' > 0$ such that

$$W_p(P_{\psi_{A,h_n}}, \widehat{P}_{n,\eta}) \leq C'(h_n^\alpha + \Gamma_n).$$

- (3) We show that the choice of the parameters m_n, h_n and λ_n gives the result.

Proof of (1)

Let Y_ψ be a random variable with density ψ_{A,h_n} and independent of X , so that the distribution of $X + Y_\psi$ is $P_{\psi_{A,h_n}}$. By definition of W_p ,

$$W_p^p(G, P_{\psi_{A,h_n}}) \leq \mathbb{E}(\|X + Y_\psi - X\|_2^p) = \mathbb{E}(\|Y_\psi\|_2^p) = h_n^p \int_{\mathbb{R}^D} \|u\|^p \psi_{A,1}(u) du,$$

and the integral is finite by point (V) of the properties of ψ_A .

Proof of (2)

If ν and μ are probability measures on \mathbb{R}^D having respective densities f and g with respect to the Lebesgue measure, letting ω be the measure with density $\min(f, g)$ with respect to the Lebesgue measure, $a \in \mathbb{R}^D$, and δ_a the Dirac measure in a , by convexity of $x \mapsto x^p$,

$$\begin{aligned} W_p^p(\mu, \nu) &\leq 2^{p-1} (W_p^p(\mu, \omega + (1 - \omega(\mathbb{R}^D))\delta_a) + W_p^p(\nu, \omega + (1 - \omega(\mathbb{R}^D))\delta_a)) \\ &\leq 2^{p-1} \int_{\mathbb{R}^D} \|x - a\|^p (f(x) - g(x) - 2 \min(f(x), g(x))) dx \\ &= 2^{p-1} \int_{\mathbb{R}^D} \|x - a\|^p |f(x) - g(x)| dx, \end{aligned}$$

so that

$$W_p^p(\mu, \nu) \leq 2^{p-1} \min_{a \in \mathbb{R}^D} \int_{\mathbb{R}^D} \|x - a\|^p |f(x) - g(x)| dx. \quad (42)$$

This entails

$$\begin{aligned} W_p^p(P_{\psi_{A,h_n}}, \widehat{P}_{n,\eta}) &\leq 2^{p-1} \min_{a \in \mathbb{R}^D} \int_{\mathbb{R}^D} \|x - a\|^p |\bar{g}(x) - c_n \widehat{g}_n^+(x) 1_{(\widehat{\mathcal{M}})_\eta}(x)| dx \\ &\leq 2^{p-1} \int_{(\widehat{\mathcal{M}})_\eta} \|x\|^p |\bar{g}(x) - c_n \widehat{g}_n^+(x)| dx + 2^{p-1} \int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_\eta} \|x\|^p \bar{g}(x) dx. \end{aligned} \quad (43)$$

For S compact subset of \mathbb{R}^D , write $M_S = \sup_{x \in S} \|x\|^p$ and $\text{Vol}(S)$ for the Lebesgue measure of S , then

$$\begin{aligned} \int_{(\widehat{\mathcal{M}})_\eta} \|x\|^p |\bar{g}(x) - c_n \widehat{g}_n^+(x)| dx &\leq M_{(\widehat{\mathcal{M}})_\eta} \int_{(\widehat{\mathcal{M}})_\eta} |\widehat{g}_n^+(x) - \bar{g}(x)| dx + M_{(\widehat{\mathcal{M}})_\eta} \frac{|c_n - 1|}{c_n} \\ &\leq M_{(\widehat{\mathcal{M}})_\eta} \text{Vol}((\widehat{\mathcal{M}})_\eta) \Gamma_n + M_{(\widehat{\mathcal{M}})_\eta} \frac{|c_n - 1|}{c_n}. \end{aligned}$$

We also have

$$\begin{aligned} \frac{|c_n - 1|}{c_n} &= \left| \frac{1}{c_n} - 1 \right| = \left| \int_{(\widehat{\mathcal{M}})_\eta} (\widehat{g}_n^+(y) - \bar{g}(y)) dy - \int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_\eta} \bar{g}(y) dy \right| \\ &\leq \|\widehat{g}_n - \bar{g}\|_{L_1((\widehat{\mathcal{M}})_\eta)} + \|\bar{g}\|_{L_1(\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_\eta)}. \end{aligned}$$

Using Hölder's inequality,

$$\|\widehat{g}_n - \bar{g}\|_{L_1((\widehat{\mathcal{M}})_\eta)} \leq \text{Vol}((\widehat{\mathcal{M}})_\eta) \Gamma_n.$$

By Lemma 13, for any $\alpha > 0$, there exists C such that

$$\int_{(\widehat{\mathcal{M}})_\eta} \|x\|^p |\bar{g}(x) - c_n \widehat{g}_n^+(x)| dx \leq 2M_{(\widehat{\mathcal{M}})_\eta} \text{Vol}((\widehat{\mathcal{M}})_\eta) \Gamma_n + M_{(\widehat{\mathcal{M}})_\eta} C h_n^\alpha.$$

For any $c > 0$, when $\widehat{\mathcal{M}} \subset (\mathcal{M}_G)_c$, one has $(\widehat{\mathcal{M}})_\eta \subset (\mathcal{M}_G)_{\eta+c}$. This inclusion entails $M_{(\widehat{\mathcal{M}})_\eta} \leq M_{(\mathcal{M}_G)_{\eta+c}}$ and $\text{Vol}((\widehat{\mathcal{M}})_\eta) \leq \text{Vol}((\mathcal{M}_G)_{\eta+c})$. Therefore, for any $c > 0$,

$$\int_{(\widehat{\mathcal{M}})_\eta} \|x\|^p |\bar{g}(x) - c_n \widehat{g}_n^+(x)| dx \leq 2M_{(\mathcal{M}_G)_{\eta+c}} \text{Vol}((\mathcal{M}_G)_{\eta+c}) \Gamma_n + M_{(\mathcal{M}_G)_{\eta+c}} C h_n^\alpha. \quad (44)$$

Again by Lemma 13, on the event where $\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_\eta$,

$$\int_{\mathbb{R}^D \setminus (\widehat{\mathcal{M}})_\eta} \|x\|^p |\bar{g}(x)| dx \leq C' h_n^\alpha. \quad (45)$$

Finally, using (43), (44) and (45), for any $\alpha \geq 1$, there exists $C > 0$ such that

$$W_p^p(P_{\psi_{A,h_n}}, \widehat{P}_{n,\eta}) \leq C(h_n^\alpha + \Gamma_n).$$

Proof of (3)

Using (1) and (2), for sequences h_n , m_n and λ_n satisfying the assumptions of Theorem 6, on the event where $\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_\eta$, for any $\alpha \geq p$, there exists $C > 0$ such that

$$W_p(G, \widehat{P}_{n,\eta}) \leq C(h_n + (h_n^\alpha + \Gamma_n)^{1/p}) \leq 2C(h_n + \Gamma_n^{1/p}).$$

We may assume $h_n \leq 1$ for all n without loss of generality. As stated in Lemma 12, for any $\delta' > 0$, there exist C' and n_0 such that for all $n \geq n_0$, with probability at least $1 - 2 \exp(-n^{1/2-\delta'})$, $\Gamma_n^{1/p} \leq C' e^{-m_n/(2p)}$ and $\mathcal{M}_G \subset \widehat{\mathcal{M}} \subset (\mathcal{M}_G)_\eta$, and therefore

$$W_p(G, \widehat{P}_{n,\eta^*}) \leq C m_n^{-1}$$

on this event, up to changing the constant C .

On the event of probability at most $2\exp(-n^{1/2-\delta'})$ where this does not hold, since the support of \widehat{P}_{n,η^*} is a subset of $\bar{B}(0, R_n)$, $W_p(G, \widehat{P}_{n,\eta^*}) \leq 2R_n$.

Therefore, taking $\delta' < \delta$ where δ is as defined in the statement of the Theorem, there exists $C > 0$ such that for $n \geq n_0$,

$$\mathbb{E}_{(G^*\mathbb{Q})^{\otimes n}} [W_p(G, \widehat{P}_{n,\eta^*})] \leq Cm_n^{-1},$$

which concludes the proof.

E.2 Proof of Theorem 7

Let \widehat{P}_n be an estimator of G . According to [27],

$$\begin{aligned} & \sup_{\substack{G \in \text{St}_{\mathcal{K}}(a,d,r_0) \cap \mathcal{L}(1,S,\mathcal{H}_1^*) \\ \mathbb{Q} \in \mathcal{Q}^{(D)}(\nu,c(\nu),E)}} \mathbb{E}_{(G^*\mathbb{Q})^{\otimes n}} [W_p(G, \widehat{P}_n)] \\ & \geq \frac{1}{2} W_p(G_0(1), G_1(1)) (1 - \|G_0(1) * \mathbb{Q} - G_1(1) * \mathbb{Q}\|_1)^n. \end{aligned}$$

Using the same two distributions $G_0(1)$, $G_1(1)$ and the same set \mathcal{H}_1^* as in Theorem 5. We have shown in Theorem 5 that there exists a constant $C > 0$ such that

$$\|G_0(1) * \mathbb{Q} - G_1(1) * \mathbb{Q}\|_{TV} \leq \frac{C}{n},$$

taking γ of the form $c \log(n)^{-1-\delta}$ for any $\delta > 0$ and c small enough, which implies that the minimax risk is lower bounded by $W_p(G_0(1), G_1(1))$. We show that there exist constants $c > 0$ and $n_0 > 0$ such that for $n \geq n_0$

$$W_p(G_0(1), G_1(1)) \geq c\gamma.$$

Let \mathcal{U}_γ be the set of $u \in \mathbb{R}$ such that $|\cos(\frac{u}{\gamma})| \geq 1/2$, that is $\mathcal{U}_\gamma = \bigcup_{k \in \mathbb{Z}} [k\pi\gamma - \frac{\pi\gamma}{3}, k\pi\gamma + \frac{\pi\gamma}{3}]$. For each $k \in \mathbb{Z}$, let $I_{k,\gamma} := [k\pi\gamma - \frac{\pi\gamma}{2}, k\pi\gamma + \frac{\pi\gamma}{2}]$. Let us also define, for any two sets A and B of \mathbb{R}^d , $d(A, B) = \inf_{x \in A, y \in B} \|x - y\|_2$. We first show that

$$d(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}), M_1(\gamma)) \geq \gamma \left(\frac{1}{4\sqrt{2}} \wedge \frac{\pi}{6} \right).$$

Let $x \in M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1})$ and $y \in M_1(\gamma)$. There exists $k \in \mathbb{Z}$ such that $x \in M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1})$. If $y \in I_{k,\gamma} \times \mathbb{R}^{D-1}$ (that is, if the first coordinate of x and y are in the same interval $I_{k,\gamma}$), then

$$\|x - y\|_2 \geq d(M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1}), M_1(\gamma) \cap (I_{k,\gamma} \times \mathbb{R}^{D-1})).$$

All points of $M_0(\gamma)$ are of the form $(u, u + \frac{1}{2}\gamma \cos(\frac{u}{\gamma}), 0, \dots, 0)^\top$ and the distance between $(u, u + \frac{1}{2}\gamma \cos(\frac{u}{\gamma}), 0, \dots, 0)^\top$ and the diagonal defined by $\mathcal{D} := \{(u, u, 0, \dots, 0)^\top : u \in \mathbb{R}\}$ is $\frac{1}{4\sqrt{2}}\gamma |\cos(\frac{u}{\gamma})|$. Since the sets $M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1})$ and $M_1(\gamma) \cap (I_{k,\gamma} \times \mathbb{R}^{D-1})$ are on opposite sides of the diagonal \mathcal{D} ,

$$\begin{aligned} & d(M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1}), M_1(\gamma) \cap (I_{k,\gamma} \times \mathbb{R}^{D-1})) \\ & \geq d(M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1}), \mathcal{D}) \\ & = \frac{1}{4\sqrt{2}}\gamma, \end{aligned}$$

so that $\|x - y\|_2 \geq \frac{1}{4\sqrt{2}}\gamma$. If now $y \notin I_{k,\gamma} \times \mathbb{R}^{D-1}$,

$$\begin{aligned} & d(M_0(\gamma) \cap ((\mathcal{U}_\gamma \cap I_{k,\gamma}) \times \mathbb{R}^{D-1}), M_1(\gamma) \cap ((\mathbb{R} \setminus I_{k,\gamma}) \times \mathbb{R}^{D-1})) \geq d(\mathcal{U}_\gamma \cap I_{k,\gamma}, \mathbb{R} \setminus I_{k,\gamma}) \\ & = \frac{\pi\gamma}{6}, \end{aligned}$$

so that $\|x - y\|_2 \geq \frac{\pi\gamma}{6}$, and thus $d(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}), M_1(\gamma)) \geq \gamma(\frac{1}{4\sqrt{2}} \wedge \frac{\pi}{6})$.

Now, let us show that $W_p(G_0(1), G_1(1)) \geq \gamma(\frac{1}{8\sqrt{2}} \wedge \frac{\pi}{12})$. Let π be a transport plan between $G_0(1)$ and $G_1(1)$, then

$$\begin{aligned} & \int_{M_0(\gamma) \times M_1(\gamma)} \|x - y\|_2^p d\pi(x, y) \\ & \geq \int_{M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}) \times M_1(\gamma)} \|x - y\|_2^p d\pi(x, y) \\ & \geq d(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}), M_1(\gamma))^p \pi(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}) \times M_1(\gamma)) \\ & = d(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}), M_1(\gamma))^p G_0(1)(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1})) \\ & = d(M_0(\gamma) \cap (\mathcal{U}_\gamma \times \mathbb{R}^{D-1}), M_1(\gamma))^p \mathbb{P}[U(1) \in \mathcal{U}_\gamma] \end{aligned}$$

since $G_1(1)$ has support $M_1(\gamma)$ and by definition of $G_0(1)$. Therefore, by taking the infimum on all transport plans between $G_0(1)$ and $G_1(1)$,

$$W_p(G_0(1), G_1(1)) \geq \gamma(\frac{1}{4\sqrt{2}} \wedge \frac{\pi}{6}) \mathbb{P}[U(1) \in \mathcal{U}_\gamma]^{1/p}.$$

$U(1)$ admits a density f_1 with respect to Lebesgue measure that is supported on $[-1, 1]$ and continuous. Let us write ω one of its modulus of continuity. We have

$$\begin{aligned} \mathbb{P}[U(1) \in \mathcal{U}_\gamma] &= \int_{[-1, 1]} f_1(x) 1_{|\mathcal{U}_\gamma}(x) dx \\ &= \sum_{k \in [-1/(\pi\gamma), 1/(\pi\gamma)]} \int_{[k\pi\gamma - \frac{\pi\gamma}{3}, k\pi\gamma + \frac{\pi\gamma}{3}]} f_1(x) dx \\ &\leq \sum_{k \in [-1/(\pi\gamma), 1/(\pi\gamma)]} \left(\int_{[k\pi\gamma - \frac{\pi\gamma}{3}, k\pi\gamma + \frac{\pi\gamma}{3}]} f_1(k\pi\gamma) dx + \frac{2\pi\gamma}{3} \omega(\pi\gamma/3) \right) \\ &\leq \sum_{k \in [-1/(\pi\gamma), 1/(\pi\gamma)]} \frac{2}{3} \int_{[k\pi\gamma - \frac{\pi\gamma}{2}, k\pi\gamma + \frac{\pi\gamma}{2}]} f_1(k\pi\gamma) dx + \frac{3}{\pi\gamma} \frac{2\pi\gamma}{3} \omega(\pi\gamma/3) \\ &\leq \sum_{k \in [-1/(\pi\gamma), 1/(\pi\gamma)]} \frac{2}{3} \int_{[k\pi\gamma - \frac{\pi\gamma}{2}, k\pi\gamma + \frac{\pi\gamma}{2}]} f_1(x) dx \\ &\quad + \frac{3}{\pi\gamma} \left(\frac{2\pi\gamma}{3} \omega(\pi\gamma/3) + \pi\gamma \omega(\pi\gamma/2) \right) \\ &\leq \frac{2}{3} \int_{[-1, 1]} f_1(x) dx + 3 \left(\frac{2}{3} \omega(\pi\gamma/3) + \omega(\pi\gamma/2) \right) \\ &\xrightarrow{\gamma \rightarrow 0} \frac{2}{3} \int_{[-1, 1]} f_1(x) dx = \frac{2}{3}. \end{aligned}$$

Therefore, there exists n_0 such that for all $n \geq n_0$, $W_p(G_0(1), G_1(1)) \geq \gamma(\frac{1}{8\sqrt{2}} \wedge \frac{\pi}{12})$.

References

- [1] Eddie Aamari, Catherine Aaron, and Clément Levrard. Minimax boundary estimation and estimation with boundary. *Bernoulli*, 29(4):3334–3368, 2023.
- [2] Eddie Aamari and Clément Levrard. Nonasymptotic rates for manifold, tangent space and curvature estimation. *Ann. Statist.*, 47(1):177–204, 2019.
- [3] Yariv Aizenbud and Barak Sober. Non-parametric estimation of manifolds from noisy data, 2021.

- [4] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [5] Clément Berenfeld, Paul Rosa, and Judith Rousseau. Estimating a density near an unknown manifold: a bayesian nonparametric approach. *The Annals of Statistics*, 52(5):2081–2111, 2024.
- [6] Jean-Daniel Boissonnat, Frédéric Chazal, and Mariette Yvinec. *Geometric and topological inference*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2018.
- [7] Victor-Emmanuel Brunel, Jason M. Klusowski, and Dana Yang. Estimation of convex supports from noisy measurements. *Bernoulli*, 27(2):772–793, 2021.
- [8] Jérémie Capitaio-Miniconi and Élisabeth Gassiat. Deconvolution of spherical data corrupted with unknown noise. *Electron. J. Stat.*, 17(1):607–649, 2023.
- [9] Antonio Cuevas and Ricardo Fraiman. A plug-in approach to support estimation. *Ann. Statist.*, 25(6):2300–2312, 1997.
- [10] Jérôme Dedecker, Aurélie Fischer, and Bertrand Michel. Improved rates for Wasserstein deconvolution with ordinary smooth error in dimension one. *Electron. J. Stat.*, 9(1):234–265, 2015.
- [11] Jérôme Dedecker and Bertrand Michel. Minimax rates of convergence for Wasserstein deconvolution with supersmooth errors in any dimension. *J. Multivariate Anal.*, 122:278–291, 2013.
- [12] Sylvain Delattre and Aurélie Fischer. Estimation via length-constrained generalized empirical principal curves under small noise, 2019.
- [13] Luc Devroye and Gary L. Wise. Detection of abnormal behavior via nonparametric estimation of the support. *SIAM J. Appl. Math.*, 38(3):480–488, 1980.
- [14] Vincent Divol. Measure estimation on manifolds: an optimal transport approach. *Probab. Theory Related Fields*, 183(1-2):581–647, 2022.
- [15] Herbert Federer. Curvature measures. *Trans. Amer. Math. Soc.*, 93:418–491, 1959.
- [16] Nicolás García Trillos, Daniel Sanz-Alonso, and Ruiyi Yang. Local regularization of noisy point clouds: improved global geometric estimates and data analysis. *J. Mach. Learn. Res.*, 20:Paper No. 136, 37, 2019.
- [17] Élisabeth Gassiat, Sylvain Le Corff, and Luc Lehericy. Deconvolution with unknown noise distribution is possible for multivariate signals. *Ann. Statist.*, 50(1):303–323, 2022.
- [18] Élisabeth Gassiat, Sylvain Le Corff, and Luc Lehericy. Supplementary to “deconvolution with unknown noise distribution is possible for multivariate signals”. *Ann. Statist.*, 2022.
- [19] Élisabeth Gassiat, Luc Lehericy, and Sylvain Le Corff. Erratum: Deconvolution with unknown noise distribution is possible for multivariate signals. preprint, 2025.
- [20] Christopher R. Genovese, Marco Perone-Pacífico, Isabella Verdinelli, and Larry Wasserman. Manifold estimation and singular deconvolution under Hausdorff loss. *Ann. Statist.*, 40(2):941–963, 2012.
- [21] Alexander Goldenshluger and Oleg Lepski. Universal pointwise selection rule in multivariate function estimation. *Bernoulli*, 14(4):1150–1190, 2008.
- [22] Alexander Meister. Estimating the support of multivariate densities under measurement error. *J. Multivariate Anal.*, 97(8):1702–1717, 2006.

- [23] G. W. Morgan. A Note on Fourier Transforms. *J. London Math. Soc.*, 9(3):187–192, 1934.
- [24] Jonathan Niles-Weed and Quentin Berthet. Minimax estimation of smooth densities in Wasserstein distance. *Ann. Statist.*, 50(3):1519–1540, 2022.
- [25] Nikita Puchkin and Vladimir Spokoiny. Structure-adaptive manifold estimation. *J. Mach. Learn. Res.*, 23:Paper No. [40], 62, 2022.
- [26] T. Tlas. Bump functions with monotone Fourier transforms satisfying decay bounds. *J. Approx. Theory*, 278:Paper No. 105742, 6, 2022.
- [27] Bin Yu. Assouad, Fano, and Le Cam. In *Festschrift for Lucien Le Cam*, pages 423–435. Springer, New York, 1997.