



HAL
open science

Are grey sets novel? Grey sets are nothing but interval-valued fuzzy sets

Didier Dubois, Henri Prade

► **To cite this version:**

Didier Dubois, Henri Prade. Are grey sets novel? Grey sets are nothing but interval-valued fuzzy sets. Bulletin pour les sous-ensembles flous et leurs applications, 1990, 44, pp.27-30. hal-04074890

HAL Id: hal-04074890

<https://hal.science/hal-04074890>

Submitted on 19 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Are grey sets novel ?

Grey sets are nothing but interval-valued fuzzy sets

Didier DUBOIS – Henri PRADE

*Institut de Recherche en Informatique de Toulouse (I.R.I.T.)
Université Paul Sabatier, 118 route de Narbonne
31062 TOULOUSE Cedex – FRANCE*

There has been a considerable amount of literature produced by Chinese researchers about so-called grey sets, grey numbers, grey systems since the mid-eighties, following an idea advocated by Prof. Deng Julong [3]. Some of these works were published in BUSEFAL (see issues n° 36, 38, 39, 40, 41, 42, 44) and there exists a publication specially devoted to this topic [2]. Usually they are presented as a fairly new, powerful tool, making reference almost only to the grey set and system literature. Especially the link with fuzzy sets is not always really acknowledged, even in a recently published paper devoted to the relationship between grey numbers and fuzzy numbers (and where fuzzy numbers are rather presented as a particular case) [22]. The aim of this short note is to clarify this point.

A grey set G on a referential X is defined by two functions $\underline{\mu}_G$ and $\bar{\mu}_G$ from X to $[0,1]$ such that $\underline{\mu}_G \leq \bar{\mu}_G$, called "lower and upper subordinate functions" [21][22]. Clearly to each $x \in X$ is attached an interval $[\underline{\mu}_G(x), \bar{\mu}_G(x)] \subseteq [0,1]$. $\underline{\mu}_G(x)$ and $\bar{\mu}_G(x)$ are called "the lower and upper subordinate degrees of x relative to G ". This interval may be understood as containing (the) possible values of the degree of membership of x to G . We have recognized a Φ -fuzzy set, first considered by Sambuc [19], i.e. an interval-valued fuzzy set. This is a well-known particular case of type 2-fuzzy set [12], a notion first introduced by Zadeh [23] (a type 2-fuzzy set is a set whose membership degrees are fuzzy sets of $[0,1]$, or more generally of a set L of valuations). See also [5] pp. 30-31 and pp. 63-66.

Φ -fuzzy sets can be encountered in the fuzzy set literature with slightly different interpretations. Let us especially mention

- a sub-specified fuzzy set, whose membership degrees, at least for some elements, are ill-known. This sub-specification may result of the appreciation of experts or of the mathematical constraint(s) defining the fuzzy set ;
- a "flou" set \mathfrak{G} [10] which is a pair of two ordinary subsets (\underline{G}, \bar{G}) of X such that $\underline{G} \subseteq \bar{G}$ where \underline{G} is understood as a set of typical elements while $\bar{G} - \underline{G}$ is interpreted as a set of elements considered as peripheral for \mathfrak{G} ;
- a sub-definite set \mathfrak{G} [14] which is a 4-tuple $(G_+, G_-, \bar{m}, \underline{m})$ where G^+ (resp. G^-) is the set of elements which are known to belong (resp. not to belong) to \mathfrak{G} and $\bar{m} \leq |X - G_-|$ and $\underline{m} \geq |G_+|$ are respectively an upper and a lower bound of the number of elements which belong to \mathfrak{G} ;
- a twofold fuzzy set G [6][7], is defined by a pair $(\underline{\mu}_G, \bar{\mu}_G)$ such that $\{x \in X, \underline{\mu}_G(x) > 0\} \subseteq \{x \in X, \bar{\mu}_G(x) = 1\}$, where $\underline{\mu}_G$ (resp. $\bar{\mu}_G$) is the membership function of the set of elements which belong more or less certainly (resp. possibly) to G (in the sense of possibility theory [24]) ;
- an intuitionistic fuzzy set [1] defined by a pair (μ_G, ϑ_G) where $\mu_G(x)$ is a degree of membership and $\vartheta_G(x)$ is a degree of non-membership for x , both degrees belonging to $[0,1]$, with the constraint $\forall x, \mu_G(x) + \vartheta_G(x) \leq 1$, or equivalently $\mu_G(x) \leq 1 - \vartheta_G(x)$, thus the pair $(\mu_G, 1 - \vartheta_G)$ is equivalent to a pair of lower and upper membership functions $(\underline{\mu}_G, \bar{\mu}_G)$;
- a rough set [16][17] and more generally a rough fuzzy set [8] is a (fuzzy) set G defined on X whose membership function on a *coarsened* referential X/R (where X/R is a quotient space which represents a partition of X induced by an equivalence relation) is bounded from below and from above respectively by

$$\forall t \in X/R, \underline{\mu}_G(t) = \inf\{\mu_G(x), x \in t\}$$

$$\bar{\mu}_G(t) = \sup\{\mu_G(x), x \in t\}$$

More generally, a fuzzy rough (fuzzy) set [8][13] is defined when considering a similarity relation or a fuzzy partition instead of R or X/R respectively.

Particular cases of grey sets are ordinary sets and fuzzy sets obtained when $\underline{\mu}_G = \bar{\mu}_G$, as well as the so-called Deng grey sets [22] where $\bar{\mu}_G$ is the characteristic function of an ordinary subset S of X and $\underline{\mu}_G = 0$ everywhere. It may be interpreted as follows, values outside S do not belong to the grey set definitely while values in S *may* belong to it. Another kind of grey set

which has a particular meaning from a possibility distribution specification point of view ([25][20][9]), obeys to the following constraints, where S is an ordinary subset of X

$$\begin{aligned} \forall x \notin S, \underline{\mu}_G(x) = 0 \quad \text{and} \quad \bar{\mu}_G(x) = 1 \\ \forall x \in S, \bar{\mu}_G(x) = 1 \end{aligned}$$

while $\underline{\mu}_G$ remains unconstrained over S . In terms of possibility distribution it means that the values x in S are *at least* possible at the degree $\underline{\mu}_G(x)$ and that nothing is known outside of S . Dually, a grey set such that $\forall x \notin S, \underline{\mu}_G(x) = 0$ and $\bar{\mu}_G(x) = 1$ and $\forall x \in S, \underline{\mu}_G(x) = 0$ corresponds to the knowledge that the value $x \in S$ is *at most* possible to the degree $\bar{\mu}_G(x)$.

Type 2-fuzzy sets have been studied from a theoretical point of view (e.g. [11]), arithmetic operations on type 2-fuzzy numbers have been defined and studied [4][5], as well as type 2-fuzzy relations [27]. Interval-valued fuzzy sets have been advocated by Zadeh himself under the name of "ultra fuzzy set" (e.g. [26]). Let us also mention the use of interval-valued fuzzy sets which has been proposed in decision analysis and in economical studies long time ago (e.g. [15][18]), among many other works. So it would be a pity that researchers in grey systems ignore results available in the fuzzy set literature. Certainly interval-valued fuzzy sets are of a considerable interest in general, however a clear interpretation of grey sets and a clarified relationship with other existing proposals are necessary from a scientific point of view.

References

- [1] Atanassov K.T. Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 20, 1986, 87–96.
- [2] Deng J.L. (Editor-in-Chief) *The Journal of Grey System*, Vol. 1, 1989, Sci-Tech Information Services & China Petroleum Industry Press.
- [3] Deng J.L. et al. *Grey System*. China Ocean Press, Beijing, 1988.
- [4] Dubois D., Prade H. Fuzzy real algebra : some results. *Fuzzy Sets and Systems*, 2, 1979, 327–348.
- [5] Dubois D., Prade H. *Fuzzy Sets and Systems – Theory and Applications*. Academic Press, New York, 1980.
- [6] Dubois D., Prade H. Twofold fuzzy sets : an approach to the representation of sets with fuzzy boundaries based on possibility and necessity measures. *J. Fuzzy Math. (China)*, 3(4), 1983, 53–76.
- [7] Dubois D., Prade H. Twofold fuzzy sets and rough sets – Some issues in knowledge representation. *Fuzzy Sets and Systems*, 23, 1987, 3–18.
- [8] Dubois D., Prade H. Rough fuzzy sets and fuzzy rough sets. *Int. J. of General Systems*, 17, 1990, 191–209.
- [9] Dubois D., Prade H. Fuzzy rules in knowledge-based systems – Modelling gradedness,

- uncertainty and preference. In : An Introduction to Fuzzy Logic Applications in Intelligent Systems (R.R. Yager, L.A. Zadeh, eds.), Kluwer Acad. Publishers, to appear.
- [10] Gentilhomme Y. Les ensembles flous en linguistique. Cahiers de Linguistique Théorique et Appliquée (Bucharest), 5, 1968, 47–63.
- [11] Gottwald S. Set theory for fuzzy sets of higher level. Fuzzy Sets and Systems, 2, 1979, 125–151.
- [12] Mizumoto M., Tanaka K. Some properties of fuzzy sets of type 2. Information and Control, 31, 1976, 312–340.
- [13] Nakamura A. Fuzzy rough sets. Note on Multiple-Valued Logic in Japan, 9(8), 1988, 1–8.
- [14] Narin'yani A.S. Sub-definite set – New data-type for knowledge representation. (In Russian) Memo n° 4-232, Computing Center, Novosibirsk, 1980.
- [15] Nojiri H. Ultrafuzzy team decisions in a dynamic environment. Information Sciences, 23, 1981, 105–122.
- [16] Pawlak Z. Rough sets. Int. J. Comput. and Information Sciences, 11(5), 1982, 341–356.
- [17] Pawlak Z. Rough classification. Int. J. of Man-Machine Studies, 20, 1984, 469–485.
- [18] Ponsard C. Hiérarchie des places centrales et graphes Φ -flous. Environ. Plann., A9, 1977, 1233–1252.
- [19] Sambuc R. Fonctions Φ -floues – Application à l'aide au diagnostic en pathologie thyroïdienne. Thèse de Doctorat en Médecine, Marseille, France, 1975.
- [20] Sanchez E. On possibility-qualification in natural languages. Information Sciences, 15, 1978, 45–76.
- [21] Wang Q.Y., Liu K.D., Wu H.Q. The conception of grey number and its property. Proc. of the 7th Annual Meeting of the North American Fuzzy Information Processing Society (NAFIPS'88), San Francisco, June 8-10, 1988, 259–262.
- [22] Wang Q.Y., Wu H.Q., Xue J.S., Lin K.D. On grey number : basic element of grey mathematics and its relations with fuzzy and real numbers. The Journal of Grey System, 1, 1988, 69–77.
- [23] Zadeh L.A. Quantitative fuzzy semantics. Information Sciences, 3, 159-176.
- [24] Zadeh L.A. Fuzzy sets as a basis for a theory of possibility. Fuzzy Sets and Systems, 1(1), 1978, 3–28.
- [25] Zadeh L.A. PRUF – a meaning representation language for natural languages. Int. J. of Man-Machine Studies, 10, 1978, 395–460.
- [26] Zadeh L.A. Fuzzy sets, usability and commonsense reasoning. In : Matters of Intelligence (L.M. Vaina, ed.), D. Reidel, Dordrecht, 1987, 289–309.
- [27] Zhang W.R., Bezdek J.C., Pettus R.O. Linguistic similarity relations. Proc. of the Inter. Symp. on Methodologies for Intelligent Systems (M. Zemankova, M.L. Emrich, eds.), Knoxville, TN., Oct. 25, 1986, 149-169.