



HAL
open science

Approaches to determine pesticides in marine bivalves

Thierno Diallo, Julia Leleu, Julien Parinet, Thierry Guérin, Hélène Thomas,
Adélaïde Lerebours

► **To cite this version:**

Thierno Diallo, Julia Leleu, Julien Parinet, Thierry Guérin, Hélène Thomas, et al.. Approaches to determine pesticides in marine bivalves. *Analytical and Bioanalytical Chemistry*, 2023, 415 (16), pp.3093-3110. 10.1007/s00216-023-04709-4 . hal-04074997

HAL Id: hal-04074997

<https://hal.science/hal-04074997v1>

Submitted on 19 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

1 Approaches to determine pesticides in marine bivalves

2

3 Thierno Diallo^{1,2§}, Julia Leleu^{1§}, Julien Parinet¹, Thierry Guérin³, Hélène Thomas², Adélaïde
4 Lerebours^{2*}

5

6

7 ¹ ANSES, Laboratory for Food Safety, F-94701 Maisons-Alfort, France

8 ² Littoral Environnement et Sociétés (LIENSs), UMR 7266, CNRS-Université de La Rochelle,
9 2 rue Olympe de Gouges, F-17042 La Rochelle Cedex 01, France.

10 ³ ANSES, Strategy and Programmes Department, F-94701 Maisons-Alfort, France

11

12

13 §: co-authors

14 *: **Corresponding author:**

15 Adélaïde Lerebours, phone number: +33 (0)5 46 50 67 83, email address:
16 adelaiderebours@univ-lr.fr

17

18

19

20

21

22

23 **Keywords:** pesticide determination, marine bivalves, recent developments, analytical
24 methods and techniques

25 **Abstract**

26 Due to agricultural runoff, pesticides end up in aquatic ecosystems and some accumulate in
27 marine bivalves. As filter feeders, bivalves can accumulate high concentration of chemicals in
28 their tissue representing a potential risk for the health of human and aquatic ecosystems. So
29 far, most of the studies dealing with pesticides contamination in marine bivalves, for example,
30 in the French Atlantic and English Channel coasts, have focused on old generation of
31 pesticides. Only a few investigated the new emerging pesticides partly due to methodological
32 challenges. A better understanding of the most sensitive and reliable methods is thus essential
33 for accurately determining a wide variety of environmentally relevant pesticides in marine
34 bivalves. The review highlighted the use of more environmentally friendly and efficient
35 materials such as sorbents and the “Quick Easy Cheap Effective Rugged Safe” extraction
36 procedure to extract pesticides from bivalve matrices, as they appeared to be the most
37 efficient while being the safest. Moreover, this method combined with high-resolution mass
38 spectrometry (MS) technique offers promising perspectives by highlighting a wide range of
39 pesticides including those that are not usually sought. Finally, recent developments in the field
40 of ultrahigh-performance liquid chromatography coupled to MS, such as two-dimensional
41 chromatography and ion mobility spectrometry will improve the analysis of pesticides in
42 complex matrices.

43 **Introduction**

44 The contamination of the aquatic environment by pesticides may have deleterious effects on
45 species of ecological and economic importance such as mussels and oysters [1–4]. Those
46 organisms filter large volumes of water (30 to 40 ml/min) [5] and can thus efficiently
47 bioaccumulate various chemicals including pesticides released by agricultural runoff in their
48 digestive gland [6–9].

49 The agricultural production makes intensive use of pesticides which comprises a wide
50 variety of molecules to protect crops against weeds (herbicides), fungi (fungicides), or insects
51 (insecticides) [10]. Those pesticides are divided into classes based on their chemical
52 properties: organochlorines (OCs), organophosphates (OPs), carbamates (Carb), pyrethroids
53 (Pyr) and neonicotinoids (Neo) [11]. Organochlorines, organophosphates and carbamates are
54 of the older generations, and pyrethroids and neonicotinoids of the newer [11–13]. Most of
55 the studies led in marine bivalves from coastal areas of France (Table S1, Text S1) and other
56 countries [14,15] mainly focused on old generation of pesticides (OCs, OPs and Carb), and
57 thus, missed the newer generation of pesticides.

58 OCs are the most frequent pesticides detected in bivalves from the French Atlantic
59 coast (Table 2), although most of them have been restricted or banned [16]. Indeed, OCs were
60 detected in bivalves from all sampling sites, highlighting a persistent contamination along the
61 French Atlantic coast (Figure 1, Table 1) [1,8,17]. However, some of them were detected at
62 very low concentration and tend to decline over the years. For instance, aldrin concentrations
63 decreased from 2.71 ng/g dry weight (dw) in the scallops (*Mimachlamys varia*) in March
64 2014 [1] to values < LOQ of 1.67 ng/g in oysters (*C. gigas*) or < LOD of 0.5 ng/g in mussels
65 (*M. edulis*) from the Pertuis Charentais littoral area, in February 2019 [18] (Table 2). This
66 decrease over the years was expected as after being widely used in the 50s–60s, this
67 insecticide was banned in France, in 1998.

68 The highest level of OC pesticides found in bivalves from the French Atlantic coast
69 was measured for metolachlor in mussels with concentrations that ranged from 349 to 398
70 ng/g dw [17]. Those concentrations seemed particularly high since this herbicide of low half-
71 life was prohibited in France in 2003. However, after its prohibition, metolachlor was
72 replaced by the active enantiomer S-metolachlor, which is actually the molecule measured by
73 classical analytical approaches (LC or GCMS). More recently, a study found lower
74 metolachlor concentrations in oysters and mussels from the Pertuis Charentais. They were
75 below 0.33 (LOQ) or up to 0.38 ng/g wet weight (ww) in Pacific oysters, and varied between
76 0.5 (LOD) and 1.67 ng/g ww (LOQ) in blue mussels [18] (Table 2). It would have been
77 interesting to know the concentration levels of metolachlor metabolites: ESA and OXA
78 metolachlor, to allow a better understanding of the fate of this pesticide in marine biota.
79 However, studies on the contamination of the marine environment by pesticide metabolites
80 are rare. Contamination of bivalves by the OC pesticide acetochlor, came second after
81 metolachlor with higher concentrations found in mussels from Chausey Island coast (125 ng/g
82 dw) than in mussels from the Seine estuary (53 ng/g dw) [17] or from the Pertuis seas (< LOD
83 of 0.5 ng/g) [18]. The concentrations of endosulfan beta and metazachlor in mussels from the
84 Seine estuary were lower and reached 15.2 and 24 ng/g dw respectively [17]. A few years
85 later, the endosulfan beta and metazachlor concentrations in mussels from the Pertuis seas,
86 were even lower, < LOD of 1 and 0.5 ng/g ww, respectively [18] (Table 2). Therefore,
87 bivalves from the Chausey archipelago were more contaminated with OCs pesticides than
88 those from the Seine estuary and the Pertuis seas. Other OCs, such as lindane (banned in
89 1998) and DDT (banned in 2010), which are highly persistent pollutants in the environment,
90 were monitored in previous studies. Lindane concentrations found in mussels and oysters
91 ranged from 0.2 to 2.2 ng/g dw in the Pertuis seas region [8,18] (Table 2). The concentrations
92 of 2,4- and 4,4-DDT are now very low, in the range of 0.4 - 0.9 ng/g dw [8,18] in bivalves

93 from the Gironde estuary and the Pertuis seas. However, significant accumulation of its
94 metabolites were found in oysters collected in 2008 in the same region, with concentrations
95 reaching 3.7 and 5.5 ng/g dw of 4,4'- and 2,4'- DDE, respectively [8].

96 Another class of widely studied pesticides are the OPs such as ethoprophos (31 ng/g)
97 and tributyl phosphate (14 ng/g) which were quantified in bivalves from the Vendée region
98 [19] (Table 2). Despite their significant accumulation in marine bivalves and their authorized
99 status, there is only one recent study available in the literature on those two molecules
100 regarding the French Atlantic coast. A recent study reported much lower concentrations of
101 parathion (between 0.56 and 0.69 ng/g) banned in 2001, chlorpyrifos (between 0.79 and 0.93
102 ng/g) banned in 2020, and chlorfenvinphos (between 1.65 and 2.12 ng/g) banned in 2007, in
103 the Pacific oysters sampled in the Pertuis Charentais littoral area [18] (Table 2). As banned
104 molecules found today at very low concentrations, those pesticides represent a minor concern
105 for marine bivalves and human health.

106 Interestingly, more recent pesticides such as bifenthrin (48 ng/g), cypermethrin (10.9
107 ng/g) and deltamethrin (11.2 ng/g) were quantified for the first time in bivalves from the
108 French Atlantic coast in 2015 by Menet-Nedelec et al. (Table 2) [17]. Other emerging
109 pesticides such as neonicotinoids could not be identified due to their low concentrations in
110 marine bivalves from the French Atlantic and English Channel coasts [17]. However, their
111 toxicity has been evidenced in bivalves exposed under laboratory conditions [20]. Altogether,
112 those results highlighted the presence of old generation of pesticides in marine biota and
113 significant concentrations of their metabolites, but data on the new generation of pesticides
114 such as pyrethroids and neonicotinoids are lacking. Those new molecules deserve to be
115 monitored further. Indeed, contrary to the old generation of pesticides, most of the new
116 generation of pesticides are currently authorized. Secondly, by considering only old
117 pesticides, studies underestimate the contamination by the environmentally realistic “cocktail”

118 of pesticides that includes the new generation of pesticides. Actually, quantifying those
119 molecules at environmental concentrations remains a methodological and technical challenge.
120 Therefore, this underlines the need to apply the most efficient procedures to accurately
121 determine the wide variety of environmentally relevant pesticides that accumulate in marine
122 bivalves.

123 The present review firstly compares the latest methods used for tissue preparation and
124 extraction, and to clean-up samples. Advices to overcome the matrix effect specific to bivalve
125 tissues are also provided. Finally, the review discusses the most relevant and promising
126 analytical techniques to use and highlights the good validation procedures to apply.

127 **Methodologies**

128 *Sample pretreatment*

129 In most studies, samples were homogenized, freeze-dried or lyophilized and stored at -20°C
130 before proceeding to the next steps (Table 1). Research focused on whole individuals or
131 specific tissues such as the digestive gland, a key tissue to assess chemical contaminant in
132 bivalves [7]. Then, due to low concentrations in bivalves, effective extraction procedures are
133 used to isolate and concentrate pesticides (Table 1).

134 *Extraction of pesticides from bivalve samples*

135 The Matrix Solid-Phase Dispersion (MSPD) refers to one of the older extraction procedures
136 that is less used nowadays, with 9.1% of the studies employing it since 2015 (Table 1),
137 although it is a suitable sample preparation technique for solid and semi-solid biological
138 samples. Carro *et al.*, (2005) was evaluated MSPD procedure using an asymmetrical
139 experimental design [21]. The authors tested and validated the optimal condition using
140 multispecies biological reference material from the International Atomic Energy Agency
141 (IAEA-406), with adequate precision and accuracy. The validated procedure was applied to
142 the analysis of several samples of aquaculture feed and marine species. MSPD consists in

143 mixing a sample with a solid support (usually a silica-based material), to obtain an
144 homogeneous mixture [22]. The resulting mixture is placed in a solid-phase extraction (SPE)
145 column or cartridge, followed by elution with suitable organic solvents (with different elution
146 powers and polarities) and collection of the analytes by fractionation. The advantages of this
147 method include the consumption of small amounts of sample and solvents, simple application,
148 low cost and simultaneous extraction and clean-up in one step [21,22]. However, the MSPD is
149 fairly labor intensive, requiring the sample to be ground up with the solid matrix and packed
150 into a column for extraction since automation cannot be done [22]. The Pressurized Liquid
151 Extraction (PLE), also known as the Accelerated Solvent Extraction (ASE), and the Selective
152 Pressurized Liquid Extraction (SPLE) were used to extract pesticides in 27.3% of the studies
153 (Tables 1 and 2) [8,23]. PLE is an instrument-based extraction procedure that extracts
154 analytes (including POPs, pesticides or pharmaceuticals) from solid and semi-solid
155 matrices at high temperature (40–200°C) and pressure (500–3000 psi) [24–26]. High
156 temperatures enhance the solubility of non-thermolabile analytes [27]. The advantages of PLE
157 are: less solvent used, time per sample reduced and processes fully automated. However, the
158 disadvantage is the presence of coextracted non-targeted interfering compounds - such as
159 proteins or lipids and the degradation of thermolabile analytes. Thus, the PLE
160 technique requires a clean-up before quantification [28]. Selective PLE (SPLE) (or enhanced
161 PLE, in-cell clean-up, in-situ clean-up) combines the extraction and clean-up
162 processes [27]. Solid-phase adsorbents can be layered in the PLE cell to avoid interference.
163 They produce clean extracts which, in most cases, are directly injected in the analytical
164 system. Thus, no post-extraction clean-up is required. Adsorbents - for in-cell clean-up PLE -
165 are commonly used to extract non-polar analytes, such as OCs in animal tissues. In this
166 regard, PLE utilises a layer of fat-removal adsorbents such as Florisil, alumina (basic, acidic
167 or neutral), silica gel and sulphuric acid-silica gel [29]. Another extraction procedure that has

168 gained considerable importance in the recent years because of its "greener" (it uses less
169 solvent) and more efficient extraction capacity is microwave-assisted extraction (MAE)
170 [30,31]. This approach simply involves placing the sample with the solvent in specialized
171 vessels and heating the solvent using microwave energy. MAE has many advantages, such as
172 reduced extraction time and solvent consumption, as well as the ability to analyse multiple
173 samples [32]. However, the need for special equipment, the low selectivity and the
174 unavoidable high temperature reaction are considered as disadvantages of MAE [30]. Since
175 2003, an alternative extraction procedure, Quick, Easy, Cheap, Effective, Rugged and Safe
176 (QuEChERS) was introduced by Anastassiades *et al.*, (2003) [33] but the first publication on
177 the extraction of pesticides from bivalves using the QuEChERS procedure appeared in 2016
178 [34]. In the QuEChERS procedure, the most commonly used solvent is acetonitrile. The use
179 of this solvent allows the extraction of a wide polarity (non-polar and medium polar) of
180 pesticides [24]. Indeed, acetonitrile leads to much less interferences from compounds that can
181 be co-extracted such as lipophilic compounds, lipids and pigments than other extraction
182 solvents such as acetone and ethyl acetate [33]. The solid sample is usually ground and
183 homogenized using an Ultra-Turrax homogenizer [35]. The $MgSO_4$ and $NaCl$ are the original
184 salts used for liquid-liquid phase separation by binding to water [33]. $MgSO_4$, as well as
185 Na_2SO_4 , has a good water absorption and allows the release of acetonitrile (the most
186 commonly used extraction solvent), while $NaCl$ affects the ionic force of the extraction
187 solvents and, therefore, increases the selectivity of the extraction [24]. To avoid ionization or
188 degradation of pesticides due to pH changes during extraction, two modifications of the
189 original QuEChERS procedure were developed: (1) AOAC Method (acetate buffer) [36] and
190 (2) EN Method 15662 (citrate buffer) [37]. Both methods stabilize the pH at 5–5.5, which
191 facilitates the extraction of pesticides sensitive to low pH or those with stability problems.
192 This procedure is currently replacing those traditional time-consuming extractions, requiring

193 larger amount of toxic reagents, organic solvents and samples. Because the QuEChERS
194 method allows a fast, reliable extraction and uses less toxic solvents, it is a better method than
195 MAE, PLE or MSPD. However, the number of studies using the QuEChERS procedure in
196 bivalves is still low: 18 against 353 studies investigating other matrices in 2020. In the
197 principles of green analytical chemistry, other extraction techniques such as magnetic solid
198 phase extraction (MSPE) and stir bar sorptive extraction (SBSE) should be considered. MSPE
199 is based on the addition of magnetic nanoparticles such as C18 functionalized to the sample
200 solution for preconcentration of moderate and non-polar pesticides [38]. In addition to being
201 environmentally friendly, MSPE is a technique that has the advantage of using a limited
202 number of steps (limiting cross-contaminations), a small amount of sorbent, as well as the
203 reuse of sorbent [39]. However, MSPE procedure requires vortex, shaker, magnetic stirrer and
204 selection of suitable sorbents [39]. In SBSE, a stir bar coated with polydimethylsiloxane
205 (PDMS) phase is added to a vial containing the sample. SBSE consists of two steps:
206 extraction and desorption. An aqueous sample is extracted by stirring for a predetermined
207 time with a PDMS-coated stir-bar. The stir-bar is subsequently removed from the aqueous
208 sample and the absorbed compounds are then thermally desorbed and analysed by GCMS
209 [40]. The advantage of the SBSE is its simplicity, robustness, cost effectiveness and
210 environmental friendliness [39]. However, its important limitation is only commercial
211 availability of PDMS coating material exists at present. Besides, due to the non-polar
212 character of PDMS, SBSE technique is mainly applied to extract non-polar and weakly polar
213 compounds (logarithm of octanol-water partitioning coefficient lower than 3) but failed in the
214 extraction of strongly polar compounds [41]. In addition, SBSE procedure can only be applied
215 to medium-high volatility and medium-high thermo-stability analytes if a thermal desorption
216 has to be employed [42].

218 Bivalves contain lipids (between 2 and 10%) and proteins (between 10 and 18%) that can
219 affect the analysis of target compounds, especially at low concentrations [35]. Matrix effect is
220 caused by interactions (via van der Waals, dipolar-dipolar or electrostatic forces) between
221 pesticides and co-extractives that could suppress or enhance the ionisation of the analytes
222 [43]. This may result in a higher or lower signal, which affects the accuracy of the
223 quantitative results. Thus, this matrix effect needs to be reduced to avoid bias in the overall
224 analysis. To overcome this challenge, several interesting approaches have been used and
225 deserved attention to minimise the matrix components such as using internal standards,
226 improving the chromatography separation (reducing coelution with matrix components),
227 making serial dilution, or performing extensive sample clean-up [44,45], as follows. For
228 clean-up of sample extracts, the SPE method, previously mentioned for extracting pesticides,
229 can be used [46]. In SPE method, the sorbents are chosen according to their properties
230 (affinity, capacity and selectivity) towards the analytes of interest and matrix interferences [46].
231 The most popular clean-up strategy to avoid interference involves the SPE with a reverse-
232 phase octadecylsilane (C18) for isolation and preconcentration of nonpolar compounds [47]
233 whereas silica-SPE retains polar compounds in the sample [8,48,49]. The primary secondary
234 amine (PSA) is used to remove interferences such as fatty acids, organic acids, pigments,
235 sugars and metal ions [34,50]. This SPE method offers the best compromise between time,
236 difficulty and performances. Other clean-up procedures are less frequently used and are more
237 complicated to set up, such as those using adsorption columns [8,51] and gel permeation
238 chromatography [48]. The dispersive SPE (dSPE) and cartridge SPE (cSPE) methods can also
239 be used for the clean-up. They are much more rapid, easy to set up and efficient. dSPE is
240 based on the dispersion of a sorbent in the liquid samples [44,52]. After centrifugation, the
241 sorbent separates the analytes from the matrix, eliminating interfering compounds and

242 reducing wastewater [53,54]. dSPE provides faster, simpler and more adaptable clean-up than
243 traditional techniques and has been widely applied [45,47,50,53,55], however the
244 centrifugation step may lengthen the procedure. Moreover, during sorbent adsorption or
245 concentration, pesticides might be lost. In the cSPE procedure, the analytes are eluted from
246 the cartridge and interfering compounds are eluted before the analytes or remained on the
247 cartridge [54]. Compared to dSPE, the cSPE is time-consuming, adding experimental steps
248 such as cartridge conditioning, sample loading, interference elution and analyte elution [54].
249 Therefore, the dSPE associated to the QuEChERS method appeared to be the most reliable
250 and efficient, although cSPE is automatable and remains the most used procedure in the
251 routine laboratory. Indeed, this combination presents the best compromise between cost, time,
252 and easiness to use, especially if looking after a large set of pesticides potentially present in
253 marine bivalves. As with the SPE method, the most commonly used dSPE sorbents for extract
254 clean-up in bivalves are C18 (especially for fat removal) and PSA, often in the presence of
255 $MgSO_4$. The latter is commonly used as a drying salt for acetonitrile extraction as it
256 effectively removes residual water before injection into chromatographic systems. The
257 zirconia based sorbent, Z-Sep is based on zirconium oxide and Z-Sep/C18 is composed of
258 zirconium oxide and C18. This mixture reduces fat and pigment levels in matrices and it is
259 recommended for matrices containing more than 18% of lipid [56]. In 2015, Agilent
260 Technology introduced for the first time Enhanced Matrix Removal of lipids (EMR-lipid).
261 Unlike other d-SPE sorbents, EMR-lipid is a novel sorbent designed specifically for lipid
262 removal. The composition of EMR-Lipid is a proprietary secret, and it does not function as a
263 conventional sorbent, but it dissolves to saturation the sample extract solution, and it selects
264 compounds according to their size and hydrophobic properties [57]. More recently, novel
265 materials based on the principle of green chemistry are being used as sorbents. Despite their
266 effectiveness, these sorbents are not always commercially available, so they have to be

267 synthesized in the laboratory. These include molecularly imprinted polymers (MIPs), metal-
268 organic frameworks (MOFs) and carbon-based sorbents. The principle of MIPs is based on
269 the affinity of a polymer with specific cavities for chosen template molecules. Therefore,
270 MIPs are highly selective for those target molecules or structural analogs. The target
271 molecules are recognized via hydrophobic interactions, hydrogen bonding, van der Waals
272 forces, or electrostatic interactions [58]. The MIPs sorbent was also used for pigment removal
273 [59]. This sorbent, in combination with PSA, demonstrated their effectiveness in clean-up
274 bivalve extracts, with matrix effects within the range of $\pm 20\%$ for 100% of the herbicides
275 analysed [59]. The MOFs as sorbents are based on three-dimensional structures of
276 coordination networks consisting of a metal ion bound to organic ligands [60]. The main
277 advantage of MOFs is their large specific surface area, enormous porosity and the possibility
278 to easily adjust their pore size and shape from microporous to mesoporous scale, which can
279 lead to high extraction efficiency [61]. In addition, the high capacity to functionalize the pores
280 with functional groups such as $-\text{NH}_2$, $-\text{CO}$, $-\text{CHO}$, $-\text{COOH}$ and $-\text{OH}$, make them good
281 candidates for adsorption of interferents through hydrophilic or hydrophobic interactions [61].
282 However, the main limitations of MOFs is their low selectivity and stability in water [62].
283 Moreover, the limited availability of MOF-based adsorbents prevents the widespread
284 application of MOFs in the analysis and preconcentration of pesticides. The main reason is
285 that only a few chemical companies (e.g. the German chemical company BASF) have started
286 to produce MOFs on a commercial scale [63]. Furthermore, not all research laboratories have
287 the capacity to produce MOFs. Finally, carbon-based materials (e.g., activated carbon, carbon
288 nanotube and graphene) are used as sorbents due to their large surface area, high porosity and
289 surface reactivity, and they can be easily functionalized [64,65].

290 **Matrix effects and challenge to overcome them**

291 Although matrix effects are not always considered among the performance parameters for
292 method validation, they must be considered as they can influence the experimental results.
293 Despite the clean-up of extracts and LC or GC separation step, it is impossible to eliminate all
294 matrix effects. Matrix effects generally lead to signal suppression in LC-MS/MS applications.
295 This is the case in the study by Álvarez-Muñoz *et al.*, (2019) where the authors noted ion
296 suppression after extraction, purification and UHPLC-HRMS analysis of 16, 42 and 51% for
297 mussels, oysters and cockles, respectively [66]. In this work, matrix matched calibration
298 together with isotopically labelled internal standards were performed to minimise interference
299 and to avoid underestimation during quantification of targeted compounds. In contrast to LC-
300 MS/MS, matrix effects in GC-MS/MS mainly result in signal enhancement. Using GC-
301 MS/MS, Yu *et al.* achieved less than 20% of matrix effects for the 14 herbicides quantified in
302 mussels, scallops, and cockles after purification of extracts using PSA and MIPs as dSPE
303 sorbents [59]. In the study performed by Cruzeiro *et al.*, (2016), an average signal suppression
304 of 89% was obtained for 80% of the studied pesticides [34]. To overcome matrix effects, the
305 authors evaluated the addition of two analyte protectors such as 3-ethoxy-1,2-propanediol and
306 D-sorbitol before injection. Both proved to enhance the signal by more than 50% for all the
307 selected compounds.

308 **Detection and identification techniques**

309 Several analytical techniques enable the determination of contaminants in seafood samples
310 (Table 1). Liquid (LC) or gas chromatography (GC) separation method with tandem mass
311 spectrometry (MS/MS) detection are suitable confirmatory techniques for organic residues
312 and contaminants in foodstuffs [67]. The majority of the studies used GC or LC, coupled with
313 MS/MS, to determine organic compounds in bivalves (Table 2). GC-ECD (electron capture

314 detection) comes second (Table 2) [51,68]. Unlike MS, which is a universal detector, ECD is
315 a specific detector for halogenated organic molecules such as organochlorine (OC) pesticides
316 [8,51]. Thus, GC-ECD can only detect halogenated pesticides, whereas MS can detect all
317 types of pesticides provided they are ionised. In addition, GC allows a greater selectivity and
318 resolution than LC in determining contaminants in complex matrices, such as farmed fish and
319 shellfish [46]. Ziarrusta *et al.*, (2015) implemented GC, coupled to triple quadrupole MS
320 (GC–MS/MS), to determine pesticides in farmed mollusk bivalves representing complex
321 matrices such as blue mussels, mangrove oysters, radial purse oysters or ark clams [49]. The
322 study reported lower limits of detection (LODs) and quantification (LOQs) using GC–MS/MS
323 than GC-MS (49). Other analytical configurations include ultra-high-performance LC
324 (UHPLC) (Table 1), which uses particles of smaller diameter (around 1.7 μm) in the
325 stationary phase and short columns [59]. High pressures induce narrow LC peaks (5–10 s)
326 with shorter analysis [23,50,69]. Compared to the HPLC-based methods, UHPLC uses less
327 solvent in general. Furthermore, coupled with MS, UHPLC triggers greater throughput and
328 improves resolution [46]. The UHPLC method separation as well as the previously described
329 HPLC and GC coupled to triple quadrupole MS operating in multiple reaction monitoring
330 (MRM) mode have two major advantages: selectivity and sensitivity. However, the use of
331 triple quadrupole instruments has proven to have some limitations, such as a limited number
332 of compounds targeted during the analysis [70,71]. For large screening purposes, an
333 alternative to MS/MS is the application of full scan techniques based on high-resolution mass
334 spectrometry (HRMS), using Orbitrap MS-based instruments and time-of-flight (ToF) MS
335 systems [72]. This analytical technique can perform simultaneous target, suspect, and non-
336 target downstream data analyses to increase the number of analytes that can be included in
337 multi-residue methods. Moreover, it allows retrospective analysis by reprocessing acquired
338 data (as new knowledge is gained) and the ability to use samples in multiple studies without

339 having to reinject them [73]. Despite the advantages described, only two studies have used
340 HRMS for the determination of pesticides in bivalves. One performed target analysis of 13
341 pesticides in mussels, oysters, cockles and razor shell [66]. More recently, Diallo *et al.*,
342 (2022) carried out a simultaneous targeted analysis of 204 pesticides in seven different
343 commercial bivalve species and a suspect screening analysis on real samples of oysters and
344 mussels collected in south-west France [35]. However, studies using HRMS for “non-target”
345 pesticide analysis in bivalve tissues are still scarce despite offering promising perspectives for
346 detecting a wide range of pesticides in marine bivalves. This is especially relevant to
347 characterize an environmentally realistic exposure to a cocktail of contaminants. Furthermore,
348 the performance of HRMS systems, including automatic data processing, has improved
349 sufficiently to make their implementation in routine analysis a realistic option, particularly
350 when used in conjunction with conventional GC and LC-MS/MS methods. Moreover, the
351 sensitivity of recent HRMS systems has now increased. It should be noted that the objective
352 of HRMS is not to replace conventional methods but to perform a diagnosis for the presence
353 of unexpected contaminants in the sample, which can then be confirmed by LC-MS/MS. This
354 could have prevented recent food safety crises such as the presence of fipronil (which was not
355 expected) in European eggs in summer 2017 [74].

356 Along with the advent of HRMS, there have been several developments in the field of
357 UHPLC-MS in recent years. The most important of these are two-dimensional
358 chromatography (2D-GC or 2D-LC) and ion mobility spectrometry (IMS). The 2D
359 chromatography use both the comprehensive (GC) and heart cutting (LC) approaches [75,76].
360 In the comprehensive approach, the entire effluent from the first column is transferred to the
361 second one. The heart cutting technique involves selecting a fraction of the co-eluting zone
362 from the first column in a sample and re-inject it to the second column for separation and
363 detection in the MS instrument. Compared to 2D-LC, 2D-GC is more frequently used for the

364 determination of a wide mixture of pesticides [77–79]. The advantages over conventional
365 chromatography are increased peak capacity, generation of unique structured chromatograms,
366 better resolution of thousands of peaks in a chromatographic run and the possibility to reduce
367 matrix interference [79]. The HRMS technique also offers the possibility of performing non-
368 targeted analyses of various compounds in complex matrices [79]. As an alternative to 2D
369 chromatography, IMS can be considered for the analysis of very complex mixtures. In IMS,
370 ions are transported and separated according to their collisions with a neutral drift gas, such as
371 helium or nitrogen, allowing the separation of ions according to their three-dimensional
372 structure [80]. By coupling chromatographic separations, ion mobility separations (IMS) and
373 HRMS, the overall peak capacity of the system is increased and allows for the separation of
374 compounds (isomeric and isobaric) that would otherwise be detected as a single component in
375 a system without IMS [81]. One difference between IMS and 2D chromatography is that
376 when the latter is used, the user can optimize the peak capacity of the method by selecting the
377 appropriate separation selectivity of the two separation dimensions involved. When IMS is
378 used, the user loses this flexibility as drift times or derived collision cross section (CCS) are
379 compound dependent and relative drift times can only be altered to a certain degree by
380 modifying the drift gas composition. UHPLC coupled with travelling-wave (UHPLC-
381 TWIMS-QTOF-MS) was successfully used by Regueiro *et al.*, (2017), for qualitative
382 screening of 156 pesticides in salmon feed [82]. The authors found that the use of the CCS as
383 an additional identification point greatly improved the method selectivity while decreasing the
384 number of false positive detections (from 42 to only 1).

385 **Method validation**

386 To ensure the reliability of analytical information, a validation protocol must be
387 followed in every study. This protocol evaluates precision, trueness, specificity, recovery,
388 LOD and LOQ and sensitivity as established by the European Commission in 2021

389 (SANTE/11312/2021) [83]. The precision is actually evaluated through the repeatability and
390 reproducibility of the analytical response by estimation of the relative standard deviation
391 ($RSD \leq 20\%$) for each spike level tested [83]. The trueness is expressed in terms of bias,
392 which is the average recovery (70-120%) for each spike level tested. The recovery is
393 expressed in percentage and represents the proportion of analyte remaining at the point of the
394 final determination. The specificity is the ability of the detector to provide signals that
395 effectively identify the analyte [83].

396 A satisfactory validation protocol in marine bivalves was applied by Ziarrusta *et al.*,
397 (2015) using the SANCO/10684/2009 guideline (one of the former SANTE/11312/2021
398 guideline) [49]. The authors assessed the repeatability by preparing four replicates of mussels
399 at 50 ng/g and analysing them in the same day. Then, for reproducibility, eight replicates were
400 analysed during three different days. The RSD values varied from 1% to 17% for
401 repeatability, and from 4% to 33% for reproducibility. The LOD and LOQ values of several
402 organophosphorus and organochlorine pesticides ranged from 0.34 to 13 ng/g and 0.83 to 29
403 ng/g respectively [49]. These values were acceptable and comparable to the values found in
404 other studies that focused on the MSPD extraction of trace organic pollutants [4,84–86]. To
405 evaluate the accuracy and precision, Rodrigues *et al.* used two different concentrations, $3 \times$
406 MQL (method quantification limit) and $10 \times$ MQL (European Commission 2002/657/CE)
407 [23]. Another reliable method to evaluate accuracy and precision was used by Cruzeiro *et al.*,
408 (2016) who made three different pools of bivalve homogenate and injected them at three
409 concentrations: LOQ, $2 \times$ LOQ, and $10 \times$ LOQ (SANCO/825/00 rev 8.1) [34].

410 **Recommendations and future perspectives**

411 A large variety of pesticides can bioaccumulate in bivalves potentially affecting their survival
412 and human health. It is thus essential to refine the methods and techniques used to accurately

413 characterize the contamination of bivalves by organic chemicals to sustain a good ecological
414 status.

415 Recommendations for sample preparation are the use of QuEChERS using MgSO₄ or
416 Na₂SO₄ and enhanced by AOAC or EN method, innovative techniques respecting the
417 principles of green analytical chemistry, such as SBSE and MSPE and the use of new
418 materials as sorbents. Finally, we recommend dSPE and analytical protectors to reduce matrix
419 effect.

420 Recommendations for detecting of a wide range of pesticides and their residues in
421 bivalves are the use of LC/GC-HRMS and UHPLC-MS followed by 2D-GC, 2D-LC or IMS.
422 Specifically, HRMS can perform a diagnosis for the presence of unexpected contaminants in
423 the sample, which can then be confirmed by GC or LC-MS/MS.

424 Those approaches will allow a more rigorous characterisation of the real cocktail of
425 contaminants existing in the coastal environment.

426 **Acknowledgements**

427 This research is part of the AQUAECOs project (Amélioration de la QUALité
428 Environnementale sur les zones CONchylicoles des Pertuis Charentais), funded by a
429 partnership with the CRC17 (Comité Régional de la Conchyliculture de la Charente
430 Maritime), the AFB (Agence Française de la Biodiversité) and the Nouvelle Aquitaine region.
431 The researchers thank the ANR (French research agency) for their financial contribution
432 through the ANR JCJC AlimOmic ANR-19-CE21-0002.

433 **Compliance with ethical standards**

434 The author declares that there is no conflict of interest.

435 436 **Table and Figure Legend.**

437 **Table 1.** Analytical methods used to evaluate pesticide concentrations in bivalve species.

438 **Table 2.** Minimum (min), mean, maximum (max), and pesticide concentrations in ng/g dw
439 (dry weight), quantified in bivalves (collected at least at one location) collected along the
440 French Atlantic coast. The class of pesticides, the year of sampling and the number of samples
441 are presented. *ww (wet weight)

442 **Figure 1.** Map indicating the location of the sampling sites along the French Atlantic and
443 English Channel coasts (source: CAPENA).

444
445

446 **References**

- 447 1. Breitwieser M, Viricel A, Graber M, Murillo L, Becquet V, Churlaud C, et al. Short-
448 Term and Long-Term Biological Effects of Chronic Chemical Contamination on
449 Natural Populations of a Marine Bivalve. *PLoS One*. 2016;11(3):e0150184.
450 <https://doi.org/10.1371/journal.pone.0150184>
- 451 2. Burgos-Aceves MA, Faggio C. An approach to the study of the immunity functions of
452 bivalve haemocytes: Physiology and molecular aspects. *Fish Shellfish Immunol*.
453 2017;67:513- 7. <https://doi.org/10.1016/j.fsi.2017.06.042>
- 454 3. Prosser RS, de Solla SR, Holman EAM, Osborne R, Robinson SA, Bartlett AJ, et al.
455 Sensitivity of the early-life stages of freshwater mollusks to neonicotinoid and
456 butenolide insecticides. *Environ Pollut*. 2016;218:428- 35.
457 <https://doi.org/10.1016/j.envpol.2016.07.022>
- 458 4. Souza MR dos R, Moreira CO, de Lima TG, Aquino A, Dórea HS. Validation of a
459 matrix solid phase dispersion (MSPD) technique for determination of pesticides in
460 lyophilized eggs of the chicken *Gallus gallus domesticus*. *Microchem J*.
461 2013;110:395- 401. <https://doi.org/10.1016/j.microc.2013.05.001>
- 462 5. Riisgård HU. On measurement of filtration rates in bivalves — the stony road to
463 reliable data: review and interpretation. *Mar Ecol Prog Ser*. 2001;211:275- 91.
464 <https://doi.org/10.3354/meps211275>
- 465 6. Breitwieser M, Vigneau E, Viricel A, Becquet V, Lacroix C, Erb M, et al. What is the
466 relationship between the bioaccumulation of chemical contaminants in the variegated
467 scallop *Mimachlamys varia* and its health status? A study carried out on the French
468 Atlantic coast using the Path ComDim model. *Sci Total Environ*.
469 2018;640- 641:662- 70. <https://doi.org/10.1016/j.scitotenv.2018.05.317>
- 470 7. Faggio C, Tsarpali V, Dailianis S. Mussel digestive gland as a model tissue for
471 assessing xenobiotics: An overview. *Sci Total Environ*. 2018;636:220- 9.
472 <https://doi.org/10.1016/j.scitotenv.2018.04.264>
- 473 8. Luna-Acosta A, Budzinski H, Le Menach K, Thomas-Guyon H, Bustamante P.
474 Persistent organic pollutants in a marine bivalve on the Marennes-Oléron Bay and the
475 Gironde Estuary (French Atlantic Coast)—Part 1: Bioaccumulation. *Sci Total Environ*.
476 2015;514:500- 10. <https://doi.org/10.1016/j.scitotenv.2014.08.071>
- 477 9. Signa G, Di Leonardo R, Vaccaro A, Tramati CD, Mazzola A, Vizzini S. Lipid and
478 fatty acid biomarkers as proxies for environmental contamination in caged mussels
479 *Mytilus galloprovincialis*. *Ecol Indic*. 2015;57:384- 94.
480 <https://doi.org/10.1016/j.ecolind.2015.05.002>
- 481 10. Désert M, Ravier S, Gille G, Quinapallo A, Armengaud A, Pochet G, et al. Spatial and
482 temporal distribution of current-use pesticides in ambient air of Provence-Alpes-Côte-
483 d'Azur Region and Corsica, France. *Atmos Environ*. 2018;192:241- 56.
484 <https://doi.org/10.1016/j.atmosenv.2018.08.054>
- 485 11. Abreu-Villaça Y, Levin ED. Developmental neurotoxicity of succeeding generations of
486 insecticides. *Environ Int*. 2017;99:55- 77. <https://doi.org/10.1016/j.envint.2016.11.019>

- 487 12. Moncaleano-Niño AM, Luna-Acosta A, Gómez-Cubillos MC, Villamil L, Ahrens MJ.
488 Cholinesterase activity in the cup oyster *Saccostrea* sp. exposed to chlorpyrifos,
489 imidacloprid, cadmium and copper. *Ecotoxicol Environ Saf.* 2018;151:242- 54.
490 <https://doi.org/10.1016/j.ecoenv.2017.12.057>
- 491 13. Stehle S, Bub S, Schulz R. Compilation and analysis of global surface water
492 concentrations for individual insecticide compounds. *Sci Total Environ.*
493 2018;639:516- 25. <https://doi.org/10.1016/j.scitotenv.2018.05.158>
- 494 14. Girones L, Oliva AL, Marcovecchio JE, Arias AH. Spatial Distribution and Ecological
495 Risk Assessment of Residual Organochlorine Pesticides (OCPs) in South American
496 Marine Environments. *Curr Environ Health Rep.* 2020;7(2):147- 60.
497 <https://doi.org/10.1007/s40572-020-00272-7>
- 498 15. Miranda JAT, Fogaça FHS, Cunha SC, Alonso MB, Torres JPM, Fernandes JO.
499 Agrochemical Residues in Fish and Bivalves from Sepetiba Bay and Parnaíba River
500 Delta, Brazil. *Int J Environ Res Public Health.* 2022;19(23):15790.
501 <https://doi.org/10.3390/ijerph192315790>
- 502 16. European Commission. Commission Delegated Regulation (EU) 2022/643 of 10
503 February 2022 amending Regulation (EU) No 649/2012 of the European Parliament
504 and of the Council as regards the listing of pesticides, industrial chemicals, persistent
505 organic pollutants and mercury and an update of customs codes (Text with EEA
506 relevance). 2022. http://data.europa.eu/eli/reg_del/2022/643/oj
- 507 17. Menet-Nedelec F, Halm-Lemeille M-P, Maheux F, Pierre-Duplessix O, Simon B,
508 Gonzalez J-L, et al. Etude d'outils d'évaluation de la contamination Chimique dans les
509 eaux de la Manche - ECUME. 2018. <https://archimer.ifremer.fr/doc/00434/54582/>
- 510 18. Lerebours A, Bathie M, Receveur J, Jézéquel R, Dubillot E, Brunello P, et al.
511 Pesticides, nonylphenols and polybrominated diphenyl ethers in marine bivalves from
512 France: A pilot study. *Mar Pollut Bull.* 2021;172:112956.
513 <https://doi.org/10.1016/j.marpolbul.2021.112956>
- 514 19. Lebourg A, Morin N, Glize P. Rapport technique CAPEMOULES (2015-2017).
515 Caractérisation des performances des élevages de moules et évaluation de la qualité du
516 milieu en pays de la Loire. Comité régional de la conchyliculture des pays de la Loire..
517 2018. [https://www.wikimer.org/projets/capemoules-caracterisation-des-performances-
518 des-elevages-de-moules-et-evaluation-de-la-qualite-du-milieu-en-pays-de-la-loire/](https://www.wikimer.org/projets/capemoules-caracterisation-des-performances-des-elevages-de-moules-et-evaluation-de-la-qualite-du-milieu-en-pays-de-la-loire/)
- 519 20. Stara A, Pagano M, Capillo G, Fabrello J, Sandova M, Vazzana I, et al. Assessing the
520 effects of neonicotinoid insecticide on the bivalve mollusc *Mytilus galloprovincialis*.
521 *Sci Total Environ.* 2020;700:134914. <https://doi.org/10.1016/j.scitotenv.2019.134914>
- 522 21. Carro AM, Lorenzo RA, Fernández F, Rodil R, Cela R. Multi-residue screening of
523 chlorinated and brominated compounds from aquaculture samples using matrix solid-
524 phase dispersion—gas chromatography—mass spectrometry. *J Chromatogr A.* 15 avr
525 2005;1071(1):93- 8. <https://doi.org/10.1016/j.chroma.2004.12.005>
- 526 22. Tu X, Chen W. A Review on the Recent Progress in Matrix Solid Phase Dispersion.
527 *Molecules.* 2018;23(11). <https://doi.org/10.3390/molecules23112767>

- 528 23. Rodrigues ET, Pardal MÂ, Salgueiro-González N, Muniategui-Lorenzo S, Alpendurada
529 MF. A single-step pesticide extraction and clean-up multi-residue analytical method by
530 selective pressurized liquid extraction followed by on-line solid phase extraction and
531 ultra-high-performance liquid chromatography-tandem mass spectrometry for complex
532 matrices. *J Chromatogr A*. 2016;1452:10- 7.
533 <https://doi.org/10.1016/j.chroma.2016.05.036>
- 534 24. Kim L, Lee D, Cho H-K, Choi S-D. Review of the QuEChERS method for the analysis
535 of organic pollutants: Persistent organic pollutants, polycyclic aromatic hydrocarbons,
536 and pharmaceuticals. *Trends Environ Anal Chem*. 2019;22:e00063.
537 <https://doi.org/10.1016/j.teac.2019.e00063>
- 538 25. Subedi B, Aguilar L, Robinson EM, Hageman KJ, Björklund E, Sheesley RJ, et al.
539 Selective pressurized liquid extraction as a sample-preparation technique for persistent
540 organic pollutants and contaminants of emerging concern. *TrAC Trends Anal Chem*.
541 2015;68:119- 32. <https://doi.org/10.1016/j.trac.2015.02.011>
- 542 26. Vazquez-Roig P, Picó Y. Pressurized liquid extraction of organic contaminants in
543 environmental and food samples. *TrAC Trends Anal Chem*. 2015;71:55- 64.
544 <https://doi.org/10.1016/j.trac.2015.04.014>
- 545 27. Hoff RB, Pizzolato TM. Combining extraction and purification steps in sample
546 preparation for environmental matrices: A review of matrix solid phase dispersion
547 (MSPD) and pressurized liquid extraction (PLE) applications. *TrAC Trends Anal*
548 *Chem*. 2018;109:83- 96. <https://doi.org/10.1016/j.trac.2018.10.002>
- 549 28. Ghosh R, Hageman KJ, Björklund E. Selective pressurized liquid extraction of three
550 classes of halogenated contaminants in fish. *J Chromatogr A*. 2011;1218(41):7242- 7.
551 <https://doi.org/10.1016/j.chroma.2011.08.052>
- 552 29. Wardrop P, Morrison PD, Hughes JG, Clarke BO. Comparison of in-cell lipid removal
553 efficiency of adsorbent mixtures for extraction of polybrominated diphenyl ethers in
554 fish. *J Chromatogr B*. 2015;990:1- 6. <https://doi.org/10.1016/j.jchromb.2015.02.045>
- 555 30. Merdassa Y, Liu J, Megersa N. Development of a one-step microwave-assisted
556 extraction method for simultaneous determination of organophosphorus pesticides and
557 fungicides in soils by gas chromatography–mass spectrometry. *Talanta*.
558 2013;114:227- 34. <https://doi.org/10.1016/j.talanta.2013.04.035>
- 559 31. Uluturhan E, Darılmaz E, Kontas A, Bilgin M, Alyuruk H, Altay O, et al. Seasonal
560 variations of multi-biomarker responses to metals and pesticides pollution in *M.*
561 *galloprovincialis* and *T. decussatus* from Homa Lagoon, Eastern Aegean Sea. *Mar*
562 *Pollut Bull*. 2019;141:176- 86. <https://doi.org/10.1016/j.marpolbul.2019.02.035>
- 563 32. Carpentieri S, Soltanipour F, Ferrari G, Pataro G, Donsì F. Emerging Green Techniques
564 for the Extraction of Antioxidants from Agri-Food By-Products as Promising
565 Ingredients for the Food Industry. *Antioxidants*. 2021;10(9):1417.
566 <https://doi.org/10.3390/antiox10091417>
- 567 33. Anastassiades M, Lehotay SJ, Stajnbaher D, Schenck FJ. Fast and easy multiresidue
568 method employing acetonitrile extraction/partitioning and « dispersive solid-phase

- 569 extraction » for the determination of pesticide residues in produce. *J AOAC Int.*
570 2003;86(2):412- 31.
- 571 34. Cruzeiro C, Rodrigues-Oliveira N, Velhote S, Pardal MÂ, Rocha E, Rocha MJ.
572 Development and application of a QuEChERS-based extraction method for the analysis
573 of 55 pesticides in the bivalve *Scrobicularia plana* by GC-MS/MS. *Anal Bioanal Chem.*
574 2016;408(14):3681- 98. <https://doi.org/10.1007/s00216-016-9440-0>
- 575 35. Diallo T, Makni Y, Lerebours A, Thomas H, Guérin T, Parinet J. Development and
576 validation according to the SANTE guidelines of a QuEChERS-UHPLC-QTOF-MS
577 method for the screening of 204 pesticides in bivalves. *Food Chem.* 2022;386:132871.
578 <https://doi.org/10.1016/j.foodchem.2022.132871>
- 579 36. Lehotay SJ. Determination of pesticide residues in foods by acetonitrile extraction and
580 partitioning with magnesium sulfate: collaborative study. *J AOAC Int.* avr
581 2007;90(2):485- 520.
- 582 37. Anastassiades M, Scherbaum E, Taşdelen B, Štajnbaher D. Recent Developments in
583 QuEChERS Methodology for Pesticide Multiresidue Analysis. In: *Pesticide Chemistry.*
584 John Wiley & Sons, Ltd; 2007. p. 439- 58.
585 <https://doi.org/10.1002/9783527611249.ch46>
- 586 38. Chang Y-C, Chen D-H. Adsorption Kinetics and Thermodynamics of Acid Dyes on a
587 Carboxymethylated Chitosan-Conjugated Magnetic Nano-Adsorbent. *Macromol*
588 *Biosci.* 2005;5(3):254- 61. <https://doi.org/10.1002/mabi.200400153>
- 589 39. Câmara JS, Perestrelo R, Berenguer CV, Andrade CFP, Gomes TM, Olayanju B, et al.
590 Green Extraction Techniques as Advanced Sample Preparation Approaches in
591 Biological, Food, and Environmental Matrices: A Review. *Molecules.* 2022;27(9).
592 <https://doi.org/10.3390/molecules27092953>
- 593 40. Camino-Sánchez FJ, Rodríguez-Gómez R, Zafra-Gómez A, Santos-Fandila A, Vílchez
594 JL. Stir bar sorptive extraction: Recent applications, limitations and future trends.
595 *Talanta.* 2014;130:388- 99. <https://doi.org/10.1016/j.talanta.2014.07.022>
- 596 41. Raccary B, Loubet P, Peres C, Sonnemann G. Life cycle assessment of sample
597 preparation in analytical chemistry: a case study on SBSE and SPE techniques. *Adv*
598 *Sample Prep.* 2022;1:100009. <https://doi.org/10.1016/j.sampre.2022.100009>
- 599 42. Cordero C, Sgorbini B, Liberto E, Bicchi C, Rubiolo P. Stir Bar Sorptive Extraction
600 (SBSE) and Headspace Sorptive Extraction (HSSE): An Overview. *LCGC N Am.*
601 2009;27(5):376–390- 376–390.
- 602 43. Baduel C, Mueller JF, Tsai H, Gomez Ramos MJ. Development of sample extraction
603 and clean-up strategies for target and non-target analysis of environmental
604 contaminants in biological matrices. *J Chromatogr A.* 2015;1426:33- 47.
605 <https://doi.org/10.1016/j.chroma.2015.11.040>
- 606 44. Anumol T, Lehotay SJ, Stevens J, Zweigenbaum J. Comparison of veterinary drug
607 residue results in animal tissues by ultrahigh-performance liquid chromatography
608 coupled to triple quadrupole or quadrupole–time-of-flight tandem mass spectrometry
609 after different sample preparation methods, including use of a commercial lipid

- 610 removal product. *Anal Bioanal Chem.* 2017;409(10):2639- 53.
611 <https://doi.org/10.1007/s00216-017-0208-y>
- 612 45. Masiá A, Suarez-Varela MM, Llopis-Gonzalez A, Picó Y. Determination of pesticides
613 and veterinary drug residues in food by liquid chromatography-mass spectrometry: A
614 review. *Anal Chim Acta.* 2016;936:40- 61. <https://doi.org/10.1016/j.aca.2016.07.023>
- 615 46. Justino CIL, Duarte KR, Freitas AC, Panteleitchouk TSL, Duarte AC, Rocha-Santos
616 TAP. Contaminants in aquaculture: Overview of analytical techniques for their
617 determination. *TrAC Trends Anal Chem.* 2016;80:293- 310.
618 <https://doi.org/10.1016/j.trac.2015.07.014>
- 619 47. Chiesa LM, Nobile M, Malandra R, Pessina D, Panseri S, Labella GF, et al. Food
620 safety traits of mussels and clams: distribution of PCBs, PBDEs, OCPs, PAHs and
621 PFASs in sample from different areas using HRMS-Orbitrap® and modified
622 QuEChERS extraction followed by GC-MS/MS. *Food Addit Contam Part A.*
623 2018;35(5):959- 71. <https://doi.org/10.1080/19440049.2018.1434900>
- 624 48. Cloutier P-L, Fortin F, Groleau PE, Brousseau P, Fournier M, Desrosiers M.
625 QuEChERS extraction for multi-residue analysis of PCBs, PAHs, PBDEs and
626 PCDD/Fs in biological samples. *Talanta.* 2017;165:332- 8.
627 <https://doi.org/10.1016/j.talanta.2016.12.080>
- 628 49. Ziarrusta H, Olivares M, Delgado A, Posada-Ureta O, Zuloaga O, Etxebarria N.
629 Multiscreening determination of organic pollutants in molluscs using matrix solid
630 phase dispersion. *J Chromatogr A.* 2015;1391:18- 30.
631 <https://doi.org/10.1016/j.chroma.2015.02.072>
- 632 50. Álvarez-Ruiz R, Picó Y. Sequential window acquisition of all theoretical fragments
633 versus information dependent acquisition for suspected-screening of pharmaceuticals in
634 sediments and mussels by ultra-high pressure liquid chromatography-quadrupole time-
635 of-flight-mass spectrometry. *J Chromatogr A.* 2019;1595:81- 90.
636 <https://doi.org/10.1016/j.chroma.2019.02.041>
- 637 51. Luna-Acosta A, Bustamante P, Budzinski H, Huet V, Thomas-Guyon H. Persistent
638 organic pollutants in a marine bivalve on the Marennes–Oléron Bay and the Gironde
639 Estuary (French Atlantic Coast)—Part 2: Potential biological effects. *Sci Total*
640 *Environ.* 1 mai 2015;514:511- 22. <https://doi.org/10.1016/j.scitotenv.2014.10.050>
- 641 52. Lehotay SJ. QuEChERS Sample Preparation Approach for Mass Spectrometric
642 Analysis of Pesticide Residues in Foods. In: Zweigenbaum J, éditeur. *Mass*
643 *Spectrometry in Food Safety: Methods and Protocols.* Totowa, NJ: Humana Press;
644 2011. p. 65- 91. (Methods in Molecular Biology). https://doi.org/10.1007/978-1-61779-136-9_4
- 646 53. Han L, Sapozhnikova Y, Lehotay SJ. Streamlined sample cleanup using combined
647 dispersive solid-phase extraction and in-vial filtration for analysis of pesticides and
648 environmental pollutants in shrimp. *Anal Chim Acta.* 2014;827:40- 6.
649 <https://doi.org/10.1016/j.aca.2014.04.005>

- 650 54. Oshita D, Jardim ICSF. Evaluation of dispersive and cartridge SPE clean-up procedures
651 using the modified QuEChERS method for the analysis of pesticides in strawberries.
652 *Anal Methods*. 2015;7(3):982- 9. <https://doi.org/10.1039/C4AY01867B>
- 653 55. Núñez M, Borrull F, Fontanals N, Pocurull E. Determination of pharmaceuticals in
654 bivalves using QuEChERS extraction and liquid chromatography-tandem mass
655 spectrometry. *Anal Bioanal Chem*. 2015;407(13):3841- 9.
656 <https://doi.org/10.1007/s00216-015-8617-2>
- 657 56. Álvarez-Ruiz R, Picó Y, Sadutto D, Campo J. Development of multi-residue extraction
658 procedures using QuEChERS and liquid chromatography tandem mass spectrometry
659 for the determination of different types of organic pollutants in mussel. *Anal Bioanal*
660 *Chem*. 2021;413(15):4063- 76. <https://doi.org/10.1007/s00216-021-03363-y>
- 661 57. Slámová T, Sadowska-Rociek A, Fraňková A, Surma M, Banout J. Application of
662 QuEChERS-EMR-Lipid-DLLME method for the determination of polycyclic aromatic
663 hydrocarbons in smoked food of animal origin. *J Food Compos Anal*. 2020;87:103420.
664 <https://doi.org/10.1016/j.jfca.2020.103420>
- 665 58. Gao M, Gao Y, Chen G, Huang X, Xu X, Lv J, et al. Recent Advances and Future
666 Trends in the Detection of Contaminants by Molecularly Imprinted Polymers in Food
667 Samples. *Front Chem*. 2020;8.
668 <https://www.frontiersin.org/article/10.3389/fchem.2020.616326>
- 669 59. Yu X, Zhang R, Liu H, Zhang Z, Shi X, Sun A, et al. Highly-selective complex
670 matrices removal via a modified QuEChERS for determination of triazine herbicide
671 residues and risk assessment in bivalves. *Food Chem*. 2021;347:129030.
672 <https://doi.org/10.1016/j.foodchem.2021.129030>
- 673 60. Raynie D. Next-Generation Sorbent-Based Extractions with Metal-Organic
674 Frameworks. *LCGC N Am*. 2021;39(3):120- 7.
- 675 61. Bazargan M, Ghaemi F, Amiri A, Mirzaei M. Metal–organic framework-based sorbents
676 in analytical sample preparation. *Coord Chem Rev*. 2021;445:214107.
677 <https://doi.org/10.1016/j.ccr.2021.214107>
- 678 62. Bagheri AR, Aramesh N, Bilal M. New frontiers and prospects of metal-organic
679 frameworks for removal, determination, and sensing of pesticides. *Environmental*
680 *Research*. 2021;194:110654. <https://doi.org/10.1016/j.envres.2020.110654>
- 681 63. Musarurwa H, Chimuka L, Tavengwa NT. Sorptive extraction of pesticides from food
682 and environmental samples using metal organic framework-based adsorbents. *Trends*
683 *Environ Anal Chem*. 2021;32:e00141. <https://doi.org/10.1016/j.teac.2021.e00141>
- 684 64. Fallah Z, Zare EN, Ghomi M, Ahmadijokani F, Amini M, Tajbakhsh M, et al. Toxicity
685 and remediation of pharmaceuticals and pesticides using metal oxides and carbon
686 nanomaterials. *Chemosphere*. 2021;275:130055.
687 <https://doi.org/10.1016/j.chemosphere.2021.130055>
- 688 65. Xiang Y, Xu Z, Wei Y, Zhou Y, Yang X, Yang Y, et al. Carbon-based materials as
689 adsorbent for antibiotics removal: Mechanisms and influencing factors. *J Environ*
690 *Manage*. 2019;237:128- 38. <https://doi.org/10.1016/j.jenvman.2019.02.068>

- 691 66. Álvarez-Muñoz D, Rambla-Alegre M, Carrasco N, Lopez de Alda M, Barceló D. Fast
692 analysis of relevant contaminants mixture in commercial shellfish. *Talanta*.
693 2019;205:119884. <https://doi.org/10.1016/j.talanta.2019.04.085>
- 694 67. 2002/657/EC: Commission Decision of 12 August 2002 implementing Council
695 Directive 96/23/EC concerning the performance of analytical methods and the
696 interpretation of results (Text with EEA relevance) (notified under document number
697 C(2002) 3044). *OJ L*, 32002D0657, 2002. <http://data.europa.eu/eli/dec/2002/657/oj/eng>
- 698 68. de Souza RV, Garbossa LHP, Campos CJA, Vianna LF de N, Vanz A, Rupp GS.
699 Metals and pesticides in commercial bivalve mollusc production areas in the North and
700 South Bays, Santa Catarina (Brazil). *Mar Pollut Bull*. 2016;105(1):377- 84.
701 <https://doi.org/10.1016/j.marpolbul.2016.02.024>
- 702 69. Cruz JM. Etude de la contamination par les pesticides des milieux eau, air et sols :
703 développement de nouveaux outils et application à l'estuaire de la Gironde [phdthesis].
704 Université de Bordeaux; 2015. <https://theses.hal.science/tel-01359188>
- 705 70. Gavage M, Delahaut P, Gillard N. Suitability of High-Resolution Mass Spectrometry
706 for Routine Analysis of Small Molecules in Food, Feed and Water for Safety and
707 Authenticity Purposes: A Review. *Foods*. 2021;10(3):601.
708 <https://doi.org/10.3390/foods10030601>
- 709 71. Hermes N, Jewell KS, Wick A, Ternes TA. Quantification of more than 150
710 micropollutants including transformation products in aqueous samples by liquid
711 chromatography-tandem mass spectrometry using scheduled multiple reaction
712 monitoring. *J Chromatogr A*. 2018;1531:64- 73.
713 <https://doi.org/10.1016/j.chroma.2017.11.020>
- 714 72. Saito-Shida S, Hamasaka T, Nemoto S, Akiyama H. Multiresidue determination of
715 pesticides in tea by liquid chromatography-high-resolution mass spectrometry:
716 Comparison between Orbitrap and time-of-flight mass analyzers. *Food Chem*.
717 2018;256:140- 8. <https://doi.org/10.1016/j.foodchem.2018.02.123>
- 718 73. Rajski Ł, Martínez-Bueno MJ, Ferrer C, Fernández-Alba AR. LC-ESI-QOrbitrap™
719 MS/MS within pesticide residue analysis in fruits and vegetables. *TrAC Trends Anal*
720 *Chem*. 2019;118:587- 96. <https://doi.org/10.1016/j.trac.2019.06.032>
- 721 74. Reich H, Triacchini GA. Occurrence of residues of fipronil and other acaricides in
722 chicken eggs and poultry muscle/fat. *EFSA J*. 2018;16(5):e05164.
723 <https://doi.org/10.2903/j.efsa.2018.5164>
- 724 75. Dunmore RE, Hopkins JR, Lidster RT, Mead MI, Bandy BJ, Forster G, et al.
725 Development of a Combined Heart-Cut and Comprehensive Two-Dimensional Gas
726 Chromatography System to Extend the Carbon Range of Volatile Organic Compounds
727 Analysis in a Single Instrument. *Separations*. 2016;3(3):21.
728 <https://doi.org/10.3390/separations3030021>
- 729 76. López-Ruiz R, Romero-González R, Garrido Frenich A. Ultrahigh-pressure liquid
730 chromatography-mass spectrometry: An overview of the last decade. *TrAC Trends*
731 *Anal Chem*. 2019;118:170- 81. <https://doi.org/10.1016/j.trac.2019.05.044>

- 732 77. Buah-Kwofie A, Humphries MS. Validation of a modified QuEChERS method for the
733 analysis of organochlorine pesticides in fatty biological tissues using two-dimensional
734 gas chromatography. *J Chromatogr B*. 2019;1105:85- 92.
735 <https://doi.org/10.1016/j.jchromb.2018.12.010>
- 736 78. Keshet U, Goldshlag P, Amirav A. Pesticide analysis by pulsed flow modulation
737 GCxGC-MS with Cold EI—an alternative to GC-MS-MS. *Anal Bioanal Chem*.
738 2018;410(22):5507- 19. <https://doi.org/10.1007/s00216-017-0757-0>
- 739 79. Pico Y, Alfarhan AH, Barcelo D. How recent innovations in gas chromatography-mass
740 spectrometry have improved pesticide residue determination: An alternative technique
741 to be in your radar. *TrAC Trends Anal Chem*. 2020;122:115720.
742 <https://doi.org/10.1016/j.trac.2019.115720>
- 743 80. Gabelica V. CHAPTER 1 Ion Mobility–Mass Spectrometry: an Overview. 2021;1- 25.
744 <https://doi.org/10.1039/9781839162886-00001>
- 745 81. Mullin L, Cleland G, Burgess JA. Chapter 8 - Recent Advances in HRMS Analysis of
746 Pesticide Residues Using Atmospheric Pressure Gas Chromatography and Ion
747 Mobility. In: Romero-González R, Frenich AG, éditeurs. *Applications in High*
748 *Resolution Mass Spectrometry*. Elsevier; 2017. p. 233- 64.
749 <https://doi.org/10.1016/B978-0-12-809464-8.00008-7>
- 750 82. Regueiro J, Negreira N, Hannisdal R, Berntssen MHG. Targeted approach for
751 qualitative screening of pesticides in salmon feed by liquid chromatography coupled to
752 traveling-wave ion mobility/quadrupole time-of-flight mass spectrometry. *Food*
753 *Control*. 2017;78:116- 25. <https://doi.org/10.1016/j.foodcont.2017.02.053>
- 754 83. European Commission. SANTE/11312/2021. Analytical quality control and method
755 validation procedures for pesticide residues analysis in food and feed. 2021.
756 [https://food.ec.europa.eu/system/files/2022-](https://food.ec.europa.eu/system/files/2022-02/pesticides_mrl_guidelines_wrkdoc_2021-11312.pdf)
757 [02/pesticides_mrl_guidelines_wrkdoc_2021-11312.pdf](https://food.ec.europa.eu/system/files/2022-02/pesticides_mrl_guidelines_wrkdoc_2021-11312.pdf)
- 758 84. García de Llasera MP, Reyes-Reyes ML. A validated matrix solid-phase dispersion
759 method for the extraction of organophosphorus pesticides from bovine samples. *Food*
760 *Chem*. 2009;114(4):1510- 6. <https://doi.org/10.1016/j.foodchem.2008.11.006>
- 761 85. Llompart M, Celeiro M, Pablo Lamas J, Sanchez-Prado L, Lores M, Garcia-Jares C.
762 Analysis of plasticizers and synthetic musks in cosmetic and personal care products by
763 matrix solid-phase dispersion gas chromatography–mass spectrometry. *J Chromatogr*
764 *A*. 2013;1293:10- 9. <https://doi.org/10.1016/j.chroma.2013.03.067>
- 765 86. Villaverde-de-Sáa E, Valls-Cantenys C, Quintana JB, Rodil R, Cela R. Matrix solid-
766 phase dispersion combined with gas chromatography–mass spectrometry for the
767 determination of fifteen halogenated flame retardants in mollusks. *J Chromatogr A*.
768 2013;1300:85- 94. <https://doi.org/10.1016/j.chroma.2013.05.064>
- 769 87. de Souza RM, Seibert D, Quesada HB, de Jesus Bassetti F, Fagundes-Klen MR,
770 Bergamasco R. Occurrence, impacts and general aspects of pesticides in surface water:
771 A review. *Process Saf Environ Prot*. 2020;135:22- 37.
772 <https://doi.org/10.1016/j.psep.2019.12.035>

- 773 88. PubChem. o,o'-DDT. <https://pubchem.ncbi.nlm.nih.gov/compound/154395>
- 774 89. UNEP. STOCKHOLM CONVENTION ON POPs. Text and Annexes. Stockh. Conv.
775 Persistent Org. Pollut. 2009.
776 http://chm.pops.int/portals/0/repository/convention_text/unep-pops-cop-convtext-
777 [full.english.pdf](http://chm.pops.int/portals/0/repository/convention_text/unep-pops-cop-convtext-full.english.pdf)
- 778 90. Barletta M, Lima ARA, Costa MF. Distribution, sources and consequences of nutrients,
779 persistent organic pollutants, metals and microplastics in South American estuaries. *Sci*
780 *Total Environ.* 2019;651:1199- 218. <https://doi.org/10.1016/j.scitotenv.2018.09.276>
- 781 91. Commendatore MG, Franco MA, Costa PG, Castro IB, Fillmann G, Bigatti G, et al.
782 Butyltins, polyaromatic hydrocarbons, organochlorine pesticides, and polychlorinated
783 biphenyls in sediments and bivalve mollusks in a mid-latitude environment from the
784 Patagonian coastal zone. *Environ Toxicol Chem.* 2015;34(12):2750- 63.
785 <https://doi.org/10.1002/etc.3134>
- 786 92. Fukuto TR. Mechanism of action of organophosphorus and carbamate insecticides.
787 *Environ Health Perspect.* 1990;87:245- 54. <https://doi.org/10.1289/ehp.9087245>
- 788 93. PubChem. Ethoprophos. <https://pubchem.ncbi.nlm.nih.gov/compound/3289>
- 789 94. Tang W, Wang D, Wang J, Wu Z, Li L, Huang M, et al. Pyrethroid pesticide residues
790 in the global environment: An overview. *Chemosphere.* 2018;191:990- 1007.
791 <https://doi.org/10.1016/j.chemosphere.2017.10.115>
- 792 95. Werner I, Hitzfeld B. 50 Years of ecotoxicology since Silent Spring - A review. *Gaia.*
793 2012;21(3):217- 24. <https://doi.org/10.14512/gaia.21.3.13>
- 794 96. Li H, Cheng F, Wei Y, Lydy MJ, You J. Global occurrence of pyrethroid insecticides
795 in sediment and the associated toxicological effects on benthic invertebrates: An
796 overview. Vol. 324, *Journal of Hazardous Materials.* Elsevier B.V.; 2017. 258- 271 p.
797 <https://doi.org/10.1016/j.jhazmat.2016.10.056>
- 798
- 799
- 800

801 Table 1.

Species	Sample	Pretreatment	Sample mass (g,ww)	Extraction method	Clean-up	Detection	LODs (ng/g)	LOQs (ng/g)	Recovery (%)	Reference
Peppery furrow shell (<i>Scrobicularia plana</i>)	-	Homogenized	5	QuEChERS	PSA dSPE	GC-MS/MS (IT) in SIM	0.22-3.4	0.33-10.3	78-119	[34]
Blue mussel (<i>Mytilus edulis</i>), Oysters (<i>Crassostrea rhizophorae</i> and <i>Isognomon radiatus</i>), Clams (<i>Polymesoda solida</i> , <i>Anadara tuberculosa</i> and <i>Anadara grandis</i>)	Pool	Homogenized and freeze-dried	0.5	MSPD	Silica SPE	GC(EI)-MS(QqQ) in SRM and SIM	0.34-13	0.83-29	55-117	[49]
						GC(EI)-MS/MS(QqQ) in SRM and SIM	0.010-2.7	0.017-3.9	64-122	
Pacific oyster (<i>Crassostrea gigas</i>)	Pool	Homogenized and freeze-dried	0.5	ASE, PLE	Acidic silica gel column	Quantified: GC-ECD Confirmed: GC-MS/MS	0.1-0.2	0.3-0.6	72-116	[8,51]
Mediterranean Mussel (<i>Mytilus galloprovincialis</i>), oyster (<i>Crassostrea gigas</i>), cockle (<i>Cerastoderma edule</i>)	Pool	Homogenized and freeze-dried	1.0	QuEChERS	C18 dSPE	UHPLC-QTOF MS/MS	0.01-0.5	0.02-1.00	96-107	[66]

Brown mussel (<i>Perna perna</i>), Pacific oyster (<i>C. gigas</i>)	-	-	-	-		LC-MS/MS	0.01-0.7	-	-	[68]
Mussels (<i>M. galloprovincialis</i> , <i>M. edulis</i> , <i>Mytilus chilensis</i> and <i>Perna canaliculus</i>), Clams (<i>Venerupis philippinarum</i> , <i>Tapes semidecussatus</i> , <i>Meretrix meretrix</i> and <i>Meretrix lyrata</i>)	Pool	Homogenized and stored at -20°C	5	QuEChERS EN method	500mg Z-Sep	GC-MS/MS	-	5	70-120	[47]
Blue mussel (<i>M. edulis</i>), Salmon (<i>Salmo salar</i>)	-	-	-	QuEChERS	GPC followed by silica SPE	GC-HRMS	11-680	-	90-110	[48]
Blue mussel (<i>M. edulis</i>)		Lyophilised	0.5 - 1	1) SPE acetonitril	Florisil	GC-MS/MS	-	10	-	[19]
	-	-	-	2) SPE dichloromethane/ethyl acetate	cSPE oasis HLB	LC-MS/MS	-	10	-	
	-	-	-	3) SPE dichloromethane/ethyl acetate in acid environment		-	-	-	-	
Blue mussel (<i>M. edulis</i>), Pacific oyster (<i>C. gigas</i>)	Individual	Homogenized	1 - 3	QuEChERS		GC-MS/MS		0.1-0.5	0.33-1.67	[18]

Peppery furrow shell (<i>S. plana</i>)	Pool	Homogenized and stored at - 20°C	0.5	SPLE	The Oasis [®] on-line columns	UPLC-ESI- MS/MS	3.7-97	11-290	68-125	[23]
Clams (<i>T. decussatus</i>)	Pool	Homogenized and freeze- dried	3 - 5	MAE	Florisil	GC	-	-	63-98	[31]

802

803 Table 2.

Pesticide	Class	CAS Number	Species	Concentration (ng/g, dw)			Sites	Reference	Year	Sample number
				Min	Mean	Max				
Acetochlor	OC	34256-82-1	Mediterranean mussel (<i>M. galloprovincialis</i>)		125		Chausey archipelago	[17]	2015	1
					67.4		Bay of Veys			1
				29.7		53.0	Seine estuary			1
Aldrin	OC	309-00-2	Variegated scallops (<i>Mimachlamys varia</i>)		2.24		Loix en Ré	[1]	2014	1
					2.71		Les Minimes			1
				<LOD		<LOQ	Les Palles			1
			Mediterranean mussel (<i>M. galloprovincialis</i>)		1.8		Chausey archipelago	[17]	2015	1
					3.3		Bay of Veys			1
				1.2		4.7	Seine estuary			1
4,4'-DDD	OC	2-54-8	Mediterranean mussel (<i>M. galloprovincialis</i>)		<0.1		Chausey archipelago	[17]	2015	1
					<0.1		Bay of Veys			1
				0.2		0.3	Seine estuary			1
				1.5	2.3	Bouin	[8]	2008	2	
				1.9	2.8	Boyard			2	
	2.0	3.7	Les Palles	2						
		1.0	1.9	Mus du loup			2			
2,4'-DDE	OC	3424-82-6	Mediterranean mussel (<i>M. galloprovincialis</i>)		<1		Chausey archipelago	[17]	2015	1
					<1		Bay of Veys			1
				<1		2.4	Seine estuary			1
			Pacific oyster (<i>C.</i>)	2.2	2.5		Bouin	[8]	2008	2

			<i>gigas</i>)	2.4	5.5	Boyard			2
				1.5	2.4	Les Palles			2
				nd	0.3	Cordouan			2
4,4'DDE + Dieldrin	-	-	Pacific oyster (<i>C. gigas</i>)	1.2	1.4	Bouin			2
				3.6	3.8	Boyard			2
				2.4	3.7	Les Palles	[8]	2008	2
				0.8	0.9	Mus du loup			2
				1.9	3.0	Cordouan			2
4,4'-DDT	50-29-3		Mediterranean mussel (<i>M. galloprovincialis</i>)		0.2	Chausey archipelago			1
				<0.1		Bay of Veys	[17]	2015	1
				2.5	2.9	Seine estuary			1
Dieldrin	OC	60-57-1	Variegated scallops (<i>M. varia</i>)	6.4		Loix en Ré			1
				12.4		Les Minimes	[1]	2014	1
				nd		Port-neuf			1
				4.2		Les Palles			1
				<1		Chausey archipelago			1
				2.7		Bay of Veys	[17]	2015	1
2.1	2.3	Seine estuary			1				
Pacific oyster (<i>C. gigas</i>)				<0.2*	0.27*	Le Martray			11
				<0.2*		Charente estuary	[18]	2019	5
				<0.2*		Seudre estuary			5
alpha-Endosulfan	OC	33213-65-9	Mediterranean mussel (<i>M. galloprovincialis</i>)	1.5		Chausey archipelago			1
				1.7		Bay of Veys	[17]	2015	1
				1.1	1.4	Seine estuary			1
beta-Endosulfan	OC	959-98-8	Mediterranean mussel (<i>M. galloprovincialis</i>)	<1		Chausey archipelago			1
				14		Bay of Veys	[17]	2015	1
				15.2		Seine estuary			1


Heptachlor	OC	76-44-8	Pacific oyster (<i>C. gigas</i>)	nd	0.3	Bouin	[8]	2008	2
				0.8	1.6	Boyard			2
				0.6	1.9	Les Palles			2
				nd	0.5	Mus du loup			2
				nd	0.7	Cordouan			2
Heptachlor endo-epoxide	OC	1024-57-3	Mediterranean mussel (<i>M. galloprovincialis</i>)	4.1		Chausey archipelago	[17]	2015	1
					1.1	Bay of Veys			1
				1.4	2.3	Seine estuary			1
Heptachlor exo-epoxide	OC	1024-57-3	Mediterranean mussel (<i>M. galloprovincialis</i>)	1.0		Chausey archipelago	[17]	2015	1
					2.3	Bay of Veys			1
				1.1	1.2	Seine estuary			1
Lindane	OC	58-89-9	Pacific oyster (<i>C. gigas</i>)	2.1	2.2	Bouin	[8]	2008	2
				0.5	0.6	Boyard			2
				0.4	1.3	Les Palles			2
				nd	0.6	Mus du loup			2
				nd	0.7	Cordouan			2
alpha-BHC	OC	319-84-6	Pacific oyster (<i>C. gigas</i>)	0.6*	0.62*	Le Martray	[18]	2019	11
					0.62*	Charente estuary			5
					0.72*	Seudre estuary			5
					0.26*	Le Martray			5
					0.31*	Charente estuary			4
Metazachlor	OC	67129-08-2	Mediterranean mussel (<i>M. galloprovincialis</i>)	12.3		Chausey archipelago	[17]	2015	1
					10.8	Bay of Veys			1
				11.8	24.0	Seine estuary			1
Metolachlor	OC	51218-45-2	Mediterranean mussel (<i>M. galloprovincialis</i>)	398		Chausey archipelago	[17]	2015	1
					151	Bay of Veys			1

				284	349	Seine estuary			1
			Pacific oyster (<i>C. gigas</i>)	<0.33*	0.38*	Le Martray			11
				<0.33*		Charente estuary	[18]	2019	5
				<0.33*		Seudre estuary			5
Phtalimide	OC	-	Blue mussel (<i>M. edulis</i>)		30	Banc du Nord			8
						Bay of Bourgneuf	[19]	2015-2017	6
						Bay of Aiguillon			12
trans-Nonachlor	OC	5103-73-1	Pacific oyster (<i>C. gigas</i>)	nd	0.6	Bouin			2
				0.4	0.7	Cordouan	[8]	2008	2
Ethoprophos	OP	13194-48-4	Blue mussel (<i>M. edulis</i>)		31	Banc du Nord			8
						Bay of Aiguillon	[19]	2015-2017	12
Chlorfenvinphos	OP	470-90-6	Pacific oyster (<i>C. gigas</i>)	1.65*	1.89*	Le Martray			11
					1.72*	Charente estuary	[18]	2019	5
					2.12*	Seudre estuary			5
Chlorpyrifos	OP	2921-88-2	Pacific oyster (<i>C. gigas</i>)	0.79*	0.80*	Le Martray			11
					0.81*	Charente estuary	[18]	2019	5
					0.93*	Seudre estuary			5
Parathion	OP	56-38-2	Pacific oyster (<i>C. gigas</i>)	0.56*	0.69*	Le Martray			11
					0.63*	Charente estuary	[18]	2019	5
					0.61*	Seudre estuary			5
Tributyl phosphate	OP	126-73-8	Blue mussel (<i>M. edulis</i>)		14	Bay of Bourgneuf			6
						Bay of Aiguillon	[19]	2015-2017	12
Bifenthrin	Pyr	82657-04-3	Blue mussel (<i>M. edulis</i>)		48	Bay of Bourgneuf			6
						Bay of Aiguillon	[19]	2015-2017	12
Cypermethrin	Pyr	52315-07-8	Mediterranean mussel (<i>M. galloprovincialis</i>)		10.9	Chausey archipelago			1
				<10		Bay of Veys	[17]	2015	1
				<10		Seine estuary			1

Deltamethrin	Pyr	52918-63-5	Mediterranean mussel (<i>M. galloprovincialis</i>)	<10	11.2	13.0	Chausey archipelago Bay of Veys Seine estuary	[17]	2015	1 1 1
--------------	-----	------------	--	-----	------	------	---	------	------	-------------

804

805 Figure 1.



806

807

808 **Supplementary material.**

Pesticide	Class	Statute in Europe	Log Kow	K_{oc} (L/kg)	Water solubility (mg/L)	Vapor pressure at 25°C (mPa)	DT₅₀ soil (days)	PNEC (mg/L)
------------------	--------------	----------------------------------	--------------------	----------------------------------	--	---	--	------------------------

809

810 Table S1. Physicochemical properties of pesticides found in bivalves from the French Atlantic
811 and English Channel coasts.

812

813

814

Acetochlor	OC	Banned	4.14	208.93	233	0.22 (20 °C)	20–100	2.6.10 ⁻⁶
Aldrine	OC	Banned	6.50	49,000	0.027	0.12	9490	-
DDT	OC	Banned	6.91	204,000	0.0055	2.1.10 ⁻¹¹	4–30	-
Dieldrin	OC	Banned	5.40	12,000	0.195	0.78 .10 ⁻³	>7	-
Endosulfan	OC	Banned	3.83 (alpha); 3.62 (beta)	11,500 (1.06)	0.32	0.83	50	5.10 ⁻⁷
Heptachlor	OC	Banned	6.10	24,000	0.18	4.0.10 ⁻⁴	250	3.10 ⁻⁸
Lindane	OC	Banned	3.72	1,480	7.3	4.5.10 ⁻⁵	15	2.10 ⁻⁶
Metazachlor	OC	Banned	2.49	-	450	9.10 ⁻⁷ (20 °C)	0.1–25.3	2.10 ⁻⁶
Metolachlor	OC	Banned	3.13	288.40	488	4.18.10 ⁻⁴	11-31	-
Phtalimide	OC		1.15	-	360	1.33.10 ⁻⁴		-
trans-Nonachlor	OC	Banned	6.20	56,000	8.2e-3	1.26.10 ⁻¹⁰	-	-
Ethoprophos	OP	Banned	3.59	70.79	750	0.05	3-56	-
Tributyl phosphate	OP	-	4.0	-	280	1.5.10 ⁻⁷	-	0.0082
Bifenthrin	Pyr	Banned	6.00	236,610	0.001	0.0178	26	1.9.10 ⁻⁹
Cypermethrin	Pyr	Approved	6.60	156,250 (2.19)	0.009	0.00023	60	-
Deltamethrin	Pyr	Approved	6.20	10,240,000	0.0002	1.24.10 ⁻⁵	13	1.10 ⁻⁷
Naphtalene	PAH	Banned	3.30	1.1.10 ⁻⁵	31	1.1.10 ⁻⁵	-	0.000325

815

816

817

818 Text S1. OCs are synthetic organic compounds containing at least one chlorine atom [87].
819 Due to their hydrophobic nature, OCs are persistent, i.g., dichlorodiphenyltrichloroethane
820 (DDT) with a Log Kow of 6.91 [88] . They tend to accumulate in sediments, bioaccumulate in
821 human and animal tissue, and biomagnify in food chains. OCs are considered as persistent
822 organic pollutants (POPs) (DT 50 up to 9490 days) [6,51,89,90]. From 1940 to the 1970s,
823 they were widely used, before being formally banned [91].

824 OPs and Carb – second generation insecticides – act similarly. They inhibit the
825 acetylcholinesterase (AChE) by releasing esters. AChE is an enzyme that catalyzes the
826 hydrolysis of the neurotransmitter acetylcholine (ACh) [87,92]. Therefore, OPs and Carb are
827 highly toxic to insects and mammals [87,92] and are as persistent in the environment as OCs
828 [87]. For instance, ethoprophos has a 81-day half-life of open seawater [93]. OPs and Carb,
829 including diazinon, parathion-methyl and carbofuran are water-soluble [13].

830 Pyr and Neo – third-generation of pesticides – were respectively produced in the 1970/80s
831 and 1990s [11,13,94]. They have a lower bioaccumulation capacity than OCs [95] . However,
832 these pesticides are extremely toxic to non-target aquatic invertebrates and fish species [96] .
833 Pyr are characterized by their low solubility in water and their high organic carbon-water
834 partition coefficients (K_{OC}) (see Log Kow values and Koc in Table S1) [13]. Finally, Neo are
835 water-soluble [13].