



# New observation of juvenile *Stichopus herrmanni* within a dense population of mixed ages in Vanuatu

Frédéric Ducarme

## ► To cite this version:

Frédéric Ducarme. New observation of juvenile *Stichopus herrmanni* within a dense population of mixed ages in Vanuatu. SPC Beche-de-mer Information Bulletin, 2023, 43, pp.76-78. hal-04078782

**HAL Id: hal-04078782**

**<https://hal.science/hal-04078782>**

Submitted on 24 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# New observation of juvenile *Stichopus herrmanni* within a dense population of mixed ages in Vanuatu

Frédéric Ducarme<sup>1</sup>

**Species:** *Stichopus herrmanni* (Semper, 1868; syn. *S. variegatus*), also known as “curryfish”.

**Context:** *Stichopus herrmanni* is one of the increasingly targeted beche-de-mer species in the Indo-Pacific (Purcell et al. 2012; Di Simone et al. 2022), and listed as Vulnerable (A2bd) by the International Union for Conservation of Nature since 2013 (Conand et al. 2014). Hence, improving the knowledge on its biology, particularly its ecology, reproduction and growth is paramount (Wolfe and Byrne 2017a). Although considered scarce in many areas, this species is also known to occur in high densities in some places, especially inside some marine protected areas (Eriksson et al. 2010).

Juvenile beche-de-mer (Holothuroidea) are rarely observed and particularly difficult to identify because many species exhibit very different morphologies and color patterns from the adults, hence constituting a challenge for scientists (Shiell 2004). Observations of large, single-species aggregations of individuals of mixed ages, sizes and morphological features can enable a better understanding of these variations and provide better *in situ* identification of isolated juveniles.

**Place:** This observation took place at the “waterfront” of Port Vila, the capital and main city of Vanuatu (South Pacific) during a survey of beche-de-mer (see Ducarme et al. 2023, p.4 in this issue). It was observed on the shallow reef at the north of Port Vila’s main bay, about 100–300 m west of Kumul Highway, in depths of 1–3 m (approximate coordinates 17°43’56.2”S, 168°18’33.1”E). This area is considered polluted because it is in the center of a Port Vila with a population of 60,000 inhabitants. Port Vila has very little in the way of sewage treatment, and the water is often milky or murky. The bottom is a detritic sandy bay with dead coral, sparse live reef elements, and an irregular rubble matrix. The water receives a high nutrient input due to its proximity to the city and a small estuary. This site appears quite different from previously reported habitats for juveniles (i.e. seagrass beds) as reported on in Conand (1993), “coralline algae and associated bacterial films” in Eriksson et al. (2013) and “piecrust reef” in Wolfe and Desbiens (2022).

**Date:** 13 April 2020

**Observation:** A couple of hundred meters from the seashore at shallow depths, we observed a dense population of *Stichopus herrmanni* exhibiting many different sizes (hence ages), ranging from 6 cm to 40 cm in length. The smallest specimens had various mottled colour patterns, mostly yellowish-brown with dark or light areas, and black wrinkles. Juveniles also had more elevated papillae than adults (making them looking like species from the *Stichopus horrens* complex), with the same distinctive concentric wrinkles around them. The largest specimens tended to have the smoothest tegument, and uniform colouration. These observations are consistent with the appearance of juveniles reported from New Caledonia by Conand (1993) and Réunion Island by Bourjon and Morcel (2016).

The size of the smaller specimens suggests an age of one or two months if the growth rate of these are similar to the ones described by Hu and al. (2010), hence likely born from a summer spawning event, consistent with Conand (1993) and Wolfe and Byrne (2017b).

The co-occurrence of juveniles and large adults is surprising as this species is known to experience ontogenic migrations towards deeper waters (Conand 1993; Palazzo et al. 2016; Wolfe and Byrne 2017b). Some ecological parameters (such as food availability) may prevent this behaviour from occurring here.

This population was part of a densely populated ecosystem, including important populations of synaptids (*Synapta maculata* and *Opheodesoma* sp.), as well as other sea cucumber species (*Holothuria hilla* and *H. difficilis*, along with scarcer *H. atra*, *H. coluber* and *Chiridota* sp.). Apart from sea cucumbers, the bottom was also densely populated by other echinoderms, especially the sea urchins *Mespilia globulus*, *Parasalenia gratiosa* and *Laganum laganum*, and diverse ophiotrichid brittle stars. *Stichopus chloronotus* was not observed around the site, although this species was associated with juvenile *S. herrmanni* in Wolfe and Desbiens (2022).

<sup>1</sup> Centre d’Ecologie et des Sciences de la Conservation (CESCO, UMR 7204), Muséum national d’Histoire naturelle, Paris, France. Frederic.Ducarme@gmail.com





One large adult and four juvenile *Stichopus herrmanni*.



A small *Stichopus herrmanni*.

## References

- Bourjon P. and Morcel E. 2016. New observations of holothurian juveniles on Réunion reefs. SPC Beche-de-mer Information Bulletin 36:41–44. <https://purl.org/spc/digilib/doc/qttxq>
- Conand C. 1993. Ecology and reproductive biology of *Stichopus variegatus* an Indo-Pacific coral reef sea cucumber (Echino-dermata: Holothuroidea). Bulletin of Marine Science 52:970–981.
- Conand C., Polidoro B.A., Mercier A., Gamboa R.U., Hamel J-F and Purcell S.W. 2014. The IUCN Red List assessment of aspidochirotid sea cucumbers and its implications. SPC Beche-de-mer Information Bulletin 34:3–7. <https://purl.org/spc/digilib/doc/mhke9>
- Di Simone M., Horellou A., Ducarme F. and Conand C. 2022. Identification guide – Commercial sea cucumbers. Paris : CITES/PatriNat. <https://cites.org/eng/virtual-college/guide-didentification-concombres-de-mer-commercialises>
- Ducarme F., Dumas P. and Kaku R. 2023. Field inventory and population assessment of sea cucumbers from Vanuatu. SPC Beche-de-mer Information Bulletin 43:4–10.
- Eriksson H., Fabricius-Dyg J., Lichtenberg M., Perez-Landa V. and Byrne M. 2010. Biology of a high-density population of *Stichopus herrmanni* at One Tree Reef, Great Barrier Reef, Australia. SPC Beche-de-Mer Information Bulletin 30:41–45. <https://purl.org/spc/digilib/doc/y2tgd>
- Eriksson H., Thorne B.V. and Byrne M. 2013. Population metrics in protected commercial sea cucumber populations (curryfish: *Stichopus herrmanni*) on One Tree Reef, Great Barrier Reef. Marine Ecology Progress Series 473:225–234, doi: 10.3354/meps10054
- Hu C., Xu Y., Wen J., Zhang L.P., Fan S. and Su T. 2010. Larval development and juvenile growth of the sea cucumber *Stichopus* sp. (Curry fish). Aquaculture 300:73–79. 10.1016/j.aquaculture.2009.09.033
- Palazzo L., Wolfe K. and Byrne M. 2016. Discovery and description of *Stichopus herrmanni* juvenile nursery sites on Heron Reef, Great Barrier Reef. SPC Beche-de-mer Information Bulletin 36:36–40. <https://purl.org/spc/digilib/doc/cv2zh>
- Purcell S.W., Samyn Y. and Conand C. 2012. Commercially important sea cucumbers of the world, FAO Species Catalogue for Fishery Purposes, No. 6. 233 p. Rome: FAO.
- Shiell G. 2004. Field observations of juvenile sea cucumbers. SPC Beche-de-mer Information Bulletin 20:6–11. <https://purl.org/spc/digilib/doc/3yr9j>
- Wolfe K. and Byrne M. 2017a. Population biology and recruitment of a vulnerable sea cucumber, *Stichopus herrmanni*, on a protected reef. Marine Ecology 38:e12397. <https://doi.org/10.1111/maec.12397>
- Wolfe K. and Byrne M. 2017b. Biology and ecology of the vulnerable holothuroid, *Stichopus herrmanni*, on a high-latitude coral reef on the Great Barrier Reef. Coral Reefs 36:1143–1156. <https://doi.org/10.1007>
- Wolfe K. and Desbiens A. 2022. Observations of juvenile *Stichopus herrmanni* and confirmation of “pie crust” nursery grounds. SPC Beche-de-mer Information Bulletin 42:93–94. <https://purl.org/spc/digilib/doc/ee9d3>