



# Crystal structure and Hirshfeld analysis of trans -diiodidobis[(methylsulfanyl)benzene- $\kappa$ S ]platinum(II)

Annika Schmidt, Isabelle Jourdain, Michael Knorr, Carsten Strohmann

## ► To cite this version:

Annika Schmidt, Isabelle Jourdain, Michael Knorr, Carsten Strohmann. Crystal structure and Hirshfeld analysis of trans -diiodidobis[(methylsulfanyl)benzene- $\kappa$  S ]platinum(II). Acta crystallographica Section E: Crystallographic communications [2015-..], 2023, 79 (5), pp.516-520. 10.1107/S2056989023003717 . hal-04086541

**HAL Id: hal-04086541**

**<https://hal.science/hal-04086541>**

Submitted on 3 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Crystal structure and Hirshfeld analysis of *trans*-diiodidobis[(methylsulfanyl)benzene- $\kappa$ S]-platinum(II)

Annika Schmidt,<sup>a</sup> Isabelle Jourdain,<sup>b</sup> Michael Knorr<sup>b</sup> and Carsten Strohmann<sup>a\*</sup>

Received 5 April 2023

Accepted 24 April 2023

Edited by J. Reibenspies, Texas A & M University, USA

**Keywords:** crystal structure; Hirshfeld surface analysis; dithioether.

**CCDC reference:** 2258409

**Supporting information:** this article has supporting information at journals.iucr.org/e


<sup>a</sup>TU Dortmund University, Institute for Inorganic Chemistry, Otto-Hahn-Strasse 6, 44227 Dortmund, Germany, and <sup>b</sup>Institut UTINAM CNRS UMR 6213, Equipe "Matériaux et Surfaces Fonctionnels", Université de Franche-Comté, Faculté des Sciences et des Techniques La Bouloie - 16 Route de Gray, 25030 BESANÇON CEDEX, France. \*Correspondence e-mail: carsten.strohmann@tu-dortmund.de

The title complex, [PtI<sub>2</sub>(C<sub>7</sub>H<sub>8</sub>I<sub>2</sub>)<sub>2</sub>], represents a further example of a square-planar Pt<sup>II</sup>-dithioether complex. It crystallizes in the monoclinic space group *P*2<sub>1</sub>/*c*. Additional Hirshfeld analyses indicate a C—H... $\pi$  interaction along the [010] axis to be the most important packing factor.

## 1. Chemical context

Dithioethers are a quite useful class of ligands for various transition-metal complexes and their coordination chemistry is well documented (Murray & Hartley, 1981). As a result of the soft character of the sulfur center, they preferably bond to soft transition metals like the coinage metals (Cu, Ag, Au), mercury(II), or catalytically active noble metals such as rhodium(I), iridium(I), palladium(II) or platinum(II). Apart from structural aspects (Marangoni *et al.*, 1995) and the investigation of inversion dynamics occurring at the coordinated sulfur atoms (Abel *et al.*, 1984), these complexes have been reported to have several applications in homogeneous catalysis (Masdeu-Bulto *et al.*, 2003; Arrayás & Carretero, 2011). They can form interesting luminescent cluster-like structures (Knorr *et al.*, 2014; Peindy *et al.*, 2007) and even coordination polymers by coordination to Cu<sup>I</sup> and Ag<sup>I</sup> (Raghuvanshi *et al.*, 2017; Awaleh *et al.*, 2006). Depending on the metal coordination sphere and the remaining ligands, the preparation of dithioether complexes may yield different isomers. In particular, the isomerism of chalcogenoether complexes with palladium and platinum has been intensively investigated (Vigo *et al.*, 2006) and the presence of both *trans*- and *cis*-isomers in solution and the solid state were proven. The clarification of the *trans*–*cis* isomerism is therefore of importance.

In the past, our groups have investigated the coordination of chelating dithioethers such as the vinylic ferrocenyl-dithioether Z-[(ArS)(H)C=C(SAr)-Fc] or the silylated compounds (PhSCH<sub>2</sub>)<sub>2</sub>SiPh<sub>2</sub> and PhSCH<sub>2</sub>Si(Me)-Si(Me)-CH<sub>2</sub>SPh<sub>2</sub> yielding [Fc-{C(S-*p*-tolyl)=C(S-*p*-tolyl)(H)}PtCl<sub>2</sub>], *cis*-[PtCl<sub>2</sub>{(PhSCH<sub>2</sub>)<sub>2</sub>SiMe<sub>4</sub>}] and *cis*-[PtCl<sub>2</sub>{(PhSCH<sub>2</sub>)<sub>2</sub>-SiPh<sub>2</sub>}] and converted them *via* metathesis in the presence of NaI to their corresponding diiodo derivatives (Clement *et al.*, 2007; Knorr *et al.*, 2004; Peindy *et al.*, 2006). We have also shown that the tetrakis(thioether) (PhSCH<sub>2</sub>)<sub>4</sub>Si can be ligated



OPEN ACCESS

Published under a CC BY 4.0 licence





Figure 1  
Synthesis scheme for *trans*-PtI<sub>2</sub>(SMePh)<sub>2</sub> (**1**).

on HgBr<sub>2</sub> in a chelating manner (Peindy *et al.*, 2005). In a similar manner, we also prepared, as shown in Fig. 1, the complex *cis*-[PtI<sub>2</sub>{(PhSCH<sub>2</sub>)<sub>2</sub>Si(CH<sub>2</sub>SPh)<sub>2</sub>}. When attempting to recrystallize this poorly soluble compound from hot toluene, partial cleavage of the Si—CH<sub>2</sub>Ph bond occurred, yielding *trans*-[PtI<sub>2</sub>(SMePh)<sub>2</sub>] **1**, albeit in a quite low yield of 10%. Alternatively, this air-stable complex could be prepared in a much improved yield of 80% by reaction of bis(benzonitrile)diodoplatinum with 2 equivalents of methyl phenyl sulfide (thioanisole) MeSPh using dichloromethane as solvent. This compound was characterized by NMR spectroscopy in solution and exhibits a singlet resonance for the two magnetically equivalent methyl groups at  $\delta$  3.01 ppm, flanked by <sup>195</sup>Pt satellites due to a <sup>3</sup>J<sub>PtH</sub> coupling of 48 Hz. Furthermore, we report herein on the solid-state structure and structural analysis of *trans*-diiodidobis[(methylsulfanyl)benzene- $\kappa$ S]-platinum(II) (**1**). In addition, the results of a Hirshfeld analysis of the intermolecular interactions are presented.



## 2. Structural commentary

*trans*-Diiodidobis[(methylsulfanyl)benzene- $\kappa$ S]platinum(II) (**1**) crystallizes from dichloromethane in the monoclinic crystal system, space group *P*2<sub>1</sub>/*c*. The molecular structure of **1** is

Table 1  
Selected geometric parameters (Å, °).

|                         |             |           |             |
|-------------------------|-------------|-----------|-------------|
| Pt1—I1                  | 2.6121 (2)  | S1—C2     | 1.782 (2)   |
| Pt1—S1                  | 2.3183 (5)  | S1—C1     | 1.800 (2)   |
| I1 <sup>i</sup> —Pt1—I1 | 180.0       | S1—Pt1—I1 | 85.641 (14) |
| S1—Pt1—I1 <sup>i</sup>  | 94.359 (14) | C2—S1—C1  | 103.46 (11) |

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .




Figure 2  
Molecular structure of **1** in the unit cell.

presented in Figs. 2 and 3 and selected bond lengths and bond angles are given in Table 1. The asymmetric unit contains half a molecule of **1**, which shows *C*<sub>2h</sub> symmetry. The distance from the coordinating iodine center I1 to Pt1 is 2.61205 (15) Å, showing a slight elongation with respect to its educt structure




Figure 3  
Asymmetric unit of **1** with labeled atoms.


*trans*-[PtI<sub>2</sub>(NCPh)<sub>2</sub>] (**2**) (2.6052 (8) Å; Viola *et al.*, 2018). The distance from the coordinating sulfur atom S1 to Pt1 is 2.3183 (5) Å. The S1—Pt1 bond is 0.015 Å longer than in the analogous chlorine compound *trans*-[PtCl<sub>2</sub>(SMePh)<sub>2</sub>] (**3**) reported by Ahlgrén (CSD LEQSUW; Vigo *et al.*, 2006). This elongation may be explained by the thermodynamic *trans*-effect of the opposite halide ligand. Therefore, similar compounds with iodo ligands such as *trans*-[PtI<sub>2</sub>(SMe<sub>2</sub>)<sub>2</sub>] (**4**) (CSD RAYNOU; Löqvist *et al.*, 1996) and *trans*-[PtI<sub>2</sub>(tetrahydrothiophene)<sub>2</sub>] (**5**) (CSD SIRPAK; Oskarsson *et al.*, 1990) show S—Pt bond lengths in the same range at 2.310 (2) and 2.310 (1) Å, respectively. Both complexes also have similar bond lengths for the Pt—I bond [2.6039 (8) Å in **4** and 2.606 (1) Å in **5**]. The chelate complexes *cis*-diiodo-[1,2-bis-(phenylsulfanyl)ethane]platinum(II) (CSD ZAJWUC; Marangoni *et al.*, 1995) and *cis*-(1,4-dithiane-*S,S'*)diiodoplatinum(II) (CSD HUFQAA; Johansson & Engelbrecht, 2001) are reported to display Pt—I bond lengths of 2.606 (1) and 2.6035 (5) Å, respectively, and somewhat shorter mean Pt—S bond lengths of 2.265 (2) and 2.2751 (16) Å, respectively.

All further bonds have characteristic dimensions (Allen *et al.*, 1987). The coordination sphere around the platinum center is square-planar. The angles I1—Pt1—I1 and S1—Pt1—


S1 are 180°. However, the angle I1—Pt1—S1 of 85.641 (14)° is somewhat more acute. This slight deviation from the ideal angle of 90° is also reported for the chlorido derivative **3** and the dimethyl sulfide analog **4**, as well as in the tetrahydrothiophene analog **5**. The sulfur center shows a distorted tetrahedral environment with angles C1—S1—Pt1 = 111.00 (8)°, C2—S1—Pt1 = 104.52 (7)° and C2—S1—C1 = 103.46 (11)°.

### 3. Supramolecular features

While a repetition of the molecular structure of **1** can be seen along the [100] axis and the [001] axis, as shown in Fig. 4, the crystal packing along the [010] axis is defined by C—H... $\pi$  interactions of the C2—C7 phenyl ring and H1B<sup>i</sup> [symmetry code: (i)  $x, \frac{1}{2} - y, -\frac{1}{2} + z$ ] with a distance between the phenyl ring and H1B<sup>i</sup> of 2.5377 (10) Å (Fig. 5). This interaction can also be visualized by a Hirshfeld surface analysis (Spackman & Jayatilaka, 2009) generated by *CrystalExplorer21* (Spackman *et al.*, 2021). The Hirshfeld surface mapped over  $d_{\text{norm}}$  in the range from -0.0074 to 1.1829 a.u. is shown in Fig. 6, with the close contact between H1B<sup>i</sup> and the C2—C7 plane indicated by the red spot. The contributions of the different types of intermolecular interactions for **1** are shown in the two-dimensional fingerprint plots (McKinnon *et al.*, 2007) in Fig. 7. The contribution of the H...H interactions, with a value of 39.8%, has the largest share of the crystal packing of **1**. The remaining hydrogen–heteronuclear interactions have a smaller share with a 15.7% contribution for I...H, a 14.4% contribution for C...H and a 3.6% contribution for S...H.




**Figure 4**  
The packing of the solid-state structure of **1** along the [100] axis.




**Figure 5**  
The packing of the solid-state structure of **1** along the [010] axis [symmetry code: (i)  $x, \frac{1}{2} - y, -\frac{1}{2} + z$ ].





**Figure 6**  
Hirshfeld surface analysis of **1** showing close contacts in the crystal. The  $\pi$ -interaction between hydrogen atom H1b' and the phenyl ring C2–C7 is indicated by the red spot [symmetry code: (i)  $x, \frac{1}{2} - y, -\frac{1}{2} + z$ ].

The heteronuclear  $I \cdots H$  and  $C \cdots H$  interactions appear as spikes.



**Figure 7**  
Two-dimensional fingerprint plots for compound **1**, showing (a) all contributions, and (b)–(d) delineated into the contributions of atoms within specific interacting pairs (blue areas).

#### 4. Database survey

By a search in the Cambridge Crystallographic Database (WebCSD, November 2022; Groom *et al.*, 2016), various structures of dihalide transition-metal complexes with the same ligand motif as **1** were found. To compare the most similar structures, only dihalide transition metal complexes with the bis[(methylsulfanyl)benzene] ligand and its oxidized derivative are focused on now. The already compared structure *trans*-dichloro-bis[methyl(phenyl)sulfanyl]platinum (LEQSUW; Vigo *et al.*, 2006) has been published, as well as its palladium derivative with (SARWEP; Oilunkaniemi *et al.*, 2006) and without (LEQSOQ; Vigo *et al.*, 2006) inserted benzene. In addition, *cis*-dichlorobis(methylphenylsulfoxide)-palladium has been published independently by two different research groups [JISWUD (Antolini *et al.*, 1991) and JISWUD01 (Gama de Almeida *et al.*, 1992)]. Further examples of  $PtI_2$  thioether complexes are *cis*-diiodo-(1,4,7-trithia-cyclononane-*S,S'*)platinum(II) (ACUXAX; Grant *et al.*, 2001), diiodo-(2,9-dimethyl-1,10-phenanthroline)(dimethylsulfide)-platinum(II) (BERTIC; Fanizzi *et al.*, 2004), and diiodo-(5-phenyl-1-thia-5-phosphacyclo-octane-*P,S*)platinum(II) (KEJHEM; Toto *et al.*, 1990).

Similar complexes were also structurally characterized by our research groups and include *cis*-[ $PtBr_2\{(PhSCH_2)_2SiPh_2\}$ ] (ECOHAG; Knorr *et al.*, 2004) and *cis*-[ $PtI_2\{(PhSCH_2)_2SiPh_2\}$ ].DCM (ECOHIO, Knorr *et al.*, 2004), which were determined in order to investigate the *trans*-influence of different halide ligands on the Pt–S bond. Further examples of dithioether complexes stemming from our laboratories are *cis*-[ $PtCl_2\{(PhSCH_2)_2Si_2Me_4\}$ ] (MEDYOK; Peindy *et al.*, 2006) and *cis*-[ $PtI_2\{(PhSCH_2)_2Si_2Me_4\}$ ].DCM (MEDZIF; Peindy *et al.*, 2006).

#### 5. Synthesis and crystallization

*trans*-Diiodobis[(methylsulfanyl)benzene]platinum (**1**) was synthesized by adding methylphenyl sulfide (37 mg, 0.30 mmol, 1.50 eq.) dissolved in 0.5 mL of dichloromethane *via* a microsyringe to a solution of bis(benzonitrile)diiodo-platinum (65 mg, 0.10 mmol, 1.00 eq.) in dichloromethane (3 mL) and stirring overnight at room temperature. *trans*-Diiodobis[(methylsulfanyl)benzene]platinum (**1**, 557 mg, 0.80 mmol, 80%) was isolated as red crystals after layering with heptane.

Calculated for  $C_{14}H_{16}I_2PtS_2$  (697.30 g mol<sup>−1</sup>): C, 24.11; H, 2.32; S, 9.20. Found: C, 23.92; H, 2.21; S, 9.05%.

<sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  = 3.01 (*s*, <sup>3</sup>*J*<sub>PtH</sub> = 48 Hz, 6H; CH<sub>3</sub>), 7.05–7.73 (*m*, 10H; phenyl) ppm.

#### 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms were positioned geometrically (C–H = 0.95–1.00 Å) and were refined using a riding model, with  $U_{iso}(H) = 1.2U_{eq}(C)$  for CH<sub>2</sub> and CH hydrogen atoms and  $U_{iso}(H) = 1.5U_{eq}(C)$  for CH<sub>3</sub> hydrogen atoms.

**Table 2**  
Experimental details.

|                                                                                                                |                                                                    |
|----------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|
| Crystal data                                                                                                   |                                                                    |
| Chemical formula                                                                                               | [PtI <sub>2</sub> (C <sub>7</sub> H <sub>8</sub> I) <sub>2</sub> ] |
| <i>M<sub>r</sub></i>                                                                                           | 697.28                                                             |
| Crystal system, space group                                                                                    | Monoclinic, <i>P</i> <sub>2</sub> <sub>1</sub> / <i>c</i>          |
| Temperature (K)                                                                                                | 100                                                                |
| <i>a</i> , <i>b</i> , <i>c</i> (Å)                                                                             | 9.5796 (3), 9.5104 (3), 9.7960 (3)                                 |
| β (°)                                                                                                          | 107.645 (1)                                                        |
| <i>V</i> (Å <sup>3</sup> )                                                                                     | 850.48 (5)                                                         |
| <i>Z</i>                                                                                                       | 2                                                                  |
| Radiation type                                                                                                 | Mo <i>K</i> α                                                      |
| μ (mm <sup>−1</sup> )                                                                                          | 12.11                                                              |
| Crystal size (mm)                                                                                              | 0.31 × 0.25 × 0.20                                                 |
| Data collection                                                                                                |                                                                    |
| Diffractometer                                                                                                 | Bruker APEXII CCD                                                  |
| Absorption correction                                                                                          | Multi-scan ( <i>SADABS</i> ; Krause <i>et al.</i> , 2015)          |
| <i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub>                                                              | 0.263, 0.498                                                       |
| No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections                             | 27554, 4125, 4033                                                  |
| <i>R</i> <sub>int</sub>                                                                                        | 0.038                                                              |
| (sin θ/λ) <sub>max</sub> (Å <sup>−1</sup> )                                                                    | 0.833                                                              |
| Refinement                                                                                                     |                                                                    |
| <i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i> | 0.021, 0.053, 1.17                                                 |
| No. of reflections                                                                                             | 4125                                                               |
| No. of parameters                                                                                              | 89                                                                 |
| H-atom treatment                                                                                               | H-atom parameters constrained                                      |
| Δρ <sub>max</sub> , Δρ <sub>min</sub> (e Å <sup>−3</sup> )                                                     | 2.26, −1.80                                                        |

Computer programs: *APEX2* and *SAINT* V8.38A (Bruker, 2018), *SHELXT2014/5* (Sheldrick, 2015a), *SHELXL2019/2* (Sheldrick, 2015b), *OLEX2* (Dolomanov *et al.*, 2009) and *publCIF* (Westrip, 2010).

## References

- Abel, E. W., Bhargava, S. K. & Orrell, K. G. (1984). *Progr. Inorg. Chem.* **32**, 1–118.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Almeida, S. G. de, Hubbard, J. L. & Farrell, N. (1992). *Inorg. Chim. Acta*, **193**, 149–157.
- Antolini, L., Folli, U., Iarossi, D., Schenetti, L. & Taddei, F. (1991). *J. Chem. Soc. Perkin Trans. 2*, pp. 955–961.
- Arrayás, R. G. & Carretero, J. C. (2011). *Chem. Commun.* **47**, 2207–2211.
- Awaleh, M. O., Badia, A., Brisse, F. & Bu, X.-H. (2006). *Inorg. Chem.* **45**, 1560–1574.
- Bruker (2018). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clement, S., Guyard, L., Knorr, M., Villafane, F., Strohmman, C. & Kubicki, M. M. (2007). *Eur. J. Inorg. Chem.* pp. 5052–5061.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Fanizzi, F. P., Margiotta, N., Lanfranchi, M., Tiripicchio, A., Pacchioni, G. & Natile, G. (2004). *Eur. J. Inorg. Chem.* pp. 1705–1713.
- Grant, G. J., Brandow, C. G., Galas, D. F., Davis, J. P., Pennington, W. T., Valente, E. J. & Zubkowski, J. D. (2001). *Polyhedron*, **20**, 3333–3342.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst. B* **72**, 171–179.
- Johansson, M. H. & Engelbrecht, H. P. (2001). *Acta Cryst. E* **57**, m114–m116.
- Knorr, M., Khatyr, A., Dini Aleo, A., El Yaagoubi, A., Strohmman, C., Kubicki, M. M., Rousselin, Y., Aly, S. M., Fortin, D., Lapprand, A. & Harvey, P. D. (2014). *Cryst. Growth Des.* **14**, 5373–5387.
- Knorr, M., Peindy, H. M., Guyon, F., Sachdev, H. & Strohmman, C. (2004). *Z. Anorg. Allg. Chem.* **630**, 1955–1961.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.
- Löqvist, K. C., Wendt, O. F. & Leipoldt, J. G. (1996). *Acta Chem. Scand.* **50**, 1069–1073.
- Marangoni, G., Pitteri, B., Bertolasi, V. & Gilli, P. (1995). *Inorg. Chim. Acta*, **234**, 173–179.
- Masdeu-Bultó, A. M., Diéguez, M., Martín, E. & Gómez, M. (2003). *Coord. Chem. Rev.* **242**, 159–201.
- McKinnon, J. J., Jayatilaka, D. & Spackman, M. A. (2007). *Chem. Commun.* pp. 3814–3816.
- Murray, S. G. & Hartley, F. R. (1981). *Chem. Rev.* **81**, 365–414.
- Oilunkaniemi, R., Laitinen, R. S. & Ahlgrén, M. (2006). *Main Group Chem.* **5**, 125–136.
- Oskarsson, Å., Norén, B., Svensson, C. & Elding, L. I. (1990). *Acta Cryst. B* **46**, 748–752.
- Peindy, H. M., Guyon, F., Jourdain, I., Knorr, M., Schildbach, D. & Strohmman, C. (2006). *Organometallics*, **25**, 1472–1479.
- Peindy, H. M., Guyon, F., Khatyr, A., Knorr, M. & Strohmman, C. (2007). *Eur. J. Inorg. Chem.* pp. 1823–1828.
- Peindy, H. N., Guyon, F., Knorr, M., Smith, A. B., Farouq, J. A. A., Islas, S. A., Rabinovich, D., Golen, J. A. & Strohmman, C. (2005). *Inorg. Chem. Commun.* **8**, 479–482.
- Raghuvanshi, A., Strohmman, C., Tissot, J.-B., Clément, S., Mehdi, A., Richeter, S., Viau, L. & Knorr, M. (2017). *Chem. Eur. J.* **23**, 16479–16483.
- Sheldrick, G. M. (2015a). *Acta Cryst. A* **71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst. C* **71**, 3–8.
- Spackman, M. A. & Jayatilaka, D. (2009). *CrystEngComm*, **11**, 19–32.
- Spackman, P. R., Turner, M. J., McKinnon, J. J., Wolff, S. K., Grimwood, D. J., Jayatilaka, D. & Spackman, M. A. (2021). *J. Appl. Cryst.* **54**, 1006–1011.
- Toto, S. D., Olmstead, M. M., Arbuckle, B. W., Bharadwaj, P. K. & Musker, K. W. (1990). *Inorg. Chem.* **29**, 691–699.
- Vigo, L., Risto, M., Jahr, E. M., Bajorek, T., Oilunkaniemi, R., Laitinen, R. S., Lahtinen, M. & Ahlgrén, M. (2006). *Cryst. Growth Des.* **6**, 2376–2383.
- Viola, E., Donzello, M. P., Ercolani, C., Rizzoli, C. & Lever, A. B. P. (2018). *Inorg. Chim. Acta*, **480**, 101–107.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2023). E79, 516–520 [https://doi.org/10.1107/S2056989023003717]

## Crystal structure and Hirshfeld analysis of *trans*-diiodidobis[(methylsulfanyl)benzene- $\kappa$ S]platinum(II)

Annika Schmidt, Isabelle Jourdain, Michael Knorr and Carsten Strohmann

### Computing details

Data collection: *APEX2* (Bruker, 2018); cell refinement: *SAINT* V8.38A (Bruker, 2018); data reduction: *SAINT* V8.38A (Bruker, 2018); program(s) used to solve structure: *SHELXT2014/5* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2019/2* (Sheldrick, 2015b); molecular graphics: *Olex2* 1.5 (Dolomanov *et al.*, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

### *trans*-Diiodidobis[(methylsulfanyl)benzene- $\kappa$ S]platinum(II)

#### Crystal data

|                                                                    |                                                         |
|--------------------------------------------------------------------|---------------------------------------------------------|
| [PtI <sub>2</sub> (C <sub>7</sub> H <sub>8</sub> I) <sub>2</sub> ] | $F(000) = 632$                                          |
| $M_r = 697.28$                                                     | $D_x = 2.723 \text{ Mg m}^{-3}$                         |
| Monoclinic, $P2_1/c$                                               | Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$ |
| $a = 9.5796(3) \text{ \AA}$                                        | Cell parameters from 9775 reflections                   |
| $b = 9.5104(3) \text{ \AA}$                                        | $\theta = 3.1\text{--}36.3^\circ$                       |
| $c = 9.7960(3) \text{ \AA}$                                        | $\mu = 12.11 \text{ mm}^{-1}$                           |
| $\beta = 107.645(1)^\circ$                                         | $T = 100 \text{ K}$                                     |
| $V = 850.48(5) \text{ \AA}^3$                                      | Prism, red                                              |
| $Z = 2$                                                            | $0.31 \times 0.25 \times 0.20 \text{ mm}$               |

#### Data collection

|                                                       |                                                            |
|-------------------------------------------------------|------------------------------------------------------------|
| Bruker APEXII CCD                                     | $T_{\min} = 0.263$ , $T_{\max} = 0.498$                    |
| diffractometer                                        | 27554 measured reflections                                 |
| Radiation source: microfocus sealed X-ray tube,       | 4125 independent reflections                               |
| Incoatec I $\mu$ s                                    | 4033 reflections with $I > 2\sigma(I)$                     |
| HELIOS mirror optics monochromator                    | $R_{\text{int}} = 0.038$                                   |
| Detector resolution: $10.4167 \text{ pixels mm}^{-1}$ | $\theta_{\max} = 36.3^\circ$ , $\theta_{\min} = 2.2^\circ$ |
| $\omega$ and $\phi$ scans                             | $h = -15 \rightarrow 15$                                   |
| Absorption correction: multi-scan                     | $k = -15 \rightarrow 15$                                   |
| (SADABS; Krause <i>et al.</i> , 2015)                 | $l = -16 \rightarrow 16$                                   |

#### Refinement

|                                  |                                                   |
|----------------------------------|---------------------------------------------------|
| Refinement on $F^2$              | Hydrogen site location: inferred from             |
| Least-squares matrix: full       | neighbouring sites                                |
| $R[F^2 > 2\sigma(F^2)] = 0.021$  | H-atom parameters constrained                     |
| $wR(F^2) = 0.053$                | $w = 1/[\sigma^2(F_o^2) + (0.0193P)^2 + 1.6484P]$ |
| $S = 1.17$                       | where $P = (F_o^2 + 2F_c^2)/3$                    |
| 4125 reflections                 | $(\Delta/\sigma)_{\max} = 0.002$                  |
| 89 parameters                    | $\Delta\rho_{\max} = 2.26 \text{ e \AA}^{-3}$     |
| 0 restraints                     | $\Delta\rho_{\min} = -1.79 \text{ e \AA}^{-3}$    |
| Primary atom site location: dual |                                                   |

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>    | <i>y</i>    | <i>z</i>    | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|-------------|-------------|-------------|----------------------------------|
| Pt1 | 0.500000    | 0.500000    | 0.500000    | 0.01069 (3)                      |
| I1  | 0.67448 (2) | 0.70668 (2) | 0.48539 (2) | 0.01650 (4)                      |
| S1  | 0.66656 (6) | 0.45396 (6) | 0.72234 (6) | 0.01359 (8)                      |
| C3  | 0.7382 (3)  | 0.1669 (2)  | 0.7476 (2)  | 0.0159 (3)                       |
| H3  | 0.668801    | 0.154647    | 0.798352    | 0.019*                           |
| C7  | 0.8623 (2)  | 0.3176 (2)  | 0.6213 (3)  | 0.0168 (4)                       |
| H7  | 0.876638    | 0.407868    | 0.586351    | 0.020*                           |
| C2  | 0.7595 (2)  | 0.2989 (2)  | 0.6957 (2)  | 0.0134 (3)                       |
| C1  | 0.5743 (3)  | 0.3994 (3)  | 0.8486 (2)  | 0.0188 (4)                       |
| H1A | 0.518020    | 0.478370    | 0.868893    | 0.028*                           |
| H1B | 0.646855    | 0.368646    | 0.937513    | 0.028*                           |
| H1C | 0.507882    | 0.321330    | 0.807968    | 0.028*                           |
| C4  | 0.8199 (3)  | 0.0527 (3)  | 0.7244 (3)  | 0.0200 (4)                       |
| H4  | 0.806058    | −0.037609   | 0.759612    | 0.024*                           |
| C5  | 0.9216 (3)  | 0.0705 (3)  | 0.6501 (3)  | 0.0216 (4)                       |
| H5  | 0.976478    | −0.007512   | 0.633993    | 0.026*                           |
| C6  | 0.9426 (3)  | 0.2037 (3)  | 0.5991 (3)  | 0.0209 (4)                       |
| H6  | 1.012472    | 0.216130    | 0.549005    | 0.025*                           |

*Atomic displacement parameters ( $\text{\AA}^2$ )*

|     | $U^{11}$    | $U^{22}$    | $U^{33}$     | $U^{12}$      | $U^{13}$     | $U^{23}$      |
|-----|-------------|-------------|--------------|---------------|--------------|---------------|
| Pt1 | 0.01172 (5) | 0.00914 (5) | 0.01237 (5)  | 0.00010 (3)   | 0.00539 (3)  | 0.00002 (3)   |
| I1  | 0.01712 (6) | 0.01357 (6) | 0.02033 (7)  | −0.00397 (4)  | 0.00794 (5)  | 0.00006 (4)   |
| S1  | 0.0149 (2)  | 0.0124 (2)  | 0.01360 (19) | −0.00016 (15) | 0.00455 (16) | −0.00055 (15) |
| C3  | 0.0176 (8)  | 0.0150 (8)  | 0.0162 (8)   | 0.0008 (7)    | 0.0067 (7)   | 0.0012 (7)    |
| C7  | 0.0142 (8)  | 0.0170 (9)  | 0.0205 (9)   | −0.0018 (7)   | 0.0073 (7)   | −0.0004 (7)   |
| C2  | 0.0135 (8)  | 0.0132 (8)  | 0.0134 (8)   | −0.0002 (6)   | 0.0039 (6)   | −0.0010 (6)   |
| C1  | 0.0226 (10) | 0.0214 (10) | 0.0147 (8)   | 0.0047 (8)    | 0.0092 (7)   | 0.0018 (7)    |
| C4  | 0.0229 (10) | 0.0164 (9)  | 0.0206 (10)  | 0.0038 (7)    | 0.0063 (8)   | 0.0029 (7)    |
| C5  | 0.0194 (10) | 0.0210 (10) | 0.0244 (11)  | 0.0073 (8)    | 0.0065 (8)   | −0.0001 (8)   |
| C6  | 0.0149 (9)  | 0.0243 (11) | 0.0257 (11)  | 0.0010 (8)    | 0.0094 (8)   | −0.0012 (8)   |

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

|                     |            |        |           |
|---------------------|------------|--------|-----------|
| Pt1—I1              | 2.6121 (1) | C7—C2  | 1.403 (3) |
| Pt1—I1 <sup>i</sup> | 2.6120 (1) | C7—C6  | 1.383 (3) |
| Pt1—S1 <sup>i</sup> | 2.3183 (5) | C1—H1A | 0.9800    |
| Pt1—S1              | 2.3183 (5) | C1—H1B | 0.9800    |



|                                      |              |             |              |
|--------------------------------------|--------------|-------------|--------------|
| S1—C2                                | 1.782 (2)    | C1—H1C      | 0.9800       |
| S1—C1                                | 1.800 (2)    | C4—H4       | 0.9500       |
| C3—H3                                | 0.9500       | C4—C5       | 1.392 (4)    |
| C3—C2                                | 1.393 (3)    | C5—H5       | 0.9500       |
| C3—C4                                | 1.397 (3)    | C5—C6       | 1.398 (4)    |
| C7—H7                                | 0.9500       | C6—H6       | 0.9500       |
|                                      |              |             |              |
| I1 <sup>i</sup> —Pt1—I1              | 180.0        | C7—C2—S1    | 115.60 (17)  |
| S1 <sup>i</sup> —Pt1—I1              | 94.358 (14)  | S1—C1—H1A   | 109.5        |
| S1—Pt1—I1 <sup>i</sup>               | 94.359 (14)  | S1—C1—H1B   | 109.5        |
| S1—Pt1—I1                            | 85.641 (14)  | S1—C1—H1C   | 109.5        |
| S1 <sup>i</sup> —Pt1—I1 <sup>i</sup> | 85.643 (14)  | H1A—C1—H1B  | 109.5        |
| S1—Pt1—S1 <sup>i</sup>               | 180.0        | H1A—C1—H1C  | 109.5        |
| C2—S1—Pt1                            | 104.52 (7)   | H1B—C1—H1C  | 109.5        |
| C2—S1—C1                             | 103.46 (11)  | C3—C4—H4    | 119.8        |
| C1—S1—Pt1                            | 111.00 (8)   | C5—C4—C3    | 120.4 (2)    |
| C2—C3—H3                             | 120.3        | C5—C4—H4    | 119.8        |
| C2—C3—C4                             | 119.3 (2)    | C4—C5—H5    | 120.1        |
| C4—C3—H3                             | 120.3        | C4—C5—C6    | 119.7 (2)    |
| C2—C7—H7                             | 120.2        | C6—C5—H5    | 120.1        |
| C6—C7—H7                             | 120.2        | C7—C6—C5    | 120.4 (2)    |
| C6—C7—C2                             | 119.6 (2)    | C7—C6—H6    | 119.8        |
| C3—C2—S1                             | 123.88 (17)  | C5—C6—H6    | 119.8        |
| C3—C2—C7                             | 120.5 (2)    |             |              |
|                                      |              |             |              |
| Pt1—S1—C2—C3                         | −107.26 (19) | C1—S1—C2—C7 | −168.90 (18) |
| Pt1—S1—C2—C7                         | 74.82 (17)   | C4—C3—C2—S1 | −178.09 (18) |
| C3—C4—C5—C6                          | 0.4 (4)      | C4—C3—C2—C7 | −0.3 (3)     |
| C2—C3—C4—C5                          | 0.0 (4)      | C4—C5—C6—C7 | −0.5 (4)     |
| C2—C7—C6—C5                          | 0.2 (4)      | C6—C7—C2—S1 | 178.21 (19)  |
| C1—S1—C2—C3                          | 9.0 (2)      | C6—C7—C2—C3 | 0.2 (3)      |

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .