



On Stable Capillary Hypersurfaces with Planar Boundaries

Rabah Souam

► To cite this version:

Rabah Souam. On Stable Capillary Hypersurfaces with Planar Boundaries. The Journal of Geometric Analysis, 2023, 33 (6), pp.196. 10.1007/s12220-023-01257-2 . hal-04087042

HAL Id: hal-04087042

<https://hal.science/hal-04087042>

Submitted on 2 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON STABLE CAPILLARY HYPERSURFACES WITH PLANAR BOUNDARIES

RABAH SOUAM

ABSTRACT. We study stable immersed capillary hypersurfaces Σ in domains $\mathcal{B}$ of $\mathbb{R}^{n+1}$ bounded by hyperplanes. When $\mathcal{B}$ is a half-space, we show Σ is a spherical cap. When $\mathcal{B}$ is a domain bounded by k hyperplanes $P_1, \dots, P_k$, $2 \leq k \leq n+1$, having independent normals, and Σ has contact angle θ_i with P_i and does not touch the edges of $\mathcal{B}$, we prove there exists $\delta > 0$, depending only on $P_1, \dots, P_k$, so that if $\theta_i \in (\frac{\pi}{2} - \delta, \frac{\pi}{2} + \delta)$ for each i , then Σ has to be a piece of a sphere.

1. INTRODUCTION

A capillary hypersurface in a domain $\mathcal{B}$ of a Riemannian manifold is a constant mean curvature (CMC) hypersurface in $\mathcal{B}$ meeting $\partial\mathcal{B}$ at a constant angle. Capillary hypersurfaces are critical points for an energy functional for variations which preserve the *enclosed volume*. More precisely, given an angle $\theta \in (0, \pi)$, for a compact hypersurface Σ inside $\mathcal{B}$ such that $\partial\Sigma \subset \partial\mathcal{B}$ and $\partial\Sigma$ bounds a compact domain W in $\partial\mathcal{B}$, the energy of Σ is by definition the quantity

$$E(\Sigma) := |\Sigma| - \cos \theta |W|.$$

Here and throughout this paper, we use the notation $|M|$ to denote the volume of a Riemannian manifold M . The energy functional E can be defined for general immersed hypersurfaces, not only those with boundary bounding a compact domain in $\partial\mathcal{B}$. Details can be found in [1]. The stationary hypersurfaces of E for variations preserving the *enclosed volume* are precisely the CMC hypersurfaces making a constant angle θ with $\partial\mathcal{B}$.

Existence and uniqueness problems of capillary hypersurfaces are interesting on their own right. In the minimal case, when $\mathcal{B}$ is the unit ball in $\mathbb{R}^3$ and the angle of contact is $\frac{\pi}{2}$, these questions proved to be related to deep problems in Geometric Analysis, see for instance the work of Fraser-Schoen [4]. When $\mathcal{B}$ is a domain in $\mathbb{R}^3$, capillary surfaces correspond to models of incompressible liquids inside a container $\mathcal{B}$ in the absence of gravity. The free interface of the liquid (locally) minimizes the energy functional E . It is then a natural problem to study the stable ones, that is, those for which the second variation of the energy functional is non-negative for all volume preserving variations. Stability issues for capillary hypersurfaces have been recently actively addressed in different ambient

1991 *Mathematics Subject Classification.* 49Q10, 53A10.

Key words and phrases. Capillary hypersurfaces, constant mean curvature hypersurfaces, stability.

manifolds and domains, see for instance [1, 3, 5, 7, 8, 9, 10]. One of the achievements in this direction is the proof by Wang-Xia [10] that in a unit ball in a space form, totally umbilical hypersurfaces are the only stable capillary hypersurfaces (Nunes [8] previously solved the free boundary case in a unit ball in $\mathbb{R}^3$).

In this paper, we are interested in domains $\mathcal{B}$ in $\mathbb{R}^{n+1}$ bounded by a finite family of hyperplanes. The first case we consider is when $\mathcal{B}$ is a half-space in $\mathbb{R}^{n+1}$. In this case, it was shown by Ainouz-Souam [1] that a stable capillary hypersurface with contact angle $\theta < \frac{\pi}{2}$ is a spherical cap provided each of its boundary components is embedded. Choe-Koiso [3] obtained the same result when $\theta > \frac{\pi}{2}$ under the assumption that the boundary is convex and Marinov [7] treated the case $n = 2$ assuming the boundary is embedded. In our main result (Theorem 3.2) we remove the boundary embeddedness assumption and characterize the spherical caps as the only stable immersed capillary hypersurfaces supported by a hyperplane in $\mathbb{R}^{n+1}$, $n \geq 2$. The second case we study is when $\mathcal{B}$ is a domain in $\mathbb{R}^{n+1}$ bounded by a finite number of hyperplanes $P_1, \dots, P_k$ with linearly independent normals. Li-Xiong [5] showed that, in this situation, a stable capillary hypersurface meeting each P_i with a constant angle $\theta_i \in [\pi/2, \pi]$ and not touching the edges of $\mathcal{B}$ is a piece of a sphere under the assumption that its boundary is embedded for $n = 2$ and that each of its boundary components is convex for $n \geq 3$. The case $k = 2$, that is, when $\mathcal{B}$ is a wedge, was previously obtained by Choe-Koiso [3]. We here prove (Theorem 3.4) the existence of a positive number $\delta > 0$ such that if Σ is a stable immersed capillary hypersurface in $\mathcal{B}$ not touching the edges of $\mathcal{B}$ and making an angle $\theta_i \in (\frac{\pi}{2} - \delta, \frac{\pi}{2} + \delta)$ with each P_i , then Σ is a piece of a sphere. We emphasize that in our results we do not assume the boundary of the hypersurfaces is embedded.

2. PRELIMINARIES

We here recall briefly some basic facts about capillary hypersurfaces and refer to [1] for more details.

Let $n \geq 2$ be an integer and $\mathcal{B}$ a domain in $\mathbb{R}^{n+1}$ bounded by a finite number $P_1, \dots, P_k$ of hyperplanes, $k \geq 1$. A capillary hypersurface in $\mathcal{B}$ is a compact and constant mean curvature (CMC) hypersurface contained in $\mathcal{B}$ with boundary on $\partial\mathcal{B}$ and meeting each P_i at a constant angle $\theta_i \in (0, \pi)$. Consider a capillary hypersurface defined by a smooth immersion $\psi : \Sigma \rightarrow \mathcal{B}$ and let N be a global unit normal to Σ along ψ chosen so that its (constant) mean curvature satisfies $H \geq 0$. Write $\partial\Sigma = \bigcup_{i=1}^k \Gamma_i$, as a union of compact submanifolds with $\psi(\Gamma_i) \subset P_i$. By an edge of $\mathcal{B}$ we mean a non empty intersection $P_i \cap P_j$ with $i \neq j$. We will suppose that the hypersurface does not touch the edges of $\mathcal{B}$ so that $\psi(\partial\Sigma)$ lies on the smooth parts of $\partial\mathcal{B}$ where the unit normal is well defined (see however the comments following Theorem 3.4).

Capillary immersions in $\mathcal{B}$ are critical points of an energy functional for deformations through immersions in $\mathcal{B}$, mapping $\partial\Sigma$ into $\partial\mathcal{B}$ and preserving the *enclosed volume*. When ψ is an embedding with each $\psi(\Gamma_i)$ bounding a domain $W_i \subset P_i$, the energy writes:

$$E(\psi) = |\psi(\Sigma)| - \sum_{i=1}^k \cos \theta_i |W_i|$$

and the enclosed volume is the volume of the domain in $\mathbb{R}^{n+1}$ bounded by $\psi(\Sigma)$ and $\cup_{i=1}^k W_i$. This energy functional and the notion of enclosed volume can be extended in a suitable way to include general immersions, details can be found in [1].

We denote by $\mathbf{n}_i$ the exterior unit normal to P_i . The angle of contact along Γ_i is the angle $\theta_i \in (0, \pi)$ between N and $\mathbf{n}_i$. We let ν_i be the exterior unit normal to Γ_i in Σ and $\bar{\nu}_i$ be the unit normal to Γ_i in $\partial\mathcal{B}$ chosen so that $\{N, \nu_i\}$ and $\{\mathbf{n}_i, \bar{\nu}_i\}$ have the same orientation in $(T\partial\Sigma)^\perp$. The angle between ν_i and $\bar{\nu}_i$ is also equal to θ_i .

The index form $\mathcal{I}$ of ψ is the symmetric bilinear form defined on $\mathcal{C}^\infty(\Sigma)$ by

$$\mathcal{I}(f, g) = - \int_{\Sigma} f (\Delta g + |\sigma|^2 g) dA + \sum_{i=1}^k \int_{\Gamma_i} f \left(\frac{\partial g}{\partial \nu_i} - q_i g \right) ds,$$

where Δ stands for the Laplacian of the metric induced by ψ , σ is the second fundamental form of ψ and

$$q_i = \cot \theta_i \sigma(\nu_i, \nu_i).$$

The capillary immersion ψ as above is said to be stable if the second variation of the energy E is nonnegative for all volume preserving deformations. This condition is equivalent to

$$\mathcal{I}(f, f) \geq 0 \quad \text{for all } f \in \mathcal{C}^\infty(\Sigma) \quad \text{with} \quad \int_{\Sigma} f dA = 0,$$

see [1]. We note that we can take this as a definition of stability for capillary immersions supported by $\partial\mathcal{B}$ but not necessarily contained in $\mathcal{B}$.

Let D denote the standard covariant differentiation on $\mathbb{R}^{n+1}$. A basic well-known fact we will use is the following (see for instance [1]):

Proposition 2.1. *Let $\mathcal{B}$ be as above and $\psi : \Sigma \rightarrow \mathcal{B}$ a capillary immersion into $\mathcal{B}$. Then, the unit conormal ν_i along Γ_i is a principal direction of ψ , that is, $D_{\nu_i} N = -\sigma(\nu_i, \nu_i) \nu_i$.*

3. THE RESULTS

A key ingredient we will use is the following fact obtained in [1], we include the proof for completeness.

Proposition 3.1. *Let $\psi : \Sigma \rightarrow \mathbb{R}^{n+1}$ be a smooth immersion in the Euclidean space $\mathbb{R}^{n+1}$ of a smooth compact orientable n -dimensional manifold Σ with or*

without boundary. Let $N : \Sigma \rightarrow \mathbb{S}^n \subset \mathbb{R}^{n+1}$ be a global unit normal of ψ and ν the exterior unit conormal to $\partial\Sigma$ in Σ . Then

$$(3.1) \quad n \int_{\Sigma} N dA = \int_{\partial\Sigma} \{ \langle \psi, \nu \rangle N - \langle \psi, N \rangle \nu \} ds$$

where dA and ds denote the volume elements of Σ and $\partial\Sigma$, respectively.

In particular, if Σ has no boundary, then

$$\int_{\Sigma} N dA = 0.$$

Proof. Let a be a constant vector field on $\mathbb{R}^{n+1}$. Consider the following vector field on Σ

$$X = \langle a, N \rangle \psi^T - \langle \psi, N \rangle a^T,$$

where, $\psi^T = \psi - \langle \psi, N \rangle N$ (resp. $a^T = a - \langle a, N \rangle N$) is the projection of ψ (resp. of a) on $T\Sigma$. Using the following well known formulas that can be thoroughly checked:

$$\operatorname{div} \psi^T = n + nH \langle \psi, N \rangle, \quad \operatorname{div} a^T = nH \langle a, N \rangle,$$

we compute the divergence of X :

$$\begin{aligned} \operatorname{div} X &= \langle a, N \rangle \operatorname{div} \psi^T + \langle a, D_{\psi^T} N \rangle - \langle \psi, N \rangle \operatorname{div} a^T - \langle a^T, N \rangle - \langle \psi, D_{a^T} N \rangle \\ &= n \langle a, N \rangle + nH \langle \psi, N \rangle \langle a, N \rangle + \langle a^T, D_{\psi^T} N \rangle - nH \langle a, N \rangle \langle \psi, N \rangle - \langle \psi^T, D_{a^T} N \rangle \\ &= n \langle a, N \rangle, \end{aligned}$$

where we used the symmetry of the Weingarten map, that is, $\langle a^T, D_{\psi^T} N \rangle = \langle \psi^T, D_{a^T} N \rangle$. Integrating on Σ and using the divergence theorem we get

$$n \int_{\Sigma} \langle a, N \rangle dA = \int_{\partial\Sigma} \{ \langle a, N \rangle \langle \psi, \nu \rangle - \langle \psi, N \rangle \langle a, \nu \rangle \} ds$$

Since this is true for any a , (3.1) follows. $\square$

We now prove our first result characterizing the spherical caps as the only stable immersed capillary hypersurfaces in half-spaces in $\mathbb{R}^{n+1}$. This was proved under additional assumptions on the boundary and the angle of contact in [1], [3] and [7]. We note here that it is not necessary to assume the hypersurface is contained in a half-space, we only need the fact that it meets a hyperplane at a constant angle, so we state the result in this more general form.

Theorem 3.2. *Let $\psi : \Sigma \rightarrow \mathbb{R}^{n+1}$ be a capillary immersion of a compact and connected orientable manifold Σ of dimension n in $\mathbb{R}^{n+1}$, $n \geq 2$, supported by a hyperplane. If the immersion is stable then $\psi(\Sigma)$ is a spherical cap.*

Proof. We denote, as usual, by $e_1, \dots, e_{n+1}$ the vectors of the canonical basis of $\mathbb{R}^{n+1}$. Without loss of generality, we may suppose the supporting hyperplane is the hyperplane $\{x_{n+1} = 0\}$ oriented by the unit normal $-e_{n+1}$. The starting point is to derive a suitable test function inspired by the work of Barbosa-do Carmo [2] proving that round spheres are the only compact and closed stable CMC hypersurfaces in $\mathbb{R}^{n+1}$.

Integrating the equation $\operatorname{div} \psi^T = n + nH \langle \psi, N \rangle$, we get

$$(3.2) \quad \int_{\partial\Sigma} \langle \psi, \nu \rangle ds = n \int_{\Sigma} \{1 + H\langle \psi, N \rangle\} dA.$$

On $\partial\Sigma$, we have $\cos \theta N + \sin \theta \nu = -e_{n+1}$, $\langle \psi, \nu \rangle = \cos \theta \langle \psi, \bar{\nu} \rangle$ and $\langle \psi, N \rangle = -\sin \theta \langle \psi, \bar{\nu} \rangle$. Proposition 3.1 gives in this case

$$(3.3) \quad n \cos \theta \int_{\Sigma} N dA = - \left(\int_{\partial\Sigma} \langle \psi, \nu \rangle ds \right) e_{n+1}.$$

From (3.2) and (3.3), we conclude that:

$$\int_{\Sigma} \{1 + H\langle \psi, N \rangle + \cos \theta \langle N, e_{n+1} \rangle\} dA = 0.$$

So we may use $\phi := 1 + H\langle \psi, N \rangle + \cos \theta \langle N, e_{n+1} \rangle$ as a test function in the stability inequality. Set $u = \langle \psi, N \rangle$ and $v = \langle N, e_{n+1} \rangle$. These functions satisfy the following well-known formulas

$$(3.4) \quad \Delta u + |\sigma|^2 u = -nH,$$

and

$$(3.5) \quad \Delta v + |\sigma|^2 v = 0.$$

Using these equations, we compute:

$$\begin{aligned} \Delta \phi &= H(-nH - |\sigma|^2 u) - \cos \theta |\sigma|^2 v \\ &= -nH^2 - |\sigma|^2 (Hu + \cos \theta v) \\ &= -nH^2 - |\sigma|^2 (\phi - 1). \end{aligned}$$

Therefore

$$\Delta \phi + |\sigma|^2 \phi = (|\sigma|^2 - nH^2).$$

Moreover:

$$\frac{\partial u}{\partial \nu} = \langle \nu, N \rangle + \langle \psi, D_{\nu} N \rangle = -\sigma(\nu, \nu) \langle \psi, \nu \rangle$$

and

$$\frac{\partial v}{\partial \nu} = \langle D_{\nu} N, e_{n+1} \rangle = -\sigma(\nu, \nu) \langle \nu, e_{n+1} \rangle = \sigma(\nu, \nu) \sin \theta,$$

Now, taking into account that, along $\partial\Sigma$: $\langle \psi, \nu \rangle = \cos \theta \langle \psi, \bar{\nu} \rangle$ and $\langle \psi, N \rangle = -\sin \theta \langle \psi, \bar{\nu} \rangle$, one can check after direct computations that:

$$(3.6) \quad \frac{\partial \phi}{\partial \nu} = \cot \theta \sigma(\nu, \nu) \phi.$$

It follows that

$$(3.7) \quad \mathcal{I}(\phi, \phi) = - \int_{\Sigma} (|\sigma|^2 - nH^2) \phi dA = - \int_{\Sigma} (|\sigma|^2 - nH^2) (1 + H\langle \psi, N \rangle + \cos \theta \langle N, e_{n+1} \rangle) dA.$$

We now introduce the following function:

$$F = H|\psi|^2 + 2\langle \psi, N \rangle.$$

One has $\Delta|\psi|^2 = 2n(1 + H\langle\psi, N\rangle)$ and $\Delta\langle\psi, N\rangle = -|\sigma|^2\langle\psi, N\rangle - nH$, so that

$$\begin{aligned}\Delta F &= 2nH(1 + H\langle\psi, N\rangle) + 2(-|\sigma|^2\langle\psi, N\rangle - nH) \\ &= 2(nH^2 - |\sigma|^2)\langle\psi, N\rangle\end{aligned}$$

Integrating we get

$$2 \int_{\Sigma} (nH^2 - |\sigma|^2)\langle\psi, N\rangle dA = \int_{\Sigma} \Delta F dA = \int_{\partial\Sigma} \frac{\partial F}{\partial \nu} ds,$$

that is,

$$(3.8) \quad \int_{\Sigma} (nH^2 - |\sigma|^2)\langle\psi, N\rangle dA = \int_{\partial\Sigma} (H - \sigma(v, v))\langle\psi, v\rangle ds.$$

Denote by $H_{\partial\Sigma}$ the mean curvature of the immersion $\psi|_{\partial\Sigma} : \partial\Sigma \rightarrow \mathbb{R}^n \times \{0\}$ computed with respect to the unit normal $\bar{v}$. We claim that, on $\partial\Sigma$:

$$(3.9) \quad H - \sigma(v, v) = -(n-1)(H + \sin\theta H_{\partial\Sigma}).$$

Indeed, let $i \in \{1, \dots, k\}$ and $\{v_1, \dots, v_{n-1}\}$ be a local orthonormal frame on $\partial\Sigma$. Then:

$$\sigma(v, v) = nH - \sum_{j=1}^{n-1} \sigma(v_j, v_j).$$

Now, considering the unit normal $\bar{v}$ to $\partial\Sigma$ in $\mathbb{R}^n \times \{0\}$ along ψ , as chosen in Sec. 2, we have $N = -\sin\theta \bar{v} - \cos\theta e_{n+1}$. We can thus write

$$\sigma(v_j, v_j) = -\langle \nabla_{v_j} N, v_j \rangle = \sin\theta \langle \nabla_{v_j} \bar{v}, v_j \rangle.$$

Therefore the following relation holds on $\partial\Sigma$

$$\sigma(v, v) = nH + (n-1) \sin\theta H_{\partial\Sigma}.$$

This proves (3.9). Using (3.9) and the fact that $\langle\psi, v\rangle = \cos\theta\langle\psi, \bar{v}\rangle$, we can write (3.8) as follows:

$$(3.10) \quad \int_{\Sigma} (nH^2 - |\sigma|^2)\langle\psi, N\rangle dA = -(n-1) \cos\theta \int_{\partial\Sigma} (H + \sin\theta H_{\partial\Sigma})\langle\psi, \bar{v}\rangle ds.$$

We claim that

$$(3.11) \quad \int_{\partial\Sigma} (H + \sin\theta H_{\partial\Sigma})\langle\psi, \bar{v}\rangle ds = 0.$$

To see this, we first apply Minkowski formula to the immersion $\psi|_{\partial\Sigma} : \partial\Sigma \rightarrow \mathbb{R}^n \times \{0\}$, obtaining: $\int_{\partial\Sigma} H_{\partial\Sigma} \langle\psi, \bar{v}\rangle ds = -|\partial\Sigma|$. Thus

$$\int_{\partial\Sigma} (H + \sin\theta H_{\partial\Sigma})\langle\psi, \bar{v}\rangle ds = H \int_{\partial\Sigma} \langle\psi, \bar{v}\rangle ds - \sin\theta |\partial\Sigma|.$$

Integrating the relation $\Delta\psi = nHN$, we obtain

$$\int_{\partial\Sigma} v ds = nH \int_{\Sigma} N dA.$$

Therefore

$$\int_{\partial\Sigma} \langle v, e_{n+1} \rangle ds = nH \int_{\Sigma} \langle N, e_{n+1} \rangle dA,$$

that is,

$$(3.12) \quad -\sin \theta |\partial \Sigma| = nH \int_{\Sigma} \langle N, e_{n+1} \rangle dA.$$

Along $\partial \Sigma$, we have $\langle \psi, \nu \rangle = \cos \theta \langle \psi, \bar{\nu} \rangle$ and $\langle \psi, N \rangle = -\sin \theta \langle \psi, \bar{\nu} \rangle$. So (3.1) gives

$$(3.13) \quad n \int_{\Sigma} \langle N, e_{n+1} \rangle dA = - \int_{\partial \Sigma} \langle \psi, \bar{\nu} \rangle ds.$$

(3.12) and (3.13) show that

$$H \int_{\partial \Sigma} \langle \psi, \bar{\nu} \rangle ds - \sin \theta |\partial \Sigma| = 0.$$

This proves (3.11). So (3.10) gives

$$\int_{\Sigma} (nH^2 - |\sigma|^2) \langle \psi, N \rangle = 0.$$

Going back to (3.7), we get

$$\mathcal{I}(\phi, \phi) = - \int_{\Sigma} (|\sigma|^2 - nH^2) (1 + \cos \theta \langle N, e_{n+1} \rangle).$$

As $\theta \in (0, \pi)$, we have $1 + \cos \theta \langle N, e_{n+1} \rangle > 0$. Also $|\sigma|^2 - nH^2 \geq 0$ with equality only at umbilics. Hence $\mathcal{I}(\phi, \phi) \leq 0$. However, by the stability assumption, $\mathcal{I}(\phi, \phi) \geq 0$. We conclude that necessarily $|\sigma|^2 - nH^2 \equiv 0$ and so $\psi(\Sigma)$ is totally umbilical, that is, a spherical cap. $\square$

Remark 3.3. More generally we can consider capillary immersions in $\mathbb{R}^{n+1}$ supported by a hyperplane so that the angle of contact is locally constant, that is, this angle is constant along each boundary component. It is an open problem to decide if spherical caps are the only stable capillary immersions of this type. Using the argument of Theorem 3.1 in [1] (see also [7]) one shows this is the case for surfaces of genus zero in $\mathbb{R}^3$.

In the proof of Theorem 3.2 we strongly used in our computations that the origin lies on the supporting hyperplane. This explains the hypothesis in our second result dealing with a domain bounded by a finite family of hyperplanes. It was inspired by the work of Li-Xiong [5].

Theorem 3.4. *Let $\mathcal{B}$ be a domain in $\mathbb{R}^{n+1}$ bounded by k hyperplanes $P_1, \dots, P_k$, $2 \leq k \leq n+1$, having linearly independent normals. Then there exists a constant $\delta > 0$ so that if $\psi : \Sigma \rightarrow \mathcal{B}$ is a stable compact and connected orientable immersed capillary hypersurface in $\mathcal{B}$ not touching the edges of $\mathcal{B}$ and having a contact angle $\theta_i \in (\frac{\pi}{2} - \delta, \frac{\pi}{2} + \delta)$ with P_i , then $\psi(\Sigma)$ is a part of a sphere.*

Proof. Let $\mathbf{n}_i$ denote, as above, the exterior unit normal to $\mathcal{B}$ along P_i . The independence of $\mathbf{n}_1, \dots, \mathbf{n}_k$ ensures that $\cap_{i=1}^k P_i \neq \emptyset$ and so we may assume, without loss of generality, that the origin lies in $\cap_{i=1}^k P_i$. Following [5], we consider a linear combination $a := \sum_{i=1}^n c_i \mathbf{n}_i$ verifying: $\langle a, \mathbf{n}_i \rangle = -\cos \theta_i$, $i = 1, \dots, k$. The vector a exists and is unique by the independence of $\mathbf{n}_1, \dots, \mathbf{n}_k$. We consider the function

$$\phi = 1 + H \langle \psi, N \rangle + \langle a, N \rangle.$$

Proceeding as in the proof of Theorem 3.2, one shows that $\int_{\Sigma} \phi dA = 0$ and

$$\mathcal{I}(\phi, \phi) = - \int_{\Sigma} (|\sigma|^2 - nH^2) \phi dA = - \int_{\Sigma} (|\sigma|^2 - nH^2) (1 + H\langle \psi, N \rangle + \langle a, N \rangle) dA.$$

We claim that

$$(3.14) \quad \int_{\Sigma} (|\sigma|^2 - nH^2) H \langle \psi, N \rangle dA = 0.$$

To prove this, write $\partial\Sigma = \cup_{i=1}^k \Gamma_i$, with $\psi(\Gamma_i) \subset P_i$. As in the proof of Theorem 3.2, considering the function $F = H|\psi|^2 + 2\langle \psi, N \rangle$, one shows that

(3.15)

$$\int_{\Sigma} (nH^2 - |\sigma|^2) \langle \psi, N \rangle dA = -(n-1) \sum_{i=1}^k \cos \theta_i \int_{\Gamma_i} (H + \sin \theta_i H_{\Gamma_i}) \langle \psi, \bar{v}_i \rangle ds,$$

where H_{Γ_i} is the mean curvature of the immersion $\psi|_{\Gamma_i} : \Gamma_i \rightarrow P_i$ with respect to the unit normal $\bar{v}_i$. As before, Proposition 3.1 together with the relation $\langle \psi, v_i \rangle = \cos \theta \langle \psi, \bar{v}_i \rangle$ on Γ_i , give

$$(3.16) \quad n \int_{\Sigma} N dA = \sum_{i=1}^k \left(\int_{\Gamma_i} \langle \psi, \bar{v}_i \rangle dA \right) \mathbf{n}_i.$$

Integrating the relation $\Delta\psi = nHN$, we obtain

$$nH \int_{\Sigma} N dA = \sum_{i=1}^k \int_{\Gamma_i} v_i ds = \sum_{i=1}^k \int_{\Gamma_i} (\cos \theta_i \bar{v}_i + \sin \theta_i \mathbf{n}_i) ds.$$

By Proposition 3.1 applied to the immersion $\psi|_{\Gamma_i} : \Gamma_i \rightarrow P_i$, one has $\int_{\Gamma_i} \bar{v}_i ds = 0$. So,

$$(3.17) \quad nH \int_{\Sigma} N dA = \sum_{i=1}^k \sin \theta_i |\Gamma_i| \mathbf{n}_i.$$

From (3.16), (3.17) and the independence of the $\mathbf{n}_i$'s, we infer that

$$(3.18) \quad \int_{\Gamma_i} H \langle \psi, \bar{v}_i \rangle ds = \sin \theta_i |\Gamma_i|.$$

Applying Minkowski formula to the immersion $\psi|_{\Gamma_i} : \Gamma_i \rightarrow P_i$, we get

$$(3.19) \quad |\Gamma_i| = - \int_{\Gamma_i} H_{\Gamma_i} \langle \psi, \bar{v}_i \rangle ds.$$

(3.18) and (3.19) give

$$\int_{\Gamma_i} (H + \sin \theta_i H_{\Gamma_i}) \langle \psi, \bar{v}_i \rangle ds = 0.$$

Together with (3.15) this proves (3.14). Summarizing, we have proved that

$$\mathcal{I}(\phi, \phi) = - \int_{\Sigma} (|\sigma|^2 - nH^2) (1 + \langle a, N \rangle) dA.$$

As in the proof of Theorem 3.2, we can conclude that $\psi(\Sigma)$ is totally umbilical provided $|a| < 1$. Now, recall that a is the unique solution, in the linear space spanned by $\mathbf{n}_1, \dots, \mathbf{n}_k$, to the system of linear equations $\langle a, \mathbf{n}_i \rangle = -\cos \theta_i$, $i = 1, \dots, k$, and depends continuously on the θ_i 's. When

$\theta_i = \frac{\pi}{2}$, for $i = 1, \dots, k$, the solution to the system is $a = 0$. Therefore there exists $\delta > 0$ such that if $\theta_i \in (\frac{\pi}{2} - \delta, \frac{\pi}{2} + \delta)$, $i = 1, \dots, k$, then $|a| < 1$ and $\psi(\Sigma)$ is totally umbilical. Note that $\psi(\Sigma)$ cannot be planar because otherwise it would not be disjoint from the edges of $\mathcal{B}$. Therefore $\psi(\Sigma)$ is a piece of a sphere. $\square$

Conversely, a part of a sphere as above is stable and even a minimizer of the energy functional E among all embedded hypersurfaces in $\mathcal{B}$ enclosing the same volume, see [11].

In the free boundary case, that is, when $\theta_i = \frac{\pi}{2}$, for each $i = 1, \dots, k$, the only possibility for a piece of a sphere to meet orthogonally the hyperplanes P_i is to be centered at a point in the intersection $\cap_{i=1}^k P_i$. The conclusion of Theorem 3.4 is therefore that a stable hypersurface with free boundary has to touch some of the edges of $\mathcal{B}$. The case of a wedge, i.e. when $k = 2$, was considered by Lopez [6].

It is possible to restate Theorem 3.4 for immersions not necessarily contained in the domain bounded by the hyperplanes and possibly touching its edges. More precisely, let $P_1, \dots, P_k$, $2 \leq k \leq n+1$ be a family of hyperplanes in $\mathbb{R}^{n+1}$, oriented by given unit normals $\mathbf{n}_1, \dots, \mathbf{n}_k$, respectively. We consider capillary immersions $\psi : \Sigma \rightarrow \mathbb{R}^{n+1}$ with $\partial\Sigma$ decomposed into a union $\cup_{i=1}^k \Gamma_i$ of (not necessarily connected) submanifolds with $\psi(\Gamma_i) \subset P_i$ and $\angle(N, \mathbf{n}_i) = \theta_i$ along Γ_i , $i = 1, \dots, k$. As above, N denotes the unit normal to the immersion for which the mean curvature verifies $H \geq 0$. We note that this definition excludes, for instance, the part of a round sphere in a wedge and centered at its edge. Then the conclusion of Theorem 3.4 is valid for such immersions.

REFERENCES

- [1] A. Ainouz and R. Souam. Stable capillary hypersurfaces in a half-space or a slab. *Indiana Univ. Math. J.* **65** (2016), no. 3, 813–831.
- [2] J-L. Barbosa and M. do Carmo. Stability of hypersurfaces with constant mean curvature. *Math. Z.* **185** (1984), no. 3, 339–353.
- [3] J. Choe and M. Koiso. Stable capillary hypersurfaces in a wedge. *Pacific J. Math.* **280** (2016), no. 1, 1–15.
- [4] A. Fraser and R. Schoen. Sharp eigenvalue bounds and minimal surfaces in the ball. *Invent. Math.* **203** (2016), no. 3, 823–890.
- [5] H. Li and C. Xiong. Stability of capillary hypersurfaces with planar boundaries. *J. Geom. Anal.* **27** (2017), no. 1, 79–94.
- [6] R. Lopez. Capillary surfaces with free boundary in a wedge. *Adv. Math.* **262** (2014), 476–483.
- [7] P. Marinov. Stability of capillary surfaces with planar boundary in the absence of gravity. *Pacific J. Math.* **255** (2012), no. 1, 177–190.
- [8] I. Nunes. On stable constant mean curvature surfaces with free boundary. *Math. Z.* **287** (2017), no. 1-2, 473–479.
- [9] R. Souam. Stable constant mean curvature surfaces with free boundary in slabs. *J. Geom. Anal.* **31** (2021), no. 1, 282–297.
- [10] G. Wang and C. Xia. Uniqueness of stable capillary hypersurfaces in a ball. *Math. Ann.* **374** (2019), no. 3-4, 1845–1882.
- [11] R. K. P. Zia ; J. E. Avron and J. E. Taylor. The summertop construction: crystals in a corner. *J. Statist. Phys.* **50** (1988), no. 3-4, 727–736.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, CNRS ET UNIVERSITÉ DE
PARIS, BÂTIMENT SOPHIE GERMAIN, CASE 7012, 75205 PARIS CEDEX 13, FRANCE

Email address: `rabah.souam@imj-prg.fr`