



HAL
open science

Which sensations should you imagine?

François Le Jeune, Léa Driessens, Emile Savalle, Léa Pillette

► **To cite this version:**

François Le Jeune, Léa Driessens, Emile Savalle, Léa Pillette. Which sensations should you imagine?. Collectif pour la Recherche Transdisciplinaire sur les Interfaces Cerveau-Ordinateur CORTICO 2023, May 2023, Institut du Cerveau, Paris, France. hal-04087104v2

HAL Id: hal-04087104

<https://hal.science/hal-04087104v2>

Submitted on 4 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Which sensations should you imagine?

François Le Jeune
Inria, Univ Rennes, IRISA, CNRS
F35000 Rennes, France

Léa Driessens
Inria, Univ Rennes, IRISA, CNRS
F35000 Rennes, France

Emile Savalle
Inria, Univ Rennes, IRISA, CNRS
F35000 Rennes, France

Léa Pillette
Univ Rennes, Inria, CNRS, IRISA
F35000 Rennes, France

During motor imagery-based brain computer interfaces (MI-BCI) training, users are most often instructed to perform kinesthetic motor imagery, i.e., imagine the sensations related to a movement [1]. However, there is a great variety of sensations associated with motor imagery that can either be interoceptive or exteroceptive [2]. Interoceptive sensations related to movement arise from within the body, such as muscle contraction, whereas exteroceptive sensations are sensed through the skin, such as touch or vibration. Among those different sensations, we do not know which to advise MI-BCI users to imagine [3].

Thus, our experiment aims at studying the influence of imagining sensations on neurophysiological activity and user experience. It will consist in a two hours session during which participants (thirty expected) will be equipped with an electroencephalographic headset while performing five MI tasks: interoceptive, exteroceptive and both, with either a unique or multiple exteroceptive sensations, i.e., pressure and vibration (Fig. 1). After all MI conditions, participants will be asked to fill a user-experience questionnaire [4]. The kinesthetic and visual imagery abilities of participants will also be considered in our analyses [5].

Based on previous results, we expect that imagining both exteroceptive and interoceptive sensations leads to a stronger desynchronization over the sensorimotor cortex [6] than imagining interoceptive sensations only [7, 8], in turn stronger than imagining exteroceptive sensations only. Imagining several exteroceptive sensations should also enable a stronger desynchronization than imagining only one [9]. We also expect a correlation between participants' kinesthetic and visual imagery abilities, their neurophysiological results [10, 11] and their preferences.

Our results should provide useful insights on which instructions to give to participants during MI-BCI user training and potentially how to adapt them to the participants' profile.




Figure 1: Experimental conditions according to the type of sensation(s) and the number of exteroceptive sensation(s) to imagine.

References

- Roc, A., Pillette, L., Mladenovic, J., Benaroch, C., N’Kaoua, B., Jeunet, C., & Lotte, F. (2021). A review of user training methods in brain computer interfaces based on mental tasks. *Journal of Neural Engineering*, 18(1), 011002.
- Kessner, S. S., Bingel, U., & Thomalla, G. (2016). Somatosensory deficits after stroke: a scoping review. *Topics in stroke rehabilitation*, 23(2), 136-146.
- Pillette, L., Lotte, F., N’Kaoua, B., Joseph, P. A., Jeunet, C., & Glize, B. (2020). Why we should systematically assess, control and report somatosensory impairments in BCI-based motor rehabilitation after stroke studies. *NeuroImage: Clinical*, 28, 102417.
- Hart, S. G., & Staveland, L. E. (1988). Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In *Advances in psychology* (Vol. 52, pp. 139-183). North-Holland.
- Malouin, F., Richards, C. L., Jackson, P. L., Lafleur, M. F., Durand, A., & Doyon, J. (2007). The Kinesthetic and Visual Imagery Questionnaire (KVIQ) for assessing motor imagery in persons with physical disabilities: a reliability and construct validity study. *Journal of neurologic physical therapy*, 31(1), 20-29.
- Park, S., Ha, J., Kim, D. H., & Kim, L. (2021). Improving motor imagery-based brain-computer interface performance based on sensory stimulation training: an approach focused on poorly performing users. *Frontiers in Neuroscience*, 15, 732545.
- Edwards, L. L., King, E. M., Buetefisch, C. M., & Borich, M. R. (2019). Putting the “sensory” into sensorimotor control: the role of sensorimotor integration in goal-directed hand movements after stroke. *Frontiers in integrative neuroscience*, 16.
- Melnik, A., Hairston, W. D., Ferris, D. P., & König, P. (2017). EEG correlates of sensorimotor processing: independent components involved in sensory and motor processing. *Scientific Reports*, 7(1), 4461.
- Wriessnegger, S. C., Bauernfeind, G., Kurz, E. M., Raggam, P., & Müller-Putz, G. R. (2018). Imagine squeezing a cactus: Cortical activation during affective motor imagery measured by functional near-infrared spectroscopy. *Brain and cognition*, 126, 13-22.
- Vuckovic, A., & Osuagwu, B. A. (2013). Using a motor imagery questionnaire to estimate the performance of a brain-computer interface based on object-oriented motor imagery. *Clinical Neurophysiology*, 124(8), 1586-1595.
- Marchesotti, S., Bassolino, M., Serino, A., Bleuler, H., & Blanke, O. (2016). Quantifying the role of motor imagery in brain-machine interfaces. *Scientific reports*, 6(1), 1-12.