



**HAL**  
open science

# Nonexistence of global weak solutions to semilinear wave equations involving time-dependent structural damping terms

Mokhtar Kirane, Ahmad Fino, Sebti Kerbal

## ► To cite this version:

Mokhtar Kirane, Ahmad Fino, Sebti Kerbal. Nonexistence of global weak solutions to semilinear wave equations involving time-dependent structural damping terms. 2023. hal-04091015

**HAL Id: hal-04091015**

**<https://hal.science/hal-04091015v1>**

Preprint submitted on 6 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Nonexistence of global weak solutions to semilinear wave equations involving time-dependent structural damping terms

Mokhtar Kirane<sup>a,1,1</sup>, Ahmad Z. Fino<sup>b</sup>, Sebti Kerbal<sup>c</sup>, Aymen Laadhari<sup>a</sup>

<sup>a</sup>Department of Mathematics, College of Arts and Sciences, Khalifa University of Science and Technology, Abu Dhabi, United Arab Emirates

<sup>b</sup>College of Engineering and Technology, American University of the Middle East, Egaila 54200, Kuwait

<sup>c</sup>Department of Mathematics, Sultan Qaboos University, FracDiff Research Group (DR/RG/03), Al-Khoud 123, Muscat, P.O. Box 36, Oman

---

## Abstract

We consider a semilinear wave equation involving a time-dependent structural damping term of the form  $\frac{1}{(1+t)^\beta}(-\Delta)^{\sigma/2}u_t$ . Our results show the influence of the parameters  $\beta, \sigma$  on the nonexistence of global weak solutions under assumptions on the given system data.

**Keywords:** Blow-up, damped wave equations, variable coefficients, Fractional Laplacian

**2020 MSC:** Primary: 35B44, 35A01, Secondary: 35L20, 35L71

---

## 1. Introduction

The aim of the paper is to establish a nonexistence of global weak solutions to the Cauchy problem for the following semilinear structurally damped wave equation

$$\begin{cases} u_{tt} - \Delta u + \frac{b_0}{(1+t)^\beta}(-\Delta)^{\sigma/2}u_t = |u|^p & x \in \mathbb{R}^n, t > 0, \\ u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x), & x \in \mathbb{R}^n, \end{cases} \quad (1)$$

where  $0 < \sigma < 2, n \geq 1, p > 1, b_0$  is a positive constant, and  $\beta \in \mathbb{R}$ . Without loss of generality, we assume that  $b_0 = 1$ .

Prior to our main blow-up results, we would like to emphasize that the special case of Eq. (1) for  $b_0 = 0$  is governed by the classical semilinear wave equation, where the Strauss conjecture for this case states that: if  $p \leq p_S$  then there is no global solution for (1) under suitable sign assumptions on the initial data, and for  $p > p_S$  a global solution exists for small initial data; see e.g. [10, 14, 16, 18] among many other references, where  $p_S$  is the positive solution of the following quadratic equation  $(n-1)p^2 - (n+1)p - 2 = 0$ , and is given by

$$p_S = p_S(n) := \frac{n+1 + \sqrt{n^2 + 10n - 7}}{2(n-1)}.$$

In order to have more general initial data, but still with compact support, Kato [11] obtained with a slightly less sharp blow-up result an exponent of the form  $(n+1)/(n-1)$  which is less than the Strauss exponent  $p_S$ , for  $n \geq 2$ .

We would like to emphasize here that the test function method, introduced by [17] and used by [8, 9, 12], plays a similar role as of Kato's method in the proof of blow-up results. In fact, the test function is effective in the case of parabolic equations as it provide exactly the critical exponent  $p_c$ , but in the case of hyperbolic equations (cf. [12]) we

---

\*Corresponding author

Email addresses: mokhtar.kirane@ku.ac.ae (Mokhtar Kirane), ahmad.fino@aum.edu.kw (Ahmad Z. Fino), skerbal@sq.edu.om (Sebti Kerbal), aymen.laadhari@ku.ac.ae (Aymen Laadhari)

get the so-called Kato's exponent  $p^*$  which is less than  $p_c$ . This is one of the weakness of the test function method but in general it can be applied to a more general equation and system with no positive condition on solutions.

When  $\beta = 0$ ,  $\sigma \rightarrow 2$ , and  $b_0 = 1$ , problem (1) is reduced to

$$u_{tt} - \Delta u - \Delta u_t = |u|^p, \quad x \in \mathbb{R}^n, t > 0, \quad (2)$$

which is called the viscoelastic damping case. D'Ambrosio and Lucente [3, Theorem 4.2] proved that the solution of (2) blows-up in finite time when  $1 < p \leq (n+1)/(n-1)_+$ , where  $(\cdot)_+ := \max\{0, \cdot\}$ , by applying the test function method. Similar result has been obtained recently by Fino [6] in the case of an exterior domain. On the other hand, D'Abbicco-Reissig [2] proved that there exists a global solution for (2) when  $p > 1 + \frac{3}{n-1}$  ( $n \geq 2$ ) for sufficiently small initial data. Therefore, the exact value of the critical exponent is still an open question.

Recently, problem (1) with  $\sigma \rightarrow 2$ ,

$$u_{tt} - \Delta u - \frac{b_0}{(1+t)^\beta} \Delta u_t = |u|^p, \quad x \in \mathbb{R}^n, t > 0, \quad (3)$$

has been studied by Fino & Hamza [7]. They proved that if  $\int_{\mathbb{R}^n} u_1(x) dx > 0$ ,

$$\begin{cases} p \in (1, \infty) & \text{for } n = 1, 2, \\ p \in (1, \frac{n}{n-2}] & \text{for } n \geq 3, \end{cases}$$

and

$$\begin{cases} 1 < p \leq \frac{n+1}{(n-1)_+} & \text{if } \beta \geq -1, \\ 1 < p \leq \frac{n(1-\beta)+2}{(n(1-\beta)-2)_+} & \text{if } \beta \leq -1, \end{cases}$$

then the mild solution of (3) blows-up in finite time.

The goal of this paper is to prove the nonexistence of global weak solutions of (1) which is a generalization of the results of [7] to the case of fractional Laplacian damping term by using a recent version of the test function method that relies on Lemma 1 below. The novelty of this paper is the choice of this test function which is consistent with the suppression of positivity usually assumed by many authors.

In the case of  $\beta \neq 0$  and  $b_0 = 1$ , we give an intuitive observation for understanding the influence of the damping term  $(\frac{1}{(1+t)^\beta}(-\Delta)^{\sigma/2} u_t)$  by scaling argument. Let  $u(t, x)$  be a solution of the linear strong damped wave equation

$$u_{tt}(t, x) - \Delta u(t, x) + \frac{1}{(1+t)^\beta} (-\Delta)^{\sigma/2} u_t(t, x) = 0. \quad (4)$$

When  $\sigma \geq 1 - \beta$ , we put

$$u(t, x) = v(\lambda(1+t), \lambda x), \quad \lambda(1+t) = s, \quad \lambda x = y, \quad (5)$$

with a parameter  $\lambda > 0$ , we have

$$v_{ss}(s, y) - \Delta v(s, y) + \frac{\lambda^{\sigma+\beta-1}}{s^\beta} (-\Delta)^{\sigma/2} v_s(s, y) = 0.$$

Thus, when  $\sigma = 1 - \beta$  we notice that Eq. (4) is invariant, while when  $\sigma > 1 - \beta$ , letting  $\lambda \rightarrow 0$ , we obtain the wave equation without damping

$$v_{ss}(s, y) - \Delta v(s, y) = 0.$$

We note that  $\lambda \rightarrow 0$  is corresponding to  $t \rightarrow +\infty$ .  
On the other hand, when  $\sigma < 1 - \beta$ , we put

$$u(t, x) = v(\lambda^{\frac{\sigma}{1-\beta}}(1+t), \lambda x), \quad \lambda^{\frac{2}{1-\beta}}(1+t) = s, \quad \lambda x = y,$$

with a parameter  $\lambda > 0$ , we have

$$v_{ss}(s, y) + \frac{1}{s^\beta}(-\Delta)^{\sigma/2}v_s(s, y) - \lambda^{\frac{2(1-\beta-\sigma)}{1-\beta}}\Delta v(s, y) = 0.$$

In this case, letting  $\lambda \rightarrow 0$ , we obtain the pseudo-parabolic equation

$$v_{ss}(s, y) + \frac{1}{s^\beta}(-\Delta)^{\sigma/2}v_s(s, y) = 0.$$

This means that the asymptotic behaviour of solutions depends on the behaviour of the coefficient of the damping term.

This paper is organized as follows. We start in Sec. 2 by stating the main theorem of our work and we prove it in Sec. 4. Sec. 3 is to collect some preliminaries.

## 2. Main results

This section is aimed to state our main results.

**Theorem 1 (Blow-up).** *We assume that*

$$(u_0, u_1) \in (L^1(\mathbb{R}^n) \cap H^1(\mathbb{R}^n)) \times (L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n))$$

satisfying the following condition:

$$\int_{\mathbb{R}^n} u_1(x) dx > 0. \tag{6}$$

If

$$\begin{cases} 1 < p < \min \left\{ \frac{n+1}{(n-1)_+}; \frac{n}{(n-\sigma)_+} \right\} & \text{if } \sigma \geq 1 - \beta, \\ 1 < p < \min \left\{ \frac{n(1-\beta) + \sigma}{(n(1-\beta) - \sigma)_+}; \frac{n}{(n-\sigma)_+} \right\} & \text{if } \sigma \leq 1 - \beta, \end{cases} \tag{7}$$

or

$$\begin{cases} p = \frac{n+1}{n-1}, & \text{if } \sigma \geq 1 - \beta, \sigma > \frac{2n}{n+1}, n \geq 2, \\ p = \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}, & \text{if } n(1+\beta) < \sigma \leq 1 - \beta, n \geq 1, \end{cases}$$

where  $(\cdot)_+ := \max\{0, \cdot\}$ , then problem (1) has no global weak solutions.

**Remark 1.** *We note that, by taking the limit case  $\sigma \rightarrow 2$  we recover the same results of [7].*

**Remark 2.** *We stress that the exponent  $\frac{n+1}{(n-1)_+}$  appearing in (7) was introduced first in [11] to prove the nonexistence of global solutions to the semilinear wave equation with the nonlinearity  $|u|^p$ , subject to small initial data with compact support.*

### 3. Preliminaries

#### Definition 1. [13]

Let  $\mathcal{S}$  be the Schwartz space of rapidly decaying  $C^\infty$  functions in  $\mathbb{R}^n$  and  $s \in (0, 1)$ . The fractional Laplacian  $(-\Delta)^s$  in  $\mathbb{R}^n$  is a non-local operator defined on  $\mathcal{S}$  by

$$\begin{aligned} (-\Delta)^s v(x) &:= C_{n,s} \text{ p.v.} \int_{\mathbb{R}^n} \frac{v(x) - v(y)}{|x - y|^{n+2s}} dy \\ &= \begin{cases} C_{n,s} \int_{\mathbb{R}^n} \frac{v(x) - v(y)}{|x - y|^{n+2s}} dy, & \text{if } 0 < s < 1/2, \\ C_{n,s} \int_{\mathbb{R}^n} \frac{v(x) - v(y) - \nabla v(x) \cdot (x - y) \chi_{|x-y| < \delta}(y)}{|x - y|^{n+2s}} dy, \quad \forall \delta > 0, & \text{if } 1/2 \leq s < 1, \end{cases} \end{aligned}$$

where p.v. stands for Cauchy's principal value, and  $C_{n,s} := \frac{s 4^s \Gamma(\frac{n}{2} + s)}{\pi^{\frac{n}{2}} \Gamma(1 - s)}$ .

In fact, we are rarely going to use the fractional Laplacian operator in the Schwartz space; it can be extended to less regular functions as follows: for  $s \in (0, 1)$ ,  $\varepsilon > 0$ , let

$$\mathcal{L}_{s,\varepsilon}(\Omega) := \begin{cases} L_s(\mathbb{R}^n) \cap C^{0,2s+\varepsilon}(\Omega) & \text{if } 0 < s < 1/2, \\ L_s(\mathbb{R}^n) \cap C^{1,2s+\varepsilon-1}(\Omega), & \text{if } 1/2 \leq s < 1, \end{cases}$$

where  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $C^{0,2s+\varepsilon}(\Omega)$  is the space of  $2s + \varepsilon$ - Hölder continuous functions on  $\Omega$ ,  $C^{1,2s+\varepsilon-1}(\Omega)$  the space of functions of  $C^1(\Omega)$  whose first partial derivatives are Hölder continuous with exponent  $2s + \varepsilon - 1$ , and

$$L_s(\mathbb{R}^n) = \left\{ u : \mathbb{R}^n \rightarrow \mathbb{R} \text{ such that } \int_{\mathbb{R}^n} \frac{u(x)}{1 + |x|^{n+2s}} dx < \infty \right\}.$$

#### Proposition 1. [13, Proposition 2.4]

Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $s \in (0, 1)$ , and  $f \in \mathcal{L}_{s,\varepsilon}(\Omega)$  for some  $\varepsilon > 0$ . Then  $(-\Delta)^s f$  is a continuous function in  $\Omega$  and  $(-\Delta)^s f(x)$  is given by the pointwise formulas of Definition 1 for every  $x \in \Omega$ .

**Remark:** A simple sufficient condition for function  $f$  to satisfy the conditions in Proposition 1 is that  $f \in L_{loc}^1(\mathbb{R}^n) \cap C^2(\Omega)$ .

Using [4, Lemma 2.11] and its proof, we have the following.

**Lemma 1.** Let  $\langle x \rangle := (1 + (|x| - 1)^4)^{1/4}$  for all  $x \in \mathbb{R}^n$ ,  $n \geq 1$ . Let  $s \in (0, 1]$  and  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function defined by

$$\phi(x) = \begin{cases} 1 & \text{if } |x| \leq 1, \\ \langle x \rangle^{-n-2s} & \text{if } |x| \geq 1. \end{cases} \quad (8)$$

Then

$$\phi \in C^2(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n) \cap H^2(\mathbb{R}^n), \quad \partial_x^2 \phi \in L^\infty(\mathbb{R}^n),$$

and the following estimate holds:

$$\max \{ |\Delta \phi(x)|, |(-\Delta)^s \phi(x)| \} \leq C \phi(x) \quad \text{for all } x \in \mathbb{R}^n.$$

**Lemma 2.** Let  $h$  be a smooth function satisfying  $\partial_x^2 h \in L^\infty(\mathbb{R}^n)$ . For any  $R > 0$ , let  $h_R$  be a function defined by

$$h_R(x) := h(R^{-1}x) \quad \text{for all } x \in \mathbb{R}^n.$$

Then,  $(-\Delta)^s(h_R)$ ,  $s \in (0, 1]$ , satisfies the following scaling property

$$(-\Delta)^s h_R(x) = R^{-2s} (-\Delta)^s h(R^{-1}x), \quad \text{for all } x \in \mathbb{R}^n.$$

**Lemma 3.** [4, Lemma 2.14]

Let  $s, \tilde{s} \in (0, 1)$ ,  $R > 0$  and  $p > 1$ . Then, the following estimate holds

$$\int_{\mathbb{R}^n} (\phi_R(x))^{-\frac{1}{p-1}} |(-\Delta)^{\tilde{s}} \phi_R(x)|^{\frac{p}{p-1}} dx \leq C R^{-\frac{2\tilde{s}p}{p-1} + n},$$

where  $\phi_R(x) := \phi(x/R)$  and  $\phi$  as defined in (8).

#### 4. Proof of Theorem 1

The proof of Theorem 1 relies mainly on the concept of weak solution of the Cauchy problem (1) and the use of the test function method. Let

$$X_{\delta, T} = \{\varphi \in C([0, \infty), H^2(\mathbb{R}^n)) \cap C^1([0, \infty), H^\delta(\mathbb{R}^n)) \cap C^2([0, \infty), L^2(\mathbb{R}^n)), \text{ such that } \text{supp}\varphi \subset Q_T\},$$

where  $Q_T := [0, T] \times \mathbb{R}^n$ , and the homogeneous fractional Sobolev space  $H^\delta(\mathbb{R}^n)$ ,  $\delta \in (0, 2)$ , is defined by

$$H^\delta(\mathbb{R}^n) = \begin{cases} \{u \in L^2(\mathbb{R}^n); (-\Delta)^{\delta/2} u \in L^2(\mathbb{R}^n)\}, & \text{if } \delta \in (0, 1), \\ H^1(\mathbb{R}^n), & \text{if } \delta = 1, \\ \{u \in H^1(\mathbb{R}^n); (-\Delta)^{\delta/2} u \in L^2(\mathbb{R}^n)\}, & \text{if } \delta \in (1, 2), \end{cases}$$

endowed with the norm

$$\|u\|_{H^\delta(\mathbb{R}^n)} = \begin{cases} \|u\|_{L^2(\mathbb{R}^n)} + \|(-\Delta)^{\delta/2} u\|_{L^2(\mathbb{R}^n)}, & \text{if } \delta \in (0, 1), \\ \|u\|_{L^2(\mathbb{R}^n)} + \|\nabla u\|_{L^2(\mathbb{R}^n)}, & \text{if } \delta = 1, \\ \|u\|_{L^2(\mathbb{R}^n)} + \|\nabla u\|_{L^2(\mathbb{R}^n)} + \|(-\Delta)^{\delta/2} u\|_{L^2(\mathbb{R}^n)}, & \text{if } \delta \in (1, 2). \end{cases}$$

The weak formulation associated with (1) reads as follows:

**Definition 2.** (Weak solution)

Let  $T > 0$ , and  $u_0, u_1 \in L^2(\mathbb{R}^n)$ . A function

$$u \in L^1((0, T), L^2(\mathbb{R}^n)) \cap L^p((0, T); L^{2p}(\mathbb{R}^n)),$$

is said to be a weak solution of (1) on  $[0, T] \times \mathbb{R}^n$  if

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dt dx + \int_{\mathbb{R}^n} u_1(x) \varphi(0, x) dx \\ & + \int_{\mathbb{R}^n} u_0(x) (-\Delta)^{\sigma/2} \varphi(0, x) dx - \int_{\mathbb{R}^n} u_0(x) \varphi_t(0, x) dx \\ & = \int_{Q_T} u \varphi_{tt} dt dx - \int_0^T \frac{1}{(1+t)^\beta} \int_{\mathbb{R}^n} u (-\Delta)^{\sigma/2} \varphi_t dx dt \\ & - \int_{Q_T} u \Delta \varphi dt dx + \int_0^T \frac{\beta}{(1+t)^{\beta+1}} \int_{\mathbb{R}^n} u (-\Delta)^{\sigma/2} \varphi dx dt, \end{aligned}$$

holds for all  $\varphi \in X_{\sigma, T}$ . We denote the lifespan for the weak solution by

$$T_w(u_0, u_1) := \sup\{T \in (0, \infty] \text{ for which there exists a unique weak solution } u \text{ to (1)}\}.$$

Moreover, if  $T > 0$  can be arbitrary chosen, i.e.  $T_w(u_0, u_1) = \infty$ , then  $u$  is called a global weak solution of (1).

**Proof of Theorem 1.** Let  $u$  be a global weak solution of (1), that is

$$\begin{aligned}
& \int_{Q_T} |u|^p \varphi \, dt \, dx + \int_{\mathbb{R}^n} u_1(x) \varphi(0, x) \, dx \\
& + \int_{\mathbb{R}^n} u_0(x) (-\Delta)^{\sigma/2} \varphi(0, x) \, dx - \int_{\mathbb{R}^n} u_0(x) \varphi_t(0, x) \, dx \\
& = \int_{Q_T} u \varphi_{tt} \, dt \, dx - \int_0^T \frac{1}{(1+t)^\beta} \int_{\mathbb{R}^n} u (-\Delta)^{\sigma/2} \varphi_t \, dx \, dt \\
& - \int_{Q_T} u \Delta \varphi \, dt \, dx + \int_0^T \frac{\beta}{(1+t)^{\beta+1}} \int_{\mathbb{R}^n} u (-\Delta)^{\sigma/2} \varphi \, dx \, dt,
\end{aligned} \tag{9}$$

for all  $T > 0$ , and all  $\varphi \in X_{\sigma, T}$ .

Let  $T > 0$ . Now, we introduce the following test function

$$\varphi(t, x) = \psi^\eta(t) \phi_{T^d}(x) = \psi^\eta(t) \phi\left(\frac{x}{T^d}\right) \tag{10}$$

where  $\phi$  is defined in (8),  $\psi(t) = \Psi(\frac{t}{T})$ ,  $\eta \gg 1$ ,  $d > 0$  are constants that will be determined later, and  $\Psi \in C^\infty(\mathbb{R}_+)$  is a cut-off non-increasing function such that

$$\Psi(r) = \begin{cases} 1 & \text{if } 0 \leq r \leq 1/2, \\ \searrow & \text{if } 1/2 \leq r \leq 1, \\ 0 & \text{if } r \geq 1. \end{cases}$$

From the formulation (9), we get the following inequality

$$\begin{aligned}
& \int_{Q_T} |u|^p \varphi \, dt \, dx + \int_{\mathbb{R}^n} u_1(x) \varphi(0, x) \, dx \\
& \leq \int_{\frac{T}{2}}^T \int_{\mathbb{R}^n} |u| |\varphi_{tt}| \, dx \, dt + \int_{Q_T} |u| |\Delta \varphi| \, dt \, dx \\
& + \int_{\frac{T}{2}}^T \frac{1}{(1+t)^\beta} \int_{\mathbb{R}^n} |u| |(-\Delta)^{\sigma/2} \varphi_t| \, dx \, dt + \int_0^T \frac{\beta}{(1+t)^{\beta+1}} \int_{\mathbb{R}^n} |u| |(-\Delta)^{\sigma/2} \varphi| \, dx \, dt \\
& + \int_{\mathbb{R}^n} |u_0| (|(-\Delta)^{\sigma/2} \varphi(0, x)| + |\varphi_t(0, x)|) \, dx \\
& =: I_1 + I_2 + I_3 + I_4 + I_5.
\end{aligned} \tag{11}$$

Let  $\varepsilon > 0$ . By applying  $\varepsilon$ -Young's inequality

$$AB \leq \varepsilon A^p + C(\varepsilon, p) B^{p'}, \quad A \geq 0, \quad B \geq 0, \quad p + p' = pp', \quad C(\varepsilon, p) = (p-1)(\varepsilon p^p)^{-1/(p-1)},$$

we obtain the estimation for the first integral

$$\begin{aligned}
I_1 & \leq \int_{Q_T} |u| \varphi^{1/p} \varphi^{-1/p} \phi_{T^d} |(\psi^\eta)_{tt}| \, dt \, dx \\
& \leq \varepsilon \int_{Q_T} |u|^p \varphi \, dt \, dx + C \int_{Q_T} \phi_{T^d} \psi^{-\eta/(p-1)} |(\psi^\eta)_{tt}|^{p'} \, dt \, dx.
\end{aligned} \tag{12}$$

As  $(\psi^\eta)_{tt} = \eta \psi^{\eta-1} \psi_{tt} + \eta(\eta-1) \psi^{\eta-2} |\psi_t|^2$ , the inequality (12) becomes

$$I_1 \leq \varepsilon \int_{Q_T} |u|^p \psi \, dt \, dx + C \int_{Q_T} \phi_{T^d} \psi^{\eta-p'} |\psi_{tt}|^{p'} \, dt \, dx + C \int_{Q_T} \phi_{T^d} \psi^{\eta-2p'} |\psi_t|^{2p'} \, dt \, dx. \tag{13}$$

Proceeding similarly as for (13), we get

$$I_2 \leq \varepsilon \int_{Q_T} |u|^p \varphi \, dt \, dx + C \int_{Q_T} \psi^\eta \phi_{T^d}^{-1/(p-1)} |\Delta \phi_{T^d}|^{p'} \, dt \, dx. \quad (14)$$

In the same way, we write

$$\begin{aligned} I_3 &= \int_{\frac{T}{2}}^T \frac{1}{(1+t)^\beta} \int_{\mathbb{R}^n} |u| \varphi^{1/p} \varphi^{-1/p} |(-\Delta)^{\sigma/2} \phi_{T^d}| |(\psi^\eta)_t| \, dx \, dt \\ &\leq \varepsilon \int_{Q_T} |u|^p \varphi \, dt \, dx \\ &\quad + C \int_{\frac{T}{2}}^T \frac{1}{(1+t)^{\beta p'}} \int_{\mathbb{R}^n} \phi_{T^d}^{-1/(p-1)} |(-\Delta)^{\sigma/2} \phi_{T^d}|^{p'} \psi^{\eta-p'} |\psi_t|^{p'} \, dx \, dt. \end{aligned} \quad (15)$$

Clearly,

$$\frac{1}{(1+t)^{\beta p'}} \leq C T^{-\beta p'}, \quad \forall t \in \left(\frac{T}{2}, T\right),$$

therefore,

$$I_3 \leq \varepsilon \int_{Q_T} |u|^p \varphi \, dt \, dx + C T^{-\beta p'} \int_{Q_T} \phi_{T^d}^{-1/(p-1)} |(-\Delta)^{\sigma/2} \phi_{T^d}|^{p'} \psi^{\eta-p'} |\psi_t|^{p'} \, dt \, dx. \quad (16)$$

In the same manner,

$$\begin{aligned} I_4 &\leq C \int_0^T \frac{1}{(1+t)^{\beta+1}} \int_{\mathbb{R}^n} |u| \varphi^{1/p} \varphi^{-1/p} \psi^\eta |(-\Delta)^{\sigma/2} \phi_{T^d}| \, dx \, dt \\ &\leq \varepsilon \int_{Q_T} |u|^p \varphi \, dt \, dx + C \int_0^T \frac{1}{(1+t)^{(\beta+1)p'}} \int_{\mathbb{R}^n} \phi_{T^d}^{-1/(p-1)} |(-\Delta)^{\sigma/2} \phi_{T^d}|^{p'} \psi^\eta \, dx \, dt. \end{aligned} \quad (17)$$

Finally, it remains only to control the term  $I_5$ . By exploiting the identity  $\varphi_t(0, x) = \eta \psi_t(0) \phi_{T^d}(x)$ , we infer

$$I_5 \leq C \int_{\mathbb{R}^n} |u_0| \left( |(-\Delta)^{\sigma/2} \phi_{T^d}| + |\psi_t(0) \phi_{T^d}| \right) \, dx. \quad (18)$$

Plugging (11) together with (13)-(18) and choosing  $\varepsilon$  small enough, we deduce that

$$\begin{aligned} &\int_{Q_T} |u|^p \varphi \, dx \, dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) \, dx \\ &\leq C \int_{Q_T} \phi_{T^d} \psi^{\eta-p'} |\psi_{tt}|^{p'} \, dt \, dx + C \int_{Q_T} \phi_{T^d} \psi^{\eta-2p'} |\psi_t|^{2p'} \, dt \, dx \\ &\quad + C \int_{Q_T} \psi^\eta \phi_{T^d}^{-1/(p-1)} |\Delta \phi_{T^d}|^{p'} \, dt \, dx \\ &\quad + C T^{-\beta p'} \int_{Q_T} \phi_{T^d}^{-1/(p-1)} |(-\Delta)^{\sigma/2} \phi_{T^d}|^{p'} \psi^{\eta-p'} |\psi_t|^{p'} \, dt \, dx \\ &\quad + C \int_0^T \frac{1}{(1+t)^{(\beta+1)p'}} \int_{\mathbb{R}^n} \phi_{T^d}^{-1/(p-1)} |(-\Delta)^{\sigma/2} \phi_{T^d}|^{p'} \psi^\eta \, dx \, dt \\ &\quad + C \int_{\mathbb{R}^n} |u_0| \left( |(-\Delta)^{\sigma/2} \phi_{T^d}| + |\psi_t(0) \phi_{T^d}| \right) \, dx. \end{aligned}$$

Taking account of the expression of  $\varphi$  given by (10) and Lemmas 1-3, we infer that

$$\begin{aligned} &\int_{Q_T} |u|^p \varphi \, dx \, dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) \, dx \\ &\leq C T^{-2p'+1+nd} + C T^{-2dp'+1+nd} + C T^{-\beta p' - p' - \sigma dp' + 1 + nd} \\ &\quad + C T^{-\sigma dp' + nd} \int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} \, dt + C (T^{-\sigma d} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| \, dx. \end{aligned} \quad (19)$$



Since  $0 < \sigma < 2$ , we notice that the cases of  $\sigma \geq 1 - \beta$  and  $\sigma \leq 1 - \beta$  are equivalent to

$$\sigma \geq 1 - \beta, \quad \text{for all } \beta > -1, \quad \text{and} \quad \sigma < 1 - \beta, \quad \text{for all } \beta < 1,$$

therefore, we distinguish two cases:

**I. Case:  $\sigma \geq 1 - \beta$ , for all  $\beta > -1$ .**

In this case, we choose  $d = 1$ .

Subcritical case  $p < \min\{\frac{n+1}{(n-1)_+}; \frac{n}{(n-\sigma)_+}\}$ .

Note that,

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C \begin{cases} T^{1-\frac{(\beta+1)p}{p-1}} & \text{if } \beta p < -1, \\ \ln T & \text{if } \beta p = -1, \\ 1 & \text{if } \beta p > -1. \end{cases}$$

We have two cases to distinguish.

a) If  $\beta \geq 0$ . In this case, we have  $\beta p > -1$  and so

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C.$$

Therefore, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_T(x) dx \\ & \leq C T^{-2p'+1+n} + C T^{-(\beta+1)p' - \sigma p' + 1 + n} + C T^{-\sigma p' + n} + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

Using the fact  $\beta p > -1 \implies -(\beta+1)p' - \sigma p' + 1 + n < -\sigma p' + n$ , we conclude that

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_T(x) dx \\ & \leq C T^{-2p'+1+n} + C T^{-\sigma p' + n} + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned} \quad (20)$$

Note that, we can easily see that

$$-2p' + 1 + n < 0 \iff p < \frac{n+1}{(n-1)_+} \quad \text{and} \quad -\sigma p' + n < 0 \iff p < \frac{n}{(n-\sigma)_+}.$$

Letting  $T \rightarrow \infty$ , and using the Lebesgue dominated convergence theorem together with  $u_1 \in L^1(\mathbb{R}^n)$ , we conclude that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

b) If  $-1 < \beta < 0$ . We have three cases for  $n$ .



Note that when  $n = 1$ , we have  $n = 1 < \frac{1-\beta}{1+\beta}$ .

i) If  $n \geq \frac{\sigma}{1+\beta}$ , then  $n \geq \frac{1-\beta}{1+\beta}$  and therefore

$$p < \min \left\{ \frac{n+1}{n-1}; \frac{n}{n-\sigma} \right\} \leq -\frac{1}{\beta}, \quad \text{for all } n \geq 2,$$

i.e.  $\beta p > -1$ , which implies a contradiction by following the same calculations as in part a).

ii) If  $\frac{1-\beta}{1+\beta} < n < \frac{\sigma}{1+\beta}$  when  $\sigma > 1 - \beta$ , then we have


$$n < \frac{\sigma}{1+\beta} < \frac{\sigma}{2-\sigma},$$

and therefore

$$p < \min \left\{ \frac{n+1}{n-1}; \frac{n}{n-\sigma} \right\} = \frac{n+1}{n-1} < -\frac{1}{\beta}, \quad \text{for all } n \geq 2,$$

i.e.  $\beta p > -1$ , which implies a contradiction by following the same calculations as in part a).

iii) If  $n \leq \frac{1-\beta}{1+\beta}$ , then  $n \leq \frac{\sigma}{1+\beta}$ , i.e.



• If  $p \leq -1/\beta$ , i.e.  $\beta p \geq -1$ , then

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C \ln T, \quad \text{for all } T \gg 1.$$

Then, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_T(x) dx \\ & \leq C T^{-2p'+1+n} + C T^{-(\beta+1)p' - \sigma p' + 1+n} + C T^{-\sigma p' + n} \ln T + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \end{aligned}$$

for all  $T \gg 1$ . As  $\beta p \geq -1 \implies (\beta+1)p' \geq 1$ , we conclude that

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_T(x) dx \\ & \leq C T^{-2p'+1+n} + C T^{-\sigma p' + n} \ln T + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \quad \forall T \gg 1. \end{aligned} \quad (21)$$

Note that, we can easily see that

$$-2p' + 1 + n < 0 \iff p < \frac{n+1}{(n-1)_+} \quad \text{and} \quad -\sigma p' + n < 0 \iff p < \frac{n}{(n-\sigma)_+}.$$

Letting  $T \rightarrow \infty$ , using the fact that  $\ln T \leq T^{\frac{\sigma p' - n}{2}}$  (because  $p < \frac{n}{(n-\sigma)_+}$ ) and the Lebesgue dominated convergence theorem, we conclude that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

• If  $p > -1/\beta$ , i.e.  $p\beta < -1$ , then

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C T^{1-\frac{(\beta+1)p}{p-1}}, \quad \text{for all } T > 0.$$

Then, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_T(x) dx \\ & \leq C T^{-2p'+1+n} + C T^{-(\beta+1)p' - \sigma p' + 1+n} + C(T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \end{aligned}$$

for all  $T \gg 1$ . As  $\sigma \geq 1 - \beta \implies n \leq \frac{1-\beta}{1+\beta} \leq \frac{\sigma}{2-\sigma}$ , we conclude that

$$p < \min \left\{ \frac{n+1}{(n-1)_+}; \frac{n}{(n-\sigma)_+} \right\} = \frac{n+1}{(n-1)_+} \leq \frac{n+1}{(n-\beta-\sigma)_+} \implies -(\beta+1)p' - \sigma p' + 1+n < 0,$$

and

$$p < \min \left\{ \frac{n+1}{(n-1)_+}; \frac{n}{(n-\sigma)_+} \right\} = \frac{n+1}{(n-1)_+} \iff -2p' + 1+n < 0.$$

Note that  $\sigma \geq 1 - \beta > 1$ , and  $2 > \sigma > \sigma + \beta$ , therefore

$$\begin{cases} (n-\beta-\sigma)_+ = (n-1)_+ = (n-\sigma)_+ = 0, & \text{when } n = 1, \\ (n-\beta-\sigma)_+ = n-\beta-\sigma, (n-1)_+ = n-1, (n-\sigma)_+ = n-\sigma, & \text{when } n \geq 2, \end{cases}$$

Letting  $T \rightarrow \infty$ , using the Lebesgue dominated convergence theorem, we conclude that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

Critical case:  $p < \infty$  and  $\sigma \geq 1$  when  $n = 1$ , or  $p = \frac{n+1}{n-1}$  and  $\sigma > \frac{2n}{n+1}$  when  $n \geq 2$ .

Note that when  $2 \leq n < \frac{\sigma}{2-\sigma}$ , we have

$$p = \frac{n+1}{n-1} < \frac{n}{n-\sigma} \quad \text{i.e.} \quad -\sigma p' + n < 0,$$

and when  $\sigma \geq 1$  and  $n = 1$ , we also have  $-\sigma p' + n < 0$ .

We have two cases to distinguish.

a) If  $\beta \geq 0$ . In this case, we have  $\beta p > -1$  and so

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C.$$

From the subcritical case, we can see that we have


$$u \in L^p((0, \infty); L^p(\mathbb{R}^n)). \quad (22)$$

On the other hand, by applying Hölder's inequality instead of Young's inequality, we get

$$\begin{aligned} \int_{\mathbb{R}^n} u_1(x)\phi_T(x) dx &\leq C \left( \int_{\frac{T}{2}}^T \int_{\mathbb{R}^n} |u|^p \varphi dx dt \right)^{1/p} + C \left( \int_0^T \int_{|x|\geq T} |u|^p \varphi dx dt \right)^{1/p} \\ &\quad + C T^{-\sigma p' + n} + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

Letting  $T \rightarrow \infty$  and taking into consideration (22), we get a contradiction.

b) If  $-1 < \beta < 0$ . We have three cases for  $n$ .



Note that when  $n = 1$ , we have  $n = 1 < \frac{1-\beta}{1+\beta}$ .

i) If  $n > \frac{1-\beta}{1+\beta}$ , then

$$p = \frac{n+1}{n-1} < -\frac{1}{\beta},$$

i.e.  $\beta p > -1$ , which implies a contradiction by following the same calculations as in part a).

ii) If  $n = \frac{1-\beta}{1+\beta}$ , then

$$p = \frac{n+1}{n-1} = -\frac{1}{\beta},$$

i.e.  $\beta p = -1$ , and therefore

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C \ln T, \quad \text{for all } T > 0.$$

From the subcritical case (20), we can see easily that we have

$$u \in L^p((0, \infty); L^p(\mathbb{R}^n)). \quad (23)$$

On the other hand, by applying Hölder's inequality instead of Young's inequality and using  $p = \frac{n+1}{n-1}$ , we get

$$\begin{aligned} \int_{\mathbb{R}^n} u_1(x)\phi_T(x) dx &\leq C \left( \int_{\frac{T}{2}}^T \int_{\mathbb{R}^n} |u|^p \varphi dx dt \right)^{1/p} + C \left( \int_0^T \int_{|x|\geq T} |u|^p \varphi dx dt \right)^{1/p} \\ &\quad + C T^{-\sigma p' + n} \ln T + C (T^{-\sigma} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

Letting  $T \rightarrow \infty$  and taking into consideration (23) and the fact that  $\ln T \leq T^{\frac{\sigma p' - n}{2}}$  (because  $p < \frac{n}{n-\sigma}$ ), we get

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

iii) If  $n < \frac{1-\beta}{1+\beta}$ , then

$$p = \frac{n+1}{(n-1)_+} > -\frac{1}{\beta},$$

i.e.  $p\beta < -1$ . In this case, we change the test function  $\psi$  by  $\psi(t) = \Psi\left(\frac{t}{K^{-1}T}\right)$  where  $K \geq 1$  is independent of  $T$ . Then

$$\int_0^{K^{-1}T} (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C K^{-1+(\beta+1)p'} T^{1-(\beta+1)p'}, \quad \text{for all } T > 0.$$

From the subcritical case (20), we can see easily that we have

$$u \in L^p((0, \infty); L^p(\mathbb{R}^n)). \quad (24)$$

On the other hand, by applying Hölder's inequality in (19) instead of Young's inequality, we get

$$\begin{aligned} \int_{\mathbb{R}^n} u_1(x)\phi_T(x) dx &\leq C K^{2p'-1} \left( \int_{\frac{K^{-1}T}{2}}^{K^{-1}T} \int_{\mathbb{R}^n} |u|^p \varphi dx dt \right)^{1/p} + C K^{-1} \left( \int_0^{K^{-1}T} \int_{|x| \geq T} |u|^p \varphi dx dt \right)^{1/p} \\ &\quad + C K^{-1+(\beta+1)p'} T^{-(\beta+1)p' - \sigma p' + 1 + n} + C (K^\sigma T^{-\sigma} + KT^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

As  $\sigma \geq 1 - \beta$  and  $2 > \sigma > \sigma + \beta$ , we conclude that

$$p = \frac{n+1}{n-1} \leq \frac{n+1}{n-\beta-\sigma} \implies -(\beta+1)p' - \sigma p' + 1 + n \leq 0, \quad \text{when } n \geq 2,$$

and

$$-(\beta+1)p' - \sigma p' + 1 + n < 0, \quad \text{when } n = 1,$$

so

$$\begin{aligned} \int_{\mathbb{R}^n} u_1(x)\phi_T(x) dx &\leq C K^{2p'-1} \left( \int_{\frac{K^{-1}T}{2}}^{K^{-1}T} \int_{\mathbb{R}^n} |u|^p \varphi dx dt \right)^{1/p} + C K^{-1} \left( \int_0^{K^{-1}T} \int_{|x| \geq T} |u|^p \varphi dx dt \right)^{1/p} \\ &\quad + C K^{-1+(\beta+1)p'} + C (K^\sigma T^{-\sigma} + KT^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

Letting  $T \rightarrow \infty$  and taking into consideration (24), we get

$$\int_{\mathbb{R}^n} u_1(x) dx \leq C K^{-1+(\beta+1)p'}.$$

Letting  $K \rightarrow \infty$  and using  $p\beta < -1 \implies -1 + (\beta+1)p' < 0$ , we infer that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

**II. Case:  $\sigma < 1 - \beta$ , for all  $\beta < 1$ .**

In this case we take  $d = \frac{1-\beta}{\sigma} > 1$ .

Subcritical case  $p < \min \left\{ \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}; \frac{n}{n-\sigma} \right\}$ .

Note that,

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C \begin{cases} T^{1-\frac{(\beta+1)p}{p-1}} & \text{if } \beta p < -1, \\ \ln T & \text{if } \beta p = -1, \\ 1 & \text{if } \beta p > -1. \end{cases}$$

We have two cases to distinguish.

a) If  $0 \leq \beta < 1$ . In this case, we have  $\beta p > -1$  and so

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C.$$

Therefore, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C T^{-\frac{2(1-\beta)}{\sigma}p'+1+\frac{n(1-\beta)}{\sigma}} \\ & \quad + C T^{-(1-\beta)p'+\frac{n(1-\beta)}{\sigma}} + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

As

$$\sigma < 1 - \beta \implies -\frac{2(1-\beta)}{\sigma}p' + 1 + \frac{n(1-\beta)}{\sigma} < -2p' + 1 + \frac{n(1-\beta)}{\sigma},$$

we conclude that

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C T^{(1-\beta)[-p'+\frac{n}{\sigma}]} + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \end{aligned}$$

Note that, we can easily see that


$$(1-\beta) \left[ -p' + \frac{n}{\sigma} \right] < 0 \iff p < \frac{n}{(n-\sigma)_+} \quad \text{and} \quad -2p' + 1 + \frac{n(1-\beta)}{\sigma} < 0 \iff p < \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}.$$

Note that  $n(1-\beta) - \sigma \geq 1 - \beta - \sigma > 0$ , for all  $n \geq 1$ . Letting  $T \rightarrow \infty$ , and using the Lebesgue dominated convergence theorem together with  $u_1 \in L^1(\mathbb{R}^n)$ , we conclude that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

b) If  $\beta < 0$ . We have two cases for  $n$ .



Note that  $n(1-\beta) - \sigma \geq 1 - \beta - \sigma > 0$  and  $\sigma < \beta + 1 < 1 \leq n \implies n - \sigma > 0$ , for all  $n \geq 1$ .

i) If  $n \geq \frac{\sigma}{(1+\beta)_+}$ , we have

$$p < \min \left\{ \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}; \frac{n}{n-\sigma} \right\} = \frac{n}{n-\sigma} \leq -\frac{1}{\beta},$$

i.e.  $\beta p > -1$ , which implies a contradiction by following the same calculations as in part a).


ii) If  $n < \frac{\sigma}{(1+\beta)_+}$ , we get

$$p < \min \left\{ \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}; \frac{n}{n-\sigma} \right\} = \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma},$$

and

$$-\frac{1}{\beta} < \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma},$$

i.e.



- If  $p \leq -1/\beta$ , i.e.  $p\beta \geq -1$ , then

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C \ln T, \quad \text{for all } T \gg 1.$$

Then, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C T^{-\frac{2(1-\beta)}{\sigma}p'+1+\frac{n(1-\beta)}{\sigma}} \\ & \quad + C T^{-(1-\beta)p'+\frac{n(1-\beta)}{\sigma}} \ln T + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \end{aligned}$$

for all  $T \gg 1$ . As

$$\sigma < 1 - \beta \implies -\frac{2(1-\beta)}{\sigma}p'+1 + \frac{n(1-\beta)}{\sigma} < -2p'+1 + \frac{n(1-\beta)}{\sigma},$$

we conclude that

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C T^{(1-\beta)[-p'+\frac{n}{\sigma}]} \ln T + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx, \end{aligned}$$

Note that, we can easily see that

$$(1-\beta) \left[ -p' + \frac{n}{\sigma} \right] < 0 \iff p < \frac{n}{n-\sigma} \quad \text{and} \quad -2p'+1 + \frac{n(1-\beta)}{\sigma} < 0 \iff p < \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}.$$

Letting  $T \rightarrow \infty$ , using the fact that  $\ln T \leq T^{\frac{(1-\beta)(p'-\frac{n}{\sigma})}{2}}$  (because  $p < \frac{n}{n-\sigma}$ ) and the Lebesgue dominated convergence theorem, we conclude that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6).

- If  $p > -1/\beta$ , i.e.  $p\beta < -1$ , then

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C T^{1-\frac{(\beta+1)p}{p-1}}, \quad \text{for all } T > 0.$$

Then, (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C T^{-\frac{2(1-\beta)}{\sigma}p'+1+\frac{n(1-\beta)}{\sigma}} + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

As

$$\sigma < 1 - \beta \implies -\frac{2(1-\beta)}{\sigma}p' + 1 + \frac{n(1-\beta)}{\sigma} < -2p' + 1 + \frac{n(1-\beta)}{\sigma},$$

we conclude that

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi \, dx \, dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) \, dx \\ & \leq C T^{-2p'+1+\frac{n(1-\beta)}{\sigma}} + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| \, dx, \end{aligned}$$

Note that, we can easily see that

$$-2p' + 1 + \frac{n(1-\beta)}{\sigma} < 0 \iff p < \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}.$$

Letting  $T \rightarrow \infty$ , using the Lebesgue dominated convergence theorem, we conclude that

$$\int_{\mathbb{R}^n} u_1(x) \, dx \leq 0.$$

This contradicts our assumption (6).

Critical case:  $p = \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma}$  and  $\sigma > n(1+\beta)$ .

Note that when  $n(1+\beta) < \sigma$ , we have

$$p = \frac{n(1-\beta) + \sigma}{n(1-\beta) - \sigma} < \frac{n}{n-\sigma} \quad \text{i.e.} \quad -\sigma p' + n < 0,$$

We have two cases to distinguish.

a) If  $0 \leq \beta < 1$ . In this case, we have  $\beta p > -1$  and so

$$\int_0^T (1+t)^{-\frac{(\beta+1)p}{p-1}} \, dt \leq C.$$

From the subcritical case (20), we can see easily that we have

$$u \in L^p((0, \infty); L^p(\mathbb{R}^n)). \tag{25}$$

On the other hand, by applying Hölder's inequality instead of Young's inequality, (19) implies

$$\begin{aligned} \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) \, dx & \leq C \left( \int_{\frac{T}{2}}^T \int_{\mathbb{R}^n} |u|^p \varphi \, dx \, dt \right)^{1/p} + C \left( \int_0^T \int_{|x| \geq T^d} |u|^p \varphi \, dx \, dt \right)^{1/p} \\ & \quad + C T^{(1-\beta)[-p'+\frac{n}{\sigma}]} + C (T^{-(1-\beta)} + T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| \, dx. \end{aligned}$$

Letting  $T \rightarrow \infty$  and taking into consideration (25) we get

$$\int_{\mathbb{R}^n} u_1(x) \, dx \leq 0.$$

This contradicts our assumption (6).



b) If  $\beta < 0$ . As  $n(1 + \beta) < \sigma \implies n < \frac{\sigma}{(1 + \beta)_+}$ , so

$$p = \frac{n(1 - \beta) + \sigma}{n(1 - \beta) - \sigma} > -\frac{1}{\beta},$$

i.e.  $p\beta < -1$ . In this case, we change the test function  $\psi$  by  $\psi(t) = \Psi\left(\frac{t}{K^{-1}T}\right)$  where  $K \geq 1$  is independent of  $T$ . Then

$$\int_0^{K^{-1}T} (1+t)^{-\frac{(\beta+1)p}{p-1}} dt \leq C K^{-1+(\beta+1)p'} T^{1-(\beta+1)p'}, \quad \text{for all } T > 0.$$

From the subcritical case (20), we can see easily that we have

$$u \in L^p((0, \infty); L^p(\mathbb{R}^n)). \quad (26)$$

On the other hand, by applying Hölder's inequality in (19) instead of Young's inequality and using  $p = \frac{n(1 - \beta) + \sigma}{n(1 - \beta) - \sigma}$ , (19) implies

$$\begin{aligned} & \int_{Q_T} |u|^p \varphi dx dt + \int_{\mathbb{R}^n} u_1(x) \phi_{T^d}(x) dx \\ & \leq C K^{2p'-1} \left( \int_{\frac{K^{-1}T}{2}}^{K^{-1}T} \int_{\mathbb{R}^n} |u|^p \varphi dx dt \right)^{1/p} + C K^{-1} \left( \int_0^{K^{-1}T} \int_{|x| \geq T^d} |u|^p \varphi dx dt \right)^{1/p} \\ & \quad + C K^{-1+(\beta+1)p'} + C (K^{1-\beta} T^{-(1-\beta)} + K T^{-1}) \int_{\mathbb{R}^n} |u_0(x)| dx. \end{aligned}$$

Letting  $T \rightarrow \infty$  and taking into consideration (26), we get

$$\int_{\mathbb{R}^n} u_1(x) dx \leq C K^{-1+(\beta+1)p'}.$$

Letting  $K \rightarrow \infty$  and using  $p\beta < -1 \implies -1 + (\beta + 1)p' < 0$ , we infer that

$$\int_{\mathbb{R}^n} u_1(x) dx \leq 0.$$

This contradicts our assumption (6). □

## Acknowledgments

M.K. gratefully acknowledges the financial support by KUST through the grant FSU-2021-015. A.L. gratefully acknowledges the financial support by KUST through the grant FSU-2021-027.

## References

- [1] T. Cazenave and A. Haraux *Introduction aux problèmes d'évolution semi-linéaires*, Ellipses, Paris, 1990.
- [2] M. D'Abicco and M. Reissig Semilinear structural damped waves, *Math. Meth. Appl. Sci.*, **37** (2014), 1570–1592.
- [3] L. D'Ambrosio and S. Lucente Nonlinear Liouville theorems for Grushin and Tricomi operators, *Journal Differential Equations*, **123** (2003), 511–541.
- [4] T. A. Dao and A. Z. Fino Blow up results for semi-linear structural damped wave model with nonlinear memory, *Mathematische Nachrichten*, (2022), 1–14.
- [5] M. Bonforte and J. L. Vázquez Quantitative local and global a priori estimates for fractional nonlinear diffusion equations, *Adv. Math.*, **250** (2014), 242–284.

- [6] A. Z. Fino Finite time blow up for wave equations with strong damping in an exterior domain, *Mediterranean Journal of Mathematics*, **17** (2020), no. 6, Paper No. 174, 21 pp.
- [7] A. Z. Fino and Hamza, M.A. Blow-up of solutions to semilinear wave equations with a time-dependent strong damping, *J. Evolution Equations & Control Theory*, doi: 10.3934/eect.2022006.
- [8] A. Fino and G. Karch Decay of mass for nonlinear equation with fractional Laplacian, *J. Monatsh. Math.*, **160** (2010), 375–384.
- [9] A. Z. Fino and M. Kirane Qualitative properties of solutions to a time-space fractional evolution equation, *J. Quarterly of Applied Mathematics*, **70** (2012), 133–157.
- [10] F. John Blow-up of solutions of nonlinear wave equations in three space dimensions, *Manuscripta Math.*, **28** (1979), no. 1-3, 235–268.
- [11] T. Kato Blow-up of solutions of some nonlinear hyperbolic equations, *Comp. Pure Appl. Math.*, **33** (1980), 501–505.
- [12] E. Mitidieri and S. I. Pohozaev A priori estimates and blow-up of solutions to nonlinear partial differential equations and inequalities, *Proc. Steklov. Inst. Math.*, **234** (2001), 1–383.
- [13] L. Silvestre Regularity of the obstacle problem for a fractional power of the Laplace operator, *Comm. Pure Appl. Math.*, **60** (2007), no. 1, 67–112.
- [14] W. A. Strauss Nonlinear scattering theory at low energy, *J. Functional Analysis*, **41** (1981), no. 1, 110–133.
- [15] Y. Wakasugi *On the diffusive structure for the damped wave equation with variable coefficients*, Ph.D thesis, Osaka University, 2014.
- [16] B. Yordanov and Q.S. Zhang Finite time blow up for critical wave equations in high dimensions, *J. Functional Analysis*, **231** (2006), 361–374.
- [17] Qi S. Zhang A blow up result for a nonlinear wave equation with damping: the critical case, *C. R. Acad. Sci. Paris*, **333** (2001), no. 2, 109–114.
- [18] Y. Zhou Blow up of solutions to semilinear wave equations with critical exponent in high dimensions, *Chin. Ann. Math. Ser. B*, **28** (2007), no. 2, 205–212.
- [19] A. Pazy *Semi-groups of linear operators and applications to partial differential equations*, Appl. Math. Sci 44. Springer, New-York, 1983.